Cross-talk between the methylerythritol phosphate and mevalonic acid pathways of isoprenoid biosynthesis in poplar

Hui Wei[1,†], Ali Movahedi[1,†], Chen Xu[1,2], Weibo Sun[1], Pu Wang[1], Dawei Li[1], and Qiang Zhuge[1,*]

1 Co-Innovation Center for Sustainable Forestry in Southern China, Key Laboratory of Forest Genetics & Biotechnology, Ministry of Education, College of Biology and the Environment, Nanjing Forestry University, Nanjing 210037, China; 15850682752@163.com (H.W.); ali_movahedi@njfu.edu.cn (A.M.); xuchenidea@hotmail.com (C.X.); czswb@njfu.edu.cn (W.S.); 18705155218@163.com (P.W.); dwli@njfu.edu.cn (D.L.).

2 Jiangsu Provincial Key Construction Laboratory of Special Biomass Resource Utilization, Nanjing Xiaozhuang University, Nanjing 211171, China

* Correspondence: qzhuge@njfu.edu.cn; Fax: +86-25-8542-8701

† These authors are contributed equally to the first author.

Abstract

Plants use two distinct isoprenoid biosynthesis pathways: the methylerythritol phosphate (MEP) pathway and the mevalonic acid (MVA) pathway. 1-deoxy-D-xylulose5-phosphate synthase (DXS) and 1-deoxy-D-xylulose5-phosphate reductoisomerase (DXR) are the rate-limiting enzymes in the MEP pathway, and 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR) is a key regulatory enzyme in the MVA pathway. Previously, overexpression of *Populus trichocarpa PtDXR* in Nanlin 895 (*Populus × euramericana* cv. 'Nanlin 895') poplar was found to upregulate resistance to salt and drought stresses, and the transgenic poplars showed improved growth. In the present study, *PtHMGR* overexpressors (OEs) exhibited higher expression levels of *DXS*, *DXR*, 1-hydroxy-2-methyl-2-(E)-butenyl-4-diphosphate synthase (*HDS*), and 1-hydroxy-2-methyl-2-(E)-butenyl-4-diphosphate reductase (*HDR*) and lower expression levels of 2-C-methyl-d-erythritol4-phosphate cytidylyltransferase (*MCT*), 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase (*CMK*), and 3-hydroxy-3-methylglutaryl-coenzyme A synthase (*HMGS*) than non-transgenic poplars (NT). However, the poplar *PtDXR*-OEs showed upregulated expression levels of MEP-related genes and downregulated expression of MVA-related genes. Moreover, overexpression of *PtDXR* and *PtHMGR* in poplars caused changes in MVA-derived trans-zeatin-riboside (TZR), isopentenyl adenosine (IPA), castasterone (CS) and 6-deoxocastasterone (DCS), as well as MEP-derived

carotenoids, gibberellins (GAs), and abscisic acid (ABA). In *PtHMGR-OE*s, greater accumulation of geranyl diphosphate synthase (*GPS*) and geranyl pyrophosphate synthase (*GPPS)* transcript levels in the MEP pathway led to an accumulation of MEP-derived isoprenoids, while upregulation of farnesyl diphosphate synthase (*FPS*) expression in the MVA pathway contributed to increased levels of MVA-derived isoprenoids. Similarly, in *PtDXR*-OEs, increased *GPS* and *GPPS* transcript levels in the MEP pathway boosted MEP-derived isoprenoid levels and changes in *FPS* expression affected MVA-derived isoprenoid yields. From these results, we can conclude that cross-talk exists between the MVA and MEP pathways.

Keywords: MEP; MVA; poplar; terpenoids; cross-talk

1. Introduction

In plants, terpenoids including abscisic acid (ABA), gibberellins (GAs), carotene, lycopene, cytokinins (CKs), strigolactones (GRs), and brassinosteroids (BRs) are produced via the methylerythritol phosphate (MEP) and mevalonic acid (MVA) pathways (van *et al.*, 2006; Xie *et al.*, 2008; Henry *et al.*, 2015). These MEP- and MVA-derived terpenoids are involved in plant growth and development, as well as in the response to environmental changes (Kirby & Keasling, 2009; Bouvier *et al.*, 2009). The conversion of isopentenyl diphosphate (IPP) into dimethylallyl diphosphate (DMAPP) catalyzed by isopentenyl diphosphate isomerase (IDI) provides the fundamental materials for production of all isoprenoids (Hemmerlin *et al.*, 2012; Lu *et al.*, 2012; Zhang *et al.*, 2019). Besides, IPP and DMAPP play roles in communication (cross-talk) between the MEP and MVA pathways, and previous research has shown that cross-talk occurs between the MEP and MVA pathways (Huchelmann *et al.*, 2014; Liao *et al.*, 2016).

The MVA pathway occurs in the cytoplasm, ER, as well as peroxisomes and produces sesquiterpenoids and sterols. 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR), which is a rate-limiting enzyme in the MVA pathway can catalyze 3-hydroxy-3-methylglutary-CoA (HMG-CoA) to form MVA (Cowan *et al.*, 1997; Roberts *et al.*, 2007). In addition, the MEP pathway occurs in the chloroplast and is responsible for producing carotenoids, ABA, GAs, and diterpenoids. 1-deoxy-D-xylulose5-phosphate synthase (DXS) and 1-deoxy-D-xylulose5-phosphate reductoisomerase (DXR) are rate-limiting enzymes in the MEP pathway that catalyze D-glyceraldehyde3-phosphate (D-3-P) and pyruvate to form 2-C-methyl-D-erythritol4-phosphate (MEP) (Wang *et al.*, 2012; Yamaguchi *et al.*, 2018).

Terpenoids in plant cells are involved in diverse cellular processes. Sesquiterpene derivatives

such as phytoalexin and volatile oils play important roles in plant growth, development, and

disease resistance (Hain *et al*., 1993; Ren *et al*., 2008). α-Tocopherol (vitamin E) is an

antioxidant that improves human health (Weber *et al*., 1997). Sterols are an important

component of biological membranes, which are linked to metabolism (Huang *et al*., 2008).

Photosynthetic pigments participate in plant photosynthesis and enable the conversion of basic

carbon sources for plant growth (Esteban *et al*., 2015). Plant hormones (such as ABA and GAs)

are connected to plant growth and the responses to biotic and abiotic stresses (Kazan *et al*.,

2015). Paclitaxel is one of the most effective chemotherapy agents for cancer treatment, and

artemisinin is an anti-malarial drug (Kong *et al*., 2015; Kim *et al*., 2016).

Previous metabolic engineering studies have proposed strategies for improving the yields

of specific metabolites in plants. For example, the industrial exploitation of hairy roots has

been used to enhance secondary metabolite production (Zhang *et al*., 2004; Rahman *et al*.,

2009). 3-Hydroxy-3-methylglutaryl-coenzyme A synthase (HMGS) is a second enzyme that

has been genetically engineered to improve terpenoid content. For example, overexpression of

Brassica juncea HMGS in tomato plants leads to upregulated carotenoid and phytosterol levels

(Liao *et al*., 2018). HMGR can be metabolically engineered to regulate the content of

terpenoids (Aharoni *et al*., 2005; Dueber *et al*., 2009). In *Arabidopsis thaliana*, HMGR1 has

been shown to play an important role in the biosynthesis of sterols and triterpenoids through

investigation of *A. thaliana* mutant lines (Suzuki *et al*., 2005). Moreover, overexpression of

PgHMGR1 from ginseng in *A. thaliana* can lead to increased production of sterols and

triterpenes (Kim *et al*., 2014). Transgenic tobacco overexpressing the *HMGR* of *Hevea*

brasiliensis has elevated levels of phytosterol (Schaller *et al*., 1995). Dai *et al*. (2011)

expressed *SmHMGR2* in *Salvia miltiorrhiza*, resulting in the improvement of squalene and

tanshinone contents. Interestingly, transgenic T_0 tomato fruits display elevated levels of

phytosterol with overexpression of *HMGR1* from *A. thaliana*, but only certain phytosterols

increase in mature T_2 fruits (Enfissi *et al*., 2005). DXR can be genetically engineered to

regulate the content of terpenoids and expressed *DXR* in *Arabidopsis* and observed enhanced

flux through the MEP pathway (Carretero-Paulet *et al*., 2006). Overexpression of *A. thaliana*

DXR in *Salvia sclarea* hairy roots increases levels of anthiolimine, a diterpene (Vaccaro *et al*.,

2014). Overexpression of *DXR* in peppermint affects the regulation of monoterpenes in leaf

tissues (Mahmoud *et al*., 2001). Farnesyl diphosphate is the synthetic precursor of numerous

primary metabolites including steroids, polyterpenoid alcohols, and ubiquinone, and represents the branching point of many terpenoid metabolic pathways (Wong *et al*., 2018). In summary, overexpression of genes in the MEP and MVA pathways through metabolic engineering can change the expression levels or activities of related enzymes and metabolic products, which opens up a new direction for plant breeding and acquisition of metabolic products.

To date, most studies have found that the MVA and MEP pathways are located in different parts of plants. The MVA pathway occurs in the cytoplasm and endoplasmic reticulum and involves the biosynthesis of secondary metabolites such as sterols, sesquiterpenes, and triterpenes. Meanwhile, the MEP pathway takes place in plastids and mainly involves the biosynthesis of diterpenes, monoterpenes, carotenoids, and isoprene. Previous studies have shown that the MVA and MEP pathways do not operate in isolation and that some metabolic intermediates that link them can be exchanged through plastid membranes (Laule *et al*., 2003; Liao *et al*., 2006). In this study, to comprehensively investigate such cross-talk in poplar, we determined the transcript levels of MEP- and MVA-related genes in *PtHMGR*-OEs and *PtDXR*-OEs. The results showed that not only were the transcript levels of MVA-related genes (such as acetoacetyl CoA thiolase [*AACT*], mevalonate kinase [*MVK*], and mevalonate 5-diphosphate decarboxylase [*MVD*]) upregulated in *PtHMGR*-OEs, but also the expression levels of MEP-related genes (such as *DXS*, *DXR*, *HDS*, and *HDR*) were improved. In *PtDXR*-OEs, the transcript levels of MVA-related genes were downregulated, while the expression levels of MEP-related genes were increased. Based on these results, we propose that the MEP pathway is dominant and the MVA pathway is subordinate. Overexpression of *PtDXR* in poplar increases the expression levels of MEP-related genes, which may result in the accumulation of terpenoids, and then feedback with the MVA pathway may lead to the inhibited expression of MVA-related genes. In addition, the transcript levels of MEP-related genes and products, such as MEP-derived isoprenoids, were affected in *PtHMGR*-OEs. These results demonstrate that cross-talk occurs between the MEP and MVA pathways.

2. Results

2.1. Isolation of the *PtHMGR* gene and characterization of transgenic poplars

The *PtHMGR* gene (Potri.004G208500.1) was isolated from *P. trichocarpa*. The amino acid sequence of PtHMGR contains similar domains, including HMG-CoA-binding motifs (EMPVGYVQIP' and 'TTEGCLVA) and NADPH-binding motifs (DAMGMNMV' and 'VGTVGGGT) (Ma *et al*., 2012) (Supplemental Figure 1).

In addition, the results of Southern blotting indicated a copy number of at least three

copies for *PtHMGR* in the poplar genome (Figure 1A). As shown in Figure 1B, probe 2 could

specifically recognize *PtHMGR* (Potri.004G208500.1), indicating that the probe 2 used for

qRT-PCR of *PtHMGR* could be applied to the follow-up study of *PtHMGR* expression levels.

Based on the whole genome of 10 species achieved from Phytozome, an evolution tree was

constructed and showed the evolutionary relationship of those species (Figure 2A). Also, the

phylogenetic tree illustrated that the function and evolution of the HMGR family are relatively

strongly conserved among all plants (Figure 2B). Also, the HMGR originated from a single

ancestor and evolved into two different groups, one each in monocotyledons and dicotyledons

(Ferrero *et al.*, 2015; Li *et al.*, 2014). This result shows that the evolutionary tree constructed

from HMGR is consistent with that constructed from the whole genomes of various species

(Figure 2).

Analyses of gel electrophoresis illustrated that the *PtHMGR* gene was inserted into the

genome of "Nanlin 895" plants obtained through screening with kanamycin (Kan) using the

primer for the cauliflower mosaic virus (CaMV) 35S promoter (Supplemental Table 1) as the

forward primer and that for *PtHMGR*-R (Supplemental Table 1) as the reverse primer

(Supplemental Figure 1). Besides, the transcript levels of *PtHMGR* were higher in transgenic

lines than in NT plants, showing that *PtHMGR* could be stably expressed in Nanlin 895 plants

(Supplemental Figure 2). Overall, *PtHMGR* was inserted into poplar chromosomes and

expressed in poplar cells.

2.2. Effects of *PtHMGR* overexpression on the expression levels of MVA- and MEP-related

genes in poplar *PtHMGR*-OE seedlings

MVA-related genes (*AACT*, *HMGS*, *MVK*, and *MVD*) were significantly upregulated in

PtHMGR-OE3 and *PtHMGR*-OE7 relative to NT poplars (Figure 3). The expressions of

MEP-related genes (*DXS*, *DXR*, *HDS*, *HDR*, *MCT*, and *CMK*) were also increased in

PtHMGR-OE3 and *PtHMGR*-OE7 relative to NT poplars (Figure 3). Moreover, the expression

levels of genes (*FPS*, *IDI*, *GPS*, and *GPPS*) involved in the biosynthesis of C10 and C20

universal isoprenoid precursors were significantly elevated in *PtHMGR*-OE3 and

PtHMGR-OE7 (Figure 3).

2.3. *PtDXR* overexpression affects expression levels of MVA- and MEP-related genes in

transgenic poplars

When determining the transcript levels of MVA- and MEP-related genes in

160 *PtDXR*-OE1–1 and *PtDXR*-OE3–1 via qRT-PCR, we found that the expression levels of *DXS*,

161 *HDS*, *HDR*, *MCT*, and *CMK* genes involved in the MEP pathway were significantly greater in

162 transgenic poplars. Notably, MVA-related genes (*AACT*, *HMGS*, *HMGR*, *MVK*, *MVD*) were

163 downregulated in *PtDXR*-OE seedlings (Figure 4), the opposite of the MVA-related gene

164 expression results obtained from *PtHMGR*-OEs. In addition, the expression levels of *FPS*, *IDI*,

165 *GPS*, and *GPPS* were significantly elevated in both *PtDXR*-OE1–1 and *PtDXR*-OE3–1

166 compared to the NT poplars (Figure 4).

167 2.4. Overexpression of *PtHMGR* leads to accumulation of MVA-derived TZR, IPA, and DCS

168 as well as MEP-derived carotenoids, ABA, and GAs

169  Analyses of carotenoids, ABA, GAs, zeatin, TZR, IPA, CS, and DCS, which are involved

170 in plant growth and development as well as the responses to biotic and abiotic stresses, would

171 help to elucidate plant growth and the stress response process and could guide poplar breeding.

172 The HPLC-MS/MS chromatograms of ABA, GA, zeatin, TZR, IP, IPA, CS, and DCS standards

173 are provided in Supplemental Figures 3–13.

174  The expression levels of ABA synthesis-related genes including 9-cis-epoxycarotenoid

175 dioxygenase (*NCED*) and zeaxanthin epoxidase (*ZEP*) in *HMGR*-OE and NT leaves were

176 analyzed via qRT-PCR, and ABA contents of *PtHMGR*-OE and NT leaves were analyzed via

177 HPLC-MS/MS. The transcript levels of *NCED1*, *NCED3*, *NCED5*, *NCED6*, *ZEP1*, *ZEP2*, and

178 *ZEP3* in *PtHMGR*-OE poplars were elevated compared to that in NT poplars (Figure 3). In

179 addition, ABA levels were significantly elevated in lipid extracts from *HMGR*-OE poplar

180 leaves (Figure 5). The contents of β-carotene and lycopene, which are intermediate products in

181 the synthesis of ABA, were identified via HPLC, and the results showed significant

182 enhancement of each in *HMGR*-OE leaves (Figure 5) Lutein content was also significantly

183 elevated in *HMGR*-OE leaves (Figure 5). In addition, the total level of GA3, a downstream

184 product of MEP, increased to 0.2–0.35 ng/g in *PtHMGR*-OE leaves, compared to 0.08–0.1 ng/g

185 in NT leaves (Figure 6A). Among MVA-related factors, the results of HPLC-MS/MS showed

186 that the TZR and IPA contents were higher in *PtHMGR*-OEs than in the NT (Figure 6B, and C).

187 The DCS content of *PtHMGR*-OEs increased to 0.31–0.41 ng/g relative to 0.74–1.12 ng/g in

188 the NT (Figure 6D), representing an average 3-fold increase in *PtHMGR*-OEs. However, the

189 CS content showed the opposite result in the comparison between *PtHMGR*-OEs and NT

190 (Figure 6E). Together, these results illustrate that the *HMGR* gene not only causes changes in

191 MVA-related genes and their products but also affects MEP-related genes and products.

Meanwhile, zeatin, IP, GA1, GA4, and GA7 were not detected through HPLC-MS/MS

(Supplemental Figure 14).

2.5. Enhanced TZR, IPA, DCS, carotenoid, ABA, and GA3 levels in *PtDXR*-OE poplars

The levels of β-carotene, lycopene, and ABA were much higher in *PtDXR*-OE poplars

than in NT plants, and lutein, which is considered a MEP-derived hormone, showed a similar

result (Figure 7). When the molecular mechanism underlying the improvement of ABA in

PtDXR-OEs was investigated, the expression levels of *NCED1*, *NCED3*, *NCED5*, *NCED6*,

ZEP1, *ZEP2*, and *ZEP3* were significantly higher in *PtDXR*-OEs than in NT plants (Figure 7).

In addition, a significant increase in GA3 in *PtDXR*-OEs was observed through HPLC-MS/MS

analyses (Figure 8A). In addition to upregulating MEP-derived products, overexpression of

PtDXR also affected the contents of MVA-derived products. The TZR content of *PtDXR*-OEs

increased to 0.24–0.37 ng/g compared to 0.0274–0.033 ng/g in NT plants (Figure 8B),

representing an average 10-fold increase in *PtDXR*-OEs. The content of IPA in *PtDXR*-OEs

increased to 0.84–1.12 ng/g, compared to 0.32–0.41 ng/g in NT poplars (Figure 8C),

representing an average 3-fold increase in *PtDXR*-OEs. The DCS content of *PtDXR*-OEs was

also significantly higher. Its level increased to 3.06–3.62 ng/g, with only 1.47–1.52 ng/g in NT

plants (Figure 8D), representing an average 3-fold increase in *PtDXR*-OEs (Figure 8D). By

contrast, the content of CS in *PtDXR*-OEs significantly decreased compared to that in NT

poplars (Figure 8C), indicating significant downregulation in *PtDXR*-OEs (Figure 8E). We

were unable to determine the contents of zeatin, IP, GA1, GA4, and GA7 through

HPLC-MS/MS (Supplemental Figure 15).

3. Discussion

3.1. Characterization and evolutionary history of HMGR

Previous studies have shown that *NtHMGR2* is a stress-responsive gene, whereas

NtHMGR1 is a housekeeping gene (Hemmerlin *et al*., 2004; Merret *et al*., 2007). *LcHMGR1* is

most highly expressed during the early stages of fruit development and is involved in the

regulation of fruit size. The level of *LcHMGR1* expression is higher and lasts longer in larger

fruits, while *LcHMGR2* is expressed most highly during the late stages of fruit development

and is related to the biosynthesis of isoprenoid substances required for cell elongation during

that time (Rui *et al*., 2012). The expression of *CaHMGR1* is temporary and tissue-specific,

whereas that of *CaHMGR2* is constitutive. *CaHMGR1* is only expressed in fruit tissues (pulp,

endosperm, and endocarp), flower buds, and leaves during the initial stages of development,

while *CaHMGR2* is expressed in all tissues (flower buds, leaves, branches, and roots) and fruit tissues at various stages of development (Tiski *et al*., 2012). In this study, we found that there is a copy number of at least three for *PtHMGR* genes in the poplar genome and that the *PtHMGR* gene (Potri.004G208500.1) has high homology with other known *HMGR1* genes based on multiple alignment analyses. In addition, the whole genome and HMGR protein sequences of various species were used to construct evolutionary trees, and the resulting evolutionary relationships among species were similar. Because *HMGR* in plants is a relatively conserved gene in the MVA pathway, it should be considered an important factor in studies of biological genetic differentiation and molecular evolution, and can be used as a reference for determining relationships among species.

3.2. HMGR regulates isoprenoid biosynthesis genes in *PtHMGR*-OE plants

Liao *et al*. (2018) showed that overexpression of *BjHMGS1* affects the expression levels of MEP- and MVA-related genes and slightly increases the transcript levels of *DXS* and *DXR* in transgenic plants. However, the expression levels of *DXS*, *DXR*, *HDS*, and *HDR* are significantly improved in *PtHMGR*-OE poplars, while those of *MCT* and *CMK* are downregulated. In addition, overexpression of *BjHMGS1* in tomato causes the expression levels of *GPS* and *GPPS* to significantly increase (Liao *et al*., 2018). In the present study, we also investigated the expression levels of *FPS*, *GPS*, and *GPPS*, and overexpression of *PtHMGR* increased their expression levels, which may have enhanced the interaction between IPP and dimethylallyl diphosphate (DMAPP), thereby increasing the biosynthesis of plastidial C15 and C20 isoprenoid precursors.

Xu *et al*[46] showed that overexpression of the *HMGR* gene in *Ganoderma lucidum* causes no significant changes in farnesyl diphosphate synthase (*FPS*), squalene synthase (*SQS*), or lanosterol synthase (LS) mRNA expression, despite increasing the contents of squalene and lanosterol. Overexpression of *BjHMGS1* in tomato significantly increases transcript levels of *FPS*, *SQS*, squalene epoxidase (*SQE*), and cycloartenol synthase (*CAS*) (Liao *et al*., 2018). Notably, the transcript levels of *AACT*, *MVK*, and *MVD* are significantly upregulated in *PtHMGR*-OE poplars although the expression level of *HMGS* is reduced. The increased expression has been observed for most MVA-related genes that contribute to the biosynthesis of sesquiterpenes and other C15 and universal C20 isoprenoid precursors.

3.3. Overexpression of *PtDXR* affects MEP- and MVA-related genes in *PtDXR*-OE plants

Overexpression of *TwDXR* in *Tripterygium wilfordii* increases the expression levels of

TwHMGS, *TwHMGR*, *TwFPS*, and *TwGPPS*, but decreases that of *TwDXS* (Zhang *et al*., 2018).

Overexpression of *NtDXR1* in tobacco increases the transcript levels of eight MEP-related

genes, indicating that overexpression of *NtDXR1* could lead to an upregulated expression of

downstream genes in the MEP pathway (Zhang *et al*., 2015). Changes in the *DXR* transcript

level of transgenic *A. thaliana* do not affect *DXS* gene expression or enzyme accumulation,

although overexpression of *DXR* boosts MEP-derived isoprenoids such as carotenoids,

chlorophylls, and taxadiene in *A. thaliana* (Carretero-Paulet *et al*., 2015). Overexpression of

the potato *DXS* gene in *A. thaliana* can cause upregulation of the downstream *GGPPS* gene and

PSY (phytoene synthase) gene (Henriquez *et al*., 2016). In addition, the transformation of the

DXS gene from *A. thaliana* into *Daucus carota* significantly enhances the expression level of

the *PSY* gene (Simpson *et al*., 2016).

Our results show both similarities and differences compared to previous studies. For

example, we found that the transcript levels of MEP-related genes were higher in *PtDXR*-OE

poplars than in NT poplars. However, overexpression of *PtDXR* in poplar led to significantly

downregulated expression levels of MVA-related genes. Together, these findings illustrate that

carbon sources may be preferentially synthesized into monoterpenes, diterpenes, and

tetraterpenoids in poplar cells. Secondary metabolic pathways in plants are tightly regulated by

the transcript levels of MEP- and MVA-related genes. The dominant isoprenoid compounds

synthesized in different plants differ greatly. The regulation of the expression of multiple

enzymes largely controls whether specific secondary metabolites are synthesized and to what

extent. The diversity of biosynthetic pathways, the complexity of metabolic networks, and the

unknown regulation of gene expression lead to very different regulation patterns of MEP- and

MVA-related gene expression among species. One possible conclusion is that MEP- and

MVA-related genes often do not work alone, but instead are co-expressed with upstream and

downstream genes in the MEP and MVA pathways to carry out a specific function.

3.4. Overexpression of *HMGR* promotes the formation of ABA, GAs, and carotenoids in

plastids and accumulation of TZR, IPA, DCS in the cytoplasm

Plant HMGR, as the rate-limiting enzyme in the MVA pathway, plays a very important

role in controlling the flow of carbon in that metabolic pathway. The upregulation of the

HMGR gene significantly increases isoprenoid levels in plants. Studies on homologous and

heterologous overexpression of *HMGR* have reported significantly elevated levels of

isoprenoid in transgenic plants. In one study, *Hevea brasiliensis HMGR1* was transferred into

tobacco, and it increased the sterol content of transgenic tobacco as well as the accumulation of intermediate metabolites (Schaller *et al*., 1995). In another study, *A. thaliana HMGR* overexpression in *Lavandula latifolia* increased the levels of sterols in the MVA pathway and of MEP-derived monoterpenes and sesquiterpenes (Munoz-Bertomeu *et al*., 2007). In still another, overexpression of the *SmHMGR* gene in hairy roots increased MEP-derived diterpene tanshinone (Kai *et al*., 2011). In our study, ABA synthesis-related genes (*NCED1*, *NCED3*, *NCED5*, *NCED6*, *ZEP1*, *ZEP2*, and *ZEP3*) and the contents of ABA, GA3, and carotenoids were upregulated in *PtHMGR*-OE poplar seedlings, suggesting that overexpression of *HMGR* may indirectly affect the biosynthesis of MEP-related isoprenoids including ABA, GAs, and carotenoids. The accumulation of MVA-derived isoprenoids including TZR, IPA, and DCS was significantly elevated in *PtHMGR*-OEs, indicating that *PtHMGR* overexpression has a direct influence on the biosynthesis of MVA-related isoprenoids. Therefore, the *HMGR* gene not only directly affects the contents of MVA-derived isoprenoids but also indirectly affects the content of MEP-derived isoprenoids by changing the expression levels of MEP-related genes.

3.5. Upregulation of MEP-derived products (ABA, GA3, and carotenoid) and MVA-derived products (TZR, IPA, DCS) in *PtDXR*-OE seedlings

The production of MEP catalyzed DXR, which is the most important rate-limiting enzyme in the MEP pathway and an important regulatory step in the cytoplasmic metabolism of isoprenoid compounds (Takahashi *et al*., 1998). In Mahmoud *et al*. (2001), overexpression of *DXR* in *Mentha piperita* promoted the synthesis of monoterpenes such as mint oil in the leaves and increased the production of essential mint oil by 50%. In Hasunuma *et al*. (2008), overexpression of *Synechocystis sp*. strain PCC6803 *DXR* in tobacco resulted in increased levels of β-carotene, chlorophyll, antheraxanthin, and lutein. In another study, *A. thaliana DXR* mutant plants exhibited a lack of GAs, ABA, and photosynthetic pigments, and the phenotype of the mutant *A. thaliana* plants showed pale sepals and yellow inflorescences (Xing *et al*., 2010). In our study, ABA synthesis-related genes were dramatically increased, as was the accumulation of ABA, GA3, and carotenoids, in *PtDXR*-OE poplar seedlings, indicating an effect from *DXR* overexpression. Combined with the result described above of increased *DXS*, *HDS*, *HDR*, *MCT*, *CMK*, *FPS*, *GPS*, and *GPPS* expression levels, we can conclude that overexpression of *DXR* can directly affect the expression levels of MEP-related genes and that changes in MEP-related gene expression levels contribute to the field of ABA, GA3, and carotenoids. Although the expressions of *AACT*, *HMGS*, *HMGR*, *MVK*, and *MVD*, which are

related to the MVA pathway, were downregulated, the contents of the MVA-derived components TZR, IPA, and DCS were upregulated in *PtDXR*-OE seedlings, aside from their reduced CS content.

3.6. Cross-talk in the MVA and MEP pathways

Although the starting materials differ between the MVA and MEP pathways, IPP and DMAPP are common precursors. Blocking only the MVA or the MEP pathway does not completely block the biosynthesis of terpenes in the cytoplasm or plastids, indicating that the common products of the MVA and MEP pathways can be transported freely to different subcellular regions (Aharoni *et al.* 2003; Aharoni *et al.* 2004; Gutensohn *et al.* 2013). For example, results of green fluorescent protein fusion and confocal microscopy have shown that HDR is localized in the chloroplast, and the transfer of IPP from the chloroplast to cytoplasm can be observed through ^{13}C labeling, indicating that plentiful IPP is available for use in the MVA pathway to produce terpenoids (Ma *et al.* 2017). In addition, segregation between the MVA and MEP pathways is limited, and some metabolites can be exchanged on the plasmid membrane (Laule *et al.*, 2003). Kim *et al.* (2016) used clustered regularly interspaced short palindromic repeats (CRISPR) technology to reconstruct the lycopene synthesis pathway and control the flow of carbon in the MEP and MVA pathways. The results showed that MVA-related genes were reduced by an average of 81.6% and the lycopene yield was significantly boosted. According to analyses of gene expression levels and metabolic yield in *PtHMGR*-OEs and *PtDXR*-OEs, we found cross-talk not only between IPP and DMAPP in the MVA and MEP pathways but also between pairs of genes and between products of the MVA and MEP pathways in poplar cells. On one hand, overexpression of *PtDXR* affected the transcript levels of MEP-related genes and the contents of MEP-derived isoprenoids including ABA, GAs, and carotenoids. However, the decreased accumulation of MVA-related gene products reduces the yields of MVA-derived isoprenoids including CS, while increasing TZR, IPA, and DCS contents. We also found that differential expression of MVA-related genes in *PtDXR*-OE or post-transcriptional/post-translational regulation of the products of MVA-derived isoprenoids may have occurred. Furthermore, although the expression levels of MVA-related genes in *PtDXR*-OEs were lower than those in NT poplars, the transcript levels of *IDI* were upregulated in *PtDXR*-OEs, indicating that IPP and DMAPP produced by the MEP pathway could enter the cytoplasm to compensate for the lack of IPP and DMAPP there. On the other hand, *PtHMGR*-OE plants exhibited higher transcript levels of *AACT*, *MVK*, and *MVD* as

well as higher expression levels of *DXS*, *DXR*, *HDS*, and *HDR* compared to NT, which indicates that PtHMGR affects both MEP- and MVA-related gene expression levels. We successfully demonstrated that manipulation of *HMGR* in the MVA pathway of poplars results in dramatically enhanced yields of ABA, GAs, and carotenoids. This result illustrates that cytosolic *HMGR* overexpression could lead to a boost in plastidial GPP- and GGPP-derived products, including ABA, GAs, and carotenoids. Therefore, this study provides evidence that cross-talk between the MVA and MEP pathways increased the expression levels of *GPS* and *GPPS* in *PtHMGR*-OEs, and elevated the contents of ABA, GA3, and carotenoids. Moreover, changes in MEP- and MVA-related gene expression levels affect MVA- and MEP-derived isoprenoids. Together, these results illustrate the phenomenon of cross-talk between the MVA and MEP pathways. The mechanism underlying this exchange between the MVA and MEP pathways requires further research.

In conclusion, Isoprenoid compounds are diverse, with complex structures and different properties, and play an important role in the survival and growth of plants. The basic framework of the isoprenoid metabolic pathway has been explored, and many advances have been made in clarifying the molecular regulation of metabolic pathways and metabolic engineering. However, previous studies have been limited to *Arabidopsis*, tomato, and rice, and research on isoprenoid metabolism in perennial woody plants has just begun. In this study, the overexpression of *PtHMGR* in Nanlin 895 (*Populus× euramericana* cv. 'Nanlin 895') caused the accumulation of MVA-derived isoprenoids as well as MEP-derived substances including ABA, GAs, carotenoids, and GRs, which are essential plant hormones regulating growth and stress responses. In *PtHMGR*-OE poplars, most MEP- and MVA-related genes associated with the biosynthesis of isoprenoid precursors were upregulated. In *PtDXR*-OE poplars, elevated contents of ABA, GAs, carotenoids, and GRs were attributed to increased expression of MEP-related genes as well as plastidial *GPP* and *GGPP*. Moreover, in both *PtDXR*-OE poplars, changes in MVA-related genes caused changes in MVA-derived isoprenoids, including the yields of CKs and BRs. Together, these results show that the manipulation of *PtDXR and PtHMGR* is a novel strategy to influence poplar isoprenoids and provide a theoretical basis for the cultivation of high-quality poplar.

4. Materials and Methods

4.1. Plant materials and growth conditions

Non-transgenic (NT) *P. trichocarpa* and Nanlin 895 (*Populus × euramericana* cv.

'Nanlin 895') plants were cultured in half-strength Murashige and Skoog (1/2 MS) medium (PH 5.8) under conditions of 24°C and 74% humidity (Movahedi *et al.*, 2015a and 2015b). Subsequently, NT and transgenic poplars were cultured in 1/2 MS under long-day conditions (16 h light/8 h dark) at 24°C for 1 month (Movahedi *et al.*, 2018).

4.2. *PtHMGR* gene isolation and vector construction

Total RNA was extracted from *P. trichocarpa* leaves and reverse transcriptase (TaKaRa, Japan) was used to amplify the cDNA. Forward and reverse primers (Supplemental Table: *PtHMGR*-F and *PtHMGR*-R) were designed and polymerase chain reaction (PCR) was performed to synthesize the open region frame (ORF) of the *PtHMGR* gene. Subsequently, the product of the *PtHMGR* gene was ligated into the PEASY-T3 vector (TransGen Biotech, China) based on blue-white spot screening and the *PtHMGR* gene was inserted into the vector pGWB$_9$ (Song *et al.*, 2016) using Gateway technology (Invitrogen, USA).

To identify the copy number of *PtHMGR* in *P. trichocarpa*, total genomic DNA was extracted from leaves according to the cetyltrimethylammonium bromide (CTAB) method and Southern blotting was performed. The poplar DNA was digested with *EcoR*I at 37°C for 4 h and the digested poplar DNA was separated on a 0.8% agarose gel at 15 V. Then the separated poplar DNA was transferred to a Hybond N$^+$ nylon membrane and blotting was performed according to the manufacturer's instructions (Roche, Basel, Switzerland). The details of probe 1 and probe 2 used for Southern blotting are provided in Supplemental Table 1.

4.3. Creation of phylogenetic tree

To cluster families based on protein-coding genes, the protein data for 10 plant species were downloaded Phytozome (https://phytozome.jgi.doe.gov/pz/portal.html): *P. trichocarpa* (v3.1), *Oryza sativa* (v7), *Manihot esculenta* (v6.1), *Theobroma cacao* (v1.1), *Gossypium hirsutum* (v1.1), *Prunus persica* (v2.1), *Malus domestica* (v1.0), *Arabidopsis thaliana TAIR10*, and *Zea mays* PH207 (v1.1). Only the gene model for each gene locus that encodes the longest protein sequence was retained to remove redundant sequences. Then we employed DIAMOND software to identify potentially orthologous gene families among the filtered protein sequences of these species using a maximum e-value of 1e-5 (Buchfink *et al.*, 2015). We used the OrthoFinder (v2.3.3) pipeline to identify single-copy gene families based on the aligned results with the default parameters (Emms *et al.*, 2015).

We applied MAFFT (v7.158b) software (Katoh and Standley, 2013) for multiple sequence alignment of protein-coding sequences for each single-copy gene using the accurate option

(LINS-i). Meanwhile, trimAl (v1.2) software (Capella-Gutiérrez *et al.*, 2009) was employed to

remove poorly aligned positions and divergent regions with the default parameters, and the

alignments of all of the single-copy genes were concatenated to form a supermatrix. We used

the program RAxML (v8.2.12) (Stamatakis, 2014) with the JTT+G+F model and 1000

bootstrap replicates to construct the supermatrix phylogenetic tree. Finally, the phylogenetic

tree was visualized using FigTree (v1.4.3) (Drummond *et al.*, 2009). The amino acid sequences

of HMGR from *P. trichocarpa*, *M. esculenta*, *S. purpurea*, *T. cacao*, *G. hirsutum*, *D. persica*,

M. domestica, *A. thaliana*, and *Z. mays* were obtained from the National Center for

Biotechnology Information database (https://www.ncbi.nlm.nih.gov/) and Phytozome and

MEGA5.0 software was used to construct a phylogenetic tree using 1000 bootstrap replicates.

4.4. Generation and characterization of transgenic poplars

Agrobacterium EHA105 containing a recombinant plasmid was used for infection of

Nanlin 895 (*Populus × euramericana* cv. 'Nanlin 895') leaves and petioles (Movahedi *et al.*,

2014; Movahedi *et al.*, 2020). Poplar buds were screened on differentiation MS medium

amended with 30 μg/mL Kan and the resistant buds were planted in bud elongation MS

medium containing 20 μg/mL Kan and transplanted into 1/2 MS medium containing 10 μg/mL

Kan to generate resistant poplar trees. Subsequently, the DNA and RNA of putative transgenic

poplars were isolated and analyzed via PCR and quantitative reverse transcription PCR

(qRT-PCR).

4.5. Analyses via qRT-PCR

PtDXR-OE poplars have been obtained previously (Xu *et al.*, 2019), and in this study, we

extracted RNA from 12-month-old *PtDXR*-OE and *PtHMGR*-OE poplars. qRT-PCR was

performed to identify the expression levels of MVA- and MEP-related genes in NT, *PtDXR*-OE,

and *PtHMGR*-OE poplars. qRT-PCR was performed with a StepOne Plus Real-time PCR

System (Applied Biosystems, USA) and SYBR Green Master Mix (Roche, Germany). Poplar

Actin (*PtActin*) (XM-006370951.1) has previously been tested as an internal control for

estimating the amount of RNA in samples (Zhang *et al.*, 2013). The following conditions were

used for qRT-PCR reactions: pre-denaturation at 95°C for 10 min, followed by 40 cycles of

denaturation at 95°C for 15 s and a chain extension at 60°C for 1 min. Three independent

experiments were conducted using gene-specific primers (Supplemental Table 1).

4.6. Analyses of ABA, GAs, CKs, GRs, BRs, and carotenoids via high-performance liquid

chromatography-tandem mass spectrometry (HPLC-MS/MS)

ABA, GAs, and CKs were extracted from NT, *PtDXR*-OE, and *PtHMGR*-OE leaves, and

then HPLC-MS/MS (Qtrap6500, Agilent, USA) was used to quantitatively determine levels of

ABA, GAs, zeatin, trans-zeatin-riboside (TZR), and isopentenyl adenosine (IPA).

HPLC-MS/MS (Aglient1290, AB; SCIEX-6500Qtrap, Agilent; USA) was also used to

determine the contents of 5-deoxystrgol (5-DS), castasterone (CS), and 6-deoxyocastasterone

(DCS). In addition, the carotenoid component of poplar leaves was isolated and HPLC

(Symmetry Shield RP18, Waters, USA) was used to determine the contents of carotenoids

including β-carotene, lycopene, and lutein

Author contributions

Q.Z. conceived the study. H.W. and A.M. performed the experiments and carried out the

analysis. H.W., A.M., C.X., W.B.S., P.W., D.W.L., and Q.Z designed the experiments and

wrote the manuscript. All authors read and approved the final manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by the National Key Program on Transgenic Research

(2018ZX08020002), the National Science Foundation of China (No. 31570650), and the

Priority Academic Program Development of Jiangsu Higher Education Institutions.

Reference

Aharoni, A., Giri, A. P., Deuerlein, S., Griepink, F., de Kogel, W. J., Verstappen, F. W., ... &

Bouwmeester, H. J., 2003. Terpenoid metabolism in wild-type and transgenic Arabidopsis

plants. Plant Cell, 15, 2866-2884.

Aharoni, A., Giri, A. P., Verstappen, F. W., Bertea, C. M., Sevenier, R., Sun, Z., ... &

Bouwmeester, H. J., 2004. Gain and loss of fruit flavor compounds produced by wild and

cultivated strawberry species. Plant Cell, 16, 3110-3131.

Aharoni, A., Jongsma, M. A., & Bouwmeester, H. J., 2005. Volatile science? Metabolic

engineering of terpenoids in plants. Trends Plant Sci. 10, 594-602.

Bouvier, F., Rahier, A., & Camara, B., 2005. Biogenesis, molecular regulation, and function of

plant isoprenoids. Prog. Lipid Res. 44, 357-429.

Buchfink, B., Xie, C., & Huson, D. H., 2015. Fast and sensitive protein alignment using

DIAMOND. Nat. Methods. 12, 59.

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T., 2009. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 25, 1972-1973.

Carretero-Paulet, L., Cairo, A., Botella-Pavía, P., Besumbes, O., Campos, N., Boronat, A., &
Rodríguez-Concepción, M., 2006. Enhanced flux through the methylerythritol 4-phosphate
pathway in Arabidopsis plants overexpressing deoxyxylulose 5-phosphate reductoisomerase.
Plant Mol. Biol. 62, 683-695.

Cowan, A. K., Moore-Gordon, C. S., Bertling, I., & Wolstenholme, B. N., 1997. Metabolic
control of avocado fruit growth (isoprenoid growth regulators and the reaction catalyzed by
3-hydroxy-3-methylglutaryl coenzyme A reductase). Plant Physiol.114, 511-518.

Dai, Z., Cui, G., Zhou, S. F., Zhang, X., & Huang, L., 2011. Cloning and characterization of a
novel 3-hydroxy-3-methylglutaryl coenzyme A reductase gene from *Salvia miltiorrhiza*,
involved in diterpenoid tanshinone accumulation. J. Plant Physiol.168, 148-157.

Drummond, A. J., & Rambaut, A., 2007. BEAST: Bayesian evolutionary analysis by sampling
trees. BMC Evol. Biol. 7, 214.

Dueber, J. E., Wu, G. C., Malmirchegini, G. R., Moon, T. S., Petzold, C. J., Ullal, A. V., ... &
Keasling, J. D., 2009. Synthetic protein scaffolds provide modular control over metabolic
flux. Nat. Biotechnol. 27, 753.

Emms, D. M., & Kelly, S., 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 16, 157.

Enfissi, E. M., Fraser, P. D., Lois, L. M., Boronat, A., Schuch, W., & Bramley, P. M., 2005.
Metabolic engineering of the mevalonate and non-mevalonate isopentenyl
diphosphate-forming pathways for the production of health-promoting isoprenoids in
tomato. Plant Biotechnol. J. 3, 17-27.

Esteban, R., Barrutia, O., Artetxe, U., Fernández-Marín, B., Hernández, A., & García-Plazaola,
J. I., 2015. Internal and external factors affecting photosynthetic pigment composition in plants:
a meta-analytical approach. New Phytol. 206, 268-280.

Ferrero, S., Grados-Torrez, R. E., Leivar, P., Antolín-Llovera, M., López-Iglesias, C.,
Cortadellas, N., ... & Campos, N., 2015. Proliferation and morphogenesis of the endoplasmic
reticulum driven by the membrane domain of 3-hydroxy-3-methylglutaryl coenzyme A
reductase in plant cells. Plant Physiol. 168, 899-914.

Gutensohn, M., Orlova, I., Nguyen, T. T., Davidovich☐Rikanati, R., Ferruzzi, M. G., Sitrit,
Y., ... & Dudareva, N., 2013. Cytosolic monoterpene biosynthesis is supported by
plastid-generated geranyl diphosphate substrate in transgenic tomato fruits. Plant J. 75,

351-363.

Hain, Rüdiger, Reif, H. J., Krause, E., Langebartels, R., Kindl, H., & Vornam, B., et al., 1993. Disease resistance results from foreign phytoalexin expression in a novel plant. Nature, 361, 153-156.

Hasunuma, T., Takeno, S., Hayashi, S., Sendai, M., Bamba, T., Yoshimura, S., ... & Miyake, C., 2008. Overexpression of 1-deoxy-D-xylulose-5-phosphate reductoisomerase gene in chloroplast contributes to increment of isoprenoid production. J. Biosci. Bioeng. 105, 518-526.

Hemmerlin, A., Gerber, E., Feldtrauer, J. F., Wentzinger, L., Hartmann, M. A., Tritsch, D., ... & Bach, T. J., 2004. A review of tobacco BY-2 cells as an excellent system to study the synthesis and function of sterols and other isoprenoids. Lipids. 39, 723-735.

Hemmerlin, A., Harwood, J. L., & Bach, T. J., 2012. A raison d'être for two distinct pathways in the early steps of plant isoprenoid biosynthesis? Prog. Lipid Res. 51, 95-148.

Henriquez, M. A., Soliman, A., Li, G., Hannoufa, A., Ayele, B. T., & Daayf, F., 2016. Molecular cloning, functional characterization and expression of potato (*Solanum tuberosum*) 1-deoxy-d-xylulose 5-phosphate synthase 1 (StDXS1) in response to *Phytophthora infestans*. Plant Sci. 243, 71-83.

Henry, L. K., Gutensohn, M., Thomas, S. T., Noel, J. P., & Dudareva, N., 2015. Orthologs of the archaeal isopentenyl phosphate kinase regulate terpenoid production in plants. Proc. Natl. Acad. Sci. U. S. A. 112, 10050-10055.

Huang, Z., & Szoka Jr, F. C., 2008. Sterol-modified phospholipids: cholesterol and phospholipid chimeras with improved biomembrane properties. J. Am. Chem. Soc. 130, 15702-15712.

Huchelmann, A., Gastaldo, C., Veinante, M., Zeng, Y., Heintz, D., Tritsch, D., ... & Hemmerlin, A., 2014. S-Carvone suppresses cellulase-induced capsidiol production in *Nicotiana tabacum* by interfering with protein isoprenylation. Plant Physiol. 164, 935-950.

Kai, G., Xu, H., Zhou, C., Liao, P., Xiao, J., Luo, X., ... & Zhang, L., 2011. Metabolic engineering tanshinone biosynthetic pathway in *Salvia miltiorrhiza* hairy root cultures. Metab. Eng. 13, 319-327.

Kazan, K., 2015. Diverse roles of jasmonates and ethylene in abiotic stress tolerance. Trends Plant Sci. 20, 219-229.

Kim, S. K., Han, G. H., Seong, W., Kim, H., Kim, S. W., Lee, D. H., & Lee, S. G., 2016. CRISPR interference-guided balancing of a biosynthetic mevalonate pathway increases

terpenoid production. Metab. Eng. 38, 228-240.

Kim, M. S., Haney, M. J., Zhao, Y., Mahajan, V., Deygen, I., Klyachko, N. L., ... & Hingtgen, S.

D., 2016. Development of exosome-encapsulated paclitaxel to overcome MDR in cancer

cells. Nanomedicine. 12, 655-664.

Kim, Y. J., Lee, O. R., Ji, Y. O., Jang, M. G., & Yang, D. C., 2014. Functional analysis of

HMGR encoding genes in triterpene saponin-producing *Panax ginseng* Meyer. Plant Physiol.

165, 373-387.

Kirby, J., & Keasling, J. D., 2009. Biosynthesis of plant isoprenoids: perspectives for microbial

engineering. Annu. Rev. Plant Biol. 60, 335-355.

Kong, L. Y., & Tan, R. X., 2015. Artemisinin, a miracle of traditional Chinese medicine. Nat.

Prod. Rep. 32, 1617-1621.

Laule, O., Fürholz, A., Chang, H. S., Zhu, T., Wang, X., Heifetz, P. B., ... & Lange, M., 2003.

Crosstalk between cytosolic and plastidial pathways of isoprenoid biosynthesis in *Arabidopsis*

thaliana. Proc. Natl. Acad. Sci. U. S. A. 100, 6866-6871.

Li, W., Liu, W., Wei, H., He, Q., Chen, J., Zhang, B., & Zhu, S., 2014. Species-specific

expansion and molecular evolution of the 3-hydroxy-3-methylglutaryl coenzyme A reductase

(HMGR) gene family in plants. PloS One. 9, e94172.

Liao, P., Chen, X., Wang, M., Bach, T. J., & Chye, M. L., 2018. Improved fruit α-tocopherol,

carotenoid, squalene and phytosterol contents through manipulation of Brassica juncea

3-HYDROXY-3-METHYLGLUTARYL-COA SYNTHASE 1 in transgenic tomato. Plant

Biotechnol. J. 16, 784-796.

Liao, P., Hemmerlin, A., Bach, T. J., & Chye, M. L., 2016. The potential of the mevalonate

pathway for enhanced isoprenoid production. Biotechnol. Adv. 34, 697-713.

Liao, Z. H., Chen, M., Gong, Y. F., Miao, Z. Q., Sun, X. F., & Tang, K. X., 2006. Isoprenoid

biosynthesis in plants: pathways, genes, regulation and metabolic engineering. J Biol Sci. 6,

209-219.

Lu, X. M., Hu, X. J., Zhao, Y. Z., Song, W. B., Zhang, M., Chen, Z. L., ... & Lai, J. S., 2012.

Map-based cloning of zb7 encoding an IPP and DMAPP synthase in the MEP pathway of

maize. Mol. Plant. 5, 1100-1112.

Ma, D., Li, G., Zhu, Y., & Xie, D. Y., 2017. Overexpression and suppression of Artemisia annua

4-hydroxy-3-methylbut-2-enyl diphosphate reductase 1 gene (*AaHDR1*) differentially regulate

artemisinin and terpenoid biosynthesis. Front Plant Sci. 8, 77.

Ma, Y., Yuan, L., Wu, B., Li, X. E., Chen, S., & Lu, S., 2012. Genome-wide identification and

characterization of novel genes involved in terpenoid biosynthesis in *Salvia miltiorrhiza*. J.

Exp. Bot. 63, 2809-2823.

Mahmoud, S. S., & Croteau, R. B., 2001. Metabolic engineering of essential oil yield and

composition in mint by altering expression of deoxyxylulose phosphate reductoisomerase and

menthofuran synthase. Proc. Natl. Acad. Sci. U. S. A. 98, 8915-8920.

Merret, R., Cirioni, J. R., Bach, T. J., & Hemmerlin, A., 2007. A serine involved in

actin-dependent subcellular localization of a stress-induced tobacco BY-2

hydroxymethylglutaryl-CoA reductase isoform. FEBS Lett. 581, 5295-5299.

Movahedi, A., Wei, H., Chen, Z. H., ... & Zhuge, Q., 2020. CRISPR-Cas9 integrates

exogeneous sorting new recombinant DNA in the model tree *Populus trichocarpa*. doi:

10.1101/2020.07.04.188219

Movahedi, A., Zhang, J., Sun, W., Mohammadi, K., Yaghuti, A. A., Wei, H., ... & Zhuge, Q.,

2018. Functional analyses of *PtRDM1* gene overexpression in poplars and evaluation of its

effect on DNA methylation and response to salt stress. Plant Physiol Bioch. 64-73.

Movahedi, A., Zhang, J., Gao, P., Yang, Y., Wang, L., & Yin, T., et al., 2015a. Expression of the

chickpea *CarNAC3*, gene enhances salinity and drought tolerance in transgenic poplars. Plant

Cell, Tissue Organ Cult. 120, 141-154.

Movahedi, A., Zhang, J., Yin, T., & Zhuge, Q., 2015b. Functional Analysis of Two

Orthologous *NAC* Genes, *CarNAC3*, and *CarNAC6* from *Cicer arietinum*, involved in abiotic

stresses in Poplar. Plant Mol Biol Rep. 33(5), 1539-1551.

Movahedi, A., Zhang, J., Amirian, R., & Zhuge, Q., 2014. An efficient

Agrobacterium-mediated transformation system for poplar. Int J Mol Sci. 15(6), 10780-10793.

Muñoz-Bertomeu, J., Sales, E., Ros, R., Arrillaga, I., & Segura, J., 2007. Up-regulation of an

N-terminal truncated 3-hydroxy-3-methylglutaryl CoA reductase enhances production of

essential oils and sterols in transgenic *Lavandula latifolia*. Plant Biotechnol. J. 5, 746-758.

Rahman, L., Kouno, H., Hashiguchi, Y., Yamamoto, H., Narbad, A., Parr, A., ... & Kitamura, Y.,

2009. HCHL expression in hairy roots of *Beta vulgaris* yields a high accumulation of

p-hydroxybenzoic acid (pHBA) glucose ester, and linkage of pHBA into cell walls. Bioresour.

Technol. 100, 4836-4842.

Roberts, S. C. Production and enginee*ri*ng of terpenoids in plant cell culture., 2007. Nat. Chem.

Biol. 3, 387.

Ren, D., Liu, Y., Yang, K. Y., Han, L., Mao, G., Glazebrook, J., & Zhang, S. A

fungal-responsive MAPK cascade regulates phytoalexin biosynthesis in Arabidopsis., 2008.

Proc. Natl. Acad. Sci. U. S. A. 105, 5638-5643.

Rui, X., Caiqin, L., Wangjin, L., Juan, D., Zehuai, W., & Jianguo, L., 2012.

3-Hydroxy-3-methylglutaryl coenzyme A reductase 1 (HMG1) is highly associated with the

cell division during the early stage of fruit development which determines the final fruit size in

Litchi chinensis. Gene. 498, 28-35.

Schaller, H., Grausem, B., Benveniste, P., Chye, M. L., Tan, C. T., Song, Y. H., & Chua, N. H.,

1995. Expression of the *Hevea brasiliensis* (HBK) Mull. Arg.

3-hydroxy-3-methylglutaryl-coenzyme A reductase 1 in tobacco results in sterol

overproduction. Plant Physiol. 109, 761-770.

Schie, C. C. V. Haring, M. A., Schuurink, R. C., 2006. Regulation of terpenoid and benzenoid

production in flowers. Curr. Opin. Plant Biol. 9, 203-208.

Simpson, K., Quiroz, L. F., Rodriguez-Concepción, M., & Stange, C. R., 2016. Differential

contribution of the first two enzymes of the MEP pathway to the supply of metabolic

precursors for carotenoid and chlorophyll biosynthesis in carrot (*Daucus carota*). *Front Plant*

Sci. 7, 1344.

Song, X., Yu, X., Hori, C., Demura, T., Ohtani, M., & Zhuge, Q., 2016. Heterologous

overexpression of poplar *SnRK2* genes enhanced salt stress tolerance in *Arabidopsis thaliana*.

Front Plant Sci. 7, 612.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 30, 1312-1313.

Suzuki, M., Kamide, Y., Nagata, N., Seki, H., Ohyama, K., & Kato, H., et al., 2005. Loss of

function of 3-hydroxy-3-methylglutaryl coenzyme A reductase 1 (HMGR1) in *Arabidopsis*

leads to dwarfing, early senescence and male sterility, and reduced sterol levels. Plant J. 20,

67-77.

Takahashi, S., Kuzuyama, T., Watanabe, H., & Seto, H., 1998. A 1-deoxy-D-xylulose

5-phosphate reductoisomerase catalyzing the formation of 2-C-methyl-D-erythritol

4-phosphate in an alternative nonmevalonate pathway for terpenoid biosynthesis. Proc. Natl.

Acad. Sci. U. S. A. 95, 9879-9884.

Tiski, I., Marraccini, P., Pot, D., Vieira, L. G. E., & Pereira, L. F. P., 2011. Characterization and

expression of two cDNA encoding 3-Hydroxy-3-methylglutaryl coenzyme A reductase

isoforms in coffee (*Coffea arabica L.*). OMICS: J. Integr. Biol. 15, 719-727.

Vaccaro, M., Malafronte, N., Alfieri, M., De Tommasi, N., & Leone, A., 2014. Enhanced biosynthesis of bioactive abietane diterpenes by overexpressing *AtDXS* or *AtDXR* genes in *Salvia sclarea* hairy roots. Plant Cell, Tissue Organ Cult. 119, 65-77.

Wang, H., Nagegowda, D. A., Rawat, R., Bouvier-Navé, P., Guo, D., Bach, T. J., & Chye, M. L., 2012. Overexpression of Brassica juncea wild-type and mutant HMG-CoA synthase 1 in Arabidopsis up-regulates genes in sterol biosynthesis and enhances sterol production and stress tolerance. Plant Biotechnol. J. 10, 31-42.

Weber, P., Bendich, A., & Machlin, L. J., 1997. Vitamin E and human health: rationale for determining recommended intake levels. Nutrition. 13, 450-460.

Wong, C. W., Czarny, B., Metselaar, J. M., Ho, C., Ng, S. R., Barathi, A. V., ... & Wong, T. T., 2018. Evaluation of subconjunctival liposomal steroids for the treatment of experimental uveitis. Sci. Rep. 8, 6604.

Xie, Z., Kapteyn, J., & Gang, D. R., 2008. A systems biology investigation of the MEP/terpenoid and shikimate/phenylpropanoid pathways points to multiple levels of metabolic control in sweet basil glandular trichomes. Plant J. 54, 349-361.

Xing, S., Miao, J., Li, S., Qin, G., Tang, S., Li, H., ... & Qu, L. J., 2010. Disruption of the 1-deoxy-D-xylulose-5-phosphate reductoisomerase (*DXR*) gene results in albino, dwarf and defects in trichome initiation and stomata closure in Arabidopsis. Cell Res. 20, 688.

Xu, C., Wei, H., Movahedi, A., Sun, W., Ma, X., Li, D., ... & Zhuge, Q., 2019. Evaluation, characterization, expression profiling, and functional analysis of *DXS* and *DXR* genes of *Populus trichocarpa*. Plant Physiol. Biochem. 142.

Xu, J. W., Xu, Y. N., & Zhong, J. J., 2012. Enhancement of ganoderic acid accumulation by overexpression of an N-terminally truncated 3-hydroxy-3-methylglutaryl coenzyme A reductase gene in the basidiomycete *Ganoderma lucidum*. Appl. Environ. Microbiol. 78, 7968-7976.

Yamaguchi, S., Kamiya, Y., & Nambara, E., 2018. Regulation of ABA and GA levels during seed development and germination in Arabidopsis. Annu. Plant Rev. 27, 224-247.

Zhang, K., Fan, W., Huang, Z., Chen, D., Yao, Z., Li, Y., ... & Qiu, D., 2019. Transcriptome analysis identifies novel responses and potential regulatory genes involved in 12-deoxyphorbol-13-phenylacetate biosynthesis of *Euphorbia resinifera*. Ind. Crops Prod. 135, 138-145.

Zhang, H., Niu, D., Wang, J., Zhang, S., Yang, Y., Jia, H., & Cui, H., 2015. Engineering a

platform for photosynthetic pigment, hormone and cembrane-related diterpenoid production in

Nicotiana tabacum. Plant Cell Physiol. 56, 2125-2138.

Zhang, J., Li, J., Liu, B., Zhang, L., Chen, J., & Lu, M., 2013. Genome-wide analysis of the

populus hsp90 gene family reveals differential expression patterns, localization, and heat stress

responses. BMC Genomics. 14.

Zhang, L., Ding, R., Chai, Y., Bonfill, M., Moyano, E., Oksman-Caldentey, K. M., ... & Kai, G.,

2004. Engineering tropane biosynthetic pathway in *Hyoscyamus niger* hairy root cultures. Proc.

Natl. Acad. Sci. U. S. A. 101, 6786-6791.

Zhang, Y., Zhao, Y., Wang, J., Hu, T., Tong, Y., Zhou, J., ... & Huang, L., 2018. Overexpression

and RNA interference of *TwDXR* regulate the accumulation of terpenoid active ingredients in

Tripterygium wilfordii. Biotechnol. Lett. 40, 419-425.

Figure legends

Figure 1. Southern blotting analyses of homologous *PtHMGR* copy number and identification

of probes specific to *PtHMGR* (Potri.004G208500.1). (A) Southern blotting was performed to

determine the copy number of homologous *PtHMGR* genes using probe 1. (B) Southern

blotting was performed using probe 2 to specifically recognize *PtHMGR*

(Potri.004G208500.1).

Figure 2. Phylogenetic analyses. (A) Construction of a phylogenetic tree based on the whole

genomes of various species. (B) Construction of a phylogenetic tree based on the HMGR

sequences of various species. Accession numbers of the HMGR obtained from GenBank are as

follows: *Arabidopsis thaliana* (NP_177775.2), *Gossypium hirsutum* (XP_016691783.1),

Malus domestica (XP_008348952.1), *Manihot esculenta* (XP_021608133.1), *Prunus persica*

(XM_020569919.1), *Oryza sativa* (XM_015768351.2), *Theobroma cacao* (XM_007043046.2),

Zea mays (PWZ28886.1).

Figure 3. qRT-PCR analyses of the transcript levels of MEP- and MVA-related genes in

*PtHMGR-OE*s. Transcript levels of MEP-related genes including *DXS*, *DXR*, *MCT*, *CMK*,

HDS, and HDR; of MVA-related genes including *AACT*, *HMGS*, *MVK*, and *MVD*; and

downstream genes including *IDI*, *GPS*, *GPPS*, and *GGPPS*. *PtActin* was used as an internal

reference. Vertical bars represent mean ± SD (n = 3). *Significant difference between

HMGR-OEs and NT poplars at P < 0.05. **Significant difference between *HMGR*-OEs and NT

poplars at P < 0.01. ***Significant difference between *HMGR*-OEs and NT poplars at P <

0.001.

Figure 4. qRT-PCR analyses of transcript levels of MEP- and MVA-related genes in

PtDXR-OEs. Transcript levels of MEP-related genes including *DXS*, *MCT*, *CMK*, *HDS*, and

HDR; of MVA-related genes including *AACT*, *HMGS*, *HMGR*, *MVK*, and *MVD*; and

downstream genes including *IDI*, *GPS*, *GPPS*, and *GGPPS*. *PtActin* was used as an internal

reference. Vertical bars represent mean ± SD (n = 3). *Significant difference between

*DXR-OE*s and NT poplars at P < 0.05. **Significant difference between *PtDXR*-OEs and NT

poplars at P < 0.01. ***Significant difference between *PtDXR*-OEs and NT poplars at P <

0.001.

Figure 5. qRT-PCR analyses of the expression levels of ABA synthesis-related genes and

HPLC-MS/MS analyses of the contents of ABA, β-carotene, lycopene, and lutein in

PtHMGR-OEs. The expression levels of ABA synthesis-related genes including *NCED1*,

NCED3, *NCED5*, *NCED6*, *ZEP1*, *ZEP2*, and *ZEP3* are shown. Vertical bars represent mean ±

SD (n = 3). *significant difference at P < 0.05. **significant difference at P < 0.01.

***significant difference at P < 0.001. Three independent experiments were performed in this

study.

Figure 6. HPLC-MS/MS analyses of the contents of GA3 (A), TZR (B), IPA(C), DCS (D), and

CS (E) in *PtHMGR*-OEs and NT poplars.

Figure 7. qRT-PCR analyses of the expression levels of ABA synthesis-related genes and

HPLC-MS/MS analyses of the contents of ABA, β-carotene, lycopene, and lutein in

PtDXR-OEs and NT poplars. The expression levels of ABA synthesis related genes including

NCED1, *NCED3*, *NCED5*, *NCED6*, *ZEP1*, *ZEP2*, and *ZEP3* are shown. Vertical bars represent

mean ± SD (n = 3). *significant difference at P < 0.05. **significant difference at P < 0.01.

***significant difference at P < 0.001. Three independent experiments were performed in this

study.

Figure 8. HPLC-MS/MS analyses of the contents of GA3 (A), TZR (B), IPA(C), DCS (D), and

CS (E) in *PtDXR*-OEs and NT poplars.

Supplemental Figure 1. Amino acid sequences alignment of PtHMGR protein and other known

HMGR proteins. *A. thaliana* (NP_177775.2), *G. hirsutum* (XP_016691783.1), *M. domestica*

(XP_008348952.1), *M. esculenta* (XP_021608133.1), *P. persica* (XM_020569919.1), *O.*

sativa (XM_015768351.2), *T. cacao* (XM_007043046.2), *Z. mays* (PWZ28886.1).

Supplemental Figure 2. Molecular identification of *PtHMGR*-OEs. (A) Identification of

PtHMGR in *PtHMGR*-OEs and NT poplars based on the results of PCR. (B) Identification of

the transcript levels of *PtHMGR* in *PtHMGR*-OEs and NT poplars based on the results of

qRT-PCR. Three independent experiments were performed. *significant difference at $P < 0.05$.

**significant difference at $P < 0.01$. ***significant difference at $P < 0.001$.

Supplemental Figure 3. Analyses of the chromatogram of ABA standards via HPLC-MS/MS.

The chromatogram of standard ABA at (A) 0.1, (B) 0.2, (C) 0.5, (D) 2, (E) 5, (F) 20, (G) 50,

and (H) 200 ng/mL concentrations. (I) Equations for the ABA standard curves.

Supplemental Figure 4. Analyses of the chromatogram of GA1 standards via HPLC-MS/MS.

The chromatogram of standard GA1 at (A) 0.1, (B) 0.2, (C) 0.5, (D) 2, (E) 5, (F) 20, (G) 50, and

(H) 200 ng/mL concentrations. (I) Equations for the GA1 standard curves.

Supplemental Figure 5. Analyses of the chromatogram of GA3 standards via HPLC-MS/MS.

The chromatogram of standard GA3 at (A) 0.1, (B) 0.2, (C) 0.5, (D) 2, (E) 5, (F) 20, (G) 50, and

(H) 200 ng/mL concentrations. (I) Equations for the GA3 standard curves.

Supplemental Figure 6. Analyses of the chromatogram of GA4 standards via HPLC-MS/MS.

The chromatogram of standard GA4 at (A) 0.2, (B) 0.5, (C) 2, (D) 5, (E) 20, (F) 50, and (G) 200

ng/mL concentrations. (H) Equations for the GA4 standard curves.

Supplemental Figure 7. Analyses of the chromatogram of GA7 standards via HPLC-MS/MS.

The chromatogram of standard GA7 at (A) 0.1, (B) 0.5, (C) 2, (D) 5, (E) 20, (F) 50, and (G) 200

ng/mL concentrations. (H) Equations for the GA7 standard curves.

Supplemental Figure 8. Analyses of the chromatogram of zeatin standards via HPLC-MS/MS.

The chromatogram of standard zeatin at (A) 0.5, (B) 1, (C) 2, (D) 5, (E) 20, (F) 50, and (G) 200

ng/mL concentrations. (H) Equations for the zeatin standard curves.

Supplemental Figure 9. Analyses of the chromatogram of TZR standards via HPLC-MS/MS.

The chromatogram of standard TZR at (A) 0.1, (B) 0.2, (C) 0.5, (D) 2, (E) 5, (F) 20, (G) 50, and

(H) 200 ng/mL concentrations. (I) Equations for the TZR standard curves.

Supplemental Figure 10. Analyses of the chromatogram of IP standards via HPLC-MS/MS.

The chromatogram of standard IP at (A) 0.1, (B) 2, (C) 5, (D) 20, (E) 50, and (F) 200 ng/mL

concentrations. (G) Equations for the IP standard curves.

Supplemental Figure 11. Analyses of the chromatogram of IPA standards via HPLC-MS/MS.

The chromatogram of standard IPA at (A) 0.2, (B) 0.5, (C) 2, (D) 5, (E) 20, (F) 50, and (G) 200

ng/mL concentrations. (H) Equations for the IPA standard curves.

Supplemental Figure 12. Analyses of the chromatogram of CS standards via HPLC-MS/MS.

The chromatogram of standard CS at (A) 0.5, (B) 5, (C) 10, (D) 20, and (E) ng/mL

concentrations. (F) Equations for the CS standard curves.

Supplemental Figure 13. Analyses of the chromatogram of DCS standards via HPLC-MS/MS.

The chromatogram of standard DCS at (A) 0.5, (B) 2, (C) 10, (D) 50, and (E) ng/mL

concentrations. (F) Equations for the DCS standard curves.

Supplemental Figure 14. HPLC-MS/MS analyses of the contents of zeatin (A), IP (B), GA1 (C),

GA4 (D), and GA7 (E) in *PtHMGR*-OEs and NT poplars.

Supplemental Figure 15. HPLC-MS/MS analyses of the contents of zeatin (A), IP (B), GA1 (C),

GA4 (D), and GA7 (E) in *PtDXR*-OEs and NT poplars.

Table legends

Primers used in this study.

The English in this document has been checked by at least two professional editors, both native

speakers of English. For a certificate, please see:

http://www.textcheck.com/certificate/P2gZwq

(A)

(B)