***barcodetrackR*: an R package for the interrogation of clonal tracking data.**

Diego A. Espinoza[*,1,2], Ryland D. Mortlock[*,2], Samson J. Koelle[2,3], Chuanfeng Wu[2],

Cynthia E. Dunbar[2,+]

[1]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[2]Translational Stem Cell Biology Branch, National Heart, Lung and Blood Institute,

National Institutes of Health, Bethesda, MD, USA

[3]Department of Statistics, University of Washington, Seattle, WA, USA

*Equal contributions

[+]Corresponding author

Correspondence to:

Cynthia E. Dunbar

Translational Stem Cell Biology Branch, NHLBI, NIH

Building 10 CRC, Room 5E-3332

10 Center Drive

Bethesda, Maryland 20892

dunbarc@nhlbi.nih.gov

Abstract

Clonal tracking methods provide quantitative insights into the cellular output of

genetically labelled progenitor cells across time and cellular compartments. However,

the absence of a toolbox by which to interrogate these data has precluded the

development of standardized analytical frameworks within the field. Thus, we developed

barcodetrackR, an R package that provides users with tools for the analysis and

visualization of clonal dynamics across time and cellular compartments in clonal

tracking experiments. Here, we demonstrate the utility of *barcodetrackR* in exploring

longitudinal clonal patterns and lineage relationships in the context of clonal tracking

studies of hematopoietic stem and progenitor cells.

Introduction

Genetic labelling permits quantitative tracking of clonal progeny via high-throughput sequencing (clonal tracking) and provides opportunities to interrogate clonal dynamics in a number of *in vitro* and *in vivo* contexts. The two most common clonal tracking approaches, cellular barcoding and viral integration site recovery, have been primarily leveraged to interrogate hematopoietic stem and progenitor cell (HSPC) or immune cell dynamics both in model animals(1–5) and in humans(6,7). In these methodologies, integrating retro- or lentiviruses are used to transduce individual HSPCs or other target populations such that individual cells each contain a unique, permanent genetic tag or integration site label that can be recovered from progeny cells via high throughput sequencing (**Fig. 1**). Measurement of each label's abundance in the pool of all recovered labels is directly associated with the abundance of that clone within the labelled population being assayed, for instance T cells, B cells or myeloid cells. These lineage abundance measurements can provide insights not only into the bias, stability, and ontogenetic relationships of HSPCs(8), but also into the dynamics of clones within cell populations whose abundances are largely independent of HSPC behavior, such as certain T cell(9) and natural killer (NK) cell subsets(10). Furthermore, such clonal tracking methods have also been leveraged to provide valuable insight into the clonal dynamics of cancer progression(11), *in vitro* differentiation(12), and clonal dynamics of CAR-T cells(13).

Given the diversity of labelling and recovery strategies, as well as underlying differences in vector and barcode constructs, a number of approaches for recovery of sequences

from raw sequencing data and determination of "true" integration site or genetic tag from

sequencing artifacts or other confounders have been developed and are largely

approach-dependent(14–17). However, tools with which to perform downstream

analyses of the clonal abundances determined by these pipelines have not been

published or made publicly available; as a result, flexible open-source tools, such as

those that exist for single-cell RNA-sequencing(18,19) have been sought after by those

in the clonal tracking field in order to derive biological meaning in an accessible manner

from these large datasets(20). Such tools would also allow direct comparisons across

datasets or meta-analyses.

Here, we present our open-source R package, *barcodetrackR*. *barcodetrackR*

encompasses a variety of flexible tools that can provide insights into clonal dynamics

and the relationships between cellular compartments starting with clonal abundance

data. We illustrate the utility of *barcodetrackR* by analyzing publicly available clonal

tracking datasets from studies in lentivirally transduced non-human primates(8,10,21),

xenograft mice transplanted with human cord blood cells(22) and blast cells(23), and

lentiviral gene therapy patients(6,24). More details on each dataset and access paths

are summarized in Table 1.

Table 1: Example Datasets

Dataset	Description	Dataset availability	Reference
Six et al	Longitudinal clonal tracking of HSPC output in six human patients	Downloadable from https://github.com/B	(6)

	receiving lentivirally modified autologous HSPC transplants. Data from the Wiskott-Aldrich syndrome patient *WAS5* are selected for analysis.	ushmanLab/HSC_diversity/tree/master/data	
Belderbos et al	Longitudinal clonal tracking of HSPC output in twelve mice receiving lentivirally barcoded HSPC transplants from eight human umbilical cord blood donors. Data from the xenograft mouse *C21* are selected for analysis.	Downloadable from the Supplementary materials section of https://doi.org/10.1016/j.bbmt.2019.08.026.	(22)
Elder et al	Longitudinal clonal tracking of primograft acute lymphoblastic leukaemia (ALL) blasts in serially-infused NSG mice receiving lentivirally barcoded ALL blast cells. Data from the ALL sample *L4951* are selected for analysis.	GEO Accession No: GSE149170	(23)
Espinoza et al	Longitudinal clonal tracking of HSPC output in one rhesus macaque in which a lentivirally mediated clonal expansion occurred after receiving a lentivirally barcoded autologous HSPC transplant.	GEO Accession No: GSE153130	(21)
Wu et al	Longitudinal clonal tracking of natural killer cell subsets in five rhesus macaques receiving lentivirally barcoded autologous	Downloadable from the SUPPLEMENTARY MATERIALS section	(10)

	HSPC transplants. Data from the macaque *ZJ31* are selected for analysis.	of https://doi.org/10.1126/sciimmunol.aat9781	
Koelle et al	Longitudinal clonal tracking of HSPC output in five rhesus macaques receiving lentivirally barcoded autologous HSPC transplants. Data from the macaque *ZH33* are selected for analysis.	Downloadable from https://github.com/dunbarlabNIH/R-code-and-tabular-data	(8)
Clarke et al	Longitudinal clonal tracking of T cells in thirteen patients receiving lentivirally modified autologous HSPC transplants. Data from the X-linked severe combined immunodeficiency (X-SCID) patient U207 are selected for analysis.	Downloadable from https://doi.org/10.5281/zenodo.1256169	(24)

Results

Inferring lineage relationships based on global clonal distributions

Pairwise comparisons of clonal abundance profiles in clonal tracking data can provide insight into the relationships between upstream progenitor pools across cellular compartments. Here, we use *barcodetrackR* to determine and visualize the correlation values and dissimilarity indices (**Fig. 2**) between samples from three clonal tracking datasets (Six et al, Belderbos et al, Elder et al, Table 1) as a means to interrogate the similarities of upstream progenitor pools contributing across cellular compartments, gaining insights into lineage relationships. The Six dataset contains clonal tracking information (representing individual viral integration site read counts) from longitudinally collected patient T cell, B cell, granulocyte (Gr), monocyte (Mo), and natural killer cell (NK) samples following autologous lentiviral HSPC gene therapy. Visualizing the pairwise correlations between sample clonal abundances, we find that the Gr and Mo samples share high correlation with one another, while the T cell, B cell, and NK samples show lower correlation with samples from other lineages, but high correlation between different timepoints within the same lineage (**Fig. 2A**). A similar pattern is observed when plotting the Bray-Curtis dissimilarity indices between samples from the Six dataset, projected into two dimensions using principal coordinates analysis (PCoA), where the first axis of variation separates NK cells from other lineages based on their clonal abundances, and the second separates T cells, B cells, and myeloid (Gr and Mo) cells (**Fig. 2B**). Altogether, these analyses of the Six dataset demonstrate that the myeloid lineages clonal are closely coupled and thus likely arise from shared pathways

originating from the same HSPC pool, in comparison to disparate generation of mature

T, B, and NK lineages.

These pairwise measures can be also used to compare clonal abundances across

anatomical compartments. The Belderbos dataset contains clonal abundance

information from a number of sorted and unsorted immune cell samples from bone

marrow (BM) sites and from the spleen of a mouse transplanted with a lentivirally

barcoded human cord blood CD34+ HSPC xenograft. Analyzing this dataset, we

observe high correlation between T cell samples across all anatomical sites, while B cell

and Gr samples show high correlation to one another only within each anatomical site

(**Fig. 2C**). This pattern is evident in the PCoA plot (**Fig. 2D**) as well. Unsorted cell

samples from the spleen and pelvic BM vary in their clonal relationships to other

samples, likely because of the underlying heterogeneity of the lineage composition of

these bulk samples (**Fig. 2C**).

In sum, these analyses support the notion that geographically isolated HSPC pools are

responsible for the clonal composition of their respective geographic niches, and that

the clonal composition of T cells across sites is largely independent from the output of

these pools, confirming the thymic-dependent developmental pathway for T cells, with

passage through the thymus. Indeed, geographic isolation of HSPC output has been

observed in another study in macaques; however, it is of note that within the macaque

study, T cell output early after transplantation utilizing total body irradiation conditioning

was instead found to be anatomically compartmentalized and dependent at least short-

term on these geographically isolated HSPC pools(25).

Comparing clonal distribution across animals from serial transplant experiments can also provide insight into the self-renewal capacity of engrafted, clonally marked cells. The Elder dataset contains clonal abundance information from serial xenograft mouse transplants of lentiviral-transduced ALL blasts. We observed high correlation of clonal abundances between samples collected from primary, secondary, and tertiary transplant recipient mice, excluding a few sites in the primary transplant (**Fig. 2E**). This aligns with the results presented by Elder et al(23) noting equipotential functional capability of ALL cells with some variation between sites, based on random sampling of the population of engrafted ALL cells. Samples from the same "generation" of serial transplant cluster together in principal coordinate space, supporting the notion presented in the study that ALL founder cells retain self-renewal capacity over several serial transplants (**Fig. 2F**). The distinct groupings of clones based on anatomic sampling site within the primary transplanted animals suggests that ALL clonal output also appears to be geographically compartmentalized, at least initially. Altogether, these three examples illustrate the utility of *barcodetrackR* in probing the relationships between the global clonal distributions of samples collected from various lineages or time points, providing valuable insights from a number of diverse biologic contexts.

Determining clonality based on clonal counts and diversity measures

In clonal tracking studies, both clonal counts and diversity measures can provide insight into the clonality of progenitor cell pools. Here, we utilize *barcodetrackR* to assess

clonality by visualizing the detected clone counts and Shannon diversities of samples

from three datasets over time (**Fig. 3**).

When quantifying clone numbers within the Six dataset, we show hundreds of unique

integration sites retrieved across five purified peripheral blood lineage samples, with a

larger number of clones detected in B cells and T cells as compared to Gr, Mo, and NK

cells at most individual timepoints (**Fig. 3A**). Decreasing clonal diversity (Shannon

index) in the NK cell lineage, as compared to other lineages, indicates that over time, a

smaller number of clones accounts for a larger fraction of hematopoiesis in the NK cell

compartment (**Fig 3B**). This implies a more oligoclonal population of contributing

progenitors as compared to other cell types sampled by Six et al. The finding that the

mature NK cell compartment is largely composed of a few high-contributing clones post-

transplantation is in agreement with rhesus macaque barcoded autologous HSPC

transplant studies(10).

Next, we analyzed patterns in the number of detected clones and Shannon diversities

over time in peripheral blood samples from a single mouse xenograft obtained from the

Belderbos dataset. We show that more clones were detected from the bulk peripheral

blood sample at sacrifice than at the first time point (green line, **Fig. 3C**). The Shannon

diversity of bulk samples decreased after the 9 week time point before stabilizing (**Fig.**

3D), underscoring the notion that clone counts alone are not ideal measures of sample

diversity. These findings suggest that within this xenograft transplantation model, the

diversity of HSPC output becomes stable over time, in agreement with previous long

term clonal tracking studies in macaque(8) and human(7).

Finally, we use *barcodetrackR* to quantify an extreme case of minimal clonal counts and

Shannon diversities in the context of clonal hematopoiesis using the Espinoza dataset

(Table 1). In this study, multiple lentiviral insertions in an HSPC mediated dysplastic

clonal erythroid and myeloid marked expansion, while largely sparing the lymphoid

lineages. In agreement with the findings of the study, we find that the longitudinal clone

numbers contributing to the B and T cell lineages fluctuate, but that the Shannon

diversity index of these lineages remains high, especially at early time points, indicating

polyclonal contribution to the lymphoid lineages (**Fig 3E**). However, after day 266 post-

transplant, we observe a massive drop-off in both the number of unique clones detected

(**Fig 3E**) and the Shannon diversity (**Fig 3F**) within the myeloid lineages, down to the

numbers detected in the nucleated red blood cell (nRBC) lineage. This coincides

chronologically with the development of clonal hematopoiesis in the myeloid lineage. In

the Espinoza study, 6 detected genetic tags in this dataset were all in fact found to

correspond to a single lentivirally transduced cell with 9 insertions (3 of which were

virtually undetected with the available dataset's methodology due to deletions in the

insertion proviral sequences), verifying that this pattern is representative of clonal

hematopoiesis(21). While in the above examples we utilize unique detected clones at

each time point, cumulative clone counts can also be calculated and visualized for each

example (**Fig. S1**) and provide a complementary view of clone numbers over time.

Altogether, these examples emphasize the utility of diversity measures and

visualizations in interrogating the clonal output of progenitor pools in a number of

contexts.

Analyzing longitudinal clonal dynamics based on clonal abundances

Longitudinally tracking the abundance of individual clones can provide insight into clonal

dynamics within lineages. We employed *barcodetrackR* to analyze longitudinal NK cell

samples from an animal in the Wu et al study (Table 1), in which NK cell clonal

dynamics were interrogated over 3 years in rhesus macaques receiving lentivirally

barcoded autologous HSPCs(10). The detected clones in the NK cell compartment

within this study remained largely independent from the HSPC pool responsible for the

majority of non-NK hematopoiesis, enabling tracking of NK cell/progenitor clonal output

rather than multilineage HSPC clonal output. We first visualize all individual NK cell

clones from the Wu dataset in a binary heat map that depicts the presence or absence

of all clones observed at 0.01% abundance or greater in at least one NK cell sample

(**Fig 4A**). When ordering clones (y-axis) based on their first time point of detection, we

find that new NK cell clones are detected at each time point, but that the number of

newly detected clones decreases at later time points.

Next, we analyzed distinct clonal dynamics in individual NK cell clones using

barcodetrackR to generate a heat map showing the abundance of the top ten NK cell

clones from each sample over time (**Fig. 4B**). We visualize only the top clones in order

to focus on the clones responsible for the majority of this cellular compartment's clonal

composition, with stars on the heatmap indicating the top ten contributing clones in each

sample. One group of clones contributed at high levels for 3 months but subsequently

declined. Of the top ten clones from the second time point (4 months), some clones

declined in abundance while others increased in abundance at the 6-month time point

before subsequently declining. A group clones became high abundance at 8.5 months

and continued to contribute a large fraction of NK cell activity through the 20-month time

point. Finally, a set of clones became high contributors at the 17.5-month time point

after being virtually undetectable at previous timepoints. This analysis reveals the

waxing and waning patterns of high-abundance NK cell clones over time, which were

further interrogated in the Wu study.

Lastly, we utilized statistical testing within *barcodetrackR* to view changes in proportion

within the NK cell samples, marking clones with a star which had a statistically

significant change (Chi-squared test) in abundance in the labelled sample in

comparison to the previous sample (**Fig 4C**). This type of visualization and analysis

further highlights the highly dynamic clonal patterns within the NK cell compartment and

provide new insights into NK cell biology. Altogether, these results indicate that the

longitudinal tracking of highly abundant clones within datasets can provide insight into

clonal dynamics at a single-clone level.

Quantifying lineage bias based on shared clonality

Clonal tracking studies measure HSPC clonal contributions to different mature blood

cell lineages. Thus, the lineage bias of HSPCs, such as those that skew towards

myeloid or lymphoid lineages, can be interrogated on a global scale by comparing the

ratio of clonal abundances between two specific lineages. Here, we use *barcodetrackR*

to probe this concept of lineage bias in the Six clinical gene therapy trial dataset(6) and

the Koelle rhesus macaque dataset(8)(Table 1), both based on lineage-purified samples

following autologous transplantation with genetically tagged HSPCs.

We first track the density of clones, weighted by their added proportions, at each value

of lineage bias (log ratio) between the Gr and T cell lineages across multiple timepoints

in the Six dataset. We find the presence of three high-contributing sets of clones as

determined by Gr/T lineage bias: Gr-biased (rightmost peak), balanced clones (middle

peak), and T-biased (leftmost peak) (**Fig. 5A**). By systematically comparing each cell

type in the dataset, we find that three sets of clones can be found when comparing

Mo/T, Gr/B, or Mo/B lineages (**Fig. S2**) further supporting differences in upstream

progenitors accounting for myeloid versus lymphoid lineages. In contrast, when

comparing Gr/Mo or T/B lineages, we find that the majority of clones have balanced

contribution to the two lineages (**Fig. S2**). Interestingly, clones contributing to the NK

cell samples are predominantly unilineage, sharing very little clonality with other

lineages, including other lymphoid lineages such as T and B cells (**Fig. S2**). The

distinctness of NK cell hematopoiesis from other lineages is in line with clonal tracking

studies performed in a rhesus macaque animal model(2,10).

Conducting the same analysis on longitudinal samples from the Koelle dataset reveals

the presence of Gr-biased, balanced, and T-biased clones at the 4.5-month timepoint

(**Fig. 5C**), consistent with the Six dataset. However, there is an increase in abundance

of balanced clones at later timepoints, indicating a shift towards hematopoiesis from

multipotent upstream progenitors, capable of reconstituting both myeloid and lymphoid

lineages. This is also the case when comparing the T cell lineage to the Mo lineage as

the majority of clones contribute similar abundances to Gr and Mo lineages (**Fig. S3**).

We next use *barcodetrackR* to construct an abundance-weighted chord diagram

between three lineages in the Six dataset. We selected the Gr, T, and Mo lineages at

the final 55-month time point (**Fig. 5B**), finding that a large fraction of detected

hematopoiesis at this time point is shared between all three lineages (purple). However,

there also exist clones detected only in two lineages (yellow, blue, green), and biased

clones only found in one lineage, indicated by the white space around the perimeter.

Likewise, a similar pattern is observed in the Koelle dataset at the final 38-month time

point (**Fig. 5D**) with a large fraction of detected hematopoiesis arising from clones

detected in all three lineages (purple). The fraction of detected hematopoiesis arising

from T-Gr or T-Mo restricted clones (blue, yellow respectively), however, is minimal

compared to that arising from Gr-Mo restricted clones (green), supporting the notion of

upstream progenitors biased towards the myeloid lineage within this dataset. Thus,

barcodetrackR provides a number of functions useful for inferring the lineage biases of

upstream progenitors from clonal tracking data.

barcodetrackR is versatile and includes a user-friendly graphical user interface

We highlight the versatility of *barcodetrackR* by analyzing clonal tracking data collected

from TCR sequencing by visualizing the number of T cell clones detected and the

Shannon diversity of longitudinal samples from X-SCID patients treated with HSPC

gene therapy(24). In agreement with the original paper, we found that the highest

number and diversity of T cell clones were detected at the 6 month time point for patient

U207 (**Fig. S4**).

The *barcodetrackR* package includes a graphical user interface (built using *shiny*(26)) to

allow researchers without programming experience to use these advanced quantitative

tools. Once the package is downloaded locally, users can launch the graphical user

interface using the R command *barcodetrackR::launchApp().* After uploading genetic

tag count matrices and metadata, users can toggle between tabs corresponding to any

of the desired analyses described above as well as several additional tools. Within each

interactive function, users can specify calculation and display methods in order to create

reproducible analyses and publication-quality data visualizations.

Discussion

A recent gathering of over 30 researchers in the clonal tracking field (2018

StemCellMathLab workshop(20)) formalized a call for the development of open-source

tools for the analysis of clonal tracking data in order to promote rigor and reproducibility

within the field. Here, we now provide and showcase our open-source R package

barcodetrackR, which encompasses an extensive, flexible, and accessible set of tools in

order to address these needs and serve as a critical foundation on which to build further

analytical approaches in the clonal tracking field. While tools for the processing of the

raw sequencing data from clonal tracking experiments have been previously developed

to improve the accuracy of clonal abundance quantification(16), *barcodetrackR*

represents the first formal tool developed to interrogate the underlying biology

represented by these resulting clonal abundances. As shown, *barcodetrackR* is a

multifaceted toolkit and has diverse applications, underscoring the utility of using

complementary data analysis methods and visualizations to probe and/or develop

appropriate hypotheses relevant to the biological context at hand. The development and

implementation of a simple-to-use *shiny* app further adds to the utility of the package by

making it more accessible to the clonal tracking community, which continues to expand

as sequencing costs continue to decline and methodologies continue to improve.

Although our package encompasses a large number of tools and methods by which to

derive insight from clonal tracking experiments, it is by no means an exhaustive toolbox,

and we envision continuing to add to it in the future in order to address new biologic

questions that arise. While the majority of clonal tracking experimental designs in the

literature have precluded the acquisition of replicate samples and have often

encompassed a relatively low number of transplanted humans and/or animals, future

studies will likely be able to acquire biological replicates in a number of different

contexts to allow for more rigorous statistical testing of sample relationships and clonal

dynamics. And while clustering methods to identify populations of clones with similar

properties have thus far been limited to hierarchical(10) and k-means(6) in the literature,

and to hierarchical clustering within *barcodetrackR*, the growing development of

clustering frameworks of single cells in the scRNA-seq field may provide a future basis

by which to identify clusters of clones based on longitudinal behavior and distribution

across compartments(27). Furthermore, other challenges remain in the clonal tracking

field, namely, the necessity for improved sharing and aggregating of data. We believe

barcodetrackR can be of high utility to the clonal tracking field and serve as an

important step towards the goal of building a more robust and reproducible analytical

framework in the field.

Methods

Package availability

The *barcodetrackR* package is freely available from GitHub under a Creative Commons

0 license and can be found at https://github.com/dunbarlabNIH/barcodetrackR. A frozen

version of the package at the time of publication will be placed on Zenodo. The

analytical and visualization tools included in *barcodetrackR* rely on the *ggplot2*(28),

vegan(29), *proxy*(30), and *circlize*(31) packages, as well as a number of packages in

the *tidyverse*(28) suite. The user interface is built using the *shiny* package in R(26).

barcodetrackR can be installed in R using the *devtools*(28) utility and the command

devtools::install_github("dunbarlabNIH/barcodetrackR"). Within the GitHub repository,

we include a vignette illustrating the use of *barcodetrackR* on several published

barcoding datasets, which is available as an R markdown file or in html format online at:

https://dunbarlabNIH.github.io/barcodetrackR. All figures included in the manuscript

were generated using the *barcodetrackR* package, and the R markdown file

"create_all_figures.Rmd" is included in the *barcodetrackR* GitHub.

Dataset availability

The datasets used in this study are publicly available from published barcoding

experiments and detailed in Table 1. Data were pre-processed in R to create tabular

data files amenable for entry into *barcodetrackR,* and these procedures are outlined in

scripts within the inst/sample_data directory within the *barcodetrackR* package on

GitHub.

Data collection, genetic tag retrieval, and normalization

Multiple clonal tracking methodologies exist in the literature(30), with the most recent

methods relying on next-generation sequencing to retrieve lineage tracing elements.

Several analysis pipelines exist for the retrieval and error-correction of lineage tracing

elements from sequencing data(14–17). The experimental techniques utilized, the

number of cells sampled, the level of tagged cell within the population, the sequencing

platform applied, and the computational method of genetic tag extraction affect the

number and frequency of tags detected in a lineage tracing study.

The *barcodetrackR* package can operate on any dataset that contains rows as

observations and columns as samples, regardless of which experimental method for

genetic labelling and approaches for tag retrieval were used. When creating a

SummarizedExperiment (SE) object from the publicly available *Bioconductor*

repository(32) for input into *barcodetrackR*, users have the option of including a

threshold to exclude low-abundance occurrences that are more likely to come from

noise or sequencing error(2). Using a threshold of 0.005, for example, retains genetic

tags which are present at an abundance of 0.5% or greater in at least one sample.

Within this paper, the Six, Elder, Espinoza, Wu, and Koelle datasets were used with a

threshold of 0.0001. The Belderbos and Clarke datasets were used with no threshold.

Instantiating a SE through the function *create_SE* automatically creates the following

assays: *counts*: the raw values from the input dataframe, *percentages*: the per-column

proportions of each entry in each column, *ranks*: the rank of each entry in each column,

normalized: the normalized read values in counts-per million (CPM), and *logs*: the log of

the normalized values. The default normalization is counts per million, and log-

normalized values are calculated by taking the log of plus-one normalized data so that

zeros are retained. Both the scale factor and the log base can be passed as arguments

to the *create_SE* command. The use of an SE object permits the addition of custom

assays to the object to facilitate flexibility (e.g. custom normalization strategies).

Global clonal distributions

Users can view correlations between samples on a grid using the *cor_plot* function,

longitudinally using the *autocor_plot* function, or for two samples using the *scatter_plot*

function. Choices for correlation measures include *"pearson", "spearman",* or *"kendall."*

The functions include display parameters as inputs, such as whether to display

correlation values in a *cor_plot* grid as color, colored circles, or showing the actual

correlation value.

The *mds_plot* function calculates dissimilarity indices between samples using any

distance metric within the *vegdist* function from the R package *vegan*(29). At time of

publication, these include *"manhattan", "euclidean", "canberra", "clark", "bray",*

"kulczynski", "jaccard", "gower", "altGower", "morisita", "horn", "mountford", "raup",

"binomial", "chao", or *"cao".* The distance methods produce a matrix composed of the

distances between samples, given the composition of genetic tags in each sample.

Principal coordinates analysis is performed on the distance matrix using the base R

function *cmdscale* in order to display dissimilarity between samples in two dimensions.

Clonal diversity

Three measures of within-sample diversity can be calculated by the function

clonal_diversity: shannon diversity (H') , simpson diversity (λ), and inverse-simpson,

which is calculated as 1/λ. Their equations are as follows:

$$H' = -\sum_{i=1}^{R} p_i \ ln \ p_i$$

$$\lambda = \sum_{i=1}^{R} p_i^2$$

407   Where *R* is the total number of species (in this case genetic tags), and p_i is the

408   proportion of each gentic tag in the sample. Additionally, users can simply display the

409   nominal count or Shannon count. Shannon count is calculated as:

$$Sh_{count} = e^{H'}$$

To accurately compare diversity between samples, the same number of labeled cells

should be used as starting material for quantification. This is especially important when

assessing diversity based on the nominal count of genetic tags retrieved. Belderbos et

al showed that the Shannon count is stable with respect to filtering thresholds and stays

below the theoretical library size upon re-sampling(22). Therefore, in some cases, it

may be beneficial to use Shannon count rather than nominal genetic tag counts when

comparing diversity between samples.

Clonal patterns

Heat maps created by the function *barcode_ggheatmap* display clonal abundances

across samples by coloring cells based on the log-normalized abundance of each clone.

Users specify the ordering of samples along the x-axis and the number of clones

displayed per sample. By default, the top clones from each sample are marked by a

star, but the user can also choose to label each entry in the heatmap by percentage or

number of reads. The function includes aesthetic parameters, such as the label size,

percent scale, and color scale, which can also be easily controlled within the

barcodetrackR graphical user interface.

Under default settings for the heat map function, individual clones are hierarchically

clustered along the y-axis based on their log-abundance (or other specified assay)

across samples. Setting the *"dendro"* parameter to TRUE allows users to see this

clustering and setting the *"clusters"* parameter to a non-zero value will label hierarchical

clusters of clones. The dendrogram is drawn using *ggdendro*(33). Users can choose

from a number of methods to calculate the distance metric (any method included in the

proxy R package) and clustering (*"ward.D", "ward.D2", "single", "complete", "average",*

"mcquitty", "median", or *"centroid"*). Clones can also be ordered by the first sample in

which they had a non-zero abundance by setting the *"row_order"* parameter to

"emergence."

To track the emergence of clones over time, the function *barcode_binary_heatmap* will

display a heatmap indicating the presence or absence of clones in each sample. The

"threshold" parameter to the function dictates the limit of detection. Clones with

percentage abundance below this threshold in a given sample will be set to "absent."

Clones are ordered in the binary heatmap based on the first sample in which they

emerged (had a non-zero abundance).

The function *barcode_ggheatmap_stat* allows users to quantify changes in clone

abundance based on statistical testing. This function requires the sample size of cells

for each sample which cannot be calculated from the genetic tag count data. The

sample size should be the number of labeled cells before amplification, because this is

the population of cells which the clonal tracking data represent. To compare barcode

abundances between samples, users can choose from a *"chi-squared"* or *"fisher"* exact

test. The tests operate on a contingency table for each clone to determine whether the

clone changed in proportion based on the abundance and sample size. By default, each

sample is compared to the previous, but users can also specify to compare each

sample to a single reference sample (such as the initial time point) by setting the

"stat_option" parameter to *"reference."* The user can specify a p-value threshold to

assign significance and can choose to display p-values on the heat map, rather than

stars indicating statistically significant changes. Also, the user can specify to only show

clones which increase or decrease in proportion through the *"stat_display"* parameter.

Clonal bias

The *ridge_plot* function calculates bias as a continuous variable and displays the

density of clones at each level of log bias. In order to calculate log bias between clones

that are only present in one sample, log bias is calculated as followed within the ridge

plot function:

$$log\ bias_{b.c.A} = log2\left(\frac{b.c.A\ normalized_{sample\ 1}\ +\ 1}{b.c.A\ normalized_{sample\ 2}\ +\ 1} \right)$$

The normalized value is taken from the SE slot which is scaled to counts per million.

The option to only analyze clones present in both samples is also included in the

ridge_plot function through the parameter *"remove_unique"*. The density of clones at

each value of log bias is estimated using the kernel density estimator included in the

ggplot2 R package. When the *"weighted"* parameter is set to TRUE, the function

weights the density estimation by the *combined abundance* of the clone between the

two samples, calculated as:

$combined\ abundance\ =\ b.c.A\ normalized_{sample\ 1}\ +\ b.c.A\ normalized_{sample\ 2}$

Chord diagrams

The *circos_plot* function utilizes the circlize package in R(31) to display shared clones

between samples. Samples are shown as regions around a circle with their shared

clonality shown as links between regions. With each unique combination, a new link is

created with a unique color drawn from a sequential color palette. When the *"weighted"*

option is set to *"FALSE"*, the function operates on the *counts* assay. Therefore, the

length of each region around the circle represents the number of clones detected in

each sample, and the width of links between regions is proportional to the number of

shared clones. When the *"weighted"* option is set to *"TRUE"*, the function operates on

the *percentages* slot. Each region around the circle has the same length corresponding

to 100%, and the links between regions correspond to the fractional abundance of the

shared clones within each sample. Therefore, when the "*weighted"* option is set to

"TRUE", the same link can have a different width at each connection to a region.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Accessibility to all datasets used is outlined in Table 1. *barcodetrackR* is an open-source R package on GitHub at https://github.com/dunbarlabNIH/barcodetrackR and the current version of the R package will be frozen on Zenodo at the time of publication.

Competing interests

Not applicable.

Funding

DAE was supported by NIH Medical Scientist Training Program T32 GM07170. RDM, SJK, CW and CED were supported by the intramural program of the National Heart, Lung and Blood Institute.

Authors' contributions

DAE and RDM wrote the manuscript. DAE and RDM developed code and performed analysis of existing datasets. SJK and CW aided with development of visualizations. CED supervised the project and edited the manuscript.

Acknowledgements

The authors thank members of the Dunbar lab for helpful feedback in revision of this manuscript.

References

1. Lu R, Neff NF, Quake SR, Weissman IL. Tracking single hematopoietic stem cells in vivo using high-throughput sequencing in conjunction with viral genetic barcoding. Nat Biotechnol. 2011 Oct;29(10):928–33.

2. Wu C, Li B, Lu R, Koelle SJ, Yang Y, Jares A, et al. Clonal Tracking of Rhesus Macaque Hematopoiesis Highlights a Distinct Lineage Origin for Natural Killer Cells. Cell Stem Cell. 2014 Apr;14(4):486–99.

3. Radtke S, Adair JE, Giese MA, Chan Y-Y, Norgaard ZK, Enstrom M, et al. A distinct hematopoietic stem cell population for rapid multilineage engraftment in nonhuman primates. Sci Transl Med. 2017 Nov 1;9(414):eaan1145.

4. Kim S, Kim N, Presson AP, Metzger ME, Bonifacino AC, Sehl M, et al. Dynamics of HSPC Repopulation in Nonhuman Primates Revealed by a Decade-Long Clonal-Tracking Study. Cell Stem Cell. 2014 Apr;14(4):473–85.

5. Gerrits A, Dykstra B, Kalmykowa OJ, Klauke K, Verovskaya E, Broekhuis MJC, et al. Cellular barcoding tool for clonal analysis in the hematopoietic system. Blood. 2010 Apr 1;115(13):2610–8.

6. Six E, Guilloux A, Denis A, Lecoules A, Magnani A, Vilette R, et al. Clonal tracking in gene therapy patients reveals a diversity of human hematopoietic differentiation programs. Blood. 2020 Apr 9;135(15):1219–31.

7. Biasco L, Pellin D, Scala S, Dionisio F, Basso-Ricci L, Leonardelli L, et al. In Vivo Tracking of Human Hematopoiesis Reveals Patterns of Clonal Dynamics during Early and Steady-State Reconstitution Phases. Cell Stem Cell. 2016 Jul;19(1):107–19.

8. Koelle SJ, Espinoza DA, Wu C, Xu J, Lu R, Li B, et al. Quantitative stability of hematopoietic stem and progenitor cell clonal output in rhesus macaques receiving transplants. Blood. 2017 Mar 16;129(11):1448–57.

9. Brugman MH, Wiekmeijer A-S, van Eggermond M, Wolvers-Tettero I, Langerak AW, de Haas EFE, et al. Development of a diverse human T-cell repertoire despite stringent restriction of hematopoietic clonality in the thymus. Proc Natl Acad Sci USA. 2015 Nov 3;112(44):E6020–7.

10. Wu C, Espinoza DA, Koelle SJ, Yang D, Truitt L, Schlums H, et al. Clonal expansion and compartmentalized maintenance of rhesus macaque NK cell subsets. Sci Immunol. 2018 Nov 2;3(29):eaat9781.

11. Merino D, Weber TS, Serrano A, Vaillant F, Liu K, Pal B, et al. Barcoding reveals complex clonal behavior in patient-derived xenografts of metastatic triple negative breast cancer. Nature Communications. 2019 Feb 15;10(1):1–12.

550   12.   Porter SN, Baker LC, Mittelman D, Porteus MH. Lentiviral and targeted cellular
551   barcoding reveals ongoing clonal dynamics of cell lines in vitro and in vivo.
552   Genome Biology. 2014 May 30;15(5):R75.

553   13.   Sheih A, Voillet V, Hanafi L-A, DeBerg HA, Yajima M, Hawkins R, et al. Clonal
554   kinetics and single-cell transcriptional profiling of CAR-T cells in patients
555   undergoing CD19 CAR-T immunotherapy. Nature Communications. 2020 Jan
556   10;11(1):1–13.

557   14.   Berry CC, Nobles C, Six E, Wu Y, Malani N, Sherman E, et al. INSPIIRED:
558   Quantification and Visualization Tools for Analyzing Integration Site Distributions.
559   Molecular Therapy - Methods & Clinical Development. 2017 Mar;4:17–26.

560   15.   Sherman E, Nobles C, Berry CC, Six E, Wu Y, Dryga A, et al. INSPIIRED: A
561   Pipeline for Quantitative Analysis of Sites of New DNA Integration in Cellular
562   Genomes. Molecular Therapy - Methods & Clinical Development. 2017 Mar;4:39–
563   49.

564   16.   Thielecke L, Cornils K, Glauche I. genBaRcode: a comprehensive R-package for
565   genetic barcode analysis. Wren J, editor. Bioinformatics. 2020 Apr 1;36(7):2189–
566   94.

567   17.   Bramlett C, Jiang D, Nogalska A, Eerdeng J, Contreras J, Lu R. Clonal tracking
568   using embedded viral barcoding and high-throughput sequencing. Nat Protoc. 2020
569   Apr;15(4):1436–58.

570   18.   Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression
571   data analysis. Genome Biol. 2018 Dec;19(1):15.

572   19.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
573   Comprehensive Integration of Single-Cell Data. Cell. 2019 Jun;177(7):1888-
574   1902.e21.

575   20.   Lyne A-M, Kent DG, Laurenti E, Cornils K, Glauche I, Perié L. A track of the clones:
576   new developments in cellular barcoding. Experimental Hematology. 2018
577   Dec;68:15–20.

578   21.   Espinoza DA, Fan X, Yang D, Cordes SF, Truitt LL, Calvo KR, et al. Aberrant
579   Clonal Hematopoiesis following Lentiviral Vector Transduction of HSPCs in a
580   Rhesus Macaque. Molecular Therapy. 2019 Jun 5;27(6):1074–86.

581   22.   Belderbos ME, Jacobs S, Koster TK, Ausema A, Weersing E, Zwart E, et al. Donor-
582   to-Donor Heterogeneity in the Clonal Dynamics of Transplanted Human Cord
583   Blood Stem Cells in Murine Xenografts. Biology of Blood and Marrow
584   Transplantation. 2020 Jan 1;26(1):16–25.

23. Elder A, Bomken S, Wilson I, Blair HJ, Cockell S, Ponthan F, et al. Abundant and
equipotent founder cells establish and maintain acute lymphoblastic leukaemia.
Leukemia. 2017 Dec;31(12):2577–86.

24. Clarke EL, Connell AJ, Six E, Kadry NA, Abbas AA, Hwang Y, et al. T cell
dynamics and response of the microbiota after gene therapy to treat X-linked
severe combined immunodeficiency. Genome Med. 2018 Dec;10(1):70.

25. Wu C, Espinoza DA, Koelle SJ, Potter EL, Lu R, Li B, et al. Geographic clonal
tracking in macaques provides insights into HSPC migration and differentiation. J
Exp Med. 2018 Jan 2;215(1):217–32.

26. Chang W, Cheng J, Allaire JJ, Xie Y, McPherson J. shiny: Web Application
Framework for R [Internet]. 2020. Available from: https://CRAN.R-
project.org/package=shiny

27. Kiselev VY, Andrews TS, Hemberg M. Challenges in unsupervised clustering of
single-cell RNA-seq data. Nat Rev Genet. 2019 May;20(5):273–82.

28. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al.
Welcome to the Tidyverse. Journal of Open Source Software. 2019 Nov
21;4(43):1686.

29. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package [Internet]. 2019. Available from: https://CRAN.R-
project.org/package=vegan

30. Meyer D, Buchta C. proxy: Distance and Similarity Measures [Internet]. 2020.
Available from: https://CRAN.R-project.org/package=proxy

31. Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize implements and enhances
circular visualization in R. Bioinformatics. 2014 Oct 1;30(19):2811–2.

32. Morgan M, Obenchain V, Hester J, Pagès H. SummarizedExperiment:
SummarizedExperiment container [Internet]. 2020. Available from:
https://bioconductor.org/packages/SummarizedExperiment/

33. de Vries A, Ripley BD. ggdendro: Create Dendrograms and Tree Diagrams using
"ggplot2" [Internet]. 2016. Available from: https://CRAN.R-
project.org/package=ggdendro

Figure Legends

Figure 1: Clonal tracking experimental design

Clonal tracking experiments generally follow the depicted framework, which

encompasses, in order, the genetic labelling of cells to create an integrated DNA tag,

transplantation or adoptive transfer of these cells into a recipient, acquisition of cellular

progeny from the recipient following transplantation or adoptive transfer, genetic tag

retrieval from progeny cells followed by high-throughput sequencing, algorithmic

quantification of detected individual unique tags, and finally, downstream analyses,

where the *barcodetrackR* toolkit can be utilized.

Figure 2: Global clonal distributions

Pairwise Pearson correlation plots between longitudinal samples from the Six dataset

(**A**). Row and column labels indicate months post-transplant (*m*) and cell type (*Gr,*

Granulocyte; Mo, Monocyte; NK, Natural Killer cell). Bray-Curtis dissimilarity indices

between samples from the Six dataset (**B**), grouped by cell type and labeled based on

months post-transplant. The x and y-axis represent the two main axes of variation after

conducting principal coordinate analysis on the Bray-Curtis measures of dissimilarity

(*MDS)*. Pairwise Pearson correlation plots between samples from different anatomical

sites of a single transplanted mouse at euthanasia from the Belderbos dataset (**C**). Row

and columns labels describe the anatomical site (*BM, Bone Marrow)* followed by the cell

type (*U, Unsorted samples*). Bray-Curtis dissimilarity indices between samples from the

Belderbos dataset (**D**) grouped by the anatomical site and labeled by cell type. Pairwise

Pearson correlation values between samples of the same set of serial xenograft

transplants from the Elder dataset (**E**). Row and column labels describe the animal code

(*e.g. AE12*), followed by the anatomical site, then the serial transplant designation *(pri,*

Primary; sec, Secondary, ter, Tertiary), followed by the donor animal code if it is a *sec*

or *ter* sample. *AE12* is the primary recipient of ALL blast cells, *AE32* is the secondary

recipient receiving cells from the primary animal *AE12*, and *AE88, AE89, AE90* are

tertiary recipients receiving cells from *AE32*. Bray-Curtis dissimilarity indices between

samples from the Elder dataset (**F**) colored by the mouse of origin and labeled by the

anatomical site.

Figure 3: Measures of clonal diversity

The number of clones detected in each lineage (**A**) and the Shannon diversity index of

each sample (**B**) from the Six dataset. The number of clones detected in each lineage

(**C**) and the Shannon diversity index of each sample (**D**) from the Belderbos dataset.

The number of clones detected in each lineage (**E**) and the Shannon diversity index of

each sample (**F**) from the Espinoza dataset. X-axes show months, weeks or days post-

transplant with "sac" corresponding to the timepoint of euthanasia in the Belderbos

dataset. The clone count reflects the number of unique clones detected in each sample,

not the cumulative count at each timepoint. Shannon diversity is calculated on a per-

sample basis based on the clonal population of each sample, not the cumulative

number of clones. *Gr, Granulocyte; Mo, Monocyte; bulk, unsorted population; CD34p,*

CD34 positive cell; nRBC, nucleated Red Blood Cell

Figure 4: Longitudinal clonal patterns

(**A**) Binary heat map showing the presence (blue) or absence (white) of 11,799 individual clones detected at a proportion of 0.01% or greater in at least one NK cell sample from the Wu et al dataset. Columns represent samples and rows represent individual clones ordered by their first time point of detection. (**B**) Heat map showing the log normalized counts of the top ten clones from each NK cell sample from the Wu dataset. Overlaid asterisks indicate which clone is one of the top ten contributing clones for each sample, and clones are ordered on the y-axis based on hierarchical clustering of their Euclidean distances between their log normalized counts across samples. (**C**) Heat map depicting the same log normalized count values as in (**B**) but with overlaid asterisks instead indicating which clones significantly changed in proportion from the previous sample based on a p-value of < 0.05 assessed by a chi-squared test of proportions. *m, months post-transplant; NK 16, CD3-CD14-CD20-CD56-CD16+ NK cells.*

Figure 5: Lineage bias

Ridge plot shows clonal bias between Gr and T lineages at multiple timepoints of the Six dataset (**A**) and the Koelle dataset (**C**). Ridges indicate the density of clones at the value of log-bias on the x-axes, and dots indicate individual clones, sized by their overall abundance. Multiple ridge plots along the y-axes correspond to the time point of each sample in months post-transplant. Chord diagram showing shared clonality between Gr, T, and Mo lineages from Six et al (**B**) and Koelle et al (**D**) datasets. Each uniquely colored chord represents a unique combination of lineages, and the width of each chord

as it intersects with a lineage indicates the proportional contribution of that group of

clones to that lineage. The space around the perimeter without a chord indicates the

percentage contribution of unilineage clones. *Gr, Granulocyte; Mo, Monocyte.*

Figure S1: Cumulative clone counts from the Six, Belderbos, and Espinoza

datasets

The cumulative count of unique clones detected in each lineage across longitudinal

samples from the Six (**A**), Belderbos (**B**), and Espinoza (**C**) datasets. X-axes show

months, weeks or days post-transplant with "sac" corresponding to the timepoint of

euthanasia in the Belderbos dataset. *Gr, Granulocyte; Mo, Monocyte; bulk, unsorted*

population; CD34p, CD34 positive cell; nRBC, nucleated Red Blood Cell

Figure S2: Systematic analysis of lineage bias in the Six dataset

Ridge plots comparing the lineage bias at multiple time points between all pairwise

combinations of lineages from the Six dataset not shown in Fig. 5 (**A-C, E-G, I-K**).

Ridges indicate the density of clones at the value of log-bias on the x-axes, and dots

indicate individual clones, sized by their overall abundance. Multiple ridge plots along

the y-axes correspond to the time point of each sample in months post-transplant.

Chord diagram showing shared clonality between Gr, B, and Mo lineages (**D**), Gr, NK,

and Mo lineages (**H)**, and T, B, and NK lineages (**L**) from Six et al. Each uniquely

colored chord represents a unique combination of lineages, and the width of each chord

as it intersects with a lineage indicates the proportional contribution of that group of

clones to that lineage. The space around the perimeter without a chord indicates the

percentage contribution of unilineage clones. *Gr, Granulocyte; Mo, Monocyte; NK,*

Natural Killer cell.

Figure S3: Systematic analysis of lineage bias in the Koelle dataset

Ridge plots comparing the lineage bias at multiple time points between Gr and Mo

lineages (**A**) and Mo and T lineages (**B**) from the Koelle dataset. Comparison of the Gr

and T lineage is shown in Fig. 5. Ridges indicate the density of clones at the value of

log-bias on the x-axes, and dots indicate individual clones, sized by their overall

abundance. Multiple ridge plots along the y-axes correspond to the time point of each

sample in months post-transplant. *Gr, Granulocyte; Mo, Monocyte.*

Figure S4: Clonal diversity from TCR sequencing data

The number of T cell clones detected at each timepoint (**A**) and the Shannon diversity

(**B**) of T cell clones in each sample from the Clarke et al dataset (Table 1). Clone counts

represent the number of unique TCR sequences detected at each point not the

cumulative number of clones. Multiple replicates are shown at the 4, 6, 18, and 24

month time points, as described in Clarke et al(24). Each color represents an

independent replicate. The x-axes represent months post-transplant.

unique genetic labelling and transplantation or adoptive transfer

cell sampling, DNA isolation, and high throughput label recovery by sequencing

data analysis

	S1	S2	...
...			

barcodetrackR

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. S1

Fig. S2

Fig. S3