Spotting genome-wide pigmentation variation in a brown trout admixture context

T. Valette[1*], M. Leitwein [1,4*], J.-M. Lascaux[2], E. Desmarais[1], P. Berrebi[1,3] & B. Guinand[1#]

[1] : UMR ISEM - CNRS IRD UM EPHE – Montpellier, France

[2] : ECOGEA – Muret, France

[3] : GENOME – Recherche et Diagnostic – Saint Just, France

[4] : present address : Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval - Québec, Canada

* : contributed equally

: correspondence: bruno.guinand@umontpellier.fr

ORCID IDs: B. Guinand 0000-0002-6934-1677
M. Leitwein 0000-0001-5172-0970
P. Berrebi 0000-0002-9026-4987

Running title : Admixture and pigmentation in brown trout

ABSTRACT

Variation in body pigmentation attracted fish biologists for a while, but high-throughput

genomic studies investigating its molecular basis remain limited to few species and associated

conservation issues ignored. Using 75,684 SNPs, we explored the genomic basis of

pigmentation pattern variation among individuals of the Atlantic and Mediterranean clades of

the brown trout (*Salmo trutta*), a polytypic species in which Atlantic hatchery individuals are

commonly used to supplement local wild populations. Using redundancy analyses and

genome-wide association studies, a set of 337 independent "colour patterning loci" (CPLs)

significantly associated with pigmentation traits such as the number of red and black spots on

flanks, or the presence of a black spot on the pre-opercular bone was identified. CPLs map

onto 35 out of 40 brown trout linkage groups indicating a polygenic basis to pigmentation

patterns. They are mostly located in coding regions (43.4%) of 223 candidate genes, and

correspond to GO-terms known to be involved in pigmentation (e.g. calcium and ion-binding,

cell adhesion). Annotated candidates include genes with known pigmentation effects (e.g.

SOX10, *PEML, SLC45A2*), but also the Gap-junction Δ2 (*GJD2*) gene already shown

differentially expressed in trout skin. Patterns of admixture were found significantly distinct

when using either the full SNP data set or the set of CPLs, indicating that pigmentation

patterns accessible to practitioners are not a reliable proxy of genome-wide admixture.

Consequences for management are discussed.

Keywords: ddRAD sequencing, admixture mapping, multi-trait phenotype, local ancestry,

drift, salmonid, management

1. INTRODUCTION

The study of pigmentation has a peculiar place in modern biology as colour and its patterning

play important roles in natural, artificial and sexual selection (Cieslak, Reissmann, Hofreiter,

& Ludwig, 2011; Cuthill et al., 2017). The biology and evolution of animal colouration

largely benefited over the past few years in the development of high-throughput sequencing

techniques (San José & Roulin, 2017; Orteu & Jiggins, 2020). While supergenes and large

effect mutations have been shown to support pigmentation patterns in many organisms (Orteu

& Jiggins, 2020), genomic studies also generated a huge literature that underscored their

polygenic basis (e.g. skin and hair colour in human: Crawford et al., 2017; Pavan & Sturm,

2019; structural colour variation or eyespot numbers in butterflies: Brien et al., 2019; Rivera-

Colón, Westermann, van Belleghem, Monteiro, & Papa, 2020). Recent research showed that

pigmentation variation may result from effects propagated during development by loci

belonging to regulatory networks to hub or master genes often sufficient to explain most of

the causal variation with colour expression and/or patterning (e.g. Arnould et al., 2013;

Ordway, Hancuch, Johnson, Wiliams, & Rebeiz, 2014; Ding et al., 2020, Orteu & Jiggins,

2020).

If early genetic and molecular studies were interested in colour and pigmentation

patterns in fish – e.g. sex-linked colour variation (Kottler & Schartl, 2018) – the issue of body

pigmentation gained interest with the establishment of zebrafish and medaka as model species

(Parichy, 2006; Takeda & Shimada, 2010; Singh & Nüsslein-Völlard, 2015; Nüsslein-Völlard

& Singh, 2017) and from comparisons with higher vertebrates (e.g. Kelsh, Harris, Colanesi, &

Erickson, 2009). Pigment cells (i.e. chromatophores; mainly melanophores, iridophores,

leucophores and xanthophores) are distributed in the hypo- and the epidermis in fish, and

mutational or other analyses allowed for increased understanding in the mechanisms of

pigment cell fates during development and their resulting distribution (Kelsh et al., 2004,

2009; Kimura et al., 2014; Eom, Bain, Patterson, Grout, & Parichy, 2015; Nüsslein-Völlard &

Singh, 2017; Parichy & Spiewak, 2015; Singh & Nüsslein-Völlard, 2015; Salis et al., 2018;

Volkening 2020). Studies now extend to other fish species (Maan & Sfec, 2013; Irion &

Nüsslein-Völlard, 2019) and to a large array of eco-evolutionary questions regarding the

involvement of genes or regulatory pathways in fish pigmentation, its epistatic and pleiotropic

nature, its modularity and its control, its role in speciation as well as the impact of whole-

genome duplication that promoted the diversification of pigment cell lineages (Hultman,

Bahary, Zon, & Johnson, 2007; Miller et al., 2007; Braasch, Brunet, Volff, & Schartl, 2009;

Roberts, Ser & Kocher, 2009; Albertson et al., 2014; Santos et al. 2014; Ceinos, Guillot,

Kelsh, Cerdá-Reveter, & Rotlland, 2015; Yong, Peichel, & McKinnon, 2015; Kimura,

Takehana, & Naruse, 2017; Roberts, Moore, & Kocher, 2017; Lorin, Brunet, Laudet, & Volff,

2018; Kratochwil et al., 2018; Nagao et al., 2018; Cal et al., 2019; Lewis et al., 2019; Liang,

Gerwin, Meyer, & Kratochwil, 2020). Studies also engaged fish research in high-throughput

genomic approaches of pigmentation variation (Tripathi et al., 2009; Greenwood et al., 2011;

Malek, Boughman, Dworkin, & Peichel, 2012; O'Quin, Drilea, Conte, & Kocher, 2013;

Henning, Jones, Franchini, & Meyer, 2013; Albertson et al., 2014; Henning, Lee, Franchini,

& Meyer, 2014; Xu et al., 2014; Bian et al., 2016; Zhu et al., 2016; Roberts et al. 2017; Kon

et al., 2020). However, in spite of this impressive research, the genomics of pigmentation

variation remain poorly investigated in fish compared to other phenotypic traits (Peichel &

Marques, 2017). Furthermore, links between pigmentation and conservation genomic issues

remain absent (but see Boulding et al., 2008) while this topic emerges in, e.g., birds, notably

regarding the impacts of admixture (Toews et al., 2016; Hanna, Dumbacher, Bowie,

Henderson, & Wall, 2018; Billerman, Cicero, Bowie, & Carling, 2019).

Salmonids represent a large fish family with a diverse and complex body pigmentation

going from continuous colour to spotty, marbled, blotchy and striped patterns. If whole body

colouration or specific coloured elements have a genetic basis (e.g. Blanc, Poisson, & Vibert,

1982; Skaala & Jørstad, 1988; Blanc, Chevassus, & Krieg, 1994; Blanc, Poisson, & Quillet,

2006; Boulding et al., 2008; Colihueque, 2010; Nilsson et al., 2016), another part is plastic

(Colihueque, 2010; Westley, Stanley, & Fleming, 2013; Jørgensen et al., 2018). Colour

patterns support differences in coping styles (Kittilsen et al., 2009; Brännäs et al., 2016) or in

the physiological adjustments necessary to avoid predation and match with environmental

variability (e.g. Miyamoto, 2016; Jacquin et al., 2017; Zastavniouk, Weir, & Fraser, 2017). In

salmonids, pigmentation and colour are also known to interact with social hierarchies,

influence mate choice and affect fitness (e.g. O'Connor, Metcalfe, & Taylor, 2000; Wedekind,

Jacob, Evanno, Nusslé, & Müller, 2008; Marie-Orléach et al., 2014; Watt, Swanson, Miller,

Chen, & May, 2017, Auld, Noakes, & Banks, 2019). However, the genomic basis of their

pigmentation patterns remains largely unexplored, with limited insights coming from

quantitative trait loci (QTL) (Boulding et al., 2008) or few gene expression studies (Sivka,

Snoj, Palandačič, & Sušnik Bajec, 2013; Djurdjevič, Furmanek, Miyazawa, & Sušnik Bajec,

2019). Furthermore, as pigmentation patterns may reflect admixture in salmonids (Largiadèr

& Scholl, 1996; Mezzera, Largiadèr, & Scholl, 1997; Aparicio, García-Berthou, Araguas,

Martinez, & Garcia-Marin, 2005; Miyazawa, Okamoto, & Kondo, 2010; Kirczuk &

Domagała; 2012, Kocabaş, Kutluyer, & Başçinar, 2018), it appears necessary to improve

knowledge on the genotype-phenotype association supporting observed pigmentation patterns.

Indeed, these patterns may inform on individual origin and use to improve management

knowledge, decisions or policies (Largiadèr & Scholl, 1996; Poteaux & Berrebi, 1997;

Delling, Crivelli, Rubin, & Berrebi, 2000; Aparicio et al., 2005; Kocabaş et al., 2011; Marin,

Coon, & Fraser, 2017; Duchi, 2018; Lorenzoni et al., 2019). It is thus important to evaluate if

loci supporting phenotypic divergence for pigmentation patterning reflect genome-wide

admixture.

In this study, we take advantage of former genomic knowledge established in the

brown trout (Leitwein, Gagnaire, Desmarais, Berrebi, & Guinand, 2018) to search for the loci

responsible for body pigmentation variation among populations of two clades recognised in

this species (Atlantic and Mediterranean; Bernatchez, 2001; Sanz, 2018). Then, we investigate

if those loci could be a relevant proxy to monitor hybridization and genome-wide admixture

among individuals of these clades.

2. MATERIALS AND METHODS

2.1 - BROWN TROUT'S GENOMICS DATA

Double digested restriction site-associated DNA sequencing (ddRADseq) data used in this

study are from Leitwein et al. (2018). The data set consists in 75,684 genome-wide SNPs

(40,519 RAD-loci or haplotypes) for 112 trout of hatchery and wild caught origins. Eighty-

two wild caught individuals were fished in the headwaters of three rivers within the

Mediterranean Orb River catchment (France) (Gravezon, Mare and Upper Orb rivers;

Leitwein et al., 2018). This catchment has been seeded by hatchery fish of both Atlantic and

Mediterranean origins for decades. This stopped in 2004, but unintentional releases due to

flooding of private hatcheries are documented (Leitwein et al., 2018). Analyses showed that

these wild caught individuals consisted in hatchery individuals of both Mediterranean and

Atlantic origins, F1's, F2's, backcrossed individuals, and 'pure' natural Mediterranean fish

(Leitwein et al., 2018). 'Early' and 'late' backcrossed individuals correspond to

Mediterranean wild-caught fish that contained distinct distributions of Atlantic ancestry tracts

in their genome. Thirty individuals of the domestic Atlantic ($N = 15$) and Mediterranean ($N =$

15) strains were also included in the analysis. Mediterranean hatchery fish have been

randomly sampled in a local strain formerly established using mature trout from the Gravezon

River by the Fédération de Pêche de l'Hérault in 2004. Atlantic hatchery fish originated from

the Cauterets hatchery that maintains an Atlantic strain distributed worldwide (Bohling,

Haffray, & Berrebi, 2016).

2.2 - PHENOTYPIC DATA

Acquisition of phenotypic data followed Lascaux (1996) and is used by numerous angler

associations and management authorities in France. The set of variables considered in this

protocol is listed in Table S1 and illustrated on a picture in Fig. S1. Data were recorded from

individual photographs of the 112 fish considered for genomic analyses. Photographs were

taken at fishing on slightly anesthetized trout with eugenol. A camera Canon® EOS 1000D

was used. After recovery, fish were released in the wild or in hatchery tanks. Quantitative

variables ($N = 19$) and semi-quantitative variables ($N = 11$) were recorded by visual

examination of photographs of the left flank of each individual fish. Quantitative variables

refer to punctuation patterns of brown trout (e.g. number and diameter of spots) and are

recognized important for this species (Blanc et al., 1982, 1994). Semi-quantitative variables

refer to 'ornamental' appearance patterns (e.g. parr marks, fringes on fins, white ring around

spots) that are important to practitioners or anglers. Each quantitative punctuation variable

was measured independently. Ornamental variables have been coded by modalities, either

present/absent [1, 2] or, e.g., absent/partial/complete [1, 2, 3] (Tables 1 & S1). This set of

variables avoided focusing only on peculiar attributes of the pigmentation patterns. Other

pigmentation traits that can be also important for trout (e.g. background colour, brightness, or

reflectance; Colihueque, 2010) were not considered in this study. Phenotypic data and trout

pictures will be deposited on a public repository. Eleven colour patterning variables were

found not significantly correlated (Table S2) and retained for subsequent analysis (Table 1).

2.3 - GENOTYPE-PHENOTYPE ASSOCIATION

The evaluation of the genotype-phenotype association between SNPs and pigmentation/colour

traits was performed using three distinct approaches. Hereafter, we coined a 'colour

patterning locus' (CPL) any SNP or RAD-locus found significantly associated with a colour

variable considered in this study. A CPL is not a 'pigmentation locus' (or gene, if any; see

Lorin et al., 2018), but a locus that participate to produce pigmentation and colour

information.

Redundancy analysis (RDA) - We first used RDA to investigate any association between

genomic and colour trait data. RDA is a constrained ordination method in which only the

variation of response variables explained by a set of explanatory variables is displayed and

analysed (Legendre & Legendre, 2012). RDA was performed with the remaining 11

uncorrelated colour variables as the explanatory and the SNPs as the response variables.

Missing genomic data were imputed by the most commonly observed genotype. Missing

phenotypic data were imputed by the mean of observed trait for quantitative data, and the

most commonly observed phenotype for semi-quantitative data. We classified SNPs as

showing statistically significant association with individual pigmentation/ornamental traits

when they loaded with more than 2.5 standard deviations (S.D.) from the mean.

A forward model selection was used to select for the relevant phenotypic variables

structuring the RDA (Blanchet, Legendre, & Borcard, 2008). In constrained ordination

methods like RDA, increasing the number of explanatory variables becomes similar to

unconstrained ordination method (e.g. principal component analysis) as the percentage of

variation explained increases when considering more explanatory variables, while some of

them add no relevant information. Models were defined, first including the eleven

uncorrelated response variables, then reducing this number. The Aikake Information Criterion

was computed in each case to select the most appropriate model (i.e. minimizing deviance).

Permutation tests (N = 999) were performed by permuting individuals in each model. This

procedure was first established for the RDA itself, then for each successive RDA axis to

investigate if observed patterns carried significant association between relevant pigmentation

traits and SNPs. Analyses were performed with the *vegan* package (https://cran.r-

project.org/web/packages/vegan/index.html). Once relevant colour variables were identified

in models, each significant SNP was associated to the pigmentation or ornamental trait it was

the most significantly correlated.

Genome-wide association studies (GWAS) –A single-trait association was first considered by

fitting each colour variable with SNPs by linear regression. We used a penalized maximum-

likelihood least absolute shrinkage and selection operator (LASSO) model to select for SNPs

implied in each trait association, then solving:

$$\min_{(\beta_0, \beta) \in \mathbb{R}^{p+1}} \left[\frac{1}{2N} \sum_{i=1}^{N} (y_i - \beta_0 - x_i\, \beta)^2 + \lambda \sum_{j=1}^{p} |\beta_j| \right]$$

219   (1)

in which $Y \in \mathbb{R}$ represents the response colour variable, $X \in \mathbb{R}^p$ a vector of predictor variables

(i.e. SNPs), λ the penalty parameter, β_0 the y-intercept of multiple linear regression, and $\beta \in$

$\mathbb{R}^p$ a vector of β_j coefficients (Friedman, Hastie, & Tibshirani, 2010). This vector of β_j

coefficients represents the effect size β_j of the j[th] SNP conditional on the effects of all other

SNPs. The penalized term λ shrinks the regression coefficient towards zero, keeping only a

small number of SNPs with large effects in the model. A cyclical coordinate descent

procedure was retained for model selection (Friedman et al., 2010). The retained model was

determined by cross-validation. Log(λ) was estimated by minimizing the mean quadratic error.

The number of positive β_j coefficients was estimated from log(λ), with each β_j coefficient

associated to a suite of SNPs considered as involved in the association. Analyses were

performed with the *glmnet* package (https://cran.r-project.org/package=glmnet).

A multi-trait GWAS was also implemented. We used the MultiPhen package

(O'Reilly et al., 2012) to test for the linear combination of phenotypes most associated with

the genotypes at each SNP. Such one approach may potentially capture effects hidden to

single phenotype GWAS. It performs a 'reversed regression', with multiple phenotype

predictors and genetic variant as outcome (i.e. *G* SNPs $X_{(i)} = \{X_{(i1)},…, X_{(iG)}\}$ are explained by

K pigmentation variables $Y_{(i)} = \{Y_{(i1)},…, Y_{(iK)}\}$). SNPs are encoded by allele count alleles ($X_{(ig)}$

$\in \{0, 1, 2\}$). An ordinal logistic regression was considered to derive the probability than one

SNP is associated to a multi-trait phenotype (Porter & O'Reilly, 2017). Permutation tests were

performed to determine one adjusted significance threshold to detect false positives

(Dudbridge & Gusnanto, 2008). A probability $P < 5 \times 10^{-8}$ was retained to consider one SNP

as significantly implied in a multi-trait pigmentation association.

In single- and multi-trait GWAS, we controlled for population stratification by using

nine distinct trout samples recognized by the length and number of ancestry tracts (Leitwein

et al., 2018). Local ancestry tracts may integrate linkage disequilibrium patterns which are

crucial in association studies (e.g. Shriner, 2017; Li, Kemppainen, Rastas, & Merilä, 2018).

The nine groups include the two hatchery samples (Atlantic, Mediterranean), then seven

distinct samples of wild-caught trout. Following Leitwein et al. (2018), wild-caught trout were

grouped as follows: F1's, F2's, 'early' and 'late' backcrossed individuals, then samples of

'pure' wild individuals assigned to each of the three local populations. As some wild caught

trout were identified as hatchery individuals (Leitwein et al., 2018), they have been grouped

with individuals sampled in the hatchery type they have been previously assigned.

2.4 - MAPPING AND ANNOTATION

CPLs detected by RDA and GWAS have been mapped on the Atlantic salmon genome

(*Salmo salar*; Lien et al. 2016; Genbank assembly: GCA_000233375.4) and on the high

density linkage map of *S. trutta* (Leitwein et al., 2017). We searched for genes located in a

25kb window upstream or downstream of each colour-associated marker (arbitrary range).

When one association with a gene was detected, markers were assigned to coding or non–

coding (intron, upstream or downstream sequences) gene regions, or other specific entities

(transposable elements, pseudogenes). Using *S. salar* annotations as inputs, gene ontologies

(GO) for biological processes and molecular functions were derived for genes associated to

CPLs using QuickGO (https://www.ebi.ac.uk/QuickGO/). When annotations were unavailable

on the salmon genome, a search was launched on UniProtKB

(https://www.uniprot.org/uniprot/) for the protein sequence encoded by the gene. Only

annotations with similarity >90% were retained.

Because many genes associated to melanocyte development are also implied in

melanoma development when mutated (Patton, Mitchell, & Naim, 2010; Uong & Zon, 2010),

we further performed a literature search in order to know if genes detected within the 25kb

windows were previously mentioned in former studies. Queries were made using the gene ID

(or aliases taken from GeneCards, https://www.genecards.org/) and a specific keyword. The

retained keywords were: skin, melanocyte, melanophore, melanosome, melanoma (squamous,

cutaneous and uveal), pigment/-ation, keratinocyte, chromatophore, iridiophore, follicle (hair

or skin), nevus/-i, (epi)dermis, vitiligo, erythema/-tous, and sebocyte. This search was updated

until the beginning of June, 2020, each time using the Web of Science, PubMed and Google

Scholar.

2.5 - GENOMIC DIFFERENTIATION

Distributions of F_{ST} values were established using all SNPs, then for the subset of CPLs found

significant in the RDA and GWAS. Single-locus F_{ST} values were estimated considering five

distinct trout samples: the two samples of hatchery fish, then trout caught in each of the three

rivers within the Orb catchment.

2.6 – COMPARISON OF INDIVIDUAL ASSIGNMENT

In order to compare if individual assignment (probability of membership) ranked individuals

similarly or not using all loci or CPLs only, we first established the probability of

membership of individuals to the Atlantic population using independent CPLs defined in this

study. Individual admixture proportions were computed with the LEA package (Frichot &

François, 2015), using $K = 2$ (Atlantic and Mediterranean clades). We then used the

probability of assignment established by Leitwein et al. (2018) that were based on individuals

used in this study, using the same set of 40,519 RAD-loci, also considering $K = 2$. A

Wilcoxon signed-ranked test between each assignment probability distribution was

performed. Similar rankings of individuals in assignment will indicate that CPLs represent a

proxy of genome-wide admixture, while a significant difference will be indicative of the

reverse.

3. RESULTS

3.1 - REDUNDANCY ANALYSIS

Forward model selection showed that a model based on eight colour variables minimized the

deviance in RDA *(Macrost, N.PR.Tot, Zeb, Fr.An, L.Lat, Fr.P, Diam.PN, N.PN.Tot)* (Table

S3). Hereafter, results are reported for this model. Patterns of variation explained by the RDA

were found to significantly structure the association between SNPs and pigmentation

variables ($P < 0.001$). The first axis of RDA was found to represent 29.81% of the total inertia

(Fig. 1), and was significant either ($P < 0.001$). RDA axis 2 was found to explain only a tiny

fraction of observed variation for pigmentation variables (2.10%; Fig. 1), and be marginally

significant ($P = 0.051$). As RDA is a constrained ordination method, constrained inertia (the

percentage of variation explained by uncorrelated pigmentation variables) was found to

represent 35.50% of the total inertia. The remaining portion is due to other factors. Axis 1

represented approx. 84% (ratio: 29.81/35.50) of constrained inertia, while axis 2 explained

only approx. 6% (2.10/35.50) of inertia.

RDA results are represented as a triplot in which individuals are positioned according

to the relationship established between response and explanatory variables (Fig. 1). The

ornamental trait *Macrost* (hereby a large pre-opercular black stain/spot) and the pigmentation

trait *N.PR.Tot* (total number of red spots) explained 23.6% and 22.7%, respectively, of the

total loading scores of phenotypic variables onto RDA axis 1. Results showed a clear

distinction between colour patterns of Mediterranean and Atlantic hatchery fish (Fig. 1A).

Atlantic hatchery fish were characterized by the total number of red spots, while the

macrostigma spots, but also zebra marks (*Zeb*) and total number of black spots (*N.PN.Tot*)

characterized Mediterranean hatchery fish along RDA axis 1. Results confirmed that few wild

caught individuals from the Gravezon River were released Atlantic hatchery fish (Leitwein et

al., 2018). Other individuals from the Gravezon River and 'late backcrossed' individuals have

patterns more characteristic from hatchery Mediterranean fish (Fig. 1A; the hatchery was

seeded by individuals caught in this river [see Materials and Methods section]). Other wild

caught individuals are distributed between these extremes, with individuals identified as F1's

presenting intermediary positions between wild-caught and wild Mediterranean fish on one

side, and Atlantic fish on the other side (Fig. 1A). The position of the wild caught Orb River

individuals referred to the marginally significant second axis of the RDA and to specific

variables (*L. Lat*, *Fr.P*) that were mostly unobserved in other fish (Fig. 1A). F2 individuals

presenting these characters were effectively fished in the Upper Orb River, suggesting they

possibly inherited colour characters expressed in this local population.

The ordination of SNPs by RDA is further detailed in Fig.1B. A total of 1,130 distinct

loci (1.49% of 75,684 SNPs) were found significantly associated to the first two RDA axes

(i.e. > 2.5 S.D. from the mean). Only 22 loci (0.03%) were significantly associated to each of

the two axes for distinct phenotypic variables. Different numbers of SNPs were found

associated with pigmentation variables: 299 with *Fr.P*, 269 with *N.PR.Tot*, 225 with *N.PN.Tot*,

and 213 with *Macrost*. *Macrost* showed significance for loci associated to both first and

second axis of RDA (yellow circles on Fig.1B). As axis 2 was found marginally significant

and substantially affected by individuals from one single population, only SNPs associated to

the first RDA axis will be considered further. This discarded loci associated to *Fr.P* that

explained most of the variation for the second RDA axis. Three hundred and twenty SNPs

were found associated to the main RDA axis, all of them being associated to *Macrost* and

N.PR.Tot, these variables being negatively correlated (Fig. 1B).

As a complement to this analysis and because the threshold of ± 2.5 S.D. is somewhat

arbitrary, we looked back at specific genes known to be involved in colour patterning in

vertebrates and fish, and searched for their position within the distribution of loci over the

RDA axis 1. We detected eight genes representing 30 SNPs for which sequencing reads were

available (Fig. S2). One SNP from the *Mitf1* (microphthalmia-associated transcription factor

1) gene was close to the threshold considered in this study. The *Mitf1* gene is a master gene

for melanocyte differentiation (e.g. Cheli, Ohanna, Ballotti, & Bertolotto, 2010) and, with a

negative value on the first axis of the RDA, this SNP is associated to *Macrost*, thus the

expression of the black colour.

3.2 - SINGLE AND MULTI-TRAIT GWAS

Only two pigmentation variables allowed for relevant LASSO model construction in single

trait GWAS after selection of the penalized term using the cyclical descent procedure (Fig.

S3): *N.PN.Tot* (total number of black spots: 17 candidate SNPs) and *N.PR.Tot* (total number

of red spots: 9 candidate SNPs) (Table S4). These two variables were found significant in

RDA, and negatively correlated (Fig. 1). Candidate SNPs from single GWAS models are

reported in Fig. 2A.

Two multi-trait GWAS models were retained (Fig. 2). As the single-trait GWAS, the

first one jointly retained *N.PN.Tot* and *N.PR.Tot* as variables for which some SNPs were

found associated. The second selected model considered seven of the eight variables formerly

retained in the RDA (except *Zeb*). Seven and thirty-one SNPs were considered as significant

in these multi-trait GWAS, respectively (Table S4).

Despite some similarities for the pigmentation and colour variables put forward by

the different genotype-phenotype associations, only one single SNP (RAD-locus) was found

in common between RDA and the single-trait GWAS for number of black spots (Fig. 2A). No

other SNP was commonly detected by distinct genotype-phenotype association methods, and

especially no SNP shared among GWAS models. It results in a total of 384 SNPs considered

as CPLs putatively implied in pigmentation variation in brown trout (~0.51% of the total

number of SNPs; 320 coming from RDA and 64 from the single- and multi-trait GWAS

models). These CPLs corresponds to 337 independent RAD-loci (0.83% of the total number

of RAD-loci).

3 .3 - GENOMIC DIFFERENTIATION

The distributions of F_{ST} values for the 40,519 RAD-loci and the 320 CPLs associated to axis 1

of the RDA are reported in Fig. 3. The mean F_{ST} was estimated to $F_{ST} = 0.286$ [95% CI: 0.284,

0.287] for the full set of SNPs, while this estimate was $F_{ST} = 0.575$ [95% CI: 0.570-0.583] for

the 320 CPLs detected with the RDA. Observed mean F_{ST} values were $F_{ST} = 0.176$ (min. 0.02

– max: 0.38) and $F_{ST} = 0.025$ (min.: 0.00; max. 0.20) for the single- and the multi-trait GWAS,

respectively. Min/max values are reported rather than 95% CI because the low number of

SNPs associated to phenotypic variables in each GWAS. As distributions do not overlap, the

mean F_{ST} value of the 384 CPLs has no real meaning, but is $F_{ST} = 0.547$ [95% CI: 0.538 -

0.555].

3.4 - MAPPING ON *S. TRUTTA* LINKAGE GROUPS AND ANNOTATION

We considered only the 337 independent CPLs as linked SNPs provide redundant

information. Three hundred independent CPLs were mapped onto 35 of the forty brown trout

LGs defined in Leitwein et al. (2017) (Fig. 4), while 37 of them could not be adequately

positioned on the high density linkage map of Leitwein et al. (2017). Their position relative to

other RAD-loci suffered ambiguity. The distribution of these 300 CPLs on each LG varied

from zero (LG13, LG14, LG15, LG20, LG39) to 86 (LG31). The 337 CPLs have been

mapped on the Atlantic salmon genome and their annotations are given in Table S4. Among

the 337 independent CPLs, ~75.9% were found located in coding (43.4%) or so-called

"regulatory regions" (32.5%; 17.8% upstream and 14.7% downstream genes) within the 25kb

window retained around genes. This represents 245 loci associated to one gene (223 distinct

genes; Table S4). Approximately 22.0% of the CPLs did not match any gene in the 25kb

window, and ~2% were found associated to pseudogenes or long non-coding (lnc) RNA (Fig.

S4). LncRNAs might be important in fish colour patterning (Luo et al., 2019), but not

considered further. Overall 89.86% (200 out of 223) of the genes in close vicinity of the CPLs

detected in this study have been mentioned in the literature dealing with colour, or affecting

integument patterning, differentiation or structure. Table S4 provides details on gene names,

their position and the location of associated CPL; references mentioning each of these 200

CPLs are reported.

An excerpt of twenty-four of these 200 genes (12.0%) is reported in Table 2 to

summarize data. Among these genes, nine candidate genes were detected using GWAS (four

by single-, and five by multi-trait GWAS) (Table 2). The RAD-locus detected by both RDA

and single-trait GWAS for black spotting is located ~21kb downstream of *GJD2* (Gap-

junction protein Δ2 gene; located on LG9 of *S. trutta*; Table S4). This gene was formerly

found differentially expressed in trout skin (Djurdjevič et al., 2019).

3.5 - GO TERMS

GO terms for molecular functions and biological processes of the 337 CPLs are reported in

Fig. S5. Calcium and metal ion binding were found to be the most representative molecular

functions. GO-terms for biological functions highlighted cellular adhesion processes (Fig.

S5A), signal transduction and translation, and transmembrane transport (Fig. S5B).

3.6 – WILCOXON SIGNED-RANKED TEST ON INDIVIDUAL ASSIGNMENT

As expected because of large differences in mean F_{ST} values, individual rankings in

probability of membership were found to significantly differ when using the whole RAD-loci

established in Leitwein et al. (2018) and the independent CPL data sets defined in this study

(Wilcoxon signed-rank test, $P < 10^{-5}$). As CPLs were mainly associated to red and black spots

(*N.PN.Tot*, *N.PR.Tot, Macrost*) in the GWAS and the RDA, this suggests that such patterns

are not reliable indicators of genome-wide admixture, while frequently, but intuitively used

by managers.

4. DISCUSSION

Because of the importance of visual information in natural and sexual selection, we aimed to

investigate the genomic basis of body colour patterning between trout from the Mediterranean

and Atlantic clades recognized in this species. The second one is heavily used for stocking

practices and has been introduced worldwide, including Mediterranean watersheds (Elliott,

1989; Budy et al., 2013). Based on findings by Leitwein et al. (2018), admixture (ancestry)

patterns were included in our study and improved the detection of significant associations as

shown in previous studies (e.g. Shriver et al., 2003; Daya et al., 2014; Pallares, Harr, Turner,

& Tautz, 2014; Brelsford, Toews, & Irwin, 2017), including fish (Malek et al., 2012). This

information allowed for the detection of CPLs associated to candidate genes formerly

identified in pigmentation studies. These CPLs complement genome-wide gene expression

studies by Djurdjevič et al (2019) and improve knowledge on the genomic basis of

pigmentation in the brown trout. While pigmentation patterns are important to evaluate the

impact of hatchery individuals into wild trout populations (Mezzera et al., 1997; Aparicio et

al., 2005), it is premature to rely on phenotypic colour patterns as a proxy of genome-wide

admixture to promote management decisions.

4.1 - ASSOCIATIONS PROMOTE SPOTTING TRAITS…

Genotype-phenotype associations used in this study principally put forward the numbers of

red and black spots and the presence/absence of the black pre-opercular spot as the main

pigmentation variables that aimed to distinguish Mediterranean and Atlantic trout. This was

expected as these variables have been repeatedly shown to characterize pigmentation

differences between Atlantic or Mediterranean trout in studies previously conducted in France

(Poteaux & Berrebi, 1997) and Spain (Aparicio et al., 2005). These traits were not always

found in Italian populations, and others may prevail locally (e.g. patterns of the adipose fin)

(see Lorenzoni et al., 2019). While focusing on few local French populations, results

presented in this study thus reflect most of the former results obtained for these two clades at

a broader scale. Nevertheless, as for Italian populations, model selection showed that

potentially more variables might be considered at the local scale. Spotting traits do not

uncover the full panel of pigmentation variation present in a polytypic species like brown

trout. Further works participating to establish the genotype-phenotype association for colour

pattern in brown trout should integrate other tools and variables than used in this study.

Recent software developments may participate to improve this issue (Endler, Cole, & Kranz,

2018; Van Bellenghem et al., 2018; Chan, Stevens, & Todd, 2019), including additional data

on hue, reflectance or brightness.

4.2 - … BUT DISTINCT LOCI INVOLVED IN PIGMENTATION VARIATION

If the different genotype-phenotype association methods used in our study retrieved the same

pigmentation variables, CPLs detected by RDA and GWAS appeared clearly distinct with

exception of *GJD2* formerly studied in trout (Djurdjevič et al., 2019). Indeed, the identity of

loci considered associated to a single and/or multi-trait phenotype, their number, and their

respective levels of genomic differentiation did not overlap. This is expected as models

behind methods are different and may direct the analyses towards specific findings. For

example, we already mentioned that the shrinkage procedure in LASSO aimed to keep only

SNPs with large effects while discarding others, also explaining why far less CPLs come from

GWAS. It is also widely known that results from single-trait GWAS lack power when few

individuals or SNPs are considered (e.g. Schielzeth & Husby, 2014), especially when

considering wild populations (Kardos, Husby, McFarlane, Qvanström, & Ellegren, 2018;

Santure & Garant, 2018) and polygenic variation (Berg & Coop, 2014). These drawbacks

were moderated by consideration of population stratification based on ancestry patterns. In the

opposite, GWAS retrieved loci with low and sometimes null F_{ST} values that RDA ignored

while balancing selection is crucial to explain pigmentation variation (e.g. Croucher, Oxford,

Lam, & Gillespie, 2011; Lindtke et al., 2017; Schweizer et al., 2018). We do not expand

further. A sub-section below is devoted to RDA that, despite caveats, remains potentially

neglected in association studies.

We prefer to adopt a pragmatic view: RDA and GWAS detected a large proportion

of markers formerly mentioned in the literature on skin and pigmentation. Over the 337

independent CPLs retained overall, 245 loci mapped within a 25kb window downstream or

upstream 223 distinct genes. Two hundreds of them were formerly mentioned in our literature

survey regarding issues and processes dealing with colour patterning or skin-related issues.

The GO terms highlighted by our study for both molecular functions and biological processes

are crucial to pigmentation patterning. Calcium binding is known to be important for

melanogenesis (Bush & Simon, 2007; Bellono & Oancea, 2014; Jia et al., 2020). For

example, the *SLC45A2* gene detected in this study that encodes for a transporter protein

involved in the production and regulation of melanin is regulated by calcium binding levels

(Ginger et al., 2008; Bellono & Oancea, 2014). Melanin is also known to deeply interact with

metal ions at various steps of melanogenesis (Hong & Simon, 2007). Metal ion binding is

necessary in the integrin-mediated signaling pathways whose importance in colour patterning

is well-known (Kelsh et al., 2009; Klotz et al., 2018). Zinc binding is critical to control

melanocyte migration to the epidermis (Denecker et al., 2014) and microtubule activity for

melanosome trafficking within melanocytes (Aspengren, Hedberg, Sköld, & Wallin, 2009).

Other functions such ATP- or DNA- binding activities participate to melanosome/melanocyte

biology and regulation (Levy et al., 2006; Heimerl, Bosserhof, Langmann, Ecker, & Schmitz,

2007). DNA-binding especially concerns transcription factors that promote growth, survival

and differentiation of melanocytes. The main biological functions highlighted in this study

(cellular adhesion processes, signal transduction and translation, transmembrane transport) are

involved in chromatophore or melanocyte development, interactions among chromatophores

or between melanocytes and other cell types (fibroblasts, keratinocytes) (Yamaguchi,

Brenner, & Hearing, 2007; Raposo & Marks, 2007; Kelsh et al., 2009; Mahalwar, Singh,

Fadeev, Nüsslein-Vollard, & Irion, 2016; Nüsslein-Völlard & Singh, 2017). Cell adhesion

processes are illustrated by the numerous calcium-dependent (proto)cadherin genes found in

this study (*CDH4*, *FAT1*, and *PCDH* genes), as well as integrins (*ITGAV*) (Tables 2 and S4).

Protocadherins (e.g. *PCDH10A*, *PCDHAC2*) are essential to fish skin patterning (e.g.

Williams, Hsu, Rossi, & Artinger, 2018; Du et al., 2019).

Each association method reported classical candidate genes associated to

pigmentation. The RDA reported the above-mentioned (proto)cadherins, but also the *ZIC4* or

SLC7A11 genes. *ZIC4* participates to the dorso-ventral patterning of medaka (Ohtsuka et al.,

2004). The observed trait association with *SLC7A11* – a gene controlling pheomelanin

production in mammals (Chintala et al., 2005) - is another report in fish after carp (*Cyprinus*

carpio; Xu et al., 2014; Jiang et al., 2014) and the 'red' tilapia (Zhu et al., 2016; Wang et al.,

2019). GWAS detected some classical players of pigmentation and body colouration in

vertebrates (e.g. *SOX10, PEML, SLC45A2*), including fish (Schonthaler et al., 2005; Hou,

Arnheiter, & Pavan, 2006; Kelsh et al., 2009; Singh & Nüsslein-Vollard, 2015). Indeed,

SOX10 influences colour patterning during zebrafish development (Elworthy, Lister, Carney,

Raible, & Kelsh, 2003; Greenhill, Rocco, Vibert, Nikaido, & Kelsh, 2011; Nüsslein-Vollard

& Singh, 2017). *SLC45A2* is involved in proton-transport and osmoregulation of

melanosomes resulting in colour dilution as observed in medaka (Fukamachi, Shimada, &

Shima, 2001) and other organisms (e.g. Xu et al., 2013; Adhikari et al., 2019; Sevane, Sanz,

& Dunner, 2019). *PEML* (premelanosome protein) contributes to eumelanin deposition and

colour dilution in tetrapods (Raposo & Marks, 2007; Ishishita et al., 2018), and found to be

involved in colouration of the koi carp (Luo et al., 2019) and a cichlid (Ahi & Sfec, 2017).

Finally, few CPLs are associated to genes involved in background colour adaptation

(*GABRA2*, gamma-aminobutyric acid receptor subunit alpha-2-like; Bertolesi, Vazhappill,

Hehr, & McFarlane, 2016), or light-induced colour change (*MTRR*, methionine synthase

reductase-like; Jones et al., 2018; *FGFR1*, fibroblast growth factor receptor-like 1; Czyz,

2019). *FGF* signaling is essential to the developmental patterning of skin and scales in fish

(Aman, Fulbright, & Parichy, 2018), but also eyespots of butterfly wings (Özsu & Monteiro,

2017). This raises interesting questions regarding the conservation of mechanisms involved in

spot formation in different developmental contexts.

Overall, results established for the brown trout identified both key genes and an array

of far less-known genes involved in biological and molecular processes related to the

regulation of colour and skin pigmentation patterning.

4.3 - A HIGHLY POLYGENIC ARCHITECTURE WITH A ROLE FOR DRIFT

With 337 independent CPLs distributed over 35 LGs and a fraction of them with well-

established relationships in modulating colour and pigmentation patterns, a polygenic

architecture of body colour patterning is satisfied in the case of brown trout. Informative

CPLs participating to genomic architecture of pigmentation in trout are shared in both coding

and regulatory regions. This partitioning is now often reported in the literature (e.g. She &

Jarosz, 2018), but still remains relatively poorly investigated in fish (Peichel & Marques,

2017, but see Jones et al., 2012). Within pigmentation studies, literature emphasizes both the

relative roles of coding (Protas & Patel, 2008; Uy et al., 2016) and regulatory variation to

limit pleiotropic costs during colour pattern establishment and evolution (Larter, Dunbar-

Wallis, Berardi, & Smith, 2018; Toomey et al., 2018). More coding changes than loci

dispersed in regions that might be the basis of regulatory changes were detected in trout, but,

due to the reduced representation of the genome provided by ddRAD sequencing, it is

difficult to appreciate the relevance of this observation. It might be simply due to the choice

of restriction enzymes that returned a peculiar genomic distribution of CPLs. Furthermore, we

provided only a loose definition of upstream or downstream regulatory regions (a 25kb

window) that, undoubtedly, has to be refined in further investigations to better decipher, e.g.,

cis-regulatory processes.

This polygenic architecture contrasts with former studies that supported oligogenic

models or showed the likely presence of few major QTLs in trout and salmonids (Blanc et al.,

1982, 1994; Boulding et al., 2008; Colihueque, 2010), but also in other fish species ($n < 8$

QTLs; Borowsky & Wilkens, 2002; Miller et al., 2007; Magalhaes & Seehausen, 2010;

Greenwood et al., 2011; Malek et al., 2012; O'Quin et al., 2013; Yong et al., 2015). However,

few QTLs are considered not sufficient to explain all causal variation for pigmentation

differences (Greenwood et al., 2011) and comparison with our study should be carefully

considered. First, we did not use fish from controlled crossing experiments and our ability to

detect any QTL is null. Second, we did not focus on a given body part (e.g. fin) or a

pigmentation character (stripe, bars, belly spots) as done in other studies (e.g. Ahi & Sfec,

2017; Roberts et al., 2017), including trout (red spots in Blanc et al., 1994). We explored the

complexity in trout pigmentation in order to detect the most important body pattern-related

variables relevant to our samples. This automatically increased the panel of loci involved in

body colour pigmentation, and, as in this study, when more pigmentation traits were

considered, the number of pigmentation-related QTLs was found far higher and distributed

over more LGs than when studying only one or few traits ($n > 20$; Tripathi et al., 2009;

Albertson et al., 2014).

One increased genomic basis to pigmentation patterns was also reported when drift

was present (Protas, Conrad, Gross, Tabin, & Borowsky, 2007), and drift is common in trout

because of small local effective population sizes (N_e) (e.g. Palm, Laikre, Jorde, & Ryman,

2003; Charlier, Laikre, & Ryman, 2012). This relies on the classical contrast between the

quantitative trait nucleotide (QTNs) and the infinitesimal models of trait variation (e.g.

Rockman, 2012). When stochastic processes are of prime importance, small infinitesimal

segregate in many genomic regions and might be recruited in the mechanisms and

programmes of leading to trait variation (Rockman, 2012). This may translate to a

genomically diffuse information to specific phenotypes at the local population level as

observed in trout. This is illustrated by loci associated to RDA axis 2 that are population

specific (Upper Orb River) and concern peculiar trait associations. The highly polytypic

brown trout may illustrate one intermediate situation, in which some master genes are

detected at the clade level by both GWAS and RDA and are associated to recognised spotting

traits, while 'small players' mostly detected by RDA illustrate patterns specific to local

populations, more locally expressed by other traits *(L. Lat*, *Fr.P)*. If the big picture provided

by master genes should be investigated further, 'small players' cannot be ignored.

4.4 - RDA: A NEGLECTED METHOD TO SCREEN FOR GENOTYPE-PHENOTYPE ASSOCIATIONS

Our findings illustrate the ability to investigate genotype-phenotype associations with a

constrained ordination method like RDA associated to accurate modeling of phenotypic

variables. Jombart, Pontier and Dufour (2009) already mentioned that RDA was neglected in

association studies despite now recognized desirable properties to limit false positives

(Capblanc, Luu, Blum, & Bazin, 2018; Forester et al., 2018) and robustness to recombination

rate variation (Lotterhos, 2019). Effectively, the use of RDA to investigate genotype-

phenotype associations remains seldom (Talbot et al., 2017; Vangestel, Eckert, Wegrzyn, St.

Clair, & Neale, 2018; Carvalho et al., 2020), while it became a standard in genotype-

environment association studies (Forester, Lasky, Wagner, & Urban, 2018), including trout

(Bekkevold et al., 2019). Basically, RDA allowed for distinguishing the part of variation

collectively explained by RAD-loci and independent pigmentation variables (~35% of

observed trait variation). A large portion of observed pigmentation might be noise or be

environment-dependent. Studies in salmon (Jørgensen et al. 2018) and trout (Westley et al.,

2013) showed that plasticity may indeed prevail in explaining pigmentation patterns. While it

discards markers under balancing selection, a beneficial feature of RDA is to possibly focus

on SNPs that may explain for causal variation with at least one phenotypic variable, and not

only using outlier loci with the highest F_{ST} values, but without demonstrated association to a

phenotype. For example, in a study dealing with pigmentation variation, Neethiraj, Hornett,

Hill, and Wheat (2017) used genome scans tools to identify outlier loci, then *a posteriori*

searched for association of these loci with pigmentation. If RDA remains affected by some

false positives as are genome scans, it certainly targets more relevant loci as illustrated by the

high number of CPLs retrieved from the literature in this study. Gautier (2015) proposed a

model based on Bayesian outlier detection model that may use a color variable as a covariate,

but it is limited so far to a single covariate when model-based selection prior to RDA might

allow for considering several ones concurrently. Adding on Jombart et al. (2009), RDA or

related methods (e.g canonical correlation analysis, Porter & O'Reilly, 2017) deserve more

attention in association studies.

4.5 - IMPLICATIONS FOR MANAGEMENT

Contrary to birds (Hanna et al., 2018; Billerman et al., 2019) or mammals (e.g. Anderson et

al., 2009; Fulgione et al., 2016), how differences in body pigmentation are related to

differentiation and admixture was poorly studied in fish and was not related to conservation

and management issues (Malek et al., 2012; Meier et al., 2018, but see Boulding et al., 2008).

Using mitochondrial and allozymic markers, Aparicio et al. (2005) paved the way to use

pigmentation as a proxy of admixture between Atlantic and Mediterranean trout and to

address such issues. Indeed, body pigmentation patterns are often important to anglers and

managers to characterize local trout populations and their "integrity". Earlier studies

addressing this issue using old-generation markers indicated that pigmentation might

represent such a proxy (Mezzera et al., 1997; Aparicio et al., 2005; see Delling et al., 2000 for

S. t. marmoratus), but this issue was not addressed with SNPs. In this study, RDA showed

that 'late backcrossed' individuals clustered very close from the local Mediterranean hatchery

strain and wild individuals of the Gravezon River that were used to seed this strain. This

suggests that the pigmentation pattern of these latter individuals came back close to the local

Mediterranean patterns and that counter selection of Atlantic pigmentation alleles or other

changes (e.g. Endler, Betancourt, Nolte, & Schlötterer, 2016) occurred. However, different

rankings in assignment between this study for CPLs and the full RAD-loci data set (Leitwein

et al., 2018) also suggest that this set of CPLs is not a proxy of genome-wide admixture in this

catchment. Managers could build a policy based on wild-caught individuals that exhibit the

'local pigmentation' whereas trout remain largely admixed with foreign alleles. If this

mismatch between rankings is due to neutral alleles, admixture would hopefully not constrain

the adaptive potential of populations, and pigmentation-based management policies that limit

the genotyping costs be sufficient. On the contrary, if a mismatch occurs and that CPLs do not

mirror admixture at locally-adapted fitness effect loci, using pigmentation as a proxy of local

genetic integrity alone might be at risk despite apparent counter-selection. Distinguishing

between these situations is a difficult task, but, at that time, heavy reliance on the '*local is*

best' paradigm (i.e. considering that individuals presenting the 'local' phenotype are more

adapted to local conditions) might be harmful for population fitness (Broadhurst et al., 2008;

Kronenberger et al., 2018). We are aware that the list of CPLs we propose is a list of

candidate markers that has to be refine for further testing against genome-wide admixture, but,

in the meantime, managers should adopt a precautionary approach when evaluating

population 'integrity' based on body pigmentation.

4.6 - LIMITS AND FUTURE DIRECTIONS

While providing strong results, comparisons using individuals from more natural populations,

clades and/or subspecies within the *S. trutta* complex remain necessary in order to better

investigate and elucidate the evolutionary history of trout regarding pigmentation patterns and

to refine the CPL list. DdRADseq used in this study provides a reduced genome

representation targeting only a tiny portion of the genome that is considered either to

successfully target (Belsford et al., 2017), or miss important loci responsible for pigmentation

patterns (Gauthier et al., 2020). Thus, further RNA sequencing or gene expression studies,

together with more standard association studies in control settings that, as for other traits (e.g.

Sinclair-Waters et al., 2020), may provide estimates of QTL number, variation in locus effect

sizes and more accurately pointing key master genes seem necessary to improve findings.

However, the particular settings of many natural trout populations (low N_e, high genetic drift

and reduced gene flow) have to be taken into account to explain pigmentation diversity

observed in the wild. Pigmentation issues represent one outstanding case to investigate

genotype-phenotype-environment relationships, and increasing knowledge on such

relationships is crucial in salmonid conservation (Waples et al., 2020). As thousands of trout

pictures are taken each year and tissue samples might be easily collected, the development of

conservation phenogenomics programmes involving scientists and practitioners should be

motivated.

ACKNOWLEDGEMENTS

This work benefited from the Montpellier Bioinformatics Biodiversity (MBB) platform from

the LabEx CeMEB, an ANR "Investissements d'Avenir" program (ANR-10-LABX-04-01). J.

Pouzadoux and A. Jourdan are acknowledged for inputs. ML was partly supported by a grant

from LabEx CeMEB.

DATA ACCESSIBILITY

Genomic data are from Leitwein et al. (2018) and available at NCBI Short Read Archive

under the study accession SRP136716. The final VCF and the PLINK files for *S. trutta* LGs

are available under the study accessions SRZ187687 and SRZ187688. Upon acceptance, trout

pictures and phenotypic data will be deposited on Dryad or another repository.

AUTHORS' CONTRIBUTION

TV, ML and BG analyzed data and drafted the manuscript. JML acquired phenotypic data.

ED provided support to molecular data analysis. ML, PB and BG designed the study. All

authors revised drafts and approved the final version.

REFERENCES

Adhikari, K., Mendoza-Revilla, J., Sohail A., Fuentes-Guajardo, M., Lampert, J., Chacon-Duque, J. C., … Ruiz-
Linares, A. (2019). A GWAS in Latin Americans highlights the convergent evolution of lighter skin
pigmentation in Eurasia. *Nature Communications,* 10(1), *358.* doi: 10.1038/s41467-018-08147-0

Ahi, E. P., & Sefc. K. M. (2017). A gene expression study of dorso-ventrally restricted pigment pattern in adult
fins of *Neolamprologus meeli*, an African cichlid species. *PeerJ, 5,* e2843. doi: 10.7717/peerj.2843

Albertson, R. C., Powder, K. E., Hu, Y., Coyle, K. P., Roberts, R. B., & Parsons K. J. (2014). Genetic basis of
continuous variation in the levels and modular inheritance of pigmentation in cichlid fishes. *Molecular
Ecology, 23(21),* 5135-5150. doi: 10.1111/mec.12900

Aman, A. J., Fulbright, A. N., & Parichy D. M. (2018). Wnt/β-catenin regulates an ancient signaling network
during zebrafish scale development. *Elife, 7,* e37001. doi: 10.7554/eLife.37001

Amsen, E. M., Pham, N., Pak, Y., & Rotin, D. (2006). The guanine nucleotide exchange factor CNrasGEF
regulates melanogenesis and cell survival in melanoma cells. *Journal of Biological Chemistry, 281(1),* 121-
128. doi: 10.1074/jbc.M507595200

Anderson, T. M., vonHoldt, B. M., Candille, S. I., Musiani, M., Greco, C., Stahler, D. R., … Barsh, G. S. (2009).
Molecular and evolutionary history of melanism in North American gray wolves. *Science, 323,* 1339-1343.
doi: 10.1126/science.1165448

Aparicio, E., García-Berthou, E., Araguas, R. M., Martinez, P., & Garcia-Marin, J. L. (2005). Body
pigmentation pattern to assess introgression by hatchery stocks in native *Salmo trutta* from Mediterranean
streams. *Journal of Fish Biology, 67(4),* 937-949. doi: 10.1111/j.0022-1112.2005.00794.x

Arnoult, L., Su, K. F., Manoel, D., Minervino, C., Magriña, J., Gompel, J., & Prud'Homme, B. (2013).
Emergence and diversification of fly pigmentation through evolution of a gene regulatory module. *Science,
339,* 1423-1426. doi: 10.1126/science.1233749

Aspengren, S., Hedberg, D., Sköld, H. N., & Wallin, M. (2009). New insights into melanosome transport in
vertebrate pigment cells. *International Review of Cell and Molecular Biology, 272,* 245-302. doi:
10.1016/S1937-6448(08)01606-7

Auld, H.L., Noakes, D.L.G., & Banks, M.A. (2019). Advancing mate choice studies in salmonids. *Reviews in
Fish Biology and Fisheries, 29,* 249–276. doi: 10.1007/s11160-019-09551-5

Beauvais., A, Erickson, C. A., Goins, T., Craig, S. E., Humphries, M. J., Thiery, J-P., & Dufour, S. (1995).
Changes in the fibronectin-specific integrin expression pattern modify the migratory behavior of sarcoma
S180 cells in vitro and in the embryonic environment. *Journal of Cell Biology, 128(4),* 699-713. doi:
10.1083/jcb.128.4.699

Bekkevold, D., Höjesjö, J, Nielsen, E. E., Aldvén, D., Damm Als, T., Sodeland, M., … Hansen, M. M. (2019).
Northern European *Salmo trutta* (L.) populations are genetically divergent across geographical regions and
environmental gradients. *Evolutionary Applications, 13(2),* 400-416. doi: 10.1111/eva.12877

Bellono, N. W., & Oancea, E. V. (2014). Ion transport in pigmentation. *Archives of Biochemistry and Biophysics,
563,* 35-41. doi: 10.1016/j.abb.2014.06.020

Berg, J. J., & Coop, G. 2014. A population genetic signal of polygenic adaptation. *PLoS Genetics, 10(8),*
e1004412. doi: 10.1371/journal.pgen.1004412

Bernatchez, L. (2001). The evolutionary history of brown trout (*Salmo trutta* L.) inferred from phylogeographic,
nested clade, and mismatch analyses of mitochondrial DNA variation. *Evolution,* 55, 351-379. doi:
10.1111/j.0014-3820.2001.tb01300.x

Bertolesi, G. E., Vazhappilly, S. T., Hehr, C. L., & McFarlane, S. (2016). Pharmacological induction of skin
pigmentation unveils the neuroendocrine circuit regulated by light. *Pigment Cell and Melanoma Research,*
29, 186-198. doi: 10.1111/pcmr.12442

Blanc, J-M., Poisson, H., & Vibert R. (1982). Variabilité génétique de la ponctuation noire sur la truitelle Fario
(*Salmo trutta* L.). *Annales de Génétique et de Sélection Animale,* 14, 225-236

Blanc, J-M., Chevassus, B., & Krieg, F. (1994). Inheritance of the number of red spots on the skin of the brown
trout. *Aquatic Living Resources, 7,* 133-136. doi: 10.1051/alr:1994016

Blanc, J-M., Poisson, H., & Quillet, E. (2006). A blue variant in the rainbow trout, *Oncorhynchus mykiss*
Walbaum. *Journal of Heredity, 97,* 89–93. doi: 10.1093/jhered/esj010

Bian, C., Hu, Y., Ravi, V., Kuznetsova , I. S., Shen, X., Mu, X.,… Shi, Q. (2016). The Asian arowana
(*Scleropages formosus*) genome provides new insights into the evolution of an early lineage of teleosts.
Scientific Reports, 6, 24501. doi:10.1038/srep24501

Billerman, S. M., Cicero, C., Bowie, R. C. K., & Carling, M. D. (2019). Phenotypic and genetic introgression
across a moving woodpecker hybrid zone. *Molecular Ecology, 28,* 1692-1708. doi: 10.1111/mec.15043

Blanchet, F. G., Legendre, P., & Borcard, D. (2008). Forward selection of explanatory variables. *Ecology, 89,*
2623-2632. doi: 10.1890/07-0986.1

Bohling, J., Haffray, P., & Berrebi, P. (2016). Genetic diversity and population structure of domestic brown trout
(*Salmo trutta*) in France. *Aquaculture* 462: 1-9. doi: 10.1016/j.aquaculture.2016.04.013

Borowsky, R., & Wilkens, H. 2002. Mapping a cave fish genome: polygenic systems and regressive evolution.
Journal of Heredity, 93(1), 19-21. doi: 10.1093/jhered/93.1.19

Boulding, E. G., Culling, M., Glebe, B., Berg, P.R., Lien, S., & Moen T. (2008). Conservation genomics of
Atlantic salmon: SNPs associated with QTLs for adaptive traits in parr from four trans-Atlantic backcrosses.
Heredity, 101, 381-391. doi: 10.1038/hdy.2008.67

Braasch, I., Brunet, F., Volff, J-N., & Schartl, M. (2009). Pigmentation pathway evolution after whole-genome
duplication in fish. *Genome Biology and Evolution*, *1,* 479-493. doi: 10.1093/gbe/evp050

Brännäs, E., Backström, T., Nilsson, J., Carlberg, H., Stien, L. H., & Magnhagen, C. (2016). Distinguishing
Arctic charr with different stress coping styles by visual screening of spottiness – reliability and consistency
over time. *Journal of Zoology, 300,* 213-220. doi: 10.1111/jzo.12374

Brelsford, A., Toews, D. P. L., & Irwin D. E. (2017). Admixture mapping in a hybrid zone reveals loci
associated with avian feather coloration. *Proceedings: Biological Sciences, 284,* 20171106. doi:
10.1098/rspb.2017.1106

Brien, M. N., Enciso-Romero, J., Parnell, A. J., Salazar, P. A., Morochz, C., Chala, D., … Nadeau, N. J. (2019).
Phenotypic variation in *Heliconius erato* crosses shows that iridescent structural colour is sex-linked and
controlled by multiple genes. *Interface Focus, 9,* 20180047. doi:10.1098/rsfs.2018.0047

Broadhurst, L. M., Lowe, A., Coates, D. J., Cunningham, S. A., McDonald, M., Vesk, P. A., & Yates, C. (2008).
Seed supply for broadscale restoration: maximizing evolutionary potential. *Evolutionary Applications, 1:*
587–597. doi: 10.1111/j.1752-4571.2008.00045.x

Budy, P., Thiede G. P., Lobón-Cerviá, J., Gonzales, G., McHugh, P., McIntosh, A., … Jellyman, P. (2013).
Limitation and facilitation of one of the world's most invasive fish: an intercontinental comparison. *Ecology,*
94, 356–367. doi: 10.2307/23435983

Bush, W. D., & Simon, J. D. (2007). Quantification of Ca^{2+} binding to melanin supports the hypothesis that
melanosomes serve a functional role in regulating calcium homeostasis. *Pigment Cell Research, 20*, 134-139.
doi: 10.1111/j.1600-0749.2007.00362.x

Cal, L., Suarez-Bregua, P., Braasch, I., Irion, U., Kelsh, R., Cerda-Reveter, M., & Rotllant, J. (2019). Loss-of-
function mutations in the melanocortin 1 receptor cause disruption of dorso-ventral countershading in teleost
fish. *Pigment Cell and Melanoma Research, 32,* 817-828. doi: 10.1111/pcmr.12806

Capblancq, T., Luu, K., Blum, M. G. B., & Bazin, E. (2018). Evaluation of redundancy analysis to identify
signatures of local adaptation. *Molecular Ecology Resources, 18,* 1223-1233. doi:10.1111/1755-0998.12906

Carvalho, C. S., Forester, B. R., Mitre, S. K., Alves, A., Imperatriz-Fonseca, V. L., Ramos, S. J., … Jaffé, R.
(2020). Combining genotype, phenotype, and environmental data to delineate site-adjusted provenance
strategies for ecological restoration. *Molecular Ecology Resources* , doi:10.1111/1755-0998.13191

Ceinos, R. M., Guillot, R., Kelsh, R. N., Cerdá-Reverter, J. M., & Rotllant, J. (2015). Pigment patterns in adult
fish result from superimposition of two largely independent pigmentation mechanisms. *Pigment Cell and*
Melanoma Research, 28(2), 196-209. doi: 10.1111/pcmr.12335

Chan, I. Z. W., Stevens, M., & Tood, P. A. (2019). PAT-GEOM: a software package for analysis of animal colour
patterns. *Methods in Ecology and Evolution, 10(4),* 591-600. doi: 10.1111/2041-210X.13131

Charlier, J., Laikre, L., & Ryman, N. (2012). Genetic monitoring reveals temporal stability over 30 years in a
small, lake-resident brown trout population. *Heredity, 109(4),* 246-253. doi:10.1038/hdy.2012.36

Cheli, Y., Ohanna, M., Ballotti, R., & Bertolotto, C. (2010). Fifteen-year quest for microphthalmia-associated
transcription factor target genes. *Pigment Cell and Melanoma Research, 23(1),* 27-40. doi: 10.1111/j.1755-
148X.2009.00653.x

Chintala, S., Li, W., Lamoreux, M. L., Ito, S., Wakamatsu, K., Sviderskaya, E. V., … Swank, R. T. (2005).
Slc7a11 gene controls production of pheomelanin pigment and proliferation of cultured cells. *Proceedings*
of the National Academy of Sciences of the USA, 102(31), 10964-10969. doi:10.1073/pnas.0502856102

Cichorek, M., Wachulska, M., Stasiewicz, A., & Tymińska, A. (2013). Skin melanocytes: biology and
development. *Postepy Dermatology and Alergology, 30(1),* 30-41. doi: 10.5114/pdia.2013.33376

Cieslak, M., Reissmann, M., Hofreiter, M., & Ludwig, A. (2011). Colours of domestication. *Biological Reviews*
of the Cambridge Philosophical Society, 86(4), 885-899. doi: 10.1111/j.1469-185X.2011.00177.x

Colihueque, N. (2010). Genetics of salmonid skin pigmentation: clues and prospects for improving the external
appearance of farmed salmonids. *Reviews in Fish Biology and Fisheries, 20,* 71–86. doi: 10.1007/s11160-
009-9121-6

Crawford, N. G., Kelly, D. E., Hansen M. E. B., Beltrame, M. H. Fan, S., Bowman, S. L., … Tishkoff, S. A.
(2017). Loci associated with skin pigmentation identified in African populations. *Science, 358,* eaan8433.
doi:10.1126/science.aan8433

Croucher, P. J., Oxford, G. S., Lam, A., & Gillespie, R. G. 2011. Stabilizing selection maintains exuberant
colour polymorphism in the spider *Theridion californicum* (Araneae, Theridiidae). *Molecular Ecology,*
20(2), 206-218. doi:10.1111/j.1365-294X.2010.04941.x

Cuthill, I. C., Allen, W. L., Arbuckle, K., Caspers, B., Chaplin, G., Hauber, M. E., … Caro, T. (2017). The
biology of color. *Science, 357,* ean0221. doi:10.1126/science.aan0221

Czyz, M. (2019). Fibroblast growth factor receptor signaling in skin cancers. *Cells, 8(6),* 540. doi:
10.3390/cells8060540

da Silva Romero, A. R., Siquera, F., Santiago, G. G., Correia de Almeida Regitano, L., Dorta de Souza Junior,
M., de Ameida Torres Junior, R. A., … Barufatti Grisolia, A. (2018). Prospecting genes associated with
navel length, coat and scrotal circumference traits in Canchim cattle. *Livestock Science, 210,* 33-38. doi:
10.1016/j.livsci.2018.02.004

Daya, M., van der Merwe, L., Gignoux, C. R., van Helden, P. D., Möller, M., & Hoal, E. G. (2014). Using multi-
way admixture mapping to elucidate TB susceptibility in the South African coloured population. *BMC*
Genomics, 15(1), 1021. doi: 10.1186/1471-2164-15-1021

Delling, B., Crivelli, A. J., Rubin, J-F., & Berrebi, P. (2000). Morphological variation in hybrids between *Salmo*
marmoratus and alien *Salmo* species in the Volarja stream, Soca River basin, Slovenia. *Journal of Fish*
Biology, 57, 1199-1212. doi: 10.1006/jfbi.2000.1383

Denecker, G., Vandamme, N., Akay, O., Koludrovic, D., Taminau, J., Lemaire, K., … Berx, G. (2014).
Identification of a *ZEB2-MITF-ZEB1* transcriptional network that controls melanogenesis and melanoma
progression. *Cell Death and Differentiation, 21(8),* 1250-1261. doi:10.1038/cdd.2014.44

Ding, B., Patterson, E. L., Holalu, S. V., Li, J., Johnson, G. A., Stanley, L. E., … Yuan, Y-W. (2020). Two MYB
proteins in a self-organizing activator-inhibitor system produce spotted pigmentation patterns. *Current*
Biology, 30(5), 802-814.e8. doi:10.1016/j.cub.2019.12.067

Djurdjevič, I., Furmanek, T., Miyazawa, S., & Sušnik Bajec, S. (2019). Comparative transcriptome analysis of
trout skin pigment cells. *BMC Genomics, 20(1),* 359. doi:10.1186/s12864-019-5714-1

Du, J., Chen, X., Wang, J., Chen, H., Yue, W. Lu, G., & Wang, C. (2019). Comparative skin transcriptome of
two Oujiang color common carp (*Cyprinus carpio* var. color) varieties. *Fish Physiology and Biochemistry,*
45(1), 177-185. doi: 10.1007/s10695-018-0551-8

Dudbridge, F., & Gusnanto, A. (2008). Estimation of significance thresholds for genomewide association scans.
Genetic Epidemiology, 32(3), 227-234. doi: 10.1002/gepi.20297

Duchi, A. (2018). Flank spot number and its significance for systematics, taxonomy and conservation of the
near-threatened Mediterranean trout *Salmo cettii*: evidence from a genetically pure population. *Journal of*
Fish Biology, 92, 254-260. doi:10.1111/jfb.13492

Elworthy, S., Lister, J.A., Carney, T. J., Raible, D. W., & Kelsh, R. N. (2003). Transcriptional regulation of *mitfa*
accounts for the *sox10* requirement in zebrafish melanophore development. *Development, 130(12),* 2809-
2818. doi: 10.1242/dev.00461

Elliott, J. M. (1989). Wild brown trout *Salmo trutta*: an important national and international resource.
Freshwater Biology, 21, 1-5. doi: 10.1111/j.1365-2427.1989.tb01343.x

Endler, J. A., Cole, G. L., & Kranz, A. M. (2018). Boundary strength analysis: combining colour pattern
geometry and coloured patch visual properties for use in predicting behaviour and fitness. *Methods in*
Ecology and Evolution, 9, 2334-2448. doi: 10.1111/2041-210X.13073

Endler, L., Betancourt, A. J., Nolte, V., & Schlötterer, C. (2016). Reconciling differences in pool-GWAS
between populations: A case study of female abdominal pigmentation in *Drosophila melanogaster*.
Genetics, *202*, 843–855.

Eom, D. S., Bain, E. J., Patterson, L. B., Grout, M. E., & Parichy, D. M. (2015). Long-distance communication
by specialized cellular projections during pigment pattern development and evolution. *Elife,* 4, e12401.
doi:10.7554/eLife.12401

Forester, B. R., Lasky, J. R., Wagner, H. H., & Urban, D. L. (2018). Comparing methods for detecting
multilocus adaptation with multivariate genotype-environment associations. *Molecular Ecology, 27(9),*
2215-2233. doi: 10.1111/mec.14584

Frichot, E., & Francois, O. (2015). LEA: an R package for Landscape and Ecological Association studies.
Methods in Ecology and Evolution, 6(8), 925-929. doi: 10.1111/2041-210X.12382

Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization paths for generalized linear dodels via
coordinate descent. *Journal of Statistical Softwares, 33(1),* 1-22. PMID: 20808728

Fukamachi, S., Shimada, A., & Shima, A. (2001). Mutations in the gene encoding B, a novel transporter protein,
reduce melanin content in medaka. *Nature Genetics, 28(4),* 381-385. doi: 10.1038/ng584

Fulgione, D., Rippa, D., Buglione, M., Trapanese, M., Petrelli, S., & Maselli, V. (2016). Unexpected but
welcome. Artificially selected traits may increase fitness in wild boar. *Evolutionary Applications, 9(6),* 769-
776. doi: 10.1111/eva.12383

Gallagher, S. J., Rambow, F., Kumasaka, M., Champeval, D., Bellacosa, A., Delmas, M., & Larue, L. (2013).
Beta-catenin inhibits melanocyte migration but induces melanoma metastasis. *Oncogene, 32(17),* 2230-2238.
doi: 10.1038/onc.2012.229

Gautier, M. (2015). Genome-wide scan for adaptive divergence and association with population-specific
covariates. *Genetics, 201(4),* 1555-1579. doi: 10.1534/genetics.115.181453

Gauthier, J., de Silva, D. L., Gompert, Z., Whibley, A., Houssin, C., Le Poul, Y., ... Elias, M. 2020. Contrasting
genomic and phenotypic outcomes of hybridization between pairs of mimetic butterfly taxa across a suture
zone. *Molecular Ecology, 29(7),* 1328-1343. doi: 10.1111/mec.15403

Ginger, R. S., Askew, S. E., Ogborne, R. M., Wilson, S., Fernadino, D., Dadd, T., … Green, M. R. (2008).
SLC24A5 encodes a trans-Golgi network protein with potassium-dependent sodium-calcium exchange
activity that regulates human epidermal melanogenesis. *Journal of Biological Chemistry, 283(9),* 5486-5495.
doi:10.1074/jbc.M707521200

Gramann, A. K., Venkatesan, A. M., Guerin, M., & Ceol, C. J. (2019). Regulation of zebrafish melanocyte development by ligand-dependent BMP signaling. *Elife*, *8,* e50047. doi: 10.7554/eLife.50047

Greenhill, E. R., Rocco, A., Vibert, L., Nikaido, M., & Kelsh, R. N. (2011). An iterative genetic and dynamical modelling approach identifies novel features of the gene regulatory network underlying melanocyte development. *PLoS Genetics, 7(9),* e1002265. doi: 10.1371/journal.pgen.1002265

Greenwood, A. K., Jones, F. C., Chan, Y. F., Brady, S. D., Absher, D. M., Grimwood, J., … Peichel, C. L. (2011). The genetic basis of divergent pigment patterns in juvenile threespine sticklebacks. *Heredity, 107(2),* 155-166. doi: 10.1038/hdy.2011.1

Hanna, Z. R., Dumbacher, J. P., Bowie, R. C. K., Henderson, J. B., & Wall, J. D. (2018). Whole-genome analysis of introgression between the spotted owl and barred owl (*Strix occidentalis* and *Strix varia*, respectively; Aves: Strigidae) in western North America. *G3, 8(12),* 3945-3952. doi: 10.1534/g3.118.200754

Heimerl, S., Bosserhoff, A. K., Langmann, T., Ecker, J., & Schmitz, G. (2007). Mapping ATP-binding cassette transporter gene expression profiles in melanocytes and melanoma cells. *Melanoma Research, 17(5),* 265-273. doi: 10.1097/CMR.0b013e3282a7e0b9

Hellström, A. R., Watt, B., Fard, S. S., Tenza, D., Mannström, P., Narfström, K., ... Andersson, L. (2011). Inactivation of *Pmel* alters melanosome shape but has only a subtle effect on visible pigmentation. *PLoS Genetics, 7(9),* e1002285. doi: 10.1371/journal.pgen.1002285

Henning, F., Jones, J. C., Franchini, P., & Meyer, A. (2013). Transcriptomics of morphological color change in polychromatic Midas cichlids. *BMC Genomics*, *14,* 171. doi: 10.1186/1471-2164-14-171

Henning, F., Lee, H. J., Franchini, P., & Meyer, A. (2014). Genetic mapping of horizontal stripes in Lake Victoria cichlid fishes: benefits and pitfalls of using RAD markers for dense linkage mapping. *Molecular Ecology, 23(21),* 5224-5240. doi: 10.1111/mec.12860

Hirobe, T. (2011). How are proliferation and differentiation of melanocytes regulated? *Pigment Cell and Melanoma Research, 24(3),* 462-478. doi: 10.1111/j.1755-148X.2011.00845.x

Hong, L., & Simon, J. D. (2007). Current understanding of the binding sites, capacity, affinity, and biological significance of metals in melanin. *Journal of Physical Chemistry, B111(28),* 7938-7947. doi: 10.1021/jp071439h

Hou, L., Arnheiter, H., & Pavan, W. J. (2006). Interspecies difference in the regulation of melanocyte development by *SOX10* and *MITF*. *Proceedings of the national Academy of Sciences of the USA, 103(24),* 9081-9085. doi: 10.1073/pnas.0603114103

Huang, J. M., & Hornyak, T. J. (2015). Polycomb group proteins - epigenetic repressors with emerging roles in melanocytes and melanoma. *Pigment Cell and Melanoma Research, 28(3),* 330-339. doi: 10.1111/pcmr.12341

Hultman, K. A., Bahary, N., Zon, L. I., & Johnson, S. L. (2007). Gene duplication of the zebrafish *kit ligand* and partitioning of melanocyte development functions to *kit ligand a*. *PLoS Genetics, 3(1),* e17. doi: 10.1371/journal.pgen.0030017

Irion, U., & Nüsslein-Volhard, C. (2019). The identification of genes involved in the evolution of color patterns in fish. *Current Opinion in Genetics and Development, 57,* 31-38. doi: 10.1016/j.gde.2019.07.002

Ishishita, S., Takahashi, M., Yamaguchi, K., Kinoshita, K., Nakano, M., Nunome, M., … Matsuda, Y. (2018). Nonsense mutation in *PMEL* is associated with yellowish plumage colour phenotype in Japanese quail. *Scientific Reports, 8(1),* 16732. doi: 10.1038/s41598-018-34827-4

Jacquin, L., Gauthey, Z., Roussille, V., Le Hénaff, M., Tentelier, C., & Labonne, J. (2017). Melanin in a changing world: brown trout coloration reflects alternative reproductive strategies in variable environments. *Behavioral Ecology, 28(6),* 1423–1434. doi: 10.1093/beheco/arx102

Jia, Q., Hu, S., Jiao, D., Li, X., Qi, S., & Fan, R. (2020). Synaptotagmin-4 promotes dendrite extension and melanogenesis in alpaca melanocytes by regulating Ca^{2+} influx via *TRPM1* channels. *Cell Biochemistry and Functions, 38(3),* 275-282. doi: 10.1002/cbf.3465

Jiang., Y, Zhang, S., Xu, J., Feng, J. Mahboob, S., Al-Ghanim, K.A.,… Xu, P. (2014). Comparative transcriptome analysis reveals the genetic basis of skin color variation in common carp. *PLoS One, 9(9),* e108200. doi: 10.1371/journal.pone.0108200

Jin, Y., Andersen, G., Yorgov, D., Ferrara, T. M., Ben, S., Brownson, K. M., … Spritz, R. A. (2016). Genome-
wide association studies of autoimmune vitiligo identify 23 new risk loci and highlight key pathways and
regulatory variants. *Nature Genetics, 48(11),* 1418-1424. doi: 10.1038/ng.3680

Jombart, T., Pontier, D., & Dufour, A. B. (2009). Genetic markers in the playground of multivariate analysis.
Heredity, 102(4), 330-341. doi: 10.1038/hdy.2008.130

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., … Kingsley, D. M. (2012). The
genomic basis of adaptive evolution in threespine sticklebacks. *Nature, 484,* 55-61. doi:
10.1038/nature10944

Jones, P., Lucock, M., Veysey, M., Jablonski, N., Chaplin, G., & Beckett, E. (2018). Frequency of folate-related
polymorphisms varies by skin pigmentation. *American Journal of Human Biology, 30(2),*
10.1002/ajhb.23079. doi: 10.1002/ajhb.23079

Jørgensen, K. M., Solberg, M. F., Besnier, F., Thorsen, A., Fjelldal, P. J., Skaala, Ø., ..., Glover, K. A. (2018).
Judging a salmon by its spots: environmental variation is the primary determinant of spot patterns in *Salmo
salar*. *BMC Ecology, 18(1),* 14. doi: 10.1186/s12898-018-0170-3

Kang, X., Liu, G., Liu, Y., Xu, Q., Zhang, M., & Fang, M. (2013). Transcriptome profile at different
physiological stages reveals potential mode for curly fleece in Chinese tan sheep. *PLoS One, 8(8),* e71763.
doi: 10.1371/journal.pone.0071763

Kardos, M., Husby, A., McFarlane, S. E., Qvarnström, A., & Ellegren, H. (2016). Whole-genome resequencing
of extreme phenotypes in collared flycatchers highlights the difficulty of detecting quantitative trait loci in
natural populations. *Molecular Ecology Resources, 16(3),* 727-741. doi: 10.1111/1755-0998.12498

Kelsh, R. N., Inoue, C., Momoi, A., Kondo, H., Furutani-Seiki, M., Ozato, K., & Wakamatsu, Y. (2004). The
Tomita collection of medaka pigmentation mutants as a resource for understanding neural crest cell
development. *Mechanisms of Development, 121(7-8),* 841-859. doi: 10.1016/j.mod.2004.01.004

Kelsh, R. N., Harris, M. L., Colanesi, S., & Erickson, C. A. (2009). Stripes and belly-spots -- a review of
pigment cell morphogenesis in vertebrates. *Seminars in Cell and Developmental Biology, 20(1),* 90-104.
doi: 10.1016/j.semcdb.2008.10.001

Kido, K., Sumimoto, H., Asada, S., Okada, S. M., Yaguchi, T., Kawamura, N., … Kaxakami, Y. (2009).
Simultaneous suppression of MITF and BRAF^{V600E} enhanced inhibition of melanoma cell proliferation.
Cancer Science, 100(10), 1863-1869. doi: 10.1111/j.1349-7006.2009.01266.x

Kimura, T., Nagao, Y., Hashimoto, H., Yamamoto-Shiraishi, Y., Yamamoto, S., Yabe, T., … Naruse, K. (2014).
Leucophores are similar to xanthophores in their specification and differentiation processes in
medaka. *Proceedings of National Academy of Sciences of the USA, 111(20),* 7343-7348. doi:
10.1073/pnas.1311254111

Kimura, T., Takehana, Y., & Naruse, K. (2017). *pnp4a* is the causal gene of the medaka iridophore
mutant *guanineless*. *G3* 7(4): 1357-1363. doi:10.1534/g3.117.040675

Kirczuk, L., & Domagała, J. (2012). Colouration of reciprocal hybrids of salmon (*Salmo salar* L., 1758) and sea
trout (*Salmo trutta m.* trutta L, 1758) aged 0+ and 1+ in the natural conditions. *Electronic Journal of Polish
Agricultural University, 15(1),* 02. available online: http://www.ejpau.media.pl/volume15/issue1/art-02.html

Kittilsen, S., Schjolden, J., Beitnes-Johansen, I., Shaw, J. C., Pottinger, T. G., Sørensen, C., … Øverli Ø. (2009).
Melanin-based skin spots reflect stress responsiveness in salmonid fish. *Hormones and Behavior, 56,* 292–
298. doi: 10.1016/j.yhbeh.2009.06.006

Klotz, B., Kneitz, S., Regensburger, M., Hahn, L., Dannemann, M., Kelso, J., … Schartl, M. (2018). Expression
signatures of early-stage and advanced medaka melanomas. *Comparative Biochemistry and Physiology C:
Toxicology and Pharmacology, 208,* 20-28. doi: 10.1016/j.cbpc.2017.11.005

Kocabaş, M., Kayim, M., Can, E., Ateş, M., Kutluyer F., & Aksu, Ö. (2011). Spotting pattern features in the
brown trout (*Salmo trutta macrostigma,* T., 1954) population. *Scientific Research and Essays, 6(23),* 5021-
5024.

Kocabaş, M., Kutluyer F., & Başçinar N. (2018). Phenotypic differentiation analysis: A case study in hybridizing
Çoruh trout (*Salmo coruhensis*), Rize trout (*Salmo rizeensis*) and brown trout (*Salmo trutta fario*). *Acta
Zoologica, 99(3),* 274-280. doi: 10.1111/azo.12211

Kon, T., Omori, Y., Fukuta, K., Wada, H., Watanabe, M., Chen, Z., … Furukawa, T. (2020). The genetic basis
of morphological diversity in domesticated goldfish. *Current Biology, 30(12),* 2260-2274.e6.
doi:10.1016/j.cub.2020.04.034

Kottler, V. A., & Schartl, M. (2018). The colorful sex chromosomes of teleost fish. *Genes, 9(5),* 233. doi:
10.3390/genes9050233

Kratochwil, C. F., Liang, Y., Gerwin, J., Woltering, J. M., Urban, S., Henning, F., … Meyer, A. (2018). Agouti-
related peptide 2 facilitates convergent evolution of stripe patterns across cichlid fish radiations. *Science,*
362, 457-460. doi: 10.1126/science.aao6809

Kronenberger, J. A., Gerberich, J. C., Fitzpatrick, S. W., Broder, E. D., Angeloni, L. M., & Funk, W. C. (2018).
An experimental test of alternative population augmentation scenarios. *Conservation Biology, 32(4),* 838-
848. doi: 10.1111/cobi.13076

Lamason, R. L., Mohideen, M. A., Mest, J. R., Wong, A. C., Norton, H. R., Aros, M. C., … Cheng, K. C. (2005).
SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science, 310,* 1782-
1786. doi: 10.1126/science.1116238

Largiadèr, C. R., & Scholl, A. (1996). Genetic introgression between native and introduced brown trout *Salmo trutta*
L. populations in the Rhône River Basin. *Molecular Ecology, 5,* 417–426. doi: 10.1046/j.1365-
294X.1996.00099.x

Larter, M., Dunbar-Wallis, A., Berardi, A .E., & Smith, S. D. (2018). Convergent evolution at the pathway level:
predictable regulatory changes during flower color transitions. *Molecular Biology and Evolution, 35(9),*
2159-2169. doi: 10.1093/molbev/msy117

Lascaux, J-M. (1996). *Analyse de la variabilité morphologique de la truite commune (Salmo trutta L.) dans les*
cours d'eau du bassin pyrénéen méditerranéen (Unpublished doctoral dissertation). Ecole Nationale
Supérieure Agronomique de Toulouse, France.

Lee, J. L., & Streuli, C. H. (2014). Integrins and epithelial cell polarity. *Journal of Cell Science, 127(Pt 15),*
3217-3225. doi: 10.1242/jcs.146142

Legendre, P., & Legendre, L. (2012). *Numerical Ecology, 3rd Edition*. Amsterdam: Elsevier.

Leitwein, M., Guinand, B., Pouzadoux, J., Desmarais, E., Berrebi, P., Gagnaire, P-A. (2017). A dense brown
trout (*Salmo trutta*) linkage map reveals recent chromosomal rearrangements in the *Salmo* genus and the
impact of selection on linked neutral diversity. *G3, 7(4),* 1365-1376. doi:10.1534/g3.116.038497

Leitwein, M., Gagnaire, P-A., Desmarais, E., Berrebi, P., & Guinand B. (2018). Genomic consequences of a
recent three-way admixture in supplemented wild brown trout populations revealed by local ancestry tracts.
Molecular Ecology 27(17): 3466-3483. doi: 10.1111/mec.14816

Levy, C., Khaled, M., & Fisher, D. E. (2006). *MITF*: master regulator of melanocyte development and
melanoma oncogene. *Trends in Molecular Medicine, 12(9),* 406-414. doi: 10.1016/j.molmed.2006.07.008

Lewis, V. M., Saunders, L. M., Larson, T. A., Bain, E. J., Sturiale, S. L., Gur, D., … Parichy, D. M.. (2019). Fate
plasticity and reprogramming in genetically distinct populations of *Danio* leucophores. *Proceedings of the*
National Academy of Sciences of the USA, 116(24), 11806-11811. doi: 10.1073/pnas.1901021116

Li, Z., Kemppainen, P., Rastas, P., & Merilä, J. (2018). Linkage disequilibrium clustering-based approach for
association mapping with tightly linked genomewide data. *Molecular Ecology Resources, 18(4),* 809-824.
doi: 10.1111/1755-0998.12893

Liang, Y., Gerwin, J., Meyer, A., & Kratochwil, C. F. (2020). Developmental and cellular basis of vertical bar
color patterns in the East African cichlid fish *Haplochromis latifasciatus. Frontiers in Cell and*
Developmental Biology, 8, 62. doi: 10.3389/fcell.2020.00062

Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Kent, M. P., Nome, T., ... Davidson, W. S. (2016). The
Atlantic salmon genome provides insights into rediploidization. *Nature, 533,* 200-205. doi:
10.1038/nature17164

Lindtke, D., Lucek, K., Soria-Carrasco, V., Villoutreix, R., Farkas, T. E., Riesch, R., … Nosil, P. (2017). Long-
term balancing selection on chromosomal variants associated with crypsis in a stick insect. *Molecular*
*Ecol*ogy, *26(22),* 6189-6205. doi:10.1111/mec.14280

Lorenzoni, M., Carosi, A., Giovannotti, M., La Porta, G., Splendiani, A., & Caputo Barucchi, V. (2019). Morphological survey as powerful detection tool of pure and local phenotypes in *Salmo trutta* complex. *Knowledge and Management of Aquatic Ecosystems, 420,* 48. doi: 10.1051/kmae/2019041

Lorin, T., Brunet, F. G., Laudet, V., & Volff, J-N. (2018). Teleost fish-specific preferential retention of pigmentation gene-containing families after whole genome duplications in vertebrates. *G3, 8(5),* 1795-1806. doi: 10.1534/g3.118.200201

Lotterhos, K. E. (2019). The effect of neutral recombination variation on genome scans for selection. *G3, 9(6),* 1851-1867. doi: 10.1534/g3.119.400088

Luo, M., Wang, L., Yin, H., Zhu, W., Fu, J., & Dong, Z. (2019). Integrated analysis of long non-coding RNA and mRNA expression in different colored skin of koi carp. *BMC Genomics, 20(1),* 515. doi: 10.1186/s12864-019-5894-8

Maan, M. E., & Sefc, K. M. (2013). Colour variation in cichlid fish: developmental mechanisms, selective pressures and evolutionary consequences. *Seminars in Cell and Developmental Biology, 24(6-7),* 516-528. doi: 10.1016/j.semcdb.2013.05.003

Magalhaes, I. S., & Seehausen, O. (2010). Genetics of male nuptial colour divergence between sympatric sister species of a Lake Victoria cichlid fish. *Journal of Evolutionary Biology, 23(5),* 914-924. doi: 10.1111/j.1420-9101.2010.01960.x

Mahalwar, P., Walderich, B., Singh, A. P., & Nüsslein-Volhard, C. (2014). Local reorganization of xanthophores fine-tunes and colors the striped pattern of zebrafish. *Science, 345,* 1362-1364. doi: 10.1126/science.1254837

Mahalwar, P., Singh, A. P., Fadeev, A., Nüsslein-Volhard, C., & Irion, U. (2016). Heterotypic interactions regulate cell shape and density during color pattern formation in zebrafish. *Biology Open, 5(11),* 1680-1690. doi: 10.1242/bio.022251

Malek, T. B., Boughman, J. W., Dworkin, I., & Peichel, C. L. (2012). Admixture mapping of male nuptial colour and body shape in a recently formed hybrid population of threespine stickleback. *Molecular Ecology, 21(21),* 5265-5279. doi: 10.1111/j.1365-294X.2012.05660.x

Marie-Orléach, L., Roussel, J-M., Bugeon, J., Tremblay, J., Ombredane, D., & Evanno, G. (2014). Melanin-based coloration of sneaker male Atlantic salmon is linked to viability and emergence timing of their offspring. *Biological Journal of the Linnean Society, 111(1),* 126-135. doi: 10.1111/bij.12187

Marin, K., Coon, A., & Fraser, D. J. (2017). Traditional ecological knowledge reveals the extent of sympatric lake trout diversity and habitat preferences. *Ecology and Society, 22(2),* 20. doi:10.5751/ES-09345-220220

Meier, J. I., Marques, D. A., Wagner, C. E., Excoffier, L., Seehausen, O. (2018). Genomics of parallel ecological speciation in Lake Victoria cichlids. *Molecular Biology and Evolution, 35(6),* 1489-1506. doi: 10.1093/molbev/msy051

Mezzera, M., Largiadèr, C. R., & Scholl A. (1997). Discrimination of native and introduced brown trout in the River Doubs (Rhône drainage) by number and shape of parr marks. *Journal of Fish Biology, 50,* 672-677. doi: 10.1111/j.1095-8649.1997.tb01959.x

Miller, C. T., Beleza, S., Pollen, A. A., Schluter, D., Kittles, R. A., Shriver, M. D., & Kingsley, D. M. (2007). *Cis*-regulatory changes in *Kit ligand* expression and parallel evolution of pigmentation in sticklebacks and humans. *Cell, 131(6),* 1179-1189. doi: 10.1016/j.cell.2007.10.055

Miyamoto, K. (2016). Effects of body color luminance and behavioral characteristics on predation risk in salmonid fishes. *Hydrobiologia, 783,* 249–256. doi: 10.1007/s10750-015-2573-x

Miyazawa, S., Okamoto, M., & Kondo, S. (2010). Blending of animal colour patterns by hybridization. *Nature Communications, 1(6),* 66. doi: 10.1038/ncomms1071

Nagao, Y., Takada, H., Miyadai, M., Adachi, T., Seki, R., Kamei, Y., … Hashimoyo, H. (2018). Distinct interactions of *Sox5* and *Sox10* in fate specification of pigment cells in medaka and zebrafish. *PLoS Genetics, 14(4),* e1007260. doi: 10.1371/journal.pgen.1007260

Nazari-Ghadikolaei, A., Mehrabani-Yeganeh, H., Miarei-Aashtiani, S.R., Staiger, E.A., Rashidi, A., & Huson H. J. (2018). Genome-wide association studies identify candidate genes for coat color and mohair traits in the Iranian Markhoz goat. *Frontiers in Genetics, 9,* 105. doi: 10.3389/fgene.2018.00105

Neethiraj, R., Hornett, E.A., Hill, J.A., & Wheat, C.W. (2017). Investigating the genomic basis of discrete
phenotypes using a Pool-Seq-only approach: new insights into the genetics underlying colour variation in
diverse taxa. *Molecular Ecology*, *26(19),* 4990-5002. doi: 10.1111/mec.14205

Nilsson, J., Backström, T., Stien, L. H., Carlberg, H. Jeuthe, H., Magnhagen, C., & Brännäs, E. (2016). Effects
of age and rearing environment on genetic parameters of growth and body weight and heritability of skin
pigmentation in Arctic charr (*Salvelinus alpinus* L.). *Aquaculture, 453,* 67-72. doi:
10.1016/j.aquaculture.2015.11.035

Nüsslein-Volhard, C., & Singh, A. P. (2017). How fish color their skin: A paradigm for development and
evolution of adult patterns: multipotency, plasticity, and cell competition regulate proliferation and
spreading of pigment cells in zebrafish coloration. *Bioessays, 39(3),* 10.1002/bies.201600231.
doi:10.1002/bies.201600231

O'Connor, K. I., Metcalfe, N. B., & Taylor, A. C. (2000). Familiarity influences body darkening in territorial
disputes between juvenile salmon. *Animal Behaviour, 59(6),* 1095-1101. doi: 10.1006/anbe.2000.1401

Ohtsuka, M., Kikuchi, N., Yokoi, H., Kinoshita, M., Wakamatsu, Y., Ozato, K., … Kimura, M. (2004). Possible
roles of *zic1* and *zic4*, identified within the medaka Double anal fin (Da) locus, in dorsoventral patterning of
the trunk-tail region (related to phenotypes of the Da mutant). *Mechanisms of Development, 121(7-8),* 873-
882. doi: 10.1016/j.mod.2004.04.006

Ordway, A.J., Hancuch, K. N., Johnson, W., Wiliams, T. M., & Rebeiz, M. (2014). The expansion of body
coloration involves coordinated evolution in *cis* and *trans* within the pigmentation regulatory network of
Drosophila prostipennis. *Developmental Biology, 392,* 431-440. doi: 10.1016/j.ydbio.2014.05.023

O'Quin, C. T., Drilea, A. C., Conte, M. A., & Kocher T. D. (2013). Mapping of pigmentation QTL on an
anchored genome assembly of the cichlid fish, *Metriaclima zebra*. *BMC Genomics, 14,* 287. doi:
10.1186/1471-2164-14-287

O'Reilly, P. F., Hoggart, C. J., Pomyen, Y., Calboli, F. C. F., Elliott, P., Jarvelin, M-R., & Coin, L. J. M. (2012).
MultiPhen: joint model of multiple phenotypes can increase discovery in GWAS. *PLoS One, 7(5),* e34861.
doi: 10.1371/journal.pone.0034861

Orteu, A., & Jiggins, C. D. (2020). The genomics of coloration provides insights into adaptive evolution. *Nature
Reviews Genetics, 21,* 461-475. doi:10.1038/s41576-020-0234-z

Özsu, N., & Monteiro, A. (2017). Wound healing, calcium signaling, and other novel pathways are associated
with the formation of butterfly eyespots. *BMC Genomics, 18(1),* 788. doi: 10.1186/s12864-017-4175-7

Pallares, L. F., Harr, B., Turner, L. M., & Tautz, D. (2014). Use of a natural hybrid zone for genomewide
association mapping of craniofacial traits in the house mouse. *Molecular Ecology, 23(23),* 5756-5770. doi:
10.1111/mec.12968

Palm, S., Laikre, L., Jorde, P.E., & Ryman, N. (2003). Effective population size and temporal genetic change in
resident brown trout (*Salmo trutta* L.). *Conservation Genetics, 4(3),* 249-264. doi:
10.1023/A:102406913094

Parichy, D. M. 2006. Evolution of danio pigment pattern development. *Heredity, 97(3),* 200-210. doi:
10.1038/sj.hdy.6800867

Parichy, D. M., & Spiewak, J. E. (2015). Origins of adult pigmentation: diversity in pigment stem cell lineages
and implications for pattern evolution. *Pigment Cell and Melanoma Research, 28(1),* 31-50. doi:
10.1111/pcmr.12332

Patton, E.E., Mitchell, D.L., & Nairn, R.S. (2010). Genetic and environmental melanoma models in fish. *Pigment
Cell and Melanoma Research, 23(3),* 314-337. doi: 10.1111/j.1755-148X.2010.00693.x

Pavan, W.J., & Sturm, R.A. (2019). The genetics of human skin and hair pigmentation. *Annual Review in
Genomics and Human Genetics, 20,* 41-72. doi: 10.1146/annurev-genom-083118-015230

Peichel, C. L., & Marques, D. A. (2017). The genetic and molecular architecture of phenotypic diversity in
sticklebacks. *Philosophical Transactions of the Royal Society of London B: Biological Sciences, 372(1713),*
20150486. doi: 10.1098/rstb.2015.0486

Porter, H. F., & O'Reilly, P. F. (2017). Multivariate simulation framework reveals performance of multi-trait
GWAS methods. *Scientific Reports, 7,* 38837. doi: 10.1038/srep38837

Poteaux, C., Berrebi, P. (1997). Intégrité génomique et repeuplements chez la truite commune du bassin méditerranéen. *Bulletin Français de la Pêche et de la Pisciculture, 344/345,* 309–322. doi : 10.1051/kmae:1997031

Protas, M., Conrad, M., Gross, J. B., Tabin, C., & Borowsky, R. (2007). Regressive evolution in the Mexican cave tetra, *Astyanax mexicanus. Current Biology, 17(5),* 452-454. doi: 10.1016/j.cub.2007.01.05

Protas, M. E., Patel, N. H. (2008). Evolution of coloration patterns. *Annual Review in Cell and Developmental Biology, 24,* 425-446. doi: 10.1146/annurev.cellbio.110707.175302

Rammah, C., Gadin, S., Cormier, S., Souilhol, C., Cohen-Tannoudji, M., Panthier, J-J., & Aubin-Houzelstein, G. (2009). Implication of notchless and strawberry notch homolog 2 genes in melanocyte stem cells homeostasis. *Pigment Cell and Melanoma Research, 22,* 671. doi: 10.1111/j.1755-148X.2009.00612.x 652 a

Raposo, G., & Marks, M. S. (2007). Melanosomes - dark organelles enlighten endosomal membrane transport. *Nature Reviews in Molecular and Cell Biology, 8(10),* 786-797. doi: 10.1038/nrm2258

Rivera-Colón, A. G., Westerman, E. L., Van Belleghem, S. M., Monteiro, A., & Papa, R. (2020). Multiple loci control eyespot number variation on the hindwings of *Bicyclus anynana* butterflies. *Genetics, 214(4),* 1059-1078. doi: 10.1534/genetics.120.303059

Roberts, R. B., Moore, E. C., & Kocher, T. D. (2017). An allelic series at *pax7a* is associated with colour polymorphism diversity in Lake Malawi cichlid fish. *Molecular Ecology, 26(10),* 2625-2639. doi: 10.1111/mec.13975

Roberts, R. B., Ser, J. R., & Kocher, T. D. (2009). Sexual conflict resolved by invasion of a novel sex determiner in Lake Malawi cichlid fishes. *Science, 326,* 998-1001. doi: 10.1126/science.1174705

Rockman, M. V. (2012). The QTN program and the alleles that matter for evolution: all that's gold does not glitter. *Evolution. 66(1),* 1-17. doi:10.1111/j.1558-5646.2011.01486.x

Salis, P., Roux, N., Soulat, O., Lecchini, D., Laudet, V., & Frédérich, B. (2018). Ontogenetic and phylogenetic simplification during white stripe evolution in clownfishes. *BMC Biology, 16(1),* 90. doi:10.1186/s12915-018-0559-7

San-José, L. M., & Roulin, A. (2017). Genomics of coloration in natural animal populations. *Philosophical Transactions of the Royal Society of London B: Biological Sciences, 372(1724),* 20160337. doi: 10.1098/rstb.2016.0337

Santos, M. E., Braasch, I., Boileau, N., Meyer, B. S., Sauteur, L., Böhne, A., … Salzburger,W. (2014). The evolution of cichlid fish egg-spots is linked with a *cis*-regulatory change. *Nature Communications, 5,* 5149. doi: 10.1038/ncomms6149

Santure, A. W., & Garant, D. (2018). Wild GWAS-association mapping in natural populations. *Molecular Ecology Resources, 18(4),* 729-738. doi: 10.1111/1755-0998.12901

Sanz, N. (2018). Phylogeographic history of brown trout: a review. In: J. Lobón-Cerviá & N. Sanz (Eds.), *Brown Trout: Biology, Ecology and Management*. Pp. 17-63. Oxford: Wiley Ltd.

Schielzeth, H., & Husby, A. (2014). Challenges and prospects in genome-wide quantitative trait loci mapping of standing genetic variation in natural populations. *Annals of the New York Academy of Sciences, 1320,* 35-57. doi: 10.1111/nyas.12397

Schonthaler, H. B., Lampert, J. M., von Lintig, J., Schwarz, H., Geisler, R., & Neuhauss, S. C. (2005). A mutation in the silver gene leads to defects in melanosome biogenesis and alterations in the visual system in the zebrafish mutant fading vision. *Developmental Biology, 284(2),* 421-436. doi: 10.1016/j.ydbio.2005.06.001

Schweizer, R.M., Durvasula, A., Smith, J., Vohr, S.H., Stahler, D.R, Galaverni, M., … Wayne, R.K. (2018). Natural selection and origin of a melanistic allele in North American gray wolves. *Molecular Biology and Evolution, 35(5),* 1190-1209. doi:10.1093/molbev/msy031

Sevane, N., Sanz, C.R., & Dunner, S. (2019). Explicit evidence for a missense mutation in exon 4 of *SLC45A2* gene causing the pearl coat dilution in horses. *Animal Genetics, 50(3),* 275-278. doi: 10.1111/age.12784

She, R., & Jarosz, D.F. (2018). Mapping causal variants with single-nucleotide resolution reveals biochemical drivers of phenotypic change. *Cell, 172(3),* 478-490.e15. doi: 10.1016/j.cell.2017.12.015

Schouwey, K., & Beermann, F. (2008). The Notch pathway: hair graying and pigment cell homeostasis.
Histology and Histopathology, 23(5), 609-619. doi: 10.14670/HH-23.609

Shriner, D. (2017). Overview of admixture mapping. *Current Protocols in Human Genetics, 94,* 1.23.1-1.23.8.
doi: 10.1002/cphg.44

Shriver, M. D., Parra, E.J., Dios, S., Bonilla, C., Norton, H., Jovel, C., ... Kittles, R. A. (2003). Skin
pigmentation, biogeographical ancestry and admixture mapping. *Human Genetics, 112(4),* 387-399. doi:
10.1007/s00439-002-0896-y

Sinclair-Waters, M., Ødegård, J., Korsvoll, S. A., Moen, T., Lien, S., Primmer, C. R., & Barson, N. J. (2020).
Beyond large-effect loci: large-scale GWAS reveals a mixed large-effect and polygenic architecture for age
at maturity of Atlantic salmon. *Genetics Selection Evolution, 52(1),* 9 doi: 10.1186/s12711-020-0529-8

Singh, A. P., & Nüsslein-Volhard, C. (2015). Zebrafish stripes as a model for vertebrate colour pattern formation.
Current Biology, 25(2), R81-R92. doi: 10.1016/j.cub.2014.11.013

Singh, S. K., Abbas, W. A., & Tobin, D. J. (2012). Bone morphogenetic proteins differentially regulate
pigmentation in human skin cells. *Journal of Cell Science, 125(18),* 4306-4319. doi10.1242/jcs.102038

Sivka, U., Snoj, A., Palandačić, A., & Sušnik Bajec, S. (2013). Identification of candidate genes involved in
marble color pattern formation in genus *Salmo. Comparative Biochemistry and Physiology D: Genomics
and Proteomics, 8(3),* 244-249. doi: 10.1016/j.cbd.2013.06.003

Skaala, Ø., & Jørstad, K. E. (1988). Inheritance of the fine-spotted pigmentation pattern of brown trout. *Polskie
Archiwum Hydrobiologii, 35,* 295–304.

Soong, J., & Scott, G. (2013). *Plexin B1* inhibits MET through direct association and regulates *Shp2* expression
in melanocytes. *Journal of Cell Science, 126(Pt 2),* 688-695. doi: 10.1242/jcs.119487

Soong, J., Chen, Y., Shustef, E. M., & Scott, G. A. (2012). *Sema4D*, the ligand for *Plexin B1*, suppresses c-Met
activation and migration and promotes melanocyte survival and growth. *Journal of Investigative
Dermatology, 132(4),* 1230-1238. doi: 10.1038/jid.2011.414

Talbot, B., Chen, T. W., Zimmerman, S., Joost, S., Eckert, E. J., Crow, T. M., … Manel, S. (2017). Combining
genotype, phenotype, and environment to infer potential candidate genes. *Journal of Heredity, 108(2),* 207-
216. doi: 10.1093/jhered/esw077

Takeda, H., & Shimada, A. (2010). The art of medaka genetics and genomics: what makes them so unique?
Annual Review of Genetics, 44, 217-241. doi: 10.1146/annurev-genet-051710-151001

Toews, D. P. L., Taylor, S. A., Vallender, R., Brelsford, A., Butcher, B. G., Messer, P. W., & Lovette I. J. (2016).
Plumage genes and little else distinguish the genomes of hybridizing warblers. *Current Biology, 26,* 2313-
2318. doi: 10.1016/j.cub.2016.06.034

Toomey, M. B., Marques, C. I., Andrade, P., Araujo, P. M., Sabatino, S., Gazda, M. A., … Carneiro, M. (2018).
A non-coding region near *Follistatin* controls head colour polymorphism in the Gouldian finch.
Proceedings: Biological Sciences, 285(1888), 20181788. doi: 10.1098/rspb.2018.1788

Tripathi, N., Hoffmann, M., Willing, E. M., Lanz, C., Weigel, D., & Dreyer, C. (2009). Genetic linkage map of
the guppy, *Poecilia reticulata*, and quantitative trait loci analysis of male size and colour variation.
Proceedings: Biological Sciences, 276(1665), 2195-2208. doi: 10.1098/rspb.2008.1930

Uong, A., & Zon, L. I. (2010). Melanocytes in development and cancer. *Journal of Cell Physiology, 222(1),* 38-
41. doi: 10.1002/jcp.21935

Uy, J. A., Cooper, E. A., Cutie, S., Concannon, M. R., Poelstra, J. W., Moyle R. G., & Filardi, C. E. (2016).
Mutations in different pigmentation genes are associated with parallel melanism in island flycatchers.
Proceedings: Biological Sciences, 283(1834), 20160731. doi: 10.1098/rspb.2016.0731

Van Belleghem, S.M., Papa, R., Ortiz-Zuazaga, H., Hendrickx, K., Jiggins, C.D., McMillan, W.O., &
Counterman, B.A. (2018). patternize: An R package for quantifying colour pattern variation. *Methods in
Ecology and Evolution* 9(2): 390-398. doi: 10.1111/2041-210X.12853

Vangestel, C., Eckert, A. J., Wegrzyn, J. L., St.Clair, J. B., & Neale, D. B. (2018). Linking phenotype, genotype
and environment to unravel genetic components underlying cold hardiness in coastal Douglas-fir
(*Pseudotsuga menziesii* var. *menziesii*). *Tree Genetics and Genomes, 14,* 10. doi: 10.1007/s11295-017-1225-
x

Volkening, A. (2020). Linking genotype, cell behavior, and phenotype: multidisciplinary perspectives with a
basis in zebrafish patterns. *Current Opinion in Genetics and Development, 63,* 78-85.
doi:10.1016/j.gde.2020.05.010

Wade, N., Bryant, N.J., Connolly, L. M., Simpson R. J., Luzio, J. P., … James, D. E.. (2001). *Syntaxin 7*
complexes with mouse *Vps10p tail interactor 1b, syntaxin 6, vesicle-associated membrane protein (VAMP)8,*
and *VAMP7* in b16 melanoma cells. *Journal of Biological Chemistry, 276(23),* 19820-19827. doi:
10.1074/jbc.M010838200

Wang, L. M., Bu, H. Y., Song, F. B., Zhu, W. B., Fu, J. J., & Dong, Z. J. (2019). Characterization and functional
analysis of *slc7a11* gene, involved in skin color differentiation in the red tilapia. *Comparative Biochemistry*
and Physiology A: Molecular and Integrative Physiology, 236, 110529. doi: 10.1016/j.cbpa.2019.110529

Wang, Z., Jinnin, M., Kobayashi, Y., Kudo, H., Inoue, K., Nakayama, W., … Ihn, H. (2015). Mice
overexpressing integrin *αv* in fibroblasts exhibit dermal thinning of the skin. *Journal of Dermatological*
Science, 79(3), 268-278. doi: 10.1016/j.jdermsci.2015.06.008

Waples, R. S., Naish, K. A., & Primmer, C. R. (2020). Conservation and management of salmon in the age of
genomics. *Annual Review of Animal Biosciences, 8,* 117-143. doi: 10.1146/annurev-animal-021419-083617

Watt, C., Swanson, C., Miller, D., Chen, M. & May, C. (2017). Social hierarchies override environmental
conditions in determining body coloration of brook trout. *Journal of Freshwater Ecology, 32(1),* 575-580.
doi: 10.1080/02705060.2017.1360218

Wedekind, C., Jacob, A., Evanno, G., Nusslé, S, & Müller, R. (2008). Viability of brown trout embryos
positively linked to melanin-based but negatively to carotenoid-based colours of their fathers. *Proceedings:*
Biological Sciences, 275(1644), 1737-1744. doi: 10.1098/rspb.2008.0072

Westley, P. A., Stanley, R., & Fleming, I. A. (2013). Experimental tests for heritable morphological color
plasticity in non-native brown trout (*Salmo trutta*) populations. *PLoS One, 8(11),* e80401. doi:
10.1371/journal.pone.0080401

Williams, J. S., Hsu, J. Y., Rossi, C. C., & Artinger, K. B. (2018). Requirement of zebrafish *pcdh10a* and
pcdh10b in melanocyte precursor migration. *Developmental Biology, 444(Suppl. 1),* S274-S286. doi:
10.1016/j.ydbio.2018.03.022

Xu, P., Zhang, X., Wang, X., Li, J., Liu, G., Kuang, Y., … Sun, X. (2014). Genome sequence and genetic
diversity of the common carp, *Cyprinus carpio. Nature Genetics, 46(11),* 1212-1219. doi: 10.1038/ng.3098

Xu, X., Dong, G. X., Hu, X.S., Miao, L. Zhang, X. L., Zhang D. L., … Luo, S. J. (2013). The genetic basis of
white tigers. *Current Biology, 23(11),* 1031-1035. doi: 10.1016/j.cub.2013.04.054

Yamaguchi, Y., Brenner, M., & Hearing, V. J. (2007). The regulation of skin pigmentation. *Journal of*
Biological Chemistry, 282(38), 27557-27561. doi: 10.1074/jbc.R700026200

Yong, L., Peichel, C. L., & McKinnon, J. S. (2015). Genetic architecture of conspicuous red ornaments in female
threespine stickleback. *G3, 6(3),* 579-588. doi: 10.1534/g3.115.024505

Zastavniouk, C., Weir, L. K., & Fraser, D. J. (2017). The evolutionary consequences of habitat fragmentation:
body morphology and coloration differentiation among brook trout populations of varying size. *Ecology and*
Evolution, 7(17), 6850-6862. doi: 10.1002/ece3.3229

Zhu, W., Wang, L., Dong, Z., Chen, X., Song, F., Liu, N., … Fu, J. (2016). Comparative transcriptome analysis
identifies candidate genes related to skin color differentiation in red tilapia. *Scientific Reports, 6,* 31347. doi:
10.1038/srep31347

1275

1276   **Captions of the Tables**

1277

1278   **Table 1:** Uncorrelated pigmentation variables retained to build the forward selection model in

1279   RDA, and also used as a basis to multi-GWAS modelling. The full list of variables is reported

1280   in Table S1; correlation among variables in Table S2.

1281

Table 2: List of the most indicative pigmentation- or colour-related genes associated to some

CPLs ($N = 24$; 30 distinct loci) found in this study. The method of association that detected

them is reported, together with their locations in the Atlantic salmon (*Salmo salar*) genome

(Lien et al., 2016) and brown trout (*S. trutta*) high density linkage map of Leitwein et al.

(2017). Their position in coding sequence or in a 25kb window downstream or upstream the

gene is reported. Significant GO-terms are reported. For each gene, relevant references with

link to colour and/or skin features and pigmentation are provided. The full list of CPLs and

other relevant information is given in Table S4.

Captions of the Figures

1291

Fig. 1: RDA triplots for significant or nearly significant canonical axes 1 and 2, respectively.

Pigmentation-related variables retained by forward modelling (Blanchet et al., 2008) are

represented by arrows; description of the retained model in Table S4. Length of arrows is

proportional to the strength of correlation of each variable with individual axis. Arrows

pointing to different direction indicate negatively correlated variables (*e.g. N.PR.Tot* and the

large pre-opercular black stain *Macrost* for RDA axis 1). The percentage of variance

associated to each axis is reported, as well as the eigenvalue graph for constrained axis. **(A):**

Individual trout are positioned on the map with positioning of SNPs as a grey block to the

center of the factor map. Individuals have been coloured according to the nine clusters defined

by Leitwein, Gagnaire, Desmarais, Berrebi, & Guinand (2018) (top left); Clusters reflect the

origin (hatchery *vs* wild) of trout, but also the degree of admixture of each of them as

measured by co-ancestry tracts. **(B):** Zoom on the centre of the map to illustrate the position

of SNPs. SNPs departing ± 2.5 standard deviates from the mean either on RDA axis 1 or RDA

axis 2 have been coloured; a colour being associated to each pigmentation-related variable

included in the model (insert). These SNPs are coloured with the variables they have been

found the most significantly associated in the model. SNPs given as grey spots are within the

± 2.5 S.D. interval and not considered in this study. Percentages of variation explained by

each RDA axis are reported. Fish are representatives of hatchery Mediterranean (left) and

Atlantic (right) individuals used in this study.

1311

Figure 2: Manhattan plots of multi-trait GWAS models selected using the MultiPhen package

(O'Reilly et al., 2012), plus results associated to the single-trait GWAS model used in this

study. The abscissa axis represents the forty brown trout LGs (Leitwein et al., 2017), the

ordinate axis reports values of $-log_{10}(P)$, with P being the probability of one association

between a variable and a SNP (adjusted significance threshold based on permutation,

Dudbridge & Gusnanto, 2008). The red line indicates the 5×10^{-8} significance threshold

retained in this study. **A:** Results for the multi-trait GWAS for the model using two variables:

N. PN.Tot and *N.PR.Tot*. The seven SNPs found significant in this model are represented by

circles. The results from single trait GWAS have been added to this panel to provide a full

summary of results. In this case, approximate position of the RAD-loci associated to the *N.*

PN.Tot (17 loci) and *N.PR.Tot* (9 loci) variables are indicated by black and red arrows,

respectively. One asterisk indicates two loci to close of each other to be indicated by separate

arrows. The locus associated to the *GJD2* gene on LG9 and also detected by the RDA is

indicated by a longer arrow. **B:** Multi-trait GWAS based on the variables retained with the

RDA except *Zeb* (31 candidate SNPs). The significance with each variable of the model is

indicated by a different symbol. Each vertical line of symbols points to a single SNP. Some

overlapping is possible. The full list of candidate SNPs detected using single and multi-trait

GWAS are reported in Table S4. Variables labelled as in Table 1.

1330

Figure 3: Distribution of F_{ST} values for the 40,519 RAD-loci (blue) and for CPLs found

significantly associated with RDA axis 1 ($N = 320$ loci, reported in red in the insert). For each

data set, mean F_{ST} values and confidence intervals are provided as full and hatched lines,

respectively. Mean values for the single- and multi-trait GWAS approaches are illustrated by

the green and the orange lines, respectively. No confidence intervals are reported for GWAS

because of the low number of CPLs detected with these approaches. Details in the main text.

1337

Figure 4: Positioning of the 300 mapped CPLs detected in this study on the forty LGs of the

brown trout. Loci found associated at least one single phenotypic trait are indicted by green,

red and yellow circles when detected with the RDA, single- or multi-GWAS analyses,

respectively. Horizontal bars indicate the distribution of the full set of SNPs over LGs. The

number of CPLs detected on each LG is indicated. As numerous loci are close to each other,

circles may overlap.

Table 1

1345

Phenotypic variables	Description	Nature* (coding)
N. PR. Tot	Total number of red spots over flanks	Q
N. PN. Tot.	Total number of black spots over flanks	Q
Diam. PN	Mean diameter of black spots	Q
Oc. PN	White ring around black spots	SQ (1= no ring; 2 = attenuated/thin ring, 3 = marked/thick ring)
Fr. D	Fringe on the dorsal fin	SQ (1= no fringe; 2 = white fringe, 3 = black & white fringe)
Fr. An	Fringe on the anal fin	SQ (1= no fringe; 2 = white fringe, 3 = black & white fringe)
Fr. P	Fringe on the pelvic fins	SQ (1= no fringe; 2 = white fringe, 3 = black & white fringe)
L. Lat	Visibility of the lateral line	SQ (1 = not visible; 2 = partially visible; 3 = fully visible)
Parr	Presence of parr marks	SQ (1 = absence; 2 = presence)
Macrost.	Macrostigma spot (large black dot on the pre-opercular bone)	SQ (1 = absence; 2 = partially visible/diluted; 3 = fully visible)
Zeb	Zebras on flank	SQ (1 = absence; 2 = presence)

1346 * : Q = quantitative variable; SQ = semi-quantitative variable (coding in brackets)

1347

1348 **Table 2**

1349

Method of Association	LG - S. trutta	LG - S. salar	Position on S. salar genome (pb)	Position of the 'color-associated' locus (coding vs upstream/downstream) and distance to the nearest gene (in bp).	Protein Name (or lncRNA or pseudogene)	Gene Id	GO terms - Molecular Functions	GO terms - Biological processes	Associated reference(s)	Note
RDA	1	29	16654071	Downstream 1942 bp	notchless protein homolog 1	NLE1			Schouwey and Beermann (2008) Rammah et al (2009)	Notch signalling is essential in some pigmentation and epidermal differentiation (e.g.)
Single_GWAS_N.PR.Tot	1	29	31985710	coding	methionine synthase reductase-like	MTRR	FMN binding, oxidoreductase activity	oxidation reduction process	Jones et al. (2018)	Folate-related genes (UV radiation sensitive) associated to skin pigmentation
MultiGWAS_7	1	29	34327580	coding	netrin-G1-like	NTNG1			Kang et al. (2013) Silva Romero et al. (2018) da	Coat color variation
MultiGWAS_2	4	20	17551935	Downstream 7871 bp	P2Y purinoceptor 2-like	P2RY2	signal transducer activity, G-protein coupled receptor activity	signal transduction, G-protein coupled receptor signaling pathway	Ganesan et al. (2008) Jin et al. (2014)	Impact melanogenesis
MultiGWAS_7	6	12	15186741	Downstream 1234 bp	transcription factor SOX-10	SOX10	DNA binding	Upstream 4864 pb de UPF0103 protein EVG1	Nüsslein-Vollard and Singh (2017) Mahalwar et al (2014) Zhu et al. (2016)	Essential to pigmentation patterning in fish
Single_GWAS_N.PN.Tot	9	1	17000287	Upstream 24905 bp	absent in melanoma 1 protein-like	SLC45A2			Bellono & Oancea (2014) Zhu et al. (2010) Sevane et al. (2019)	Ion transport in melanosomes
-	9	1	17132468	coding	-	SLC45A2	-	-	-	-
RDA	9	1	29826721	Upstream 12812 bp	Bone morphogenetic protein 4	BMP4	growth factor activity	regulation of signaling receptor activity	Singh et al. (2012) Gramann et al. (2019)	Melanin synthesis, melanocyte development (zebrafish)
RDA and Single_GWAS_N.PN.Tot	9	1	30355164	Upstream 21540 bp	Gap junction delta-2 protein	GJD2		action potential, synaptic transmission	Djurdjevic et al. (2019)	Differentially expressed in trout skin
RDA	10	13	98943518	Upstream 14366 bp	semaphorin-4D	SEMA4D			Soong et al. (2012)	Melanocyte survival and growth
RDA	11	5	35618944	coding	zinc finger protein ZIC 4	ZIC4	nucleic acid binding, DNA binding, metal ion binding	regulation of transcription by RNA polymerase II	Ohtsuka et al. (2004)	Dorso-ventral patterning in medaka
RDA	11	5	41442470	coding	X-linked interleukin-1 receptor accessory protein-like 2	IL1RAPL1		signal transduction	Jin et al. (2016)	Role in propagation of vitiligo and hypopigmentation
RDA	16	1	114314991	Upstream 2269 bp	vesicle-associated membrane protein 8-like	VAMP8		vesicle-mediated transport	Wade et al. (2001)	Regulate fusion events within the late endosomal pathway, critical role in melanogenesis
MultiGWAS_2	17	15	78637963	coding	E3 ubiquitin-protein ligase Itchy-like	ITCH	ubiquitin-protein transferase activity, transferase activity, ligase activity, ubiquitin protein ligase activity	ubiquitin-dependent protein catabolic process, protein ubiquitination	Raposo & Marks (2007) Nazari-Ghadikolaei et al. (2019)	Melanocyte-based coat colour variation
Single_GWAS_N.PR.Tot	17	15	87874821	Upstream 11038 bp	melanocyte protein PMEL-like	PMEL			Schonthaler et al. (2005) Raposo & Marks (2007) Hellström et al. (2011) Luo et al. (2019)	Color variation (silvering/ dilution), including zebrafish and koi carp
RDA	22	4	34326380	coding	pro-neuregulin-2 membrane-bound isoform-like	NRG2			Kido et al. (2009) Gallagher et al. (2013)	Regulated by MITF1 (a master pigmentation gene) and role in the migration of melanosome
RDA	24	21	53809138	Downstream 1115 bp	integrin alpha-V	ITGAV		cell adhesion, integrin-mediated signaling pathway	Beauvais et al. (1995) Wang et al. (2015) Lee and Streuli (2014)	Cell migratory behavior from neural crest cells from which melanophore originate, also dermal thickness and epithelial cell polarity
MultiGWAS_7	24	21	53857363	coding	-	ITGAV				
RDA	26	22	48336761	Downstream 4303 bp	plexin-B1-like	PLXNB1	semaphorin receptor activity	signal transduction, semaphorin-plexin signaling pathway	Soong et al. (2012) Soong and Scott (2013)	Survival, growth of melanocytes
RDA	31	9	53901926	Upstream 7837 bp	rap guanine nucleotide exchange factor 2-like	RAPGEF2	guanyl-nucleotide exchange factor activity, GTPase activator activity, Rap guanyl-nucleotide exchange factor activity	signal transduction, small GTPase mediated signal transduction, regulation of cell proliferation, positive regulation of GTPase activity	Anssen et al. (2006)	Regulation of melanogenesis
RDA	31	9	53902142	Upstream 7621 bp		RAPGEF2				
RDA	31	9	57854367	coding	gamma-aminobutyric acid receptor subunit alpha-2-like	GABRA2	transmembrane signaling receptor activity, GABA-A receptor activity, ion channel activity, extracellular ligand-gated ion channel activity	ion transport, chloride transport, gamma-aminobutyric acid signaling pathway, ion transmembrane transport	Bertolesi et al. (2016)	Implied in light-regulation of skin color (Xenopus)
RDA	31	9	57854882	coding		GABRA2				
RDA	31	9	58319392	Downstream 6340 bp	polycomb group RING finger protein 1-like	PCGF1	metal ion binding	transcription DNA-templated, regulation of transcription DNA-templated, aging, histone H2A-K119 monoubiquitination, developmental growth	Huang & Hornyak (2015)	Post-translational modification in melanocyte development and melanogenesis
RDA	31	9	61830541	Upstream 1170 bp	protocadherin 10	PCDH10	calcium ion binding	cell adhesion, homophilic cell adhesion via plasma membrane adhesion molecules	Williams et al. (2018)	Melanocyte migration from neural crest cells (zebrafish)
RDA	31	9	61851844	Downstream 9983 bp	-	PCDH10	-	-	-	-
RDA	31	9	61851990	Downstream 10129 bp	protocadherin-10	PCDH10				
RDA	31	9	62603472	Downstream 1358 bp	cystine/glutamate transporter-like	SLC7A11	transmembrane transporter activity	transmembrane transport	Chintala et al. (2005) Xu et al. (2014) Wang et al. (2019)	Melanogenesis regulation and pheomelanin production, found in carp and tilapia
RDA	31	9	66501417	coding	protocadherin alpha-C2-like	PCDHAC2	calcium ion binding	cell adhesion, homophilic cell adhesion via plasma membrane adhesion molecules	Djurdjevic et al. (2019)	Differentially expressed in trout skin
RDA	31	9	74389966	coding	fibroblast growth factor receptor-like 1	FGFR1	nucleic acid binding	DNA integration	Harobe (2011) Cichorek et al. (2013) Czyz (2019) Aman et al. (2018)	Participate to keratinocyte-melanocyte cooperation in melanogenesis (may result in melanoma in humans). FGF signalling is essential to scale patterning

1352   **Figure 1:**

1353

1354

1355

1356

1357   **Figure 2:**

1358

1359 **Figure 3:**

1360

1361

1362

1363

1364 **Figure 4:**

1366

1367