The prevalence of *Angiostrongylus cantonensis* in rodent hosts in China and several Asian countries: a systematic review and meta-analysis

Huifang Bai[a,1*], Huiyuan Bai[b,1], Fei Bai[c,1], Yizhen Cao[a], Yunqiu Chen[a]

a Department of Pathogenic Biology, Medical College, Ji Nan University, Guangzhou 510632, PR China

b State Key Laboratory of Bioelectronics, School of Biological Science and Medical Engineering, Southeast University, Nanjing 210096, PR China

c College of Plant Science and Technology, Huazhong Agricultural University, Wuhan 430070, Hubei Province, PR China

[*] Corresponding author

E-mail address: bhfbhf92@sina.com

[1] These authors contribute equally to this work.

Abstract

Introduction

Angiostrogylus cantonensis (*A.cantonensis*) is a zoonotic parasitic nematode, with a worldwide distribution, causing eosinophilic meningitis or meningoencephalitis in human. Human infection with *A.cantonensis* typically occur from intentional or unintentional ingestion of infected intermediate or paratenic hosts, sometimes due to contaminated produce. Although the biology of *A.cantonensis* is relatively well known, little is understood about the transmission level in different zoogeographical regions, especially in Asia. Here, to evaluate the prevalence of *A.cantonensis* in

rodent hosts in China and several Asian countries, we conducted a systematic review

registered with PROSPERO (CRD42020161665).

Methods

Records were selected systematically from 7 database (Medline via to Pubmed, Web

of science, Scopus, Google Scholar, CNKI, Wangfang, CBM). Forest plots and

random-effects model were used to display pooled estimates. The Freeman-Tukey

double arcsine transformation of R software was used to conduct meta-analysis and

statistical significance was set at 0.05.

Results

A total of 67 studies met the inclusion criteria and were included in the systematic

review and meta-analysis. The pooled prevalence estimates of *A.cantonensis* infection

in rodents was 0.1003 (95%CI: 0.0765, 0.1268). There was significant heterogeneity

in reported outcomes (p<0.0001). And we considered that there was no publication

bias in the included studies.

Conclusion

The *A.cantonensis* infection rate among rodent hosts were still high in Asia,

particularly in China, especially in *Rattus norvegicus*, and thus comprehensive

measures should be taken for rodent hosts control to avoid an angiostrongyliasis

outbreak. Due to the wide distribution and movement of rodent hosts, people in all

regions of China, even in other Asia area live at risk of an infection. Hence, the

development of more reliable diagnostic tests will be key for an effective

identification of cases as well as improved patient care. Consequently, further studies

are required to updated strategies for controlling *A.cantonensis* infection among

human population.

Author summary

Angiostrongylus cantonensis (*A.cantonensis*) is a zoonotic parasitic nematode, with a

worldwide distribution, causing eosinophilic meningitis or meningoencephalitis in

human. Human infection with *A.cantonensis* typically occur from intentional or

unintentional ingestion of infected intermediate or paratenic hosts, sometimes due to

contaminated produce. Information about angiostrongyliasis burden, including its

prevalence and incidence and geographic distribution in Asia, which is necessary to

guide in planning appropriate control measures. In this study, we performed a

systematic review and meta-analysis of published studies in China and several Asian

countries on *A.cantonensis*. Our approach allowed evidence-based approximation to

the estimations of the burden of *A.cantonensis* infection, for different rodent hosts

groups, geographic regions, publication years and so on. At the same time, our review

showed the *A.cantonensis* infection rate among rodent hosts were still high in Asia,

particularly in China, especially in *Rattus norvegicus*. Finally, much bigger and

longer-term studies are required to updated strategies for controlling *A.cantonensis*

infection among human population.

Background

Angiostrongylus cantonensis (*A.cantonensis*) is a zoonotic parasitic nematode, with a

worldwide distribution, causing eosinophilic meningitis or meningoencephalitis in

human. Human infection with *A.cantonensis* typically occur from intentional or

unintentional ingestion of infected intermediate or paratenic hosts, sometimes due to

contaminated produce.

Traditionally, with the wider domain of climate and health, the neglected tropical

diseases (NTDs), a collection of infectious diseases affected hundreds of millions of

individuals living in tropical and subtropical countries. Recently, such dramatic

habitat changes are linked to accelerated biodiversity loss, which has been linked to

major changes in the epidemiology of human diseases: increased disease risks and the

emergence of novel pathogens resulting from increased contacts among wildlife,

domesticated animals, and humans[1-3]. In particularly, angiostrongyliasis, as a kind

of NTDs, was found in the Pacific Islands, Southeast Asia, parts of South and Central

America and the Caribbean, even in European countries[4-6]. Recently, although

researchers gradually pay more attention to investment towards reducing the burden

of NTDs, they still collectively contribute to productively loss, illness and suffering in

many countries[7,8]. Recent estimates of their overall burden suggest NTDs kill over

350,000 people per annum and cause the loss of between 27 and 56 million

disability-adjusted life years[9]. To date, more than 2000 cases of eosinophilic

meningitis have been recorded in China, even 100 cases in Taiwan and most human

cases of angiostrongyliasis (~1300 cases) have been reported in Thailand[5,10], where

between 0.3 and 2 people per 100,000 become infected annually[11,12]. These

findings indicate that the extensive distribution of *A.cantonensis* in global world is

high, human will be at high risk of *A.cantonensis* infection.

In recent decades, more than 20 species of feral rodents have been categorized as

pests in tropical and subtropical countries, and these species cause tremendous losses

and damages to crops and food stocks[13]. Feral rodents are known to transmit

diseases and act as reservoir hosts to many zoonotic parasites that pose health risks to

humans, including *A.cantonensis*, hantaviruses, *Leptospira spp.*, *Bartonella spp.*,

Trypanosoma spp., and *Babesia spp.*[14-16]. For *A.cantonensis,* at least 17 rodent

species may behave as definitive hosts, capable of passing first stage (L1) larvae in

their feces. Simultaneously, snails and slugs (intermediate hosts) ingest these

larvae, which mature into the third stage (L3) larvae (infective form) that are passed

on to the definitive hosts again when the definitive hosts consume infected

intermediate hosts. Then, the worms travel to brain, especially in the subarachnoid

space, elicits an intense eosinophilic inflammation and causes acute eosinophilic

meningitis. Finally, it will become a molluscs-rodents cycle to lead infectious.

Humans become the incidental hosts by consuming the infected intermediate or

paratenic hosts (or ingestion of snail slime)[17]. Importantly, the development of

control methods against zoonotic parasites depends on knowledge of their life cycles

and transmission patterns in different zoogeographical regions. Therefore, it is

essential to survey the relationship between parasites and rodents in order to identify

the sources of zoonotic infections[18].

Surprisingly, despite rodents being the dominant vertebrate species in many of these

altered tropical systems and the key hosts for numerous zoonotic microparasites

(viruses, bacteria, and protists), consistent patterns between habitat fragmentation and

environment change with the epidemiology of multiple parasite diseases have yet to

be determined, and studies that address the context of pathogen circulation are

lacking. Indeed, simultaneous to the technical and targeted approaches being,

recommended by WHO are much wider attempts at sustainable development, most

visible through the lens of the sustainable development goals[19]. Meanwhile, there

are little diagnostic methods and effective treatment methods. Furthermore, we need

do more research to solve this challenge. Here, in this study, we aim to identify the

prevalence of *A.cantonensis* in rodent hosts in China and several Asian countries,

provided essential recommendations for the researchers and policy makers.

Materials and methods

The protocol for this systematic review was defined in advance and registered with

PROSPERO (CRD42020161665). This study was conducted in accordance with the

PRISMA guidelines (Preferred Reporting Items for Systematic Reviews and

Meta-analysis) for systematic review and meta-analysis. The PRISMA checklist

(Table S1) was used to ensure inclusion of relevant information in the analysis.

Literature search

To evaluate the prevalence of *A.cantonensis* in rodent hosts in China and several

Asian countries, we carried out a systematic review of the literature (full-text)

published online in the English and Chinese between May 1946 and December 2019.

Records were selected systematically from 7 database (Medline via to Pubmed, Web

of science, Scopus, Google Scholar, CNKI, Wangfang, CBM). This range of years

was important to study the prevalence since the first study was reported. The modified

searches were performed by various combinations of the following terms using

Boolean operator "AND" and "OR": (rat lungworm OR *Angiostrongylus cantonensis*

OR *Angiostrongyliasis OR Angiostrongylidae*) AND (prevalence OR distribution OR

epidemic OR incidence OR frequency OR occurrence OR detection OR identification

OR characterization OR investigation OR survey OR rate) AND Asia. The reference

lists of selected articles also were screened manually and appropriate articles were

included. Full text articles were downloaded or obtained through library

resources.   .

Study selection

All selected articles had to meet the following inclusion criteria: (i) a cross-sectional

study or descriptive study; (ii) the study published between May 1946 and December

2019; (iii) the language was limited in Chinese or English; (iv) full-text article; (v)

infection cases in Asia, provided geographical location; (vi) reported as animal level

prevalence data; (vii) exact total numbers and positive cases numbers; (viii) rodent

hosts numbers higer than fifty.

Investigations without these criteria were excluded. Also, if the same study data were

published in both English and Chinese sources, the articles with less detailed

information would be excluded from this study. When any authors found articles

different to judge, the first author was consulted and eventual differences discussed to

reach a consensus decision on whether to include or exclude the article.

Quality assessment

The included articles were evaluated using a 10-items quality appraisal checklist (File

S1). An item would be scored '0' if it was answered 'no data available' or 'unclear';

if it was answered 'yes', then the item scored '1'. Article quality was assessed as

follows: low quality = 0–3; moderate quality = 4–7; high quality = 8–10.

Data extraction

For the data extraction, the two independent assessors applied the inclusion criteria

selected the articles and extracted the data. Discrepancies were resolved by consensus.

The detailed characteristics of each study were extracted using a pre-designed

data-collection excel form. Information was collected as followings: first author, year

of publication, location, geographical region, sample size, number of positive cases,

diagnostic methods, types of rodent hosts, quality score of each study.

Statistical analysis

The statistical software used in the analysis was R software version (3.6.3). Estimated

pooled prevalence and 95% confidence intervals (CI) were calculated with the

Freeman-Tukey double arcsine transformation. Heterogeneity testing was performed

using the I^2 and Cochran' Q statistic methods. A significant value (p<0.05) in the

analysis suggested a real effect difference. The I^2 values of 25%, 50% and 75% were

considered as low, moderate and high heterogeneity, respectively. The risk of study

bias was assessed using the funnel plots, and the Egger's regression test. Furthermore,

subgroup analysis were done if the heterogeneity were overlapping moderate I^2. Four

potential sources of heterogeneity were examined: country, animal species,

publication year, quality score. The Q and I^2 statistics values were calculated for each

subgroup to determine the effective factors on the prevalence *A.cantonensis* and

heterogeneity about all the included studies.

Results

Study search and included studies

A total of 2103 articles were screened in the initial and 67 were obtained for the final analysis. Fig 1 showed our systematic workflow for identifying, screening and including studies in this study. Hence, 59 articles from China, 1 from Japan, 4 from Thailand, 1 from India and 2 from Philippines. The complete list of included articles with the extracted data can be found in the Fig 2 and Supporting information (File S2). A total of 45196 rodents were dissected and 4954 were identified with *A.cantonensis* infection. The detailed characteristics of each study were provided in Table 1.

Fig 1. Flow-chart of systematic review of *A.cantonensis*

Fig 2. Map of the geographical distribution in Asian countries of the surveys

Table 1. Characterizatics of the eligible studies

Study	Location	Method	Number	Positive rate	quality score
(Zhang et al. 2017)	Fujian	NE	5266(541)	10.30%	7
(Huang et al. 2017)	Guangdong, Shenzhen	NE	238(22)	9.24%	8
(Hu et al. 2017)	Guangdong, Nan'ao island	NE, PCR	110(32)	29.10%	8
(Lin et al. 2016)	Fujian, Zhangzhou	NE	1551(142)	9.20%	9
(Eamsobhana et al. 2016)	24 provinces in Thailand	NE	669(46)	6.90%	7
(Ribas et al. 2016)	Udon Thani province	Staining	98(3)	3.10%	5
(Yang et al. 2015)	Yunnan	ELISA	304(0)	0.00%	6
(Tujan et al. 2016)	Nueva Ecija, Philippines	PCR	209(64)	29%	8
(Zhang et al. 2014)	Yunnan	NE	174（0）	0.00%	7
(Castillo et al. 2018)	Nueva Ecija, Philippines	NE	600(600)	100%	9
(Liang et al. 2014)	Guangdong, Shenzhen	NE	195(21)	10.80%	6
(Hu et al. 2018)	Guangdong, Nan'ao island	NE, PCR	110(32)	29.10%	8
(Xie et al. 2013)	Guangxi, Beihai	NE	58(1)	1.70%	5
(Huang et al. 2019)	Guangdong, Shenzhen	NE, ELISA	238(22)	9.20%	8

(Yang et al. 2012)	Guangdong, Guangzhou	NA	432(23)	5.30%	8
(Chen et al. 2011)	Guangdong	NE	1391(132)	9.50%	9
(Huang et al. 2011)	Guangdong, Shenzhen	NE	331(50)	15.10%	7
(Hu et al. 2011)	Hainan island	NA	118(13)	11.00%	6
(Vitta et al. 2011)	Phitsanulok province	NE	62(1)	1.60%	7
(Zeng et al. 2011)	Jangxi	Microscopy	502(12)	2.40%	4
(Zhu et al. 2010)	Fujian, nanping	NE	291(42)	14.40%	7
(Li et al. 2010)	Fujian	NE	4890(434)	8.90%	6
(Fu et al. 2010)	Guangxi	NE	149(5)	3.40%	5
(Deng et al. 2010)	Guangdong	NE	491(56)	11.40%	7
(Zhang et al. 2009)	Guangdong, yangchun	NE	192(11)	5.70%	4
(Renapurkar et al. 1982)	Bombay	NA	937(12)	1.30%	6
(Lan et al. 2010)	Zhejiang, Lishui	NE	1937(70)	3.61%	6
(Zhang et al. 2008)	Guangdong, Shenzhen	NE	248(22)	8.90%	7
(Li et al. 2008)	Yunnan	NE	124(4)	3.20%	4
(Ye et al. 2008)	Fujian, Fuzhou	NE	1964(269)	13.70%	6
(Wang et al. 2008)	Fujian, Ximen	NE	957(64)	6.70%	4
(Zeng et al. 2007)	Jiangxi	Microscopy	187(3)	1.60%	3
(Luo et al. 2005)	Fujian, Fuzhou	NE	1487(246)	16.50%	9
(Shen et al. 2004)	Guangdong, Zhaoqing	NE	221(12)	5.40%	8
(Yang et al. 2003)	Fujian	NE	625(52)	8.30%	6
Chen et al. 2003)	Fujian, Changle	Microscopy	112(44)	39.30%	3
Pan et al. 2002)	Zhejiang, Wenzhou	NE	87(17)	19.50%	4
(Yang et al. 2001)	Fujian	NE	579(40)	6.90%	5
(Liang et al. 2000)	Zhejiang, Wenzhou	NE	2069(316)	15.30%	9
(Zhang et al. 1999)	Fujiang, Fuzhou	NE	113(18)	19.50%	7
(Tung et al. 2013)	Taiwan, Taichung	NE	51(8)	15.70%	5
(Tokiwa et al. 2013)	Ogasawara Archipelago, Japan	NE	136(46)	33.80%	5
(Pipitgool et al. 1997)	Ubon Ratchathani	NE	272(3)	1.10%	7
(Lv et al. 2009)	China	NE	711(32)	4.50%	6
(Zhang et al. 2008)	Guangdong, Jiangmen	NE	229(10)	4.40%	8
(Deng et al. 2012)	Guangdong	NA	491(56)	11.40%	7
(Barratt et al. 2016)	Guangdong, guangzhou	NA	432(23)	5.30%	7
(Duan et al. 2008)	Hunan	NE	60(1)	1.70%	6
(Ma et al. 2008)	Guangdong, shenzhen	NE	248(22)	8.90%	8
(Lin et al. 2007)	Fujian, lianjiang	NE	178(27)	15.20%	6
(Pan et al. 2011)	Guangdong	NE	5820(496)	8.50%	8
(Lan et al. 2009)	Zhejiang, Lishui	NE	218(12)	5.50%	6

(Feng et al. 2008)	Guangdong,guangzhou	NE	113(4)	3.50%	7
(Yu et al. 2008)	Guangdong, shenzhen	NE	1510(301)	19.90%	9
(Wang et al. 2007)	Fujaing, xiamen	NE	957(64)	6.70%	5
(Lin et al. 2010)	Guangdong, qingyuan	NE	387(37)	9.60%	8
(Tan et al. 2008)	Guangdong, jiangmen	NE	229(10)	4.40%	7
(Li et al. 2008)	Guangxi, wuzhou	NE	193(4)	2.10%	7
(Chen et al. 2009)	Guangdong, maoming	NE	196(21)	10.70%	5
(Tong et al. 2011)	Hainan	NE	124(14)	11.30%	8
(Lin et al. 2011)	Guangdong, jiangmen	NE	113(30)	26.50%	7
(Geng et al. 2011)	Guangdong, shenzhen	ELISA	331(40)	12.10%	10
(Huang et al. 2010)	Guangdong	NE	293(24)	8.20%	7
(Zhang et al. 2009)	Guangdong, maoming	NE	148(15)	10.10%	7
(Lin et al. 2011)	Guangdong, qingyuan	NE	844(96)	11.40%	9
(Qu et al. 2011)	Guangdong, qingyuan	NA	288(27)	9.40%	6
(Zhang et al. 2008)	Guangdong, shenzhen	NA	307(38)	12.40%	4

Pooling and heterogeneity analysis

The pooled prevalence estimates of *A.cantonensis* infection in rodent hosts with individual studies were shown in a forest plot (Fig 3). A substantial heterogeneity was observed among studies (χ^2=4979.77, P<0.05; I^2=99.0%). When calculated using a random-effects model, the overall infection rate was 0.1003 (95%CI: 0.0765,0.1268). The estimates of infection rates for different subgroups and heterogeneities were presented in Table 2 and Fig 4. All pooled infection rates for each subgroup were calculated using a random-effects model bacause of the observed high heterogeneity among studies within subgroups. Base on study periods, the estimate had been significantly increased since 2008 (≤2008: 0.0834 (95%CI: 0.0595, 0.1107)vs＞2008: 0.111 (95%CI: 0.0758, 0.1518)). In terms of the rodent hosts species, the estimate was highest in *Rattus norvegicus* (0.1441, 95%CI: 0.1100, 0.1818), followed by *Rattus flavipectus* (0.0428, 95%CI: 0.0062, 0.0987), *Bandicota indica* (0.0409, 95%CI:

0.0000, 0.1240), *Mus musculus* (0.0101, 95%CI: 0.0041, 0.0187), *Rattus rattus*

(0.0093, 95%CI: 0.0004, 0.0299), *Rattus losea* (0.0044, 95%CI: 0.0000, 0.0176), and

the lowest was *Suncus Murinus* (0.0006, 95%CI: 0.0000, 0.0086). In terms of Asia

region, China (0.0893, 95%CI: 0.0773, 0.1020) was higher than Thailand (0.0299,

95%CI: 0.0053, 0.0700). Finally, in terms of score, the estimate was highest in high

score(0.1266, 95%CI: 0.1021, 0.1560), followed by low score (0.0892, 95%CI:

0.0420, 0.1509), the lowest was middle score (0.0728, 95%CI: 0.0583, 0.0888).

Fig. 3 Random-effects model meta-analysis of *A.cantonensis* infection in Asian

countries

Fig. 4 Subgroup froest plot of infection *A.cantonensis* in Asian countries

Table.2 Subgroup meta-analysis of *A.cantonensis* in Asian countries

variable	population	No.of infected rodents	No. of total rodents	Pooled estimate Prevalence (%)	95%CI	Heterogeneity(Q)			Egger'test	
						Q	I²(%)	P-value	t	P
Asia region										
China	42213(4179)	4179	42213	0.0893	(0.0773,0.1020)	1005.38	95.1	<0.0001	-0.5678	0.5724
Thailand	1101(53)	53	1101	0.0299	(0.0053,0.0700)	19.67	93.9	0.0002	-1.0502	0.4038
Japan	136(46)	46	136							
Philippines	809(664)	664	809							
india	937(12)	12	937							
Score										
≤4 (low)	2469(192)	192	2469	0.0892	（0.0420,0.1509）	139	96.8	<0.0001	1.0588	0.3304
＞4 to ≤7 (middle)	24578(2103)	2103	24578	0.0728	（0.0583,0.0888）	674.28	95.4	<0.0001	-0.8451	0.4035
＞7 to ≤10 (high)	18149(2659)	2659	18149	0.1266	(0.1021,0.1560)	421.26	96.5	<0.0001	0.4219	0.6781
Host species										
Rattus norvegicus	18479(3048)	3048	18479	0.1441	(0.1100,0.1818)	2320.32	98	<0.0001	-0.4383	0.663
Rattus flavipectus	7840（918）	918	7840	0.0428	(0.0062,0.0987)	2230.08	98.3	<0.0001	-0.409	0.6846

Suncus Murinus	5758（179）	179	5758	0.0006	（0.0000,0.0086）	164.89	87.2	<0.0001	-0.7805	0.4416
Mus musculus	1137（22）	22	1137	0.0101	（0.0041,0.0187）	34.88	44.6	0.2469	-2.2943	0.0292
Rattus losea	1697（69）	69	1697	0.0044	（0.0000,0.0176）	25.5	62.3	0.0841	1.8715	0.0797
Bandicota indica	800（66）	66	800	0.0409	（0.0000,0.1240）	90.17	93.1	<0.0001	0.3826	0.7109
Rattus rattus	797(19)	19	797	0.0093	（0.0004,0.0299）	35.34	79.3	0.0008	0.8389	0.4179
other species	4269(415)	415	4269							
publication year										
≤2008	14007(1602)	1602	14007	0.0834	(0.0595,0.1107)	665.2	97.1	<0.0001	-1.833	0.0798
>2008	31189(3352)	3352	31189	0.111	(0.0758,0.1518)	4311.74	99.1	<0.0001	0.6216	0.5377
Overall	49156(4954)	4954	49156	0.1003	(0.0765,0.1268)	4979.77	99.0	<0.0001	0.1075	0.9147

Publication bias

Despite the significant heterogeneity, the funnel plot displayed a symmetric spread of studies in terms of relative weight and effect size, thereby we could not directly judge whether there was publication bias in the included studies (Fig. 5). The results of Egger's test showed P=0.9147 (P>0.05) (data not shown). Therefore, we considered that there was no publication bias in the included studies. Notably, the total number of studies was reasonable, and the individual studies were of variable sample size. Moreover, Duval and Tweedie's trim and fill procedure for the detection of publication bias did not support the possibility of missing studies from the analysis.

Fig. 5 Funnel plot of the arcsine transformed infection rates of *A.cantonensis* in Asian countries

Discussion

In previous study, rodents may play an important role in the transmission of certain zoonotic parasites to humans[18]. For *A.cantonensis*, the estimate was highest in *Rattus norvegicus*, *Rattus flavipectus* and *Bandicota indica*, followed by *Mus musculus*, *Rattus rattus* , *Rattus losea*, and *Suncus Murinus*. The difference of the

infection rates among seven kinds of rats may be relative to the factors: the

susceptibility to *A. cantonensis*, rat's habitat, activity range and hunted objects, and so

on. For example, the front several rodents : (i) be considered as dominant species in

several Asian countries[20,21]; (ii) live close to human dwellings[22]; (iii) they were

also known to prey on snails not plants[23]. The outcome is comparable to the

previous report[24].

Before 2008, there were outbreaks in Beijing, Wenzhou, Guangzhou and so on in

China, particularly, 160 cases infected in Beijing[25]. Then, investigator did more

research across China by the national angiostrongyliasis control programme. They

found this phenomenon was supported by the environment change, including soil,

water and air. The climate changes influence global air and land surface temperatures.

Also, soil and water are important to the natural history of many parasite or their

vectors. In such environments, a high prevalence of parasite infection (hantavirus,

Yersinia pestis) is often found in various rodents species[26]. In normal, *Rattus spp.*

feces could be in water or soil. The major intermediate hosts (e.g. *Pomacea*

canaliculata and *Achatina fulica*)[27] maybe infected by ingesting feces. Typically,

soil and water properties' changes will affect the intermediate hosts and definitive

hosts living time. Their living stages are an essential component of the *A.cantonensis*

infection life cycle. Moreover, with the wide habitat alternation, the hosts may

adapted and mitigated quickly to modify emissions and habitats. Finally, all factors

mentioned earlier leads cases increased year by year.

Limitations

Though this study provided information for the prevalence of *A.cantonensis* in rodent

hosts in Asia, several limitations may merit attention. First, the estimations of

infection rates using a random-effect model was unavoidable, then subgroup analysis

was employed but could not fully explain the source of heterogeneity. Secondly, our

study didn't included all Asian countries because the relevant data were not available

or incomplete. Finally, in this study, we only focused on *Rattus norvegicus*, *Rattus*

flavipectus, *Bandicota indica*, *Mus musculus*, *Rattus rattus*, *Rattus losea* and *Suncus*

Murinus, because they were common and most important rodent hosts. However,

other rodents, like *Rattus tiomanicus*, *Rattus exulans* and so on, though difficulty to

collect them but should not be neglected. To fully elucidate the prevalence of

angiostrongyliasis, further studies should pay attention to the other species rodent

hosts.

Conclusion

In conclusion, our systematic review and meta-analysis estimated a national

prevalence of *A.cantonensis* in rodent hosts in Asia, particularly in China, which

provided a scientific basis for prevention a control of the spread of angiostrongyliasis.

The *A.cantonensis* infection rate among rodent hosts were still high in China,

especially in *Rattus norvegicus*, and thus comprehensive measures should be taken for

rodent hosts control to avoid an angiostrongyliasis outbreak. Due to the wide

distribution and movement of rodent hosts, people in all regions of China, even in

other Asia area live at a risk of an infection. Moreover, public health officials,

epidemiologists,   researchers,   clinical   technicians,   medical   practitioners,

parasitologists, and veterinarians, as well as the general public, should be aware of

such risks, and integrated strategies should be taken to reduce or eliminate such

risks[28]. Hence, the development of more reliable diagnostic tests will be key for an

effective identification of cases as well as improved patient care. Consequently,

further studies are required to updated strategies for controlling *A.cantonensis*

infection among human population.

Supporting information

Table S1. PRISMA checklist.

(DOC)

File S1.Quality assessment checklist.

(DOCX)

File S2. References of the selected articles included in the systematic review.

(DOCX)

Funding information

290 This work has received no specific funding.

Declaration of interests

292 We declare no competing interests.

Author Contributions

294 Conceptualization: Huifang Bai, Huiyuan Bai, Fei Bai.

295 Data curation: Huifang Bai, Huiyuan Bai, Fei Bai.

296 Formal analysis:Huifang Bai, Huiyuan Bai.

297 Investigation: Huiyuan Bai, Fei Bai.

Methodology: Huifang Bai, Huiyuan Bai, Fei Bai.

Validation: Huifang Bai, Yizhen Cao, Yunqiu Chen.

Visualization: Huifang Bai, Yizhen Cao, Yunqiu Chen.

Writing-original draft: Huifang Bai, Huiyuan Bai.

Writing-review & editing: Huifang Bai, Huiyuan Bai, Fei Bai, Yizhen Cao, Yunqiu Chen.

References

1. Luis AD, Hayman DT, O'Shea TJ, Cryan PM, Gilbert AT, Pulliam JR, et al. A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special? Proc Biol Sci. 2013; 280(1756): 20122753. pmid:23378666

2. Civitello DJ, Cohen J, Fatima H, Halstead NT, Liriano J, McMahon TA, et al. Biodiversity inhibits parasites: Broad evidence for the dilution effect. Proc Natl Acad Sci U S A. 2015; 112(28): 8667-8671. pmid:26069208

3. Olival KJ, Hosseini PR, Zambrana-Torrelio C, Ross N, Bogich TL, Daszak P. Host and viral traits predict zoonotic spillover from mammals. Nature. 2017; 546(7660): 646-650. pmid:28636590

4. Rosen L, Chappell R, Laqueur GL, Wallace GD, Weinstein PP. Eosinophilic meningoencephalitis caused by a metastrongylid lung-worm of rats. Jama. 1962; 179(620-624. pmid:14493905

5. Wang QP, Lai DH, Zhu XQ, Chen XG, Lun ZR. Human angiostrongyliasis. Lancet Infect Dis. 2008; 8(10): 621-630. pmid:18922484

6. Thiengo SC, Simões Rde O, Fernandez MA, Maldonado A, Jr. Angiostrongylus

cantonensis and rat lungworm disease in Brazil. Hawaii J Med Public Health. 2013;

72(6 Suppl 2): 18-22. pmid:23901376

7. Molyneux D, Sankara DP. Guinea worm eradication: Progress and challenges-

should we beware of the dog? PLoS Negl Trop Dis. 2017; 11(4): e0005495.

pmid:28426663

8. Conteh L, Engels T, Molyneux DH. Socioeconomic aspects of neglected tropical

diseases. Lancet. 2010; 375(9710): 239-247. pmid:20109925

9. Hotez PJ. Ten global "hotspots" for the neglected tropical diseases. PLoS Negl

Trop Dis. 2014; 8(5): e2496. pmid:24873825

10. Wang QP, Wu ZD, Wei J, Owen RL, Lun ZR. Human Angiostrongylus

cantonensis: an update. Eur J Clin Microbiol Infect Dis. 2012; 31(4): 389-395.

pmid:21725905

11. Suankratay C, Wilde H, Berger S. Thailand: country survey of infectious diseases.

J Travel Med. 2001; 8(4): 192-203. pmid:11703902

12. Eamsobhana P, Lim PE, Yong HS. Genetic diversity of the rat lungworm,

Angiostrongylus cantonensis, the major cause of eosinophilic meningitis. Hawaii J

Med Public Health. 2013; 72(6 Suppl 2): 15-17. pmid:23901375

13. Claveria FG, Causapin J, de Guzman MA, Toledo MG, Salibay C. Parasite

biodiversity in Rattus spp caught in wet markets. Southeast Asian J Trop Med

Public Health. 2005; 36 Suppl 4(146-148. pmid:16438200

14. Tangkanakul W, Smits HL, Jatanasen S, Ashford DA. Leptospirosis: an emerging

health problem in Thailand. Southeast Asian J Trop Med Public Health. 2005;

36(2): 281-288. pmid:15916031

15. Truc P, Büscher P, Cuny G, Gonzatti MI, Jannin J, Joshi P, et al. Atypical human

infections by animal trypanosomes. PLoS Negl Trop Dis. 2013; 7(9): e2256.

pmid:24069464

16. Watt G, Pachirat O, Baggett HC, Maloney SA, Lulitanond V, Raoult D, et al.

Infective endocarditis in northeastern Thailand. Emerg Infect Dis. 2014; 20(3):

473-476. pmid:24572588

17. Sohal RJ, Lui F (2020) Angiostrongylus Cantonensis (Angiostrongliasis).

StatPearls. Treasure Island (FL): StatPearls Publishing

Copyright © 2020, StatPearls Publishing LLC.

18. Tung KC, Hsiao FC, Wang KS, Yang CH, Lai CH. Study of the endoparasitic

fauna of commensal rats and shrews caught in traditional wet markets in Taichung

City, Taiwan. J Microbiol Immunol Infect. 2013; 46(2): 85-88. pmid:22516746

19. Griggs D, Stafford-Smith M, Gaffney O, Rockström J, Ohman MC, Shyamsundar

P, et al. Policy: Sustainable development goals for people and planet. Nature. 2013;

495(7441): 305-307. pmid:23518546

20. Lv S, Zhang Y, Liu HX, Zhang CW, Steinmann P, Zhou XN, et al.

Angiostrongylus cantonensis: morphological and behavioral investigation within

the freshwater snail Pomacea canaliculata. Parasitol Res. 2009; 104(6): 1351-1359.

pmid:19172296

21. Salibay C. Distribution of Native and Non-native Rats (Rattus sp) Along an

elevational gradient in the tropical Rainforest of Southern Luzon, Philippines.

Ecotropica. 2008; 14(129-136.

22. Hu QA, Lv S, Guo YH, Liu HX, Zhang Y. Genetic diversity of Angiostrongylus

cantonensis in Nan'ao Island of Guangdong Province. Chin J Parasitol Parasit Dis.

2017; 35(2): 130-135.

23. Hu QA, Zhang Y, Guo YH, Lv S, Xia S, Liu HX, et al. Small-scale spatial

analysis of intermediate and definitive hosts of Angiostrongylus cantonensis. Infect

Dis Poverty. 2018; 7(1): 100. pmid:30318019

24. Chen D, Zhang Y, Shen H, Wei Y, Huang D, Tan Q, et al. Epidemiological survey

of Angiostrongylus cantonensis in the west-central region of Guangdong Province,

China. Parasitol Res. 2011; 109(2): 305-314. pmid:21301874

25. MacDonald R. Eating raw snails leads to angiostrongylus meningitis in Beijing.

The Lancet Infectious Diseases. 2006; 6(10): 622.

26. McCauley DJ, Salkeld DJ, Young HS, Makundi R, Dirzo R, Eckerlin RP, et al.

Effects of land use on plague (Yersinia pestis) activity in rodents in Tanzania. Am J

Trop Med Hyg. 2015; 92(4): 776-783. pmid:25711606

27. Lv S, Zhang Y, Steinmann P, Zhou XN. Emerging angiostrongyliasis in Mainland

China. Emerg Infect Dis. 2008; 14(1): 161-164. pmid:18258099

28. Zhang RL, Chen MX, Gao ST, Geng YJ, Huang DN, Liu JP, et al. Enzootic

angiostrongyliasis in Shenzhen, China. Emerg Infect Dis. 2008; 14(12): 1955-1956.

pmid:19046534

Figure

Figure

Figure

Figure

Figure