Non-random segregation of sister chromosomes by *Escherichia coli* MukBEF axial cores

Jarno Mäkelä, Stephan Uphoff, David J. Sherratt

Department of Biochemistry, University of Oxford, Oxford OX1 3QU, UK
Correspondence to: jarno.makela@bioch.ox.ac.uk; david.sherratt@bioch.ox.ac.uk

Summary

The *Escherichia coli* structural maintenance of chromosomes complex, MukBEF, forms axial cores to chromosomes that determine their spatio-temporal organization. Here, we show that axial cores direct chromosome arms to opposite poles and generate the translational symmetry between newly replicated sister chromosomes. MatP, a replication terminus (*ter*) binding protein, prevents chromosome rotation in the longitudinal cell axis by displacing MukBEF from *ter*, thereby maintaining the linear shape of axial cores. During DNA replication, MukBEF-MatP action directs lagging strands towards the cell center, marked by accumulation of DNA-bound β_2-clamps in the wake of the replisome, in a process necessary for translational symmetry. Finally, the ancestral template DNA strand, propagated from previous generations, is preferentially inherited by the cell forming at the old pole, dependent on MukBEF-MatP. The work demonstrates how chromosome organization-segregation fosters non-random inheritance of genetic material and provides a framework for understanding how chromosome conformation and dynamics shape subcellular organization.

Highlights

- Linear MukBEF axial cores are required for *left-oriC-right* chromosome organization
- MatP prevents longitudinal nucleoid rotation by linearizing the axial cores
- MukBEF axial cores direct lagging strands towards the cell center
- The ancestral DNA strand is preferentially segregated to the older pole cell

Introduction

Faithful chromosome propagation and inheritance underpin all replicative life. Organisms have evolved a vast range of mechanisms to ensure timely replication and segregation of genetic material. Eukaryotic cells orchestrate replication and segregation in discrete stages where newly replicated sister chromosomes are first individualized by condensin and held together by cohesin before being pulled apart by action of the mitotic spindle and cleavage of cohesion (reviewed in (Uhlmann, 2016)). In prokaryotes, chromosomes replication and segregation are not temporally separated, limiting bacteria to a single replicon and segregon per chromosome (Kuzminov, 2014). Because divergent bacterial species have evolved different solutions to the same problem, understanding the contributions of different mechanisms and physical constraints underlying robust chromosome segregation remains a

challenge (Badrinarayanan et al., 2015; Surovtsev and Jacobs-Wagner, 2018; Wang et al., 2013).

Genetic studies have identified two major classes of proteins implicated in chromosome segregation in bacteria. First, structural maintenance of chromosomes (SMC) complexes, MukBEF, MksBEF and Smc-ScpAB; with MukBEF initially identified in a screen for *Escherichia coli* mutants that generated anucleate cells as a consequence of a failure to segregate newly replicated chromosomes to daughter cells (Hiraga et al., 1989; Nolivos and Sherratt, 2014). Second, studies of low copy plasmid stability identified ParAB*S* systems, which subsequently were shown to have roles in chromosome segregation in many organisms (Surovtsev and Jacobs-Wagner, 2018). While many bacteria encode one or both of these systems, some, for example *Pseudomonas aeruginosa*, encode two different SMCs and a ParAB*S* system (Petrushenko et al., 2011; Vallet-Gely and Boccard, 2013). Nevertheless, deletion of SMC or ParAB proteins has frequently modest if any consequences for chromosome segregation. Consistent with this, it has been proposed that large bacterial chromosomes can utilise repelling entropic effects to facilitate separation of bacterial chromosomes (Jun and Mulder, 2006), unlike much smaller low copy number plasmids that require a functional ParAB*S* system for faithful segregation (Surovtsev and Jacobs-Wagner, 2018). Whatever roles entropic forces may play, studies in diverse bacterial species have demonstrated that chromosomal loci are not positioned randomly in cells (Fogel and Waldor, 2005; Umbarger et al., 2011; Vallet-Gely and Boccard, 2013; Wang et al., 2005, 2006, 2014), and that in *E. coli*, MukBEF complexes play an important role in correct positioning of replication origins and other loci by forming an axial core to the chromosome (Danilova et al., 2007; Mäkelä and Sherratt, 2020). Absence of MukBEF leads to formation of anucleate cells during growth and loss of viability at temperatures higher than 22 °C in rich media (Danilova et al., 2007; Niki and Jaffe, 1991).

In new-born *E. coli* cells growing under conditions in which replication cycles do not overlap, origins of replication (*oriC*) are positioned close to the cell center, and the left and right chromosome arms are linearly organized in separate cell halves. Chromosome replication-segregation leads to generation of daughter cells with a chromosome organization identical to their mother cell. Most cell adopt a *left-oriC-right-left-oriC-right* (*L-R-L-R*) translational symmetry prior to division (Wang et al., 2006), which requires that either the leading or lagging strand templates are symmetrically segregated to the cell poles (Toro and Shapiro, 2010; Wang et al., 2005). In agreement, an elegant chromosome degradation experiment showed that the leading strand templates are segregated towards the cell poles in most cells (White et al., 2008). In theory, cells could also control the fate of the old template strand by non-random segregation, designating the destination for each strand. During each replication cycle, there is a risk of the new strand not faithfully copying the information from the template strand. 'Immortal' strand retention was originally proposed as a strategy to maintain DNA purity in stem cells while the copied strands, potentially carrying mutations from replication, were segregated to non-stem cell progeny (Cairns, 1975). Whether this strategy is actually utilized by stem cells remains controversial (Lansdorp, 2007; Rando, 2007; Wakeman et al., 2012). Ancestral strand segregation has also been tested in *C. crescentus* (Marczynski et al., 1990; Osley and Newton, 1974) and *B.*

subtilis (Errington and Wake, 1991), however, none of these studies showed any
segregational strand preference between daughter cells.
We lack a mechanistic understanding of how chromosome conformation and
orientation is maintained inside a bacterial cell. It also remains unknown how progressive
chromosome segregation facilitates non-random sister chromosome inheritance in an
otherwise apparently symmetrical organism. In this study, we address these questions in *E.*
coli utilizing microfluidics time-lapse imaging, high-throughput microscopy and quantitative
analysis. We first demonstrate that in the absence of MukBEF, anucleate cells arise
predominantly from the mother cell's new pole as a consequence of the failure to segregate
new replicated origins in a timely fashion. We show that newly replicated lagging strands
and their templates are directed towards cell centers, a process that is required for the
observed translational *L-R-L-R* segregational symmetry; and which is perturbed in the
absence of MukBEF. Furthermore, we show directly that the ancestral DNA strand, inherited
from previous generations, is preferentially segregated to the old cell pole dependent on
both MukBEF and its partner MatP. Lack of MatP does not perturb immediate asymmetry
after replication-segregation; rather it leads to loss of chromosome orientation during cell
cycles, consistent with the observed loss of ancestral strand retention at old poles. Taken
together, the results explain how MukBEF axial cores and their MatP-driven depletion from
the *ter* region, lead to asymmetric strand and chromosome segregation. The possible
functional and evolutionary consequences of this are explored.
105
Results

In the absence of MukBEF, anucleate cells arise from the newer mother cell pole

To understand how anucleate *E. coli* cells form in the absence of MukBEF, we followed
successive cell cycles of *ΔmukB* cells with *oriC* and *ter* (*ori1* and *ter3*) regions fluorescently
labeled by FROS markers. We used a 'mother machine' microfluidics device (Uphoff, 2018;
Wang et al., 2010) to follow thousands of cell generations and identify changes in
chromosome organization that correlate with chromosome mis-segregation. Under these
conditions, 15.7 ± 0.4% (±SD) of divisions led to the formation of an anucleate daughter cell
(Fig. 1A, Fig. S1).
In *ΔmukB* cells *ori1* was of often mis-localized towards the old pole at birth, rather
than at midcell (Fig. 1B), as previously observed for wild type (WT) cells (Wang et al., 2006).
As the cell cycle progressed, *ori1* localized preferentially towards the old cell pole, with
newly replicated sister *ori1* loci frequently remaining in close proximity. In contrast, *ter3*
migrated from the new-born cell pole to midcell similar to what has been previously
reported (Fig. 1B)(Wang et al., 2005). ~80% of anucleate cells were generated when
duplicated *ori1* loci in mother cells remained together in the region of the old pole prior to
cell division (Fig. 1C). In contrast, in ~70% of mother cells where chromosome segregation
was faithful, *ori1* loci were visible as separate foci (Fig. 1C). Delayed separation of newly
replicated *ori1* loci could be a consequence of delayed decatenation, since the decatenase
TopoIV, is no longer recruited by MukBEF to *oriC*-proximal regions (Zawadzki et al., 2015).
Indeed, modest over-expression of TopoIV leads to a reduction in cohesion time of newly
replicated *oriC* from ~14 min to ~5 min (Wang et al., 2008). In anucleate cell divisions,

daughter cells that inherited both chromosomes divided normally after a modest increase in
generation time (Fig. S1).
Prior to anucleate cell formation, mother cells divided nearly symmetrically (2.1 ± 0.2
μm and 2.4 ± 0.2 μm, respectively (±SD); two-sample t-test p-value 0.17), with the divisome
being placed close to midcell. While the average anucleate cell length at birth did not
significantly differ from that of the sister, the growing sister was systematically longer than
the anucleate sister (Fig. S1), the bias for the longer growing sister increasing with mother
cell division size. Importantly, we showed that anucleate cells formed preferentially at the
newer mother cell pole (74.4 ± 1.8% (±SD), Fig. 1D). Therefore, anucleate cell formation is
associated with the nucleoid being preferentially retained at the old pole of *ΔmukB* mother
cells, while in the case of WT cells the nucleoid is localized closer to the newer pole of the
dividing cell (Fisher et al., 2013). We conclude that mis-localization of *ori1* towards the old
pole, accompanied by delayed segregation of newly replicated *ori1* loci, directs the
formation of anucleate cells to the mother cell's new pole.
142
MukBEF and MatP direct left-oriC-right chromosome organization

Next we investigated how MukBEF directs chromosome organization and segregation in
growing cells. MukBEF and MatP have been proposed to be major factors in dictating *left-*
oriC-right (*L-R*) chromosome organization in *E. coli.* MukBEF complexes form axial cores that
linearly organize the chromosome outside the 800 kb *ter* region, from where MatP bound to
matS sites displaces MukBEF (Fig. 2A)(Mäkelä and Sherratt, 2020). Complete axial cores
were most easily visualized in cells in which MukBEF occupancy on the chromosome was
increased ~3.5-fold, while cells with WT MukBEF abundance on chromosomes exhibited
more granular structures (Mäkelä and Sherratt, 2020). To characterize how MukBEF axial
cores influence chromosome organization during the cell cycle, we used strains that allowed
us to test the requirements for *left* and *right* chromosome arm organization in relation to
oriC and *ter* in MatP⁺ and *ΔmatP* cells with WT levels of MukBEF, and in *ΔmukB* cells (Fig. 2).
The *left* and *right* chromosome arms were labeled at *L3* and *R3* (128° and 122° from *oriC*,
respectively) with FROS markers, as were *ori1* and *ter3* loci (Fig. 2A, B).
New-born WT cells exhibited the distinctive *left-oriC-right* (*L3-R3*) chromosome
organization (Fig. 2C, D and E, Fig. S2), where *oriC* remained at the cell center and the
chromosome arms (*L3* and *R3*) resided in opposite cell halves (97.8 ± 0.6% (±SD), Fig. 2F and
G) (Nielsen et al., 2006; Wang et al., 2006). During replication-segregation, the pattern was
extended into a translationally symmetric *left-oriC-right-left-oriC-right* (*L3-R3-L3-R3*) pattern
in 73.1 ± 3.9% (±SD) of WT cells (Fig. 2G). In the absence of MukBEF, we expected
chromosome organization to be more 'relaxed' as lengthwise chromosome compaction is
relieved (Mäkelä and Sherratt, 2020). Indeed, the localization of all four chromosomal
markers was less precise, with a wide distribution of *L3-R3* distances (Fig. 2F), fewer *L3* and
R3 foci localized in opposite cell halves (56.6 ± 3.2% (±SD), Fig. 2G), and a random chance of
observing the *L3-R3-L3-R3* organization (47.7± 0.2% (±SD), versus *L3-R3-R3-L3* or *R3-L3-L3-*
R3. The impaired chromosome organization in *ΔmukB* cells is frequently accompanied by
the chromosome arms being aligned together along the long cell axis with *ori1* towards the
old cell pole (Fig. 2B and D)(Danilova et al., 2007), an organization reminiscent of the

situation in *C. crescentus* (Wang et al., 2013). We conclude that absence of lengthwise
compaction by MukBEF axial cores causes loss of both the distinctive *L-R* chromosome
organization prior to replication and the *L-R-L-R* organization after replication.
Absence of MatP leads to circular MukBEF axial core formation, rather than linear
ones (Fig. 2A). We observed that *ΔmatP* cells exhibited chromosome locus localization
patterns strikingly different from that of WT and *ΔmukB* cells (Fig. 2 C and E). The average
distance between *L3* and *R3* was reduced two-fold (Fig. 2F), consistent with MukBEF-
mediated lengthwise compaction of *ter* in the absence of MatP (Mäkelä and Sherratt, 2020).
Lengthwise compaction of *ter* reduced the efficiency of *L3* and *R3* being directed into
opposite cell halves (65.7 ± 0.8% (±SD), Fig. 2G). Concomitantly, it also led to *L3* and *R3* foci
being preferentially localized closer to cell center than segregated sister *ori1* loci during
replication, in contrast to WT cells where *L3* and *R3* localize towards the cell poles (Fig. 2E).
Surprisingly, despite these substantial perturbations, the normal *L3-R3-L3-R3* organization
was retained in *ΔmatP* cells prior to cell division (80.2 ± 1.9% (±SD)).
Because the circular MukBEF axial cores of *ΔmatP* cells lead to rotational
chromosome symmetry (Mäkelä and Sherratt, 2020), we hypothesized that non-replicating
chromosomes in new born cells are free to rotate around the longitudinal axis of the cell.
This would switch the configuration from *L3-R3* to *R3-L3* (or *vice versa*); while bi-lobed
replication intermediates could prevent the rotation as replication progresses. To test this
hypothesis, we followed *ΔmatP* cells under the microscope to observe how often the *L3-R3-*
L3-R3 orientation flips to *R3-L3-R3-L3* (or *vice versa*) in consecutive generations (Fig. S2).
Indeed, *ΔmatP* cells retained the chromosome orientation only in 32.2 ± 4.6% (±SD) of
daughter cells, while WT cells predominantly retained the orientation (91.4 ± 5.2% (±SD),
Fig. 2H). We propose that the *L3-R3-L3-R3* organization arises from an interplay between
bidirectional replication and the action of MukBEF axial cores.
196

DnaN marks lagging strand segregation to the cell center

Translational symmetry of sister chromosomes arises at least in part from the symmetric
segregation of lagging strands during DNA replication towards midcell (and leading strands
towards the cell poles) (White et al., 2008). While this work used an elegant genetic system
to infer the segregation pattern, we sought directly to visualise the positioning of lagging
strands in WT, *ΔmatP* and *ΔmukB* cells.
During replication, ~20 DNA-bound β_2-clamps, which ensure DNA polymerase III
processivity, accumulate on the lagging strand and its template because of a ~3 min
residence time before clamps are unloaded from completed Okazaki fragments (Fig. 3A)
(Moolman et al., 2014). We reasoned that since β_2-clamps could potentially cover >100kb of
newly replicated lagging strand DNA, they could serve as a marker to monitor lagging strand
segregation (Fig. 3B). As a reference, for the localization of replication forks, we imaged
fluorescent DNA polymerase III ε-subunits (DnaQ). Strikingly, while DnaN localized in the
vicinity of cell center, DnaQ foci were more spread towards cell poles (Fig. 3C). By directly
measuring the distance from each DnaQ focus to the closest DnaN focus, we found that 41.2
± 5% (±SD) of DnaQ foci do not colocalize with DnaN foci during replication (Fig. 3D).
Differential location of bulk DnaN and replication forks was confirmed by measurement of

the distances from replicative helicase (DnaB) foci to their closest DnaN focus (47.1 ± 6.1%
(±SD) not colocalizing) (Fig. S3B, C). Since DnaN and DnaQ colocalize during early and late
replication, when sister replisomes are necessarily close together, we also show focus
localization in mid-replication cycle (Fig. 3E), when independently tracking replication forks
are more frequently spatially separate. The pattern of DnaQ foci that did not colocalize with
DnaN foci (Fig. S3F) underlines the conclusion that spatially separate sister replisomes in
different cell halves have a different cellular location from the bulk DnaN. Our results are
consistent with previous independent measurements of DnaQ and DnaN localization, and
the observation that DnaN foci often do not spatially separate (Mangiameli et al., 2017;
Reyes-Lamothe et al., 2008; Wallden et al., 2016). They provide the first evidence that the
replisome and β$_2$-clamps frequently do not colocalize during replication.
Our direct visualization of the segregation of lagging strands during replication by
imaging DnaN, supports the previously shown symmetric segregation of leading strands
towards the cell poles (White et al., 2008). To analyze how MukBEF and MatP contribute to
lagging strand segregation, we measured DnaN localization in *ΔmatP* and *ΔmukB* cells. The
DnaN distribution in *ΔmatP* cells was much broader than in WT cells (Fig. 3E and F),
indicative of spatially less precise lagging strand segregation, but still directed towards cell
centers, as predicted by the *L3-R3-L3-R3* organization. The DnaQ distribution in mid-cycle
ΔmatP cells was more central than that of DnaN (Fig. 3E, 50.8 ± 1.3% (±SD) colocalization
with DnaN), most likely because of less separated chromosome arms. Both DnaQ and DnaN
exhibited a broader distribution at shorter cell lengths (Fig. 3F), presumably because of a
more random chromosome conformation (Fig. 2). *ΔmukB* cells exhibited a bimodal
distribution of DnaN and DnaQ localizations towards cell poles, with the DnaQ and DnaN
profiles being almost identical (Fig. 3G). This shows that lagging strands and their templates
cannot be directed to cell centers in a timely manner in the absence of MukBEF function, a
result consistent with impaired *L-R* and *L-R-L-R* organization in *ΔmukB* cells. By measuring
the distance from each DnaQ focus to the closest DnaN focus, we found that lagging strands
did not leave the vicinity of the replisome during the DnaN dwell time on chromosomes of
~3 min (78.4 ± 0.5% (±SD) of foci; Fig. S3G). We hypothesize that this is a consequence of
delayed decatenation by TopoIV in the absence of MukBEF (Zawadzki et al., 2015), since
lagging strand templates can only be segregated from the leading strands once
decatenation has occurred.
Finally, a dynamin-like protein CrfC (aka YjdA) has been proposed to bind β$_2$-clamps
and tether the nascent strands of sister-chromosomes together (Ozaki et al., 2013).
However, upon deletion of *crfC*, we observed no changes to DnaN localization along the
long cell axis, or any decrease in the frequency of the *L3-R3-L3-R3* configuration (Fig. S3G,
H). This result indicates that CrfC is not necessary for WT chromosome conformation and
segregation.
252
Ancestral DNA strands are preferentially retained at older cell poles

Previously it has been hypothesized that a symmetrical segregation of lagging strands to the
cell center leads to sister chromosomes' translational symmetry and, in consequence, the
ancestral template DNA strand is not randomly segregated to daughter cells over

subsequent generations (discussed in (Toro and Shapiro, 2010)). To address this theory, we
have developed a novel pulse-chase assay. It allowed us to visualize relative age of DNA
strands between sister chromosomes and relate their position to the older pole in individual
cells without the need for cell synchronization or tracking (Fig. 4A).
The assay comprises pulse labelling of newly replicated DNA and identifying relative
pole age by chemoreceptor accumulation at cell poles. The newly synthesized DNA was
labelled by a 15 min EdU (5-Ethynyl-2'-deoxyuridine) pulse, after which cells were washed,
and allowed to grow for 3 h (generation time ~150 min). To avoid EdU-mediated growth
defects, thymidine was added to the medium to outcompete EdU. We observed no
detrimental effects on growth rate or cell size from the low concentration of EdU used in
the pulse (Fig. S4). After the growth period, which was longer than a single generation time,
most cells have completed an additional round of replication resulting in only one of the
two sister chromosomes remaining EdU-labelled (Fig. 4A). Cells were fixed and EdU was
visualized by click-chemistry using Alexa 488 azide. In cells with segregated chromosomes
just before division, the chromosome with new strand was fluorescently labeled, while the
one with the ancestral strand was not.
To identify the older cell pole, we exploited the fact that the serine chemoreceptor,
Tsr, accumulates near-linearly with time at the cell poles. Hence the older pole can be
distinguished from the new pole by a higher quantity of fluorescently labeled Tsr (Ping et al.,
2008). A functional HaloTag fusion of the endogenous *tsr* gene was labeled with TMR dye,
and nucleoids were stained by DAPI (Fig. 4B). This allowed us to determine if the older
strand chromosome was segregated towards older or newer pole in each cell (Fig. 4C). We
limited analysis only to cells with segregated chromosomes (i.e. two visibly separate
nucleoid regions) only one of which displayed EdU fluorescence. This does not bias the
analysis towards a specific pole. In a time-lapse control experiment we confirmed that Tsr-
mYpet forms higher intensity foci at the older pole in 99.2 ± 0.5% (±SD) of cells.
We observed that 71.3 ± 3.9% (±SD) of WT cells contained EdU foci in the
chromosome closer to the new pole (Fig. 5A). Because EdU was incorporated into the new
template strand, this indicates that the ancestral strand is preferentially retained at the
older pole. The result deviates significantly from random retention, where the older pole
would have a 50% chance of inheriting either strand (binomial two-tailed test p-value < 10^{-5}
^{5}). We also compared the dispersion (SD) of our data to a binomial distribution with
different sample sizes to estimate reliability of our experiment (Fig. 5B). We found excellent
agreement showing that our measurements are robust for the given sample size, with no
additional noise sources, and increasing sample size would give diminishing returns.
How is ancestral strand retention related to chromosome organization? For this, we
tested the contributions of MukBEF and MatP to ancestral strand retention. Upon deletion
of *mukB*, we observed a random segregation of the ancestral strand (48.5 ± 3.8% (±SD), Fig.
5A), indicating a clear requirement for functional MukBEF for ancestral strand retention at
older poles. Deletion of *matP* also abolished the preferential segregation of the ancestral
strand (46.2 ± 1.1% (±SD), Fig. 5A). While MatP has not been implicated in early
chromosome segregation, when the segregation pattern(s) emerge, the influence of MatP
on MukBEF action is crucial as it prevents longitudinal chromosome rotation (Fig. 2H), which
would disrupt ancestral-strand-older-pole association. MatP-*matS* also interacts with the

divisome through ZapB and this interaction has been proposed to partially anchor *ter* to the inner cell membrane (Espéli et al., 2012). This interaction could plausibly contribute to the ancestral strand retention by anchoring the chromosome and thereby preventing chromosome rotation. However, upon replacing the native *matP* with a non-divisome interacting *matPΔC20* mutant or deleting *zapB*, we did not observe any difference to WT with regard to ancestral strand retention (71.0 ± 3.1% and 71.8 ± 1.0%, respectively (±SD); Fig. 5A). This confirms that the loss of ancestral strand retention in *ΔmatP* cells is related to the proposed longitudinal rotation of the chromosome over generations.

Finally, since MukBEF and MatP have co-evolved with a group of proteins (including Dam and SeqA) that are related to Dam DNA methyltransferase activity (Brézellec et al., 2006), we tested the influence of these proteins in the ancestral strand retention. Dam methylates adenines in the sequence GATC, which transiently distinguishes strands after replication because of delayed methylation of the newly replicated strands. Prior to Dam methylation, SeqA binds to hemimethylated GATC sites, negatively regulating replication initiation and possibly contributing to chromosome segregation (reviewed in (Waldminghaus and Skarstad, 2009)). Deletion of either *dam* or *seqA* did not influence ancestral strand retention at older poles (69.9 ± 1.5% and 67.6 ± 3.3%, respectively (±SD); Fig. 5A), indicating GATC methylation patterns do not affect the observed asymmetry and consequently, overall *L-R* chromosome organization.

Discussion

Our results demonstrate how MukBEF axial cores direct nucleoid organization and non-random segregation of sister chromosomes in *E. coli*. This work extends that describing chromosome organization by MukBEF axial cores (Mäkelä and Sherratt, 2020), and uncovers important principles of nucleoid organization and replication-segregation. The main features of nucleoid organization are illustrated in Fig. 6 and outlined below.

1. MukBEF axial cores, linearized by *matS*-MatP-mediated depletion of MukBEF from *ter*, are required for *left-oriC-right* chromosome locus organization along the long cell axis.
2. The linearity of axial cores prevents longitudinal nucleoid rotation, thereby retaining the *left-oriC-right* chromosome orientation over generations.
3. Translational symmetry of the sister chromosomes results from the interplay of axial cores and segregation of lagging strand towards the cell center.
4. The ancestral DNA strand is preferentially segregated to the older pole cell over generations, a process dependent on MukBEF and MatP.
5. Absence of MukBEF leads to anucleate cell formation predominantly at the newer mother cell pole, largely because of failure to segregate newly replicated origins from the vicinity of the older pole.

Our analyses provide molecular mechanisms underlying the *E. coli* chromosome organization and segregation, and complement previous studies that have rigorously quantified the nucleoid dynamics in mechanical terms (Cass et al., 2016; Fisher et al., 2013;

Fritsche et al., 2012). The model succinctly explains previous observations and provide a conceptual foundation for understanding how nucleoid conformation and dynamics shape the subcellular organization.

E. coli chromosome organization

Stiff nucleoid 'bundles' that are radially confined by cell dimensions and exhibit a contour length of the scale of cell dimensions were characterized in live-imaging studies of WT *E. coli* (Fisher et al., 2013). Bundles were also identified in cells with increased volume, which allowed visualisation of non-replicating toroidal chromosomes (Wu et al., 2019). We propose that MukBEF axial cores correspond to these bundles; the duality of the bundles observed in WT cells likely relate to the bimodal nature of axial cores in non-replicating MatP$^+$ cells. We hypothesize that individual dimer of dimer MukBEF complexes within axial cores form DNA loops using energy from ATP, most likely through progressive loop enlargement (Davidson et al., 2019; Ganji et al., 2018; Goloborodko et al., 2016). Unloading and re-loading of MukBEF complexes will impact nucleoid spatio-temporal dynamics and could be dependent on intra-chromosomal stress (Fisher et al., 2013). The correlative timescale of bundle dynamics is of the order of a minute, similar to the dwell time of individual MukBEF complexes on the chromosome (Badrinarayanan et al., 2012; Fisher et al., 2013; Mäkelä and Sherratt, 2020). MukBEF forms a linear axial core to the chromosome by compacting the chromosome lengthwise outside of *ter*, where *matS* bound MatP displaces MukBEF (Mäkelä and Sherratt, 2020). The absence of MatP leads to formation of circular symmetric MukBEF axial cores. Theoretical studies concluded that lengthwise compaction of the chromosome (by MukBEF) would maintain the linear organization and forms rigid chromosome bundles (Marko and Siggia, 1997). We propose that the linearity of the axial core reduces the dimensions of free chromosome movement, thereby preventing longitudinal chromosome rotation and therefore explaining how chromosome loci can be robustly positioned along the long cell axis in a colinear manner outside of *ter* (Fig. 6).

Previous attempts to explain chromosome locus positioning by a randomly oriented polymer, or by transcription factor-mediated DNA loops fail to explain nucleoid organization in Δ*mukB* or Δ*matP* cells (Fritsche et al., 2012; Jun and Mulder, 2006). The presence of MukBEF axial cores explains how loss of a flexible *ter* region in the absence of MatP results in chromosome arms being closer together, as shown here. A flexible *ter* region might be required for efficient chromosome segregation during fast growth as Δ*matP* cells exhibit more frequent anucleate cell production than MatP$^+$ cells (Mercier et al., 2008). Broadly, we propose that the rigid linear axial core removes the requirement for membrane anchoring to orient and/or position the chromosome and could be a requirement for robust segregation during overlapping replication cycles. Membrane tethering is generally found in organisms in which MukBEF has been replaced by Smc-ScpAB complexes and which carry a *parABS* segregation system; e.g. through PopZ in *C. crescentus* (Ebersbach et al., 2008), HubP in *V. cholera* (Yamaichi et al., 2012), and RacA/DivIVA in sporulating *B. subtilis* (Ben-Yehuda et al., 2003; Wu and Errington, 2003). Membrane anchoring typically uses ParB bound to *oriC*-proximal *parS* sites as an intermediary. Intriguingly, some bacteria, such as *V. cholera* or *P. aeruginosa*, not only encode MukBEF/MksBEF, but also specify a *parABS*

387   system (David et al., 2014; Vallet-Gely and Boccard, 2013). Whether organisms that encode
388   MukBEF orthologs, but not typical Smc-ScpAB complexes, and which lack ParABS systems,
389   generally have life cycles that encompass overlapping replication cycles, similar to *E. coli*,
390   remains to be determined.
391
Sister chromosome replication and segregation
Chromosome organization specified by the MukBEF axial core directs the translational
symmetric (*L-R-L-R*) segregation of sister chromosomes during replication (Fig. 6), which
alone is insufficient for this organization (Fig. 6). Lengthwise compaction of newly replicated
DNA promotes individualization of sister chromosomes through excluded volume
interactions and by maximization of conformational entropy that leads to repulsion
between sister chromosomes (Goloborodko et al., 2016; Marko and Siggia, 1997). The
presence of the *L-R-L-R* segregation pattern of sister chromosomes in both WT and Δ*matP*
cells indicates that the pre-replication organization of the chromosome, where chromosome
arms are directed into separate cell halves only in WT cells, is not required for establishing
this pattern. While symmetrically lengthwise compacted chromosomes in Δ*matP* cells (Fig.
2A)(Mäkelä and Sherratt, 2020) are prone to change *L-R* orientation in non-replicating cells,
bi-loped replication intermediates prevent rotation during late replication (Fig. 6).
405   The relationship between *L-R-L-R* chromosome organization and symmetrical
segregation of leading/lagging strands has been discussed previously in relation to the
observed leading strand spatial segregation pattern (Toro and Shapiro, 2010; White et al.,
2008). Consistent with this, we observed accumulation of β_2-clamps, present primarily on
lagging strands, towards cell centers of replicating cells, when compared to both DNA
polymerase III and helicase localization. Differential positioning of the replisome and β_2
clamps resolves the conundrum that emerged from studies that favoured a model of a
single replication 'factory' containing two replisomes, based on clamp labelling (Mangiameli
et al., 2017). The results support the model of independent tracking of the two often
spatially separated replisomes in cells undergoing a single round of replication (Japaridze et
al., 2020; Reyes-Lamothe et al., 2008). A dynamin-like protein YjdA (aka CrfC), a possible
candidate for directing β_2-clamps to the cell centre, was not required for this action.
Because clamp localization at the cell center is dependent on the formation of linear axial
cores, we hypothesize that MukBEF could plausibly differentiate the strands, e.g. leaving
lagging strands less compacted (Fig. 6). However, the different contributions of MukBEF
activity are difficult to delineate, as in the absence of MukBEF, cohesion time is increased
and *L-R* organization prior to replication is impaired (Fig. 1).
422   Our results also show that the lifetime of individual chromosome-associated clamps
(estimated to be ~3 min (Moolman et al., 2014)) must be longer than the daughter
chromosome cohesion time for chromosomal regions outside of *oriC* and *ter* (estimated ~14
min and ~9 min, respectively) (Nolivos et al., 2016; Wang et al., 2008). Cohesion time is
largely determined by the time required for TopoIV to remove replicative catenanes
(Nolivos et al., 2016; Wang et al., 2008; Zawadzki et al., 2015). In addition, slower *oriC*
segregation may additionally require accumulation of sufficient newly replicated DNA that
leads to abrupt separation of newly replicated sister chromosomes (Cass et al., 2016; Fisher

et al., 2013). Cohesion time is influenced by the absence of MukBEF, which promotes
TopoIV activity (Nolivos et al., 2016; Zawadzki et al., 2015), although tethering of *ter* to the
divisome through MatP-ZapB interactions may also influence cohesion time in this region
(Monterroso et al., 2019). Precise measurements of cohesion times for the rest of the
chromosome have been refractory to precise experimental determination.
435
Ancestral strand retention at the older pole

We have directly shown preferential ancestral strand retention towards the older poles in *E.*
coli orchestrated by the translational symmetry of the sister chromosomes (*L-R-L-R*), along
with efficient maintenance of chromosome orientation over generations. Intriguingly, loss of
preferential segregation in the absence of MukBEF is correlated with longitudinal *oriC-ter*
chromosome organization along the long cell axis where chromosome arms are co-aligned,
similar to *C. crescentus* and sporulating *B. subtilis*. However, as *E. coli* lacks the properties of
cell differentiation, development and regeneration of a multicellular organism, it is not clear
why it has evolved a chromosome organization that preferentially segregates the ancestral
DNA strand to the older pole cell. *E. coli* older pole containing cells exhibit a constant
growth rate for hundreds of generations (Wang et al., 2010). However, the death rate was
found to increase with replicative cell age, which was attributed to growth-independent
accumulation of protein damage (Wang et al., 2010). Increasing cellular maintenance
processes through the general stress response reduced the death rate while its absence
increased it (Yang et al., 2019). The older pole accumulates more membrane proteins (e.g.
chemoreceptors, efflux pumps) than the new pole and in fluctuating or poor environment,
these can significantly contribute to cell growth (Bergmiller et al., 2017; Ping et al., 2008).
For example, the main multidrug efflux pump of *E. coli*, AcrAB-TolC, exhibits a strong
partitioning bias for the older cell poles (Bergmiller et al., 2017). Consequently, older pole
cells display increased efflux activity relative to new cell pole daughters giving the older pole
cell a growth advantage under subinhibitory antibiotic concentrations and possibly
protection against other toxic compounds. AcrAB-TolC pump activity is also required for
acquiring a resistance gene from mobile genetic elements in the presence of antibiotics, as
it reduces antibiotics concentrations inside the cell (Nolivos et al., 2019). Ancestral strand
retention at older pole cells could therefore be beneficial to structured growth of *E. coli* in
colonies where younger bacteria progress to a new terrain while the older ones stay closer
to the colony center. Finally, older strand retention could simply be an evolutionary by-
product of the *left-oriC-right* chromosome organization where chromosome orientation is
retained. Nevertheless, since ancestral strand retention occurs in only ~70% of older-pole
cells, this gives opportunities for selection in fluctuating or harmful environments
independent of whether older or newer cell pole cells thrive better.

Fig. 1. Anucleate cell formation in absence of MukB is biased towards mother cell newer poles (**A**) Schematic of mother machine microfluidics device and representative cells in a channel. *ΔmukB* cells contain *ori1* and *ter3* FROS markers and a segmentation marker (grey). A non-growing anucleate cell lacking FROS markers is indicated. Scale bar: 1 μm. (**B**) *ori1* and *ter3* localization as a function of cell length in *ΔmukB* cells. Sample numbers with different cell lengths are normalized. 221057 cells. (**C**) Number of *ori1* foci prior to anucleate (one of the daughters is anucleate; 2444 cells) and normal cell division (10468 cells). Two-sample t-test between mean *ori1* numbers prior to anucleate and normal division p-value < 10^{-5}. (**D**) Percentage of anucleate cells forming at a mother cell's old and newer poles (2269 divisions). Percentage of anucleate cell divisions also shown (14392 divisions). All data from 3 repeats.

479
480
481

Fig. 2. MukBEF and MatP action generates and retains *left-oriC-right E. coli* chromosome organization. (**A**) *E. coli* chromosome circular map with *ori1*, *ter3*, *L3*, and *R3* loci illustrates uniform MukBEF occupancy except for 800 kbp *ter*, from which *matS* bound MatP displaces MukBEF. A folded chromosome conformation by linear MukBEF axial cores is shown with and without MatP (for more details see (Mäkelä and Sherratt, 2020). (**B**) Representative images of WT, *ΔmukB* and *ΔmatP* cells with *ori1* and *ter3*, or *L3* and *R3* FROS markers. Note an atypical *R3-L3-L3-R3* configuration in WT (white arrow) in comparison to standard *L3-R3-L3-R3*. Scale bars: 1 μm. (**C**) *L3* and *R3* localizations and (**D**) *ori1* localizations along the long cell axis as a function of cell length in WT (*L3-R3* 57509 cells, *ori1* 42612 cells), *ΔmukB* (*L3-R3* 27984 cells, *ori1* 54820 cells) and *ΔmatP* (*L3-R3* 46679 cells, *ori1* 51350 cells). Sample numbers with different cell lengths are normalized. Cells are oriented to place *L3* more towards the negative pole (towards figure bottom) or, in the *ori1* data, *ter3* is oriented more towards the negative pole (see Fig. S2). White lines denote cell borders. (**E**) Overlay of *ori1* and *L3-R3* localization data in WT and *ΔmatP* from (C) and (D). (**F**) Distance between *L3* and *R3* markers in WT (47376 cells), *ΔmukB* (15615 cells) and *ΔmatP* (41625 cells) in single *L3* and *R3* focus cells. Mean and dispersion (SD) between cells are shown for each distribution. * and ** denote two-sample t-test of *L3-R3* distances between WT and *ΔmukB* (p-value 0.0081) and WT and *ΔmatP* (p-value 5×10^{-4}), respectively. (**G**) (left) Percentage of cells with *L3* and *R3* in opposite cell halves in single *L3* and *R3* focus cells (WT 47376 cells, *ΔmukB* 15615 cells, *ΔmatP* 41625 cells). (right) Percentage of cells with *L3-R3-L3-R3* (or *R3-L3-R3-L3*) configuration (versus *L3-R3-R3-L3* or *R3-L3-L3-R3*) in double *L3* and *R3* focus cells (WT 10352 cells, *ΔmukB* 2535 cells, *ΔmatP* 6297 cells). Scale bars: 1 μm. (**H**) Percentage of cells retaining *L3-R3-L3-R3* orientation (versus flipping to *R3-L3-R3-L3*) from a mother cell to a daughter cell in WT (859 pairs) and *ΔmatP* (1054 pairs). Scale bars: 1 μm. Data from 3 repeats of each experiment.

Fig. 3. DnaN visualizes the lagging strands during replication. (**A**) Schematic of
accumulation of β₂-clamps (DnaN) on the lagging strand and its template during replication
(Moolman et al., 2014). The DNA polymerase ε-subunit (DnaQ) marks the location of the
replisome. (**B**) Representative images of WT cells with fluorescently labeled DnaN and DnaQ.
Scale bars: 1 μm. (**C**) DnaQ and DnaN localization in WT cells as a function of cell length
(37720 cells). White lines denote cell borders. Shaded areas denote intermediate cell
lengths for localization data in (E). (**D**) Distance from a DnaQ focus to the closest DnaN
focus. DnaQ and DnaN colocalize in 58.8 ± 5% (±SD) of focus pairs (38855 pairs) as defined
by a threshold (red lines) below which two proteins colocalize (dictated by a diffraction limit
of 300 nm). Inset shows the same data as a cumulative distribution. Same data as in (C). (**E**)
DnaQ or DnaN localization with intermediate cell lengths (3.3-3.7 μm) in WT (DnaN 7104,
DnaQ 8006 spots), *ΔmatP* (DnaN 11925, DnaQ 8025 spots) and *ΔmukB* (DnaN 5060, DnaQ
4205 spots) cells (see C, F, G). Full width at half maximum (FWHM) (±SD) of the distribution
in WT: DnaN 0.67 ± 0.06 μm, DnaQ 1.67 ± 0.08 μm; and in *ΔmatP*: DnaN 1.85 ± 0.04 μm,
DnaQ 1.14 ± 0.14 μm. Grey areas denote cell poles. (**F**) DnaQ and DnaN localization in
ΔmatP cells (51956 cells) and in (**G**) *ΔmukB* cells (22902 cells) as a function of cell length.
White lines denote cell borders. Data from 3 repeats in all analyses.

Fig. 4. Visualization of ancestral DNA strand retention in *E. coli*. (A) Ancestral DNA strand retention following an EdU pulse and the subsequent growth shown with cell cycle stages. After the 2nd round of replication only one of the chromosomes inherits the EdU label. Note that only a part of the chromosome is labelled with EdU. The 15 min EdU pulse and growth period are also shown relative to a schematic of cell cycle stages (B, C and D periods, generation time ~150 min). **(B)** Representative EdU^{Alexa488}, Tsr^{TMR} and DAPI images of a cell at stage (ii) (see (A)) after the EdU pulse. Note that each chromosome has two EdU foci because pulse-labeled chromosome arms are separated. Scale bars: 1 μm **(C)** Representative EdU^{Alexa488}, Tsr^{TMR} and DAPI images after the complete pulse-chase protocol (stage (iv), see (A)). The older pole is indicated with an arrow. Scale bars: 1 μm. **(D)** Accuracy of the older pole classification using Tsr prior to cell division. Shaded areas denote SD. Data from 2505 cells and 3 repeats.

Fig. 5. Preferential retention of the ancestral strand at older cell poles requires functional MukBEF and MatP. Percentage of ancestral strands retained at the older pole in WT (988 cells), *ΔmukB* (427 cells)*, ΔmatP* (1050 cells), non-divisome interacting *matPΔC20* mutant (1617 cells), *ΔzapB* (969 cells), *Δdam* (717 cells) and *ΔseqA* (473 cells). Dashed line shows random retention. p-value from two-proportion two-tailed z-test was used to test if binomial distributions significantly differ from WT; indicated by n.s. (> 0.01, non-significant) and * (< 0.01, significant) (p-values <10^{-5}, <10^{-5}, 0.51, 0.83, 0.26, and 0.03, respectively). Shaded areas denote SD. Data from 3 repeats.

Fig. 6. Chromosome organization and segregation by MukBEF and MatP. (**A**) MukBEF dimer of dimer complexes form DNA loops dependent on ATP hydrolysis. Ubiquitous loop formation outside of *ter* compacts the chromosome lengthwise forming a linear axial core to the chromosome. Axial cores form stiff linear nucleoid bundles that localize chromosomal loci linearly along the long cell axis and maintain chromosome orientation by preventing longitudinal cell axis rotation. (**B**) Absence of functional MukBEF increases the effective contour length of chromosomes, leading to mis-localization of chromosome loci and loss of *left-oriC-right* organization of the chromosome. (**C**) MukBEF forms a complete circular axial core to the chromosome in the absence of MatP. Consequently, chromosome arms cannot be efficiently directed to opposite cell halves, *left-oriC-right* organization in non-replicating cells is impaired and chromosomes rotate in the longitudinal cell axis. During replication and prior to division, WT (**D**) and *ΔmatP* cells (**E**) exhibit translational symmetric (*L-R-L-R*) segregation of sister chromosomes. This is accompanied by the symmetric segregation of lagging strands and their templates during replication, as visualized by accumulation of β_2-clamps. Relative FROS marker, replisome and β_2-clamps localizations are derived from the data generated here.

Methods

Bacterial strains and growth conditions

Bacterial strains and primers are listed in Table S1 and S2, respectively. All strains were derivatives of *E. coli* K12 AB1157 (Bachmann, 1996). *kan*, *cat*, *gen*, and *hyg* refer to insertions conferring resistance to kanamycin (Km[r]), chloramphenicol (Cm[r]), gentamycin (Gm[r]) and hygromycin B (Hyg[r]), respectively. The insertions are flanked by Flp site-specific recombination sites (*frt*) that allow removing the resistance gene using Flp recombinase from plasmid pCP20 (Datsenko and Wanner, 2000). *tsr-HaloTag-kan* and *tsr-mYpet-kan* were inserted into the native *tsr* chromosomal locus using λ-red recombination (Datsenko and Wanner, 2000). The generated gene loci were transferred by phage P1 transduction to AB1157 yielding strains JM122 (*tsr-HaloTag-kan*) and JM133 (*tsr-mYpet-kan*). Deletion strains of JM122 were constructed by P1 transduction, first removing the *kan* resistance gene using Flp recombinase. *L3-R3* deletion strains were constructed from RRL66 using P1 transduction. The microfluidics strain (JM09) was constructed from RRL189 by introducing *Δflhd-kan* and *ΔmukB-kan* by consecutive rounds of P1 transduction and Flp recombination. GFPmut2 cell marker was inserted at an *attTn7* site by a plasmid transformation as described in (McKenzie and Craig, 2006). The DnaQ and DnaN labeled strain (JM141) was constructed from RRL388 using P1 transduction from RRL36. Deletion strains of JM141 were constructed by P1 transduction, first removing the *kan* resistance gene using Flp recombinase. JM142 and JM143 were constructed by P1 transduction from JW4070. All genetic modifications were verified by PCR and/or sequencing and behavior in quantitative imaging. *mukB* deletions were verified by temperature-sensitivity in rich media, as described in (Nolivos et al., 2016).

Cells were grown in M9 minimal medium supplemented with 0.2% (v/v) glycerol, 2 μg ml^{-1} thiamine, and required amino acids (threonine, leucine, proline, histidine and arginine; 0.1 mg ml^{-1}) at 30 °C. For microscopy, cells were grown overnight, diluted 1000-fold and grown to an A_{600} of ~0.1. Cells were then pelleted, spotted onto an M9 glycerol 1% (w/v) agarose pad on a slide and covered by a coverslip. In mother machine microfluidics experiments, cells were first grown in M9 minimal medium with 0.2% (v/v) glycerol at 30 °C (as above), and then placed inside the microfluidics device, when the media was changed to M9 minimal medium supplemented with 0.2% (v/v) glucose, 2 μg ml^{-1} thiamine, MEM amino acids (Gibco, #11130-036), 0.1 mg ml^{-1} proline, and 0.85 mg ml^{-1} Pluronic F127 (Sigma-Aldrich, P2443), and the temperature was set to 37 °C.

EdU pulse labelling

Cells grown until A_{600} of ~0.1 were labelled with 10 μM EdU (5-Ethynyl-2′-deoxyuridine, Thermofisher, C10337) for 15 min after which cells were washed, introduced to fresh media containing 60 μg/ml thymidine and allowed to grow for 3 h. Following this, cells were fixed with 4% PFA (v/v) for 30 min and permealized with 0.5% Triton X-100 (v/v) for 30 min. EdU click-chemistry reaction was conducted following the instructions (Thermofisher, #C10337) using Alexa 488 azide in a final volume of 50 μl for 30 min at room temperature, followed by washing. Cells were then labelled with TMR HaloTag ligand as in (Banaz et al., 2019). Briefly, cells were incubated with 2 μM TMR ligand for 30 min and washed several times. Finally,

nucleoids were labelled with 1 μg/ml DAPI for 15 min and washed, after which the cells were
ready for imaging.

Epifluorescence microscopy
Fluorescence images were acquired on an inverted fluorescence microscope (Ti-E, Nikon)
equipped with a perfect focus system, a 100× NA 1.4 oil immersion objective, a motorized
stage, an sCMOS camera (Orca Flash 4, Hamamatsu), and a temperature chamber (Okolabs).
Exposure times were 300 ms for TMR, Alexa 488 and mCherry, mYpet; 150 ms for mCerulean,
and 100 ms for DAPI using an LED excitation source (Lumencor SpectraX). Phase contrast
images were collected for cell segmentation. Microscopy data was collected automatically
from the sample area. Time-lapse images were collected every 10 min for 3 h with a shorter
exposure time of 150 ms, except fluorescence images from the microfluidics device were
collected every 5 min.

Microfluidic devices
The microfluidic single-cell imaging device ("mother machine") was prepared as in (Uphoff,
2018). The device was designed using Autodesk AutoCAD software. The dimensions of the
cell channels were 1.2 μm x 1.2 μm x 20 μm and the media flow channels were 100 μm x 25
μm. The structures were fabricated on a silicon wafer (Kavli Nanolab, Delft University)
(Moolman et al., 2013) and a negative polydimethylsiloxane (PDMS) mold was created from
the silicon wafer using a 5:1 mixture of monomer and curing agent (Dow Corning Sylgard
184 Kit). After removing air bubbles using vacuum, the chip was cured at 65°C for 1.5 hours.
The mould was treated with Trichloro(1H,1H,2H,2H perfluorooctyl)silane (Sigma) in vacuum
overnight. The PDMS device was generated from the negative mold using a 10:1 mixture of
monomer and curing agent and cured at 65°C for 1.5 hours. Media flow holes were punched
through the device with 0.75 mm diameter. Cover slips were cleaned by sonication in
acetone for 20 min, washing with dH2O, sonication in isopropanol for 20 min, and dried
with nitrogen. The PDMS device was washed with isopropanol and dried with nitrogen. The
device and a cover slip were bonded using air plasma (Plasma Etch PE-50) and placed in an
oven at 95°C for 30 min. Cells were pipetted into the device and the device was centrifuged
at 4000 rpm for 10 min to place cells into the channels. The media supplemented by
Pluronic F127 was fed into the device through silicon tubing (Tygon ND 100-80 microbore,
VWR) using a motorized infusion pump (New Era Pump Systems). Initially, a high flow rate of
1.5 ml/hr was applied to flush the excess cells and then lowered to 0.5 ml/hr. After this, cells
were allowed to grow for ~2 h before starting the time-lapse imaging.

Image analysis
Cell based information, including cell outlines, lineages, pole ages, per pixel fluorescence
intensities, and fluorescent marker localization, was extracted using SuperSegger (Stylianidou
et al., 2016) in MATLAB (MathWorks). SuperSegger uses an image-curvature method to
identify foci to avoid the identification of false positive foci due to background intensity from
cytoplasmic fluorescence and uses a gaussian fit to find the subpixel resolution location of
foci. Focus quality is determined by a combination of intensity and fitting parameters and bad
quality foci were filtered out. A threshold value was confirmed by visual inspection and the

same threshold was used for all compared data sets. The channels were aligned prior to analysis.

Mother machine analysis

From the lineage data, cells were classified as 'normal' growing cells, anucleate cells, mothers of anucleate cells and sisters of anucleate cells. Cells that disappeared early or didn't have a tracked lineage were excluded from the analysis. Anucleate cells were considered as cells that didn't divide, didn't elongate and lacked an *ori1* marker present, while its sister cell elongated and divided normally, and had *ori1* marker(s). If neither of the sister cells divided normally, cells were excluded from analysis. The older pole of the anucleate cell was traced back ≥2 generations to determine whether the anucleate cell formed on the older or newer pole of the mother. The cell size at birth (Fig. S1B) was determined at the first frame of each cell and the number of *ori1* foci prior to division (Fig. 1C) at the last frame of a cell.

Fluorescent marker localization

For *ori1, ter3, L3* and *R3* markers, intensity profiles with different cell lengths were normalized, as the expectation is that a cell will have at least a single focus at all times. As cell orientation is random relative to the pole age, cells were oriented to place *L3* more towards the negative pole than *R3* and, in the *ori1* data, *ter3* was oriented more towards the negative pole. To determine flipping frequency of *L3-R3-L3-R3* markers from time-lapse imaging, first mother cells that contained at *L3-R3-L3-R3* or *R3-L3-R3-L3* were identified. Next, their daughter cells with *L3-R3-L3-R3* or *R3-L3-R3-L3* were identified. The angle (Fig. S2) between vectors pointing from the more polar *L3* to the more polar *R3* was calculated between mother and daughter cells. If the angle exceeded 90°, the chromosome orientation was considered flipped. To measure width of a unimodal distribution and to avoid inaccuracy from binning the data, the data was fitted by a kernel distribution in MATLAB and full width at half maximum (FWHM) was calculated from the fitted distribution.

EdU pulse labelling

A functional HaloTag fusion of the endogenous *tsr* gene was used in the EdU labeling, as click-chemistry reaction conditions are detrimental to conventional fluorescent proteins. To measure EdU association with the older pole, the following criteria were used to select cells from an asynchronous cell population (see Fig. S4). First, Otsu's thresholding (Otsu, 1979) was used to segment nucleoid area(s) from the cellular background and only cells that have two separate, large-enough nucleoid areas were analysed. Second, a cell must exhibit a clear difference in polar Tsr intensity. The center line of a cell was extracted to find coordinates of cell poles by fitting a cell mask to a second-order curve. The intersection of the cell mask border and the curve was used to define cell poles. Median Tsr intensity of the cell area was subtracted from all Tsr pixel intensities and a sum of 9 brightest pixels from each pole were used to quantify the pole intensity. To minimize effects of noise and discrete pixel size in segmentation, only cells with >1.5-fold difference in polar Tsr intensity were analysed. The pole that had a higher intensity of Tsr was designated as the older pole. Third, only one of the nucleoids must be labelled by EdU. EdU with short incorporation times appear as distinct foci (see Fig. 4 and Fig. S4). The foci below a fixed threshold for the score were discarded. The foci

701 were mapped to the nucleoids by projecting coordinates of both on the center line of the cell.
702 With these criteria, the processed microscopy data from SuperSegger was automatically
703 analysed to extract the result of EdU association with older cell pole. To avoid segmentation
704 errors of the cell area, correct cell segmentation was visually inspected and inaccurately
705 segmented cells were removed.
706
Tsr time-lapse
The accuracy of Tsr-based identification of the old cell pole in our growth conditions was
estimated by tracking cells with a functional mYpet fusion to the endogenous *tsr* gene over
generations under a microscope. Only cells that both were born and divided during the
time-lapse were analysed. Tsr intensity at each pole was calculated with same criteria as in
the EdU experiment. The accuracy of Tsr as the older pole marker was determined for each
frame separately by comparing results between Tsr intensity analysis and lineage tracking.
The accuracy (Fig. 4D) was shown for the last frame prior to division to mimic the EdU
experiment where only cells with segregated chromosomes were analysed.
716

Acknowledgements

718 We thank other members of the Sherratt and Uphoff group, and Katarzyna Ginda-Mäkelä
719 for insightful discussions. The research was supported by a Wellcome Investigator Award to
720 DJS (200782/Z/16/Z). SU was in receipt of a Henry Dale-Wellcome Fellowship.

Author contributions

723 JM and DJS conceived the project and directed it. JM undertook and analyzed experiments.
724 SU developed the microfluidics, advised on its use and was generally involved in project
725 discussions. JM, SU and DJS wrote the paper.
726

Declaration of Interests

728 The authors declare no competing interests.
729

Data and code availability

731 All data required to understand and assess the conclusions of the research are available in
732 the main text and supplementary materials. All materials and codes are available upon
733 reasonable request.
734

References

Bachmann, B.J. (1996). Derivation and Genotypes of Some Mutant Derivatives of *Escherichia coli* K-12. Escherichia Coli Salmonella Typhimurium Cell. Mol. Biol. *ASM*, 2460–2488.

Badrinarayanan, A., Reyes-Lamothe, R., Uphoff, S., Leake, M.C., and Sherratt, D.J. (2012). In Vivo Architecture and Action of Bacterial Structural Maintenance of Chromosome Proteins. Science *338*, 528–531.

Badrinarayanan, A., Le, T.B.K., and Laub, M.T. (2015). Bacterial Chromosome Organization and Segregation. Annu. Rev. Cell Dev. Biol. *31*, 171–199.

Banaz, N., Mäkelä, J., and Uphoff, S. (2019). Choosing the right label for single-molecule tracking in live bacteria: Side-by-side comparison of photoactivatable fluorescent protein and Halo tag dyes. J. Phys. D. Appl. Phys. *52*, 064002.

Ben-Yehuda, S., Rudner, D.Z., and Losick, R. (2003). RacA, a bacterial protein that anchors chromosomes to the cell poles. Science *299*, 532–536.

Bergmiller, T., Andersson, A.M.C., Tomasek, K., Balleza, E., Kiviet, D.J., Hauschild, R., Tkačik, G., and Guet, C.C. (2017). Biased partitioning of the multidrug efflux pump AcrAB-TolC underlies long-lived phenotypic heterogeneity. Science *356*, 311–315.

Brézellec, P., Hoebeke, M., Hiet, M.S., Pasek, S., and Ferat, J.L. (2006). DomainSieve: A protein domain-based screen that led to the identification of dam-associated genes with potential link to DNA maintenance. Bioinformatics *22*, 1935–1941.

Cairns, J. (1975). Mutation selection and the natural history of cancer. Nature *255*, 197–200.

Cass, J.A., Kuwada, N.J., Traxler, B., and Wiggins, P.A. (2016). *Escherichia coli* Chromosomal Loci Segregate from Midcell with Universal Dynamics. Biophys. J. *110*, 2597–2609.

Danilova, O., Reyes-Lamothe, R., Pinskaya, M., Sherratt, D., and Possoz, C. (2007). MukB colocalizes with the oriC region and is required for organization of the two *Escherichia coli* chromosome arms into separate cell halves. Mol. Microbiol. *65*, 1485–1492.

Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. Proc Natl Acad Sci U S A *97*, 6640– 6645.

David, A., Demarre, G., Muresan, L., Paly, E., Barre, F.X., and Possoz, C. (2014). The Two Cis-Acting Sites, parS1 and oriC1, Contribute to the Longitudinal Organisation of *Vibrio cholerae* Chromosome I. PLoS Genet. *10*, e1004448.

Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.-M. (2019). DNA loop extrusion by human cohesin. Science *3418*, 1–13.

Ebersbach, G., Briegel, A., Jensen, G.J., and Jacobs-Wagner, C. (2008). A Self-Associating Protein Critical for Chromosome Attachment, Division, and Polar Organization in *Caulobacter*. Cell *134*, 956–968.

Errington, J., and Wake, R.G. (1991). Chromosome strand segregation during sporulation in *Bacillus subtilis*. Mol. Microbiol. *5*, 1145–1149.

Espéli, O., Borne, R., Dupaigne, P., Thiel, A., Gigant, E., Mercier, R., and Boccard, F. (2012). A MatP-divisome interaction coordinates chromosome segregation with cell division in *E. coli*. EMBO J. *31*, 3198–3211.

Fisher, J.K., Bourniquel, A., Witz, G., Weiner, B., Prentiss, M., and Kleckner, N. (2013). Four-Dimensional Imaging of *E. coli* Nucleoid Organization and Dynamics in Living Cells. Cell *153*, 882–895.

Fogel, M.A., and Waldor, M.K. (2005). Distinct segregation dynamics of the two *Vibrio cholerae* chromosomes. Mol. Microbiol. *55*, 125–136.

Fritsche, M., Li, S., Heermann, D.W., and Wiggins, P.A. (2012). A model for *Escherichia coli* chromosome packaging supports transcription factor-induced DNA domain formation.

Nucleic Acids Res. *40*, 972–980.

Ganji, A.M., Shaltiel, I.A., Bisht, S., Kim, E., Kalichava, A., Haering, C.H., and Dekker, C. (2018).
Real-time imaging of DNA loop extrusion by condensin. Science *7831*, 1–9.

Goloborodko, A., Imakaev, M. V., Marko, J.F., and Mirny, L. (2016). Compaction and
segregation of sister chromatids via active loop extrusion. Elife *5*, 1–16.

Hiraga, S., Niki, H., Ogura, T., Ichinose, C., Mori, H., Ezaki, B., and Jaffe, A. (1989).
Chromosome partitioning in *Escherichia coli*: novel mutants producing anucleate cells. J.
Bacteriol. *171*, 1496–1505.

Japaridze, A., Gogou, C., Kerssemakers, J.W.J., Nguyen, H.M., and Dekker, C. (2020). Direct
observation of independently moving replisomes in *Escherichia coli*. Nat. Commun. 3109.

Jun, S., and Mulder, B. (2006). Entropy-driven spatial organization of highly confined
polymers: lessons for the bacterial chromosome. Proc. Natl. Acad. Sci. U. S. A. *103*, 12388–
12393.

Kuzminov, A. (2014). The precarious prokaryotic chromosome. J. Bacteriol. *196*, 1793–1806.

Lansdorp, P.M. (2007). Immortal Strands? Give Me a Break. Cell *129*, 1244–1247.

Mäkelä, J., and Sherratt, D.J. (2020). Organization of the *Escherichia coli* Chromosome by a
MukBEF Axial Core. Mol. Cell *78*, 250–260.

Mangiameli, S.M., Veit, B.T., Merrikh, H., and Wiggins, P.A. (2017). The Replisomes Remain
Spatially Proximal throughout the Cell Cycle in Bacteria. PLOS Genet. *13*, e1006582.

Marczynski, G.T., Dingwall, A., and Shapiro, L. (1990). Plasmid and chromosomal DNA
replication and partitioning during the *Caulobacter crescentus* cell cycle. J. Mol. Biol. *212*,
709–722.

Marko, J.F., and Siggia, E.D. (1997). Polymer models of meiotic and mitotic chromosomes.
Mol. Biol. Cell *8*, 2217–2231.

McKenzie, G.J., and Craig, N.L. (2006). Fast, easy and efficient: Site-specific insertion of
transgenes into Enterobacterial chromosomes using Tn7 without need for selection of the
insertion event. BMC Microbiol. *6*, 39.

Mercier, R., Petit, M.A., Schbath, S., Robin, S., El Karoui, M., Boccard, F., and Espéli, O.
(2008). The MatP/*matS* Site-Specific System Organizes the Terminus Region of the *E. coli*
Chromosome into a Macrodomain. Cell *135*, 475–485.

Monterroso, B., Zorrilla, S., Sobrinos-Sanguino, M., Robles-Ramos, M.Á., Alfonso, C.,
Söderström, B., Meiresonne, N.Y., Verheul, J., Den Blaauwen, T., and Rivas, G. (2019). The
bacterial DNA binding protein matp involved in linking the nucleoid terminal domain to the
divisome at midcell interacts with lipid membranes. MBio *10*, 1–14.

Moolman, M.C., Huang, Z., Krishnan, S.T., Kerssemakers, J.W.J., and Dekker, N.H. (2013).
Electron beam fabrication of a microfluidic device for studying submicron-scale bacteria. J.
Nanobiotechnology *11*, 12.

Moolman, M.C., Krishnan, S.T., Kerssemakers, J.W., van den Berg, A., Tulinski, P., Depken,
M., Reyes-Lamothe, R., Sherratt, D.J., and Dekker, N.H. (2014). Slow unloading leads to DNA-
bound beta2-sliding clamp accumulation in live *Escherichia coli* cells. Nat Commun *5*, 5820.

Nielsen, H.J., Ottesen, J.R., Youngren, B., Austin, S.J., and Hansen, F.G. (2006). The
Escherichia coli chromosome is organized with the left and right chromosome arms in
separate cell halves. Mol. Microbiol *62*, 331–338.

Niki, H., and Jaffe, A. (1991). The new gene mukB codes for a 177 kd protein with coiled coil
domains involved in chromosome partitioning of *E. coli*. EMBO J. *10*, 183–193.

Nolivos, S., and Sherratt, D. (2014). The bacterial chromosome: Architecture and action of
bacterial SMC and SMC-like complexes. FEMS Microbiol. Rev. *38*, 380–392.

Nolivos, S., Upton, A.L., Badrinarayanan, A., Müller, J., Zawadzka, K., Wiktor, J., Gill, A.,
Arciszewska, L., Nicolas, E., and Sherratt, D. (2016). MatP regulates the coordinated action
of topoisomerase IV and MukBEF in chromosome segregation. Nat. Commun. *7*, 10466.
Nolivos, S., Cayron, J., Dedieu, A., Page, A., Delolme, F., and Lesterlin, C. (2019). Role of
AcrAB-TolC multidrug efflux pump in drug-resistance acquisition by plasmid transfer.
Science *364*, 778–782.
Osley, M.A., and Newton, A. (1974). Chromosome segregation and development in
Caulobacter crescentus. J. Mol. Biol. *90*, 359–370.
Otsu, N. (1979). A threshold selection method from gray-level histograms. IEEE Trans. Sys.
Man. Cyber. *9*, 62–66.
Ozaki, S., Matsuda, Y., Keyamura, K., Kawakami, H., Noguchi, Y., Kasho, K., Nagata, K.,
Masuda, T., Sakiyama, Y., and Katayama, T. (2013). A replicase clamp-binding dynamin-like
protein promotes colocalization of nascent DNA strands and equipartitioning of
chromosomes in *E.coli*. Cell Rep. *4*, 985–995.
Petrushenko, Z.M., She, W., and Rybenkov, V. V. (2011). A new family of bacterial
condensins. Mol. Microbiol. *81*, 881–896.
Ping, L., Weiner, B., and Kleckner, N. (2008). Tsr-GFP accumulates linearly with time at cell
poles, and can be used to differentiate "old" versus "new" poles, in *Escherichia coli*. Mol.
Microbiol. *69*, 1427–1438.
Rando, T.A. (2007). The Immortal Strand Hypothesis: Segregation and Reconstruction. Cell
129, 1239–1243.
Reyes-Lamothe, R., Possoz, C., Danilova, O., and Sherratt, D.J. (2008). Independent
Positioning and Action of *Escherichia coli* Replisomes in Live Cells. Cell *133*, 90–102.
Soubry, N., Wang, A., and Reyes-Lamothe, R. (2019). Replisome activity slowdown after
exposure to ultraviolet light in *Escherichia coli*. Proc. Natl. Acad. Sci. U. S. A. *116*, 11747–
11753.
Stylianidou, S., Brennan, C., Nissen, S.B., Kuwada, N.J., and Wiggins, P.A. (2016).
SuperSegger: robust image segmentation, analysis and lineage tracking of bacterial cells.
Mol. Microbiol. *102*, 690–700.
Surovtsev, I. V., and Jacobs-Wagner, C. (2018). Subcellular Organization: A Critical Feature of
Bacterial Cell Replication. Cell *172*, 1271–1293.
Toro, E., and Shapiro, L. (2010). Bacterial Chromosome Organization and Segregation. Cold
Spring Harb Perspect Biol *2*, a000349.
Uhlmann, F. (2016). SMC complexes: From DNA to chromosomes. Nat. Rev. Mol. Cell Biol.
17, 399–412.
Umbarger, M.A., Toro, E., Wright, M.A., Porreca, G.J., Baù, D., Hong, S.H., Fero, M.J., Zhu,
L.J., Marti-Renom, M.A., McAdams, H.H., et al. (2011). The three-dimensional architecture
of a bacterial genome and its alteration by genetic perturbation. Mol. Cell *44*, 252–264.
Uphoff, S. (2018). Real-time dynamics of mutagenesis reveal the chronology of DNA repair
and damage tolerance responses in single cells. Proc. Natl. Acad. Sci. U. S. A. *115*, E6516–
E6525.
Vallet-Gely, I., and Boccard, F. (2013). Chromosomal Organization and Segregation in
Pseudomonas aeruginosa. PLoS Genet. *9*, e1003492.
Wakeman, J.A., Hmadcha, A., Soria, B., and McFarlane, R.J. (2012). The immortal strand
hypothesis: Still non-randomly segregating opinions. Biomol. Concepts *3*, 203–211.
Waldminghaus, T., and Skarstad, K. (2009). The *Escherichia coli* SeqA protein. Plasmid *61*,
141–150.

Wallden, M., Fange, D., Lundius, E.G., Baltekin, O., and Elf, J. (2016). The Synchronization of
Replication and Division Cycles in Individual *E. coli* Cells. Cell *166*, 729–739.
Wang, P., Robert, L., Pelletier, J., Dang, W.L., Taddei, F., Wright, A., and Jun, S. (2010).
Robust growth of *Escherichia coli*. Curr. Biol. *20*, 1099–1103.
Wang, X., Possoz, C., and Sherratt, D.J. (2005). Dancing around the divisome: asymmetric
chromosome segregation in *Escherichia coli*. Genes Dev *19*, 2367–2377.
Wang, X., Liu, X., Possoz, C., and Sherratt, D.J. (2006). The two *Escherichia coli* chromosome
arms locate to separate cell halves. Genes Dev. *20*, 1727–1731.
Wang, X., Reyes-lamothe, R., and Sherratt, D.J. (2008). Modulation of *Escherichia coli* sister
chromosome cohesion by topoisomerase. Genes Dev. 2426–2433.
Wang, X., Llopis, P.M., and Rudner, D.Z. (2013). Organization and segregation of bacterial
chromosomes. Nat. Rev. Genet. *14*, 191–203.
Wang, X., Montero Llopis, P., and Rudner, D.Z. (2014). *Bacillus subtilis* chromosome
organization oscillates between two distinct patterns. Proc. Natl. Acad. Sci. U. S. A. *2014*, 1–
6.
White, M.A., Eykelenboom, J.K., Lopez-Vernaza, M.A., Wilson, E., and Leach, D.R.F. (2008).
Non-random segregation of sister chromosomes in *Escherichia coli*. Nature *455*, 1248–1250.
Wu, L.J., and Errington, J. (2003). RacA and the Soj-Spo0J system combine to effect polar
chromosome segregation in sporulating *Bacillus subtilis*. Mol. Microbiol. *49*, 1463–1475.
Wu, F., Japaridze, A., Zheng, X., Wiktor, J., Kerssemakers, J.W.J., and Dekker, C. (2019).
Direct imaging of the circular chromosome in a live bacterium. Nat. Commun. *10*, 1–9.
Yamaichi, Y., Bruckner, R., Ringgaard, S., Möll, A., Ewen Cameron, D., Briegel, A., Jensen,
G.J., Davis, B.M., and Waldor, M.K. (2012). A multidomain hub anchors the chromosome
segregation and chemotactic machinery to the bacterial pole. Genes Dev. *26*, 2348–2360.
Yang, S., Kim, S., Kim, D.K., Jeon An, H., Bae Son, J., Hedén Gynnå, A., and Ki Lee, N. (2019).
Transcription and translation contribute to gene locus relocation to the nucleoid periphery
in E. coli. Nat. Commun. *10*, 1–12.
Zawadzki, P., Stracy, M., Ginda, K., Zawadzka, K., Lesterlin, C., Kapanidis, A.N., and Sherratt,
D.J. (2015). The Localization and Action of Topoisomerase IV in *Escherichia coli* Chromosome
Segregation Is Coordinated by the SMC Complex, MukBEF. Cell Rep. *13*, 2587–2596.
906

 Supplementary Figures

Fig. S1. (**A**) Generation time in normally dividing cells (12103 cells) and in sisters of
anucleate cells (1605 cells). Two-sample t-test of mean generation time p-value 0.2176.
Data from 3 repeats. (**B**) Difference in cell length at birth between anucleate and growing
sister cells at anucleate cell division. Black dashed line indicates symmetric division and solid
line shows a linear fit to the data. Black circles show binned mean. 2266 cell pairs from 3
repeats.

Fig. S2. (**A**) *ter3* localization along long cell axis in WT (26926 cells), Δ*mukB* (48770 cells) and
Δ*matP* (45532 cells). From same data as in Fig. 2D; *ter3* is oriented more towards the
negative pole than *ori1*. Data from 3 repeats. (**B**) Representative time-lapse images of WT
(top) and Δ*matP* (bottom) cells with *L3* and *R3* markers. (top) *L3-R3-L3-R3* orientation is
maintained over a generation while (bottom) *L3-R3-L3-R3* orientation is flipped. From *L3-R3-
L3-R3* cells, angle between vectors pointing from the more polar *L3* to the more polar *R3* is
calculated between mother cell (red arrow) and daughter cells (orange arrow) and, if the
angle exceeds 90°, the chromosome orientation is considered flipped (blue arrow). Scale
bars: 1 μm. Angle between mother and daughter cell *L3-R3-L3-R3* vectors in (**C**) WT (859
pairs) and (**D**) Δ*matP* (1054 pairs) cells. Data from 3 repeats.

Fig. S3. (**A**) Representative images of WT cells with labelled DnaN and DnaB. Scale bars: 1 µm. (**B**) DnaB and DnaN localization in WT cells as a function of cell length (16134 cells). White lines denote cell borders. (**C**) Distance from a DnaB locus to the closest DnaN locus. DnaB and DnaN colocalize in 52.9 ± 6.1% (±SD) of pairs (11714 pairs) as defined by a threshold (red lines) below which two proteins colocalize (dictated by a diffraction limit of 300 nm). Inset shows the same data as a cumulative distribution. Same data as in (B). (**D**) DnaQ (4567 spots) or DnaN (5393 spots) localization with early replication cells in WT (cell lengths 2.5-2.9 µm) (same data as in Fig. 3C). (**E**) Distance from a DnaQ locus to the closest DnaN locus in *ΔmatP* cells. DnaQ and DnaN colocalize in 50.8 ± 1.3% (±SD) of pairs (46330 pairs). Inset shows the same data as a cumulative distribution. Same data as in (Fig. 3F). (**F**) DnaQ localization as a function of cell length in WT cells in which DnaQ foci are spatially separate from DnaN (16158 cells). Same data as in Fig. 2C and D. (**G**) Distance from a DnaQ locus to the closest DnaN locus in *ΔmukB* cells. DnaQ and DnaN colocalize in 78.4 ± 0.5%

(±SD) of pairs (32603 pairs). Inset shows the same data as a cumulative distribution. Same

data as in (Fig. 3G). (**H**) DnaN localization in *ΔcrfC* cells as a function of cell length (49955

cells). Shaded areas denote intermediate cell lengths for localization data in (I). White lines

denote cell borders. (**I**) DnaN localization with intermediate cell lengths (3.3-3.7 µm) in WT

(8006 spots) and *ΔcrfC* (10691). Data from (H) and Fig. 3E. (**J**) Percentage of *ΔcrfC* cells (8393

cells) with *L3-R3-L3-R3* (or *R3-L3-R3-L3*) configuration (versus *L3-R3-R3-L3* or *R3-L3-L3-R3*) in

double *L3* and *R3* focus cells. All data from 3 repeats.

Fig. S4. (**A**) Cell growth following a 15 min EdU pulse compared to no EdU. Cell length at
different time intervals (**B**) without or (**C**) with EdU pulse. n.s. indicates two-sample t-test of
mean cell length compared to control p-value > 0.01. Data from 3 repeats. (**D**) Image
analysis of EdU experiment. A representative cell after EdU protocol showing Tsr[TMR],
EdU[Alexa488] and DAPI labelling. Red line is the cell border and dashed line shows the center
line of the cell. Black line in Tsr[TMR] channel shows the pole areas from where the Tsr
intensity is calculated. Black cross in EdU[Alexa488] channel indicates a detected EdU focus.
Black lines in DAPI channel indicate segmented nucleoid areas. For more information see
Methods. Scale bars: 1 µm (**E**) Accuracy of the retention measurement as function of sample
size. Different sample sizes were drawn from a binomial distribution with 50% (dashed line)
or 70% (solid line) success rate and SD was calculated between them (10^5 repeats for each
value). The data from Fig. 5 are shown with dots.

 Table S1. Strain list.

Strain	Relevant genotype	Source or reference
AB1157	F⁻, λ⁻, rac⁻, *thi-1, hisG4, Δ(gpt-proA)62, argE3, thr-1, leuB6, kdgK51, rfbD1, araC14, lacY1, galK2, xylA5, mtl-1, tsx-33, supE44(glnV44), rpsL31*(str^R), *qsr'-0, mgl-51*	Coli Genetic Stock Center (CGSC) #1157 (Bachmann, 1996)
AU2101	AB1157, *lacO240* at *ori1* (3908) (*hyg*), *tetO240* at *ter3* (1644) (*gen*), *ΔleuB::Plac-lacI-mCherry-frt, ΔgalK::Plac-tetR-mCerulean-frt, ΔmukB::kan*	(Nolivos et al., 2016)
JM09	AB1157, *lacO240* at *ori1* (3908) (*hyg*), *tetO240* at *ter3* (1644) (*gen*), *ΔleuB::Plac-lacI-mCherry-frt, ΔgalK::Plac-tetR-mCerulean-frt, attTn7-GFPmut2, Δflhd-frt, ΔmukB::kan*	This study
JM122	AB1157, *tsr-HaloTag-kan*	This study
JM127	AB1157, *tsr-HaloTag-frt, ΔmatP::kan*	This study
JM128	AB1157, *tsr-HaloTag-frt, ΔseqA::kan*	This study
JM130	AB1157, *tsr-HaloTag-frt, Δdam::kan*	This study
JM131	AB1157, *tsr-HaloTag-frt, ΔmukB::kan*	This study
JM133	AB1157, *tsr-mYpet-kan*	This study
JM135	AB1157, *lacO240* at *L3* (2268) (*hyg*), *tetO240* at *R3* (852) (*gen*), *ΔleuB::Plac-lacI-mCherry-frt, ΔgalK::Plac-tetR-mCerulean-frt, ΔmatP::kan*	This study
JM136	AB1157, *tsr-HaloTag-frt, matPΔC20-kan*	This study
JM137	AB1157, *tsr-HaloTag-frt, ΔzapB::kan*	This study
JM140	AB1157, *lacO240* at *L3* (2268) (*hyg*), *tetO240* at *R3* (852) (*gen*), *ΔleuB::Plac-lacI-mCherry-frt, ΔgalK::Plac-tetR-mCerulean-frt, ΔmukB::kan*	This study
JM141	AB1157, *frt-mCherry-dnaN, dnaQ-Ypet-kan*	This study
JM142	AB1157, *frt-mcherry-dnaN, ΔyjdA::kan*	This study
JM143	AB1157, *lacO240* at *L3* (2268) (*hyg*), *tetO240* at *R3* (852) (*gen*), *ΔleuB::Plac-lacI-mCherry-frt, ΔgalK::Plac-tetR-mCerulean-frt, ΔyjdA::kan*	This study
JM150	AB1157, *frt-mCherry-dnaN, dnaQ-Ypet-frt, ΔmatP::kan*	This study
JM152	AB1157, *frt-mCherry-dnaN, dnaQ-Ypet-frt, ΔmukB::kan*	This study
JW4070	*ΔyjdA::kan*	Coli Genetic Stock Center (CGSC) #10929
RRL36	AB1157, *dnaQ-Ypet-kan*	(Reyes-Lamothe et al., 2008)
RRL66	AB1157, *lacO240* at *L3* (2268) (*hyg*), *tetO240* at *R3* (852) (*gen*), *ΔleuB::Plac-lacI-mCherry-frt, ΔgalK::Plac-tetR-mCerulean-frt*	(Reyes-Lamothe et al., 2008)

RRL189	AB1157, *lacO*240 at *ori1* (3908) (*hyg*), *tetO*240 at *ter3* (1644) (*gen*), Δ*leuB::Plac-lacI-mCherry-frt*, Δ*galK::Plac-tetR-mCerulean-frt*	(Reyes-Lamothe et al., 2008)
RRL388	AB1157, *frt-mCherry-dnaN*	(Soubry et al., 2019)
RRL396	AB1157, *frt-Ypet-dnaB, kan-mCherry-dnaN*	(Soubry et al., 2019)
SN192	AB1157, *lacO*240 at *ori1* (3908) (*hyg*), *tetO*240 at *ter3* (1644) (*gen*), Δ*leuB::Plac-lacI-mCherry-frt*, Δ*galK::Plac-tetR-mCerulean-frt, mukB-mYpet-frt*	(Nolivos et al., 2016)
SN302	AB1157, *lacO*240 at *ori1* (3908) (*hyg*), *tetO*240 at *ter3* (1644) (*gen*), Δ*leuB::Plac-lacI-mCherry-frt*, Δ*galK::Plac-tetR-mCerulean-frt, mukB-mYpet-frt, ΔmatP::cat*	(Nolivos et al., 2016)

967 **Table S2.** Primer list

Name	Sequence	Construct
JMP48_Fw	CGCCGCGTAAAATGGCCGTGGCAGATAGCGAGGA GAACTGGGAAACATTTTCGGCTGGCTCCGCTGC	λ-red attachment of *HaloTag-kan* or *mYpet-kan* to *tsr* at the endogenous locus.
JMP49_Rv	AATCTCCTTATGCCCGATAACATTTTGCTTATCGGGCA TTTTCATGGCGATATGAATATCCTCCTTAGTTCCTAT	