Manuscript title

Identification and characterisation of the gene encoding an amastine-like surface protein (ALSP) in the *Leishmania donovani* genome: a putative anti-leishmanial drug target

Bapi Biswas[1], Bhakti Laha[1], Monidipa Ghosh[1]*

[1]Department of Biotechnology, National Institute of Technology Durgapur, Durgapur-713209, West Bengal, India.

* To whom correspondence should be addressed. Tel: +91-9434789001; Fax: +91-343-2754027; Email: monidipa.ghosh@bt.nitdgp.ac.in

ABSTRACT

The host-parasite interaction is a complex molecular cross-talk, where the parasite employs a repertoire of surface and secretory effector molecules for host cell manipulation and successful parasitism. The current study identifies an Amastine like Surface Protein (ALSP), expressed in abundant amount in the intracellular amastigotes form of *Leishmania donovani*, the causative agent of visceral leishmaniasis. Characterization and subcellular localization of the protein indicates its stage specific presence in the cytosol of the amastigotes. MALDI-TOF MS analysis reveals its molecular mass to be 10.147 kDa. The trypsin digested purified native protein shows 100% identity with the leishmanial ALSP through MALDI-TOF-MS-MS peptide analysis. Prediction of its presence and expression was done by immunolocalization and transcript level observations. Based on *in-silico* study, ALSP has been hypothesized to serve as a source of energy for the intracellular parasite during parasitism by conversion of triglycerides into glycerol and fatty acid, assigning a role in virulence through triglyceride lipase activity. The intracellular survival of the parasite with the intervention of the target protein may help in harnessing the candidate molecule, as an appealing target for design of novel chemotherapeutics against visceral leishmaniasis in future.

Keywords: Amastine like Surface Protein, host-parasite interaction, *Leishmania donovani*, parasitism.

INTRODUCTION

Leishmaniasis is a vector-borne disease, which is spread by protozoan parasites of the genus *Leishmania*. The causative agent *Leishmania* exists in 20 species all over the globe, which are transmitted to the human, by the bites of ~30 different species of infected female phlebotomine sandflies and of *Lutzomyia* in the Old World and in the New World respectively (1). Clinical manifestation of *Leishmania* typically has four forms: cutaneous leishmaniasis (CL), mucocutaneous leishmaniasis, visceral leishmaniasis, or kala-azar (VL) and post-kala-azar dermal leishmaniasis (PKDL) (2,3). The most fatal form of the disease, if left untreated, is visceral leishmaniasis, which is caused by *Leishmania donovani* in East Africa and the Indian subcontinent and *Leishmania infantum* in North Africa, Latin America and Europe (4,5). The *Leishmania*-affected cells' morphology consists of two distinct forms: an extracellular flagellated promastigotes form found in the midgut of female phlebotomine sandflies and an obligate intracellular non-flagellated amastigotes form in the mammalian host (6,7). They comprise of the ninth-most inherited diseases among the 13 parasitic and bacterial neglected tropical diseases worldwide (8,9). 98 countries and three territories across five continents are affected by these diseases (10). Based on the World Health Organization (WHO) reports, 12 million people suffer from leishmaniasis worldwide, among which 20,000–40,000 people succumb to death every year (10,11). Few drugs such as antimonials, amphotericin B, and miletefosine, are used to treat VL but they are not much effective due to their lack of efficacy, adverse effects, increasing drug resistance, and high cost.

Moreover, there are no available vaccines against Leishmaniasis and chemotherapeutic treatment of infected people is one of the main strategies to control the disease. The development of a novel chemotherapeutic to control Leishmaniasis therefore of utmost importance. However, lack of well-validated molecular targets in *L. donovani* has hindered the development of future chemotherapeutics. Hence identification and functional validation of novel drug targets in the trypanosomatid parasite is urgently required and to this end, a much better understanding of parasite biology is needed.

In our previous comparative genomics analysis between *L. donovani* and other *Leishmania species*, a repertoire of fifty seven species-specific genes in *L. donovani* was identified, which

indicates their essentiality in virulence and/or pathogenesis in VL leishmaniasis (12). Among them, the putative functions of thirty eight genes have been assigned through *in-silico* analysis (12). In the present study, we have studied the expression and functional implication of amastin like surface protein (ALSP) (Gene ID: LdBPK_301490.1) in *L. donovani* parasitism through cell biology, proteomics and bioinformatics approaches. Amastin is a transmembrane glycoprotein found on cell surfaces of leishmanial species (12,13). Our previous studies have documented the presence of the amastin-encoding genes at multiple chromosomal locations in leishmania-causing parasites, of which one amastin gene copy on chromosome 34, specific for *L. donovani*, was considered responsible for visceralization of the pathogen (12). Rochette et al suggest that amastin genes are largely expressed in *L. donovani* amastigotes from patients of visceral leishmaniasis (14). We observed that ALSP was expressed only in amastigotes but absent in promastigotes form of the parasite. ALSP was present in the cytoplasm of host cells and is involved in the parasitism through triglyceride lipase activity. The study supports that *Leishmania* needs energy to stay alive in the host cells during parasitism and ALSP provides the necessary energy to the amastigotes for their intracellular survival through the production of glycerol from triglycerides.

MATERIAL AND METHODS

Cell and parasite culture

L. donovani (AG83) parasites were maintained as promastigotes at 23°C in M199 media supplemented with 10% FBS, penicillin (50 U/mL) and streptomycin (50 μg/mL) (Gibco, US). Promastigotes in their late log phase (10 million promastigotes/mL) were harvested by centrifugation at 1000*g* for 10 min and washed with phosphate buffer saline (PBS) pH 7 (15). Human monocytic cell line THP1 was maintained in RPMI (Gibco, US) media similarly supplemented with FBS, penicillin and streptomycin (16). The promastigotes were transformed into amastigotes by infecting THP1 cells with *L. donovani* promastigotes at a ratio of 1:10 (cell: parasite). The amastigotes were isolated following the method described with some modifications (17).

Accession numbers

The IDs and annotations of the protein sequence are as follows:

(CBZ37742.1 LDBPK_342650), GeneDB[i] (LdBPK_301490.1.1: pep), Gene ID: 13392833.

Production of customized antibody and ELISA

The production of customized antisera against the ALSP (Amastine like surface protein) was performed by using the best probable peptide epitopic domain from that protein (SSPFSSTRSSSSSRS –C, the addition of a Cystine residue at C terminal end required for Keyhole limpet hemocyanin (KLH) conjugation) by employing bioinformatics tool outsourced from BioBharati LifeScience Pvt. Ltd. (Kolkata, India). A New Zealand (NZ) White Rabbit model was immunized with the help of a KLH conjugation solution containing KLH used for coupling with the peptide epitopic domain. The serum was collected from both batches of the NZ White Rabbit model before (pre bleed) and after giving the 7th immunogenic injections (6th Booster dosage). It took 2 months 15 days to collect the final antisera.

Determinations of antibody titer and affinity purification of antisera were done by Indirect ELISA. In both cases, 500 pg of purified ALSP per well have been used to check their titer of various dilutions of antisera containing the specific antibody of both the batches (1:500, 1:1000, 1:2000, 1:5000, 1:10000, 1:20000, 1:400000 dilutions). The equal dilution of pre-immune sera taken from the same rabbit was also used to check the specificity of the antibody.

Affinity-based immune-precipitation assay for the native protein purification

To detect the molecular weight of the native ALSP, 10 million promastigotes and amastigotes of *L.donovani* were resuspended in 1x PBS (pH 7.4) buffer and incubated at 4^0c for 10 min. Rapid freeze-thawing was done in the presence of liquid nitrogen and the insoluble part removed by the centrifugation at 200g for promastigotes and 100g for amastigotes forms. Finally supernatants were taken for immunoprecipitation along with the pre-immune sera and antisera of the NZ white rabbit. The immunoprecipitation was carried out with Protein A- Sepharose bead which were added to the pre-immune sera and antisera in a 1:2 volume ratio and incubated at 4^0c overnight with gentle inversion rotation. The unbound preimmune sera and antisera were washed with PBS (pH 7.4) at room temperature thrice by centrifugation at 200g for promastigotes and 100g for amastigotes forms. The soluble supernatants of both forms were added to the specific antibody loaded Protein A- sepharose column and incubated overnight at 4^0 c with gentle inversion rotation. The unbound protein was removed similarly

like before. Hence the bound protein with antisera and pre-immune sera through Protein A-sepharose bead in both forms was eluted by boiling with SDS-PAGE loading buffer for 15 min at 100^0 c and collected by centrifugation (same parameters). The elutes were analyzed through immunoblotting after probing with Anti-ALSP polyclonal antibody (18).

MALDI-TOF

The dialysate of the native protein was observed in MALDI-TOF to detect its purity and molecular weight. The dissolved protein was spotted on a target MALDI plate using α-cyano-hydroxycinnamic acid (CHCA) as a matrix and analyzed using a MALDI-TOF mass spectrometer (Applied Biosystems, USA). Spectra were calibrated using the matrix. This study was done at CSIR-IICB Kolkata using the central instrumentation facility for mass spectrometry (MALDI) (19).

Two-step reverse transcription PCR

The presence of ALSP transcripts in promastigotes and amastigotes forms of *L. donovani* was checked using isolated whole-cell RNA from both structural forms of the parasite with TRI Reagent (Sigma T9424). Then, the whole-cell RNA was used to synthesize cDNA using reverse transcriptase enzyme at 42 °C for 60 min with the gene-specific ALSP-F and ALSP-R primers (Thermo Scientific RevertAid First Strand cDNA Synthesis Kit). Subsequently, PCR was employed to check the expression level of the native ALSP, maintaining a Tm value of 55°C and by adding 5 µl of cDNA (20).

Fluorescence microscopy

L.donovani promastigotes and amastigotes were harvested by centrifugation (4000g for promastigotes and 100g for amastigotes) and the pellets were washed with PBS two times by centrifugation. Cells were fixed in chilled methanol for 2 min at 4^0c and washed with PBS. Following permeabilisation with permeabilisation buffer for 30 sec-1min at 4^0c, cell pellets were washed with PBS and incubated with preimmune sera and immune sera against the native ALSP protein (1:25 dilution) in the presence of 3% bovine serum albumin for 45 min. Unbound proteins were removed by washing three times in PBS and probed with fluorescein isothiocyanate (FITC)-conjugated goat-derived IgG (Thermo Fisher SCINETIFIC, Waltham)

(1:500 dilution) for 30 min. Further cells were mounted with mounting media along with DAPI (16) and its presence observed under a fluorescence microscope (Zeiss, UK) at 10× magnification and oil immersion. The parasites and infected cells were stained with the pre-immune sera as negative controls and similar optical filters were used for DAPI ($\lambda = 461$ nm) and FITC ($\lambda = 591$ nm).

Determination of native protein conformation through sequencing

The MALDI-TOF-MS/MS analysis was done using trypsin digested native ALSP sample to check its sequence identity based on MS/MS Ions Search on the MASCOT database server (http://www.matrixscience.com/search_form_select.html). Parameters considered were: carbamidomethyl (C) for fixed modification, oxidation (M) for variable modification, trypsin as enzyme, peptide mass tolerance (100 ppm), and fragment mass tolerance (+0.2).

This study was carried out at CSIR-IICB Kolkata using the central instrumentation facility for mass spectrometry (MALDI-TOF-MS/MS).

Protein 3D structure and function prediction through I-TASSER server

The amino acid sequence of native ALSP was uploaded to the I-TASSER server (https://zhanglab.ccmb.med.umich.edu/I-TASSER/). For each target, I-TASSER simulations generate a large ensemble of structural conformations, called decoys. To select the final models, I-TASSER uses the SPICKER program to cluster all the decoys based on pair-wise structure similarity and reports up to five models which correspond to the five largest structure clusters. The confidence of each model is quantitatively measured by the C-score that is calculated based on the significance of threading template alignments and the convergence parameters of the structure assembly simulations. C-score is typically in the range of [-5, 2], where a C-score of a higher value signifies a model with a higher confidence and vice-versa. TM-score and RMSD are estimated based on C-score and protein length following the correlation observed between these qualities. Since the top 5 models are ranked by their cluster sizes, it is possible that the lower-rank models have a higher C-score in rare cases. Although the first model has a better quality in most cases, it is also possible that the lower-rank models have a better quality than the higher-rank models as seen in our benchmark tests.

The function of the ALSP has been annotated using COFACTOR and COACH based on the I-TASSER structure prediction. While COFACTOR deduces protein functions (ligand-binding sites, EC and GO) using structure comparison and protein-protein networks, COACH is a meta-server approach that combines multiple function annotation results (on ligand-binding sites) from the COFACTOR, TM-SITE and S-SITE programs.

RESULTS

Production of antiserum against the best probable epitopic ALSP domain

The best probable antigenic domain from the ALSP with KLH conjugation was used for immunisation of NZ White Rabbit model. Antibody titers was measured and affinity purification of antisera was performed through indirect ELISA. 500 pg of purified ALSP per well was taken to check the titers of different antisera dilutions while the same dilutions of pre-immune sera were taken from the same rabbit to check the specificity of the antibody. Both the batches showed a decrease of titer value along with the increase of dilution, though the titer of pre-immune sera remains very less in comparison with the immunized batches which implies that the raised antisera is specific against the antigen. In comparison with two batches, the Batch 1 showed more titer than Batch 2 (Fig. 1a). Therefore Batch 1 has been selected to purify ALSP-antibody through true affinity purification (Fig. 1b) and used for further analysis.

Specificity check of immunized antibody

The specificity was checked of raised antibody against the most probable peptide epitopic domain from ALSP through Indirect ELISA (Fig.1a and 1b) and Immunoprecipitation (IP) (Supplementary Figure S1).

Immunolocalization of the native protein in *Leishmania donovani*

Stage-specific localization of the native protein in both the forms of *L. donovani* was evaluated by fluorescence microscopic analysis. The native protein was visualized in the amastigotes (Fig. 2. ii), after probing with antisera, while it was absent in promastigotes (Fig. 2.i). FITC-tagged IgG was used as secondary antibody for the semi-confocal microscopic localization of the ALSP protein. The protein stained with pre immune sera in promastigote

(Fig. 2. iii) and amastigote (Fig. 2. iv), was taken into account as negative control. Through in-silico analysis the exact location of the protein was identified to be in the cytosol.

Stage-specific expression and characterization of the native protein in *L. donovani*

The stage-specific expressions of ALSP gene in the promastigotes and amastigotes forms of *L.donovani* were analysed by reverse transcription PCR, using cDNA obtained from the whole- cell RNA from both its structural forms. PCR with the specific primers for ALSP gene showed a clear band near 255 bp (Fig. 3.A, L3) for the amastigotes whereas no bands were found for the promastigotes (Fig. 3. A, L2). This clearly showed that the expression of ALSP mRNA was absent in the promastigotes while amastigotes have abundant amount of expressed mRNA for the native protein.

The presence of the native protein in both the stages of *L.donovani* was also checked through fluorescence microscopy. The protein was found only in the amastigote stage (Fig. 3. B. b) while in promastigotes it was not deteted (Fig. 3. B. a). The analysis of the stage speific localization of the native ALSP by immunofluroscence, coupled with gene expression validation by RT PCR, revealed a selective expression of ALSP only in intracellular amastigote stage (absent in extracellular promastigotes), indicating its potential role in infection and parasitism.

Purification and sequence analysis of the native protein from *L. donovani* amastigotes

Physical characterization of the native protein from the amastigote stage was done through its purification from the intracellular extract while the crude extract from promastigotes was used as negative control (since expression of ALSP was not observed in promastigotes) . The antisera with sufficient titre of antibody against ALSP was used to purify the native protein from the amastigote lysate. The antisera and preimmune sera were cross- linked with protein A-Sepharose bead (Invitrogen-101041) to avoid the contamination of heavy and light chains of the antibody with the bound antigen, viz our target native protein, while being eluted. The purification was done in the presence of another negative control: albumin like protein in amastigote form of *L.donovani* (21). Both the negative controls exhibited an

absence of the native protein, whereas the lysate of amastigotes exhibited a sharp band near 10 kDa (Fig. 4. A) after incubation with protein-A-sepharose bead cross-linked with the antisera. The precised molecular weight of the native protein obtained through MALDI-TOF MS analysis was 10.147 kDA (Fig. 4. B). Few extra peaks appeared in the MALDI-TOF MS analysis data, due to the impurity of the sample. Further MALDI-TOF-MS-MS analysis supported its presence through sequence study, employing the peptide fragments of the tripsin digested native protein (from amastigotes)(Fig. 4. C).

3D structure and function prediction from the sequence of the native protein

The 3D structure of native ALSP was predicted via the I-TASSER online tool through homology modeling, by utilizing its amino acid sequence. The top five models were predicted by I-TASSER using the SPICKER program based on the structural conformations. The confidence of each model was measured by the C-score, calculated based on the significance of the threading template alignments and the convergence parameters of the structure assembly simulations. The range of the C-score is [-5, 2], wherein a C-score of a higher value signifies a model with greater confidence and vice-versa. C-scores of top 5 models were estimated as: -3.58 for model 1, -4.36 for model 2, -4.77 for model 3, -4.56 for model 4 and -4.95 for model 5. Owing to its higher C-score, the structure of Model 1 is depicted here (Fig. 5).

The function and cellular component of the native ALSP was also predicted by COFACTOR and COACH based on the I-TASSER structure prediction. The predictable function of the protein is triglyceride lipase activity (GO: 0004806) wherein the biological process involves lipid catabolism (GO: 0016042) in the cellular compartment of cytosol (GO: 0005829).

DISCUSSION

The present study predicts for the presence of a hypothetical species-specifically expressed gene, encoding an amastin like surface protein in *L.donovani,* which is potentially conferring the pathogenesis of visceral leishmaniasis. To characterize the gene, a polyclonal antibody was raised against the most antigenic site of the leishmanial native protein, which is present only in the amastigote stage of *L.donovani*. Immuno-fluroscence analysis confirmed the presence of amastin like surface protein in the cytosolic compartment of amastigotes but not in the promastigotes, which was supported by the transcript level studies. This indicates its selective stage specific expression in the intracellular amastigote form of *L.donovani*.

Further characterization of the native protein was carried out by purifying the whole cell lysate of amastigotes through immune-precipitation. The antisera, which was raised against the best possible epitopic site of the native leishmanial protein was cross-linked with protein A-Sepharose bead. Loading of the amastigote lysate in the protein-A-antibody column prevented antibody co-elution and led to localization of a protein band with molecular weight of 10.147 kDa after MALDI-TOF MS analysis. Further MALDI-TOF-MS-MS analysis was carried out with the peptide fragments of that protein band after trypsin digestion, following which an amino acid sequence was detected which exhibited 100% identity with the leishmanial ALSP (CBZ37742.1).

Subsequently the 3D structure, sub-cellular localization and function of the native ALSP was predicted by homology modelling using its amino acid sequence via the I-TASSER online tool. The predicted function of the protein is triglyceride lipase activity (GO: 0004806) and it is localised in the cytosol (GO: 0005829). Based on *in-silico* study, ALSP is predicted to serve as a source of energy for the intracellular parasite during parasitism by conversion of triglycerides into glycerol and fatty acid. Phospholipids and sphingolipids form important constituents of leishmanial cell membranes which are essential for virulence and viability of the pathogen (22,23). A plausible explanation of the role of ALSP might be that the metabolism of lipid moieties leads to membrane remodelling in *L. donovani*, which in turn plays a role in the entry of the extracellular promastigote form of the pathogen into the host cell and helps in its subsequent transformation into the intracellular non-flagellated amastigote form. Previous investigations suggest that amastins help in survival of the *L. donovani* parasites during their differentiation, from promastigote to amastigote stage transformation, inside the acidic environment of the phagolysosome. This is reportedly achieved by alteration of pH via transport of water- soluble protons and ions across the cell membrane (14, 24).

The complex molecular processes underlying the interaction between *L. donovani* and the host cell, followed by endocytosis of the pathogen, deploy a range of receptors and surface proteins which help to trigger the signalling pathways, required for prolonged survival and host-to-host transmission of the pathogen. This host-parasite dialogue can thus potentially affect the outcome of parasitic infection. Hence such parasite specific molecular target deployed by the parasite to manipulate the host behaviour may serve as an attractive future drug target against parasitic diseases. To this end, investigation of the precise molecular conformation in which the amastin protein exists, its functional interactions with other proteins in the parasite genome and its binding dynamics with potent drug molecules might help

researchers to identify new drug candidates. Understanding the mechanistic basis of stage specific expression of ALSP in obligatory intracellular amastigote form of *L. donovani*, together with its putative role as triglyceride lipase to generate glycerol from triglycerides and necessary energy for intracellular survival, may help in harnessing this protein as an appealing target for design of novel chemotherapeutics against visceral leishmaniasis in future.

AVAILABILITY

I-TASSER (Iterative Threading ASSEmbly Refinement) is a hierarchical approach to protein structure and function prediction (https://zhanglab.ccmb.med.umich.edu/I-TASSER/)

COFACTOR is a structure, sequence, and protein-protein interaction (PPI) based method for biological function annotation of protein molecules (https://zhanglab.ccmb.med.umich.edu/COFACTOR/)

COACH is a meta-server approach to protein-ligand binding site prediction (https://zhanglab.ccmb.med.umich.edu/COACH/)

SPICKER is a clustering algorithm to identify the near-native models from a pool of protein structure decoys (https://zhanglab.ccmb.med.umich.edu/SPICKER/)

SUPPLEMENTARY DATA

Supplementary Figure S1. Illustration of ALSP-GST fusion protein purification, Immunoblotting, and Immunoprecipitation. I. ALSP-GST protein purification on 15% SDS-PAGE in respect of 1kb Pre-stained Protein Marker (Lane 1). Lane 2: Purified BL21 cells with empty vector, Lane 3: Purified BL2 1 cells without Construct; Lane 4, 5 and 6: 1st, 2nd, and 3rd elution of Purified ALSP-GST fusion protein. II. Illustrate the conformation of the purified ALSP-GST fusion protein (about 35kDa) in induced (Lane 2) and un-induced form (Lane 3) by Western Blotting with Anti-GST-tagged antibody in respect of 1kb Pre-stained Protein Ladder (Lane 1). III. Immuno-precipitation (in Classical method) of the ALSP-GST fusion protein. The purified induced and un-induced ALSP-GST fusion protein pulled down with Antisera and Preimmune sera of an NZ White Rabbit by probing customized anti -ALSP antibody and anti-Rabbit-HRP conjugate Secondary antibody. Lane 2 and 3: Preimmune Sera and antisera with the purified fusion protein; Lane 4 and 5: Preimmune Sera and antisera with the uninduced fusion protein; and Lane 1: 1kb Pre-stained Protein Marker.

ACKNOWLEDGEMENT

The authors are thankful to Dr. Nahid Ali of Indian Institute of Chemical Biology (IICB) for providing pure *L. donovani (*AG83 strain) parasite culture for this project. The authors would also like to acknowledge Ms. Anandi Batabyal and Mr. Pratyay Sengupta for their sincere help to prepare the manuscript.

FUNDING

This work was funded by the Ministry of Human Resource Development, Government of India.

CONFLICT OF INTEREST

The authors do not declare any conflict of interest.

REFERENCES

1. Pearson, R.D. and de Queiroz Sousa, A. (1996) Clinical spectrum of leishmaniasis. *Clinical infectious diseases*, pp.1-11.

2. Dedet, J.P., Pratlong, F. (2009) Leishmaniasis. In: Cook, G.C., Zumla, A. (Eds.), Manson'sTropical Diseases. Saunders, Edinburgh, pp. 1341–1365.

3. WHO Expert Committee on the Control of the Leishmaniases. Meeting and World Health Organization. (2010) Control of the Leishmaniases: Report of a Meeting of the WHO Expert Committee on the Control of Leishmaniases, Geneva, 22-26 March 2010 (Vol. 949). World Health Organization.

4. McCall, L.I., Zhang, W.W. and Matlashewski, G. (2013) Determinants for the development of visceral leishmaniasis disease. *PLoS Pathog*, *9*(1), p.e1003053.

5. Lukeš, J., Mauricio, I.L., Schönian, G., Dujardin, J.C., Soteriadou, K., Dedet, J.P., Kuhls, K., Tintaya, K.W.Q., Jirků, M., Chocholová, E. and Haralambous, C. (2007)Evolutionary and geographical history of the Leishmania donovani complex with a revision of current taxonomy. *Proceedings of the National Academy of Sciences*, *104*(22), pp.9375-9380.

6. Sunter, J. and Gull, K. (2017) Shape, form, function and Leishmania pathogenicity: from textbook descriptions to biological understanding. *Open biology*, *7*(9), p.170165.

7. Rittig, M.G. and Bogdan, C. (2000) Leishmania–host-cell interaction: complexities and alternative views. *Parasitology Today*, *16*(7), pp.292-297.

8. Molyneux, D.H., Hotez, P.J. and Fenwick, A. (2005) "Rapid-impact interventions": how a policy of integrated control for Africa's neglected tropical diseases could benefit the poor. *PLoS Med*, *2*(11), p.e336.

9. Hotez, P.J., Molyneux, D.H., Fenwick, A., Ottesen, E., Sachs, S.E. and Sachs, J.D. (2006) Incorporating a rapid-impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis, and malaria. *PLoS Med*, *3*(5), p.e102.

10. Alvar, J., Vélez, I.D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jannin, J., den Boer, M. and WHO Leishmaniasis Control Team. (2012) Leishmaniasis worldwide and global estimates of its incidence. *PloS one*, *7*(5), p.e35671.

11. Alvar, J., Yactayo, S. and Bern, C. (2006) Leishmaniasis and poverty. *Trends in parasitology*, *22*(12), pp.552-557.

12. Sengodan SK, Gokulasuriyan RK, Ghosh M (2014) Comparative in- silico genome analysis of Leishmania (Leishmania) donovani: a step towards its species specificity. MGENE 2:782–798.

13. Jackson, A.P. (2010) The evolution of amastin surface glycoproteins in trypanosomatid parasites. *Molecular biology and evolution*, *27*(1), pp.33-45.

14. Rochette, A., McNicoll, F., Girard, J., Breton, M., Leblanc, É., Bergeron, M.G. and Papadopoulou, B. (2005) Characterization and developmental gene regulation of a large gene family encoding amastin surface proteins in Leishmania spp. *Molecular and biochemical parasitology*, *140*(2), pp.205-220.

15. Roychoudhury, J. and Ali, N. (2008) Sodium stibogluconate: Therapeutic use in the management of leishmaniasis.

16. Das, A., Dasgupta, A., Sharma, S., Ghosh, M., Sengupta, T., Bandopadhyay, S. and Majumder, H.K. (2001) Characterisation of the gene encoding type II DNA topoisomerase from Leishmania donovani: a key molecular target in antileishmanial therapy. *Nucleic acids research*, *29*(9), pp.1844-1851.

17. Lúcia M, Moreno V, De M et al (1998) In vitro method for isolation of amastigote forms of Leishmania amazonensis from J774G8 macro- phage induced by temperature shifting. Mem Inst Oswaldo Cruz 93: 99–102.

18. Kowal, R. and Parsons, R.G. (1980) Stabilization of proteins immobilized on Sepharose from leakage by glutaraldehyde crosslinking. *Analytical biochemistry*, *102*(1), pp.72-76.

19. Raghwan, Chowdhury R. (2014) Host cell contact induces fur-dependent expression of virulence factors CagA and VacA in Helicobacter pylori. *Helicobacter,* 19:17–25.

20. Couvreur, B., Bollen, A., Le Ray, D. and Dujardin, J.C. (2003) Reverse transcription-polymerase chain reaction construction of plasmid-based, full-length cDNA libraries from Leishmania infantum for in vitro expression screening. *Memórias do Instituto Oswaldo Cruz*, *98*(4), pp.477-480.

21. Laha, B., Verma, A.K., Biswas, B., Sengodan, S.K., Rastogi, A., Willard, B. and Ghosh, M. (2019) Detection and characterization of an albumin-like protein in Leishmania donovani. *Parasitology research*, *118*(5), pp.1609-1623.

22. Zhang, K. and Beverley, S.M. (2010) Phospholipid and sphingolipid metabolism in Leishmania. *Molecular and biochemical parasitology*, *170*(2), pp.55-64.

23. Messaoud, H.B.B., Guichard, M., Lawton, P., Delton, I. and Azzouz-Maache, S. (2017) Changes in lipid and fatty acid composition during intramacrophagic transformation of Leishmania donovani complex promastigotes into amastigotes. *Lipids*, *52*(5), pp.433-441.

24. Cruz, M.C., Souza-Melo, N., da Silva, C.V., DaRocha, W.D., Bahia, D., Araújo, P.R., Teixeira, S.R. and Mortara, R.A. (2012) Trypanosoma cruzi: role of δ-amastin on extracellular amastigote cell invasion and differentiation. *PLoS One*, *7*(12), p.e51804.

FIGURE LEGENDS:

Fig. 1 Characterisation of polyclonal antibody against L.donovani ALSP protein. (A) Binding capacities of polyclonal antibodies raised against both Batch 1 and Batch 2 before and after immunisation of an NZ White Rabbit model, measured by indirect ELISA method. The titre of pre immune serum with different dilutions is indicated by the green and blue curves while the binding capacities of antisera in batch 1 and batch 2 are indicated by red and black curves respectively. (B) True affinity purification of ALSP-Ab from batch 1 of the NZ white rabbit model with varying dilutions.

Fig 2. Sub-cellular localisation of the native ALSP in the both structural forms of *L.donovani* through semiconfocal fluorescence microscopy using filters at 461 nm for DAPI and 591 nm for FITC. Visualization of the L.donovani promastigotes (i) and amastigotes (ii), are probed with antisera and FITC conjugated goat- derived anti-rabbit IgG. Likewise, L.donovani promastigotes (iii) and amastigotes (iv) are probed with pre immune sera as negative controls.

Fig. 3 Stage-specific expression of native ALSP in L.donovani promastigotes and amastigotes. A. Reverse transcription-based amplification of ALSP-encoding gene transcripts using whole-cell RNA obtained from promastigote and amastigote forms of L.donovani. Lane 1, 1 kb DNA marker; Lane 2 and 3, RT-PCR products of whole-cell RNA from promastigotes and amastigotes respectively. Lane 3 contains a sharp band near 255 bp; no detection of bands in Lane 2. B. Immunocytochemical localisation by probing with antisera and FITC conjugated goat-derived anti-rabbit IgG secondary antibody in promastigote (a) and amastigote (b), using fluorescence microscopy. For detection, filters were used at 461 nm for DAPI and at 591 nm for FITC.

Fig. 4 MALDI-TOF based analysis of protein expression level and determination of molecular weight of the native purified protein in L.donovani amastigotes. Confirmation by sequencing performed via MALDI-TOF-MS-MS technique. A. Purification of native ALSP from cell lysate of promastigotes and amastigotes using antisera cross-linked with protein A-sepharose bead. Immunoblotting using antiserum and anti-rabbit HRP conjugated secondary antibody. Lane 1, protein marker (10-250 kDa); Lane 2, purified native protein expressed in promastigotes, taken as negative control; Lane 3, native ALSP band detected near 10 kDa; Lane 4, purified albumin-like protein expressed in amastigotes, taken as negative control. B. Determination of molecular weight of purified native ALSP. Exact molecular weight determined after being singly (M) charged with the lesser. The m peak at 10.147 kDa represents the accurate molecular mass of the purified native protein. C. Sequencing of the native protein by MALDI-TOF-MS-MS analysis after digestion with trypsin. Peptide of the protein exhibited 100% identity with Amastine like surface protein (ALSP).

Fig. 5 Prediction of three-dimensional structure of the native ALSP by homology modelling based on amino acid sequence. C-score=-3.58; estimated TM-score = 0.32±0.11 and estimated RMSD =11.6±4.5Ã.

A

B

DAPI	FITC	DAPI/ FITC

a

b

B

C