The essential role of *Dnmt1* in gametogenesis in the large milkweed bug

Oncopeltus fasciatus

Joshua T. Washington[1], Katelyn R. Cavender[1], Ashley U. Amukamara[1], Elizabeth C. McKinney[1], Robert J. Schmitz[2], Patricia J. Moore[1, *]

[1] Department of Entomology, University of Georgia

[2] Department of Genetics, University of Georgia

*Patricia J. Moore, 413 Biological Sciences Building, Department of Entomology, University of Georgia, Athens, GA, 30602; (706) 542-0169

Email: pjmoore@uga.edu

Patricia J. Moore: https://orcid.org/0000-0001-9802-7217

Robert J. Schmitz: https://orcid.org/0000-0001-7538-6663

Keywords

DNA methylation, epigenetics, *Dnmt1*, spermatogenesis, germ cells, *Oncopeltus fasciatu*

Abstract

Given the importance of DNA methylation in protection of the genome against transposable

elements and transcriptional regulation in other taxonomic groups, the diversity in both levels

and patterns of DNA methylation in the insects raises questions about its function and evolution.

We show that the maintenance DNA methyltransferase, DNMT1, affects meiosis and is essential

to fertility in milkweed bugs, *Oncopeltus fasciatus*, while DNA methylation is not required in

somatic cells. Our results support the hypothesis that *Dnmt1* is required for the transition of germ

cells to gametes in *O. fasciatus* and that this function is conserved in male and female

gametogenesis. They further suggest that DNMT1 has a function independent of DNA

methylation in germ cells. Our results raise the question of how a gene so critical in fitness

across multiple insect species can have diverged widely across the insect tree of life.

Introduction

Despite the apparent ubiquity of DNA methylation across the eukaryotic tree of life (Schmitz et al. 2019, Lewis et al. 2020), in the insects there is considerable variation both in the presence and extent of DNA methylation and even the presence and number of the DNA methyltransferases (Bewick et al. 2016, Lyko 2018, Glastad et al. 2019). Thus, the functional role of DNA methylation and its associated DNA methyltransferases in the insects is unclear. While some studies have associated levels of methylation with gene expression, most studies in the insects find no functional association (Bewick et al. 2016, Glastad et al. 2019).

One role emerging from knockdown and silencing studies across several insect species is that *Dnmt1* is required for oogenesis (Schulz et al. 2018, Bewick et al. 2019, Gegner et al. 2019, Amukamara et al. 2020). For example, *Dnmt1* is required for maintenance of DNA methylation following cell division in the milkweed bug, *Oncopeltus fasciatus* (Bewick et al. 2019, Amukamara et al. 2020), the downregulation of *Dnmt1* using RNAi results in a reduction in methylation and also the cessation of oogenesis. However, it is unclear whether the effect on oogenesis is mediated by the reduction in DNA methylation (Amukamara et al. 2020). While the expected reduction in DNA methylation is seen throughout the organism following *Dnmt1* knockdown, the only phenotypic consequence is to the germ cells. Evidence from other species supports a function independent of DNA methylation. *Dnmt1* is also essential to egg production in *T. castaneum* (Schulz et al. 2018), a beetle that has no DNA methylation at all (Zemach et al. 2010). This suggests that *Dnmt1* can have a function specific to germ cells in insects that is independent of its function in maintaining DNA methylation. This led us to hypothesize that *Dnmt1* plays a role in meiosis in insects (Amukamara et al. 2020).

In this study, we tested the hypothesis that *Dnmt1* is essential to gametogenesis in *O. fasciatus*.

Furthermore, if *Dnmt1* has a role in gametogenesis, including meiosis, it should be conserved

across the sexes. Testing the function of *Dnmt1* in males allows us to capitalize on the well-

characterized process of spermatogenesis in *O. fasciatus*, specifically (illustrated in Figure 1;

Economopoulos & Gordon 1971), and the conserved features of insect spermatogenesis,

generally (Dallai 2014). In *O. fasciatus,* as in many insects, there are two points in development

where meiosis can occur. The first stage where meiosis will occur is during the larval stages.

Testis development and spermatogenesis is initiated during larval development in *O. fasciatus*

(Economopoulos & Gordon 1971). During the first three instars the testes consist of seven

globular follicles that will develop into the testis tubules (Schmidt et al. 2002). Meiosis is

initiated in the 4^{th} instar, and by the end of the 4^{th} instar, cysts containing spermatids are present.

Differentiation of the spermatids commences in the 5^{th} instar, and males emerge with up to

250,000 spermatids that continue to differentiate during sexual maturation. Thus, we can target

the developmental time point at which meiosis occurs. In addition, *O. fasciatus* males can

produce gametes throughout their adult lives and therefore meiosis can occur in the adult testis.

If *Dnmt1* is required for meiosis and gametogenesis we predicted that it will be required both

during larval development and as adult males replenish sperm stores following mating. We

therefore compared the testis phenotypes of adults developing from nymphs in which *Dnmt1*

expression was downregulated either before or after the major wave of meiosis that occurs in

testis development (Economopoulos & Gordon 1971, Schmidt et al. 2002, Ewen-Campen et al

2013). We also treated sexually mature adults, and examined fertility in males following sperm

depletion, testing for the ability of knockdown males to replenish sperm stores following

multiple matings. Our results demonstrated that expression of *Dnmt1* is required for the

development of sperm both during larval and adult spermatogenesis and that the impact of

Dnmt1 knockdown was greatest if it occurred prior to the onset of meiotic divisions in the

developing testes. These results establish that *Dnmt1* plays a critical conserved function across

the sexes during gametogenesis in *O. fasciatus* and that it is required for germ cell development.

Results

The pattern of *Dnmt1* expression during testis development and across tissues suggests a

role in spermatogenesis.

We measured *Dnmt1* expression during larval development and sexual maturation. Our

prediction was that if *Dnmt1* is required during gametogenesis, particularly meiosis, then its

expression should mirror those of two genes known to be involved in germ cell development,

Boule and *Vasa* (Shah et al. 2010, Yajima & Wessel 2011). Thus, we predicted that *Dnmt1*

expression would be highest during the stages when gametogenesis is occurring. As predicted,

testis-specific expression of *Dnmt1* is highest during the 4^{th} and 5^{th} instars of development

(Figure 2A; ANOVA, F = 27.426, d.f. = 3, 38, p < 0.001), stages when meiosis and

spermatogenesis is most active (Economopoulos & Gordon 1971). The expression is lowest in

virgin adult males. This expression pattern mirrored that observed for *Boule*, a gene with a well-

characterized role in meiosis (Figure 2B; ANOVA, F = 36.346, d.f. = 3, 38, p < 0.001). *Dnmt1*

expression pattern was also similar to *Vasa*, a highly conserved marker of the germline (Figure

2C; ANOVA, F = 20.444, d.f. = 3, 38, p < 0.001). While *Dnmt1* is expressed across all adult

tissues, as would be expected given its role in maintenance of DNA methylation, expression is

highest in testes compared to gut, head, and muscle tissue (Figure S2; ANOVA, F = 67.311, d.f.

= 3, 36, p < 0.001), also suggesting a special role during gametogenesis.

***Dnmt1* knockdown reduced levels of DNA methylation in the testis genome**.

To confirm the effectiveness of our RNAi treatment, we measured expression of *Dnmt1* in a

subset of ds-RED and ds-*Dnmt1* injected males (Figure 3). Quantitative PCR demonstrated

reduced levels of *Dnmt1* mRNA in ds-*Dnmt1* injected males compared to ds-RED injected males

in both 3rd instar (ANOVA; F = 83.126, d.f. = 1, 31, p < 0.001) and 5th instar (ANOVA; F =

49.793, d.f. = 1, 22, p < 0.001) treated males 6 to 10 days following injection with dsRNA.

Treatment of males with ds-*Dnmt1* had the predicted effect on *Dnmt1* expression. We performed

whole genome bisulfite sequencing to evaluate the impact of the ds-*Dnmt1* on DNA methylation

genome wide. The reduction in expression of *Dnmt1* in the RNAi individuals had the expected

phenotypic effect on DNA methylation with a reduction of genome methylation in the testes for

ds-*Dnmt1* treated males, but not control males (Supplementary materials, Figure S2). Genomic

DNA from the testes of control treated males had approximately 12.5% CpG methylation

regardless of stage they were treated. Knockdown of *Dnmt1* at the earlier stage of development

led to a greater percentage reduction of methylation. Treatment with ds-*Dnmt1* at the 5th instar

reduced the percent CpG methylation from around 12.5% to around 5%. Treatment with ds-

Dnmt1 at the 3rd instar reduced the level of methylation even further, to around 2%, as predicted

given the greater numbers of cell divisions that were expected between treatment and sampling

between these two treatments.

Dnmt1 knockdown affected testis size and structure.

Knockdown of *Dnmt1* prior to meiosis, in the 3rd instar stage, had a significant effect on testis

size in virgin males, while knockdown of *Dnmt1* during the developmental stage after which

meiosis is initiated, the 5th instar stage, had no effect on testis size (Figure 4; ANOVA, F =

20.360, d.f. = 4, 58, p < 0.001) compared to either uninjected controls or ds-RED treated males.

Knockdown of *Dnmt1* prior to meiosis affected testis tubules to a greater extent than knockdown

following the initiation of meiosis during larval testis development (Figure 5). Control males

injected in the 3^{rd} or 5^{th} instar stage of development showed the expected structure of the testis

tubule (Figure 5A and C). At the anterior end of the testis tubule, spermatogonia and primary

spermatocytes had the characteristic nuclear structure and there was clear evidence of mitotic

division within the spermatogonia (Figure 5C, arrowhead), using α-phosphohistone H3 (pHH3)

to stain for chromosome condensation in preparation for mitosis and meiosis. We also observed

α-pHH3 staining in spermatocysts at the border between primary and secondary spermatocytes

in the control testis tubules in both 3^{rd} and 5^{th} instar treated males. Following this band of

relatively synchronous meiotic activity, the posterior testis tubule showed spermatids and

developing spermatozoa as they matured. We confirmed the use of this band of α-pHH3 stained

spermatocysts as a landmark of meiosis using knockdown of Boule (Supplementary materials,

Figure S3).

The structure of the testis tubules of males treated with ds-*Dnmt1* in the 3^{rd} instar was highly

disrupted and both the anterior and posterior testis tubule was affected (Figure 5B). There were

fewer spermatocysts in both the region of spermatogonia and primary spermatocytes. While

there were occasional α-pHH3 positive nuclei, these were not well organized into spermatocysts

and were spread throughout the testis tubule rather than organized at the junction between

primary and secondary spermatocytes. There was evidence of mitotic activity in the

spermatogonia (Figure 5B arrowhead), although these were less frequent in ds-*Dnmt1* treated

males. The testis tubules of males treated with ds-*Dnmt1* in the 5^{th} instar stage

of development had a structure much more similar to control males (Figure 5D). Mitotic activity

was apparent in the spermatogonia (Figure 5D arrowhead) and most testis tubules had evidence

of mature sperm and α-pHH3 stained spermatocysts at the junction between primary and

secondary spermatocytes (Figure 5D arrow). Unorganized spermatocysts below this junction

were frequently observed, however, and ds-*Dnmt1* males treated at the 5th instar had variable

phenotypes posterior to the primary spermatocytes, presumably depending on when treatment

occurred following the wave of meiosis along the testis tubule axis.

***Dnmt1* knockdown in adult males prevented replenishment of sperm stores**

Downregulating *Dnmt1* expression in adult males resulted in a loss of fecundity over time. The

third and final females mated to ds-*Dnmt1* treated males ran out of sperm to fertilize eggs more

rapidly than those mated to control males. Clutches of eggs laid by females mated to ds-*Dnmt1*

treated males were not fertilized and failed to hatch at an earlier collection day than those laid by

females mated to control males (Figure 6; Wilcoxon $\chi^2 = 13.978$, d. f. = 1, p < 0.001).

The loss of fertility in males was associated with smaller testis size. Males treated with ds-*Dnmt1*

at seven days post-emergence and then allowed to mate for three weeks prior to dissection had

statistically significantly smaller testis area than the testes of control males after the same mating

treatment (ANOVA, F = 29.084, d.f. = 1, 51, p < 0.001; Supplementary materials, Figure S4).

The smaller testis area of ds-*Dnmt1* treated adult males was associated with a breakdown in testis

tubule structure and the loss of sperm from the testis tubules (Figure 7). In control males, three

weeks post injection the regions of spermatogenesis were recognizable. Small spermatocysts

with spermatogonia undergoing mitotic divisions were observable at the anterior tip of the testis

tubule (Figure 7A and B). Posterior to the spermatogonia were the primary spermatocytes with

more diffuse nuclear structure. We often observed the band of α-pHH3 positive spermatocysts

that indicated the first meiotic division to form the secondary spermatocytes. At the most

posterior end of the testis tubule the spermatids developed into spermatozoa. Three weeks after

Dnmt1 knockdown testis tubule structure is significantly disrupted. There appeared to be fewer

spermatogonia and those that remained had a more condensed nuclear structure than in the testis

tubules of control males (Figure 7C and D). The most noticeable effect was seen in the region

containing the spermatocytes. There were few primary spermatocytes in the testis tubule and α-

pHH3 positively stained spermatocysts were rarely observed, indicating few spermatocysts

undergoing meiosis.

Discussion

The function of DNA methylation and the DNA methylation enzymes across the insect tree of

life has been widely debated, particularly in light of the extreme variation in presence of this

chromatin modification and the diversification of the enzymatic toolkit required to methylate

DNA *de novo* and maintain DNA methylation patterns (Bewick et al. 2016, Lyko 2018, Lewis et

al. 2020). This evolutionary pattern is particularly intriguing given that in two species on

divergent branches of the insect phylogeny the maintenance methyltransferase DNMT1 is

essential to gametogenesis, and thus is tightly tied to fitness, even in a species where DNA

methylation itself is absent (Schulz et al. 2018). While functional studies have not been done in

other species, a summary of expression studies on the DNA methyltransferases across a diversity

of insects suggest that these enzymes could be involved in gametogenesis across a number of

species, ranging from fire ants, *Solenopsis invicta* (Kay et al. 2018), to jewel wasps, N*asonia*

vitripennis (Zwier et al. 2012), to migratory locusts, *Locusta migratoria* (Robinson et al. 2016),

and the brown plant hopper, *Nilaparvata lugens* (Zhang et al. 2015). Here we showed that *Dnmt1*

expression affects germ cell development. The effect of downregulating *Dnmt1* in males, as in

females (Amukamara et al. 2020), was specific to the germ cells. The downregulation of *Dnmt1*

did result in reduction of DNA methylation, but there were no obvious morphological impacts to

the RNAi-treated individuals. Our results show that the function of *Dnmt1* in germ cells is

conserved across the sexes within a species, and other results document that *Dnmt1* expression is

required for oogenesis among species. This raises a conundrum when we look across the insect

tree of life. How can a gene required for such a fundamental fitness activity in examples from

such a diverse group of insects also be so evolutionarily labile? This suggests that the function

Dnmt1 in germ cell development is easily lost or replaced.

Dnmt1 knockdown causes inhibition of spermatogenesis consistent with a role in meiosis

If our hypothesis on the role of *Dnmt1* in meiosis is correct, then spermatogenesis should be

affected by *Dnmt1* expression as well as oogenesis, as it is a shared process across

gametogenesis during sexual reproduction. In the testes *O. fasciatus* males, we observed both

spermatogonia and primary spermatocytes that stain positively for phosphorylation of the serine

10 residue of histone protein H3. While the pattern of histone H3 phosphorylation across meiosis

has not been specifically studied in *O. fasciatus*, phosphorylation of the serine 10 is typically

associated with chromosome condensation during meiosis (Hans and Dimitrov 2001) and mitosis

(Prigent and Dimitrov 2003). Moreover, the pattern of α-pHH3 staining along the axis of the

testis tubule is consistent with this histone modification occurring in both mitosis and meiosis.

We observed α-pHH3 positive cells both within spermatocysts containing the mitotically

dividing spermatogonia and also at the boundary between primary and secondary spermatocytes.

We have confirmed this landmark using *Boule* knockdown males. In *Drosophila*, *Boule* is

required for progression of spermatocytes through meiosis. Germ cells lacking functional *Boule*

become arrested at meiotic prophase (Eberhart et al. 1996). In the testes of *O. fasciatus* males

treated with ds-*Boule*, the affected cells were all posterior to the band of synchronous α-pHH3

positive spermatocytes. Thus, α-pHH3 staining provided a landmark for entry into meiosis.

We did not see the band of α-pHH3 positive spermatocysts at the border between primary and

secondary spermatocytes in the *Dnmt1* knockdown males treated at the 3rd instar stage of

development. This could be interpreted as the primary spermatocytes have not initiated

chromosome condensation. However, we did observe individual α-pHH3 positive nuclei in the

posterior testis tubule. So there could have been arrest after chromatin condensation with a

breakdown in spermatocyst structure. Interestingly, the testis phenotype of our *Dnmt1*

knockdown males resembled the phenotypic effect of *Vasa* knockdown reported by Ewen

Campen et al. 2013. *Vasa* RNAi results in defects in cyst integrity and Ewen Campen et al.

(2013) propose that *Vasa* plays a specific role in the onset or synchrony of meiosis. *Vasa* has also

been proposed to be required for the correct progression through meiosis in mice (Tanaka et al.

2000) and humans (Medrano et al. 2012). Thus, the phenotypic similarity between *Dnmt1*

knockdown and *Vasa* knockdown testes in *O. faciatus* support the hypothesis that *Dnmt1* may be

required for the successful initiation and completion of meiosis in spermatocytes. More work

remains to determine exactly what stage of sperm development is impacted by the decrease in

Dnmt1 expression.

The phenotype of knockdown of *Dnmt1* is exacerbated when knockdown occurs prior to the

stage of testis development when meiosis occurs. This result is consistent with what was

observed in female *O. fasciatus* (Amukamara et al. 2020) in which knockdown of *Dnmt1*

expression prior to the stage of oocyte development inhibited all oocyte formation. One

interpretation of this result is that *Dnmt1* is required for successful progression through meiosis.

However, it is clear that reducing *Dnmt1* expression had an impact beyond simply reducing the

ability to enter or complete meiosis. A block to progression through meiosis, as demonstrated by

the *Boule* knockdown, spermatogonia and primary spermatocytes would continue to be born but

remain viable after a failure to complete meiosis. In the *Dnmt1* knockdowns, however, there

were fewer germ cells of any type, and those that remained in the testis tubule often appeared to

have abnormal and condensed nuclei. It remains to be determined if *Dnmt1* is required for

meiosis or if the association with meiosis is a correlation; *Dnmt1* could be required for viability

of germ cells and act at the stage of development at which they would be undergoing meiosis.

Previous studies on the function of *Dnmt1* in insect spermatogenesis have not documented any

effect on male fertility. Knockdown of *Dnmt1* in the red flour beetle, *T. castaneum* (Schulz et al.

2018) had no effect on the fecundity of female mating partners was measured in the first nine

days post mating and the authors conclude that *Dnmt1* is not required for spermatogenesis in *T.*

castaneum. We suspect that the function of *Dnmt1* in spermatogenesis is not unique to *O.*

fasciatus, given the expression patterns of *Dnmt1* in other insects (Robinson et al. 2016, Kay et

al. 2018). In the *T. castaneum* study, the authors did not examine testis structure or design a

mating strategy that would exhaust the sperm stores of the males and so may have missed a

fertility effect. In holometabolous insects, including beetles, spermatocytes are typically formed

by the end of larval development (Economopoulos & Gordon 1971), thus males treated as pupae

would still emerge with large numbers of sperm in their testes. Here we showed that male *O.*

fasciatus treated with ds-*Dnmt1* RNA as sexually mature adults become sperm limited more

rapidly than control males. There is a potential alternative explanation, that *Dnmt1* knockdown

males could continue to produce sperm but that sperm was of low quality and unable to support

development. Given that the testes of ds-*Dnmt1* treated males were significantly smaller than

control males and had altered testis tubule structure, which indicated that spermatogenesis was

significantly impacted, we concluded that *Dnmt1* knockdown males are unable to replenish

sperm supplies. The functional role of *Dnmt1* in testes across other species remains to be tested.

DNA methylation and germ cell development

The knockdown of *Dnmt1* resulted in a reduction of DNA methylation within the testes, as predicted. The effect was greater in the nymphs that were treated in an earlier stage of development, as would be expected given the greater number of cell divisions that would occur between treatment and collection of the testes between the 3^{rd} and 5^{th} instar treatments. The more extreme phenotype in the 3^{rd} instar treated males could be explained by the greater reduction in methylated CpG. However, the percent methylation seen in the testes of the 5^{th} instar ds-*Dnmt1* treated males was similar to that seen in females treated at the 4^{th} instar stage of development (Amukamara et al. 2020), greater than a two-fold reduction in the percent methylated CpG residues. In *Dnmt1* knockdown females, there was a compete loss of oocyte production when DNA methylation was reduced to this level while in males the phenotype of the 5^{th} instar *Dnmt1* knockdowns was close to normal. This disconnect between reduction of DNA methylation and phenotypic effects mirrors what has been seen in the beetle, *T. castaneum* (Schulz et al. 2018). In this species, the reduction of DNA methylation is an evolved difference rather than an experimental effect, but the results are similar to what we have observed in *O. fasciatus*; DNA methylation is not required for function of somatic cells, but downregulation of *Dnmt1* expression leads to specific germ cell effects. The lack of correlation between extent of methylation and function in germ cells strengthens our hypothesis that there may be a pleiotropic function for *Dnmt1* in germ cells that acts independently of DNA methylation levels. Alternatively, DNA methylation may have a specific role in the germ cells that it does not play in somatic cells. Perhaps DNA methylation is required for proper gene expression in germ cells or it may be required for transposon silencing only in germ cells. However, demethylation of ovarian tissue in *O. fasciatus* showed little relationship between methylation levels and gene

expression, with few differentially transcribed genes (Bewick et al. 2019). Another possibility is

that DNA methylation state may be a prerequisite for successful completion of meiosis and not

mitosis. For example, *O. fasciatus* has holocentric chromosomes and undergoes inverted meiosis

(Viera et al. 2009). During mitosis chromosomes have a kinetocore that extends across the

majority of the chromosome but in meiosis, the kinetochore plate is missing and microtubules

extend into the chromosome (Comings & Okada 1972). One could imagine that DNA

methylation might be required for this interaction. However, the complete lack of methylation

but a functional role of *Dnmt1* in *T. casteneum* oogenesis alongside our results argues for a

pleiotropic function for *Dnmt1* in gametogenesis that is independent of its role in DNA

methylation.

Conclusion

Dnmt1 expression is required for germ cell development in both male and female *O. fasciatus*.

The block to germ cell development in *Dnmt1* knockdowns appeared to be associated with

meiosis, although it was not a simple block to progression through meiosis as germ cells are lost

from the testes. Thus, *Dnmt1* may be required for germ cell viability. It still remains to be

determined if the block in gametogenesis depends on DNA methylation or an alternative

function of *Dnmt1*, as suggested by the requirement for *Dnmt1* during gametogenesis in an insect

with a non-methylated genome (Schultz et al. 2018). Whatever the specific function of *Dnmt1* in

gametogenesis, the requirement for this enzyme in such a critical fitness function as the

production of gametes in representatives of different groups of insects raises important questions

as to how this enzyme and the entire methylation toolkit has evolved across the insect tree of life

(Lyko et al 2018, Provataris et al. 2018). Functional analysis of *Dnmt1* in the insect groups

where it is found, and study of how these functions are replaced in the species where it is no

longer found, will be essential for understanding the evolution of this important base

modification.

Materials and Methods

Animal care

All experimental animals were from colonies of laboratory reared *O. fasciatus* (Carolina

Biologicals, Burlington, NC) and were reared under standard rearing conditions of 12 hr:12 hr

light/dark at 27° C. To collect animals of known age and social conditions, eggs were removed

from the mass colonies and allowed to hatch in plastic storage containers containing *ad libitum*

deionized water and organic, raw sunflower seeds. For the nymph injections, nymphs were

pulled from mixed sex nymph colonies at the 3rd instar or 5th instar. For adult injections, nymphs

were separated by sex at the 4th instar and housed in single sex colonies. These were checked

daily and newly emerged adults. All experimental animals were placed into individual petri

dishes with food and water.

Developmental expression

Given our hypothesis that *Dnmt1* is required for development of germ cells and meiosis we

examined the expression levels of *Dnmt1* across testis development in males. Groups of nymphs

were staged and sexed. Testes were dissected from 4th and 5th instar nymphs and flash frozen in

liquid nitrogen and stored at -80° C. We also collected testes from virgin males on the day of

adult emergence and after sexual maturation at 7 days post-adult-emergence. We also collected

head, gut, and muscle samples from the sexually mature adult male individuals for tissue specific

expression.

Total RNA was extracted using a Qiagen RNA easy kit with Qiazol (Qiagen, Venlo, The

Netherlands) and complementary DNA (cDNA) synthesized from 500 ng RNA with aScript

cDNA Super-Mix (Quanta Biosciences, Gaithersburg, MD). Quantitative real-time PCR (qRT-

PCR) was used to determine expression levels of *Dnmt1* and two genes with known functions in

spermatogenesis, *Boule* and *Vasa*. Primers are described in Amukamara et al. 2020. As in

Amukamara et al. 2020, actin and GAPDH were used as reference genes. We have previously

validated these reference genes and they are accepted as robust reference genes in *O. fasciatus*

(Meinzer et al. 2019). We used a Roche LightCycler 480 with the SYBR Green Master Mix

(Roche Applied Science Indianapolis, IN). All samples were run with 3 technical replicates using

10 μL reactions. There were 10 biological replicates for each stage. Each biological replicate of

2^{nd}, 3^{rd}, 4^{th}, and 5^{th} instar nymphs consisted of pools of 10, 5, 4, and 3 individuals, respectively.

Adult replicates consisted of individual animals. Sample size was based on past experience

balanced by the cost of qRT-PCR. Primer efficiency calculations, genomic contamination

testing, and endogenous control gene selection were performed as described in Cunningham et

al. 2014. We used the ΔΔCT method to compare levels of gene expression across the samples

(Livak and Schmittgen 2001). Gene expression was standardized per individual to account for

different numbers of individuals within each group at each developmental stage. Differences in

expression levels were analyzed using ANOVA in JMP Pro v14. If there was a significant

overall effect, we compared means using Tukey-Kramer HSD.

RNAi preparation

Double-stranded RNAs were prepared as described in Amukamara et al. 2020. Briefly, DNA

templates were prepared by PCR using gene-specific primers (Amukamara et al. 2020). Sense

and anti-sense RNA were transcribed together with an Ambion MEGAscript kit (ThermoFisher

Sci, Waltham, MA) and allowed to anneal to form a 404 bp ds-*Dnmt1* RNA. The concentration

of dsRNA was adjusted to 3 μg/μL in injection buffer (5 mM KCl, 0.1 mM NaH_2PO_4).

333   Nymph injections, testis size and morphology

To examine the effect of *Dnmt1* knockdown prior to or following the wave of meiosis initiated in

the 4th instar stage, nymphs were injected with ds-*Dnmt1* or control ds-RED injections at either

the 3rd instar or 5th instar stage of development. For further information on controls and testing

for potential off-target effects please see Amukamara et al. 2020. Nymphs were anaesthetized at

4° C for 20 minutes prior to injection. Nymphs were injected in the abdomen using pulled glass

capillary needles (Sutter Instrument Company micropipette puller model P-97, Novato, CA)

between the third and fourth abdominal segments (Chesebro et al. 2009). Nymphs were injected

with 2 μL volume for all injections. Following injections, nymphs were placed in individual petri

dishes and monitored for development. Date of adult emergence was recorded. We did not do a

power analysis, but based on preliminary data on the strength of the effect we aimed for 25

individual males for each treatment. Males were randomly assigned to a treatment group. Not all

males survived to age of dissection, resulting in final sample sizes of 22 and 19 for 3rd instar

males injected with buffer and ds-*Dnmt1*, respectively and 25 and 20 for 5th instar males injected

with buffer and ds-*Dnmt1*, respectively.

Testis size: At 7 to 10 days post-adult emergence, virgin males were dissected and their testes

were removed. Whole testes were allowed to settle into 1 mL Phosphate Buffered Saline (PBS)

and were imaged with a Leica M60 Stereomicroscope with Leica Application Suite software

(LAS v4). Testis area was measured on one of the pair from each male with the LAS by

outlining the whole testis with all 7 testis tubules. Differences in testis area were analyzed using

ANOVA in JMP Pro v14.

Testis tubule structure: *O. fasciatus* testes contain 7 individual testis tubules surrounded by a

relatively impermeable, autofluorescent membrane. Individual testis tubules were removed from

the outer membranous sheath for fixation and staining. Males from each treatment were dissected

across 4 dissection days. Testis tubules from individual males within a treatment were pooled for

staining. Thus each day one tube was a replicate with tubules from several individual males.

Tubules were fixed for 30 minutes in 4% formaldehyde in PBS plus 0.1% Triton-X100 (PBT)

and stained for evidence of cell division using an α-phosphohistone H3 Ser10 (pHH3) primary

antibody (Millipore antibody 06-570, Sigma-Aldrich, St. Louis, MO). α-phosphohistone H3

(pHH3) stains for chromosome condensation in preparation for mitosis and meiosis (Hans &

Dimitrov 2001, Prigent & Dimitrov 2003). The secondary antibody was an Alexa Fluor goat-

anti-rabbit 647 (ThermoFisher Scientific, Waltham, MA). Following antibody staining the

tubules were stained with DAPI (0.5 μg/mL PBT) to visualize nucleic acids. Stained tubules

were mounted in Mowiol 4-88 mounting medium (Sigma-Aldrich, St. Louis, MO) and visualized

with an Olympus BX51 Fluorescent microscope. Images were taken of every testis tubules

present on each slide. Representative images are presented in the figures.

Quantitative Real Time PCR

While we had evidence that all our RNAi treatments successfully knocked down expression in

females (Amukamara et al. 2020), to confirm that our RNAi treatment was effective in males,

total RNA and genomic DNA was extracted from flash frozen testes of a subsample of males 6

to 10 days following injection with the dsRNA using a Qiagen Allprep DNA/RNA Mini Kit

(Qiagen, Venlo, The Netherlands). Expression levels for *Dnmt1* was analyzed using qRT-PCR as

described above.

Quantification of DNA methylation

The DNA from each prep used for qRT-PCR was used to prepare MethylC-seq libraries as

described in Urich et al. 2015 and Amukamara et al. 2020. Samples were sequenced on a

NextSeq500 and qualified reads were aligned to the *O. fasciatus* genome assembly according to

previously published methods (Schmitz et al. 2013). The percent DNA methylation was

calculated by dividing the total number of methylated CpG sites by the total number of CpG sites

(Schultz et al. 2012). Spiked in Lambda DNA, which is fully demethylated, was used as a control

for the sodium bisulfite conversion rate of unmodified cytosines. Three individuals were

sequenced for each treatment.

Adult injections

RNAi treatment: Sexually mature virgin males (7 days post-adult-emergence) were injected with

3 μL ds-*Dnmt1* RNA or control injections using a pulled glass capillary needle between the third

and fourth abdominal segments (Chesebro et al. 2009). Control injections for the male fecundity

experiments were injection buffer alone. We used ds-RED control injections for the testis

imaging experiment. Previous studies from our lab have shown no difference in buffer alone

controls or non-specific ds-RNA (Bewick et al. 2019). Following injection males were placed

into individual petri dishes and provided with *ad libitum* food and water. We did not do a power

analysis, but based on preliminary data on the strength of the effect we aimed for 30 individual

males for each treatment. Males were randomly assigned to a treatment group.

Male fecundity: Preliminary experiments had shown that two previous matings were required to

deplete sperm stores. Therefore, in order to allow males to deplete sperm stores acquired during

nymphal development and sexual maturation, males were placed in mating trials with three 7- to

10-day-old virgin females, one provided each week. The first female was placed in the male's

petri on the day of injection with cotton wool as an oviposition substrate. The female and all eggs

were removed at the end of the week (7 days post-injection) and discarded and replaced with a

second 7- to 10-day-old virgin female and fresh cotton wool. Again, at the end of the week (14

days post-injection), the female and all resulting eggs were discarded. A third, focal, 7- to 10-

day-old virgin female was placed with the male with fresh cotton wool. The 3rd female and

experimental male were given one week to mate before the male was removed from the petri

dish for analysis of testis size and structure. The female was maintained for her lifetime,

provided with *ad libitum* food and water. The eggs produced by the 3rd female were collected

from the petri dish twice a week at 3-4 day intervals and the oviposition substrate replaced with

fresh cotton wool. The eggs collected were placed in a separate container and allowed to develop

to hatching (approximately 7-10 days following collection) and then frozen at -20° C until

assayed. We recorded the first collection date for which no eggs hatched. We analyzed the time

to end of fertilized eggs due to sperm depletion relative to treatment using a survival analysis

(Wilcoxon Rank Sum test) using JMP Pro v14.1. Not all males survived to date of dissection, or

their mates died during the course of egg collection, requiring these males to be removed from

the analysis, resulting in a sample size of 21 control and 20 ds-*Dnmt1* treated males in the final

fecundity analysis.

Testis size: At the end of the one-week mating trial with the 3rd female (21 days post-injection),

males were dissected and their testes removed into 1 mL Phosphate Buffered Saline (PBS).

Whole testes were photographed and measured as described above. Not all males survived to

date of dissection, resulting in a final sample size of 28 control and 25 ds-*Dnmt1* treated males.

Testis tubule structure: Testis tubules from mated males were isolated, fixed and stained as

described above.

Acknowledgments

The authors acknowledge Allen J. Moore and other members of the Moore lab for many helpful
discussions. We acknowledge Luvika Gupta, who's undergraduate research provided preliminary
data used in our experimental design. We also would like to acknowledge Tyler Earp for
preparing and analyzing low throughput whole genome bisulfite sequencing data.

Competing Interests

The authors have no competing interests in regards to this work.
429

References

Amukamara A.U., J. T. Washington, Z. Sanchez, E.C. McKinney, A. J. Moore, R. J. Schmitz, P. J. Moore. More than DNA methylation: does pleiotropy drive the complex pattern of evolution of Dnmt1? *Frontiers in Ecology and Evolution* 8, 4 (2020).

Bewick, A.J., K. J. Vogel, A. J. Moore, R. J. Schmitz. Evolution of DNA methylation across insects. *Molecular Biology and Evolution* 34, 654-66 (2016).

Bewick, A.J., Z. Sanchez, E. C. Mckinney, A. J. Moore, P. J. Moore, R. J. Schmitz. *Dnmt1* is essential for egg production and embryo viability in the large milkweed bug, *Oncopeltus fasciatus*. *Epigenetics & Chromatin* 12, 6 (2019).

Chesebro, J., S. Hrycaj, N. Mahfooz, A. Popadić. Diverging functions of Scr between embryonic and post-embryonic development in a hemimetabolous insect, *Oncopeltus fasciatus*. *Developmental Biology* 329, 142-151 (2009).

Comings, D.E., T. A. Okada. Holocentric chromosomes in *Oncopeltus*: kinetochore plates are present in mitosis but absent in meiosis. *Chromosoma* 37, 177-192 (1972). https://doi.org/10.1007/BF00284937

Cunningham, C.B., L. Ji, R. A. W. Wiberg, J. Shelton, E. C. McKinney, D. J. Parker, R. B. Meagher, K. M. Benowitz, E. M. Roy-Zokan, M. G. Ritchie, S. J. Brown, R. J. Schmitz, A. J. Moore. The genome and methylome of a beetle with complex social behavior, *Nicrophorus vespilloides* (Coleoptera: Silphidae). *Genome Biology and Evolution* 7, 3383-3396 (2015).

Dallai, R. Overview on spermatogenesis and sperm structure of Hexapoda. *Arthropod Structure & Development* 43, 257-290 (2014). https://doi.org/10.1093/gbe/evy066

Eberhart, C.G., J. Z. Maines, S. A. Wasserman. Meiotic cell cycle requirement for a fly homologue of human Deleted in Azoospermia. *Nature* 381, 783-785 (1996).

Economopoulos, A.P. H. T. Gordon. Growth and differentiation of the testes in the large milkweed bug, *Oncopeltus fasciatus* (Dallas). *Journal of Experimental Zoology* 177, 391-405 (1971).

Ewen-Campen, B., T. E. Jones, C. G. Extavour. Evidence against a germ plasm in the milkweed bug *Oncopeltus fasciatus*, a hemimetabolous insect. *Biology Open* 2, 556-568 (2013).

Gegner, J., T.Gegner, H. Vogel, A. Vilcinskas. Silencing of the DNA methyltransferase 1 associated protein 1 (DMAP1) gene in the invasive ladybird *Harmonia axyridis* implies a role of the DNA methyltransferase 1-DMAP1 complex in female fecundity. *Insect Molecular Biology* 29, 148-159 (2020). https://doi.org/10.1111/imb.12616

Glastad, K.M., B. G. Hunt, M. A. Goodisman. Epigenetics in insects: genome regulation and the generation of phenotypic diversity. *Annual Review of Entomology* 64, 185-203 (2019). doi.org/10.1146/annurev-ento-011118-111914

Hans, F., S. Dimitrov. Histone H3 phosphorylation and cell division. *Oncogene* 20, 3021-3027 (2001).

Kay. S., D. Skowronski, B. G. Hunt. Developmental DNA methyltransferase expression in
the fire ant Solenopsis invicta. Insect Science 25, 57-65 (2018).

Lewis S.H., L. Ross, S. A. Bain, E. Pahita, S. A. Smith, R. Cordaux, et al. Widespread
conservation and lineage-specific diversification of genome-wide DNA methylation
patterns across arthropods. *PLoS Genetics* 16, e1008864 (2020).
https://doi.org/10.1371/journal.pgen.1008864

Livak, K.J., T. D. Schmittgen. Analysis of relative gene expression data using real-time
quantitative PCR and the 2− ∆∆CT method. *Methods* 25, 402-408 (2001).

Lyko, F. The DNA methyltransferase family: a versatile toolkit for epigenetic regulation.
Nature Reviews Genetics 19, 81-92 (2018).

Medrano, J.V., C. Ramathal, H. N. Nguyen, C. Simon, R. A. Reijo Pera. Divergent RNA-
binding proteins, DAZL and VASA, induce meiotic progression in human germ cells
derived in vitro. *Stem cells* 30, 441-451 (2012).

Meinzer, F., S. Dobler, A. Donath, J. N. Lohr. Robust reference gene design and validation
for expression studies in the large milkweed bug, *Oncopeltus fasciatus*, upon cardiac
glycoside stress. *Gene* 710, 66-75 (2019).

Prigent, C. S. Dimitrov. Phosphorylation of serine 10 in histone H3, what for?. *Journal of
Cell Science* 116, 3677-3685 (2003).

Provataris, P., K. Meusemann, O. Niehuis, S. Grath, B. Misof. Signatures of DNA
methylation across insects suggest reduced DNA methylation levels in holometabolan.
Genome Biology and Evolution 10, 1185-1197 (2018).

Robinson, K.L., D. Tohidi-Esfahani, F. Ponton, S. J. Simpson, G. A. Sword, N. Lo.
Alternative migratory locust phenotypes are associated with differences in the
expression of genes encoding the methylation machinery. *Insect Molecular Biology* 25,
105-115 (2016).

Schmidt, E. D., E. Sehn, A. Dorn. Differentiation and ultrastructure of the spermatogonial
cyst cells in the milkweed bug, *Oncopeltus fasciatus*. *Invertebrate Reproduction &
Development* 42, 163-178 (2002). DOI:10.1080/07924259.2002.9652773

Schmitz, R. J., Z. A. Lewis, M. G. Goll. DNA methylation: shared and divergent features
across eukaryotes. *Trends in Genetics* 35, 818-827 (2019).
doi.org/10.1016/j.tig.2019.07.007

Schmitz, R.J., Y. He, O. Valdés-López, S. M. Khan, T. Joshi, M. A. Urich, J. R. Nery, B.
Diers, D. Xu, G. Stacey, J. R. Ecker. Epigenome-wide inheritance of cytosine
methylation variants in a recombinant inbred population. *Genome Research* 23, 1663-
1674 (2013).

Schultz, M.D., R. J. Schmitz, J. R. Ecker. 'Leveling' the playing field for analyses of single-
base resolution DNA methylomes. *Trends in Genetics* 28, 583-585 (2012).
DOI:https://doi.org/10.1016/j.tig.2012.10.012

Schulz, N.K., C. I. Wagner, J. Ebeling, G. Raddatz, M. F. Diddens-de Buhr, F. Lyko, J.
Kurtz. Dnmt1 has an essential function despite the absence of CpG DNA methylation

509 in the red flour beetle *Tribolium castaneum*. *Scientific Reports* 8, 1-10 (2018).
510 https://doi.org/10.1038/s41598-018-34701-3

511 Shah, C., M. J. VanGompel, V. Naeem, Y. Chen, T. Lee, N. Angeloni, Y. Wang, Y. E. Y.
512 Xu. Widespread presence of human BOULE homologs among animals and
513 conservation of their ancient reproductive function. *PLoS Genetics* 6, e1001022 (2010).
514 doi.org/10.1371/journal.pgen.1001022

515 Tanaka, S.S., Y. Toyooka, R. Akasu, Y. Katoh-Fukui, Y. Nakahara, R. Suzuki, M.
516 Yokoyama, T. Noce. The mouse homolog of Drosophila Vasa is required for the
517 development of male germ cells. *Genes & Development* 14, 841-853 (2000).

518 Ulrich, M. S., J. R. Nery, R. Lister, R. J. Schmitz, J. R. Ecker. MethylC-seq library
519 preparation for base-resolution whole-genome bisulfite sequencing. *Nature Protocols*
520 10, 475–483 (2015). doi: 10.1038/nprot.2014.114

521 Viera, A., J. Page, J. Rufas. Inverted meiosis: the true bugs as a model to study. In:
522 Benavente, R., Volff J.-N. (eds): Meiosis. Genome Dynamics, Basel, Karger, 2009, vol
523 5, pp 137-156 (2009). https://doi.org/10.1159/000166639

524 Yajima, M., G. M. Wessel. The multiple hats of Vasa: its functions in the germline and in
525 cell cycle progression. *Molecular reproduction and development* 78, 861-867 (2011)

526 Zemach, A., I. E. McDaniel, P. Silva, D. Zilberman. Genome-wide evolutionary analysis of
527 eukaryotic DNA methylation. *Science* 328, 916-919 (2010).

528 Zhang, J. Y. Xing, Y. Li, C. Yin, C. Ge, F. Li. DNA methyltransferases have an essential role
529 in female fecundity in brown planthopper, Nilaparvata lugens. Biochemical and
530 Biophysical Research Communications 464, 83-88 (2015).

531 Zwier, M. V., E. C. Verhulst, R. D. Zwahlen, L. W. Beukeboom, L. van de Zande. DNA
532 methylation plays a crucial role during early Nasonia development. *Insect Molecular*
533 *Biology* 21, 129-138 (2012).
534

Figure Legends

Figure 1. Progression of spermatogenesis in *O. fasciatus*. In *O. fasciatus* spermatogenesis progresses from the tip of the testis tubule. (A) Dapi-stained testis tubule of stock male. (B) Diagram of the stages of spermatogenesis. At the apical tip (region I) of each of the seven testis tubules there is a "rosette" of niche cells (blue) surrounded by the germline stem cells (B, green; Schmidt et al. 2002). The rosette is not typically visible in confocal images of testis tubules and markers for the rosette have not been identified yet. As spermatogonia (B, light gray) arise from division of the germline stem cells, they are enclosed by cyst cells (B, yellow). In region II, spermatogonia undergo mitotic transit amplification divisions to form spermatocysts containing 64 spermatogonia (Economopoulos & Gordon 1971, Ewen-Campen et al. 2013). Spermatocytes (B, dark gray) in region III divide meiotically. *Oncopeltus fasciatus* undergoes inverted meiosis (Viera et al. 2009). Primary spermatocytes undergo the first meiotic division to produce diploid secondary spermatocytes. The meiotic division of the secondary spermatocytes produce the haploid spermatids (B, dark blue) in region IV that then differentiate into spermatozoa in the region V at the terminal end of the testis tubule.

Figure 2. Expression pattern of *Dnmt1* within the developing testes mirrored that the two gametogenesis genes *Boule* and *Vasa*. *Dnmt1* expression (A) was highest in the larval stages where spermatogenesis is initiated compared to the testes of newly emerged (day 0) and virgin sexually mature (day 7-10) adults. This is similar to the expression patterns of *Boule* (B), in which expression was higher in the 4th and 5th larval stages during which the transition to meiosis occurs compared to newly emerged and virgin sexually mature adults. Expression of *Vasa* (C) demonstrated a similar pattern, with the highest expression in the 4th instar stage and lowest in sexually mature virgin males. Black dots and bars represent mean and SE. Grey dots represent

data points for each individual tested. Lower case letters designate significant differences ($p <$

0.05) among pairwise comparisons using *post-hoc* Tukey-Kramer HSD test.

Figure 3. Expression of *Dnmt1* was reduced in the testes of adult males treated with ds-*Dnmt1* at

both stages of development. Relative gene expression is standardized to expression levels in

control treatments. Black dots and bars represent mean and SE. Grey dots represent data points

for each individual tested.

Figure 4. Downregulation of *Dnmt1* during the 3rd instar stage of development significantly

reduced the size of the testis in sexually mature males. There was no effect of downregulation of

Dnmt1 on testis size when treatment occurred during the 5th instar stage of development after

meiosis has been initiated. Black dots and bars represent mean and SE. Grey dots represent data

points for each individual tested.

Figure 5.

Timing of treatment during development determined the effect on testis structure of adults that

develop from *Dnmt1* (Panels B and D) knockdown males. In control testis tubules from males

injected with ds-RED (A and C), spermatogonia divided mitotically in region II to form

spermatocysts as is observed in untreated males. Mitotic spermatogonia are labeled with anti-

phosophohistone 3 antibodies in region II of the testis tubule (C, arrowhead). Meiotic divisions

also occured as in untreated males in ds-RED injected males. In our ds-RED control samples, a

band of synchronously dividing spermatocysts, identified by positive anti-phosphohistone H3

antibody staining, was present at the interface between primary and secondary spermatocytes (A,

C). In testes from males with *Dnmt1* knockdown in the 3rd instar (B), the anterior tip of the testis

tubule showed evidence of mitotic activity (arrowhead). However, there were fewer

spermatocytes present and the spermatocysts in this region were disorganized and there was little

evidence of the band of positive anti-phosphohistone H3 stained meiotic spermatocytes. The

testis tubule structure from males treated with ds-*Dnmt1* following meiosis at the 5th instar stage

of development (D) was more similar to the controls than those treated at the 3rd instar stage of

development. In ds-*Dnmt1* males treated at the 5th instar stage, there were positive anti-

phosphohistone H3 stained spermatogonia (arrowhead) and spermatocytes (arrow). There were

differences between males treated as 5th instars with ds-RED males and ds-*Dnmt1*, however,

including spermatocysts containing cells with highly condensed nuclei that were not present in

the controls and fewer spermatocysts containing spermatids or spermatozoa. All images taken at

10X magnification.

Figure 6. Females mated to control males lay fertilized eggs longer than females mated to

Dnmt1 knockdown males. Eggs were collected twice per week and eggs allowed to develop to

hatching. Eggs that did not hatch showed no sign of development, indicating that they had not

been fertilized. Both treatments demonstrate a decrease in proportion of eggs that hatch over

time, but the ds-*Dnmt1* treatment group shows a faster decrease.

Figure 7. Testis structure breaks down in *Dnmt1* knockdown males treated as adults after having

sperm replenishment induced by mating activity. The regions of spermatogenesis were apparent

in mated males following 3 weeks of mating activity in control testis tubules (A and B) and the

evidence of both mitotic division in spermatogonia (B, arrowhead) and the band of meiotic

divisions was clear from anti-phosphohistone H3 staining. In *Dnmt1* knockdown males at low

magnification (C), the anterior tip of the testis tubule looked relatively normal. However, the

region containing both the primary and secondary spermatocytes was disorganized.

Spermatocyst structure was broken down, and the nuclei of the primary and secondary

spermatocytes had lost their characteristic structure (Ewen-Campen et al. 2013). Finally, there

were fewer mature spermatids in the posterior end of the testis tubule. At higher magnification

(D), it was apparent that nuclear structure in the anterior tip was also affected by the knockdown,

both in the spermatogonia and spermatocytes. Spermatogonia nuclei in the *Dnmt1* knockdown

testis tubules (D) were more condensed than in the control testis tubules (B), although they still

seemed to be organized into spermatocysts. Spermatocyte nuclei, however, were fewer in

number than in controls, did not have their characteristic shape (Ewen-Campen et al. 2013,

Economopoulos & Gordon 1971), and were not organized in spermatocytes. A and C:10X

magnification, B and D: 20X magnification.

Figure 1.

Figure 2.

Figure 3

Figure 4.

Figure 5.

Figure 6.

Figure 7.