On Rank Deficiency in Phenotypic Covariance Matrices

F. Robin O'Keefe[1*]

Julie A. Meachen[2]

P. David Polly[3]

1. Marshall University, Biological Sciences, Huntington, WV

2. Des Moines University, Anatomy Department, Des Moines, IA

3. Indiana University, Earth and Atmospheric Sciences, Bloomington, IN

*corresponding author: F. Robin O'Keefe, Professor, Marshall University, College of

Science 265, One John Marshall Drive, Hunington, WV 25755. Phone: +1 304 696 2427.

Email: okeefef@marshall.edu

Running Head: Rank Deficiency in Phenotypic Covariance

Conflict of Interest Statement: The authors state that they have no conflict of interest to

disclose.

Keywords: Dire wolf, *Canis dirus*, geometric morphometrics, phenotypic covariance,

modularity and integration, information entropy, effective rank, effective dispersion,

relative dispersion, latent dispersion.

ABSTRACT

This paper is concerned with rank deficiency in phenotypic covariance matrices: first to establish it is a problem by measuring it, and then proposing methods to treat for it. Significant rank deficiency can mislead current measures of whole-shape phenotypic integration, because they rely on eigenvalues of the covariance matrix, and highly rank deficient matrices will have a large percentage of meaningless eigenvalues. This paper has three goals. The first is to examine a typical geometric morphometric data set and establish that its covariance matrix is rank deficient. We employ the concept of information, or Shannon, entropy to demonstrate that a sample of dire wolf jaws is highly rank deficient. The different sources of rank deficiency are identified, and include the Generalized Procrustes analysis itself, use of the correlation matrix, insufficient sample size, and phenotypic covariance. Only the last of these is of biological interest.

Our second goal is to examine a test case where a change in integration is known, allowing us to document how rank deficiency affects two measures of whole shape integration (eigenvalue standard deviation and standardized generalized variance). This test case utilizes the dire wolf data set from Part 1, and introduces another population that is 5000 years older. Modularity models are generated and tested for both populations, showing that one population is more integrated than the other. We demonstrate that eigenvalue variance characterizes the integration change incorrectly, while the standardized generalized variance lacks sensitivity. Both metrics are impacted by the inclusion of many small eigenvalues arising from rank deficiency of the covariance matrix. We propose a modification of the standardized generalized variance, again based on information entropy, that considers only the eigenvalues carrying non-redundant

information. We demonstrate that this metric is successful in identifying the integration

change in the test case.

The third goal of this paper is to generalize the new metric to the case of arbitrary

sample size. This is done by normalizing the new metric to the amount of information

present in a permuted covariance matrix. We term the resulting metric the 'relative

dispersion', and it is sample size corrected. As a proof of concept we us the new metric to

compare the dire wolf data set from the first part of this paper to a third data set

comprising jaws of *Smilodon fatalis*. We demonstrate that the *Smilodon* jaw is much

more integrated than the dire wolf jaw. Finally, this information entropy-based measures

of integration allows comparison of whole shape integration in dense semilandmark

environments, allowing characterization of the information content of any given shape, a

quantity we term 'latent dispersion'.

[introduction]

The study of integration in biological systems has a long history, stretching back to

Darwin (1859), given a modern footing by Olson and Miller (1958), quantified by Van

Valen (1974) and Cheverud (1982, 1996), and maturing into a broad topic of modern

inquiry encompassing genetics, development, and the phenotype (see Klingenberg, 2013,

and Goswami and Polly, 2010, for reviews). In a biological context, 'integration' means

that traits of a whole organism covary to a significant degree. This covariance is critical

in evolving populations, because traits are not free to respond to selection without

impacting dependent traits, and the directionality of selection response is constrained by

these dependencies (Grabowski and Porto, 2017, Figure 1). Yet trait dependency can also

remove constraint by giving a population access to novel areas of adaptive space

(Goswami et al., 2014, Figure 5). Consequently phenotypic integration is central to the

evolvability of biological systems, because it defines the achievable adaptive space.

Background

The use of landmark data to characterize biological shape, and to quantify shape change,

has become ubiquitous since the introduction of geometric morphometrics by Bookstein

(1997; reviewed in Zelditch et al., 2012). Analysis of geometric morphometric landmark

data begins with Generalized Procrustes Analysis (GPA), wherein the centroid of each

specimen is translated to the origin, and the coordinates are rotated and scaled to a mean

shape so that the intra-landmark variance among specimens is minimized (Bookstein

1997). Given a matrix $\mathbf{LM_{n,v}}$ of v two- or three-dimensional landmark coordinates and n

specimens, a matrix resulting from Procrustes superimposition is:

$$\mathcal{X}_{n,v} = GPA(LM_{n,v})$$

There are myriad examples in the literature of the application of principal

components analysis to GPA-transformed landmark data (for example, Segura et al.,

2020, for canid dentaries; Brannick et al., 2015, for dire wolf dentaries, and O'Keefe et

al. 2014, for dire wolf crania). The usual analysis progression is to calculate the principal

component scores for the first few PCs, and plot the specimens into the principal

component space. Principal components analysis is therefore a powerful tool for

dimensionality reduction, allowing the visualization and biological interpretation of a

large proportion of the variance on one or a few axes. Analyses of this type typically

utilize the first few eigenvectors, so the magnitude of their associated eigenvalues is not a

central concern beyond ensuring their significance. However the eigenvalue vector also

carries biological signal, and analysis of its distribution has given rise to the concept of

whole-shape phenotypic integration.

Whole-Shape Integration Measures

"The 'generalized variance', $|\Sigma|$, the determinant of the variance-co-variance matrix, is

related to the area (or hypervolume) of the equi- probability ellipses (ellipsoids) of the

distribution."—VanValen, 1974, p. 235.

The distribution of the eigenvalue vector has formed the basis of several metrics intended

to quantify the strength of covariation in a whole shape. As stated by VanValen in the

quote above, the shape space containing the objects of interest can be conceptualized as a

hyperellipse in the original variable space. Principal components analysis moves this

hyperellipse to the origin, rotates it, and defines its principal axes. VanValen realized that

this hyperellipse had two salient properties: 1) its dimensionality, and 2) its variance, or

dispersion, on each axis, which he called 'tightness'. In subsequent work the dispersion

of the eigenvalues of a phenotypic covariance matrix—this tightness— has received a

great deal of attention, while its dimensionality has not (reviewed in Najarzadeh, 2019).

Cheverud (1983) introduced the first modern measure of trait integration, defining it as

$$I = 1 - \left(\prod_{i=1}^{n} \lambda_i \right)^{1/n}$$

or the geometric mean of the eigenvalues of the correlation matrix subtracted from unity.

This quantity is identical to one minus the nth root of the generalized variance (Wilks,

1932) proposed by VanValen in 1974. In a wider statistical context this quantity is

termed the 'standardized generalized variance', or SGV (SenGupta, 1987). The SGV is

the geometric mean of the eigenvalue variance, and so may be thought of as the mean

diameter of the axes in the hyperellipse. Both Cheverud (1983) and SenGupta (1987)

state that the SGV is comparable among spaces of different dimensionality, and this

assumption is widely accepted, although it has never been tested (e.g. Najarzadeh, 2019).

While both the Cheverud integration and the equivalent SVG have been used as measures

of dispersion, a consensus has emerged that the standard deviation of the eigenvalues is a

better measure statistically (Pavlicev et al., 2009). Several other metrics have been

proposed for quantifying the dispersion of phenotypic matrices, summarized by Pavlicev

et al., 2009, while the sample size requirements of various metrics are evaluated by

Grabowski and Porto, 2017.

This paper explores the performance of the eigenvalue dispersion and the SGV in

rank-deficient geometric morphometric matrices, and we define these metrics here.

Eigenvalue dispersion is measured as the standard deviation of the eigenvalue vector of

the correlation matrix of GPA landmarks, standardized to the mean eigenvector

(Goswami and Polly, 2010, modified from Equations 7 and 8):

$$\lambda\sigma = \frac{\sqrt{\frac{\sum_{i=1}^{n}\left(\frac{\lambda_i}{\mu\lambda} - 1\right)^2}{n}}}{\sqrt{n-1}} \qquad \textit{Eq. 1}$$

In geometric morphometric data sets, either four (2D) or five(3D) degrees of

freedom are lost to the Procrustes analysis, so the number of non-zero eigenvalues will be

reduced by this amount (Zelditch et al., 2012). This will also be true of the SGV,

calculated as the n^{th} root of the first n-4 eigenvectors of the correlation matrix. Because

we use two-dimensional data sets with 14 landmarks in this paper, the correct formulation

of the SVG is:

$$SGV_{24} = \sqrt[24]{\prod_{i=1}^{24} \lambda_i} \qquad \textit{Eq. 2}$$

Integration and Modularity

"However, if the structure of a dataset is strongly modular, with several different groups

of strongly co-varying traits, then variance will be distributed more evenly across a

number of principal components, and eigenvalue variance will be relatively low. Thus,

the dispersion of eigenvalues provides a simple measure for comparison of the relative

integration or modularity of the structures described in a matrix."—Goswami and Polly,

2010, p. 226.

In the 21[st] century the study of GPA covariance matrices has grown to include the field of

modularity and integration (Goswami et al. 2014; Klingenberg, 2013). This conception of

integration is similar to Van Valen's and Cheverud's in that it studies covariance

structure, but differs in attempting to identify variable subsets that covary relative to

others. Rather than relying on a single overall measure of dispersion, modularity studies

test specific models of landmark covariation against a null model in an attempt to

characterize subsets with high intraset connectivity and low interset connectivity

(Goswami and Polly, 2010). Because these sets are primary covariance structures in the

data, they will exist on the first several principal components, and the entire eigenvalue

distribution is not directly relevant. This approach to modularity has yielded powerful

hypotheses of evolutionary plasticity and constraint linking development to phenotype,

and to evolutionary changes in modularity over deep time (Goswami et al., 2015).

Modularity models were first assessed using the RV coefficient introduced by

Klingenberg (2008). This coefficient has been superseded by the similar covariance ratio

statistic (CR, Adams, 2016), which is more robust to differences in sample size and data

dimensionality. Another common approach to assessing modularity model significance is

Partial Least Squares (PLS; Goswami and Polly, 2010); this approach is more involved

mathematically but has the same goal as the CR statistic (for a recent example on dog and

wolf crania see Curth et al., 2017). As stated in the quote above, the modularity concept

can also be applied to a whole shape. In whole-shape modularity, a shape would be more

integrated if it had greater eigenvalue dispersion, and would be more modular if it

possessed lesser dispersion. This is a specific prediction: in a population where

modularity is evolving, an increase in modularity should lead to a decrease in integration

and hence dispersion, and thus a decrease in dispersion metrics like SVG and eigenvalue

standard deviation.

Rank Deficiency in Phenotypic Covariance Matrices

It [PCA] is used to obtain a more economic description of the N-dimensional dispersion

of the original data by a smaller number of "principal components", which are formally

the eigenvectors of the dispersion matrix... Hence the number of principal components

that contribute significantly to the variation of the sample is the actual "dimensionality"

of the dispersion. – G. P. Wagner, 1984, pp. 92-93.

The rank of a matrix is the dimensionality of the vector space spanned by its variables. If

all variables are completely independent then the matrix is of "full rank," equal to the

number of variables, but when one variable depends completely on another, the matrix

has "deficient rank". The full rank value of a matrix is the number of columns, while the

"real rank" is the maximum number of linearly independent columns. The derivation of

the covariance or correlation matrix **K** directly from **X**,

$$K_{v_i v_j} = cov(x_i x_j)$$

often results in a covariance matrix that is overdetermined; the high degree of

correlation in biological systems often results in matrices whose full rank is much higher

than the real rank (Van Valen, 1974; Adams, 2016). Significant rank deficiency in **K** will

impact the eigenvalue distribution, because eigenvalue distributions of rank deficient

matrices will have tails of very small non-zero eigenvalues. This tail of small eigenvalues

may impact integration metrics that rely on the eigenvalue distribution, but this subject

has not been treated explicitly. Characterizing this problem of rank deficiency, and

developing treatments for it, are the goals of this paper.

In geometric morphometric covariance matrices, rank deficiency can arise from at

least three sources. These are the Generalized Procrustes Analysis itself, lack of matrix

information, and phenotypic covariance. Of these, only phenotypic covariance is of direct

biological interest, while the first two may confound attempts to measure it. The

magnitude of rank deficiency due to Procrustes analysis is known; the translation,

scaling, and rotation performed during the analysis remove four degrees of freedom from

two-dimensional data (Zelditch et al., 2012). A covariance matrix derived from **X** should

therefore possess $v - 4$ non-zero eigenvalues, or in other words be deficient by four ranks

from the full rank.

The rank deficiency resulting from the information intrinsic to **X** is harder to

quantify. This lack of information can have two sources: insufficient sample size and

landmark oversampling. Lack of information due to sample size arises from too few

sampled specimens. Grabowski and Porto (2017) studied the sample size requirements

for a range of integration metrics and concluded that an *n* of greater than 100 was

required to achieve values stable to changes in n. Integration studies with sample sizes of

less than 100 are therefore information poor, and this should increase rank deficiency.

What we term landmark oversampling refers to the dense landmarking of a relatively

simple shape. A simple shape with few modes of variation requires few landmarks to

characterize fully (a circle requires one landmark). Use of more than the necessary

number of landmarks will result in covariance among the variables, and increased rank

deficiency. This source of rank deficiency has also not been quantified. However it is

tractable, as we outline in the Discussion.

220   Another factor impacting matrix rank is use of the correlation matrix rather than

the covariance matrix. Use of the correlation matrix artificially inflates matrix rank,

because each coordinate of each variable is awarded a full rank. This tacitly assumes that

the coordinates within a landmark are uncorrelated, and that variance magnitudes in each

coordinate direction are equivalent. However landmark displacements that are dominated

by a vector parallel to one axis will have trivial variation on the other, and in a correlation

matrix this trivial variation will be awarded a full rank and expanded to a variance of one.

A correlation matrix of **X** will therefore possess a degree of noise from inflation of trivial

variance (for discussion of the desirability of the covariance matrix in morphometric

applications see Goswami and Polly, 2010, p. 217). Therefore use of the correlation

matrix is unwise with morphometric data unless some procedure is used to concatenate

the landmark coordinates into a single measure before analysis (Goswami and Polly,

2010, and previous authors have used the congruence coefficient, although this has been

criticized by Klingenberg 2008).

Information Entropy and Effective Rank

The first to point out the problem of redundancy in the full and real ranks of **K** in a

morphometric context was Van Valen (1974), who tailored an information metric to

correct for it. The problem of rank deficiency can be framed in terms of the eigenvalues

of a matrix, with the true rank represented by the subset of eigenvalues carrying

significant variance. This differs from the set of eigenvalues that carry non-zero variance,

which is how matrix rank is usually defined. The question of the number of 'significant'

eigenvalues is a general question in data analysis and there are many techniques for

determining this number, summarized by Cangelosi and Goriely (2007) in the context of

cDNA microarray data. They include the broken stick model, Cattell's SCREE test, and

Bartlett's sphericity test among many others (see also Hine and Blows, 2006; Bunea et

al., 2011). All these techniques share fundamental shortcomings, such as an integer value

and recourse to an *ad hoc* criterion to assess the cutoff for significance. Yet the problem

of redundancy in eigenvalue distributions is general, and has been solved in signal

processing contexts by the use of "information entropy" (Shannon, 1948). Information

entropy is a metric used to represent the information content of a system by determining

how much a signal can be compressed without loss of information. At least two metrics

based on Shannon's information entropy have been proposed to characterize the

dimensionality of covariance matrices in a biological context (for a thorough

mathematical development see Cangelosi and Goriely 2007). Those authors develop a

dimensionality metric for cDNA microarray data they termed the 'information

dimension'. A very similar metric was developed by Roy and Vetterli (2007); those

authors term their metric the 'effective rank', and use it synonymously with 'effective

dimensionality'. The following application of the Shannon entropy to geometric

morphometrics follows the Roy and Vetterli (2007) development of effective rank.

Given a GPA covariance matrix $\mathbf{K}$ with eigenvalues $\mathbf{\Lambda_V}$:

$$\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_v \geq 0$$

263   The Shannon entropy E_S of $\mathbf{K}$ is defined as

$$E_s = -1 * \sum_{i=1}^{v} \left(\frac{\lambda_i}{\sum \Lambda_V}\right)\left(ln \frac{\lambda_i}{\sum \Lambda_V}\right) \qquad Eq.\ 3$$

266   Note that the eigenvalues are standardized to the trace of $\mathbf{\Lambda}$; this is necessary

267   because the information entropy was originally defined in a probability context, and the

268   terms being evaluated must therefore sum to unity (Shannon, 1948). Roy and Vetterli

269   define their effective rank as e raised to the power E_s; the 'effective rank' of $\mathbf{K}$ is defined

270   in the same way:

$$R_e = e^{Es} \qquad Eq.\ 4$$

This effective rank, R_e, of $\mathbf{K}$ is a continuous metric, is based on solid theoretical

grounds from information theory, and may be thought of as the signficant number of

dimensions of the shape space represented by $\mathbf{K}$, or equivalently as the numner of non-

redundant eigenvalues in $\mathbf{K}$.

Study Design

This paper has three main components, with the later dependent on the results of the

earlier. Therefore we divide the Materials and Methods section into an independent

Materials section, followed by three sections of Methods and Results. This organization

allows sequential presentation of results as they become necessary for further

methodological development. The following sections are:

284   *Materials*—This section introduces the three data sets used in this paper. These

285   comprise two samples of dire wolf jaws from Rancho La Brea (RLB), originating from

286   tar pits deposited about 5000 years apart. The third data set consists of jaws from the

287   sabertooth cat, *Smilodon fatalis*, assembled from multiple RLB pits ($n = 81$). All three

288   data sets comprise 14 two-dimensional landmarks taken from photographs of jaws in

289   lateral view.

290   *Methods and Results, Part 1*—In this section, we use the concept of effective rank

to investigate the rank deficiency of the Pit 61/67 dire wolf population. We demonstrate

that the covariance matrix based on these data is highly rank deficient. Through use of a

permutation test, we characterize the magnitude of the rank deficiency arising from lack

of matrix information. When combined with the rank deficiency expected from the GPA,

this allows us to quantify the rank deficiency due to phenotypic covariance. Lastly, we

quantify the rank inflation arising from use of the correlation matrix.

297   *Methods and Results, Part 2*—Here we evaluate the impact of the observed rank

deficiency on whole shape integration metrics. We do this by introducing the Pit 13 dire

wolf sample, and demonstrating that it is more integrated than the Pit 61/67 sample. This

requires testing each population for candidate modularity models, and demonstrating that

Pit 61/67 wolves are significantly modular (best described by a two-parameter model),

while the Pit 13 wolves are more integrated (best described by a one-parameter model).

Current whole shape integration metrics fail to capture this change in integration;

therefore we develop a modified version of the SVG based on the effective rank. Because

the Pit 61/67 sample size is larger than Pit 13, we can jackknife the Pit 61/67 sample size

down to that of Pit 13. This jackknife yields both a confidence interval for hypothesis

testing, and normalizes the rank degeneracy due to matrix information content, allowing

us to demonstrate that the new metric is successful in capturing the increase in integration

in Pit 13 wolves. We term this metric the 'effective dispersion'.

Methods and Results, Part 3—In this section, we generalize the concept of

effective dispersion to the case of arbitrary matrix information content. The general

metric developed here, termed the 'relative dispersion', allows comparison of whole

shape integration among data sets of different sample sizes. It also allows for comparison

among data sets of arbitrary landmark number, because it is normalized to the

information content of the input variables. This normalization relies on the computation

of the effective rank of a permuted data matrix. We demonstrate the utility of the relative

dispersion metric by analyzing the *Smilodon* data set, finding that the *Smilodon* jaw is

much more tightly integrated than that of *Canis dirus*.

The last analytical section is followed by a standard Discussion. One topic

considered is the extension of the relative dispersion into a dense semilandmark context,

allowing quantification of the information content of a comnplete shape. We propose the

term 'latent dispersion' for this quantity.

MATERIALS

Background

The primary data in this paper come from two populations of dire wolves (*Canis dirus*,

Figure 1) from the Rancho La Brea tar pit lagerstätten, deposited in what is now Los

Angeles during the terminal Pleistocene (Stock and Harris, 1992). The populations come

from two pit deposits, Pits 13 and 61/67. The Pit 13 wolves show elevated tooth breakage

and wear, while those from 61/67 do not (Binder et al., 2002). Elevated tooth breakage

and wear are thought to indicate nutrient stress, as animals both process carcasses more

completely, and compete more intensely, when prey resources are scarce (Van

Valkenburgh, 1988, 2009; Meloro, 2012). Subsequent work established this pattern

beyond doubt and was able to assign approximate ages to the relevant pit deposits

(Binder and Van Valkenburgh, 2010; O'Keefe et al., 2014; Brannick et al., 2015).

Phenotypic differences between pits have also been documented; morphometric analyses

of dire wolf jaws (Brannick et al., 2015) and crania (O'Keefe et al., 2014) detected clear

phenotypic differences in Pit 13 wolves relative to those in 61/67. Pit 13 wolves are

significantly smaller in overall body size, and also show significant shape differences,

including diminution of the viscerocranium relative to the neurocranium. The observed

size and shape changes were attributed to a chronic lack of nutrition during development,

producing a neotenic adult phenotype in Pit 13. The biological drivers behind this

evolutionary change are not relevant here; what is relevant is that there is a significant

modularity shift between the two pits. This creates a useful model system comprising two

populations of the same species deposited in the same place, separated only in time, and

possessing a significant change in modularity.

Data Collection

Fossils utilized in this paper are left dentaries of *Canis dirus*, from Pits 13 and

61/67, housed in the Tar Pit Museum at Hancock Park, Los Angeles. The input data for

all analyses were 14 landmarks taken from 119 dire wolf jaws, a subset of the 16

landmark data set from Brannick et al. (2015). One hundred and nineteen adult *Canis*

dirus dentaries were included in this study. The labial side of specimens from Pits 13 (n

=36), and 61/67 (n = 83) were photographed using a tripod-mounted Cannon EOS 30D

8.20-megapixel camera. All specimens were anatomical lefts and were laid flat and

photographed with a 5 cm scale-bar. While camera angle and distance were held constant

when photographing, scale-bars were used to properly size specimen images in Adobe

Photoshop CS2 v.9.0 before landmark digitization; scaling was mostly unnecessary, and

always minor.

Fourteen homologous landmarks were digitized on each specimen using the

program tpsDig2 (Rohlf, 2013). Positions of landmarks were chosen to give a general

outline of the mandible and capture information of functional relevance (Brannick et al.,

2015; Figure 1). Landmarks on the tooth row were placed on the alveolus so specimens

with missing teeth could be included. However, presence of the lower carnassial tooth

was required in order to obtain landmark 5. Two landmarks from Brannick et al. were not

used; one of these, at the angle of the mandible, was difficult to identify and showed

unacceptably large variance relative to the other landmarks. The second was an interior

landmark in the masseteric fossa; it was deleted to make the dataset amenable to outline-

based iterative semilandmarking.

Lastly, as a demonstration of the utility of the relative dispersion metric in Part 3,

we analyze a data set of *Smilodon fatalis* jaws (Meachen et al., 2014), again from Rancho

La Brea. The data set comprised 14 landmarks digitized from photographs of 81

Smilodon jaws originating from Pits 61/67 and 13 at the Tar Pit Museum, and the UCMP

collection. Pooling of the *Smilodon* sample was necessary to increase sample size.

Figure 1. Landmarks used in this study, numbered on a schematic dire wolf jaw

(A), and (B) a representative *Canis dirus* dentary from Pit 61/67, Rancho La Brea,

California.

MATERIALS AND RESULTS

Part 1: Rank Deficiency in Pit 61/67

In this section, we examine the amount of rank deficiency of covariance matrices derived

from the Pit 61/67 *Canis dirus* data set, and quantify where it is coming from. Standard

Procrustes superimposition of the landmark data was first performed using the **geomorph**

package (Adams et al., 2020) in R (R Core Team, 2014). Procrustes superimposition

yielded a matrix of transformed (GPA) landmarks **X** and a vector of centroid size.

Principal components analysis of **X** was performed on four covariance matrices. These

were the correlation and covariance matrices for the original data, and correlation and

covariance matrices for permuted data. The scree plots for the eigenvalue vectors of the

four matrices are shown in Figure 2. Also shown in Figure 2 are the effective ranks of the

four matrices, calculated as defined in Equations 3 and 4. Code for computing the

effective rank of covariance and correlation matrices, including the permutation code,

was written in R (R Core Team, 2014; see Appendix 1). The effective ranks calculated

from the correlation and covariance matrices derived from the real data are unique values.

Effective rank was also calculated for 10000 matrices in which each column was

permuted; this permuted mean is plotted and the standard deviations are reported in the

caption to Figure 2.

Figure 2. Rank deficiency and its sources in Pit 61/67 dire wolves. Scree plots for

four data treatments are plotted. Note that the last four eigenvalues are exactly zero for

both the correlation and covariance matrices computed from the real data, reflecting the

loss of four degrees of freedom to the GPA. The effective rank of permuted data shows

rank deficiency due to lack of matrix information; the true effective ranks are much

smaller than this, and this additional deficiency is due to phenotypic covariance. Effective

ranks for each matrix are: Permuted covariance = 21.028 +/- 0.226; permuted correlation

= 23.728 +/- 0.258; covariance = 11.362; correlation = 12.424.

The scree plots of both the correlation and covariance matrices of X demonstrate

that the vector of eigenvalues for each matrix contains 24 non-zero eigenvalues, the

expected number given the loss of four ranks to the GPA. Eigenvalues 25-28 are non-

zero in the permuted matrices, showing that this rank deficiency is recovered when the

matrix is randomized. Effective rank calculations show that both matrices are also highly

rank deficient. The rank of the covariance matrix is 11.362, meaning that less than half of

the 24 non-zero eigenvalues carry meaningful information. The number of ranks in the

permuted covariance matrix is about 21, indicating that the covariance matrix is

information poor by about seven ranks. Four of these are from the GPA; the rest are

probably due to insufficient sample size (Grabowski and Porto, 2017). The rank

deficiency due to phenotypic covariance is therefore 21 minus the effective rank of the

real matrix (11.36), or about 9.64 ranks. Therefore 12.64 ranks, or over half, of the

eigenvalue distribution is redundant. The small values of the higher eigenvalues are

caused mainly by phenotypic covariance, but they do not carry information about it, and

lack of matrix information acerbates this deficiency. The eigenvalue vector is

overdetermined with respect to its information, and its distribution will be affected as a

result.

427   Figure 2 also shows that the rank of the correlation matrix is 12.424. Use of the

correlation matrix therefore adds over one full rank of information relative to the

covariance matrix. We believe this rank increase is noise, as described in the

Introduction. The pattern for the permuted matrices is similar; the correlation matrix adds

almost three ranks of noise over the value for the covariance matrix.

Part 2: Modularity Models and Whole Shape Integration Metrics

In this section, we introduce the Pit 13 dire wolves and show they are more integrated

than those from Pit 61/67. We do this by testing both populations for modularity against a

range of candidate two- and three parameter models. We then calculate the eigenvalue

distribution and SVG for the two covariance matrices, and show that both fail to capture

the modularity change. We then modify the SVG to use only the non-redundant

eigenvalues, and show that this measure does capture the modularity change.

Modularity Models—A challenge in all modularity studies is initial specification

of the models to be tested. In this paper we use a heuristic approach: examining the

multivariate allometry vectors of interlandmark distances to identify candidate models.

While important, the specification of models is tangential to our discussion of rank

deficiency, and we include it below as Appendix 2. This allows us to move directly to

testing the candidate models for the two data sets. A total of 14 models of modularity

were suggested by the multivariate allometry analyses of the interlandmark distances, and

these models were tested individually using the CR statistic (Goswami and Polly, 2010;

Figure 3; Table 1), performed in the **geomorph** package in R (Adams et al., 2020). The

models were tested both on the pooled data, and on data divided by pit. Effect-size tests

and tests against a model of zero modules were performed on 61/67 wolves only, as only

these wolves displayed significant modularity; representative tests and results are listed in

Table 1.

Model, vs. all other LMs	CR_{pooled}	P value	CR_{13}	P value	$CR_{61/67}$	P value	P_{null}
1,2	1.17	0.427	1.234	0.581	1.063	0.219	-
1,2,9,10	1.10	0.573	1.201	0.838	1.090	0.491	-
1-7	1.15	0.837	1.147	0.786	1.171	0.921	-
3-7	0.96	0.052~	1.067	0.387	0.898	0.022*	0.031*
4-7	0.92	0.046*	0.994	0.108	0.894	0.028*	0.046*
1,2,9,10; 3,4,5,6,7	1.06	0.244	1.192	0.719	1.029	0.226	-

Table 1. Results of representative modularity model tests. Representative module

models are listed on the left column, with CR statistics for three data partitions in

subsequent columns. Pit 61/67 wolves have two significant modules comprising the

check teeth vs. the rest of the jaw, while Pit 13 wolves do not show significant modularity

between these or any other modules. As shown in Figure 3, Pit 13 wolves are best fit with

a one-parameter model, while Pit 61/67 wolves are best fit with a two-parameter model.

Wolves from 61/67 are therefore significantly more modular than those from Pit 13.

Other modularity models tested that were not significant include the two-parameter

models (4,5,6,7,13,14), (4,5,6,7,8), (4,5,6,7,8,13,14), (1,2,8,9,10), and (1,2,14,9,10,11),

and the three-parameter models ((1,2,8,9,10)(4567)), ((1,2,9,10)(4,5,6,7,13,14)), and

((129,10)(4,5,6,7,8)).

There are two models with statistically significant CR values (Table 1). Both are

two parameter models that contrast the length of the post-canine tooth row with the rest

of the jaw. The effect sizes of the modularity models with significant CRs were compared

using the **compare.CR** function in **geomorph**; the effect sizes were not significantly

different. Comparison with a null model of zero modules was performed using the same

function. For Pit 61/67 wolves only, the cheek tooth model (3-7) was significantly better

than the null model, while the model containing only the molars (4-7) was marginally so.

The anterior-posterior width of the canine and of the condyloid process (1,2,9,10) do not

form a distinct module on their own; the significant CR of the three-module model is

driven by the presence of the cheek tooth module. The cheek teeth are clearly a module.

The canine does not take part in this module, and therefore has significant variance

attributable to another factor. The inability of the modularity tests to identify the canine

and condyloid process as a module was surprising. We believe it was not identified

because most of its variation covaries with size. The cheek teeth vary against size, and

this allows identification of that module. We note that Pit 13 wolves had no significant

modules, and must therefore be more integrated than Pit 61/67 wolves.

Figure 3. Candidate modularity models for the *Canis dirus* jaw tested in this paper. The

preferred modularity model for Pit 61/67 wolves is D, a two-parameter model of

(3,4,5,6,7) vs. all other landmarks. This is the model with the best statistical support, but

only in Pit 61/67 (Table 1). The preferred model for Pit 13 wolves is A, the one-

parameter model. The other models shown did not have significant modularity according

to the CR statistic; all models tested are listed in the caption to Table 1.

The magnitude of the CR differences between pits was surprising, given they are

taken from populations of the same species at the same location, sampled 5000 years

apart. Given that the sample size of Pit 13 was 36 while that of 61/67 was 83, we were

concerned that a lack of statistical power was obscuring the modularity signal in Pit 13.

To test for this, we jackknifed the 61/67 sample to 36 members to mirror the sample size

of Pit 13 before re-running the CR analysis. Although confidence intervals widened in

these subsets, the covariance structure of Pit 61/67 was still clearly evident and

significantly modular, indicating that the modularity difference between pits is not

attributable to difference in sample size. The pooled CR statistics, run on all 119 wolves,

indicate that the inclusion of the Pit 13 wolves actually degrades the global modularity

signal (Table 1). This indicates that the relative lack of modularity in Pit 13 is real.

Measures of Whole-Jaw Integration— the eigenvalue dispersion and SVG metrics

were calculated using eigenanalysis, bootstrapping, and permutation codes written in R

(R Core Team, 2014; Appendix 1). The eigenvalue standard deviation metric is

significantly less in Pit 13 wolves (Table 2: $\lambda\sigma$, $13 < 61/67$, $p = 0.00008$), while the

SGV_{24} metric is equivalent between the two groups ($13 = 61/67$, $p = 0.837$, Student's T in

this and subsequent comparisons). This implies that eigenvalue dispersion is significantly

less in Pit 13, and those wolves should be more modular than Pit 61/67. Yet this is not the

case; the modularity tests reported above clearly show that Pit 61/67 wolves are

significantly modular, while Pit 13 wolves are not. The eigenvalue dispersion is less in

Pit 13 even though the jaws are more integrated. The SGV_{24} metric fails to detect the

increase in modularity. As demonstrated above, over half of the ranks used to calculate

both metrics are redundant, and we believe the SGV$_{24}$ is essentially measuring noise. We

experimented with the use of the covariance matrix in the SGV$_{24}$ to see if the failure of

that metric was due to use of the correlation matrix. Use of the covariance matrix

required standardization of the eigenvalues before calculation to control for different

matrix variances. This modified metric is also reported in Table 2, and like the SGV$_{24}$ it

is equivalent between the two matrices. Rank inflation due to use of the correlation

matrix is therefore not the reason for failure of the SGV$_{24}$.

521   *Effective rank-scaled SGV or 'effective dispersion'.*— Calculation of the effective

rank for the two populations indicates that the rank of Pit 61/67 is less than that of Pit 13

(Figure 4: R_e, 13 > 61/67, $p = 0.00148$). To account for this difference in dimensionality

dispersion, we modify the SGV to consider only the non-redundant eigenvalues of the

covariance matrix. This metric includes only the eigenvalues that carry significant

information, i.e. those up to the effective rank of the matrix. We label this quantity

SGV$_{Re}$, and calculate it as follows:

$$SGV_{Re} = \sqrt[Re]{\prod \lambda_{Re}} = \sqrt[Re]{\left(\prod_{i=1}^{[Re]} \lambda_i\right) * \left(R_e - [R_e]\right)(\lambda_{[Re]+1})} \qquad Eq.\ 5$$

Where R_e is the effective rank of the covariance matrix, and $[R_e]$ is the integer value of

the effective rank. The SGV$_{Re}$ is the R_eth root of the product of the eigenvalues up to the

integer value of R_e, multiplied by the decimal remainder of R_e times the next smallest

eigenvalue. The value for this statistic was calculated for the different data partitions

using code written in R, with jackknifed confidence intervals for Pit 61/67 (Table 2;

Appendix 1). Using this measure, Pit 13 wolves have significantly higher dispersion than

those of Pit 61/67. Hence they are more integrated, while 61/67 wolves are more

modular. The SGV_{Re} metric successfully recovers the increase in modularity exhibited by

61/67 wolves.

Figure 4. Scree plots and effective rank for GPA landmark data of *Canis dirus*

jaws. Blue is Pit 13, black is Pit 61/67. Confidence intervals for effective rank (R_e) are

10000 jackknife replicates of the Pit 61/67 sample to an $n = 36$. Eigenvalue dispersion

appears greater in Pit 61/67 wolves, even though they have significant modularity and Pit

13 wolves do not. However the amount of total variance is greater in Pit 13, and the Pit

13 effective rank is significantly larger, so the variance is spread over a greater number of

dimensions. The permuted dimensionality is also shown on both plots; this value is

identical between pits ($\pi R_e = 16.84$), demonstrating that the rank deficiency difference

between the pits is due to phenotypic covariance only (i.e. the rank deficiency due to lack

of matrix information is equivalent). The effective rank of both matrices is very deficient,

with fewer than half of the 24 eigenvalues being meaningful. The effective rank of Pit 13

wolves is 10.414; the permuted effective rank is 16.841 +/- 0.345. The jackknifed

effective rank for Pit 61/67 wolves is 9.89 +/- 0.656, while the permuted value at $n = 36$

is 16.832 +/- 0.519.

Previous authors rejected SGV as a statistic to measure dispersion because its

distribution is not linear (Pavlicev et al., 2009); they prefered the standard deviation of

the eigenvalues because of its linearity and because it has the same units as the input data.

Because the SGV_{Re} is a measure of variance, one may utilize the definition of the

standard deviation to transform the SGV_{Re} into a similar measure:

$$D_e = \sqrt{SGV_{Re}} = \sqrt[2Re]{\left(\prod_{i=1}^{[Re]} \lambda_i\right) * (R_e - [R_e])(\lambda_{[Re]+1})} \qquad Eq.\ 6$$

562   where the square root of the SGV_{Re} is a new metric, D_e, that we call 'effective dispersion'.

563   It measures dispersion in both variance and dimensionality together, and accounts for the

564   rank deficiency that misleads the classical integration metrics. The D_e metric is

565   significantly different between the two samples ($p = 0.0049$; Table 2).

Data Partition	$\lambda\sigma$, +/-	SGV_{24}, +/-	$covSGV_{24}$, +/-	$SGV_{Re,}$ +/-	D_e +/-
C. dirus 13	0.277	0.363	0.0108	4.131e-5	0.0064

$n = 36$					
C. dirus 6167j	0.304	0.362	0.0108	3.496 e-5	0.0058
$n = 36$	0.027	0.029	0.0008	1.42 e-5	0.0012

Table 2. Whole jaw integration measures and other statistics for the *Canis dirus*

mandible data sets. All reported confidence intervals are one standard deviation of 10000

jackknife or permutation replicates. The D_e metric is significantly different between the

two samples ($p = 0.0049$). Quantities listed are eigenvalue standard deviation (correlation

matrix, 24 eigenvalues), $(\lambda\sigma)$; standardized generalized variance 24 (SGV_{24});

standardized generalized variance 24 using the covariance matrix, and standardizing the

eigenvalue vector to the sum of eigenvalues ($covSGV_{24}$); standardized generalized

variance effective rank (SGV_{Re}); and effective dispersion (D_e).

Part 3: Matrix Information and Relative Dispersion

Permuted Rank Standardization— Sample size impacts effective rank; the

effective rank of 61/67 wolves at $n = 36$ is 9.89, while the full matrix of $n = 83$ has an

effective rank of 11.362 (Figures 2 and 4). Clearly the derivation of a version of D_e that

accounts for matrix size is desirable, as this will allow comparisons between matrices of

different sizes. Pavlicev et al. (2009) accomplish a similar standardization for eigenvalue

standard deviation by dividing the observed eigenvalue standard deviation by its

maximum possible value to yield their "relative standard deviation". Because they use the

correlation coefficient in their calculations, the minimum possible correlation in a matrix

is simply the number of traits minus one, because each trait adds an additional unit of

uncorrelated variance. A relative version of effective dispersion is more difficult to

calculate because the minimal covariance in a matrix is not derivable from first

principles, because the input variance for each variable is not one. Also, because the

matrix is rank deficient this must also be considered, suggesting that an approach based

on effective rank is necessary. The quantity needed for standardization must therefore

preserve the variances of the input variables, but remove their statistical covariance, and

must also be treated for rank deficiency. One can generate a quantity with these

properties by permuting the columns in **X** and then calculating the resulting covariance

matrix. This matrix will preserve the input variances on the diagonal, but will remove

statistical covariance on the off-diagonal (although covariances will still be non-zero in a

rank-deficient matrix). The effective rank of the permuted matrix can then be calculated,

yielding πR_e. This permuted effective rank is the maximum matrix rank given the

variable input variances and no statistical correlation among them. It is the amount of

matrix information, and is driven by sample size in reasonably complex shapes. It can be

used as a basis for standardizing D_e for matrix size.

The appropriate scaling is based on consideration of what the effective dispersion

actually is, namely the square root of the geometric mean of a subset of the eigenvalues

of **K**, or alternatively, the root of the variance of one dimension of the hyperellipse.

Because D_e it is the square root of a mean eigenvalue (a variance), it has the same units as

the original data (Pavlicev et al, 2009), and we would like to conserve this property in the

scaled coefficient. Therefore there should be no variance term in the standardization on

dimensional grounds. More importantly, the other property of D_e is that it has been

limited to its non-redundant information content, and we wish to scale D_e to an

expectation of maximum possible information content per axis. We are not concerned

with scaling for the variance in the matrix, but for the amount of information in one

dimension of an uncorrelated matrix that conserves this variance. The appropriate scaling

term is therefore one rank of the permuted matrix, or $1/\pi R_e$. This yields the definition for

the "relative dispersion", or D_r:

$$D_r = \frac{D_e}{\sqrt{\dfrac{1}{\pi R_e}}} = \frac{\sqrt[Re]{\prod \lambda_{Re}}}{\sqrt{\dfrac{1}{\pi R_e}}} \qquad Eq.\ 7$$

Relative Dispersion of Canis dirus *and* Smilodon— To demonstrate the use of

relative dispersion, we turn to the third data set used in this paper, that of RLB *Smilodon*

fatalis (Meachen et al., 2014). These data comprise 14 two-dimensional landmarks on 81

jaws, hence are quite similar to the dire wolf data (11 of the 14 landmarks are

homologous between *Smilodon* and *Canis dirus*). We began by calculating the effective

rank for *Smilodon* and comparing it to Pit 61/67 dire wolves (Figure 6). Both samples

were bootstrapped for confidence intervals, and the sample size for 61/67 was held at 81

during the bootstrap. Both procedures lower effective sample size, and so reduce matrix

information and hence rank. The exact values for integration statistics are reported in

Table 4, along with the bootstrap means and standard deviations. Figure 6 plots the $n =$

81 bootstrap values for effective rank.

Figure 6. Scree plot comparison for *Canis dirus* and *Smilodon* data sets, 14

landmarks on jaws, at $n = 81$. Plotting conventions are the same as those used in Figure 2.

Canis dirus 61/67 is shown in black, *Smilodon* in red. Both data sets have been

bootstrapped for a confidence interval (10000 replicates), hence the values shown here

are less than the precise values of R_e for *C. dirus* (11.36) and *Smilodon* (12.32; Table 4).

Because bootstrapping lowers the effective sample size, this drop in rank illustrates the

strong sensitivity of the effective rank calculation to sample size. The plotted means and

standard deviations are: *Canis dirus* 61/67, $R_e = 10.33$ +/- 0.677; $\pi R_e = 20.614$ +/-0.503.

Smilodon $R_e = 11.184$ +/-0.438; $\pi R_e = 21.91$ +/- 0.503.

       Smilodon and *Canis dirus* jaws are similar in that covariance matrices for both are

highly overdetermined, although the effective rank of *Smilodon* is significantly greater.

The permuted ranks of both taxa are much closer to 24 at a sample size of 81 compared to

Canis dirus Pit 13, at a sample size of 36. This implies that the permuted rank of the

matrices should converge toward 24 at large n (probably over 100, in accord with sample

size requirements of other integration metrics as shown by Grabowski and Porto, 2017).

The relative dispersion is comparable among data sets of different sample sizes, as

demonstrated in Table 4 (the D_e of the 61/67 $n = 36$ jackknife sample is 0.0058, while

that of the full $n = 83$ data set is 0.0052; the D_r of both partitions is 0.0238). The values in

Table 4 demonstrate that relative dispersion functions as intended; it is robust to

difference is sample size, and successfully captures the modularity evolution between

dire wolf populations. The comparison for D_r is marginally significant, while that for

sample-size controlled D_e was strongly significant (Table 2); the reasons for this loss of

power were not investigated. Lastly, the D_r metric shows that the *Smilodon* jaw is much

more tightly integrated than that of *Canis dirus*. We note that the SGV_{24} successfully

recovers this difference, while the eigenvalue standard deviation is still incorrect.

Data Partition	$\lambda\sigma_{24}$, +/-	SGV_{24}	SGV_{Re}	D_e	D_r
C. dirus 13 $n = 36$	0.277	0.363	4.131e-5	0.0064	0.0263
BS estimate	0.316	--	5.029e-5	0.00690	0.0283
BS sd	0.015	--	2.283e-5	0.00162	0.0066
C. dirus 6167 $n = 83$	0.278	0.500	2.71e-5	0.0052	0.0238
BS estimate	0.296	0.421	3.373e-5	0.00567	0.0260
BS sd	0.025	0.030	1.311e-5	0.00112	0.0051
Smilodon n = 81	0.250	0.575	7.20e-5	0.0085	0.0391
BS estimate	0.270	0.480	9.098e-5	0.00940	0.0432
BS sd	0.010	0.024	3.055e-5	0.00161	0.0074

Table 3. Whole jaw integration measures for the *Canis dirus* jaw samples, and for

the *Smilodon* data. The relative dispersion difference between Pit 13 and Pit 61/67 wolves

is marginally significant (Welch's $T = 1.863$, $p = 0.068$); Those between *Smilodon* and

the two wolf data sets are highly significant (13, $T = -10.85$, $p < 0.0001$; 6167, $T = -$

17.29, $p < 0.0001$). The *Smilodon* jaw is much more tightly integrated than that of *Canis*

dirus. Note that the SGV_{24} also correctly captures the integration difference between

Smilodon and *Canis dirus*, while the eigenvalue standard deviation still fails. Quantities

listed are eigenvalue standard deviation (correlation matrix, 24 eigenvalues), $(\lambda \sigma)$;

standardized generalized variance 24 (SGV_{24}); standardized generalized variance

effective rank (SGV_{Re}); effective dispersion (D_e); and relative dispersion (D_r).

Discussion

Rank Deficiency and Its Treatment

In this paper we demonstrate that a typical GPA phenotypic covariance matrix is highly

rank deficient. This fact is obvious from the scree plot, but quantifying the rank

deficiency requires use of the Shannon, or information, entropy. Information entropy is

used in signal processing to determine the amount a signal can be compressed without

loss of information. We operationalize the Shannon entropy to a geometric morphometric

context and use it to calculate that the information, or effective, rank of the dire wolf

covariance matrix is less than half of the full rank. Geometric morphometric data sets

have several characteristics that make rank deficiency more severe than in a matrix of

linear measures. The first is the GPA procedure, which utilizes four degrees of freedom

and hence removes four ranks from the covariance matrices dealt with here. This

magnitude of this rank loss is known (Figure 1) and can be accommodated. By contrast,

The magnitude of rank deficiency due to lack of matrix information is not known. We use

a permutation approach to estimate this magnitude, allowing us to calculate the

magnitudes of rank deficiency due to phenotypic covariance vs. the other sources.

Phenotypic covariance accounts for about 9.6 ranks of deficiency in the dire wolf data set

considered in Part 1, while lack of matrix information accounts for about 3 ranks, and the

GPA accounts for 4 ranks. This non-phenotypic rank deficiency is worrisome, because

misleads whole shape integration metrics that rely on the entire eigenvalue distribution.

To demonstrate the impact of rank deficiency on current integration metrics, we

show that current whole shape integration metrics fail to identify the modularity change

between dire wolf populations. Granted this is a minor change, from a one parameter

model to a two parameter model, and the differences in whole-shape metric magnitudes

are not large. However eigenvalue standard deviation is statistically significant in the

wrong direction, while the SGV_{24} seems to lack sensitivity. The SGV_{24} does successfully

capture the modularity difference between *Smilodon* and *Canis dirus*, where the

integration difference is large. Yet eigenvalue standard deviation continues to fail. We

also show that use of the correlation matrix leads to rank inflation in coordinate data; this

rank inflation is noise, and the correlation matrix should not be used on raw coordinate

data.

The failure of current integration metrics arises from the inclusion of a long tail of

non-zero eigenvalues in the eigenvalue distribution. This tail of small values will have

low dispersion, and eigenvalue standard deviation will be pushed lower as rank

deficiency increases. The geometric mean of the eigenvalue distribution (the SGV_{24}) will

also be pushed lower, again due to the inclusion of many small eigenvalues. These facts

are true on inspection but should be demonstrated; one immediate line of future research

is a simulation study that quantifies the sensitivity of current integration metrics to

information rank deficiency. This paper is meant as a theoretical development and proof

of concept, and we do not attempt a simulation study here, although the behavior of the

effective and relative dispersion metrics should also be explored by simulation. The

seeming loss of statistical power between D_e and D_r deserves specific attention.

The use of the effective rank allows calculation of a modified form of the

geometric mean of the eigenvalues, which is simply the mean of the non-redundant

eigenvalues. We demonstrate that this new metric, D_e, successfully recovers the evolution

of increased modularity in Pit 61/67 wolves. However, matrix rank is very sensitive to

sample size (Table 3), and we held sample size constant in Part 2. Therefore we introduce

the concept of relative dispersion, D_r, to account for differences in matrix information

content in Part 3. Relative dispersion accounts for dispersion in dimensionality, as well as

dispersion in variance, and it accounts for differences in matrix size by standardizing

against the permuted information content. The metric D_r should therefore be comparable

among data sets, as we demonstrate via comparison with *Smilodon*. This comparison

utilizes data sets with an equal number of landmarks (14); the sensitivity of relative

dispersion to different landmark number requires quantification in a manner similar to the

analysis of classical integration metrics by Grabowski and Porto (2017).

Dense Semilandmarks and Latent Dispersion

The relative dispersion metric accounts for information rank deficiency, but it is still

space-specific. By this we mean that it is calculated in a shape space defined by the

landmarks, not the entire shape, and the relation between these two spaces is not clear.

For maximum utility, it is desirable that D_r should be comparable among spaces defined

by different, arbitrary sets of landmarks. In the case of two spaces defined by the same

number of landmarks, as in the *Canis dirus* and *Smilodon* data compared here, one might

suppose that relative dispersion is comparable even if the landmarks are not homologous

(11 of the 14 landmarks are in fact homologous). Yet this is not true. A landmark-defined

space with covariance matrix $\mathbf{K_V}$ and an effective rank based on the Shannon entropy

only captures the variance of the landmarks, not of the entire shape upon which those

landmarks occur. One way to represent the total covariance of the entire shape, $\mathbf{K_T}$,

would be to sample the shape with an arbitrarily large number of landmarks, so that as

the magnitude of $\mathbf{V}$ increases $\mathbf{K_V}$ would approach $\mathbf{K_T}$. Thought of in this way, a typical

set of geometric morphometric data would comprise

$$K_T = K_V + K_R$$

Where $\mathbf{K_R}$ is the residual covariance in the shape not captured by the landmarks in

$\mathbf{K_V}$. The residual matrix $\mathbf{K_R}$ is of unknown magnitude in all mainstream applications of

geometric morphometrics, and there is no guarantee that is negligible, nor that the

proportion $\mathbf{K_R/K_T}$ is of constant magnitude among spaces, even if $\mathbf{K_V}$s are of equivalent

effective rank. Therefore the relative dispersion metric defined above remains space-

specific.

A recent approach to estimating residual shape variance is the employment of

dense semilandmarks, where a curve or surface is first landmarked, and then densely

sampled with an increasing number of semilandmarks until the coefficient of interest

stabilizes (Marshall et al. 2019). This procedure can be employed here. Given a shape

with variance $\mathbf{K_T}$ and landmark variance $\mathbf{K_V}$, the residual covariance $\mathbf{K_R}$ will decrease as

$\mathbf{V}$ increases, so that

$$K_T \approx K_V, and \ K_R \approx 0$$

as the number of semilandmarks becomes large. The number of semilandmarks can be

arbitrarily large, down to the pixel or voxel resolution of the image being digitized, but in

practice semilandmarks need only be added until the coefficient stabilizes. Because the

relative dispersion metric relies on information entropy, it will scale with the information

added by each landmark, not by landmark number. Using this procedure allows

characterization of the true, or latent, dispersion of the shape;

$$D_l = \lim_{v \to \infty} D_r \qquad\qquad Eq.\ 8$$

where the latent dispersion of the shape space, D_l, is the relative disper sion of a

matrix that is semilandmarked densely enough to asymptotically stabilize D_r. Classical

integration measures are not amenable to this procedure, because they consider all

eigenvalues, and as the number of landmarks increases the rank of $\mathbf{K}$ will become

increasingly rank deficient.

The latent dispersion metric should be useful for comparing shapes across

arbitrary landmark spaces, and for assessing the fidelity with which the original landmark

data capture phenotypic shape change. It is a general statement of the information content

of a shape. Calculation of this metric would allow us to state definitively that the

Smilodon data is tighter in shape dispersion than *Canis dirus*, assuming the observed

pattern from 14 landmarks holds up. Yet given the rank deficiency in **K** demonstrated by

the permuted matrices, we doubt there is sufficient residual variation not being captured

by the landmarks to substantially change this result. *Smilodon* is a highly specialized

hypercarnivore, and intuitively it should be more tightly adapted—and less complex—

than a more generalized canid like *Canis dirus*, whose jaw should be more information

rich.

Conclusions

-- Measures of whole-shape integration are reviewed; VanValen's realization that

dimensionality dispersion is a critical property is highlighted, and the Shannon, or

information, entropy is employed to measure it as effective rank.

-- Phenotypic covariance matrices for dire wolf jaws from two populations are

introduced: Pit 13, deposited circa 19 kya, and Pit 61/67, deposited circa 14 kya. Both

data matrices comprise X, Y coordinate data for 14 identical landmarks, after Procrustes

superimposition.

-- Use of the concept of information entropy allows identification of the magnitude and

sources of rank deficiency in the Pit 61/67 sample. Rank deficiency is found to be high,

with almost half not due to phenotypic covariance.

-- Modularity model tests show that the only significant module is the ontogenetic one of

the cheek teeth relative to the jaw corpus. This module is only significant in Pit 61/67

wolves; its absence in Pit 13 means that Pit 13 wolves are more integrated (best fit by a

one parameter model), while 61/67 wolves are more modular (best fit by a two parameter

model).

-- Classical metrics of whole shape integration fail to recover the evolutionary increase in

modularity in Pit 61/67 wolves. They fail because 1) they rely on the correlation matrix,

which inflates trivial variance in geometric morphometric data; and 2) they measure only

variance dispersion, while ignoring dimensionality dispersion.

-- New metrics based on the Shannon entropy-modified SGV are defined to quantify the

dispersion of phenotypic shapes spaces. The relative dispersion (D_r) is a sample-size

corrected version of the effective dispersion (D_e); the latent dispersion is an asymptotic

extension of D_r to a dense semilandmark context. The relative dispersion is the core result

of this paper; we restate it from Equation 7:

$$D_r = \frac{\sqrt[2Re]{\prod \lambda_{Re}}}{\sqrt{\frac{1}{\pi R_e}}}$$

-- New whole shape integration measures incorporating effective rank are successful at

recovering the evolution of increased modularity in 61/67 wolves, and show promise for

the characterization of phenotypic spaces among taxa.

Supplementary Materials

Data Availability Statement: The data used in this paper are available as supplementary

materials, and are also posted to the Dryad Digital Repository:

http://dx.doi.org/10.5061/dryad.d7wm37q01

F. R. O'Keefe, J. A. Meachen, and P. D. Polly

Acknowledgements

We thank the curators and staff at the Tar Pit Museum, specifically John Harris, Emily

Lindsey, Chris Shaw, Aisling Farrell, and Gary Takeuchi, for allowing us to collect data

from the dire wolf specimens used in this project and for assistance of all kinds. We also

thank Dr. H. David Sheets for his assistance using the IMP programs, including

pcagen_7a. We thank the Graduate College for the Marshall University Summer Thesis

Award and the Marshall University Drinko Research Fellowship to FRO for funding

towards this project. Special thanks are due to Emily Lindsey, John Southon, Wendy

Binder, and Larisa DeSantis for input on early versions of this manuscript, and to Bryan

Carstens, Lauren Esposito, and two anonymous reviewers for thorough and very helpful

input during the review process. Work by FRO was funded in part by NSF EAR-SGP

1757236 (Project SABER), and by a Drinko Distinguished Research Fellowship to FRO.

We grudgingly thank Thom Yorke.

Literature Cited

Adams, D. C. 2016. Evaluating modularity in morphometric data: challenges with the RV

coefficient and a new test measure. Methods in Ecology and Evolution *7*(5): 565-

572.

Adams, D., Collyer, M., and Kaliontzopoulou, A. 2020. Geomorph: Software for

geometric morphometric analyses. R package version 3.2.1. https://cran.r-

project.org/package=geomorph.

Binder, W. J., Thompson, E. N., & Van Valkenburgh, B. 2002. Temporal variation in

tooth fracture among Rancho La Brea dire wolves. Journal of Vertebrate

Paleontology 22(2): 423-428.

Binder, W.J. and Van Valkenburgh, B. 2010. A comparison of tooth wear and breakage

in Rancho La Brea sabertooth cats and dire wolves across time. Journal of

Vertebrate Paleontology 30: 255-261.

Bookstein, F. L. 1997. Morphometric tools for landmark data: geometry and biology.

Cambridge University Press.

Brannick, A.L., Meachen, J.A., and O'Keefe, F.R. 2015. Microevolution of jaw shape in

the dire wolf, Canis dirus, at Rancho La Brea. La Brea and beyond: The

paleontology of asphalt-preserved biotas, ed. J. M. Harris. Natural History

Museum of Los Angeles County, Science Series 42: 23-32.

Bunea, F., She, Y., and Wegkamp, M. H. 2011. Optimal selection of reduced rank

estimators of high-dimensional matrices. The Annals of Statistics *39*(2): 1282-

1309.

Cangelosi, R. and Goriely, A. 2007. Component retention in principal component

analysis with application to cDNA microarray data. Biology Direct *2*(1): 1-21.

Cheverud, J. M. 1982. Phenotypic, genetic, and environmental morphological integration

in the cranium. Evolution 36: 499-516.

Cheverud, J. M. 1996. Developmental integration and the evolution of pleiotropy.

American Zoologist *36*(1): 44-50.

Curth, S., Fischer, M. S., and Kupczik, K. 2017. Patterns of integration in the canine

skull: an inside view into the relationship of the skull modules of domestic dogs

and wolves. Zoology 125: 1-9.

Darwin, C. 1859. The origin of species and The descent of man, New York. The Modern

Library.

Grabowski, M. and Porto, A. 2017. How many more? Sample size determination in

studies of morphological integration and evolvability. Methods in Ecology and

Evolution 8(5): 592-603.

Goswami, A., Smaers, J. B., Soligo, C. and Polly, P. D. 2014. The macroevolutionary

consequences of phenotypic integration: from development to deep time.

Philosophical Transactions of the Royal Society B: Biological Sciences,

369(1649): 20130254.

Goswami, A., Binder, W. J., Meachen, J., & O'Keefe, F. R. 2015. The fossil record of

phenotypic integration and modularity: A deep-time perspective on

developmental and evolutionary dynamics. Proceedings of the National Academy

of Sciences 112(16): 4891-4896.

Goswami, A. and Polly, P.D. 2010. Methods for studying morphological integration and

modularity. The Paleontological Society Papers 16: 213-243.

Hine, E., and Blows, M. W. 2006. Determining the effective dimensionality of the

genetic variance–covariance matrix. Genetics 173(2): 1135-1144.

Horvath, S. 2011. *Weighted network analysis: applications in genomics and systems*

biology. Springer Science & Business Media.

Jolicoeur, P. 1963. Note: the multivariate generalization of the allometry equation.

Biometrics 19(3): 497-499.

Klingenberg, C. P. 2008. Morphological integration and developmental modularity.

Annual Review of Ecology, Evolution and Systematics 39: 115–132.

Klingenberg CP. 2010. Evolution and development of shape: integrating quantitative

approaches. Nature Reviews Genetics 11(9): 623-35.

Klingenberg CP. 2013. Cranial integration and modularity: insights into evolution and

development from morphometric data. Hystrix, the Italian Journal of Mammalogy

24(1): 43-58.

Marshall, A. F., Bardua, C., Gower, D. J., Wilkinson, M., Sherratt, E., & Goswami, A.

2019. High-density three-dimensional morphometric analyses support conserved

static (intraspecific) modularity in caecilian (Amphibia: Gymnophiona)

crania. Biological Journal of the Linnean Society 126(4): 721-742.

Meachen, J.A., O'Keefe, F.R., and Sadleir, R.W. 2014. Evolution in the sabre-tooth cat,

Smilodon fatalis, in response to Pleistocene climate change. Journal of

Evolutionary Biology 27: 714-723.

Meloro, C. 2012. Mandibular shape correlates of tooth fracture in extant Carnivora:

implications to inferring feeding behavior of Pleistocene predators. Biological

Journal of the Linnean Society 106: 70-80.

Najarzadeh, D. 2019. Testing equality of standardized generalized variances of k

multivariate normal populations with arbitrary dimensions. Statistical Methods &

Applications *28*(4): 593-623.

O'Keefe, F. R., Rieppel, O., & Sander, P. M. 1999. Shape disassociation and inferred

heterochrony in a clade of pachypleurosaurs (Reptilia, Sauropterygia).

Paleobiology 25: 504-517.

O'Keefe, F. R., Binder, W. J., Frost, S. R., Sadlier, R. W. and Van Valkenburgh, B. 2014.

Cranial morphometrics of the dire wolf, *Canis dirus*, at Rancho La Brea: temporal

variability and its links to nutrient stress and climate. Palaeontologia Electronica.

O'Keefe F. R., Meachen J, Fet E. V., Brannick A. 2016. Ecological determinants of clinal

morphological variation in the cranium of the North American gray wolf. Journal

of Mammalogy 94(6):1223-36.

Olson, E.C. and Miller, R.L., 1958. *Morphological Integration*. University of Chicago

Press (reprint, 1999).

Pavlicev, M., Cheverud, J. M., & Wagner, G. P. 2009. Measuring morphological

integration using eigenvalue variance. Evolutionary Biology 36(1): 157-170.

R Core Team, 2014. R: A language and environment for statistical computing.

Rohlf, F. 2013. tpsDig2, Stony Brook: State University of New York.

Roy, O. and Vetterli, M. 2007. The effective rank: A measure of effective dimensionality.

In 2007 15th European Signal Processing Conference: 606-610. IEEE.

Segura, V., Cassini, G. H., Prevosti, F. J., & Machado, F. A. 2020. Integration or

Modularity in the Mandible of Canids (Carnivora: Canidae): a Geometric

Morphometric Approach. Journal of Mammalian Evolution

doi.org/10.1007/s10914-020-09502-z.

SenGupta, A., 1987. Tests for standardized generalized variances of multivariate normal

populations of possibly different dimensions. Journal of Multivariate Analysis

23(2): 209-219.

Shannon, C.E. 1948. A mathematical theory of communication. The Bell System

Technical Journal *27*(3): 379-423.

Stock, C. and Harris, J.M. 1992. Rancho La Brea: A Record of Pleistocene Life in

California. Science Series No. 37. Natural History Museum of Los Angeles

County, Los Angeles, CA.

Van Valen, L. 1974. Multivariate structural statistics in natural history. Journal of

Theoretical Biology 45(1): 235-247.

Van Valkenburgh, B. 1988. Incidence of tooth fracture among large, predatory mammals.

American Naturalist 13: 291-302.

Van Valkenburgh, B. 2009. Costs of carnivory: tooth frac- ture is Pleistocene and recent

carnivores. Biological Journal of the Linnean Society 96: 68-81.

Wagner, G. P. 1984. On the eigenvalue distribution of genetic and phenotypic dispersion

matrices: evidence for a nonrandom organization of quantitative character

variation. Journal of Mathematical Biology 21: 77-95.

Wilks, S. S. 1932. Certain generalizations in the analysis of variance. Biometrika: 471-

494.

Zelditch, M.L., Swiderski, D.L. and Sheets, H.D. (2012). Geometric morphometrics for

biologists: a primer. Academic Press.

Appendix 1
946
947   *R Code for Effective Dispersion Metrics, 1. Code for calculating permuted SVG, Effective*
948   *Rank, and Effective Dispersion*
949
950   #attach necessary libraries
951
952   library(dplyr)
953   library(broom)
954   library(readxl)
955
956   #Define a variable containing the GPA X,Y landmark data. The data is read in as a data
957   frame from an Excel file.
958
959   GPALMData <- read_excel("~/Desktop/C dirus 14 LM/CDirus6167LM.xlsx")
960
961   View(GPALMData)
962
963   EffRank=matrix(nrow=1,ncol=0)
964   SGV=matrix(nrow=1,ncol=0)
965   EffRankSGV=matrix(nrow=1,ncol=0)
966   EigenSumRec=matrix(nrow=1,ncol=0)
967
968   #Start permutation loop
969   for (i in 1:1000){
970
971   #Permute rows in each variable column vector independently for permutation test.
972   LMPrune = (GPALMData [3:30])
973   PermLM = apply(LMPrune, 2, sample, 83, replace=FALSE)
974   #sample size
975
976   #Define variable that is the covariance matrix of the above. Be sure to specify only the
977   landmark columns in the data frame.
978   CVLM= cov(PermLM)
979   #toggle for cov/cor
980
981   #Define object that is the eigenanalysis of the covariance matrix.
982   CVLMeigen=eigen(CVLM)
983
984   EigenVal<-CVLMeigen$values
985
986   #Define the summation of eigenvalues from the eigenanalysis.
987   EigenSum=sum(EigenVal)
988
989   #Create vector of scaled eigenvalues
990   PofK<-EigenVal/EigenSum

```
991
992    #Creat vector of lnPofK
993    lnPofK<-log(PofK)
994
995    Product<-lnPofK*PofK
996
997    SumProduct<-sum(Product, na.rm=TRUE)
998
999    ShannonEntropy <- -1*SumProduct
1000
1001   EffectiveRank<-exp(ShannonEntropy)
1002
1003   #Begin SGV24 calculation
1004   EigenProduct<-prod(EigenVal[1:24])
1005
1006   StandGenVar<-(EigenProduct)^(1/24)
1007
1008   SGV<-cbind(SGV, StandGenVar)
1009
1010   #Begin ReSGV calculation
1011   ERInteger<-as.integer(EffectiveRank)
1012
1013   EigenTrim<-EigenVal[1:ERInteger]
1014
1015   FractRankVar<-EigenVal[ERInteger+1]*(EffectiveRank-ERInteger)
1016
1017   ProductEffRankEigen<-prod(EigenTrim)*FractRankVar
1018
1019   EffectiveRankSGV<-(ProductEffRankEigen)^(1/EffectiveRank)
1020
1021   EffRank <-cbind(EffRank, EffectiveRank)
1022
1023   EffRankSGV <-cbind(EffRankSGV, EffectiveRankSGV)
1024
1025   EigenSumRec <-cbind(EigenSumRec, EigenSum)
1026
1027   cat("\014")
1028
1029   print(i)
1030
1031   }
1032
1033   #calculate and print st deviation and mean for each parameter
1034
1035   print(mean(EigenSumRec))
1036   print (sd(EigenSumRec))
```

```
1037
1038    print(mean(EffRank))
1039    print (sd(EffRank))
1040
1041    print (mean(SGV))
1042    print (sd(SGV))
1043
1044    print (mean(EffRankSGV))
1045    print (sd(EffRankSGV))
1046
```

1047   *R Code for Effective Dispersion Metrics, 2. Code for calculating SVG, Effective Rank,*
1048   *and Effective Dispersion*
1049
1050   #attach necessary libraries
1051
1052   library(dplyr)
1053   library(broom)
1054   library(readxl)
1055
1056   #Define a variable containing the X,Y landmark data. The data is read in as a data frame
1057   from an Excel file.
1058
1059   GPALM <- read_excel("~/Desktop/C Dirus 14 LM/CDirus13LM.xlsx")
1060   View(GPALM)
1061
#initialize variables
EffRank=matrix(nrow=1,ncol=0)
SGV=matrix(nrow=1,ncol=0)
EffRankSGV=matrix(nrow=1,ncol=0)
EigenSumRec=matrix(nrow=1,ncol=0)
EffDispersion=matrix(nrow=1,ncol=0)
1068
1069   #Start bootstrap loop
1070   for (i in 1:10000){
1071   bootsample=sample_n(GPALM, 36, replace = FALSE)
1072   #edit for n and T/F
1073
1074   #Define variable that is the covariance matrix of the above. Be sure to specify only the
1075   landmark columns in the data frame.
1076   CVLM= cov(bootsample[3:30])
1077   #toggle cov/cor
1078
1079   #Define object that is the eigenanalysis of the covariance matrix.
1080   CVLMeigen=eigen(CVLM)
1081
1082   EigenVal<-CVLMeigen$values
1083
1084   #Define the summation of eigenvalues from the eigenanalysis.
1085   EigenSum=sum(EigenVal)
1086
1087   #Create vector of scaled eigenvalues
1088   PofK<-EigenVal/EigenSum
1089
1090   #Creat vector of lnPofK and calculate effective rank
1091   lnPofK<-log(PofK)
1092

```
1093    Product<-lnPofK*PofK
1094
1095    SumProduct<-sum(Product, na.rm=TRUE)
1096
1097    ShannonEntropy <- -1*SumProduct
1098
1099    EffectiveRank<-exp(ShannonEntropy)
1100
1101    #Begin SGV24 calculation
1102    EigenProduct<-prod(EigenVal[1:24])
1103
1104    StandGenVar<-(EigenProduct)^(1/24)
1105
1106    SGV<-cbind(SGV, StandGenVar)
1107
1108    #Begin ReSGV calculation
1109
1110    ERInteger<-as.integer(EffectiveRank)
1111
1112    EigenTrim<-EigenVal[1:ERInteger]
1113
1114    FractRankVar<-EigenVal[ERInteger+1]*(EffectiveRank-ERInteger)
1115
1116    ProductEffRankEigen<-prod(EigenTrim)*FractRankVar
1117
1118    EffectiveRankSGV<-(ProductEffRankEigen)^(1/EffectiveRank)
1119
1120    #Make Effective Dispersion
1121
1122    EffDisperse<-sqrt(EffectiveRankSGV)
1123
1124    EffRank <-cbind(EffRank, EffectiveRank)
1125
1126    EffRankSGV <-cbind(EffRankSGV, EffectiveRankSGV)
1127
1128    EigenSumRec <-cbind(EigenSumRec, EigenSum)
1129
1130    EffDispersion <-cbind(EffDispersion, EffDisperse)
1131
1132    cat("\014")
1133
1134    print(i)
1135
1136    }
1137
1138    #calculate and print st deviation and mean for each parameter
```

```
1139
1140     print(mean(EigenSumRec))
1141     print (sd(EigenSumRec))
1142
1143     print(mean(EffRank))
1144     print (sd(EffRank))
1145
1146     print (mean(SGV))
1147     print (sd(SGV))
1148
1149     print (mean(EffRankSGV))
1150     print (sd(EffRankSGV))
1151
1152     print (mean(EffDispersion))
1153     print (sd(EffDispersion))
1154
1155
```

1156   *R Code for Effective Dispersion Metrics, 3. Code for calculating eigenvalue standard*
1157   *deviation*
1158
1159   #attach required packages
1160   library(dplyr)
1161   library(broom)
1162   library(readxl)
1163
1164   #Define a variable containing the X,Y landmark data. The data is read in as a data frame
1165   from an Excel file.
1166
1167   GPALM <- read_excel("~/Desktop/C dirus 14 LM/CDirus6167LM.xlsx")
1168   View(CDirus6167AllomResidLM)
1169
1170   BSSDEigen=matrix(nrow=1,ncol=0)
1171
1172   #Start bootstrap loop
1173   for (i in 1:1000){
1174   bootsample=sample_n(GPALM, 83, replace = FALSE)
1175   #edit n, bootstrap or not
1176
1177   #Define variable that is the covariance matrix of the above. Be sure to specify only the
1178   landmark columns in the data frame.
1179   CVLM= cor(bootsample[3:30])
1180   #toggle cov/cor
1181
1182   #Define object that is the eigenanalysis of the covariance matrix.
1183   CVLMeigen=eigen(CVLM)
1184
1185   EigenVal<-CVLMeigen$values
1186
1187   EigenValTrim<-EigenVal[1:24]
1188
1189   EigenMean<-(sum(EigenValTrim))/24
1190
1191   EigenValTrimMean<-EigenValTrim/EigenMean
1192
1193   EigenK <- (abs(EigenValTrimMean-1))^2
1194
1195   Eigenvar <- (sum(EigenK))/24
1196
1197   SDEigen <- sqrt(Eigenvar)/4.7958
1198
1199   BSSDEigen <-cbind(BSSDEigen, SDEigen)
1200
1201   }

```
1202
1203    #calculate and pring st deviation and mean for each parameter
1204
1205    print(mean(BSSDEigen))
1206    print (sd(BSSDEigen))
1207
1208
```

Appendix 2

Testing the dire wolf data sets for modularity requires specifying candidate models. We

use a heuristic approach based on the multivariate allometry vectors of interlandmark

distances. Initial analysis consisted of calculating the interlandmark distances from the

raw digitized landmark data **LM** (Figure A1). These distances were then evaluated for

multivariate allometry by calculating the first principal component vector of the

covariance matrix of *ln*-transformed distances; this procedure yields the multivariate

allometry vector (Jolicoeur, 1963; O'Keefe et al., 1999; O'Keefe et al., 2013). Confidence

intervals for the allometry vector coefficients where calculated by bootstrapping the input

data (10000 replicates) in the R core package (R Core Team, 2014). The eigenvalues of

the full 13-measure multivariate allometry vector were unusual, necessitating additional

runs with some variables removed; all runs are reported in Table A1.

The multivariate allometry vector for all 13 interlandmark distances is odd, with a

single variable scoring very highly (the distance between the canine and the first

premolar, landmarks 3-4), while the rest show significantly negative coefficients. This

pattern indicates that the c1-p1 distance is highly variable, and that this variance is not

correlated with any other variable. Further analysis shows that the c1-p1 distance crosses

the boundary between two modules, accounting for its unique loading. We therefore

discarded that distance and ran the analysis again with 12 distances. The resulting vectors

for both populations are strongly positively allometric for two variables, the

anteroposterior width of the canine (1-2) and the length of the condyloid process (9-10).

Lengths associated with the other teeth are negatively allometric (Table A1). The

covariation of canine width and condyloid process length suggests a functional unit, and

provides one hypothesis for a testable module in the modularity tests. The lengths of the

other teeth are all negatively allometric, as are the landmarks on the inferior margin on

the jaw that are linked to tooth location. This finding of negative dental allometry in the

cheek teeth has been documented before, both in *Canis lupus* (O'Keefe et al., 2016) and

in *Canis dirus* (O'Keefe et al., 2014). Negative dental allometry is a consequence of the

adult teeth forming early in ontogeny, erupting before the end of the first year, while full

somatic growth is not attained until the end of the second year (O'Keefe et al., 2016).

This negative dental allometry in the cheek teeth suggests a second module, comprising

the lengths of the premolar arcade, the carnassial, and m2. We therefore specify three

candidate modules for testing. They are the condyloid process and canine, the cheek

teeth, and the rest of the jaw. The modularity models tested in Table 1 are permutations

of these modules. We also note a significant allometric difference in the coronoid process

in Pit 13 wolves; this is consistent with sexual diomorphism (O'Keefe et at., 2016), and

we also tested modules including landmark 8 to test for this.

A valid criticism of our heuristic approach is that it does not test the model space

exhaustively. The space of possible modules in a shape of 14 landmarks is $2^{14} - 1$, or

16,383, small enough to allow exhaustive optimization using maximum likelihood. An

alternative approach is weighted network analysis (Horvath, 2011), an effective method

of model induction used with genomic data. We believe that the parts of the model space

tested here capture all plausible candidate models, based on our analysis of the

multivariate allometry vectors, and on subsidiary principal components analysis of the

landmark data.

1254

Distance	Pit 13	Pit 61/67	+/-, *n* = 36j	All	+/-, BS
1-2	0.6543	0.3366	0.105**	0.4889	0.056**
3-4	0.0625	0.1503	0.036**	0.1883	0.017**
4-6	0.1339	0.1721	0.037	0.1500	0.022**
6-7	0.1291	0.1561	0.071	0.1439	0.060**
7-8	0.0706	0.2486	0.034**	0.1927	0.020**
8-9	0.0982	0.2457	0.038**	0.2043	0.018**
9-10	0.4183	0.4479	0.096	0.5682	0.043**
10-11	0.2218	0.3494	0.056**	0.2711	0.033
11-12	0.1140	0.2324	0.048**	0.2105	0.023**
12-13	0.1408	0.1562	0.041	0.1531	0.029**
13-14	0.0593	0.1539	0.041**	0.1842	0.012**
14-1	0.5082	0.4540	0.133	0.2967	0.066
Isometry	0.289	0.289		0.289	

Table 1. Allometry vectors calculated from interlandmark distances on *Canis dirus* dentaries as shown in Figure 1. Three vectors are shown; that for Pit 13, for Pit 61/67, and for the pooled sample. We also jackknifed the Pit 61/67 sample to establish a confidence interval; stars in this column indicate significant differences from Pit 13. The pooled vector was also bootstrapped, and stars in this column indicate differences from isometry in that vector. The isometric coefficient value for each vector is shown at bottom; confidence intervals for coefficient values are one standard deviation of 1000

1262   bootstrap replicates of each vector. Stars are 2 sigma. The heat maps in Figure 1A are

1263   indexed to the pooled coefficients.

1264

1265

Figure A1. Interlandmark distances used in calculating multivariate allometry

vectors. Redder lines indicate more positively allometric distances, blues lines more

negative. Panel A depicts the 12-measure multivariate allometry vector, with each length

heat mapped to its allometry coefficient from Table A1. Panel B shows the same for a 10-

measure allometry vector. These different allometries suggest three different modules:

the condoloid process and the canine, the check teeth, and the rest of the jaw.

1272