Improved molecular characterization of the *Klebsiella oxytoca* complex reveals the prevalence of

the kleboxymycin biosynthetic gene cluster

Preetha Shibu[1]#†, Frazer McCuaig[2]#, Anne L. McCartney[3], Magdalena Kujawska[4], Lindsay J. Hall[4,5],

Lesley Hoyles[2,6]*

[1]Life Sciences, University of Westminster, United Kingdom

[2]Department of Biosciences, Nottingham Trent University, United Kingdom

[3]Department of Food and Nutritional Sciences, University of Reading, United Kingdom

[4]Gut Microbes & Health, Quadram Institute Bioscience, Norwich Research Park, Norwich, United

Kingdom

[5]Chair of Intestinal Microbiome, ZIEL – Institute for Food & Health, Technical University of

Munich, Freising, Germany

[6]Department of Surgery and Cancer, Imperial College London, United Kingdom

#These authors made an equal contribution to the work; shared authorship.

***Corresponding author:** Lesley Hoyles, lesley.hoyles@ntu.ac.uk

†**Present address:** Berkshire and Surrey Pathology Services, Frimley Health NHS Trust, Wexham

Park Hospital, Slough, United Kingdom.

Running title: Distribution of the kleboxymycin BGC in *Klebsiella*

Abbreviations: AAHC, antibiotic-associated haemorrhagic colitis; AMR, antimicrobial resistance;

BGC, biosynthetic gene cluster; MAG, metagenome-assembled genome; MLST, multi-locus

sequencing typing; MSA, multiple-sequence alignment; NEC, necrotizing enterocolitis; PBD,

pyrrolobenzodiazepine; rMLST, ribosomal MLST; TM, tilimycin; TV, tillivaline; VFDB, Virulence

Factors of Pathogenic Bacteria Database.

Keywords: tilimycin, tillivaline, *Klebsiella michiganensis*, antibiotic-associated haemorrhagic colitis,

necrotizing enterocolitis

Data statement: Supplementary data and material associated with this article are available from

figshare.

Data summary: Draft genome sequences for PS_Koxy1, PS_Koxy2 and PS_Koxy4 have been

deposited with links to BioProject accession number PRJNA562720 and under accession numbers

VTQC00000000, VTQB00000000 and VTQA00000000, respectively.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

IMPACT STATEMENT

Members of the *Klebsiella oxytoca* complex are difficult to speciate using phenotypic and

chemotaxonomic methods. Consequently, many genomes deposited in public databases are

misclassified as *K. oxytoca*. Here we demonstrate that the current multi-locus sequence typing

(MLST) system for the complex can be used to accurately speciate many strains, which will be of use

to clinical laboratories in resource-limited settings which rely on the MLST scheme for typing and

epidemiological tracking of isolates. In addition, extended analyses of the genomes of *Klebsiella* spp.

have revealed the kleboxymycin biosynthetic gene cluster (BGC) is restricted to species of the

Klebsiella oxytoca complex (*K. oxytoca*, *K. michiganensis*, *K. pasteurii* and *K. grimontii*). Species-

and/or gene-specific differences in the cluster's sequences may be relevant to virulence of *K. oxytoca*

and related species. The finding of the kleboxymycin BGC in the preterm infant gut microbiota may

have implications for disease presentation in a subset of neonates.

ABSTRACT

As part of ongoing studies with clinically relevant *Klebsiella* spp., we characterized the genomes of three clinical GES-5-positive ST138 strains originally identified as *Klebsiella oxytoca*. bla_{OXY} gene, average nucleotide identity and phylogenetic analyses showed the strains to be *Klebsiella michiganensis*. Affiliation of the strains to ST138 led us to demonstrate that the current multi-locus sequence typing scheme for *K. oxytoca* can be used to speciate members of this genetically diverse complex of bacteria. The strains encoded the kleboxymycin biosynthetic gene cluster (BGC), previously only found in *K. oxytoca* strains and one strain of *Klebsiella grimontii*. The finding of this BGC, associated with antibiotic-associated haemorrhagic colitis, in *K. michiganensis* led us to carry out a wide-ranging study to determine the prevalence of this BGC in *Klebsiella* spp. Of 7,170 publicly available *Klebsiella* genome sequences screened, 88 encoded the kleboxymycin BGC. All BGC-positive strains belonged to the *K. oxytoca* complex, with strains of four (*K. oxytoca*, *K. pasteurii*, *K. grimontii*, *K. michiganensis*) of the six species of the complex found to encode the complete BGC. In addition to being found in *K. grimontii* strains isolated from preterm infants, the BGC was found in *K. oxytoca* and *K. michiganensis* metagenome-assembled genomes recovered from neonates. Detection of the kleboxymycin BGC across the *K. oxytoca* complex may be of clinical relevance and this cluster should be included in databases characterizing virulence factors, in addition to those characterizing BGCs.

INTRODUCTION

Members of the *Klebsiella oxytoca* complex encode a chromosomal β-lactamase gene (bla_{OXY}) (1). Differences in the sequence of this gene allowed the establishment of phylogroups (Ko), which correspond to species: *K. michiganensis* (Ko1, with Ko5 representing a sub-lineage), *K. oxytoca* (Ko2), *K. spallanzanii* (Ko3), *K. pasteurii* (Ko4), *K. grimontii* (Ko6) and *K. huaxiensis* (Ko8). Ko7 has been described on the basis of a single isolate (1). Individual gene (*rpoB*, *gyrA*, *rrs*) sequences can be used to differentiate species of the complex (2), as can genome-based average nucleotide identity (ANI) and phylogenomic analyses (1,3). All members of the *K. oxytoca* complex can be differentiated by MALDI-TOF (1), but reference databases currently in routine clinical use lack reference spectra of the different species to allow identification beyond *K. oxytoca*.

Recent work has demonstrated genomic characterization of *K. oxytoca* strains is inadequate, with large numbers of genomes deposited in public databases erroneously assigned to *K. oxytoca* instead of *K. michiganensis* or *K. grimontii* (3–6). Consequently, *K. michiganensis* and *K. grimontii* are clinically relevant but under-reported in the literature (3,7). Given that the bla_{OXY} gene has diversified in parallel to housekeeping genes in the *K. oxytoca* complex, it is likely that the *K. oxytoca* multi-locus sequence typing (MLST) scheme (8) can be used to speciate all members of this genetically diverse group of bacteria.

Little is known about the antibiotic-resistance and virulence genes encoded by *K. oxytoca* and related species. In the course of ongoing *Klebsiella*–phage work, with three GES-5-positive ST138 strains originally described as *K. oxytoca* (9,10), we sought to determine whether widely recognized virulence factors such as enterobactin, yersiniabactin and salmochelin are encoded in the strains' genomes, and the kleboxymycin biosynthetic gene cluster (BGC), as this was until recently a little-studied BGC implicated in non-*Clostridioides difficile* antibiotic-associated haemorrhagic colitis (AAHC) (11–14). AAHC is caused by the overgrowth of cytotoxin-producing *K. oxytoca* secondary to use of antibiotics such as penicillin or amoxicillin, resulting in the presence of diffuse mucosal oedema and haemorrhagic erosions (15,16). This type of colitis is distinct from the more common form of antibiotic-associated diarrhoea caused by toxin-producing *Clostridiodes difficile*, which usually gives rise to watery diarrhoea resulting in mild to moderate disease.

Gene-based and genomic analyses of our ST138 isolates showed they were *K. michiganensis*,

not *K. oxytoca*, and that along with common virulence genes they encoded the kleboxymycin BGC.

Our findings led us to 1) determine whether the *K. oxytoca* MLST scheme could be used to speciate

members of the *K. oxytoca* complex, and 2) investigate the distribution of the kleboxymycin BGC in a

range of *Klebsiella* and related species.

METHODS

Clinical isolates. Strains PS_Koxy1 (isolated December 2014; cardiothoracic/intensive care unit),

PS_Koxy2 (isolated August 2015; haematology unit) and PS_Koxy4 (isolated September 2015;

haematology unit) had been recovered from a throat swab, urine and rectal swab, respectively,

obtained from three different adults. The strains were from the study of Eades *et al.* (9), described in

further detail by Ellington *et al.* (10) (PS_Koxy1, patient X; PS_Koxy2, patient A; PS_Koxy4, patient

B; Frances Davies, personal communication). The study of anonymized isolates beyond the diagnostic

requirement was approved by an NHS research ethics committee (number 06/Q0406/20). Full details

of Methods associated with the phenotypic and genotypic characterization of the clinical isolates can

be found in **Supplementary Material**.

ANI analysis of genome sequences. All annotated non-redundant *Klebsiella* genome assemblies

available in the NCBI Genome database on 2 September 2019 ($n = 7{,}170$; **Supplementary Table 1**)

were downloaded (17). ANI of genomes with their closest relatives and type strains of species was

assessed using FastANI (18), which uses Mashmap as its MinHash-based alignment-free

sequence mapping engine to provide ANI values for both complete and draft-quality

genomes that are related by 80–100 % ANI.

MLST analyses. Allele sequences ($\underline{n=442}$ representing seven housekeeping genes – *gapA*, *infB*, *mdh*,

pgi, *phoE*, *rpoB*, *tonB* – contributing to 354 different MLST sequence types; correct as of 19 March

2021) for the *K. oxytoca* MLST scheme (8) were used to determine the MLST profiles of all *K.*

oxytoca complex genomes included in this study (**Supplementary Table 1**). The allele sequences

were used to create BLASTN databases against which the assemblies of all genomes included in this

study were searched. Sequences with exact hits to one allele of each housekeeping gene were

retained, allowing us to identify the sequence types of the genomes included in this study

(**Supplementary Table 2**). For those genomes that returned hits to alleles across all seven

housekeeping genes, a phylogenetic tree (neighbour joining, Jukes Cantor) was generated in Geneious

Prime v2019.2.1 using the aligned (CLUSTAL W) concatenated (*gapA–infB–mdh–pgi–phoE–rpoB–*

tonB) nucleotide sequences of their housekeeping genes and those of each sequence type used in the

K. oxytoca MLST scheme (8). Support for clustering of nodes in the tree was determined by bootstrap

analysis (1,000 replications).

Characterization of the kleboxymycin BGC in genomes. The annotated reference sequence of the

kleboxymycin BGC was downloaded from GenBank (accession number MF401554 (11)) and used as

a BLASTP database for searches with the protein sequences encoded within the genomes of

PS_Koxy1, PS_Koxy2 and PS_Koxy4. Initially, Geneious Prime v2019.2.1 was used to identify

regions of the three draft genomes encoding the complete BGC, and to align them to MF401554.

The protein sequences of the annotated assemblies were searched for the kleboxymycin BGC

using the reference sequence and BLASTP v2.9.0+, and the resulting hits were filtered based on >70

% identity and >70 % coverage to identify isolates potentially carrying genes from the BGC. *K.*

grimontii (*n*=3) and *K. michiganensis* (*n*=2) and *K. oxytoca*-related metagenome-assembled genomes

(MAGs) (*n*=25) from Chen *et al.* (3) were also subject to BLASTP searches. Genomes that encoded

the full BGC (i.e. all 12 BGC genes on a contiguous stretch of DNA) were identified from the BLAST

results. The protein sequences encoded in the BGC were extracted from the annotated assemblies

using samtools v1.9 faidx (19) and concatenated into a single sequence (the sequence data are

available as supplementary material from figshare). These concatenated sequences were used to

produce a multiple-sequence alignment (MSA) in Clustal Omega v1.2.4, along with the BGC

sequences of the three *K. michiganensis* clinical isolates, the reference sequence (11), a recently

described *K. grimontii* sequence (20) and a homologous sequence found in *Pectobacterium*

brasiliense BZA12 (to be used as an outgroup in later phylogenetic analyses; identified as encoding

the complete kleboxymycin BGC through NCBI BLASTP). Phylogenetic analyses were carried out

on the MSA using the R package Phangorn v2.5.5 (21), producing a maximum-likelihood tree, which

was visualised and rooted (on *P. brasiliense* BZA12) using the Interactive Tree of Life (iTOL v5.5)

(22). To examine variation at the individual protein level, further within-species MSAs were produced

for each of the 12 protein sequences in the BGC. Each of these alignments was used as the basis for a

consensus sequence, produced using EMBOSS Cons v6.6.0.0, representing each of the four species

carrying the BGC. An MSA and per cent identity matrix were then generated for each protein

between the consensus sequences of *K. oxytoca*, *K. grimontii*, *K. michiganensis* and *K. pasteurii*,

along with the reference sequence (11).

The species affiliations of the genomes encoding the full kleboxymycin BGC were

determined using FastANI v1.2 (18) against genomes of type strains of the *K. oxytoca* and *K.*

pneumoniae complexes (1,23) and *K. aerogenes* ATCC 13048$^{\text{T}}$ (assembly accession number

GCA_003417445), with PhyloPhlAn 0.99 used to conduct a phylogenetic analysis to confirm species

affiliations. PhyloPhlAn identifies hundreds of conserved (core) proteins from a given genomic

dataset and uses them to build a complete high-resolution phylogeny.

RESULTS

Characterization of the clinical isolates

Although initial phenotypic tests (**Supplementary Material**) and genomic analyses (9,10)

identified PS_Koxy1, PS_Koxy2 and PS_Koxy4 as *K. oxytoca*, analyses of the isolates' proteomes

showed them to be *K. michiganensis* ST138 (phylogroup Ko1, *bla*$_{\text{OXY1-8}}$) (**Supplementary Figure 1**).

Full details of phenotypic characterization and genome sequencing of the clinical isolates can be

found in **Supplementary Material**. PS_Koxy1, PS_Koxy2 and PS_Koxy4 all shared 98.81 %, 98.71

% and 98.71 % ANI, respectively, with the type strain of *K. michiganensis* (W14$^{\text{T}}$, GCA_901556995),

and 99.98 to 100.00 % ANI with each other. Based on current recommendations, ANI of 95–96 %

and above with the genome of the type strain is indicative of species affiliation (24). Inclusion of the

genomes with representatives of all six species of the *K. oxytoca* complex in a phylogenetic analysis

confirmed the affiliation of PS_Koxy1, PS_Koxy2 and PS_Koxy4 with *K. michiganensis*

(**Supplementary Figure 2**).

Assigning MLST sequence types to species

184   While annotations for genomes are improving, we have previously noted and continue to

notice issues with identities attributed to *K. oxytoca* genomes in public repositories (3). Consequently,

the identity of all genomes included in this work was first confirmed by ANI analysis

(**Supplementary Table 1**), with bla_{OXY} gene and phylogenetic analyses supporting our findings

(**Supplementary Material**). Of the 178 *K. oxytoca* complex genomes identified, many had been

misassigned in GenBank: seven genomes were listed as *K. grimontii*, 106 as *K. oxytoca*, 51 as *K.*

michiganensis, 13 as *Klebsiella* sp. and one as *K. pneumoniae*. Our analyses of the 178 genomes

showed the dataset actually represented *K. michiganensis* (*n=76*), *K. oxytoca* (*n=66*), *K. grimontii*

(*n=24*), *K. pasteurii* (n=6), *K. huaxiensis* (*n=5*) and *K. spallanzanii* (*n=1*).

193   The *K. oxytoca* MLST scheme uses sequence polymorphisms among seven housekeeping

genes – *gapA*, *infB*, *mdh*, *pgi*, *phoE*, *rpoB*, *tonB* – to generate sequence types for isolates. Currently,

there are 442 allele sequences that contribute to 354 unique MLST sequence types. We first identified

nucleotide sequences within the genomes with exact matches to nucleotide sequences within the allele

reference dataset. One-hundred-and-twenty-nine genomes returned hits to known MLST profiles, and

10 isolates returned MLST profiles with no assigned sequence type (**Supplementary Table 2**). Our

clinical isolates returned the expected ST138 result.

200   Of the 66 *K. oxytoca* genomes, 59 could be assigned to known sequence types (in order of

abundance: ST2, ST176, ST199, ST36, ST19, ST30, ST53, ST101, ST18, ST31, ST34, ST48, ST58,

ST59, ST141, ST145, ST153, ST181, ST221, ST222, ST257, ST258, ST287, ST323) and one

(GCA_003937225) represented a novel sequence type. Of the 24 *K. grimontii* genomes, 13 could be

assigned to known sequence types (ST172, ST216, ST104, ST186, ST236, ST263, ST316, ST319,

ST350), with four (GCA_002856195, GCA_900451335, GCA_008120915, GCA_004343645)

representing unique novel sequence types. Of the six *K. pasteurii* genomes, three could be assigned to

known sequence types (ST47, ST311, ST351) and one (GCA_901563825) represented a novel

sequence type. Of the 79 *K. michiganensis* genomes (including our three clinical isolates), 57 could be

assigned to known sequence types (ST85, ST27, ST202, ST143, ST29, ST50, ST84, ST138, ST11,

ST88, ST317, ST28, ST40, ST52, ST82, ST92, ST98, ST108, ST127, ST144, ST146, ST157, ST170,

ST180, ST226, ST294, ST315), with four genomes (GCA_000783895, GCA_000735215,

GCA_007097185, GCA_007097115) representing three novel sequence types. None of the *K.*

huaxiensis or *K. spallanzanii* genomes returned hits to known alleles (**Supplementary Table 2**), but

the relevant individual housekeeping gene sequences are provided as **Supplementary Files** for use by

other researchers.

For those genomes that encoded known or novel sequence types, we concatenated their

housekeeping-gene sequences and used them to create a MSA with the concatenated sequences of

each of the 354 recognized MLST sequence types. This MSA was used to create a phylogenetic tree,

allowing us to visualize the relationships among species and sequence types (**Figure 1**).

Of the 354 known MLST sequence types, 342 (96.6 %) were associated with specific

members of the *K. oxytoca* complex (**Supplementary Table 2**): 115 with *K. oxytoca*, 130 with *K.*

michiganensis, 73 with *K. grimontii* and 24 with *K. pasteurii*. Eleven were associated with

unspecified members of the *K. oxytoca* complex. ST105 was associated with *Raoultella*

ornithinolytica, sharing 99.73 % sequence similarity type strain's MLST profile. *K. oxytoca*-specific

sequence types shared 98.64–100 % sequence similarity, *K. michiganensis*-specific sequence types

shared 96.62–100.00 % sequence similarity, *K. grimontii*-specific sequence types shared 98.20–

100.00 % sequence similarity, *K. pasteurii*-specific sequence types shared 99.00–100.00 % sequence

similarity and *K. huaxiensis*-specific sequence types shared 97.09–99.7 % sequence similarity. A

matrix of similarity values for the 504 sequences included in the analysis is available in

Supplementary Material, along with the MSA alignment used to generate the phylogenetic tree

shown in **Figure 1**.

Detection of the complete kleboxymycin BGC in clinical isolates

It has long been known that *K. oxytoca* gut colonization is linked with AAHC (16). Schneditz

et al. (12) showed tillivaline (TV), a pyrrolobenzodiazepine (PBD) derivative produced by *K. oxytoca*,

is one of the enterotoxins responsible for causing AAHC. This toxic product is encoded by the

heterologous expression of the kleboxymycin [also known as tilimycin (TM) (14)] BGC comprising

12 genes (11). Protein sequences of the reference sequence (11) were used to create a BLASTP

database against which the proteins encoded in the genomes of PS_Koxy1, PS_Koxy2 and PS_Koxy4

were compared. The genomes of PS_Koxy1, PS_Koxy2 and PS_Koxy4 encoded a complete

kleboxymycin BGC (**Figure 2**). All genes in each of the genomes shared >99 % identity and >99 %

query coverage with the genes of the reference sequence (12): *mfsX*, 99.76 % identity; *uvrX*, 99.87 %;

hmoX, 99.80 %; *adsX*, 99.85 %; *icmX*, 99.52 %; *dhbX*, 99.62 %; *aroX*, 99.74 %; *npsA*, 99.80 %; *thdA*,

98.68 %; *npsB*, 99.93 %; *npsC*, 98.47 %; *marR*, 99.39.

Our strains were *K. michiganensis* ST138, so we downloaded and assembled (from

BioProject PRJEB30858) available raw sequence data from 19 *K. oxytoca* ST138 strains described

recently (10) and determined whether they were in fact *K. michiganensis* and encoded the

kleboxymycin BGC. All strains were confirmed to be *K. michiganensis* on the basis of ANI analysis,

and encoded the complete kleboxymycin BGC (**Supplementary Figure 4**).

Schneditz *et al.* (12) reported *npsA*/*npsB* were functionally conserved in six sequenced strains

of *K. oxytoca* (**Table 1**), based on a BLASTP analysis. Full details of the analysis are unavailable,

with only a brief mention of presence being determined based on BLASTP sequence identities >90 %

with no indication of sequence coverage. All the genomes included in the study of Schneditz *et al.*

(12) were compared with those of the type strain of *K. oxytoca* and related species to confirm their

species affiliations (**Table 1**). While some strains were *K. oxytoca*, others belonged to *K.*

michiganensis, *K. pasteurii*, *K. grimontii* and *R. ornithinolytica*. Using thresholds of 70 % identity and

70 % query coverage in our BLASTP analyses to reduce the potential for detecting false positives, we

reanalysed the genomes included in the study of Schneditz *et al.* (12). Our results agreed with those of

Schneditz *et al.* (12) for all genomes, except we detected *npsA*/*npsB* (and all other genes encoded in

the kleboxymycin BGC) in *K. grimontii* SA2. *K. oxytoca* 10–5243, *K. pasteurii* 10–5250, *K. oxytoca*

11492-1, *K. oxytoca* 10–5248 and *K. grimontii* M5a1 also encoded the whole kleboxymycin BGC. All

genes in all matches shared greater than 90 % identity across greater than 99 % query coverage. *K.*

michiganensis 10–5242, E718 and KCTC 1686 did not encode homologues associated with the

kleboxymycin BGC. *K. oxytoca* 10–5245 encoded almost-complete homologues of four genes

[EHS96696.1 (*marA*) 98.79 % identity, 99.39 % coverage; EHS96697.1 (*npsC*) 95.38 % identity,

99.23 % coverage; (EHS96698.1 (*mfsX*) 96.68 % identity, 99.87 % coverage; EHS96699.1 (*uvrX*)

94.88 % identity, 99.76 % coverage] in contig JH603137.1.

Detection of the kleboxymycin BGC in the faecal microbiota of preterm infants

Our previous work had highlighted the preterm infant gut microbiota harbours a range of

species belonging to the *K. oxytoca* complex (3). BLASTP searches of the two *K. michiganensis*

(P049A W, GCA_008120305; P095L Y, GCA_008120085) and three *K. grimontii* (P038I,

GCA_008120465; P043G P, GCA_008120425; P079F P, GCA_008120915) strains we previously

characterized showed all three *K. grimontii* strains encoded the kleboxymycin BGC (**Supplementary**

Figure 5). All BGC genes in their genomes shared >98 % identity and >99 % query coverage with the

genes of the reference sequence (11): *mfsX*, 100 % identity; *uvrX*, 99.60–99.73 %; *hmoX*, 99.80 %;

adsX, 99.69 %; *icmX*, 100 %; *dhbX*, 100 %; *aroX*, 99.94–99.74 %; *npsA*, 99.21–99.41 %; *thdA*, 98.68

%; *npsB*, 99.31–99.38 %; *npsC*, 99.23–100 %; *marR*, 100 %. The BGC was also detected in 8/25 of

the preterm-associated *K. oxytoca* complex MAGs (3 *K. oxytoca*, 5 *K. michiganensis*) we described

previously (3). An MSA of the preterm-associated genomes' BGC against the reference sequence (11)

suggested species-specific clustering of the sequences (**Supplementary Figure 5**).

Prevalence of the kleboxymycin BGC in *Klebsiella* spp.

Given the work detailed above had detected the kleboxymycin BGC in several different but

closely related *Klebsiella* species and in a range of clinical and gut-associated isolates, and Hubbard *et*

al. (20) recently detected the BGC in a strain of *K. grimontii*, we chose to increase the scope of our

analysis to include 7,170 publicly available assembled *Klebsiella* genomes (including our three

clinical strains, and five isolates from preterm infants (3)) (**Supplementary Table 1**).

As mentioned above, we have noted issues with identities attributed to *Klebsiella* genomes in

public repositories (3), so the identity of all non-*K. oxytoca* complex genomes included in this work

was first confirmed by ANI analysis (**Supplementary Table 1**). The majority (*n*=6,245) of the

additional genomes were *K. pneumoniae*, followed by *K. variicola* subsp. *variicola* (*n*=241), *K.*

quasipneumoniae subsp. *similipneumoniae* (*n*=184), *K. aerogenes* (*n*=168), *K. quasipneumoniae*

subsp. *quasipneumoniae* (*n*=120), *K. variicola* subsp. *tropica* (*n*=19), 'K. quasivariicola' (*n*=11) and

K. africana (*n*=1). Out of 7,170 genomes, 110 (1.5 %) had one or more matches with the 12 genes

encoded within the kleboxymycin BGC reference sequence, with all except two genomes (both *K.*

pneumoniae) belonging to species of the *K. oxytoca* complex (**Supplementary Table 3**). Ninety-six

genomes – all belonging to the *K. oxytoca* complex – encoded at least 12 genes belonging to the BGC

(**Supplementary Table 3**), and were examined further.

One genome (GCA_002856195) encoding 12 BGC genes was found to encode two stretches

of the same protein with the other cluster-associated genes non-contiguous, while one

(GCA_004005605) encoded 13 BGC genes (one gene duplicated) in a non-contiguous arrangement.

Fifty-five out of 66 (83.3 %) *K. oxytoca* genomes encoded the entire kleboxymycin BGC, as did

19/24 (79.2 %) *K. grimontii*, 9/79 (11.4 %) *K. michiganensis* and 5/6 (83.3 %) *K. pasteurii* genomes

(**Figure 3a**). Phylogenetic analysis (**Figure 3b**) confirmed findings from ANI analyses

(**Supplementary Table 1**) that showed all genomes belonged to species of the *K. oxytoca* complex.

The 88 genomes confirmed to encode the complete kleboxymycin BGC included the type strain of *K.*

grimontii. The BGC cluster sequences grouped according to species, and the reference sequence (11)

clustered with *K. grimontii* sequences and was closely related to the type strain of that species (**Figure**

3c).

Species-specific consensus sequences were generated for all genes within the kleboxymycin

BGC and are available as supplementary material from figshare. Similarity values for each gene

within the BGC consensus sequences across the four species are available in **Supplementary Table**

4.

DISCUSSION

Genotypic characteristics of the three clinical *K. michiganensis* strains

The three clinical strains characterized herein had previously been included in a study of

outbreak strains encoding GES-5 and CTX-M-15 (9), the first report of GES-5-positive clinical

isolates of *K. oxytoca* ST138 in the UK. Subsequently, it has been shown that the GES-5 gene in these

strains is encoded on an IncQ group plasmid (10). The whole-genome sequence data reported on

previously (9) were not available to us. API 20E (this study; **Supplementary Material**), MALDI-

TOF and limited sequence analysis (9) had shown the strains to be *K. oxytoca*. Our previous work

with isolates recovered from preterm infants had shown that API 20E testing on its own was

insufficient to accurately identify *K. oxytoca* strains (3). The strains described by Eades *et al.* (9) were

characterized before the availability of MALDI-TOF databases capable of splitting species of the *K.*

oxytoca complex (MALDI-TOF was only able to identify as *K. oxytoca* but did not have sufficient

resolution to identify individual species within the complex) (1). As we are using PS_Koxy1,

PS_Koxy2, PS_Koxy4 in ongoing phage work, we generated draft genome sequences for the strains,

to accurately identify them and facilitate detailed host–phage studies in the future.

ANI and phylogenetic analyses confirmed all three strains belonged to the species *K.*

michiganensis, not *K. oxytoca* (**Supplementary Material**). In addition to the AMR genes GES-5 (β-

lactamase with carbapenemase activity) and CTX-M-15 (an ESBL responsible for resistance to

cephalosporins) reported previously (9), the strains encoded SHV-66, an ESBL not previously

reported in *K. oxytoca* and related species (**Supplementary Material**). SHV-66 has previously only

been reported in a minority of β-lactamase-producing *K. pneumoniae* in Guangzhou, China (25). In

this study, SHV-66 (99.65 % identity, bit-score 580 – strict CARD match) was also found in *K.*

michiganensis strains E718 (26), GY84G39 (unpublished), K1439 (unpublished) and

2880STDY5682598 (7) (accession numbers GCA_000276705, GCA_001038305, GCA_002265195

and GCA_900083915, respectively), included in the phylogenetic analysis shown in **Supplementary**

Figure 1. Moradigaravand *et al.* (7) noted in their study that 2880STDY5682598 encoded a bla_{SHV}

gene, but did not document its type nor indicate its novelty.

The three strains had identical virulence factor profiles (**Supplementary Figure 3b**),

encoding the plasminogen activating omptin Pla, the Mg^{2+} transport proteins MgtBC, Hsp60,

autoinducer-2 (LuxS), type I fimbriae, type 3 fimbriae, type 6 secretion system I, *Escherichia coli*

common pilus and enterobactin. They also encoded numerous proteins associated with capsule,

regulation of capsule synthesis (RcsAB) and LPS, with several of the latter sharing identity with

Haemophilus endotoxins (RfaD, GalU, LpxC, GmhA/LpcA, KdsA). All six proteins required for

allantoin utilization were encoded in the strains' genomes.

No capsule or O antigen types could be assigned to the strains using Kaptive, but all three

strains were best matched with KL68 [PS_Koxy1, 17/18 genes matched (*cpsACP* missing);

PS_Koxy2 and PS_Koxy4, 16/18 genes matched (*cpsACP* and KL68_18 missing)] and O1v1 [4/7

genes (*wzm*, *wzt*, *glf*, *wbbO*) matched in all strains].

MLST sequence types can be used to speciate members of the *K. oxytoca* complex

The bla_{OXY} gene diversified in parallel to housekeeping genes in the *K. oxytoca* complex, and

it is already known that *rpoB* – one of the seven genes included in the *K. oxytoca* MLST scheme (8) –

can be used to speciate members of the complex (2). Given that our three clinical strains were ST138

and belonged to *K. michiganensis*, we determined whether specific sequence types within the MLST

scheme could be assigned to species. We found that all species of the *K. oxytoca* complex are

associated with specific sequence types. In addition, we identified 10 novel MLST sequence types

that can be used to identify *K. grimontii*, *K. michiganensis*, *K. oxytoca* and *K. grimontii* genomes

(**Supplementary Table 2**).

Herzog *et al.* (27), when originally describing the *K. oxytoca* MLST scheme to characterize

clinical isolates, showed their concatenated sequence data for 74 clinical *K. oxytoca* isolates were

associated with three clusters (A, B1 and B2). Comparison of their sequence types with our

annotations shows that cluster A represents *K. oxytoca*, cluster B1 represents *K. michiganensis* and

cluster B2 represents *K. grimontii* and *K. pasteurii*.

The ability to use the *K. oxytoca* MLST scheme to speciate clinical isolates will be of

particular interest to clinical microbiologists in resource-limited settings who rely on the MLST

scheme for typing and epidemiological tracking of isolates in the absence of whole-genome sequence

data. It should also be noted that ribosomal MLST (28) (rMLST) available via the Species ID portal

of the PubMLST website allows those working with genome sequence data derived from *K. oxytoca*

complex isolates to speciate isolates. This resource uses 53 genes encoding the bacterial ribosome

protein subunits (*rps* genes) to rapidly characterize genomic data to the species level.

The identification of ST105 as belonging to *Raoultella ornithinolytica* indicates this sequence

type should be withdrawn from the *K. oxytoca* MLST scheme.

Distribution of the kleboxymycin BGC in *Klebsiella* spp.

As relatively little is known about the virulence factors of *K. oxytoca* and related species, and

the VFDB is limited with respect to the number of *Klebsiella* spp. on which it reports information, we

wanted to see whether our strains encoded the kleboxymycin BGC responsible for generating

microbiome-associated metabolites known to directly contribute to AAHC (11,12). The cytotoxic

nature of a heat-stable, non-proteinaceous component of spent media from *K. oxytoca* strains isolated

from patients with AAHC was first reported in 1990 (29). With respect to *K. oxytoca* being a

causative agent of AAHC, the bacterium has fulfilled Koch's postulates (15). While a commensal of

the gut microbiota of some individuals, it has been suggested that cytotoxic *K. oxytoca* is a transient

member of the gut microbiota (29).

TV is a PBD produced by *K. oxytoca* and is a causative agent of AAHC (12). The TV

biosynthesis genes are encoded on a non-ribosomal peptide synthase operon and include *npsA, thdA*

and *npsB*. The genes *aroX* and *aroB* are also essential for TV production (13). *npsA*, *thdA*, *npsB* and

aroX are located on a pathogenicity island (PAI). In clinical isolates, the PAI was present in 100 % of

toxin-producing isolates, but only 13 % of non-toxin-producing isolates (12). AAHC is characterized

by disruption of epithelial barrier function resulting from apoptosis of epithelial cells lining the colon.

TV exerts its apoptotic effect by binding to tubulin and stabilising microtubules, leading to mitotic

arrest (14).

A second PBD generated by the same pathway as TV has been identified (13). TM [also

called kleboxymycin (11)] has stronger cytotoxic properties than TV, having a PBD motif with a

hydroxyl group at the C11 position, while TV has an indole ring. When deprived of indole by the

inactivation of the indole-producing tryptophanase gene *tnaA*, *K. oxytoca* produces TM but not TV.

TV production is restored with the addition of indole, as indole spontaneously reacts with TM to

produce TV. Limited interconversion between TM and TV may also occur spontaneously *in vivo* (11).

TM is a genotoxin and triggers apoptosis by interacting with DNA, which leads to the activation of

damage repair mechanisms, causing DNA strand breakage (14). DNA interaction is prevented in the

case of TV by its indole ring, and both the molecular targets and apoptotic mechanisms of TM and

TV are distinct. The kleboxymycin BGC is not native to *K. oxytoca*, nor the wider

Enterobacteriaceae. Instead, the BGC is thought to have been acquired via horizontal gene transfer

from *Xenorhabdus* spp., which in turn acquired the BGC from bacteria of the phylum *Actinobacteria*

(11).

In the current study, we found the kleboxymycin BGC in our *K. michiganensis* isolates and

that it was common among four species of the *K. oxytoca* complex, with *K. oxytoca* and *K. grimontii*

strains making the largest contribution and the type strains of *K. grimontii* and *K. pasteurii* encoding

the BGC (**Figure 3**). Prior to this study, sequences from two *K. oxytoca* strains (MH43-1, GenBank

accession number MF401554 (11); AHC-6, GenBank accession number HG425356 (12)) were

available for the kleboxymycin BGC. Draft genome sequences do not appear to be publicly available

for either of these strains. However, our analysis of the kleboxymycin BGC across the *K. oxytoca*

complex has shown that MH43-1 is a strain of *K. grimontii* (**Figure 3c**). Hubbard *et al.* (20) recently

reported on a strain of *K. grimontii* that encoded the BGC, based on antiSMASH analysis.

Comparison of the AHC-6 sequence with that of MH43-1 and other sequences included in this study

shows AHC-6 is a strain of *K. oxytoca* (99.0–99.55 % nucleotide similarity with the BGCs encoded

by the three *K. oxytoca* MAG sequences included in **Supplementary Figure 5**). It is likely that as

more genomes of *K. oxytoca* complex species are deposited in public databases, the range of species

encoding the kleboxymycin BGC will increase.

All three of our *K. michiganensis* strains encoded the kleboxymycin BGC (**Figures 2 and 3**),

as did strains of *K. grimontii* we previously isolated from preterm infants and *K. oxytoca* and *K.*

michiganensis MAGs recovered from publicly available shotgun metagenomic data (**Supplementary**

Figure 5). Whether the BGCs encoded in our clinical and infant-associated strains are functional will

be the subject of future studies. The discovery of the kleboxymycin BGC in strains and MAGs

recovered from preterm infants is of particular concern. Gut colonization is linked with AAHC, with

disease caused by the overgrowth of cytotoxin-producing strains secondary to use of antibiotics (16).

AAHC presents as diffuse mucosal oedema and haemorrhagic erosions (16), and patients pass bloody

diarrhoea (30). The gut microbiota of preterm infants is shaped by the large quantity of antibiotics

these infants are given immediately after birth to cover possible early onset infection, with 'blooms'

of bacteria preceding onset of infection (3). Blood in the stool is frequently associated with

necrotizing enterocolitis (NEC) in preterm infants, which shares similar pathological hallmarks to

AAHC – i.e. intestinal necrosis. Notably, NEC is difficult to diagnose in the early stages and is often

associated with sudden serious deterioration in infant health, with treatment options limited due to

emerging multi-drug-resistant bacteria associated with disease. Previous studies have linked

Klebsiella spp. to preterm NEC (supported by corresponding clinical observations), with bacterial

overgrowth in the intestine linked to pathological inflammatory cascades, facilitated by a 'leaky'

epithelial barrier and LPS–TL4 activation. Recent work has demonstrated *K. oxytoca* complex isolates

of ST173, ST246 and a novel ST (7-32-38-44-69-25-43) recovered from infants with NEC can

produce kleboxymycin (TM) and TV (31). Using our MLST annotation scheme (**Supplementary**

Table 2), we determine these sequence types represent *K. grimontii*, *K. grimontii* and *K. pasteurii*,

respectively, with rMLST analyses of the whole-genome sequence data of Paveglio *et al.* (31)

confirming our findings (rST 124484, rST 124487 and rST 157090, respectively). Taken together

with the results from our study, we suggest specific virulence factors – i.e. kleboxymycin-related

metabolites encoded by atypical *Klebsiella* spp. – may also play a role in NEC, and this warrants

further study.

Attempts have been made to link specific subtypes of *K. oxytoca* to AAHC (32). Cytotoxic

effects were limited to *K. oxytoca*, with faecal (and to a lesser extent skin) isolates of *K. oxytoca* most

commonly associated with cytotoxicity (32). No genetic relationship was associated with cytotoxic

strains based on pulsed-field gel electrophoresis, and 31/97 strains exhibited evidence of cytotoxin

production (i.e. reduced viability of Hep2 cells). Joainig *et al.* (32) isolated genetically distinct

cytotoxin-positive and -negative strains from one AAHC patient, leading them to suggest that, when

detected in faeces, *K. oxytoca* should be considered an opportunistic pathogen able to produce disease

upon antibiotic treatment. They also found that, in patients with acute or chronic diarrhoeal diseases,

more than half of the isolates recovered were cytotoxin-positive. Given that *K. oxytoca*-related species

are not routinely screened for in such samples, it is possible that kleboxymycin-producing isolates

may make a greater contribution to diarrhoeal diseases than currently recognized, especially in

patients suffering from non-*C. difficile*-associated disease. We have shown that there are species-

specific differences in the kleboxymycin BGC (**Figure 3c**). These differences may have implications

for virulence of strains and warrant further study. It is hoped that the identification of an increased

range of strains (including type strains) encoding the kleboxymycin BGC will facilitate such studies.

ACKNOWLEDGEMENTS

Consultant microbiologist Dr Frances Davies (Imperial College Healthcare NHS Trust) is

thanked for providing access to clinical strains. Imperial Health Charity is thanked for contributing to

registration fees for the Professional Doctorate studies of PS. This work used the computing resources

of CLIMB (33), the UK MEDical BIOinformatics partnership (UK Med-Bio, supported by Medical

Research Council grant number MR/L01632X/1) and the Nottingham Trent University Hamilton

High Performance Computing Cluster. The research placement of FM was funded by Nottingham

Trent University. LJH is funded by a Wellcome Trust Investigator Award (no. 100/974/C/13/Z); a

BBSRC Norwich Research Park Bioscience Doctoral Training grant no. BB/M011216/1 (supervisor

LJH, student MK); an Institute Strategic Programme Gut Microbes and Health grant no.

BB/R012490/1 and its constituent projects BBS/E/F/000PR10353 and BBS/E/F/000PR10356; and an

Institute Strategic Programme Gut Health and Food Safety grant no. BB/J004529/1 to LJH. PS did all

phenotypic characterization work. PS did initial bioinformatics analyses (assembly, annotation,

CARD, initial BGC work), while FM and LH undertook the large-scale BGC analyses and infant-

associated BGC work; LH undertook the MLST annotation work; ALM and MK prepared and pre-

processed DNA for sequencing; LJH, ALM and LH supervised the work. We would like to thank

Dave Baker and the QIB core sequencing team for WGS library preparation and sequencing. All

authors contributed to interpretation and analyses of data, and writing of the manuscript.

REFERENCES

1.   Merla C, Rodrigues C, Passet V, Corbella M, Thorpe HA, Kallonen TVS, et al. Description of

Klebsiella spallanzanii sp. nov. and of *Klebsiella pasteurii* sp. nov. Front Microbiol.

2019;10:2360.

2.   Passet V, Brisse S. Description of *Klebsiella grimontii* sp. nov. Int J Syst Evol Microbiol. 2018

Jan;68(1):377–81.

3.   Chen Y, Brook TC, Soe CZ, O'Neill I, Alcon-Giner C, Leelastwattanagul O, et al. Preterm

infants harbour diverse *Klebsiella* populations, including atypical species that encode and

produce an array of antimicrobial resistance- and virulence-associated factors. Microb Genom.

2020 May 21;761924.

4.   Zheng B, Xu H, Yu X, Lv T, Jiang X, Cheng H, et al. Identification and genomic

characterization of a KPC-2-, NDM-1- and NDM-5-producing *Klebsiella michiganensis* isolate.

J Antimicrob Chemother. 2018 01;73(2):536–8.

5.   Pedersen T, Sekyere JO, Govinden U, Moodley K, Sivertsen A, Samuelsen Ø, et al. Spread of

plasmid-encoded NDM-1 and GES-5 carbapenemases among extensively drug-resistant and

pandrug-resistant clinical *Enterobacteriaceae* in Durban, South Africa. Antimicrob Agents

Chemother. 2018;62(5).

6.   Seiffert SN, Wüthrich D, Gerth Y, Egli A, Kohler P, Nolte O. First clinical case of KPC-3-

producing *Klebsiella michiganensis* in Europe. New Microbes New Infect. 2019

May;29:100516.

7.   Moradigaravand D, Martin V, Peacock SJ, Parkhill J. Population structure of multidrug-resistant

Klebsiella oxytoca within hospitals across the United Kingdom and Ireland identifies sharing of

virulence and resistance genes with *K. pneumoniae*. Genome Biol Evol. 2017 Mar 1;9(3):574–

84.

8.   Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb

software, the PubMLST.org website and their applications. Wellcome Open Res. 2018;3:124.

9.   Eades C, Davies F, Donaldson H, Hopkins K, Hill R, Otter J, et al. GES-5 carpabapenemase-

producing *Klebsiella oxytoca* causing clinical infection in a UK haematopoetic stem cell

transplantation unit. 26th European Congress of Clinical Microbiology and Infectious Diseases,

Amsterdam, The Netherlands; 2016.

10.   Ellington MJ, Davies F, Jauneikaite E, Hopkins KL, Turton JF, Adams G, et al. A multi-species

cluster of GES-5 carbapenemase producing *Enterobacterales* linked by a geographically

disseminated plasmid. Clin Infect Dis Off Publ Infect Dis Soc Am. 2019 Nov 20;

11.   Tse H, Gu Q, Sze K-H, Chu IK, Kao RY-T, Lee K-C, et al. A tricyclic pyrrolobenzodiazepine

produced by *Klebsiella oxytoca* is associated with cytotoxicity in antibiotic-associated

hemorrhagic colitis. J Biol Chem. 2017 24;292(47):19503–20.

12.   Schneditz G, Rentner J, Roier S, Pletz J, Herzog KAT, Bücker R, et al. Enterotoxicity of a

nonribosomal peptide causes antibiotic-associated colitis. Proc Natl Acad Sci U S A. 2014 Sep

9;111(36):13181–6.

13.   Dornisch E, Pletz J, Glabonjat RA, Martin F, Lembacher-Fadum C, Neger M, et al. Biosynthesis

of the Enterotoxic pyrrolobenzodiazepine natural product tilivalline. Angew Chem Int Ed Engl.

2017 13;56(46):14753–7.

14.   Unterhauser K, Pöltl L, Schneditz G, Kienesberger S, Glabonjat RA, Kitsera M, et al. *Klebsiella*

oxytoca enterotoxins tilimycin and tilivalline have distinct host DNA-damaging and

microtubule-stabilizing activities. Proc Natl Acad Sci U S A. 2019 26;116(9):3774–83.

15.   Högenauer C, Langner C, Beubler E, Lippe IT, Schicho R, Gorkiewicz G, et al. *Klebsiella*

oxytoca as a causative organism of antibiotic-associated hemorrhagic colitis. N Engl J Med.

2006 Dec 7;355(23):2418–26.

16.   Beaugerie L, Metz M, Barbut F, Bellaiche G, Bouhnik Y, Raskine L, et al. *Klebsiella oxytoca* as

an agent of antibiotic-associated hemorrhagic colitis. Clin Gastroenterol Hepatol Off Clin Pract

J Am Gastroenterol Assoc. 2003 Sep;1(5):370–6.

17.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinforma Oxf Engl. 2014 Jul

15;30(14):2068–9.

18.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun. 2018

30;9(1):5114.

19.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and SAMtools. Bioinforma Oxf Engl. 2009 Aug 15;25(16):2078–9.

20.   Hubbard ATM, Newire E, Botelho J, Reiné J, Wright E, Murphy EA, et al. Isolation of an

antimicrobial-resistant, biofilm-forming, *Klebsiella grimontii* isolate from a reusable water

bottle. MicrobiologyOpen. 2020 Mar 3;e1023.

21.   Schliep KP. phangorn: phylogenetic analysis in R. Bioinforma Oxf Engl. 2011 Feb

15;27(4):592–3.

22.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.

Nucleic Acids Res. 2019 Jul 2;47(W1):W256–9.

23.   Rodrigues C, Passet V, Rakotondrasoa A, Diallo TA, Criscuolo A, Brisse S. Description of

Klebsiella africanensis sp. nov., *Klebsiella variicola* subsp. *tropicalensis* subsp. nov. and

Klebsiella variicola subsp. *variicola* subsp. nov. Res Microbiol. 2019 May;170(3):165–70.

24.   Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, et al. Proposed minimal

standards for the use of genome data for the taxonomy of prokaryotes. Int J Syst Evol

Microbiol. 2018 Jan;68(1):461–6.

25.   Zuo B, Liu Z-H, Wang H-P, Yang Y-M, Chen J-L, Ye H-F. [Genotype of TEM- and SHV-type

beta-lactamase producing *Klebsiella pneumoniae* in Guangzhou area]. Zhonghua Yi Xue Za

Zhi. 2006 Nov 7;86(41):2928–32.

26.   Liao T-L, Lin A-C, Chen E, Huang T-W, Liu Y-M, Chang Y-H, et al. Complete genome

sequence of *Klebsiella oxytoca* E718, a New Delhi metallo-β-lactamase-1-producing

nosocomial strain. J Bacteriol. 2012 Oct;194(19):5454.

27.   Herzog KAT, Schneditz G, Leitner E, Feierl G, Hoffmann KM, Zollner-Schwetz I, et al.

Genotypes of *Klebsiella oxytoca* isolates from patients with nosocomial pneumonia are distinct

from those of isolates from patients with antibiotic-associated hemorrhagic colitis. J Clin

Microbiol. 2014 May;52(5):1607–16.

28.   Jolley KA, Bliss CM, Bennett JS, Bratcher HB, Brehony C, Colles FM, et al. Ribosomal

multilocus sequence typing: universal characterization of bacteria from domain to strain.

Microbiol Read Engl. 2012 Apr;158(Pt 4):1005–15.

29.   Higaki M, Chida T, Takano H, Nakaya R. Cytotoxic component(s) of *Klebsiella oxytoca* on

HEp-2 cells. Microbiol Immunol. 1990;34(2):147–51.

30.   Youn Y, Lee SW, Cho H-H, Park S, Chung H-S, Seo JW. antibiotics-associated hemorrhagic

colitis caused by *Klebsiella oxytoca*: two case reports. Pediatr Gastroenterol Hepatol Nutr. 2018

Apr;21(2):141–6.

31.   Paveglio S, Ledala N, Rezaul K, Lin Q, Zhou Y, Provatas AA, et al. Cytotoxin-producing

Klebsiella oxytoca in the preterm gut and its association with necrotizing enterocolitis. Emerg

Microbes Infect. 2020 Dec;9(1):1321–9.

32.   Joainig MM, Gorkiewicz G, Leitner E, Weberhofer P, Zollner-Schwetz I, Lippe I, et al.

Cytotoxic effects of *Klebsiella oxytoca* strains isolated from patients with antibiotic-associated

hemorrhagic colitis or other diseases caused by infections and from healthy subjects. J Clin

Microbiol. 2010 Mar;48(3):817–24.

33.   Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, et al. CLIMB (the

Cloud Infrastructure for Microbial Bioinformatics): an online resource for the medical

microbiology community. Microb Genomics. 2016;2(9):e000086.

34.   Segata N, Börnigen D, Morgan XC, Huttenhower C. PhyloPhlAn is a new method for improved

phylogenetic and taxonomic placement of microbes. Nat Commun. 2013;4:2304.

Table 1. Genomes included in analyses conducted by Schneditz *et al.* (12) with corrected

species affiliations (originally reported as *K. oxytoca*)

Assembly accession	Strain	Species	ANI with shown genome*	*npsA/npsB*
GCA_000240325.1	KCTC 1686	*K. michiganensis*	98.69 %, GCA_901556995.1	–
GCA_000247835.1	10–5242	*K. michiganensis*	97.59 %, GCA_901556995.1	–
GCA_000247855.1	10–5243	*K. oxytoca*	99.31 %, GCA_900977765.1	+
GCA_000247875.1	10–5245	*K. oxytoca*	99.13 %, GCA_900977765.1	–
GCA_000247895.1	10–5246	*Raoultella ornithinolytica*	99.21 %, GCA_001598295.1	–
GCA_000247915.1	10–5250	*K. pasteurii*	99.29 %, GCA_901563825.1	+
GCA_000252915.3	11492-1	*K. oxytoca*	99.15 %, GCA_900977765.1	+
GCA_000276705.2	E718	*K. michiganensis*	98.37 %, GCA_901556995.1	–
GCA_000427015.1	SA2	*K. grimontii*	99.33 %, GCA_900200035.1	+
GCA_001078235.1	10–5248	*K. oxytoca*	99.25 %, GCA_900977765.1	+
GCA_001633115.1	M5a1	*K. grimontii*	99.40 %, GCA_900200035.1	+

589 *GCA_901556995.1 = *K. michiganensis* W14[T]; GCA_900200035.1, *K. grimontii* 06D021[T];

590 GCA_900977765.1 = *K. oxytoca* ATCC 13182[T]; GCA_001598295.1 = *R. ornithinolytica*

591 NBRC 105727[T]; *K. pasteurii* SB3355[T] GCA_901563825.1.

Legend:
- *Raoultella* spp.
- *K. spallanzanii* (Ko3)
- *K. huaxiensis* (Ko8)
- *K. oxytoca* (Ko2)
- *K. pasteurii* (Ko4)
- *K. grimontii* (Ko6)
- *K. michiganensis* (Ko1, Ko5)

Tree scale: 0.01

Figure 1. Sequence types within the *K. oxytoca* MLST scheme can be used to speciate members of the *K. oxytoca* complex. The three *K. michiganensis* clinical isolates (all ST138) characterized in this study are shown in white. The phylogenetic tree (neighbour joining, Jukes Cantor) was generated using concatenated nucleotide sequences of housekeeping genes (*gapA–infB–mdh–pgi–phoE–rpoB–tonB*) used in the *K. oxytoca* MLST scheme (8). The purple circles represent bootstrap values ≥80 % (based on 1,000 replications); the larger the circle, the higher the bootstrap value. Scale bar, average number of nucleotide substitutions per position. The full list of MLST sequence types and their species affiliations are available in **Supplementary Table 2**.

603 **Figure 2.** Alignment of the kleboxymycin BGCs from the three clinical *K. michiganensis* strains with the complete cluster of *K. oxytoca* MH43-1 [GenBank

604 accession number MF401554 (11)]. (a) The image (alignment view) was generated via the progressiveMauve algorithm plugin of Geneious Prime v2019.2.1

605 (default settings, full alignment), with gene names for the three clinical isolates assigned manually. (b) Genes corresponding to Prokka-generated annotations.

606 Consensus identity is the mean pairwise nucleotide identity over all pairs in the column: green, 100 % identity; greeny-brown, at least 30 % and under 100 %

607 identity; red, below 30 % identity.

Figure 3. Distribution of the kleboxymycin BGC in *Klebsiella* spp. genomes. (a) Distribution of the

K. oxytoca complex genomes encoding the entire kleboxymycin BGC. (b) Unrooted maximum-

likelihood tree [generated using PhyloPhlAn v0.99 (34) and 380 protein-encoding sequences

conserved across the genomes] confirming species affiliations of the 88 genomes within the *K.*

oxytoca complex (1) encoding the kleboxymycin BGC. Type strains are shown with coloured

backgrounds corresponding to the legend. The clade associated with *K. huaxiensis* and *K. spallanzanii*

has been collapsed because of space constraints. (c) Maximum-likelihood tree generated with the

concatenated protein sequences for the kleboxymycin BGC of the 88 genomes found to encode all 12

genes of the BGC plus the reference sequence (11). The tree was rooted using the kleboxymycin-

encoding BGC of *Pectobacterium brasiliense* BZA12. Values at nodes, bootstrap values expressed as

a percentage of 100 replicates. Sources of isolates, where known, are shown to the right of the

assembly accession numbers. (b, c) Scale bar, average number of substitutions per position.