Maintenance of mitochondrial integrity in midbrain dopaminergic neurons governed by

a conserved developmental transcription factor

Federico Miozzo[1], Luca Stickely[1], Damla Tas[1,2], Nicolas Loncle[1,3], Irina Nikonenko[4],

Michaela Dorcikova[1], Eva P. Valencia-Alarcón[1], Peter Bou Dib[5,6] and Emi Nagoshi[1,*]

[1]Department of Genetics and Evolution and Institute of Genetics and Genomics of Geneva

(iGE3), University of Geneva, CH-1211, Geneva 4, Switzerland

[2]Present address: The Janssen Pharmaceutical Companies of Johnson & Johnson, Bern,

Switzerland

[3]Present address: Puma Biotechnology, Inc., Berkeley, California

[4]Department of Basic Neurosciences and the Center for Neuroscience, CMU, University of

Geneva, CH-1211, Geneva 4, Switzerland

[5]Institute of Cell Biology, University of Bern, CH-3012, Bern, Switzerland

[6]Present address: Department of Chemistry, Humboldt-Universität zu Berlin, 12489 Berlin,

Germany

[*]Corresponding author: Emi.Nagoshi@unige.ch

ABSTRACT

The degeneration of dopaminergic (DA) neurons in the substantia nigra is a hallmark of

Parkinson's Disease (PD). Dysregulation of developmental transcription factors is implicated

in dopaminergic neurodegeneration, but the underlying molecular mechanisms remain largely

unknown. *Drosophila Fer2* is a prime example of a developmental transcription factor

required for the birth and maintenance of midbrain DA neurons. Using an approach

combining ChIP-seq, RNA-seq, and genetic epistasis experiments with PD-linked genes, here

we demonstrate that *Fer2* controls a transcriptional network to maintain mitochondrial

structure and function, and thus confers dopaminergic neuroprotection against genetic and

oxidative insults. We further show that conditional ablation of *Nato3,* a mouse homolog of

Fer2, in differentiated DA neurons results in locomotor impairments and mitochondrial

abnormality in aged mice. Our results reveal the essential and conserved role of *Fer*2

homologs in the mitochondrial maintenance of midbrain DA neurons, opening new

perspectives for modelling and treating PD.

INTRODUCTION

Midbrain dopaminergic (mDA) neurons play a central role in controlling key brain functions,

including voluntary movement, associative learning, motivation and cognition[1]. The

progressive and selective degeneration of mDA neurons in the substantia nigra (SN) results in

a loss of nigrostriatal DA neurotransmission, leading to the motor symptoms of Parkinson's

disease (PD). Although PD was first described over 200 years ago and is the most prevalent

neurodegenerative movement disorder, the exact pathologic mechanisms underlying this

disease remain unclear. Treatment options are limited to symptomatic relief with DA

replacement therapy[2,3]. Thus, the identification of targets for disease-modifying therapies that

reverse or hinder neurodegeneration in PD is greatly needed.

PD is a multifactorial disorder caused by a combination of genetic and environmental

factors[4]. Increasing evidence indicates that PD risk factors lead to common pathological

processes, including excessive production of reactive oxygen species (ROS), axonal

pathology, neuroinflammation, defects in the ubiquitin-proteasome system, and

mitochondrial dysfunction[5,2]. In particular, mitochondrial impairment is integral to most other

pathological cellular processes and thought to play a central role in dopaminergic

neurodegeneration. Complex I activity is reduced in the brains of PD patients, and

mitochondrial toxins such as 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) and

rotenone potently induce SN degeneration. The findings that multiple genes linked to familial

PD, such as *PINK1*, *Parkin*, *LRRK2*, *SNCA* and *DJ-1*, directly or indirectly affect

mitochondrial physiology further support this notion[6,7].

Although several genes linked to rare monogenic forms of PD have been identified,

genetic susceptibility factors for sporadic PD, which accounts for approximately 90% of

cases, remain largely unknown. In this regard, it is noteworthy that dysregulation of

transcription factors required for mDA development is implicated in PD. Numerous

transcription factors required for various aspects of mDA neurons development, including

En1, *En2*, *Lmx1a*, *Lmx1b*, *Foxa1*, *Foxa2*, *Nurr1*, *Pitx3* and *Otx2*, remain expressed in the

adult brain and are crucial for the survival and function of differentiated mDA neurons in

mice[8–16]. Genome-wide association studies (GWAS) have shown that genetic variants of

developmental transcription factors are highly represented in sporadic PD patients[17,18].

Moreover, overexpression of *Nurr1*, *Foxa2*, *En1* or *Otx2* prevents the loss of DA neurons in

murine PD models[19–21]. Therefore, a better comprehension of the genetic networks controlled

by developmental transcription factors in differentiated mDA neurons may advance our

molecular understanding of neurodegeneration in PD and open new therapeutic options.

68   *Drosophila* offers a powerful model system to investigate molecular mechanisms of

PD pathogenesis *in vivo*, due to the abundance of genetic tools, the conservation of most PD-

related pathways and familial PD-linked genes, and its relatively simple nervous system[22,23].

The adult fly midbrain contains approximately a dozen clusters of DA neurons projecting to

different areas[24]. Although the anatomical arrangements of DA neurons in *Drosophila* and

vertebrates differ considerably, recent works have highlighted the functional homology of

some of the dopaminergic circuits[25]. The protocerebral anterior medial (PAM) cluster is the

largest subgroup containing approximately 80% of all *Drosophila* brain DA neurons. PAM

neurons are highly heterogeneous in their function and projection patterns, modulating

various behavior including associative learning, sleep, and locomotion[26,27]. Remarkably,

some of the PAM neurons are required for the startle-induced climbing, a locomotor behavior

found to be defective in multiple PD models[27–29], rendering the PAM cluster partially

analogous to the mammalian SN. Dopaminergic innervation from the PPL1 and PPM3

clusters to the central complex also appear to be functionally homologous to the nigrostriatal

pathway and is reported to be impaired in some PD models[25,30,31].

83   The notion that developmental transcription factors play critical roles in the

maintenance of adult DA neurons is shared between flies and mammals, as supported by our

recent studies of *Drosophila Fer2* gene. *Fer2* encodes a basic helix-loop-helix (bHLH)

transcription factor required for the development of DA neurons in the PAM cluster, and

fewer PAM neurons are formed in *Fer2²* hypomorphic mutants. Additionally, *Fer2²* mutant

flies display a progressive loss of PAM neurons in adulthood, locomotor impairment that can

be improved by L-DOPA administration, and increased ROS levels in the brain. PAM

neurons deficient for *Fer2* accumulate abnormal mitochondria and show autophagy

dysfunction. *Fer2* expression persists into adulthood in PAM neurons and increases in

response to oxidative stress. Importantly, the maintenance of mitochondrial integrity and the

survival of PAM neurons in aged flies require the presence of *Fer2* in PAM neurons in

adulthood[28,29]. Taken together, these lines of evidence establish the dual role of *Fer2* in

development and lifelong maintenance of DA neurons in the PAM cluster, and suggest that

molecular studies of *Fer2* function will shed light on the mechanisms of dopaminergic

neurodegeneration relevant to PD.

In the present study, we ask how *Fer2* maintains DA neurons throughout adulthood

and whether its function is conserved in mammals. We address these questions by analyzing

functional interaction between *Fer2* and familial PD-linked genes, by identifying *Fer2* target

genes and by generating mice with a conditional deletion of *Nato3*, a murine homolog of

Fer2, in differentiated DA neurons. This combined approach demonstrates that *Fer2* is a

master regulator of a transcriptional network controlling mitochondrial integrity and function,

and thereby confers dopaminergic neuroprotection in multiple genetic and toxin models of

PD. *Nato3* is known to be required for the genesis of mDA neurons[32–34] but its post-

developmental role has never been studied. We show here for the first time that selective

ablation of *Nato3* in differentiated DA neurons leads to age-dependent motor impairment and

a loss of mitochondrial integrity in mDA neurons. Taken together, our results indicate that

Fer2 and *Nato3* share an essential role in maintaining mitochondrial health and dopaminergic

function, offering new opportunities to study the selective vulnerability of mDA neurons and

to develop therapeutic interventions for PD.

RESULTS

***Fer2* overexpression protects PAM dopaminergic neurons from genetic and oxidative**

insults

To explore the genetic mechanisms underlying the role of *Fer2* in the maintenance of DA

neurons, we investigated whether *Fer2* loss or gain of function affects the survival of DA

neurons in genetic models of PD. Mutations in the *leucine-rich repeat kinase 2* (*LRRK2*) gene

are the most common cause of familial PD and also predispose carriers to sporadic PD[35].

Because pathogenic *LRRK2* variants are considered to be gain-of-function mutants, most fly

models of *LRRK2*-linked PD employ overexpression of wild-type or mutant forms of *LRRK2*

or its fly homolog. We thus overexpressed wild-type or mutant *Drosophila Lrrk* via the

GAL4/UAS system using PAM neuron-specific drivers, *Fer2-GAL4* or *R58E02-GAL4*[26,28].

The number of surviving DA neurons was assessed by immunostaining with anti-tyrosine

hydroxylase (TH) antibodies.

Forced expression of either wild-type or LRRK I1915T mutant, which is homologous

to the human pathogenic LRRK2 I2020T variant[35,36], leads to a loss of PAM neurons. A

neuronal loss of approximately 10% of was observed in 1-day-old flies, and thereafter the cell

loss progressed in an age-dependent manner, indicating that *Lrrk* gain of function impairs

both the development and the maintenance of PAM neurons (Supplementary Fig. 1a and Fig.

1a-b). Remarkably, overexpression of *Fer2* in PAM neurons prevented PAM neuron loss

induced by *Lrrk* gain of function (Fig. 1a-b). Forced expression of *Lrrk* in *Fer2²*

heterozygous background resulted in PAM neuron loss to a similar extent as in wild-type

background. *Lrrk* overexpression in *Fer2²* homozygous mutants additively enhanced PAM

neuron loss compared with either of the single genetic perturbations (Supplementary Fig. 1b).

These results suggest that *Fer2* and *Lrrk* act in independent pathways but *Fer2* can attenuate

Lrrk-induced neuronal damage.

Mutations in the *PRKN* gene, which encodes the E3 ubiquitin ligase Parkin, are linked

to autosomal recessive juvenile PD[37]. *Drosophila parkin* loss-of-function mutant *park¹*

exhibits PD-like phenotypes, including locomotor deficits, mitochondrial impairments and

loss or shrinkage of a few subclasses of DA neurons[38,39]. We examined *park¹* mutant brains

with anti-TH immunostaining and found that DA neuron loss occurs also in the PAM cluster

(Fig. 1c). A greater loss of PAM neuron was observed in *park¹*, *Fer2²* double mutants than in

either of the single mutants. The increase in cell loss was additive but not synergistic,

suggesting that *parkin* and *Fer2* do not act in the same linear pathway (Supplementary Fig.

1c). However, interestingly, the overexpression of *Fer2* in PAM neurons significantly

reduced *park¹*-induced cell loss (Fig. 1d).

Parkin mutations lead to severe defects in mitochondrial morphology and function in

all organisms studied thus far, including flies[40–43]. We therefore examined whether *Fer2*

overexpression ameliorates mitochondrial morphology in DA neurons of *park¹* flies by

expressing mitochondria-targeted GFP (mitoGFP) in PAM neurons. In contrast to the

reticular mitochondria observed in control flies, the majority of mitochondria appeared to be

more rounded in PAM neurons of *park¹* mutants (Fig. 1e). This observation was

quantitatively confirmed by the analysis of mitochondria circularity, which showed a shift

towards more globular mitochondria in *park¹* compared with control (*park¹*/+) flies (Fig. 1f).

Remarkably, the overexpression of *Fer2* in PAM neurons restored tubular mitochondrial

morphology in *park¹* flies (Fig. 1e-f).

The findings that *Fer2* overexpression cell-autonomously prevents PAM neuron loss

in two independent familial PD models and ameliorates mitochondrial morphology defects in

park[1] mutants suggests that *Fer2* gain of function might confer dopaminergic neuroprotection

against a broad spectrum of genetic and chemical insults. We therefore tested if *Fer2*

overexpression can protect DA neurons from damage caused by oxidative stress, using an

assay in which flies were exposed to a non-lethal dose of hydrogen peroxide (H_2O_2). 7-day-

old flies were fed with 5% H_2O_2 for 24h, and 7 days later DA neurons in the brains were

examined by anti-TH immunostaining. This treatment triggered the loss of approximately

10% of DA neurons in the PAM cluster in *w[1118]* flies, *Fer2[2]/+* flies and flies carrying a PAM-

specific driver. The PAM neuron loss was exacerbated in flies homozygous for the *Fer2[2]*

allele, consistent with our previous finding that *Fer2* inactivation renders PAM neurons more

vulnerable to oxidative stress[28,29]. In contrast, *Fer2* overexpression in PAM neurons

prevented H_2O_2-induced PAM neuron loss (Fig. 2a). These results indicate that *Fer2* gain of

function protects DA neurons in the PAM cluster from oxidative insults.

ROS production increases as flies age[44]. Interestingly, we found an age-dependent

increase in FER2 protein levels in PAM neurons, parallel to the augmentation of ROS levels

in the brain (Fig. 2b-e). Combined with the fact that *Fer2* loss of function causes a

progressive loss of PAM neurons, this finding suggests that FER2 is upregulated to protect

PAM neurons from age-dependent increases in endogenous oxidative stress. Taken together,

these results provide evidence that *Fer2* is a potent neuroprotective transcription factor for

PAM dopaminergic neurons, counteracting various genetic and oxidative insults.

Identification of *Fer2* direct targets by a combination of ChIP-seq and RNA-seq

approaches

To investigate how *Fer2* elicits dopaminergic neuroprotection, we wished to identify *Fer2*

transcriptional targets. To this end, we first determined the genome-wide binding sites of

FER2 using chromatin immunoprecipitation coupled to sequencing (ChIP-seq). Since FER2-

specific antibodies were unattainable, ChIP-seq was performed on the heads of 14-day-old

flies harboring a *Fer2::GFP::V5* genomic transgene in *Fer2^1* mutant background[28], using

anti-V5 antibody (Fig. 3a). *Fer2::GFP::V5* is functional and recapitulates the physiological

dosage and expression pattern of *Fer2*, as it rescues PAM neuron loss and locomotor defects

in *Fer2^1* mutants (Supplementary Fig. 2b-c) and is expressed in PAM neurons but scarcely in

other cells in the brain (Supplementary Fig. 2a), in agreement with existing single-cell RNA-

seq data[45,46]. We identified 269 FER2-bound regions, which were annotated to 279 putative

Fer2 target genes (Supplementary Table 1). FER2-bound sequences were highly enriched

with a 12-nucleotide motif containing a canonical bHLH binding site (CANNTG), validating

the specificity of the procedure (Fig. 3b).

The biding of transcription factors to chromatin does not necessarily lead to

transcriptional regulation of nearby genes. To determine which binding events regulate

proximal gene expression, we next identified transcripts that are up- or downregulated after a

transient overexpression of *Fer2* and compared these transcripts with the ChIP-seq data. The

heat-inducible *hs-GAL4* was used to ubiquitously but acutely express *Fer2,* and RNA-seq

was performed on fly heads collected 12 and 48 h after a 3-h heat shock treatment. The same

heat-shock treatment was applied to the flies carrying *hs-GAL4* but not *UAS-Fer2* to identify

and exclude from our analysis any heat-responsive gene (Fig. 3c). These experiments

identified 328 genes (12 h after heat shock) and 916 genes (48 h after heat shock)

differentially expressed in the head following *Fer2* overexpression (Fig. 3c and

Supplementary Table 2). Then, by comparing the head RNA-seq and *Fer2* ChIP-seq data, we

found 32 genes that were present in FER2 ChIP-seq and also differentially expressed 48 h

after *Fer2* overexpression. These genes are considered as potential *Fer2* transcriptional

targets. The overlap was statistically highly significant (hypergeometric probability = 0.0004).

In contrast, there was no statistically significant overlap between the ChIP-seq data and the

genes differentially expressed 12 h after heat-shock, therefore we focused our subsequent

analysis on the data set obtained at the 48 h timepoint. The ChIP-seq profiles and RNA-seq

data of two *Fer2* targets, *CG8128* and *skd*, are presented as examples in Fig. 3d. Because

Fer2 expression is largely restricted to PAM neurons, the majority of FER2 ChIP signal

represents FER2 binding within PAM neurons, whereas whole-head RNA-seq following

Fer2 induction detects genes also expressed outside the PAM cluster. Therefore, to determine

whether the *Fer2* targets identified above are expressed in PAM neurons, we profiled the

transcriptome of isolated PAM neurons. Indeed, the majority (26 out of 32) of *Fer2* target

genes identified by the ChIP-seq/head RNA-seq intersectional analysis were found to be

expressed in PAM neurons (Fig. 4a-b), validating this subset of 26 genes as *bona fide* direct

transcriptional targets of *Fer2* (Fig. 4c, Supplementary Table 3).

A further analysis of the 26 *bona fide* targets revealed that *Fer2* preferentially binds to

introns of the target genes and acts as a repressor in most instances, as the majority of direct

targets are downregulated after acute *Fer2* overexpression (Fig. 4d-e). A large fraction of the

26 *Fer2* direct targets are involved in the regulation of gene expression, including

transcription factors (*ftz-f1*, *Eip75B*, *Eip93F*, *Snoo*), transcriptional coregulators (*skd*),

components of chromatin remodeling complexes (*Smr*, *osa*, *sba*) and histone modifiers (*gpp*).

These genes show a significant interconnectivity (p-value = 0,00006) either physically or

functionally, as inferred from STRING database analysis, suggesting that they may cooperate

to regulate downstream gene expression (Fig. 4c).

Multiple *Fer2* direct target genes contribute to the maintenance of PAM dopaminergic

neurons

Next, we assessed whether *Fer2* direct target genes have roles similar to *Fer2* by

manipulating their expression levels individually in PAM neurons and monitoring neuronal

survival. To mimic the effect of *Fer2* loss of function, target genes upregulated by *Fer2* were

knocked-down via RNAi, and genes downregulated by *Fer2* were overexpressed in PAM

neurons. PAM neurons were examined at the age of 35 days with anti-TH staining. 19 out of

26 *Fer2* direct targets were tested in this assay, since transgenic lines for RNAi or

overexpression for 7 genes were unavailable or lethal when combined with the *Fer2-GAL4*

driver. The majority of the transgenic lines used in this assay have been validated in previous

studies. Remarkably, genetic manipulation of one third of the tested genes (6 out of 19)

caused a significant loss of PAM neurons compared with the driver-only control, thus

partially recapitulating the phenotype of *Fer2* loss of function (Fig. 4f and Supplementary

Table 3). These results broadly confirm that the ChIP-seq and RNA-seq intersectional

analysis identified a panel of *Fer2* direct targets that contribute to the role of *Fer2* in the

development and maintenance of PAM neurons.

Fer2 is a master regulator of mitochondrial health and function

Since many *Fer2* direct targets are transcriptional regulators, *Fer2* most likely orchestrates

the expression of a larger set of secondary targets to promote the survival of PAM neurons.

To decipher downstream biological processes controlled by *Fer2*, we profiled the

transcriptome of isolated PAM neurons following constitutive, PAM-specific *Fer2*

overexpression (Fig. 4a). We reasoned that this approach should capture the long-term

consequences of *Fer2* overexpression on gene expression in PAM neurons. A comparison of

PAM neuron transcriptomes of flies overexpressing *Fer2* and age-matched controls identified

132 differentially expressed genes (DEGs) (24 genes upregulated and 108 genes

downregulated; $p < 0.01$). These genes were distinct from the 26 direct targets and were

considered as *Fer2* secondary targets (Fig. 4a-b; Supplementary Table 4).

259   Gene ontology (GO) analysis of *Fer2* secondary targets revealed a marked enrichment

260 of nuclear-encoded mitochondrial genes, particularly among up-regulated genes. These

261 include a component of the electron transport chain (ETC) (*ND-15*), genes involved in the

262 assembly or maturation of ETC complexes (*Sdhaf3, CG7639, Cchl, CG5037*), and the

263 mitochondrial *Superoxide dismutase 2* (*Sod2*) (Fig. 4g and Supplementary Table 4.). As a

264 crucial player in the cellular antioxidant response, SOD2 is localized to the mitochondrial

265 matrix and catalyzes superoxide ($O_2\bullet^-$) to the less reactive H_2O_2[47]. Remarkably, *Sod2*

266 overexpression in PAM neurons partially but significantly prevented neuronal loss in *Fer2*[2]

267 mutant flies, which suggests that *Sod2* acts downstream of *Fer2* to contribute to

268 dopaminergic neuroprotection (Fig. 4h). Combined with the finding that *Fer2* regulates the

269 expression of numerous mitochondrial genes, these results demonstrate that *Fer2* plays a

270 prominent role in the maintenance of mitochondria.

271   To explore how *Fer2* direct targets control a large set of secondary targets, we

272 searched for transcription factor binding sites that are significantly overrepresented in the

273 promoters of DEGs in PAM RNA-seq. Seven binding motifs were found to be enriched,

274 including those for Broad, Dichate and USP, which are expressed in PAM neurons.

275 Interestingly, these genes are known to form complexes or functionally interact with FTZ-F1,

276 a *Fer2* direct target[48,49] (STRING database). Therefore, gene regulatory networks initiated by

277 *Fer2* may be mediated by FTZ-F1 together with these factors (Supplementary Fig. 3a-b). Our

278 finding that *ftz-f1* is necessary for PAM neuron maintenance (Fig. 4f) supports this prediction.

279 Another *Fer2* direct target, SMRTR, can interact with USP[50], and may thus initiate a second

280 pathway to control gene networks in PAM neurons (Supplementary Fig. 3a-b).

282 **Conditional deletion of *Nato3* in differentiated DA neurons leads to locomotor deficits in**

283 **aged mice**

Since *Fer2* is required for the maintenance of mitochondrial health and integrity in DA

neurons, we wondered whether the mammalian homolog of *Fer2* has a similar role. Blast

searches found four mammalian bHLH transcription factors with a significant similarity to

FER2, namely P48/PTF1A, NATO3 (N-TWIST, FERD3L), TCF15, and TWIST1.

Interestingly, it has been shown that *Nato3* is expressed in the progenitors of mDA neurons

and is required for dopaminergic neurogenesis, analogously to the role of *Fer2* in the genesis

of DA neurons in flies[32,33,51]. Furthermore, RNA-seq data from the Genotype-Tissue

Expression (GTEx) project (the GTEx Consortium, Nat Genetics, 2013) indicate that, among

human tissues, *Nato3* expression is relatively restricted to the SN of the adult brain (Fig. 5a).

Based upon these evident similarities in the protein sequence, the expression in mDA neurons

and the role in the development of mDA neurons, we consider *Nato3* to be a functional

homolog of *Fer2*.

To investigate whether *Nato3* is involved in the maintenance of postnatal mDA

neurons, we first determined its expression levels and specificity in the mouse brain. Reverse

Transcription quantitative PCR (RT-qPCR) analysis of dissected brain regions displayed a

significant enrichment of *Nato3* transcripts in the midbrain of 4-month-old mice (Fig. 5b).

Nato3 continues to be expressed at similar levels in the midbrain of young (1-month-old),

adult (4-month-old) and aged (14-month-old) mice (Fig. 5c). Immunohistochemistry analysis

of coronal midbrain sections of 14-month-old mice based on anti-TH and anti-NATO3

antibodies revealed the presence of NATO3 in DA neurons of both the ventral tegmental area

(VTA) and the SN, as well as in non-TH-positive cells (Fig. 5d), consistent with the RT-

qPCR data.

Having verified the expression of *Nato3* in adult mDA neurons, we sought to

conditionally delete *Nato3* in postmitotic DA neurons and investigate its consequences to

dopamine-related behavior and the integrity of mDA neurons. We generated a mouse strain

bearing the *loxP*-flanked (floxed) allele of *Nato3* and crossed it with the *DAT-Cre* mouse

expressing the Cre recombinase under the control of the dopamine transporter gene (*Slc6a3*)

regulatory sequences[52]. *DAT-Cre* expression is known to arise postnatally in differentiated

midbrain DA neurons, has no deleterious effects on DA neurons, and does not alter

locomotion or anxiety-related behaviors[52,53]. We obtained homozygous *Nato3*-floxed animals

harboring either no copies (*Nato3^{floxed}*, referred to as control) or one copy of the *DAT-Cre*

transgene (*Nato3^{DAT-Cre}*, referred to as *Nato3* cKO) (Fig. 6a). RT-qPCR analysis of the ventral

midbrain confirmed the ablation of *Nato3* in *Nato3^{DAT-Cre}* mice (Fig. 6b).

We analyzed control and *Nato3* cKO mice at different ages using assays for

dopamine-related behaviors. In the pole test, which is a sensitive assay for motor impairment

associated with reduced striatal dopamine in rodents[54], *Nato3* cKO mice took significantly

longer to turn and descend the pole than control mice at 13 months, while no difference was

observed in younger animals (Fig. 6d). This observation indicates an age-dependent decline

in motor coordination in *Nato3* cKO mice. Spontaneous activity levels measured via

voluntary wheel running over three days were similar between *Nato3* cKO and control mice

until 13 months of age. However, the activity levels of *Nato3* cKO significantly declined at

16 months (Fig. 6e). In contrast, no differences were found between *Nato3* cKO and control

mice in the open field test, which measures exploratory behavior when mice are placed in a

novel environment[55] (Supplementary Fig. 4). These results indicate that *Nato3* ablation in

postmitotic DA neurons lead to motor abnormalities in aged mice, possibly caused by an age-

dependent decline in the nigrostriatal dopaminergic function.

***Nato3* is required for the maintenance of mitochondrial integrity in midbrain DA**

neurons

To broadly assess the integrity of the nigrostriatal pathway, ventral midbrain and striatal

sections of 16- to 18-month-old mice were analyzed by anti-TH immunohistochemistry. The

number of DA neuron cell bodies in the SN and VTA (Fig. 6f-g), as well as the density of

TH$^+$ fibers in the striatum (Fig. 6h-i), did not show significant differences between control

and *Nato3* cKO mice. Therefore, locomotor deficits in the *Nato3* cKO mice likely originate

from functional impairments of DA neurons rather than an overt loss of cellular integrity.

Fer2 mutation impairs mitochondrial morphology selectively of DA neurons in the

PAM cluster, which precedes the degeneration of PAM neurons and locomotor deficits in

Drosophila[28]. To determine whether *Nato3* ablation also leads to mitochondrial abnormalities

in DA neurons in mice, we first visualized mitochondria in the SN using the antibodies

against mitochondrial cytochrome c oxidase subunit 4 (COX4). The mitochondria appeared

more globular in DA neurons of the SN in old (16-18 months) *Nato3* cKO mice compared

with age-matched controls (Fig. 7a). Quantification of mitochondrial circularity showed a

statistically significant increase (i.e. an accumulation of rounder mitochondria) in *Nato3*-

ablated DA neurons, confirming the visual impression (Fig. 7b). We further examined the

ultrastructure and morphology of mitochondria in the SN using transmission electron

microscopy. In TH-labeled neurons from control mice, most mitochondria (88%) appeared

normal, showing a dense and uniform matrix and regularly distributed cristae (Fig. 7c, top

panels, and Fig. 7d). Strikingly, a large fraction (46%) of mitochondria from *Nato3* cKO

mice displayed an aberrant morphology, showing a reduced and patchy matrix density with

few remaining cristae; in certain cases these mitochondria contained almost no cristae,

presented white inclusions and appeared dysmorphic (Fig. 7c, bottom panels, and Fig. 7d).

Mitochondria from TH-negative cells appeared normal in *Nato3* cKO mice, suggesting that

the aberrant mitochondrial morphology was cell-autonomously induced by *Nato3* deletion in

DA neurons (Fig. 7c, right bottom panel). Although mitochondrial size did not differ between

the two genotypes, mitochondria from SN neurons in *Nato3* cKO mice were rounder than

those in the control (Fig. 7e). These results demonstrate that *Nato3* is required for the

maintenance of mitochondrial integrity in the DA neurons of the SN.

DISCUSSION

Our results demonstrate that *Fer2* is a master regulator of a transcriptional network that

maintains mitochondrial integrity in DA neurons in *Drosophila*. This property renders *Fer2* a

potent dopaminergic neuroprotective factor against genetic and oxidative insults implicated

in PD pathogenesis. Furthermore, we show that *Nato3*, a mammalian homolog of *Fer2*, is

required for the maintenance of mitochondrial integrity in mDA neurons. Both *Fer2* and

Nato3 are developmental transcription factors for DA neurons that continue to be expressed

throughout life, and their loss of function in differentiated DA neurons leads to locomotor

deficits in aged animals. Our findings corroborate the critical role of developmental

transcription factors in the lifelong maintenance of DA neurons and suggest the evolutionary

conservation of transcriptional mechanisms for dopaminergic neuroprotection and

mitochondrial integrity.

Drosophila Fer2 (*48 related 2*, CG5952) and its paralogs *Fer1* and *Fer3* were

originally named for their similarity to the p48 subunit of the pancreas transcription factor 1

complex (PTF1) in the bHLH domain by the group of Yuh Nung Jan[56]. Independently, due to

the similarity to mammalian Atonal within the bHLH domain, *Fer3* (CG6913) was named

Drosophila Nephew of Atonal 3 (*Dnato3*) by the group of Nissim Ben-Arie[57]. Ben-Arie's

group also isolated a mouse homolog of *Dnato3* and named it *MNato3*[57]. Brian L Black's lab

also independently isolated the same mouse gene and named it *Neuronal-twist* (*N-twist*), due

to its high sequence similarly to *M-twist*[58]. It is puzzling that, whereas Jan's group has

reported the expression of *Fer2* mRNA in the brain and ventral nerve cord (VNC) of the

embryo and *Fer3* expression in the primordial midgut of the embryo[56], Ben-Arie's and

Black's groups have shown the expression of *Fer3* mRNA in the CNS of the embryo, which

supports the idea that *Fer3* is the homolog of *Nato3*[58]. Notwithstanding this inconsistency, we

argue that *Fer2* is a functional homolog of *Nato3* based on the following reasons. The fact

that fewer DA neurons are generated in *Fer2* loss-of-function mutants[28] is strikingly similar

to the observation that 30% fewer mDA neurons are formed in *Nato3* constitutive KO[32],

suggesting that *Fer2* bears a proneural function similar to that of *Nato3*. FER2 bHLH domain

shows a high similarity to that of NATO3 as well as to FER1 and FER3. Moreover, we have

demonstrated here and in previous work[28] that DA neurons are the primary site of *Fer2*

expression and function, and that both *Fer2* and *Nato3* are required for mitochondrial

maintenance in differentiated DA neurons. Nevertheless, the possibility that *Fer3* may also

have similar roles as *Fer2* in the development and maintenance of DA neurons cannot be

excluded and is worthy of further investigation.

This study is the first to report the function of *Nato3* in differentiated mDA neurons in

mice. *Nato3* is classically known as a neurogenic transcription factor, expressed in the floor

plate cells in the developing mesencephalon and spinal cord[32,34]. Although *Nato3* expression

in the spinal cord of postnatal and adult mice has been reported[59], we found here its presence

in the mDA neurons of adult mice. By conditionally ablating *Nato3* in postmitotic DA

neurons, we were able to distinguish the function of *Nato3* in mature mDA neurons, i.e. the

maintenance of mitochondrial integrity, from its role in neurogenesis. Importantly, human

Nato3 is expressed almost exclusively in the SN among adult tissues. Thus, it is plausible that

human *Nato3* also plays an important role in the mitochondrial maintenance in SN

dopaminergic neurons.

The genesis, specification and differentiation of mDA neurons are governed by a

complex web of transcription factors[60]. Whereas *Nato3* acts as an "early transcription factor"

required for neurogenesis[32], it can also directly regulate *Lmx1b*, *Shh* and *Foxa2* and thereby

indirectly regulates the late transcription factor *Nurr1*[33,51]. Since *Lmx1b*, *Foxa2* and *Nurr1* are

neuroprotective transcription factors[9,12,13,19], *Nato3* might contribute to dopaminergic

neuroprotection by regulating these genes. However, none of the homologs of developmental

transcription factors for mDA neurons were found among *Fer2* downstream targets in adult

PAM neurons. Therefore, the gene regulatory networks for mDA neuron development and

adult mDA maintenance may be substantially different.

Identification of *Nato3* transcriptional targets and downstream molecular events in

differentiated mDA neurons will provide crucial information to understand the role of *Nato3*

in mitochondrial maintenance. While further studies are required to obtain such data, here we

have performed homologous experiments in *Drosophila*, i.e. identification of direct and

indirect targets of *Fer2* in DA neurons in the PAM cluster. An intersectional approach

combining ChIP-seq, whole-head RNA-seq and RNA-seq of isolated PAM neurons,

unequivocally identified 26 direct and over 100 indirect *Fer2* target genes expressed in PAM

neurons. Among *Fer2* indirect targets, genes involved in mitochondrial function and structure,

including those required for ETC complexes assembly, are remarkably enriched. Malfunction

of the ETC leads to ROS overproduction, which in turn induces mitochondrial DNA

mutations, alters membrane permeability and mitochondria morphology, and further damages

the ETC complexes, in a pathogenic self-reinforcing loop. Indeed, dysfunctions of the ETC,

particularly of Complex I, have been proposed as a leading mechanism of DA

neurodegeneration[61,62]. Thus, our molecular data are in accordance with the observation that

Fer2² flies display an overt defect in mitochondrial networks[28]. It is noteworthy that *Sod2*,

which encodes mitochondrial superoxide dismutase, is also an indirect target of *Fer2*.

Consistent with this finding, *Sod2* overexpression partially rescues the loss of PAM neurons

in *Fer2²* mutants. Furthermore, *Fer2* overexpression reverses mitochondrial abnormalities in

the PAM neurons of *park¹* mutants and prevents PAM neuron loss in genetic and toxin

434 models of PD. These results are in agreement with previous reports that *Sod2* and *Sod2*-

435 mimetic agents are protective in cell and fly models of PD[63], and endorse the further

436 exploration of *Sod2* as a potential therapeutic target for PD.

437  Among *Fer2* direct targets, we discovered six genes whose dysregulation induces a

438 loss of PAM neurons. None of these genes have been previously related to the DA system or

439 implicated in neuronal development or survival. Further studies are needed to clarify how

440 these novel actors and their mammalian homologues contribute to the biology of DA neurons.

441 The number of new genes implicated in PAM neuron survival may increase in the future, as

442 our list of *Fer2* direct targets is likely not exhaustive. *Fer2* is a stress-responsive transcription

443 factor that is upregulated by oxidative insult or possibly by other stressors[28]. Under such

444 conditions FER2 may be recruited to and/or regulate different sets of target genes.

445  A large fraction of *Fer2* primary targets encode transcription factors or chromatin

446 proteins. This positions *Fer2* at the vertex of multiple transcriptional cascades and suggests

447 that *Fer2* orchestrates the expression of a broad set of secondary targets (Supplementary Fig.

448 3b). *Fer2* expression is restricted to PAM neurons and a few other cell clusters in the brain,

449 and ubiquitous or even pan-neuronal constitutive overexpression of *Fer2* causes lethality

450 (data not shown), indicating that genetic programs driven by *Fer2* are highly specific to PAM

451 neurons.

452  PD progresses from the asymptomatic (preclinical) period to a stage with a variety of

453 nonmotor symptoms and subtle motor impairments that do not meet diagnostic criteria

454 (prodromal PD), before a definite diagnosis is reached[64]. Modeling in animals this latent

455 phase of the disease would facilitate the development of disease-modifying therapies and the

456 discovery of early PD biomarkers. We propose that *Nato3* cKO mice can serve as a relevant

457 model for prodromal PD, presenting moderate motor deficits and quasi-normal mDA neurons

458 that accumulate abnormal mitochondria. There is growing evidence that neurodegeneration in

PD is manifested after several risk factors, including genetic predisposition, toxin exposure

and aging, interact to overcome protective compensatory mechanisms[65]. Future work is

needed to determine whether exposing *Nato3* cKO mice to a second genetic or environmental

insult exacerbates its PD-related phenotypes and trigger overt DA neurodegeneration.

The activation of neuroprotective transcription factors offers potent therapeutic

opportunities, allowing to modulate multiple downstream pathways in parallel. Indeed, *Nurr1*

and *Foxa2* gene delivery[19], *Nurr1* activators[66] and *En1* infusion[20] in rodent PD models have

shown encouraging results. In light of our findings that *Fer2* confers neuroprotection in

multiple fly PD models, it is tempting to speculate that the overexpression or activation of

Nato3 may protect mDA neurons from genetic and environmental insults in mammals, with

potential far-reaching implications for PD treatment.

METHODS

***Drosophila* stocks and maintenance**

The following lines were obtained from the Bloomington *Drosophila* Stock Center (BDSC)

(Indiana University, Bloomington, IN): *UAS-RedStinger* (#8546), *UAS-Sod2* (#24494), *park^1*

(#34747), and *hs-GAL4* (#1799). *Fer2::GFP::V5, PBac{RB}Fer2^{e03248}* (referred to as *Fer2^1*),

Mi{ET1}Fer2^{MB09480} (referred to as *Fer2^2*), *Fer2-GAL4, R58E02-GAL4, UAS-mitoGFP*, and

UAS-Fer2::FLAG lines were previously described[26,28,67]. *UAS-Lrrk WT* and *UAS-Lrrk*

I1915T were a kind gift from Yuzuru Imai[68]. All of the lines used to overexpress or knock-

down *Fer2* primary targets are described in the Supplementary Table 3. All flies were

maintained on standard cornmeal-agar food at 25 °C in a 12 h:12 h light-dark cycle and under

controlled humidity.

Fly brain immunohistochemistry

Immunostaining of whole fly brains was performed as previously described[28]. Briefly, the

flies were decapitated, and the heads were fixed in 4% paraformaldehyde (PFA) + 0.3%

Triton X-100 for 1 h on ice and washed three times with PBST-0.3 (PBS, 0.3% Triton X-100).

The cuticle was partially removed, and the partly opened heads were incubated with blocking

solution (5% normal goat serum, PBS, 0.3% Triton X-100) for 1 h at room temperature (RT)

on a rocking platform. Subsequently, the samples were incubated with primary antibodies for

48 h at 4 °C. After three 20 min washes with PBST-0.3, the samples were then incubated

with Alexa Fluor-conjugated secondary antibodies overnight at 4 °C. After three 20 min

washes with PBST-0.3, the remaining cuticle and trachea were removed and the brains were

mounted in Vectashield medium (Vector Laboratories). Primary antibodies and the dilution

factors were as follows: rabbit polyclonal anti-TH (ab152, Millipore) 1:100, polyclonal rabbit

anti-GFP (A6455, Invitrogen) 1:250. Neuropil was counterstained with the mouse

monoclonal antibody nc82 (Developmental Studies Hybridoma Bank) 1:100 to visualize the

entire brain and assess its integrity. Secondary antibodies and the dilution factors were as

follows: goat anti-rabbit IgG Alexa Fluor 488 conjugate (A-11034, Thermo Fisher Scientific)

1:250, goat anti-mouse IgG Alexa Fluor 633 conjugate (A-21052, Thermo Fisher Scientific)

1:250.

Mouse brain immunohistochemistry

Animals were anesthetized and perfused through the left ventricle with 0,2% Heparin in PBS

at RT followed by ice-cold 10% formalin solution (Sigma HT501128-4L). The brains were

dissected and post-fixed for 72 h in fresh 4% PFA (Thermo Fisher Scientific #28906) before

being embedded in low melting point agarose (Thermo Fisher Scientific #R0801) and

sectioned in a sliding vibratome at a thickness of 40 μm. Striatum sections were stained as

floating sections. Ventral midbrain sections were adhered to the SuperFrost plus slides

(Thermo Fisher Scientific J1800AMNZ) before staining. Slices were incubated in blocking

buffer (4% non-fat milk, 4% Bovine Serum Albumin (BSA), PBS) containing 0,1% Triton X-

100 for 30 min at RT. The sections were then incubated with primary antibodies in blocking

buffer containing 0,1% Tween for 48 h at 4 °C. After three 30 min washes in washing buffer

(0.1% Tween, PBS), sections were incubated with the corresponding secondary antibody

overnight at 4 °C, in the dark. After three 30 min washes in washing buffer, the slices were

mounted with DABCO-glycerol mounting medium (25 mg/ml DABCO (Sigma D27802-

25G), 90% glycerol, PBS; pH 8,6). For nuclear staining, prior to mounting midbrain slices

were treated with 100μg/ml RNAse A (#EN0531, Thermo Fisher Scientific) for 30 min at

37 °C, and then incubated with TO-PRO-3 iodide (T3605, Sigma) for 30 min at RT. Primary

antibodies and dilution factors were as follows: rabbit polyclonal anti-TH (Millipore ab152)

1:300, mouse anti-TH (Immunostar 22941) 1:300, rb polyclonal anti-NATO3 (Aviva

ARP37423_P050) 1:70, mouse anti-COX4 (Thermo Fisher Scientific 1D6E1A8) 1:300.

Secondary antibodies and dilution factors were the same as those for the fly brain

immunostaining.

Microscopy and image analysis

Fly brains and mouse brain slices were scanned using a Leica TCS SP5 confocal microscope.

Image analysis was performed using Fiji software[69]. All analyses were conducted blindly

with respect to genotype.

Fly brain. TH-positive neurons were counted manually from Z-stacks of confocal

images using the cell counter plugin of Fiji. PAM neuron counts were scored blindly.

FER2::V5::GFP and TH fluorescence intensity in the PAM region was measured on Z-SUM

projection images. The region of interest (ROI) corresponding to the PAM cluster was

manually drawn to include TH$^+$ neurons. Background signal measured outside the ROI was

subtracted from the ROI signal. To quantify mitochondria circularity in PAM clusters,

mitoGFP-positive objects within the PAM neurons region were analyzed using a particle

analysis plugin as described previously[29].

Mouse brain. TH immunoreactivity in striatal regions was quantified by optical

densitometry using Fiji. Three striatum images were acquired for each animal, and in every

picture fluorescence intensity was measured for four squares drawn on the dorsal striatum

(n=3-4 animals per genotype). Measures from all the squares were averaged to give the value

for each animal. To obtain the number of mDA neurons, TH-positive neurons were counted

manually from Z-stacks of confocal images using the cell counter plugin (n=3 animals per

genotype). Every fourth section covering the entire VTA and SN in one brain hemisphere

was included in the counting procedure. Neuron counts were scored blindly. To quantify

mitochondria circularity in DA neurons, COX4-positive objects within the TH-positive DA

neurons were analysed using the particle analysis plugin.

Treatment of flies with oxidative reagents

7-day-old flies were placed into empty vials and starved for 6 h. The flies were then

transferred into vials containing *Drosophila* instant food (Formula 4-24 Instant Drosophila

Medium, Carolina Biological Supply Company, USA) containing 5% H_2O_2 (Sigma, H1009),

or vials containing *Drosophila* instant food only as a control, and kept for 24 h. All vials were

kept in a humid box at 25 °C during the treatment. Following the treatment, flies were

transferred back to the vials with standard cornmeal-agar food and maintained normally.

Seven days after the treatment the flies were decapitated, and anti-TH-staining was

performed as described above.

ROS detection

The level of ROS production in the fly brain was measured using a cell-permeable probe,

2'7'-dichlorofluorescin diacetate (H2DCF), following the protocol described in[70]. The

fluorescence intensity for the entire central brain was measured using Fiji software[69] after

generating a Z-SUM projection across the corresponding Z-stacks, as previously described[28].

Startle-induced climbing assay

For each experimental group, 20 male flies were briefly anesthetized with CO_2 and placed

into a 100 ml graduated glass cylinder, which was divided into five equal parts marked 1-5

from bottom to top. Flies were left to recover from the CO_2 exposure for 1 h before the assay.

The climbing assay was performed at ZT2 (2 h after lights-on in 12 h:12 h light-dark cycles)

to avoid circadian variations in the locomotor activities of the flies, following the procedure

previously described[28]. Briefly, the cylinder was tapped on a flexible pad to collect all of the

flies at the bottom, and then the flies were allowed to climb up the wall of the cylinder for 20

s. The number of flies that climbed up to each zone within 20 s was counted manually and

used to calculate the climbing index (CI) with the following formula: $CI = (0 \times n_0 + 1 \times n_1 +$

$2 \times n_2 + 3 \times n_3 + 4 \times n_4 + 5 \times n_5) / n_{total}$, where n_x is the number of flies that reached zone X,

n_0 is the number of flies that remained in the bottom of the cylinder, and n_{total} is the total

number of flies. Each experiment consisted of three trials, and three independent experiments

were performed for each genotype. The mean CI values from three independent experiments

are shown.

Chromatin immunoprecipitation coupled to sequencing (ChIP-seq)

ChIP was performed based on previous protocols[71,72] on *Fer2[1]* mutant flies expressing a

GFP::V5-tagged *Fer2* genomic transgene (*Fer2::GFP::V5*) and on *w[1118]* flies (negative

control). Briefly, ~8000 14-day-old flies per sample were collected 2 h before lights-on

(ZT22), frozen in liquid nitrogen and maintained at -80 °C until processed. Heads were

collected using metal sieves, and then, 1 ml of heads was ground in liquid nitrogen and

homogenized 30 times in 5 ml of NE Buffer (15 mM HEPES pH 8, 10 mM KCl, 0.1 mM

EDTA, 0.5 mM EGTA, 350 mM sucrose, 0.1% Tween 20, 5 mM MgCl2, 1 mM DTT, 1 mM

PMSF plus Protease inhibitor cocktail (Roche 11 873 580 001)) with 1% PFA. Fixation was

performed for 10 min at RT and quenched with 125 mM glycine for 5 min at RT. The

homogenate was filtered through a 100 µm nylon mesh filter (Thermo Fisher Scientific

22363549), and nuclei were collected by centrifugation at 800 x g for 5 min at 4 °C. The

nuclei were washed three times, resuspended in 1,5 ml of cold RIPA buffer (25 mM Tris-HCl

pH 8, 150 mM NaCl, 0,5% sodium deoxycholate, 0,1% SDS, 1% NP-40, 0,5 mM DTT plus

Protease inhibitor cocktail) and sonicated using a Bioruptor sonicator for 18 min (30 s ON/30

s OFF). The sonicated chromatin was then centrifuged at 10000 x g for 10 min at 4 °C to

remove cell debris. The supernatant was pre-cleared with agarose beads (1 h, 4 °C) and

incubated with 50 µl of saturated anti-V5 beads (Sigma A7345) overnight at 4 °C. The beads

were washed twice with low-salt buffer (20 mM Tris pH 8, 150 mM NaCl, 2 mM EDTA,

0,1% SDS, 1% Triton X-100), twice with high-salt buffer (20 mM Tris pH 8, 500 mM NaCl,

2 mM EDTA, 0,1% SDS, 1% Triton X-100), twice with LiCl buffer (10 mM Tris pH 8, 250

mM LiCl, 1 mM EDTA, 1% sodium deoxycholate, 1% NP-40) and twice with TE buffer (10

mM Tris-HCl pH 8, 1 mM EDTA). DNA was eluted in the V5 elution buffer (10 mM HEPES,

1,5 mM MgCl2, 0,25 mM EDTA, 20% glycerol, 250 mM KCl, 0,3% NP-40, 0,5 mg/ml V5

peptide) and incubated with RNase (30 min, 37 °C) and proteinase K (overnight, 65 °C).

Subsequently, the DNA was purified with a MinElute PCR purification kit (Qiagen 28004).

Libraries were prepared with 0,5 to 0,7 ng of ChIP-enriched DNA as starting material and

processed using the Illumina TruSeq ChIP kit according to the manufacturer's specifications,

at the iGE3 Genomics Platform in Geneva. The libraries were validated on a Tapestation

2200 (Agilent) and a Qubit fluorometer (Invitrogen - Thermo Fisher Scientific). Single reads

of 50 bp were generated using the TruSeq SBS HS v3 chemistry on an Illumina HiSeq 2500

sequencer.

ChIP-seq analysis

Sequencing reads were mapped to the *Drosophila melanogaster* genome reference dm3 using

Bowtie2 with default settings[73]. *Fer2* binding sites were identified by individually comparing

each FER2 ChIP sample (N=2) to each w^{1118} ChIP sample (N=2; negative control) using

MACS2[74]. The minimum FDR (q-value) cutoff for peak detection was 0.05. The 267 shared

peaks overlapping in all the four FER2 ChIP vs w^{1118} ChIP comparisons were selected as

confident peaks. Peak calling was also performed using Homer[75] and over 90% of Homer

peaks overlapped with MACS2 peaks, confirming our results. ChIP-seq profiles were

visualized using IGV[76]. HOMER motif analysis algorithm was used to search for enriched

motifs, and identified a 12-mer containing a bHLH binding site, present in the majority of the

best ranked peaks. Peak-to-gene assignment was performed using the HTSstation[77] and

resulted in a list of 279 genes.

Whole-head RNA-seq

hs-GAL4, *UAS-Fer2::FLAG* flies (*hs>Fer2::FLAG*) were heat-shocked at 37 °C for 3 h, or

maintained at 25 °C (control), and collected 12 h or 48 h later (three independent biological

replicates per group). The same procedure was also applied to flies carrying *hs-GAL4* but not

the *UAS-Fer2,* in order to identify and exclude from our analysis the genes that responded to

heat irrespective of *Fer2* induction. The flies were frozen in liquid nitrogen and kept at -

80 °C until processed. Fly heads were collected using metal sieves and kept in Trizol (Life

Technologies 15596026) at -80 °C until processed. Total RNA was isolated from dissected

mouse brain regions using Trizol according to the manufacturer's instructions and treated

with Turbo DNase for 30 min at 37 °C (Turbo DNA-free kit, Thermo Fisher Scientific

AM1907). Libraries were prepared at the iGE3 Genomics Platform in Geneva. Single reads

of 50 bp were generated using an Illumina HiSeq 2500 sequencer. Data analysis was

performed using Galaxy[78]. Sequencing reads were mapped to the *Drosophila melanogaster*

genome reference dm3 using TopHat[79] with default settings. Differentially expressed genes

(DEG) analysis was performed with Cufflinks, Cuffmerge and Cuffdiff, to compare flies who

received the heat shock treatment with control flies kept at 25 °C. The same analysis was

performed on *hs-GAL4* flies, to identify the genes which responded to heat irrespective of

Fer2 induction. These heat-responsive genes were subtracted from DEG identified in the

hs>Fer2::FLAG experiment.

PAM-specific RNA-seq

Flies carrying *R58-GAL4* were combined with *UAS-Fer2::FLAG* to overexpress *Fer2*, and

with *UAS-RedStinger* to mark PAM neurons. Flies carrying only *R58-GAL4* and *UAS-*

RedStinger were used as a control. Cells were dissociated from the brains of ~70 14-day-old

flies 2 h before lights-on (ZT22) as previously described[67], except that the brains were treated

with papain for 25 min. After trituration in 300 μl SMactive medium containing 50 μm AP5,

20 μm DNQX and 0.1 μm TTX, the cell suspension was filtered through a sterile 70 μm

membrane and distributed to three wells of three two-well chamber glass slides (LabTek II,

Nunc) kept on ice. The second well of each slide contained ice to cool the cells while

RedStinger-positive cells were sorted on a fluorescence microscope equipped with a

motorized stage and a needle holder (CellSorter Company for Innovations). Cells were

selected manually using capillaries with 50 μm tip diameter. Collected cells were released in

a clean part of the chamber and immediately picked up again, greatly reducing the amount of

debris, before being transferred to a fresh slide with 100 µl medium spread in a 1 × 1 cm

GeneFrame (Thermo Fisher Scientific). Cells were reselected for two rounds and finally

transferred using a new capillary with 30-µm tip diameter to 100-µl lysis buffer (Absolutely

RNA Nanoprep Kit; Agilent). Cells of multiple collections were pooled if needed to obtain

200 cells per sample, and total RNA was extracted according to the manufacturer's

instructions. The RNA was converted to cDNA and amplified using the SMART-Seq4 Kit

(Takara Bio). Libraries were generated using NexteraXT (Illumina) and sequenced on an

Illumina HiSeq 2500 or 4000. DEG analysis was performed using the HTSstation[77] and

resulted in a list of 132 genes.

Bioinformatic analysis of genes found in ChIP-seq and RNA-seq

Since human databases are generally more comprehensive, fly genes were converted into

human orthologs prior to Gene Ontology (GO) analysis[80]. Human orthologs of fly genes were

predicted using DIOPT[81] and were analyzed for GO with Metascape[80]. A network

representation of GO terms was obtained using Cytoscape[82]. A search for TF binding sites

enriched among promoters of PAM-specific RNAseq DEG was performed using PSCAN[83].

Protein-protein interactions were investigated using STRING[84] and a previously reported

Drosophila TFs database [48].

Mice strains

Nato3-LoxP mice was generated using Clustered Regularly Interspaced Short Palindromic

Repeats (CRISPR) technology by Applied StemCell, Inc. (Milpitas, CA). Two copies of a

synthetic *LoxP* sequence were inserted upstream and downstream of the unique exon of

Nato3 gene. Briefly, a mixture of active guide RNA molecules (gRNAs), two single stranded

oligo donor nucleotides (ssODNs) and *Cas-9* mRNA was prepared and injected into the

cytoplasm of C57BL/6 embryos. Mice born from the microinjection were screened to identify

LoxP insertions at the designated locations using PCR and sequencing. The founder mice

were bred to evaluate the germline transmission and expand the *Nato3-LoxP* colonies. *DAT-*

Cre mice have been described previously[52] (obtained from Jean-Claude Martinou's

laboratory, University of Geneva). *DAT-Cre* and *Nato3-LoxP* mice were maintained in a

C57BL/6J genetic background. Crosses between these lines generated mice that are

homozygous for the *Nato3-LoxP* allele and harbor one copy of the *DAT-Cre* transgene

(*Nato3$^{DAT-Cre}$*). Littermates homozygous for *Nato3* floxed alleles and harboring no copy of the

DAT-Cre transgene were used as controls (*Nato3^{floxed}*). Mice were maintained in rooms with

controlled 12 h light/dark cycles, temperature and humidity, with food and water provided *ad*

libitum. Mice were housed at a maximum of five animals per cage in individually ventilated

cages. All experiments were conducted in accordance with the Institutional Animal Care and

Use Committee of the University of Geneva and with permission of the cantonal authorities

(Permit No. GE/3/19 27381 and GE/78/19 31317).

Electron microscopy image acquisition and analysis

Animals were anesthetized, perfused and embedded in agarose as described in the mouse

brain immunohistochemistry section, except that the perfusion was performed with 3% PFA

complemented with 0.2% glutaraldehyde in 0.1 M phosphate buffer (PB), pH 7.4.

Subsequently, 60 μm-thick sections were cut and processed for pre-embedding

immunoperoxidase staining, as previously described [85]. Briefly, immunostaining for DA

neurons in the SN was performed by incubating the sections with rabbit anti-TH antibodies

(Millipore ab152, 1:100) followed by goat anti-rabbit biotinylated secondary antibodies

(Vector Laboratories PK-4001, 1:200) and then avidin biotin peroxidase complex (Vector

Laboratories PK-4001). 3,3'-diaminobenzidine tetrachloride (DAB; Invitrogen Life

Technologies 750118) was applied for 1-2 minutes until the stain developed. Sections were

further embedded in EPON resin (Fluka) and trimmed around the labeled cells of interest.

Ultrathin (60 nm) sections were produced and imaged by a Tecnai G212 electron microscope

(FEI Company, Oregon, United states) equipped with a digital camera (Mega View III; Soft

Imaging Systems). ~15 labeled cells from two mice were analyzed for each genotype.

Mitochondria were classified blindly. Mitochondrial size and roundness were measured via

Fiji software using the particle analysis plugin after a manual selection of each mitochondrion.

RT-qPCR on mouse brain regions

Animals were sacrificed by cervical dislocation and decapitated. The brains were

immediately removed and all the subsequent steps were performed on ice. The olfactory

bulbs, prefrontal cortex, striatum, cortex, and cerebellum were scalpel-dissected. The ventral

and dorsal midbrain were scalpel-dissected from 1 mm-thick brain slices cut in an ice-cold

coronal matrix (Zivic Instruments). Samples were kept in Trizol (Life Technologies

15596026) at -80 °C until processed. Total RNA was isolated from dissected mouse brain

regions using Trizol according to the manufacturer's instructions and treated with Turbo

DNase for 30 min at 37 °C (Turbo DNA-free kit, Thermo Fisher Scientific AM1907).

Reverse transcription was performed from 500 to 1000 ng RNA using the Maxima reverse

transcriptase (Thermo Fisher Scientific 18090050) and oligo-dT primers, according to the

manufacturer's instructions. Primers were designed with Primer-BLAST online tool

(https://www.ncbi.nlm.nih.gov/tools/primer-blast/): *B2m* (F: 5'-

AGAATGGGAAGCCGAACATA, R: 5'-CGTTCTTCAGCATTTGGATT), *Cyclophilin*

(F: 5'-TGCCATCCAGCCAGGAGGTC, R: 5'-CCATCGTGTCATCAAGGACTT), *Hmbs*

(F: 5'-GCTGAAAGGGCTTTTCTGAG, R: 5'-TGCCCATCTTTCATCACTGT), *Hprt1* (F:

5'-TGTCAGTTGCTGCGTCCCCAGA, R: 5'-TCTACCAGAGGGTAGGCTGGCC), *Pgk1*

(F: 5'-ACCTGCTGGCTGGATGGGCT, R: 5'-CTCGACCCACAGCCTCGGCA), *Rpl13a*

(F: 5'-CGGATGGTGGTCCCTGCTG, R: 5'-GAGTGGCTGTCACTGCCTGG), *Nato3* (F:

5'-GGTGAGCCCAAGAGACACTC, R: 5'-GTGCCGGATCCCCAACTTAT). Quantitative

PCR (qPCR) was performed on the Light Cycler LC480 (Roche) using the TB Green Ex

Taq II polymerase (Takara RR820L). Each reaction was performed in triplicate. Relative

quantification and normalization were performed with a multiple reference genes model[86].

For normalization, the best reference genes for each experiment were determined using the

geNorm algorithm (https://genorm.cmgg.be) within a panel of 4-5 putative reference genes.

To compare *Nato3* levels between different brain regions, *B2m* and *cyclophilin* were used

as reference genes. To compare *Nato3* levels in the midbrain between different ages, *B2m*

and *Hprt1* were used.

Behavioral tests

Male mice of different ages (7, 10, 13 and 16 months) were evaluated for posture control and

coordination (pole test), spontaneous locomotor activity (wheel test) and exploration-driven

locomotor behavior (open field test). All behavioral tests were performed during the light

cycle by experimenters blind to the genotype. The mice were allowed to habituate to the

behavioral room for at least 45 min before each test. Behavioral equipment was cleaned with

70% ethanol after each test session to avoid olfactory cues.

753   *Pole test.* The pole test was performed as previously described[54] with minor

modifications. Mice were first trained 4-6 times by placing the animal head-down on top of a

vertical pole (diameter: 1 cm, height: 55 cm) and letting it descend. Then, animals were

trained 3-4 times in the regular turning and descending procedure, by placing the animal

head-up on top of the pole. For the actual test, mice performed eight trials and were recorded

with a video camera. The time to orient downward (T-turn) and the total time to turn and

descend the pole (T-tot) were measured, with a maximum duration of 30 s. When the mouse

was not able to turn downward and instead dropped from the pole in a lateral body position, a

value of 30 s was assigned. The average of the eight trials was used as the final score. During

the training and for the real test, mice were tested a maximum of three times per day, with an

interval of at least 5 min between trials.

Running wheel test. Mice were singly housed in cages with free access to a running

wheel (14 cm in diameter) for 6 days, under a 12 h light/dark cycles. After three days of

habituation, the number of wheel revolutions per minute was recorded with a sensor by

LabVIEW program for the following three days. The total amount of time spent on the wheel

and the total distance run were calculated for each animal.

Open field test. Open field test was performed as previously described[13] with minor

modifications. The test was performed in a 40 x 40 x 40 cm arena made of white Plexiglas

plastic, which was illuminated from above (light intensity of 100 lux on the floor). Mice were

placed individually in the center of the area and allowed to freely explore it for 10 min while

being recorded. The horizontal locomotor activity was analyzed using the Mouse Behavioral

Analysis Toolbox (MouBeAT)[87] for ImageJ. The vertical activity (rearing), defined as a

mouse rising on its hind legs, was scored manually.

Statistical analysis

Statistical comparisons were performed using GraphPad Prism software (San Diego, CA).

The statistical significance of differences between two groups was calculated using a

nonparametric two-tailed Mann-Whitney test. To compare three or more groups, ANOVA

followed by Turkey's multiple comparisons test was used. For the circularity analysis, to

compare the distribution of mitochondria particles between two or three genotypes, the

Kolmogorov-Smirnov test and the Kruskal-Wallis test followed by Dunn's test were applied,

respectively. Differences were considered significant at *$p<0.05$, **$p<0.01$, ***$p<0.001$,

****$p<0.0001$.

ACKNOWLEDGMENTS

We thank Ernst Hafen, Chi-Hon Lee, Junhai Han, the Bloomington *Drosophila* Stock Center,

the Vienna Drosophila Resource Center and the Kyoto Stock Center for fly stocks, J-C.

Martinou for the *DAT-Cre* mouse strain. We thank the Genomics Platform of iGE3 for ChIP-

seq and RNA-seq experiments, the animal facility of the University of Geneva for

maintenance of mice colonies, and the Pole Facultaire de Microscopie Electronique (Centre

Médical Universitaire, Geneva) for electron microscopy. We are grateful to J-C. Martinou

and our lab members for valuable discussions of this work. This research was supported by

the European Research Council (ERC-StG-311194), the Swiss National Science Foundation

(31003A_130387), the Georges and Antoine Claraz Foundation, the Novartis Foundation for

Medical-Biomedical Research (19A025), the Société Académique de Genève, the Plan

Stratégique Sciences Vie (PSVIE) of the University of Geneva, and the Canton of Geneva.

AUTHOR CONTRIBUTIONS

EN conceptualized the study. EN, FM, LS, DT and NL designed experiments. FM, DT, LS,

NL, IN, MD, EV and PBD performed experiments and analyzed data. FM, DT, LS, MD, and

EN produced visualizations of the results and made figures. FM and EN wrote the manuscript.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1. Klein, M. O. *et al.* Dopamine: Functions, Signaling, and Association with Neurological Diseases. *Cell. Mol. Neurobiol.* **39**, 31–59 (2019).
2. Shulman, J. M., De Jager, P. L. & Feany, M. B. Parkinson's Disease: Genetics and Pathogenesis. *Annu. Rev. Pathol. Mech. Dis.* **6**, 193–222 (2011).
3. Connolly, B. S. & Lang, A. E. Pharmacological Treatment of Parkinson Disease: A Review. *JAMA* **311**, 1670 (2014).
4. Kalia, L. V. & Lang, A. E. Parkinson's disease. *The Lancet* **386**, 896–912 (2015).
5. Deng, H., Wang, P. & Jankovic, J. The genetics of Parkinson disease. *Ageing Res. Rev.* **42**, 72–85 (2018).
6. Haelterman, N. A. *et al.* A Mitocentric View of Parkinson's Disease. *Annu. Rev. Neurosci.* **37**, 137–159 (2014).
7. Von Stockum, S., Nardin, A., Schrepfer, E. & Ziviani, E. Mitochondrial dynamics and mitophagy in Parkinson's disease: A fly point of view. *Neurobiol. Dis.* **90**, 58–67 (2016).
8. Sonnier, L. *et al.* Progressive Loss of Dopaminergic Neurons in the Ventral Midbrain of Adult Mice Heterozygote for Engrailed1. *J. Neurosci.* **27**, 1063–1071 (2007).
9. Laguna, A. *et al.* Dopaminergic control of autophagic-lysosomal function implicates Lmx1b in Parkinson's disease. *Nat. Neurosci.* **18**, 826–835 (2015).
10. Doucet-Beaupré, H. *et al.* Lmx1a and Lmx1b regulate mitochondrial functions and survival of adult midbrain dopaminergic neurons. *Proc. Natl. Acad. Sci.* **113**, E4387–E4396 (2016).
11. Kittappa, R., Chang, W. W., Awatramani, R. B. & McKay, R. D. G. The foxa2 Gene Controls the Birth and Spontaneous Degeneration of Dopamine Neurons in Old Age. *PLoS Biol.* **5**, e325 (2007).
12. Domanskyi, A., Alter, H., Vogt, M. A., Gass, P. & Vinnikov, I. A. Transcription factors Foxa1 and Foxa2 are required for adult dopamine neurons maintenance. *Front. Cell. Neurosci.* **8**, (2014).
13. Kadkhodaei, B. *et al.* Transcription factor Nurr1 maintains fiber integrity and nuclear-encoded mitochondrial gene expression in dopamine neurons. *Proc. Natl. Acad. Sci.* **110**, 2360–2365 (2013).
14. van den Munckhof, P. *et al.* Pitx3 is required for motor activity and for survival of a subset of midbrain dopaminergic neurons. *Dev. Camb. Engl.* **130**, 2535–2542 (2003).
15. Di Salvio, M. *et al.* Otx2 controls neuron subtype identity in ventral tegmental area and antagonizes vulnerability to MPTP. *Nat. Neurosci.* **13**, 1481–1488 (2010).
16. Blaudin de Thé, F. *et al.* Engrailed homeoprotein blocks degeneration in adult dopaminergic neurons through LINE 1 repression. *EMBO J.* **37**, (2018).
17. Le, W. *et al.* Mutations in NR4A2 associated with familial Parkinson disease. *Nat. Genet.* **33**, 85–89 (2003).
18. Bergman, O. *et al.* Do polymorphisms in transcription factors LMX1A and LMX1B influence the risk for Parkinson's disease? *J. Neural Transm.* **116**, 333–338 (2009).
19. Oh, S. *et al.* Combined Nurr1 and Foxa2 roles in the therapy of Parkinson's disease. *EMBO Mol. Med.* **7**, 510–525 (2015).
20. Alvarez-Fischer, D. *et al.* Engrailed protects mouse midbrain dopaminergic neurons against mitochondrial complex I insults. *Nat. Neurosci.* **14**, 1260–1266 (2011).
21. Di Giovannantonio, L. G. *et al.* Otx2 selectively controls the neurogenesis of specific neuronal subtypes of the ventral tegmental area and compensates En1-dependent neuronal loss and MPTP vulnerability. *Dev. Biol.* **373**, 176–183 (2013).
22. Hewitt, V. L. & Whitworth, A. J. Mechanisms of Parkinson's Disease. in *Current Topics in Developmental Biology* vol. 121 173–200 (Elsevier, 2017).

23. Nagoshi, E. Drosophila Models of Sporadic Parkinson's Disease. *Int. J. Mol. Sci.* **19**,
3343 (2018).
24. Landayan, D. & Wolf, F. W. Shared neurocircuitry underlying feeding and drugs of
abuse in Drosophila. *Biomed. J.* **38**, 496–509 (2015).
25. Strausfeld, N. J. & Hirth, F. Deep Homology of Arthropod Central Complex and
Vertebrate Basal Ganglia. *Science* **340**, 157–161 (2013).
26. Liu, C. *et al.* A subset of dopamine neurons signals reward for odour memory in
Drosophila. *Nature* **488**, 512–516 (2012).
27. Riemensperger, T. *et al.* A Single Dopamine Pathway Underlies Progressive Locomotor
Deficits in a Drosophila Model of Parkinson Disease. *Cell Rep.* **5**, 952–960 (2013).
28. Bou Dib, P. *et al.* A Conserved Role for p48 Homologs in Protecting Dopaminergic
Neurons from Oxidative Stress. *PLoS Genet.* **10**, e1004718 (2014).
29. Tas, D. *et al.* Parallel roles of transcription factors dFOXO and FER2 in the development
and maintenance of dopaminergic neurons. *PLOS Genet.* **14**, e1007271 (2018).
30. Lawal, H. O. *et al.* The Drosophila vesicular monoamine transporter reduces pesticide-
induced loss of dopaminergic neurons. *Neurobiol. Dis.* **40**, 102–112 (2010).
31. Sun, J. *et al.* Neural Control of Startle-Induced Locomotion by the Mushroom Bodies and
Associated Neurons in Drosophila. *Front. Syst. Neurosci.* **12**, 6 (2018).
32. Y. Ono, Nakatani, T., Minaki, Y. & Kumai, M. The basic helix-loop-helix transcription
factor Nato3 controls neurogenic activity in mesencephalic floor plate cells. *Development*
137, 1897–1906 (2010).
33. Nissim-Eliraz, E., Zisman, S., Schatz, O. & Ben-Arie, N. Nato3 Integrates with the Shh-
Foxa2 Transcriptional Network Regulating the Differentiation of Midbrain Dopaminergic
Neurons. *J. Mol. Neurosci.* **51**, 13–27 (2013).
34. Mansour, A. A., Khazanov-Zisman, S., Netser, Y., Klar, A. & Ben-Arie, N. Nato3 plays
an integral role in dorsoventral patterning of the spinal cord by segregating floor plate/p3
fates via Nkx2.2 suppression and Foxa2 maintenance. *Development* **141**, 574–584 (2014).
35. Zimprich, A. *et al.* Mutations in LRRK2 Cause Autosomal-Dominant Parkinsonism with
Pleomorphic Pathology. *Neuron* **44**, 601–607 (2004).
36. Langston, R. G., Rudenko, I. N. & Cookson, M. R. The function of orthologues of the
human Parkinson's disease gene LRRK2 across species: implications for disease
modelling in preclinical research. *Biochem. J.* **473**, 221–232 (2016).
37. Kitada, T. *et al.* Mutations in the parkin gene cause autosomal recessive juvenile
parkinsonism. *Nature* **392**, 605–608 (1998).
38. Cha, G.-H. *et al.* Parkin negatively regulates JNK pathway in the dopaminergic neurons
of Drosophila. *Proc. Natl. Acad. Sci.* **102**, 10345–10350 (2005).
39. Park, J. *et al.* Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented
by parkin. *Nature* **441**, 1157–1161 (2006).
40. Park, J., Lee, G. & Chung, J. The PINK1–Parkin pathway is involved in the regulation of
mitochondrial remodeling process. *Biochem. Biophys. Res. Commun.* **378**, 518–523
(2009).
41. Greene, J. C. *et al.* Mitochondrial pathology and apoptotic muscle degeneration in
Drosophila parkin mutants. *Proc. Natl. Acad. Sci.* **100**, 4078–4083 (2003).
42. Clark, I. E. *et al.* Drosophila pink1 is required for mitochondrial function and interacts
genetically with parkin. *Nature* **441**, 1162–1166 (2006).
43. Poole, A. C. *et al.* The PINK1/Parkin pathway regulates mitochondrial morphology. *Proc.*
Natl. Acad. Sci. **105**, 1638–1643 (2008).
44. Scialò, F. *et al.* Mitochondrial ROS Produced via Reverse Electron Transport Extend
Animal Lifespan. *Cell Metab.* **23**, 725–734 (2016).

45. Croset, V., Treiber, C. D. & Waddell, S. Cellular diversity in the Drosophila midbrain
revealed by single-cell transcriptomics. *eLife* **7**, e34550 (2018).
46. Davie, K. *et al.* A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain. *Cell*
174, 982-998.e20 (2018).
47. Flynn, J. M. & Melov, S. SOD2 in mitochondrial dysfunction and neurodegeneration.
Free Radic. Biol. Med. **62**, 4–12 (2013).
48. Shokri, L. *et al.* A Comprehensive Drosophila melanogaster Transcription Factor
Interactome. *Cell Rep.* **27**, 955-970.e7 (2019).
49. Zhu, J., Chen, L., Sun, G. & Raikhel, A. S. The Competence Factor Ftz-F1 Potentiates
Ecdysone Receptor Activity via Recruiting a p160/SRC Coactivator. *Mol. Cell. Biol.* **26**,
9402–9412 (2006).
50. Xie, X.-J. *et al.* CDK8-Cyclin C Mediates Nutritional Regulation of Developmental
Transitions through the Ecdysone Receptor in Drosophila. *PLoS Biol.* **13**, e1002207
(2015).
51. Peterson, D. J. *et al.* The Basic Helix-Loop-Helix Gene Nato3 Drives Expression of
Dopaminergic Neuron Transcription Factors in Neural Progenitors. *Neuroscience* **421**,
176–191 (2019).
52. Turiault, M. *et al.* Analysis of dopamine transporter gene expression pattern -- generation
of DAT-iCre transgenic mice. *FEBS J.* **274**, 3568–3577 (2007).
53. Engblom, D. *et al.* Glutamate receptors on dopamine neurons control the persistence of
cocaine seeking. *Neuron* **59**, 497–508 (2008).
54. Matsuura, K., Kabuto, H., Makino, H. & Ogawa, N. Pole test is a useful method for
evaluating the mouse movement disorder caused by striatal dopamine depletion. *J.*
Neurosci. Methods **73**, 45–48 (1997).
55. Gould, T. D., Dao, D. T. & Kovacsics, C. E. The open field test. in *Mood and anxiety*
related phenotypes in mice: Characterization using behavioral tests 1–20 (Humana Press,
2009). doi:10.1007/978-1-60761-303-9_1.
56. Moore, A. W., Barbel, S., Jan, L. Y. & Jan, Y. N. A genomewide survey of basic helix-
loop-helix factors in Drosophila. *Proc. Natl. Acad. Sci.* **97**, 10436–10441 (2000).
57. Segev, E., Halachmi, N., Salzberg, A. & Ben-Arie, N. Nato3 is an evolutionarily
conserved bHLH transcription factor expressed in the CNS of Drosophila and mouse.
Mech. Dev. **106**, 197–202 (2001).
58. Verzi, M. P. *et al.* N-twist, an evolutionarily conserved bHLH protein expressed in the
developing CNS, functions as a transcriptional inhibitor. *Dev. Biol.* **249**, 174–190 (2002).
59. Khazanov, S. *et al.* Floor plate descendants in the ependyma of the adult mouse Central
Nervous System. *Int. J. Dev. Biol.* **61**, 257–265 (2017).
60. Arenas, E., Denham, M. & Villaescusa, J. C. How to make a midbrain dopaminergic
neuron. *Development* **142**, 1918–1936 (2015).
61. Vilain, S. *et al.* The Yeast Complex I Equivalent NADH Dehydrogenase Rescues pink1
Mutants. *PLoS Genet.* **8**, e1002456 (2012).
62. Pogson, J. H. *et al.* The complex I subunit NDUFA10 selectively rescues Drosophila
pink1 mutants through a mechanism independent of mitophagy. *PLoS Genet.* **10**,
e1004815 (2014).
63. Biosa, A. *et al.* Superoxide dismutating molecules rescue the toxic effects of PINK1 and
parkin loss. *Hum. Mol. Genet.* **27**, 1618–1629 (2018).
64. Mahlknecht, P., Seppi, K. & Poewe, W. The Concept of Prodromal Parkinson's Disease.
J. Park. Dis. **5**, 681–697 (2015).
65. Carvey, P. M., Punati, A. & Newman, M. B. Progressive dopamine neuron loss in
Parkinson's disease: the multiple hit hypothesis. *Cell Transplant.* **15**, 239–250 (2006).

66. Spathis, A. D. *et al.* Nurr1:RXRα heterodimer activation as monotherapy for Parkinson's
disease. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 3999–4004 (2017).

67. Nagoshi, E. *et al.* Dissecting differential gene expression within the circadian neuronal
circuit of Drosophila. *Nat. Neurosci.* **13**, (2010).

68. Imai, Y. *et al.* Phosphorylation of 4E-BP by LRRK2 affects the maintenance of
dopaminergic neurons in Drosophila. *EMBO J.* **27**, 2432–2443 (2008).

69. Schindelin, J. *et al.* Fiji - an Open Source platform for biological image analysis. *Nat.*
Methods **9**, (2012).

70. Owusu-Ansah, E., Yavari, A. & Banerjee, U. A protocol for in vivo detection of reactive
oxygen species. *Protoc. Exch.* (2008) doi:10.1038/nprot.2008.23.

71. Meireles-Filho, A. C. A., Bardet, A. F., Yáñez-Cuna, J. O., Stampfel, G. & Stark, A. cis-
regulatory requirements for tissue-specific programs of the circadian clock. *Curr. Biol.*
CB **24**, 1–10 (2014).

72. Zhou, J., Yu, W. & Hardin, P. E. ChIPping away at the Drosophila clock. *Methods*
Enzymol. **551**, 323–347 (2015).

73. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–359 (2012).

74. Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X. S. Identifying ChIP-seq enrichment using
MACS. *Nat. Protoc.* **7**, (2012).

75. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–
589 (2010).

76. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

77. David, F. P. A. *et al.* HTSstation: A Web Application and Open-Access Libraries for
High-Throughput Sequencing Data Analysis. *PLOS ONE* **9**, e85879 (2014).

78. Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2018 update. *Nucleic Acids Res.* **46**, W537–W544 (2018).

79. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of
insertions, deletions and gene fusions. *Genome Biol.* **14**, R36 (2013).

80. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat. Commun.* **10**, 1523 (2019).

81. Hu, Y. *et al.* An integrative approach to ortholog prediction for disease-focused and other
functional studies. *BMC Bioinformatics* **12**, 357 (2011).

82. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

83. Zambelli, F., Pesole, G. & Pavesi, G. Pscan: finding over-represented transcription factor
binding site motifs in sequences from co-regulated or co-expressed genes. *Nucleic Acids*
Res. **37**, W247-252 (2009).

84. Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased
coverage, supporting functional discovery in genome-wide experimental datasets.
Nucleic Acids Res. **47**, D607–D613 (2019).

85. Nikonenko, I. *et al.* PSD-95 promotes synaptogenesis and multiinnervated spine
formation through nitric oxide signaling. *J. Cell Biol.* **183**, 1115–1127 (2008).

86. Hellemans, J., Mortier, G., De Paepe, A., Speleman, F. & Vandesompele, J. qBase
relative quantification framework and software for management and automated analysis
of real-time quantitative PCR data. *Genome Biol.* **8**, R19 (2007).

87. Bello-Arroyo, E. *et al.* MouBeAT: A New and Open Toolbox for Guided Analysis of
Behavioral Tests in Mice. *Front. Behav. Neurosci.* **12**, 201 (2018).

1003
1004

 FIGURE LEGENDS

 Figure 1. *Fer2* overexpression prevents the loss of PAM dopaminergic neurons and

 mitochondrial dysfunction in genetic models of PD.

 (a and b) *Fer2* overexpression prevented the loss of PAM neurons induced by the targeted

 expression of wild-type (*WT*) or pathogenic mutant *Lrrk* in PAM neurons. **(a)** Representative

 images of PAM neurons at day 14 detected by anti-TH antibodies (green). *PAM*, PAM

 neuron-specific *R58E02-GAL4* driver. *PAM>X* indicates that *UAS-transgene X* is driven by

 R58E02-GAL4. Scale bar, 20 μm. Top left, a schematic of DA neurons in the PAM (green)

 and other clusters (black). **(b)** Quantification of the number of PAM neurons per hemisphere.

 Box boundaries are the 25th and 75th percentiles, the horizontal line across the box is the

 median, and the whiskers indicate the minimum and maximum values. n=7-10 brains per

 group. **p<0.01, ***p<0.001, ****p<0.0001. n.s., not significant. One-way ANOVA

 followed by a Turkey's test for multiple group comparison. **(c and d)** *Fer2* overexpression in

 PAM neurons rescued PAM neuron loss in *Park^1* mutants. Quantification of the number of

 PAM neurons per hemisphere in the indicated age and genotypes. n=7-10 brains per group.

 *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. n.s., not significant. One-way ANOVA

 followed by a Turkey's test for multiple group comparison **(c)** and Mann-Whitney test **(d)**.

 (e) The mitochondrial network in PAM neurons was visualized using mitoGFP. Scale bar, 5

 μm. **(f)** Top, histograms showing the circularity distribution of GFP$^+$ particles measured from

 (e). Bottom, the Gaussian curve fit to the circularity histogram. Circularity ranges from 0

 (reticular mitochondria) to 1 (round mitochondria). The histogram of *PAM>mitoGFP, park^1*

 is right-shifted compared with those of *PAM>mitoGFP, park^1/+* and *PAM>mitoGFP, Fer2,*

 park^1, indicating the accumulation of rounder mitochondria in *park^1* and its rescue by *Fer2*.

 ****p<0.0001 using Kruskal-Wallis test followed by Dunn's test for multiple group

 comparison. More than 500 mitochondrial particles were analyzed from 7-10 brains per

1030   genotype.

1031

Figure 2. *Fer2* overexpression protects PAM dopaminergic neurons from oxidative

insults.

(a) w^{1118}, *Fer2²/+*, *Fer2²*, *R58E02-GAL4* (*PAM*) and PAM>*Fer2* flies were fed with H_2O_2 at

day 7 for 24 h and the number of PAM neurons was examined at day 14. *Fer2* mutation

exacerbates, whereas *Fer2* overexpression prevents, H_2O_2-induced PAM neuron loss. n=7-10

brains per group. Mann-Whitney test, **p<0.01, ***p<0.001, ****p<0.0001. **(b)** ROS levels

in w^{1118} flies at different ages were monitored using H2DCF dye. Scale bar, 25 μm. **(c)**

Quantification of ROS levels in the central brain from (**b**). Data are represented as mean ±

SEM of the fold change relative to day 1. n=7-10 brains per group. One-way ANOVA

followed by a Turkey's test for multiple group comparison, **p<0.01. **(d)** Brains of

Fer2::GFP::V5 flies of different ages were stained with anti-GFP (green) and PAM neurons

were detected by anti-TH antibodies (magenta). Scale bar, 5 μm. **(e)** GFP and TH

fluorescence intensities in the PAM region from (**d**) were quantified. Data are represented as

mean ± SEM of the fold change relative to day 1. n=7-10 brains per group. One-way

ANOVA followed by a Turkey's test for multiple group comparison, *p<0.05, ****p<0.0001.

1047

Figure 3. Identification of *Fer2* transcriptional targets through a combination of ChIP-

seq and RNA-seq approaches.

(a) ChIP-seq was performed on 14-day-old *Fer2::GFP::V5, Fer2¹* flies. w^{1118} flies were used

as a negative control. Two biological replicates per group. **(b)** A sequence containing an E-

box (CANNTG, underlined) was found enriched in FER2 ChIP-seq peaks over w^{1118} ChIP-

seq. **(c)** Scheme of the whole-head RNA-seq analysis to identify DEG after heat shock-driven

Fer2 transient overexpression. Three biological replicates per group. The number of DEGs

up- and downregulated after heat shock are shown**. (d)** Examples of ChIP-seq and RNA-seq

data (48 h) for two *Fer2* direct targets, *CG8128* and *skd*. The rectangles indicate ChIP-seq

peaks.

1058

Figure 4. *Fer2* targets contribute to PAM neurons maintenance and mitochondrial

health.

(a) Scheme of RNA-seq analysis of isolated PAM neurons to identify genes regulated by

Fer2 constitutive overexpression. Three biological replicates per group. **(b)** Intersection

between peak-associated genes identified by FER2 ChIP-seq, DEG identified by whole-head

RNA-seq 48 h after heat shock, and genes expressed in PAM neuron RNA-seq. **(c)** Network

visualization of the 26 *Fer2* direct targets expressed in PAM neurons and their interactions.

Each node represents a gene and is colored based on its function. The shape of the node

indicates whether the gene is up- or downregulated by *Fer2*. The line thickness indicates the

strength of data support. **(d)** Position of FER2 peaks and effect of *Fer2* overexpression on the

expression of *Fer2* direct targets. **(e)** Volcano plot of 48 h heat shock RNA-seq data. Orange

(downregulated) and blue (upregulated) dots indicate the 26 *Fer2* direct targets. **(f)**

Functional validation of *Fer2* direct targets regarding their effect on PAM neuron survival.

Quantification of the number of PAM neurons per hemisphere detected by anti-TH

immunostaining in 35-day-old flies, upon overexpression (OE) or knock-down (RNAi) of the

indicated gene in PAM neurons. n=7-10 per genotype. Genes that cause a statistically

significant loss of PAM neurons are highlighted in red. A second independent replicate was

assessed for these genes to confirm the significant loss of PAM neurons. One-way ANOVA

followed by a Dunnett's test for multiple group comparison,**p<0.01, ****p<0.0001. **(g)**

Network visualization of the most enriched GO terms among DEGs in PAM RNA-seq. GO

terms were grouped into clusters based on their membership similarities. Each node

represents a GO term and is colored based on its cluster ID. GO terms that did not cluster are

not shown. **(h)** Quantification of the number of PAM neurons per hemisphere, as detected by

anti-TH immunostaining for the indicated genotypes and ages. *Sod2* overexpression in PAM

neurons partially rescued PAM neuron loss in *Fer2^2* mutants. n=7-10 per group. One-way

ANOVA followed by a Turkey's test for multiple group comparison, *$p<0.05$, ****$p<0.0001$.

1085

Figure 5. *Nato3* is expressed in the midbrain DA neurons in adult mice. (a) Screenshot

from GTEx portal (gtexportal.org) indicating *NATO3* expression levels in human tissues

measured by RNA-seq. **(b)** RT-qPCR analysis of *Nato3* mRNA expression levels in different

brain regions of C57BL6 mice at 4 months. OB, olfactory bulbs. PFC, prefrontal cortex. vMB,

ventral midbrain. dMB, dorsal midbrain. Data are represented as mean ± SEM of the fold

change relative to the expression in the OB. n=3 animals per region. One-way ANOVA

followed by a Turkey's test for multiple group comparison, vMB vs any region other than

dMB ***$p<0.001$, dMB vs any other region than vMB and cerebellum **$p<0.01$, dMB vs

cerebellum #$p<0.05$. **(c)** RT-qPCR analysis of *Nato3* mRNA expression levels in the ventral

midbrain of C57BL6 mice at different ages. Data are represented as mean ± SEM of the fold

change relative to the expression at 1 month. n=3-5 animals per group. No statistically

significant difference between different ages by ANOVA. **(d)** Representative images of anti-

NATO3 (green) and anti-TH (magenta) immunostaining in coronal sections of the ventral

midbrain of 14-month-old mice. The region shown in each raw is illustrated by the scheme

on the right. Scale bars indicate 200 μm (top panels) and 50 μm (central and bottom panels).

1101

Figure 6. *Nato3* cKO in DA neurons leads to age-dependent impairment in locomotor

coordination and activity.

(a) Strategy for conditional ablation of *Nato3* in differentiated DA neurons. **(b)** RT-qPCR

analysis of *Nato3* mRNA levels in ventral midbrains of 2-month-old *Nato3^{floxed}* (ctrl) and

Nato3$^{DAT-Cre}$ (cKO) mice. Data are represented as mean ± SEM of the fold change relative to

the expression in ctrl. n=4 mice per genotype. Mann-Whitney test, *p<0.05. **(c)** The motor

coordination of control and *Nato3* cKO mice was assessed at different ages using the pole test.

The time taken to orient downward at the top of the pole (T-turn; left) and the total time to

turn and descend the pole (T-tot; right) are shown. n=7-11 mice per group. Data are

represented as mean ± SEM. Mann-Whitney test comparing ctrl and cKO of the same age,

*p<0.05, **p<0.01. **(d)** Locomotor activity assessed at different ages using the voluntary

wheel running test. The time spent running on the wheel (left) and the total distance run

(right) are shown. n=7-11 mice per group. Data are represented as mean ± SEM. Mann-

Whitney test comparing ctrl and cKO of the same age, *p<0.05. **(e)** Representative TH

immunostaining images of coronal ventral midbrain sections from 16- to 18-month-old

control and *Nato3* cKO mice. Scale bar, 200µm. **(f)** Quantification of TH$^+$ neurons in SN and

VTA from **(e)**. n=3 mice per genotype. Data are represented as mean ± SD of the fold change

relative to control. No statistically significant difference between ctrl and cKO by Mann-

Whitney test. **(g)** Representative TH immunostaining images of coronal striatal sections from

16- to 18-month-old control and *Nato3* cKO mice. Scale bar, 200µm. **(h)** Optical

densitometry from TH immunostaining of the dorsal striatum in **(g)**. n=4 mice per genotype.

Data are represented as mean ± SD of the fold change relative to control. No statistically

significant difference between ctrl and cKO by Mann-Whitney test.

1125

Figure 7. *Nato3* is essential for mitochondrial integrity in the midbrain DA neurons.

(a) Mitochondria were visualized by anti-COX4 immunostaining (magenta) in TH$^+$ neurons

(green) of SN in 16- to 18-month-old control and *Nato3* cKO mice. Scale bar, 10µm. **(b)** The

circularity of COX4$^+$ mitochondria particles in TH$^+$ neurons from **(a)** was measured and its

distribution was plotted in a histogram. The histogram for the cKO mouse is significantly

right-shifted compared with that of the control. Kolmogorov-Smirnov test, p<0.0001. More

than 700 COX4$^+$ particles per genotype were analyzed from three brains per genotype. **(c)**

Electron micrographs of mitochondria from TH-labeled DA neurons from the SN of 16- to

18-month-old ctrl and *Nato3* cKO mice. The right bottom panel illustrates the difference

between a degenerating mitochondrion in the dendrite of a *Nato3*-ablated TH$^+$ neuron, and a

normal mitochondrion in an adjacent non-DA neuron. M, normal mitochondria. DM,

degenerative mitochondria. Scale bar, 0.5 μm. **(d)** Classification criteria for mitochondria

integrity (top) and mitochondria distribution in TH$^+$ neurons for ctrl and cKO mice (bottom).

In total, 200 to 350 mitochondria from TH$^+$ neurons in SN were analyzed per genotype. **(e)**

Measurement of size (left) and roundness (right) of mitochondria. 200 to 350 mitochondria

from SN TH$^+$ neurons were analyzed per genotype. Data are represented as mean ± SEM.

Mann-Whitney test, **p<0.01.

1143

Supplementary Figure 1. *Fer2* loss of function and *Lrrk* gain of function or *park^1*

mutation additively increase PAM neuron loss.

Quantification of the number of PAM neurons per hemisphere, as detected by anti-TH

immunostaining for the indicated genotypes and ages. *Fer2>X* indicates that *UAS-transgene*

X is expressed with *Fer2-GAL4*. n=7-10 brains per group. One-way ANOVA followed by a

Turkey's test for multiple group comparison., *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001. In (**a**), where brackets are not shown, flies overexpressing *Lrrk* and *Fer2-*

Gal4 flies (ctrl) of the same age are compared. In (**b**) at day 1, §p<0.05 when comparing

Fer2-Gal4, Fer2^2/+ with *Fer2>Lrrk WT, Fer2^2/+,* and #p<0.0001 when comparing *Fer2-*

Gal4, Fer2^2/+ with all the other genotypes. At day 14 and 35, &p<0.0001 when comparing

Fer2-Gal4, Fer2^2/+ with all the other genotypes at the same age.

1155

Supplementary Figure 2. *Fer2::GFP::V5* **transgene rescues PAM neuron loss and locomotor deficits in** *Fer2^1* **mutants.**

(**a**) Brains of *Fer2::GFP::V5, Fer2^1* flies at day 14 were stained with anti-GFP (green) and anti-TH (magenta) antibodies. The right panel presents a high-magnification image of the PAM neurons in the square from the left panel. Left panel scale bar, 50 μm. Right panel scale bar, 5 μm. (**b**) Quantification of the number of PAM neurons per hemisphere, as detected by anti-TH immunostaining in *w^{1118}*, *Fer2^1* and *Fer2::GFP::V5, Fer2^1* flies at day 14. n=7-10 brains per group. One-way ANOVA followed by a Turkey's test for multiple group comparison, ****$p < 0.0001$. n.s., not significant. (**c**) Climbing index of *w^{1118}*, *Fer2^1* and *Fer2::GFP::V5, Fer2^1* flies at day 14. Data are represented as mean of three independent experiments ± SD. One-way ANOVA followed by a Turkey's test for multiple group comparison, ****$p < 0.0001$. n.s., not significant.

1168

Supplementary Fig. 3. Genetic pathways downstream of *Fer2*.

(**a**) List of transcription factors that are expressed in PAM neurons and whose binding sites are significantly overrepresented in the promoters of PAM RNA-seq DEGs. Enrichment p-value and the databases used for the analysis are indicated. Transcription factors known or predicted to interact with one or more *Fer2* direct targets are highlighted in orange. (**b**) Model of the role of *Fer2* in dopaminergic neuroprotection. *Fer2* binds to and regulates the expression of a set of direct target genes, including multiple transcription factors and chromatin regulators. *Fer2* direct targets FTZ-F1 and SMR interact with D, Br and USP and regulate a large set of genes within PAM neurons. In this manner, *Fer2* controls multiple pathways leading to regulation of mitochondrial gene expression, improved mitochondrial health, and PAM neuron survival against genetic and oxidative insults.

1180

Supplementary Figure 4. Locomotor activity measured in the open field test is not

affected in *Nato3* cKO mice.

Levels of horizontal activity (left), measured as the total distance traveled in the arena, and

vertical activity (right), measured as the number of rearing events, were not different between

Nato3 cKO and control mice. n=5-7 mice per group. Data are represented as mean ± SEM.

No statistically significant difference between ctrl and cKO by Mann-Whitney test.

1

2

a

day 7
H₂O₂ 24h

day 14
TH staining

Number of PAM neurons

w^{1118} $Fer2^2/+$ $Fer2^2$ PAM   PAM>Fer2

*** ** **** ** ns

□ ctrl   ■ H₂O₂

b

day 1   day 14   day 35

ROS

c

Fluorescence intensity

**

1   14   35
Age (days)

d

day 1   day 14   day 35

TH Fer2::GFP::V5

e

Fluorescence intensity

****
*

■ FER2   □ TH

1   14   35
Age (days)

3

a ChIP-seq

Fer2::GFP::V5, Fer2¹

PAM

w¹¹¹⁸ (neg ctrl)

PAM

267 peaks
279 associated genes

b

CTGTCAGCTGCA

p-value : e^{-42}

c Head RNA-seq

hs>Fer2
hs>Fer2 + heat-shock
$\}$ DEG after heat-shock induced Fer2 OE

hs-Gal4
hs-Gal4 + heat-shock
$\}$ DEG after heat-shock (independent on Fer2)

DEG after Fer2 OE (independent on heat-shock)

heat shock
3h 37°C
12h
48h

head RNA-seq
head RNA-seq

down 132   up 196   down 423   up 493

d

ChIP-seq
Fer2::GFP::V5, Fer2¹
Fer2::GFP::V5, Fer2¹
w¹¹¹⁸
w¹¹¹⁸

RNA-seq *hs>Fer2*
(-) heat-shock
(-) heat-shock
(-) heat-shock
48h heat-shock
48h heat-shock
48h heat-shock

CG8128

ChIP-seq
Fer2::GFP::V5, Fer2¹
Fer2::GFP::V5, Fer2¹
w¹¹¹⁸
w¹¹¹⁸

RNA-seq *hs>Fer2*
(-) heat-shock
(-) heat-shock
(-) heat-shock
48h heat-shock
48h heat-shock
48h heat-shock

skd

4

a PAM RNA-seq

PAM>RedStinger ctrl
PAM>RedStinger, Fer2 OE

PAM

Brain dissociation and isolation of PAM neurons

PAM neuron transcriptome

7134 genes expressed in PAM
132 DEG, p-value<0.01

b genes bound ChIP-seq
DEG 48h head RNA-seq

120 | 6 | 432
127 | **26**
452
6529
132 DEG
genes expressed in PAM RNA-seq

Fer2 direct targets

c
CadN
B4
Appl
RhoGAP93B
Rapgap1
gpp
sty
sba
ftz-f1
Eip93F
cdi
skd
tna
osa
tlk
Theg
CG1578
Eip75B
CG34357
CG4998
Ih
Smr
Snoo
PAN3
Msi
Ae2

○ down-regulated
◇ up-regulated

Gene function
- 8% kinase
- 12% signaling
- 35% TF/chromatin
- 8% RNA metabolism
- 4% channel
- 4% receptor
- 4% transmembrane
- 12% other
- 15% unknown

d Peak position | Gene regulation

- 15% Upstream
- 4% Promoter
- 62% Introns
- 4% Exons
- 8% Downstream
- 8% Intergenic

- 12% up-regulated
- 88% down-regulated

e
● *Fer2* direct target down-regulated
● *Fer2* direct target up-regulated
heat shock, 48h

$-\log_{10}$ (p-value)
p=0.05
$\log_2$ (fold change)

f
Number of PAM neurons (% of *Fer2-GAL4*)

**** ** **** ** ** **

Fer2-Gal4, *Appl OE*, *B4 OE*, *CadN OE*, *CG1578 RNAi*, *CG34357 OE*, *CG4998 RNAi*, *Eip75B OE*, *Eip93F OE*, *ftz-f1 OE*, *Ih OE*, *msi OE*, *osa OE*, *rapGAP1 OE*, *RhoGAP93B OE*, *Snoo OE*, *sty OE*, *Theg RNAi*, *Tlk OE*, *tna OE*

Fer2 secondary targets

g
CG5037
CG7639
Sdhaf3
ND-15
Cchl
Sod2

- mitochondrial ribosome
- mitochondrial envelope
- mitochondrial interm. space
- endoplasmic reticulum
- vacuoles lysosomes

h
Number of PAM neurons

ns
*
*
****

Age (days)
1 | 14

■ *Fer2-Gal4, Fer2²*
▧ *Fer2>Sod2, Fer2²*

5

Human *NATO3* gene expression (gtexportal.org)

a

Mouse *Nato3* gene expression

b

Nato3 mRNA

c

Nato3 mRNA (vMB)

d

NATO3	TH	MERGE

vMB

SN

VTA

6

a

b *Nato3* mRNA (vMB)

c Pole test : T-turn   Pole test : T-tot

d Wheel test : time on the wheel   Wheel test : distance run on the wheel

e   **f**   **g**   **h**

7

a

TH   COX4

ctrl

cKO

b

Percentage of COX4+ particles

p-value < 0.0001 [ctrl ■ cKO □

circularity
0 - 0.1, 0.1 - 0.2, 0.2 - 0.3, 0.3 - 0.4, 0.4 - 0.5, 0.5 - 0.6, 0.6 - 0.7, 0.7 - 0.8, 0.8 - 0.9, 0.9 - 1

c

ctrl

M   M   M   M

cKO

DM   DM   DM   DM   M   TH+ dendrite   DM   neighbouring cell

d

normal (M)   degenerative (DM)

mitochondria classification (% of the total)

ctrl   88   12   □ normal   ■ degenerative
cKO   54   46

0 10 20 30 40 50 60 70 80 90 100

e

Mitochondrial size (µm²)   ctrl   cKO

Mitochondrial roundness   **   ctrl   cKO