Title: The BMP antagonist *Gremlin1* contributes to the development of cortical

excitatory neurons, motor balance and fear responses

Running Title: *Grem1* in cortical neuron development

Mari Ichinose[1,4], Nobumi Suzuki[1,4], Tongtong Wang[1,4], Hiroki Kobayashi[1,2,4], Laura

Vrbanac[1,4], Jia Q Ng[1,4], Josephine A Wright[1,4], Tamsin R M Lannagan[1,4], Krystyna

A Gieniec[1,4], Martin Lewis[3,4], Ryota Ando[2], Atsushi Enomoto[2], Simon Koblar[1,4],

Paul Thomas[1,4], Daniel L Worthley[4], Susan L Woods[1,4]

1. School of Medicine, University of Adelaide, SA, Australia

2. Department of Pathology, Nagoya University Graduate School of Medicine,

Nagoya, Japan

3. College of Medicine and Public Health, Flinders University, Bedford Park, SA,

Australia

4. South Australian Health and Medical Research Institute, Adelaide, SA, Australia

Corresponding author: Susan L Woods

susan.woods@adelaide.edu.au

Key words

Grem1, BMP, cortical development, excitatory neuron

Summary statement

The BMP antagonist, *Grem1,* marks neuroprogenitors that give rise to deep layer

glutamatergic neurons in the embryonic mouse brain. *Grem1* conditional knockout

mice display cortical and behavioural abnormalities.

Abstract

Bone morphogenetic protein (BMP) signaling is required for early forebrain

development and cortical formation. How the endogenous modulators of BMP

signaling regulate the structural and functional maturation of the developing brain

remains unclear. Here we show that expression of the BMP antagonist, *Grem1,*

marks a neuroprogenitor that gives rise to layer Ⅴ and Ⅵ glutamatergic neurons

in the embryonic mouse brain. Lineage tracing of *Grem1*-expressing cells in the

embryonic brain was examined by administration of tamoxifen to pregnant

Grem1creERT; *Rosa26LSLTdtomato* mice at 13.5 days post coitum (dpc), followed by

collection of embryos later in gestation. In addition, at 14.5 dpc, bulk mRNA seq

analysis of differentially expressed transcripts between FACS sorted *Grem1* positive

and negative cells was performed. We also generated *Emx1-cre* mediated *Grem1*

conditional knockout mice (*Emx1-Cre;Grem1$^{flox/flox}$*) in which the *Grem1* gene was

deleted specifically in the dorsal telencephalon. *Grem1^{Emx1cKO}* animals had reduced

cortical thickness, especially layers Ⅴ and Ⅵ and impaired motor balance and

fear sensitivity compared to littermate controls. This study has revealed new roles

for Grem1 in the structural and functional maturation of the developing cortex.

Introduction

During development of the nervous system, bone morphogenetic protein (BMP) signaling has important roles in the promotion of dorsal identity, and the regulation of cell proliferation and differentiation. Yet BMP function in the maturation of neurons in vivo is poorly understood. Misregulation of BMP signaling has been suggested to contribute to human neurodevelopmental conditions such as autism spectrum disorders (Kumar et al., 2019), however a detailed understanding of the exact role for BMP signaling in these disorders is lacking, in line with our imperfect knowledge of the regulation of the BMP pathway in normal brain development.

The BMP ligands (BMP 2, 4, 5, 6 and 7) bind to type I (BmprIA, BmprIB, Acvr1) and type II (BmprII, ActrIIA, and ActrIIB) receptors. BMP ligands are expressed broadly in the telencephalon during early gestation in mice and then become more tightly localized to the choroid plexus at 13.5 dpc (Furuta et al., 1997). BMPs have also been derived from the mesenchyme of the brain, such as meninges and endothelial cells (Imura et al., 2008, Choe and Pleasure, 2018). Both type I and II BMP receptors are expressed in the telencephalon during development but this is further restricted in adulthood to BmprII in the cortex and hippocampus and ActrIIA/IIB in the dentate gyrus (Söderström et al., 1996). Aside from this spatiotemporal regulation of receptors, the BMP signaling pathway is also regulated by a family of secreted extracellular antagonists, that directly bind to the BMP ligands to prevent interactions with BMP receptors both in development and disease (Ali and Brazil, 2014). Antagonists such as Gremlin1 (Grem1), Noggin (Nog) and Chordin (Chrd) have been shown to inhibit BMP action in a range of different cell types and developmental stage-specific contexts to provide exquisite spatiotemporal regulation of the pathway. The roles of Nog and Chrd have been partially elucidated:

they are required for forebrain development (Bachiller et al., 2000), as well as to

create a niche for adult hippocampal neurogenesis (Lim et al., 2000, Sun et al., 2007).

The expression and function of Grem1 in the developing brain has not yet been

determined. Grem1 is an extracellular secreted antagonist of BMP 2,4, and 7 that

signals to intestinal stem cells in the gut (Worthley et al., 2015) and plays a crucial

role in *Xenopus* dorsalisation (Hsu et al., 1998) and limb and kidney formation

(Khokha et al., 2003). Likewise the role of Grem1 in the normal adult CNS is

unmapped territory to date, outside of prior work in the pathogenic state of glioma

(Yan et al., 2014, Guan et al., 2017, Fu et al., 2018).

BMP signaling has been implicated in the regulation of forebrain patterning during

early embryogenesis, working together with other signaling pathways such as

fibroblast growth factor (FGF), Wnt, and Notch. High levels of BMP activity

suppresses anterior neural development, whereas abrogation of BMP signaling can

promote neural specification (Wilson and Houart, 2004, Lamb et al., 1993, Bachiller

et al., 2000). *In vitro* the addition of BMP 2 and 4 to mouse neural stem cell cultures

represses cell proliferation (Mathieu et al., 2008). BMP ligands regulate neuronal

differentiation as well as determination of glial cell fate, by promoting astrocyte

differentiation at the expense of oligodendrocytic fates (Mathieu et al., 2008, Yun et

al., 2004, Sun and Xu, 2010, Gomes et al., 2003).

Cortical neural stem cells (also known as radial glial cells, RGC) begin to divide

asymmetrically to start producing neurons at around 11.5 dpc in mice, signaling the

start of cortical development. RGC daughter cells detach from the ventricle and form

the first neuronal layer of the preplate by 13.5 dpc (Martynoga et al., 2012).

Subsequently born neurons migrate along RGCs and start to form the cortical plate,

separating the preplate into Cajal-Retzius cells in the marginal zone and subplate

neurons (Dwyer et al., 2016). The neocortex develops in an inside out manner, with

deep layers emerging first and superficial layers last. These neurons differentiate

into glutamatergic pyramidal neurons, whereas inhibitory interneurons are born in

the subcortical ganglionic eminences. Deep-layer pyramidal neurons (DLPNs, layer

V and VI) have both intratelenchephalic projections to superficial cortical layers and

extracephalic projections to other brain regions (Baker et al., 2018). Multiple

molecular mechanisms regulate this corticogenesis (Martynoga et al., 2012),

however the role of BMP signaling in cortical layer formation and functional

maturation of neurons has only been reported in limited studies so far. *In utero*

electroporation of BMP7 to murine 14.5 dpc cortical ventricular cells impaired

neuronal migration, suggesting BMP signaling regulates neuronal positioning and

migration (Choe and Pleasure, 2018). BMPs also regulate dendritogenesis and

neurite growth *in vitro* (Lee‑Hoeflich et al., 2004, Matsuura et al., 2007). This is

consistent with a recent study that suggests perturbation of BMP signaling by

delivery of a dominant-negative version of BMP IB receptor affects migration,

polarity and dendritogenesis of mouse cortical neurons *in vivo* (Saxena et al., 2018).

To further establish the role of BMP signaling in forebrain development and

neuronal function, we focus here on the expression and function of the BMP

antagonist Grem1 in the developing mouse brain. We first assess *Grem1* expression

using transgenic *Grem1* reporter mice in the dorsal telencephalon and developing

neocortex. Next, to investigate the role of *Grem1* in the developing brain we use

transcriptomic analyses of sorted mouse *Grem1*-expressing cells and human single

cell RNA sequence (scRNAseq) data, combined with mouse neural stem/progenitor

cell (NSPC) culture *ex vivo*. Lastly, to examine the functional contribution of *Grem1*

to cortical development, we conditionally delete *Grem1* in the dorsal telecephalon

and undertake behavioural testing of mutant animals and littermate controls.

Results

***Grem1* expressing cells are located in the dorsal telencephalon and give rise to deep**

layer neocortical neurons

To assess *Grem1* expression in the embryonic mouse brain we utilized transgenic

Grem1creERT; Rosa26LSLTdtomato reporter mice, in which tamoxifen treatment

results in expression of TdTomato in cells where the *Grem1* enhancer and promoter

sequences are active and the progeny of those cells (Worthley et al., 2015). Pregnant

Grem1creERT; Rosa26LSLTdtomato mice were administered tamoxifen at 11.5 dpc

and embryonic brains were collected 24 h later, but no TdTomato[+] cells were

observed, suggesting that *Grem1* is not yet expressed in the brain at this early time-

point (**Suppl Fig.1A**). Next, we administered tamoxifen to pregnant dams at 13.5 dpc,

and collected embryonic brains at 14.5, 17.5 and 20.5 dpc (**Fig.1A,B**). From this

timepoint we observed cells expressing TdTomato 24 hours after tamoxifen

administration located in the lower cortical plate and subplate of the dorsal

telencephalon, with dendrites extending to the pia matter (**Fig.1B,G**). ISH confirmed

that *Grem1* RNA was detected in almost all TdTomato[+] cells at 14.5 dpc, confirming

that the reporter line recapitulates endogenous *Grem1* expression. Six days later at

20.5 dpc, TdTomato[+] cell somas had migrated toward the lateral ventricle with

dendrites extending to the pia matter, becoming layer Ⅴ pyramidal neurons

(**Fig.1B,C**). Layer Ⅵ neurons were TdTomato[+] at 17.5 dpc and expanded in number

at 20.5 dpc (**Fig.1B,C**). ISH showed *Grem1* RNA was detected in most TdTomato⁺

layer Ⅴ pyramidal neurons, but only 22% of layer Ⅵ neurons (**Fig.1C,D**). This

suggests that *Grem1* is actively expressed in layer V neurons but not layer Ⅵ

pyramidal neurons at 20.5 dpc. Cortical *Grem1* mRNA levels dramatically decrease

after birth as determined by qRT-PCR analysis of total RNA isolated from mouse

brain cortex at birth, p10 and 4w post-birth (**Fig.S1B**).

To confirm that *Grem1* + cells are actively dividing in order to give rise to their

traced progeny, we administered the thymidine analogue BrdU concomitantly with

tamoxifen to pregnant *Grem1creERT; Rosa26LSLTdtomato* dams at 13.5 dpc

(**Fig.1A**). BrdU is incorporated into newly synthesized DNA and so marks cells that

are dividing. 55% of TdTomato⁺, layer Ⅴ pyramidal neurons cells are labelled with

BrdU at 17.5 dpc, demonstrating *Grem1* expressing cells at 13.5 dpc are progenitor

cells undergoing mitosis (**Fig.1E,F**). Fewer TdTomato⁺ cells (17%) in layer Ⅵ were

BrdU positive, suggesting lower cell proliferation over the same period (**Fig.1F**). We

subsequently assessed whether TdTomato⁺ cells express a marker of post-mitotic

mature neurons, NeuN, by immunohistochemistry. At 14.5 dpc most TdTomato⁺ cells

were NeuN negative and were located in the subplate, and lower NeuN positive

cortical plate (**Fig.1G**). In contrast, later in gestation at 20.5 dpc almost all

TdTomato⁺ cells in layer V and approximately half of those in layer VI express NeuN

(**Fig.1H**). This suggests that Grem1-lineage traced cells give rise to mature neurons

in the telencephalon.

Global transcriptomic analysis defines transcript modules enriched in *Grem1-*

expressing cells

To further characterise *Grem1* expressing cells in the developing mouse brain, we

undertook mRNAseq analysis on TdTomato[+] and negative cells from the

telencephalon. Pregnant *Grem1creERT; Rosa26LSLTdtomato* mice were

administered tamoxifen at 13.5 dpc and embryonic brains were collected at 14.5 dpc.

At this time-point almost all TdTomato[+] cells express endogenous *Grem1* RNA

(**Fig.1C**). Dissociated brain cells were flow sorted and live, TdTomato[+] and

TdTomato[-] cells were collected for transcriptomic analysis. TdTomato[+] cells

accounted for $5.1 \pm 2.1\%$ of all live cells (n=5) (**Fig.2A**). Bulk mRNA sequencing of

TdTomato[+] and TdTomato[-] populations and analysis of 1845 DEG between the two

cell populations revealed that *Grem1* was significantly upregulated in TdTomato[+]

cells and BMP transcriptional target genes (Id1, Id3, Id4) were significantly

downregulated (adjusted P value (FDR) $\leq$ 0.05 (**Fig.2B**). We next undertook a

correlation analysis to identify modules of the 1845 DEGs that are coordinately

regulated in TdTomato[+] cells (**Fig.S2**). Shared membership of a module can suggest

genes that together perform a particular function. The clustering tree depicts the

topological distance between different modules, i.e. how similar or different the

expression of transcripts within the module are from other modules and highlighted

one module that contained 288 genes with particularly tightly correlated transcripts

(light green module, **Fig.S2**). Network visualization of the correlation analysis

showed that the majority of transcripts within this tightly correlated module are

upregulated in TdTomato[+] cells (magenta square, **Fig.2C**). We investigated the

known function of genes within this module using a hypogeometric test for gene set

enrichment. Significantly enriched pathways (adjusted P value (FDR) $\leq$ 0.001) were

related to neuronal differentiation and projection, calcium signaling (possibly

associated with synapse functions) and axons (**Fig.2D**). The BMP target gene, *Id1*,

was identified as one of the central hub genes in this cluster and its expression was

significantly associated with other transcripts that have functions in neuronal

maturation (such as Lrrtm3, Ryr3) (**Fig.2E**). This suggests that neuronal functions

are upregulated in the *Grem1*-expressing TdTomato⁺ cells of the developing

telencephalon, and that this biological function is linked, at least in part, to BMP

signaling.

***Grem1/GREM1* is enriched in excitatory neuronal lineage cells during brain**

development

To further characterize the *Grem1*-expressing TdTomato⁺ cell population we

undertook a candidate gene profiling approach using the bulk mRNAseq data. Gene

expression patterns of neuronal markers from a series of differentiation stages are

shown in **Fig.3A**. Immature neuronal markers, such as *Dcx, Ncam1, Neurod 2/6,*

Tbr1, were significantly upregulated at the RNA level in TdTomato⁺ cells in

comparison to the TdTomato⁻ cells at 14.5 dpc, while neural stem cell, radial glia,

and intermediate progenitor transcripts were significantly under-represented

(p<0.05). When RGCs (marked by co-expression of Sox2 and Pax6) generate

neocortical neurons, the expression of specific transcription factors can be used to

identify neurons within particular cortical layers(Martynoga et al., 2012). Our bulk

RNAseq analysis showed that transcripts encoding the layer Ⅴ marker, *Fezf2*, and

Ⅵ markers, *Tbr1 / Sox5*, were significantly upregulated in *Grem1*-expressing

TdTomato$^+$ cells, whereas the RGC markers, *Sox2* and *Pax6* were significantly

downregulated (absolute value of log2 fold change≥2.0, FDR<0.05) and did not

colocalize with Tdtomato at 14.5 dpc by immunohistochemistry (**Fig.S3**).

To assist with identification of the neuronal subtype likely generated from *Grem1-*

expressing TdTomato$^+$ cells, we also performed GSEA on DEG transcripts between

TdTomato$^+$ and TdTomato$^-$ cells. We found a significant enrichment of a glutamate

secretion gene set in the DEGs (NES = 2.16 p = 0.0049) (**Fig.3B**), whereas GABAergic

and dopaminergic gene sets were not enriched (NES = -0.93 p = 0.75, NES = -0.69 p

= 0.87 respectively) (**Fig.S4A**). Consistent with the GSEA, glutamatergic neuron

markers, such as Slc17a7 (vGlut1), Grin1 and Grin2b were significantly upregulated

in Grem1 expressing cells at 14.5 dpc using a candidate gene approach, whereas

gabaergic and dopaminergic neuron markers, such as Slc6a1 (GABA transporter 1),

Gad1, Gad2, and Th (Tyrosine hydroxylase) were significantly downregulated

(p<0.05, **Fig.S4B**). Tbr1 plays a central role in the development of early-born cortical

excitatory neurons and regulates the connectivity of layer Ⅵ neurons (Hevner et al.,

2001). To confirm that Grem1 cells generate excitatory neuronal lineages, we

undertook immunohistochemical staining with Tbr1. This revealed *Grem1-*

expressing TdTomato$^+$ cells at 14.5 dpc expressed very low to no Tbr1, however they

did give rise to Tbr1$^+$ neurons in cortical layer Ⅵ at 20.5 dpc (**Fig.3C,D**).

To understand the relevance of our mouse focused study to the human setting, we

reanalyzed publicly available human scRNAseq data generated from human brain

at mid-gestation (22-23 weeks post-conception) (Fan et al., 2018). We categorized

each human cell in the dataset as either *GREM1$^+$* or *GREM1$^-$* based on the presence

or absence of *GREM1* RNAseq counts for each cell. Next, we compared the

transcriptional profiles of our mouse embryonic brain *Grem1*-expressing TdTomato[+]

cells and TdTomato[-] cells, with the human mid-gestational brain *GREM1*[+] and

GREM1[-] cells using a multidimensional scaling plot (**Fig.S4C**). Samples within the

same group cluster tightly together, dimension 2 separates the samples along species

lines, while dimension 1 clearly shows a similar separation of samples based on

altered expression profiles depending on *Grem1/GREM1* status in mouse and human

developing brain (**Fig.S4C**). By mapping the *GREM1*[+] cells onto a tSNE plot

generated from the human scRNAseq data, this revealed that most *GREM1*-

expressing cells accumulated in the excitatory neuron cluster within the human mid-

gestational cortex (**Fig.3E**). Lastly, we analysed the expression of BMP-signaling

components in the human scRNAseq dataset (**Fig.S4D,S5**). Interestingly, of the

pathway antagonists and similar to *GREM1, Sclerostin domain-containing protein*

1 (*SOSTDC1*) transcripts were also highly enriched in the excitatory neuron cluster,

with *CHRD, NOG, BMP Binding Endothelial Regulator* (*BMPER)* and *Follistatin*

(*FST)* transcripts also enriched in specific cell types (**Fig.S4D**). Transcripts encoding

BMP2,4,7 that are antagonized by GREM1 were detected in various cell populations,

however expression of *Bmp2,7* was inversely correlated with *Grem1* expression in

the excitatory neuron cell population suggesting paracrine pathway regulation

(**Fig.S5**). Expression of the BMP target genes, *Id1,3,4* was also inversely correlated

with *Grem1*-expression in excitatory neurons, consistent with our developmental

mouse brain data (Fig 2B). This highlights the complexity of regulation of BMP

signaling during development of the cortex.

Grem1 promotes proliferation and neural differentiation in neural stem / progenitor cells *ex vivo*

Next we collected NSPCs from embryonic brains of *Grem1^flox/flox* mice at 14.5 dpc and transduced the cells ex vivo with control, *Cre*-expressing or *Grem1*-expressing lentivirus to generate control *Grem1^flox/flox*, *Grem1^-/-* and *Grem1* over-expressing (*Grem1^O/E*) primary cultures. Endogenous Grem1 was detected in *Grem1^flox/flox* control cultures by western blot, with absent or elevated protein levels in *Grem1^-/-* and *Grem1^O/E* cultures, respectively (**Fig.4A**). Control *Grem1^flox/flox* cultures were responsive to BMP pathway induction as determined by increased BRE-luciferase reporter activity and expression of the BMP target genes, *Id1/2/3/4*, following addition of recombinant human BMP2 (**Fig.4B, Fig.S6A**). This induction of BRE-reporter activity and increase in BMP target gene transcript levels was significantly enhanced in *Grem1^-/-* cells and attenuated in *Grem1^O/E* cells (**Fig.4B, Fig.S6A**). This confirmed that Grem1 acts as an antagonist of BMP2 and suppresses downstream transcriptional targets in embryonic NSPCs.

We next employed neurosphere assays to determine the effect of Grem1 modulation on proliferation of embryonic NSPCs. Only mitogen responsive cells proliferate to form clusters termed neurospheres, where sphere size correlates with proliferative ability (Mori et al., 2006, Reynolds and Rietze, 2005). *Grem1^O/E* cells form significantly more and larger neurospheres than control *Grem1^flox/flox* cells, whereas *Grem1^-/-* cells form significantly fewer (**Fig.4D,E**). To understand this phenomenon from the aspect of cell viability, we evaluated the proliferation rates of *Grem1^flox/flox*, *Grem1^-/-* and *Grem1^O/E* cultures. Over-expression of *Grem1* significantly increased the rate of NSPC proliferation compared to control *Grem1^flox/flox* NSPCs, while

conversely the number of viable $Grem1^{-/-}$ cells did not increase over the 72h time

course analysed (**Fig.4F**). These results suggested that Grem1 contributes to the

proliferation of NSPCs. Following on from this observation, we wanted to assess the

role of Grem1 in the differentiation of NSPCs to neurons, astrocytes and

oligodendrocytes. When cultured in recombinant BMP2-containing differentiation

media, addition of 1ng/ml recombinant GREM1 significantly increased the number

of NSPCs that differentiated into Tuj1$^+$ neurons and O4$^+$ oligodendrocyte lineage

cells, and decreased the number of GFAP$^+$ astrocytes, in comparison to vehicle

treated control NPSCs (**Fig.4G,H**). This suggests that Grem1 may regulate the

differentiation potential of NPSCs.

***Grem1* is required for normal cortical development**

In order to assess the functional role of *Grem1* in mouse forebrain development in

vivo, we generated tissue-specific *Grem1* conditional knockout mice using the *empty*

spiracles homeobox 1 (*Emx1*)-*cre* driver. We first verified that *Emx1-cre* generates

efficient cre-mediated recombination in the dorsal telencephalon at 14.5 dpc by

visualizing TdTomato$^+$ cells in reporter *Emx1-cre ; Rosa26LSLTdtomato* mouse

brains (**Fig.S7A**). Next we confirmed that *Emx1-cre ; Grem1$^{flox/flox}$* (*Grem1*

conditional knockout, *Grem1^{cKO}*) mice had a significant reduction in *Grem1* RNA in

the developing forebrain compared with cre negative littermate controls by ISH at

14.5 dpc (**Fig.5A**) and real time RT-PCR at postnatal day0 (**Fig.5B**). *Grem1^{cKO}* mice

were viable and fertile. To examine the morphological consequences of conditional

Grem1 loss in the developing mouse brain we performed Nissl staining on tissue

samples from *Grem1^{cKO}* and cre negative littermate controls. At 10 weeks of age,

total cortical thickness was significantly reduced in *Grem1^cKO* mice in comparison

with *Grem1^flox/flox* littermate controls both in males and females, due to significantly

thinner cortical layers Ⅴ and Ⅵ (**Fig.5C, Fig.S7B**). The position and thickness of

each layer was confirmed by immunohistochemistry using markers specific for each

layer, i.e. CDP for layer Ⅱ‑Ⅳ, Ctip2 for layer Ⅴ and Ⅵ and Foxp2 for layer Ⅵ

(**Fig.5D,E**). Cellular density was also more sparse in *Grem1^cKO* mice in comparison

with *Grem1^flox/flox* littermate controls (**Fig.S7C,D**). Conversely, the other predominant

region of *Emx1-cre* driver activity, the hippocampus, that has important functions

in memory for navigation pertinent to our behavioural testing, displayed normal

morphology and cell density in *Grem1^cKO* mice (**Fig.S7E**).

Our earlier transcriptional network analysis from bulk RNAseq data identified an

Id1 associated gene cluster that was differentially regulated in the *Grem1*-

expressing TdTomato⁺ cells from the cortex at 14.5 dpc (**Fig.2E**). From this cluster,

Lrrtm3 and *Ryr3* were selected for further analysis because of their known functions

in synapse development (Balschun et al., 1999, Um et al., 2016, Roppongi et al., 2017,

Futagi and Kitano, 2015, Linhoff et al., 2009). The expression of these two

transcripts were significantly downregulated in the cortex of *Grem1^cKO* mice

compared with littermate controls at postnatal day 10 (**Fig.S7F**). This provides

further evidence of the relationship between *Grem1* expression and these genes.

Motor coordination and fear responses were impaired in *Grem1^cKO* mice

To assess the functional consequences of embryonic *Grem1* deletion in the mouse

forebrain, we undertook behavioural testing with *Grem1^cKO* mice in comparison with

Grem1^{flox/flox} littermate controls. We compared motor balance using the Rotarod test.

Latency to fall was significantly shorter in *Grem1^{cKO}* mice both for males and females,

suggesting an impaired motor balance with loss of *Grem1* (**Fig.5G**). During the open

field test used to assess exploratory behaviours, both male and female *Grem1^{cKO}*

mice spent significantly more time in the central area away from the walls than

littermate controls (**Fig.5G**). The total distance moved and velocity of movement were

similar between *Grem1^{cKO}* and littermate controls (data not shown). Behavioral

testing using an elevated plus maze also indicated that female *Grem1^{cKO}* mice spent

significantly more time in the open maze arms and entered into the open arms more

frequently than littermate controls (**Fig.5G**). Both of the exploratory behavior tests

indicate that conditional loss of *Grem1* leads to reduced anxiety-like behavior. Lastly,

we used the Y maze test, to assess the short-term memory of *Grem1^{cKO}* and

littermate controls. We observed no difference in the number of entries made to the

novel arm of the Y maze between groups (**Fig.S8**), suggesting that *Grem1* expression

is not required for short-term memory function.

Discussion

The spatiotemporal regulation of BMP signaling in brain development is poorly

understood. Here we focus on the expression and function of the BMP antagonist

Grem1 during brain development, using transgenic lineage tracing, gene expression,

in vitro culture and conditional knock out approaches.

In the developing mouse brain *Grem1* is first expressed at 13.5 dpc in the actively

dividing dorsal telencephalon cells that retain mitotic potential (**Fig.1E,F**). *Grem1-*

expression identifies neuroprogenitors committed to the NeuN+ layer Ⅴ and Tbr1+

VI neurons. Transcriptomic profiling using our mRNAseq data from embryonic

Grem1-reporter mouse brains and publicly available human developmental brain

scRNAseq data suggests that *Grem1/GREM1* expression is primarily associated with

markers of glutamatergic excitatory neuronal lineages, rather than GABAergic or

dopaminergic neurons or other cell types of the developing brain (**Fig.3B,E**,

Fig.S4A,B). Consistent with this, *Grem1* was not expressed in the ganglionic

eminences, from which GABAergic interneurons originate; nor were GABAergic

transcripts upregulated or enriched by GSEA in *Grem1*-expressing TdTomato$^+$ cells

in the dorsal telencephalon using bulk RNAseq analysis (**Fig.S4A, B**). Likewise

Grem1-lineage traced cortical neurons expressed the excitatory marker, Tbr1

(**Fig.3C**). *Grem1*-expression labels and generates important excitatory neuronal

lineages, i.e. pyramidal neurons, that play high-level cognitive functions in the

neocortex. As Grem1 expression is required for the correct development of early-born

excitatory neurons of the neocortex (**Fig.1G**), misregulation of BMP signaling likely

affects the balance between excitatory and inhibitory neuronal activities, which has

been implicated in neurodevelopmental disorders (Dani et al., 2005).

Our analysis of scRNAseq data from human midgestational cortex (22-23 weeks

post-conception) suggests that there is coordinated regulation of BMP agonists and

antagonists at this developmental stage in humans, but that there may be also be

functional redundancies (**Fig.S4D**). Noggin is the most extensively studied of the

BMP antagonists in the CNS, where it is a recognized neural inducer during

gastrulation (Lamb et al., 1993). Addition of recombinant Noggin to neural stem cells

promotes neuronal and suppresses astrocytic differentiation *in vitro* (Mikawa and

Sato, 2011). In our study, we confirmed that Grem1 acts as a BMP antagonist in

embryonic NSPC cultures, suppressing downstream BMP transcriptional targets

(**Fig.4,S6**). Similar to Noggin, we determined that Grem1 promotes proliferation and

neural differentiation, at the expense of astrocytic lineages *in vitro* (**Fig.4**). BMP

transcriptional targets were significantly downregulated in *Grem1*-expressing

TdTomato$^+$ cells also in vivo (**Fig.2B**) and *Grem1*$^+$ excitatory neurons in the human

embryonic cortex (**Fig.S5**). Together these results suggest that Grem1 expressing

cells are fated to become neurons via antagonism of potential paracrine BMP signals

(**Fig.S4D,S5**) (Furuta et al., 1997, Imura et al., 2008, Choe and Pleasure, 2018).

Our histological analyses identified that the cortex (particularly neocortical layers

Ⅴ and Ⅵ) of *Grem1^{cKO}* animals is significantly thinner, with less cells, in

comparison to littermate controls, with no obvious morphological changes in the

hippocampus. Our in vitro experiments confirmed that Grem1 drives proliferation

and neuronal differentiation in NSPC, thus this cortical phenotype is likely due to

reduced proliferation of embryonic NSPCs in *Grem1^{cKO}* mice. Layer Ⅴ neurons

target the spinal cord, cerebellum, striatum and the thalamus (Shipp, 2007) and play

roles in movement preparation, movement guidance, and the execution of well-timed

movements (Baker et al., 2018, Li et al., 2015). The observed decrease in layer Ⅴ

neurons would lead to impaired preparation for and coordination of movement in

Grem1^{cKO} animals and explain the motor balance defect in these animals in

comparison to litter mate controls. Emerging evidence suggests that layer Ⅵ

neurons play a central role in modulating thalamic and cortical neurons during

sensory processing (Wang et al., 2016). Our behavioural testing confirmed that loss

of Grem1 expression significantly impaired fear sensitivity in *Grem1^{cKO}* animals,

compared to littermate controls, consistent with a deficiency in layer Ⅵ neurons

with roles in sensory connections. BMP signaling is known to promote

synaptogenesis (Xiao et al., 2013, Shen et al., 2004) and altered synaptogenesis can

affect both fear sensitivity and motor abilities (Ha et al., 2016, Wood and Shepherd,

2010). Network analysis of DEG between *Grem1*-expressing TdTomato[+] and

TdTomato[-] cells from the developing mouse dorsal telencephalon identified a gene

cluster of inter-related transcripts in which the BMP target gene, *Id1*, acts as a hub.

Genes in the cluster play roles in neuron morphogenesis, synapse and axon

maturation (**Fig.2D,E**). For example, *Leucine-rich-repeat transmembrane neuronal*

proteins 3/4 (*Lrrtm3/4*) genes encode synapse organizing proteins critical for the

development and function of excitatory synapses (Um et al., 2016, Roppongi et al.,

2017, Linhoff et al., 2009). Likewise, the *Ryanodine receptor3* (*Ryr3*) gene encodes a

member of a family of receptors that shape synaptic transmissions(Balschun et al.,

1999) by amplifying spike-driven calcium signals in presynaptic terminals, and

consequently enhancing the efficacy of transmitter release (Futagi and Kitano, 2015).

In *Grem1cKO* animals, the mRNA expression of *Lrrtm3* and *Ryr3* was significantly

decreased in comparison to littermate controls (**Fig.S7F**). These results suggest

Grem1 may contribute to synapse formation and function by inhibiting BMP

signaling and expression of associated gene networks, the subject of further studies.

In summary, this is the first study to reveal the important function of Grem1 in

cortical development, particularly as a marker, and maker, of excitatory

neuroprogenitors. In the future, Grem1 may hold value beyond understanding the

cellular biology of brain development and function as we develop new approaches to

help tackle complex neurodevelopmental and neurological diseases.

Materials and Methods

Mice

Grem1-CreERT transgenic mice (Worthley et al., 2015) were crossed with *R26-*

LSLTdTomato mice (B6.Cg-*Gt(ROSA)26Sor*^{tm14(CAG-tdTomato)Hze}/J, JAX 007914) to

generate a tamoxifen-induced *Grem1* reporter line. Pregnant dams were

administered 6mg of tamoxifen by oral gavage to induce embryonic *Grem1*-tracing,

with some dams bearing 13.5 dpc embryos also intraperitoneally injected with 5-

Bromodeoxyuridine (BrdU, Roche, 40 mg/kg) to track cell division. *Grem1^{flox/flox}*

mice (Gazzerro et al., 2007) were crossed with *Emx1-cre* mice (B6.129S2-

Emx1^{tm1(cre)Krj}/J, JAX 005628) to generate *Emx1-cre* mediated *Grem1* conditional

knockout mice (*Emx1-cKO*). The line was maintained by crossing *Emx1*^{-cre/+} ;

Grem1 ^{flox/flox} mice with *Grem1^{flox/flox}* mice.

All mice were on the C57BL6/J background and experimentation was conducted

following approval by SAHMRI Animal Ethics Committee (approval number

SAM284) in accordance with the Australian code for the care and use of animals for

scientific purposes, 8th edition.

Preparation of single cell suspensions and flow cytometry

Pregnant *Grem1creERT; Rosa26LSLTdtomato* mice were administered tamoxifen at

13.5 dpc. At 14.5 dpc, dorsal telencephalons of the embryos were dissected in cold

phosphate-buffered saline (PBS) and the meningeal membranes were removed.

Neurocult Enzymatic Dissociation kit (Stemcell Technologies) was used for cell

dissociation according to the manufacturer's protocol. Dissociated cells were

resuspended in FACS buffer containing DAPI (0.5µg/ml). Sorting and analyses were

carried out on a FACS Fusion flow cytometer (Becton-Dickinson). Dead cells were

excluded by gating on forward and side scatter and by eliminating DAPI-positive

events. The cells harvested from *cre* negative littermate control mice were used to

set background fluorescence levels. Viable cells were sorted into FACS buffer,

collected via centrifugation and resuspended in Trizol (Invitrogen) for RNA

extraction.

RNA extraction and mRNA seq

RNA was extracted from sorted cells in Trizol according to the manufacturer's

protocol with the exception of addition of glycogen (20µg/µl) and isopropanol

precipitation overnight at -80℃ to maximize yield. RNA quality and quantity were

analyzed using a NanoDrop and TapeStation. Total RNA was converted to strand-

specific Illumina compatible sequencing libraries using the Nugen Universal Plus

mRNA-Seq library kit from Tecan (Mannedorf) as per the manufacturer's

instructions (MO1442 v2). Briefly, 500ng of total RNA was polyA selected and the

mRNA fragmented prior to reverse transcription and second strand cDNA synthesis

using dUTP. The resultant cDNA was end repaired before the ligation of Illumina-

compatible barcoded sequencing adapters. The cDNA libraries were strand selected

and PCR amplified for 12 cycles prior to assessment by Agilent Tapestation for

quality and Qubit fluorescence assay for quantity. Sequencing pools were generated

by mixing equimolar amounts of compatible sample libraries based on the Qubit

measurements. Sequencing of the library pool was performed with an Illumina

Nextseq 500 using single read 75bp (v2.0) sequencing chemistry.

Bioinformatic analysis

RNAseq data processing

Fastq files from the sequencing run were firstly subjected to quality controls with

FastQC version 0.11.3. Raw reads with low quality were removed using Trim Galore

alignment (Phred score less than 28 and/or reads contains adaptor sequences). After

trimming, all bases with low quality scores and adaptor sequences were removed.

The trimmed reads were mapped to Ensembl mouse genome (GRCm38) with STAR

2.4.2a. No more than 1 base mismatch was allowed. Only uniquely mapped reads

were retained. Option -quantMode was enabled to generate gene level quantification.

The counts files were then merged into an expression table for downstream

differential expression analysis.

Differential expression analysis

Differentially expressed genes (DEG) between the *Tdtomato*[+] (*Grem1* positive) and

Tdtomato[-] (*Grem1* negative) populations were analyzed with edgeR packages in R

version 3.6.0 (Robinson et al., 2010) . Reads were inspected through

multidimensional scaling plot and outliers were removed. TMM normalized counts

were log2-transformed and counts per million (CPM) obtained. Paired comparisons

between the *Tdtomato*[+] (*Grem1* positive) and *Tdtomato*[-] (*Grem1* negative)

populations were performed. DEG reported as significant were selected by requiring

both adjusted P value (FDR) $\leq$ 0.05 and absolute value of log2 fold change $\geq$ 2.0.

Supervised Weighted Gene Correlation Network Analysis (WGCNA)

Only DEGs were used for subsequent network analysis. The gene network was

constructed using the R package WGCNA following the procedure as described

(Langfelder and Horvath, 2008). After the low expression genes (FPKM < 1) had been

filtered out from all gene expression libraries, Pearson's correlation based adjacency

was calculated on the basis of pairwise correlations of gene expression within

Tdtomato⁺ (*Grem1* positive) samples. Topological overlap of the correlations was

used to weight the edges of the correlation network. The higher the weight, the

stronger the interaction between two genes. Connectivity for a single gene was

calculated as the sum of weights relative to the rest of the genes, and the top 5% of

genes with the highest connectivity in the network defined as hub genes. For

visualisation, a heat map was generated using the TOMplot() function in WGCNA R

package (version 1.68), with dissimilarity topological overlap (1 - topological overlap),

employed for hierarchical clustering. To generate the network plot, the weights of

the network were cut off at 0.1. Hypergeometric enrichment tests for each module-

defined gene list were performed with the enricher() function, as part of

clusterProfiler R package version 3.13.0 (Yu et al., 2012). Bonferroni adjustment for

p value was used, only gene sets with an adjusted p value less than 0.05 were

considered for interpretation of the biological function modules.

Gene set enrichment analysis (GSEA)

To utilize the Molecular signatures database (MSigDB), we accessed all gene sets

(.gmt file, version 6.2) from the Broad Institute and chose a subset for further

analyses including BioCarta, Hallmark, Gene Ontology (GO), Kyoto Encyclopedia of

Genes and Genomes (KEGG), Pathway Interaction Database, and the Reactome

Pathway Database. To enable cross species comparisons, mouse gene ensembl IDs

were converted to human orthologous gene symbols using biomaRt R package,

version 2.41.7 (Durinck et al., 2009). Humanised gene lists for the whole

transcriptome were ranked based on logfc values by comparing TdTomato$^+$ and

TdTomato- cells. The R package clusterProfiler version 3.13.0 (Yu et al., 2012) was

used to scan through all the gene sets mentioned above, using Benjamini & Hochberg

adjusted p values.

Analysis of publicly available single cell RNA seq dataset

Single cell RNAseq raw expression data was accessed for human mid-gestational

brain (22 and 23 weeks post-conception) cortex samples from GSE103723 (Fan et al.,

2018). The cells from two individual samples were grouped together, with the sum

of reads calculated for each gene to represent the data structure of bulky

transcriptomics. The limma package(Ritchie et al., 2015) was used to remove batch

effects with removeBatchEffect() and to generate multidimensional scaling plots. To

understand the expression pattern of human *GREM1* and its possible contribution

to neural differentiation, normalized counts and tSNE coordinates were employed

for visualization using plots generated with ggplot2 package in R environment

(Wickham, 2009). To investigate cell types expressing BMP antagonists, pearson's

Chi-Square test of independence was performed in the R environment with

chisq.test() function. Testing variables were gene expression categorised into high

and low by median expression, and using cell type identifiers from the original

scRNAseq study (Fan et al., 2018). Scaled standardised residuals for each gene were

used to plot the heatmap with ComplexHeatmap package (Gu et al., 2016).

In situ hybridization (ISH)

All ISH analyses were performed on formalin-fixed and paraffin-embedded mouse

tissue samples using RNAscope technology (RNAscope 2.5 HD Detection Kit,

RNAscope Multiplex Fluorescent Reagent Kit v2, Advanced Cell Diagnostics)

following the manufacturer's instructions. Briefly, tissue sections were baked in a

dry oven (HybEZ II Hybridization System, Advanced Cell Diagnostics) at 60°C for 1

h and deparaffinized, followed by incubation with an H_2O_2 solution (Pretreat 1

buffer) for 10 min at RT. Slides were boiled in a target retrieval solution (Pretreat 2

buffer) for 15 min, followed by incubation with a protease solution (Pretreat 3 buffer)

for 30 min at 40°C. Slides were incubated with a mouse _Grem1_ probe (NM_011824.4,

region 398-1359, catalogue number 314741) for 2 h at 40°C, followed by successive

incubations with signal amplification reagents. Staining was visualized with DAB

or TSA Plus Cyanine 5 (NEL745001KT, PerkinElmer;1:750). Combined ISH and

immunofluorescence (IF)/immunohistochemistry (IHC) was undertaken by first

performing ISH, followed by IF/IHC. For IF following ISH, the sections were blocked

with blocking buffer (X0909, Dako) and then incubated with a primary antibody

(rabbit polyclonal anti-RFP, Rockland, 600-401-379) overnight at 4℃. The sections

were washed in 1x T-PBS 3 times and then incubated with Alexa Fluor 555-

conjugated secondary antibody (Thermo Fisher Scientific) for 60 min at room

temperature. The sections were mounted with ProLong Gold antifade reagent

containing 4'6-diamidino-2-phenylindole (DAPI; Thermo Fisher Scientific), and

fluorescence was examined using an inverse immunofluorescence microscope BZ-

X710 (Keyence, Japan) with optical sectioning. For IHC following ISH, after

inactivation of endogenous alkaline phosphatase with alkaline phosphatase blocking

solution for 10 minutes (SP-6000, Vector Laboratories), the slides were blocked with

blocking buffer (X0909, Dako) and incubated with a primary antibody (rabbit

polyclonal anti-RFP, Rockland, 600-401-379) at 4 °C overnight. The sections were

washed with PBS three times, followed by incubation with a horse anti-rabbit IgG

polymer (MP-5401, Vector Laboratories) for 30 min at room temperature. The slides

were washed with TBS three times, and signal developed using an alkaline

phosphatase substrate (SK-5105, Vector Laboratories). The reaction was stopped by

washing in TBS four times, and slides were counterstained with hematoxylin,

dehydrated in EtOH, cleared in xylene, and coverslipped. Cells expressing more than

1 ISH signal were regarded as positive for Grem1 RNA.

Real time RT-PCR

RNeasy mini kits (Qiagen) were used to isolate RNA from snap frozen mouse brain

tissues. RNA was reverse-transcribed into cDNA with cDNA master (Sigma). PCR

amplification was performed with Kappa Sybr qPCR mix or Kappa Probe qPCR mix

using QuantStudio7 (Applied Biosystems). The mouse GAPDH gene was used as an

endogenous control. The following primers were used:

TaqMan probes and primers, Gremlin1 (IDT, Mm.PT.58.11631114), Id1

(IDT,Mm.PT.58.6622645.g), Id2 (IDT, Mm.PT.58.13116812.g), Id3 (IDT,

Mm.PT.58.29482466.g), Id4 (IDT, Mm.PT.58.6851535)

Sybr primers, Gapdh-forward AAGGTCATCCCAGAGCTGAA, Gapdh-reverse

CTGCTTCACCACCTTCTTGA Ryr3-forward TGCTGTCGCTTCCTTTGCTA, Ryr3-

reverse CATCGATGGGGACGCTAGAC, Lrrtm3-forward

TAGCAAATCAGGCTCCAGGG, Lrrtm3-reverse GAGTTCATGATGGACCCCACA

Immunohistochemistry

To collect 14.5 dpc, 17.5 dpc and 20.5 dpc embryonic brains, euthanized pregnant

dams immediately underwent cardiac perfusion with 4% PFA. Whole embryos at

14.5 dpc, or dissected embryonic brains for 17.5 and 20.5 dpc were fixed in 4% PFA

at 4℃ overnight. Embryos and brains were then cryoprotected in 30% sucrose and

frozen in OCT embedding medium. 16 μm sections were cut using a Leica CM1900

cryostat. To make paraffin sections, samples were post fixed in 10% neutral buffered

formalin overnight and processed. 5 μm sections were cut using a Leica HM325

microtome. Sections were blocked with Protein Block Serum-Free (Dako) for 1 hour

at room temperature, incubated overnight with first antibody at 4℃, secondary

antibody for 1 hour at room temperature, and coverslipped with Vectashield Antifade

Mounting Medium (Vector laboratories). The following antibodies were used: Anti-

TBR1(Abcam, ab183032, 1:400), Anti-NeuN (Abcam, ab104225, 1:500), Anti-β Ⅲ

tubulin (Sigma, T5076, 1:400), Anti-O4 (R & D systems, MAB1326, 1:500), Anti-

GFAP (Dako, Z0334, 1:250), Anti-FOXP2 (Abcam, ab16046, 1:10000), Anti-Ctip2

(Abcam, ab18465, 1:800), Anti-CDP (Santa Cruz, sc-13024, 1:400), Anti-BrdU

(Abcam, ab6326, 1:600), Anti-RFP (Rockland, 600-401-379, 1:1000), Anti-Sox2

(Millipore, AB5603, 1:400), Anti-Pax6 (Millipore, AB2237, 1:100)

Images were acquired on a Leica SP5 spectral scanning confocal microscope.

Neural stem cell and progenitor cell culture

Before seeding cells, tissue culture dishes were coated with PDL (100μg/ml) and laminin (10μg/ml). Embryonic neural stem and progenitor cells were isolated from pregnant mice at 14.5 dpc. Dorsal telencephalons were dissected from each embryo in PBS, meningeal membranes removed and the tissue triturated to a single cell suspension. Cells were cultured in NeuroCult™ Proliferation Medium containing 20 ng/ml epidermal growth factor (EGF) (Stemcell Technologies). For differentiation assays, cells were seeded onto 4 well chamber slides (Thermofisher #NUN177399) in Neurocult Differentiation medium (Stemcell Technologies). To induce recombination or overexpression of *Grem1*, cells collected from Grem1$^{flox/flox}$ embryos were infected with plenti-EF1-*Cre-2a-sfGFP-2a-puro*, plenti-EF1-*Grem1-2a-sfGFP-2a-puro,* or plenti-EF1-*2a-sfGFP-2a-puro* lentivirus, and transduced cells were selected using puromycin for 5 days and used for experiments before passage 5. Lentivirus plasmids, psPAX2 and MD2.G were transfected to 293T cells to generate lentivirus, and viral supernatant was concentrated using Amicon-Ultra 100k spin columns. To assess neurosphere forming ability, cells were seeded in 6-well uncoated plates (1.6x10^5 cells per well) and cultured for 5 days. The number of neurospheres sized 50-100 μm and >100 μm was counted. For cell viability assays, neural stem / progenitor cells were seeded in 96-well coated plates (1x10^4 cells per well) and RealTime-Glo™ MT Cell Viability Assay (Promega) was used with the continuous-read protocol at 0, 24, 52 and 72 hours. Recombinant human BMP2 (Prospec) was added to the differentiation medium.

Luciferase assay

BMP pathway activity was measured using the BMP response element luciferase

reporter pGL3 BRE luciferase (addgene #45126), internal control pRL/TK-luciferase

reporter and the dual luciferase reporter assay kit (Promega). Neural stem /

progenitor cells were seeded in coated 24-well plates (1.6×10^5 cells per well) and

transfected with the reporter plasmids using xTremeGENE HP DNA transfection

reagent (Roche). Cells were collected 48hr after transfection using Passive lysis

buffer (Promega). The pGL3 empty vector was used as a control for BMP-

independent changes in reporter activity.

Western blot

Cell lysates were solubilized with M-PER Mammalian Protein Extraction Reagent

containing complete protease and phosphatase inhibitors. Lysates were separated

by SDSPAGE and transferred to PVDF membranes (Millipore). After blocking with

5% nonfat skim milk in PBST for 30 min at room temperature, the membranes were

incubated overnight at 4℃ with anti-Gremlin1 antibody (R & D systems, AF956,

1:1000) or anti-ßActin (Santa Cruz, sc-47778, 1:1000) in 0.5% nonfat skim milk in

PBST. Membranes were washed with PBST and incubated with alkaline

phosphatase-conjugated secondary antibodies (anti-rabbit IgG and anti-mouse IgG,

GE healthcare life sciences, 1:10000) for 1 hr at room temperature. Finally, the blot

was visualized with Immobilon HRP substrate (Millipore) using a Chemi Doc XRS1

(Bio-Rad).

Behavioral tests

Mice were submitted to Rotarod, open field, elevated plus maze, and Y maze tests at

the age of 7-10 weeks. Mice while in their home cages were acclimatized to the

behavior suite for at least 30 min prior to testing. The data was acquired blindly to

the genotype.

Rotarod. Animals were placed on the rotarod (Panlab/Harvard Apparatus) that

linearly increased rotation speed from 4 to 40 rpm during a 120 second period. An

accelerating protocol was employed to eliminate the need for habituation to the

rotarod. This procedure was repeated for a total of three trials per mouse, separated

by 15min inter-trial intervals and the mean latency to fall from the rotarod in

seconds was compared between wild type littermate controls and *Grem1^{cKO}* mice to

assess motor coordination.

Open field test. The open field test was conducted in four identical square arenas

(50x50x50cm) surrounded by walls. Mice were individually placed in a corner of a

clean arena and allowed to explore for 10 min. For the purpose of data collection, the

arena was conceptually partitioned into two zones: a virtual center zone of 23x23 cm

and a peripheral zone occupying the remaining area. Lower percentages of time

spent in the center zone was used to indicate a higher level of anxiety.

Elevated plus maze test. The elevated plus maze consisted of a central square (8×8

cm) and four arms (29 cm long × 8 cm wide, two open arms with no railing and two

closed arms enclosed by a transverse wall 20 cm in height). A mouse was placed in

the center of the central square facing the open arm and allowed to explore the maze

apparatus for 10 min. The time spent in any of the open arms was recorded and used

as a measure of anxiety.

Y maze test. The apparatus consisted of three arms with an angle of 120° between

each of the two arms. Each arm was 40 cm long × 8 cm wide × 15 cm high. Visual

cues were placed on the walls of the mazes. The Y-maze test consisted of two trials

separated by an inter-trial interval (ITI) to assess spatial recognition memory. The

first trial (training) had a 5 min duration and allowed the mouse to explore only two

arms of the maze, with the third arm (novel arm) being blocked. After 30min ITI, the

second trial was conducted, during which all three arms were accessible for 5 min.

Trials were recorded using a ceiling-mounted camera and analyzed by a video

analyzer (Ethovision XT, Noldus) to determine the period that each mouse spent in

each arm of the maze.

Statistics

All statistical analyses were performed using Graphpad prism 8, with methods and

values summarised in each figure legend.

Acknowledgements

We thank Dr. Randall Grose (SAHMRI, Australia) for flow cytometry assistance; Dr

Mark van der Hoek and the Genomics Core (SAHMRI, Australia) for RNA-seq

library preparation and sequencing, Dr. Hiroaki Fujiwara (The Institute for Adult

Diseases, Asahi Life Foundation, Tokyo, Japan) for reviewing the manuscript;

Regeneron and Prof Andrew Zannettino for providing *Grem1$^{flox/flox}$* ; *Emx1-cre* mice.

Competing interests

The authors declare no competing interests.

Funding

MI was supported by Japan Society for the Promotion of Science (JSPS) Overseas

Research Fellowships and Overseas study grant of Kanzawa Medical Research

Foundation; NS by fellowship grants from the Astellas Foundation for Research on

Metabolic Disorders and the Uehara Memorial Foundation; DLW by a National

Health & Medical Research Council (NHMRC) Career Development Fellowship;

SLW by the Cancer Council SA Beat Cancer Project on behalf of its donors and the

State Government of South Australia through the Department of Health (MCF0418

to SLW), HK by JSPS Overseas Challenge Program for Young Researchers, Takeda

Science Foundation Fellowship and the Greaton International PhD Scholarship.

This study was supported by grants from the NHMRC (APP1140236 to SLW,

APP1099283 to DLW, APP1143414 to DLW/SLW), the KAKENHI Grant-in-Aid for

Scientific Research (19K17478 to NS) and the Kanae Foundation Asia-Oceania

Collaborative Research Grants to NS.

Data availability

729   The accession number for all sequencing data reported in this paper is GEO:

730   GSEXXXXXX.

Reference

733   ALI, I. H. A. & BRAZIL, D. P. 2014. Bone morphogenetic proteins and their
734   antagonists: current and emerging clinical uses. *British Journal of*
735   *Pharmacology,* 171**,** 3620-3632.
736   BACHILLER, D., KLINGENSMITH, J., KEMP, C., BELO, J. A., ANDERSON, R. M.,
737   MAY, S. R., MCMAHON, J. A., MCMAHON, A. P., HARLAND, R. M.,
738   ROSSANT, J. & ROBERTIS, D. E. M. 2000. The organizer factors Chordin
739   and Noggin are required for mouse forebrain development. *Nature,* 403**,** 658-
740   661.

BAKER, A., KALMBACH, B., MORISHIMA, M., KIM, J., JUAVINETT, A., LI, N. &
DEMBROW, N. 2018. Specialized Subpopulations of Deep-Layer Pyramidal
Neurons in the Neocortex: Bridging Cellular Properties to Functional
Consequences. *The Journal of neuroscience,* 38, 5441-5455.

BALSCHUN, D., WOLFER, D. P., BERTOCCHINI, F., BARONE, V., CONTI, A.,
ZUSCHRATTER, W., MISSIAEN, L., LIPP, H. P., FREY, U. J. &
SORRENTINO, V. 1999. Deletion of the ryanodine receptor type 3 (RyR3)
impairs forms of synaptic plasticity and spatial learning. *The EMBO Journal,*
18, 5264-5273.

CHOE, Y. & PLEASURE, S. J. 2018. Meningeal Bmps Regulate Cortical Layer
Formation. *Brain Plasticity,* 4, 169-183.

DANI, V. S., CHANG, Q., MAFFEI, A., TURRIGIANO, G. G., JAENISCH, R. &
NELSON, S. B. 2005. Reduced cortical activity due to a shift in the balance
between excitation and inhibition in a mouse model of Rett syndrome.
Proceedings of the National Academy of Sciences of the United States of
America, 102, 12560-12565.

DURINCK, S., SPELLMAN, P. T., BIRNEY, E. & HUBER, W. 2009. Mapping
identifiers for the integration of genomic datasets with the R/Bioconductor
package biomaRt. *Nature Protocols,* 4, 1184-1191.

DWYER, N. D., CHEN, B., CHOU, S.-J., HIPPENMEYER, S., NGUYEN, L. &
GHASHGHAEI, T. H. 2016. Neural Stem Cells to Cerebral Cortex: Emerging
Mechanisms Regulating Progenitor Behavior and Productivity. *The Journal*
of Neuroscience, 36, 11394-11401.

FAN, X., DONG, J., ZHONG, S., WEI, Y., WU, Q., YAN, L., YONG, J., SUN, L.,
WANG, X., ZHAO, Y., WANG, W., YAN, J., WANG, X., QIAO, J. & TANG, F.
2018. Spatial transcriptomic survey of human embryonic cerebral cortex by
single-cell RNA-seq analysis. *Cell Research,* 28, 730-745.

FU, C., LI, D., ZHANG, X., LIU, N., CHI, G. & JIN, X. 2018. LncRNA PVT1
Facilitates Tumorigenesis and Progression of Glioma via Regulation of MiR-
128-3p/GREM1 Axis and BMP Signaling Pathway. *Neurotherapeutics,* 15,
1139-1157.

FURUTA, Y., PISTON, D. W. & HOGAN, B. L. 1997. Bone morphogenetic proteins
(BMPs) as regulators of dorsal forebrain development. *Development,* 124,
2203-2212.

FUTAGI, D. & KITANO, K. 2015. Ryanodine-receptor-driven intracellular calcium
dynamics underlying spatial association of synaptic plasticity. *Journal of*

Computational Neuroscience, 39, 329-347.

GAZZERRO, E., SMERDEL-RAMOYA, A., ZANOTTI, S., STADMEYER, L., DURANT, D., ECONOMIDES, A. N. & CANALIS, E. 2007. Conditional Deletion of Gremlin Causes a Transient Increase in Bone Formation and Bone Mass. *Journal of Biological Chemistry,* 282, 31549-31557.

GOMES, W. A., MEHLER, M. F. & KESSLER, J. A. 2003. Transgenic overexpression of BMP4 increases astroglial and decreases oligodendroglial lineage commitment. *Developmental biology,* 255, 164-177.

GU, Z., EILS, R. & SCHLESNER, M. 2016. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics,* 32, 2847-2849.

GUAN, Y., CHENG, W., ZOU, C., WANG, T. & CAO, Z. 2017. Gremlin1 promotes carcinogenesis of glioma in vitro. *Clinical and Experimental Pharmacology and Physiology,* 44, 244-256.

HA, S., LEE, D., CHO, Y., CHUNG, C., YOO, Y.-E., KIM, J., LEE, J., KIM, W., KIM, H., BAE, Y., TANAKA-YAMAMOTO, K. & KIM, E. 2016. Cerebellar Shank2 Regulates Excitatory Synapse Density, Motor Coordination, and Specific Repetitive and Anxiety-Like Behaviors. *Journal of Neuroscience,* 36, 12129-12143.

HEVNER, R. F., SHI, L., JUSTICE, N., HSUEH, Y.-P., SHENG, M., SMIGA, S., BULFONE, A., GOFFINET, A. M., CAMPAGNONI, A. T. & RUBENSTEIN, J. 2001. Tbr1 Regulates Differentiation of the Preplate and Layer 6. *Neuron,* 29, 353-366.

HSU, D. R., ECONOMIDES, A. N., WANG, X., EIMON, P. M. & HARLAND, R. M. 1998. The Xenopus dorsalizing factor Gremlin identifies a novel family of secreted proteins that antagonize BMP activities. *Molecular cell,* 1, 673-683.

IMURA, T., TANE, K., TOYODA, N. & FUSHIKI, S. 2008. Endothelial cell‐derived bone morphogenetic proteins regulate glial differentiation of cortical progenitors. *European Journal of Neuroscience,* 27, 1596-1606.

KHOKHA, M. K., HSU, D., BRUNET, L. J., DIONNE, M. S. & HARLAND, R. M. 2003. Gremlin is the BMP antagonist required for maintenance of Shh and Fgf signals during limb patterning. *Nature genetics,* 34, 303-307.

KUMAR, S., REYNOLDS, K., JI, Y., GU, R., RAI, S. & ZHOU, C. J. 2019. Impaired neurodevelopmental pathways in autism spectrum disorder: a review of signaling mechanisms and crosstalk. *Journal of Neurodevelopmental*

812   *Disorders,* 11**,** 10.

813 LAMB, T. M., KNECHT, A. K., SMITH, W. C., STACHEL, S. E., ECONOMIDES, A.

814   N., STAHL, N., YANCOPOLOUS, G. D. & HARLAND, R. M. 1993. Neural

815   induction by the secreted polypeptide noggin. *Science (New York, N.Y.),* 262**,**

816   713-718.

817 LANGFELDER, P. & HORVATH, S. 2008. WGCNA: an R package for weighted

818   correlation network analysis. *BMC Bioinformatics,* 9**,** 559.

819 LEE‐HOEFLICH, S., CAUSING, C. G., PODKOWA, M., ZHAO, X., WRANA, J. L.

820   & ATTISANO, L. 2004. Activation of LIMK1 by binding to the BMP receptor,

821   BMPRII, regulates BMP‐dependent dendritogenesis. *The EMBO Journal,*

822   23**,** 4792-4801.

823 LI, N., CHEN, T.-W., GUO, Z. V., GERFEN, C. R. & SVOBODA, K. 2015. A motor

824   cortex circuit for motor planning and movement. *Nature,* 519**,** 51-56.

825 LIM, D. A., TRAMONTIN, A. D., TREVEJO, J. M., HERRERA, D. G., GARC A-

826   VERDUGO, J. M. & ALVAREZ-BUYLLA, A. 2000. Noggin antagonizes BMP

827   signaling to create a niche for adult neurogenesis. *Neuron,* 28**,** 713-726.

828 LINHOFF, M. W., LAUR N, J., CASSIDY, R. M., DOBIE, F. A., TAKAHASHI, H.,

829   NYGAARD, H. B., AIRAKSINEN, M. S., STRITTMATTER, S. M. & CRAIG,

830   A. 2009. An Unbiased Expression Screen for Synaptogenic Proteins Identifies

831   the LRRTM Protein Family as Synaptic Organizers. *Neuron,* 61**,** 734-749.

832 MARTYNOGA, B., DRECHSEL, D. & GUILLEMOT, F. 2012. Molecular Control of

833   Neurogenesis: A View from the Mammalian Cerebral Cortex. *Cold Spring*

834   *Harbor Perspectives in Biology,* 4.

835 MATHIEU, C., SII-FELICE, K., FOUCHET, P., ETIENNE, O., HATON, C.,

836   MABONDZO, A., BOUSSIN, F. D. & MOUTHON, M.-A. 2008. Endothelial

837   cell-derived bone mo rphogenetic proteins control proliferation of neural

838   stem/progenitor cells. *Molecular and Cellular Neuroscience,* 38**,** 569-577.

839 MATSUURA, I., ENDO, M., HATA, K., KUBO, T., YAMAGUCHI, A., SAEKI, N. &

840   YAMASHITA, T. 2007. BMP inhibits neurite growth by a mechanism

841   dependent on LIM-kinase. *Biochemical and Biophysical Research*

842   *Communications,* 360**,** 868-873.

843 MIKAWA, S. & SATO, K. 2011. Noggin expression in the adult rat brain.

844   *Neuroscience,* 184**,** 38-53.

845 MORI, H., NINOMIYA, K., KINO-OKA, M., SHOFUDA, T., ISLAM, M., YAMASAKI,

M., OKANO, H., TAYA, M. & KANEMURA, Y. 2006. Effect of neurosphere
size on the growth rate of human neural stem/progenitor cells. *Journal of*
Neuroscience Research, 84**,** 1682-1691.

REYNOLDS, B. A. & RIETZE, R. L. 2005. Neural stem cells and neurospheres—re-
evaluating the relationship. *Nature Methods,* 2**,** 333-336.

RITCHIE, M. E., PHIPSON, B., WU, D., HU, Y., LAW, C. W., SHI, W. & SMYTH, G.
K. 2015. limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Research,* 43.

ROBINSON, M. D., MCCARTHY, D. J. & SMYTH, G. K. 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics, 26**,** 139-140.

ROPPONGI, R. T., KARIMI, B. & SIDDIQUI, T. J. 2017. Role of LRRTMs in synapse
development and plasticity. *Neuroscience Research,* 116**,** 18-28.

S DERSTR M, S., BENGTSSON, H. & EBENDAL, T. 1996. Expression of
serine/threonine kinase receptors including the bone morphogenetic factor
type II receptor in the developing and adult rat brain. *Cell and tissue research,*
286**,** 269-279.

SAXENA, M., AGNIHOTRI, N. & SEN, J. 2018. Perturbation of canonical and non-
canonical BMP signaling affects migration, polarity and dendritogenesis of
mouse cortical neurons. *Development,* 145.

SHEN, W., FINNEGAN, S., LEIN, P., SULLIVAN, S., SLAUGHTER, M. & HIGGINS,
D. 2004. Bone morphogenetic proteins regulate ionotropic glutamate
receptors in human retina. *European Journal of Neuroscience,* 20, 2031-2037.

SHIPP, S. 2007. Structure and function of the cerebral cortex. *Current Biology,* 17.

SUN, M., THOMAS, M. J., HERDER, R., BOFENKAMP, L. M., SELLECK, S. B. &
O'CONNOR, M. B. 2007. Presynaptic Contributions of Chordin to
Hippocampal Plasticity and Spatial Learning. *The Journal of Neuroscience,*
27**,** 7740-7750.

SUN, Y. & XU, Z. 2010. Regulation of neural stem cell by bone morphogenetic protein
(BMP) signaling during brain development. *Frontiers in Biology,* 5**,** 380-385.

UM, J., CHOI, T.-Y., KANG, H., CHO, Y., CHOII, G., UVAROV, P., PARK, D., JEONG,
D., JEON, S., LEE, D., KIM, H., LEE, S.-H., BAE, Y.-C., CHOI, S.-Y.,
AIRAKSINEN, M. S. & KO, J. 2016. LRRTM3 Regulates Excitatory Synapse
Development through Alternative Splicing and Neurexin Binding. *Cell*
Reports, 14**,** 808-822.

WANG, W., ANDOLINA, I. M., LU, Y., JONES, H. E. & SILLITO, A. M. 2016. Focal

Gain Control of Thalamic Visual Receptive Fields by Layer 6 Corticothalamic
Feedback. *Cerebral Cortex.*
WICKHAM, H. 2009. *Ggplot2 : elegant graphics for data analysis,* New York,
Springer.
WILSON, S. W. & HOUART, C. 2004. Early Steps in the Development of the
Forebrain. *Developmental Cell,* 6**,** 167-181.
WOOD, L. & SHEPHERD, G. 2010. Synaptic circuit abnormalities of motor-frontal
layer 2/3 pyramidal neurons in a mutant mouse model of Rett syndrome.
Neurobiology of Disease, 38**,** 281-287.
WORTHLEY, D. L., CHURCHILL, M., COMPTON, J. T., TAILOR, Y., RAO, M., SI,
Y., LEVIN, D., SCHWARTZ, M. G., UYGUR, A., HAYAKAWA, Y., GROSS, S.,
RENZ, B. W., SETLIK, W., MARTINEZ, A. N., CHEN, X., NIZAMI, S., LEE,
H., KANG, P. H., CALDWELL, J.-M., ASFAHA, S., WESTPHALEN, B. C.,
GRAHAM, T., JIN, G., NAGAR, K., WANG, H., KHEIRBEK, M. A., KOLHE,
A., CARPENTER, J., GLAIRE, M., NAIR, A., RENDERS, S., MANIERI, N.,
MUTHUPALANI, S., FOX, J. G., REICHERT, M., GIRAUD, A. S., SCHWABE,
R. F., PRADERE, J.-P., WALTON, K., PRAKASH, A., GUMUCIO, D.,
RUSTGI, A. K., STAPPENBECK, T. S., FRIEDMAN, R. A., GERSHON, M.
D., SIMS, P., GRIKSCHEIT, T., LEE, F. Y., KARSENTY, G., MUKHERJEE,
S. & WANG, T. C. 2015. Gremlin 1 Identifies a Skeletal Stem Cell with Bone,
Cartilage, and Reticular Stromal Potential. *Cell,* 160**,** 269-284.
XIAO, L., MICHALSKI, N., KRONANDER, E., GJONI, E., GENOUD, C., KNOTT,
G. & SCHNEGGENBURGER, R. 2013. BMP signaling specifies the
development of a large and fast CNS synapse. *Nature Neuroscience,* 16**,** 856-
864.
YAN, K., WU, Q., YAN, D. H., LEE, C. H., RAHIM, N., TRITSCHLER, I.,
DEVECCHIO, J., KALADY, M. F., HJELMELAND, A. B. & RICH, J. N. 2014.
Glioma cancer stem cells secrete Gremlin1 to promote their maintenance
within the tumor hierarchy. *Genes & Development,* 28**,** 1085-1100.
YU, G., WANG, L.-G., HAN, Y. & HE, Q.-Y. 2012. clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. *OMICS: A Journal of*
Integrative Biology, 16**,** 284-287.
YUN, K., MANTANI, A., GAREL, S., RUBENSTEIN, J. & ISRAEL, M. A. 2004. Id4
regulates neural progenitor proliferation and differentiation in vivo.
Development, 131**,** 5441-5448.
917

Figure legends

Fig.1 *Grem1*-expressing cells give rise to cortical neurons in the developing mouse

brain

(A) Schematic showing tamoxifen and BrdU administration to *Grem1creERT;*

Rosa26LSLTdtomato (*Grem1*-reporter) mice.

(B) Representative images of TdTomato⁺ (red) cells in the telencephalon at 14.5dpc

(24h post-induction), 17.5dpc (3d post-induction), and 20.5dpc (6d post-induction)

in *Grem1*-reporter mice treated with tamoxifen at 13.5 dpc, DAPI (blue). Scale bar

= 200 μm.

(C) Representative images of immunohistochemical staining of boxed region from

(B) to visualize TdTomato⁺ cells (red) and *Grem1* mRNA by ISH (green) at 14.5dpc

(I, 24h post-induction) and 20.5dpc (III, 6d post-induction) in *Grem1*-reporter mice

treated with tamoxifen at 13.5 dpc, DAPI (blue). The boxed areas were further

magnified in adjacent panels. Scale bar = 50 μm.

(D) Quantification of (C) showing the percentage of TdTomato⁺ cells that were also

Grem1 RNA⁺ in 4 HPFs of 2-3 biological replicates, t-test.

(E)   Representative   images   of   immunofluorescence   staining   of   (II)   17.5dpc

telencephalon from *Grem1*-reporter (red) mice induced with tamoxifen at dpc13.5,

BrdU (green), DAPI (blue). Scale bar = 50 μm.

(F) Quantification of (E) showing the percentage of TdTomato⁺ cells that were also

BrdU⁺ 2 HPFs of 3 biological replicates, t-test.

(G) Representative images of immunofluorescence staining of 14.5 and 20.5 dpc

neocortex from *Grem1*-reporter (red) mice induced with tamoxifen at dpc13.5, NeuN

(green), DAPI (blue). Layer V and VI boxed in（ⅰ）and（ⅱ）, respectively. Scale bar =

50 μm. MZ, marginal zone ; CP, cortical plate ; SP, subplate

(H) Quantification of (G) showing the percentage of TdTomato[+] cells that were also

NeuN[+] in 3 HPF from 3 biological replicates. One way ANOVA with Tukey's multiple

test.

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

Fig.2 Grem1 expressing cells have neuron associated gene signatures

(A) TdTomato[+] and TdTomato[-] cells were isolated from 14.5 dpc brains of *Grem1-*

CreERT; R26-TdTomato mice induced with tamoxifen at 13.5dpc. Representative

FACS plot is shown, n=5 mice sorted for bulk RNAseq analysis.

(B) Volcano plot to show differentially expressed genes (DEG) between

TdTomato[+].and TdTomato[-] cells from (A). *Grem1* was significantly upregulated,

whereas BMP target genes, *Id1,3,4* were significantly downregulated in TdTomato[+]

cells compared to TdTomato[-] cells . Absolute value of log2 fold change ≥ 2.0,

FDR<0.05

(C) Network visualization of correlated DEG modules in TdTomato[+] cells. Each dot

represents a gene, dot perimeter colour indicates module membership (see Fig S3),

dot interior colour denotes upregulation (red) or downregulation (blue) of gene

transcript in TdTomato[+] cells compared to TdTomato[-], dot size indicates the

magnitude of gene expression correlation to neighboring genes, lines connecting dots

represent topological distance. The module containing highly correlated genes that

were predominantly upregulated in TdTomato[+] cells is boxed in magenta.

(D) Significantly enriched gene sets in magenta boxed module in (C). FDR<0.001

(E) Genes associated with Id1, a hub gene in the magenta boxed module in (C). Dot

perimeter colour indicates module membership (see Fig S3), dot interior colour

denotes upregulation (red) or downregulation (blue) of gene transcript in TdTomato$^+$

cells compared to TdTomato$^-$.

Fig.3 *Grem1* is expressed in the glutamatergic excitatory neuron lineage cells

(A) Heatmap depicting unsupervised clustering of TdTomato$^+$ and TdTomato$^-$ cells

isolated from 14.5dpc *Grem1*-reporter (red) mice induced with tamoxifen at dpc13.5

based on expression of representative differentiation marker transcripts for neural

stem cell (NSC), radial glia, intermediate progenitor, immature neuron, and mature

neuron.

(B) Gene set enrichment analysis (GSEA) for Glutamate Secretion genes between

TdTomato+ and TdTomato- samples from (A). Normalised enrichment score (NES)

= 2.16, p = 0.0049.

(C) Representative images of immunofluorescence staining of 14.5 and 20.5 dpc

telencephalon from *Grem1*-reporter (red) mice induced with tamoxifen at dpc13.5,

TBR1 (green), DAPI (blue). Scale bar = 100 μm.

(D) Quantification of (C) showing the percentage of TdTomato$^+$ cells that were also

TBR1$^+$ in 3 representative fields from 3 biological replicates.

(E) tSNE plot of human scRNAseq dataset. *GREM1* expressing cells outlined in black.

Dot size represents Grem1 expression value as indicated.

Fig.4 Gremlin1, a BMP antagonist, promotes proliferation and neuronal

differentiation in neural stem / progenitor cells *ex vivo*

(A) Detection of Grem1 protein levels in control *Grem1$^{flox/flox}$, Grem1$^{-/-}$* and *Grem1$^{O/E}$*

NSPC by western blot.

(B) BRE-reporter activity relative to internal control reporter. Data obtained from 4

independent experiments each performed in triplicate.

(C) BMP target gene, *Id1*, mRNA expression normalized to *Gapdh* in *Grem1^{flox/flox}*,

Grem1^{-/-} and *Grem1^{O/E}* NSPC treated with vehicle or rBMP2 for 24h. Columns, mean;

bars, SD. One way ANOVA with Tukey's multiple test. Results from 5 independent

experiments performed in triplicate.

(D) Representative images of *Grem1^{flox/flox}*, *Grem1^{-/-}* and *Grem1^{O/E}* neurosphere

cultures. Scale bar = 100 μm.

(E) Quantification of neurosphere number and size from (D) in 4 representative fields

from 4 independent experiments. One way ANOVA with Tukey's multiple test.

(F) Cell viability normalized to 0h timepoint for each culture. Results from 3

independent experiments each performed in triplicate and analyzed using linear

regression.

(G) Representative images of immunofluorescence staining to identify Tuj1+ neurons,

GFAP+ astrocytes and O4+ oligodendrocytes in differentiated cultures from wild type

NSPCs treated with vehicle or rGrem1.

(H) Quantification of (G), showing the percentage of marker+ cells in 3 representative

fields from 3 independent experiments. t-test. Scale bar = 100 μm.

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

1010

Fig.5 Cortical development, motor balance, and fear is impaired in Grem1

conditional knock out mice

Grem1 **is required for normal cortical development**

(A) Representative *Grem1* expression in the dorsal telencephalon of *Grem1^cKO* mice

and *Grem1^flox/flox* littermate controls by ISH at 14.5 dpc. Monochrome images were

prepared for quantitation of the *Grem1* ISH signal with ImageJ. n = 3

(B) *Grem1* mRNA levels normalized to *Gapdh* were determined using real time

qRT-PCR in mouse cortical brain samples collected at postnatal day 0 from *Grem1^cKO*

mice and littermate controls n=4 biological replicates. Columns, mean; bars, SD. t-

test. ****$p<0.0001$

(C) Representative histological images of cortical layers of *Grem1^cKO* mice and

Grem1^flox/flox littermate controls using Nissl at 10 weeks of age. Scale bar = 100 μm.

(D) Quantification of cortical layer thickness from (C) compared in 8 pairs of male

Grem1^cKO mice and littermate controls at 10 weeks of age. Female, n=7 control and

n=7 *Grem1^cKO* . t-test, *$p<0.05$.

(E) Representative images of immunofluorescence staining of cortex of *Grem1^cKO*

mice   and littermate controls at 10 weeks of age to visualise layer Ⅱ-Ⅳ with

marker CDP (green), layer Ⅴ and Ⅵ with marker Ctip2 (red), and DAPI (blue). n =

3 Scale bar = 100 μm.

(F) Representative images of immunohistochemical staining of cortex of *Grem1^cKO*

mice   and littermate controls at 10 weeks of age to visualise layer Ⅵ marker Foxp2

IHC (shown in red arrowheads). n = 3 Scale bar = 100 μm.

(G) Behavioral testing was performed to compare *Grem1^cKO* and *Grem1^flox/flox* mice

using age and sex matched littermates at 7-10 weeks of age. Columns, mean; bars,

SEM.

(i) Rotarod test latency to fall. Male, n=14 control and n=11 *Grem1^cKO* , Female, n=12

control and n=13 *Grem1^cKO* . t-test, *p<0.05.

（ⅱ）Open field test, cumulative duration spent in the center area. Male, n=13 control

and n=14 *Grem1^cKO* , Female, n=13 control and n=13 *Grem1^cKO* . t-test, *p<0.05.

（ⅲ，ⅳ）Elevated plus maze test.（ⅲ）Cumulative duration spent in open arms.（ⅳ）

The number of entries to open arms. Male, n=12 control and n=11 *Grem1^cKO* , Female,

n=14 control and n=14 *Grem1^cKO* . t-test, *p<0.05.

1043

Fig.S1 *Grem1* starts to express after 12.5 dpc and is dramatically decreased after

birth

(A) Representative coronal images of immunofluorescence staining of 12.5 dpc

telencephalon from *Grem1*-reporter (red) mice induced with tamoxifen at 11.5 dpc.

Scale bar = 100 μm. LV, lateral ventricle ; NCx, neocortex

(B) *Grem1* mRNA levels normalized to *Gapdh* were determined using real time qRT-

PCR in mouse cortical brain samples collected at p0, p10 and 4 weeks post-birth

(n=4-6 mice/group, PCR conducted in triplicate) Columns, mean; bars, SD. One way

ANOVA with Tukey's multiple test. ****p<0.0001

1053

1054   **Fig.S2 Clustering of differentially expressed genes**

1055   A heat map was generated to visualise the unsupervised hierarchical clustering of

1056   correlation scores for gene expression of DEG in TdTomato⁺ cells. Yellow colour

1057   indicates closely related and red not related. The color bar between the hierarchical

1058   clustering tree and plotting fields denotes the module membership of each gene.

1059

1060 **Fig.S3 *Grem1*-expressing cells are radial glial stem cell markers, Sox2 and Pax6**

1061 **negative**

1062 Representative images of immunofluorescence staining of 14.5 dpc neocortex from

1063 *Grem1*-reporter (red) mice induced with tamoxifen at dpc13.5, (A) Sox2 (green),

1064 DAPI (blue). (B) Pax6 (green), DAPI (blue). Scale bar = 100 μm.

1065

Fig.S4 Extended transcriptome analysis of bulk mRNAseq data from *Grem1*-

expressing TdTomato+ and TdTomato- telencephalon cells at 14.5dpc

(A) Gene set enrichment analysis (GSEA) for genes involved in the regulation of (i)

GABAergic synaptic transmissions and (ii) dopamine secretion that were

differentially regulated between *Grem1*-expressing TdTomato+ and TdTomato-

cells. Normalised enrichment score (NES) = -0.93, p = 0.75 and NES = -0.69, p =

0.87.

(B) Unsupervised clustering of *Grem1*-expressing TdTomato+ and TdTomato samples

based on expression of representative markers for glutamatergic, GABAergic, and

Dopaminergic neurons, in bulk mRNAseq data. Marker transcript expression is

displayed as a heat map with high expression in red to low expression in blue.

(C) Multidimensional scaling plot comparison of transcriptome of TdTomato+ and

TdTomato- cells isolated from 14.5dpc *Grem1*-reporter mice induced with tamoxifen

at dpc13.5 and *GREM1*+ and *GREM1*- cells from scRNAseq of human mid-gestational

cortex.

(D) GREM1 expression is associated with the excitatory neuronal lineage in the

developing human cortex. Unsupervised clustering of human developmental brain

cell populations from scRNAseq GSE103723 based on expression of lineage markers

and transcripts encoding BMP antagonists (*FST, NOG, CHRD, BMPER, GREM1,*

SOSTDC1), and BMP target genes *(ID1-4)*. Cell type is indicated by colour legend as

per tSNE plot shown in Figure 4E. Chi-square testing was used to evaluate the level

of each transcript in each cell population and the resulting standardized residual

values are depicted in the heat map.

1089

Fig.S5 Grem1 suppresses intrinsic BMP signaling in developing human excitatory

neurons, likely via antagonism of paracrine BMP ligands.

(A) Heat map depicting expression of BMP ligands in human developmental brain

cell populations from scRNAseq GSE103723. Cell types as per tSNE plot shown in

Figure 4E. Chi-square testing was used to evaluate the level of each transcript in

each cell population and the resulting standardized residual values are depicted in

the heat map.

(B) GREM1 expression inversely correlates with enrichment of BMP2,7 and BMP

target genes ID1, 3, 4 expression in human excitatory neurons in mid-gestation.

Expression of transcripts encoding BMP ligands antagonised by GREM1 and ID

target genes in *GREM1⁺* and *GREM1⁻* cells is displayed from human scRNAseq

GSE103723. One way ANOVA with Tukey's multiple test. ns $p > 0.05$, *$p < 0.05$,

**$p < 0.01$, ****$p < 0.0001$

1103

Fig.S6 Grem1 acts to antagonize BMP signaling in NSPCs resulting in altered

downstream transcript levels

Transcript levels of BMP target genes, *Id2*, *Id3* and *Id4*, normalized to *Gapdh* in

Grem1ᶠˡᵒˣ/ᶠˡᵒˣ, *Grem1⁻/⁻* and *Grem1�O/E* NSPC treated with vehicle or rBMP2 for 24h.

Results from 5 independent experiments performed in triplicate. Columns, mean;

bars, SD. One way ANOVA with Tukey's multiple test. *p<0.05, ***p<0.001

1110

Fig.S7 Extended morphological, immunostaining and transcript analysis of

***Grem1^{cKO}* mice and littermate controls**

(A) Representative coronal section of TdTomato$^+$ (red) cells in the dorsal

telencephalon at 14.5dpc in *Emx1-cre ; Rosa26LSLTdtomato* mouse brain. This cre-

driver is broadly active across the telencephalon region in which *Grem1* is highly

expressed. Scale bar = 100 μm. NCx, neocortex ; H, hippocampus ; Str, striatum ; T,

thalamus

(B) Quantification of cortical layer thickness compared between age and sex matched

littermates at 10 weeks of age. Female, n=7 control and n=7 *Grem1^{cKO}*,. t-test,

*p<0.05.

(C) Representative histological images of neocortical layer 5 and 6 from *Grem1^{cKO}*

mice and *Grem1$^{flox/flox}$* littermate controls at 10 weeks of age using Nissl. Scale bar =

20um.

(D) Quantification of (C) showing the cell number per area of layer Ⅴ and Ⅵ using

Nissl in 2 HPF of 3 biological replicates. Columns, mean; bars, SD. t-test. *p<0.05.

Scale bar = 20um.

(E) Representative histological images of hippocampus from *Grem1^{cKO}* mice and

Grem1$^{flox/flox}$ littermate controls at 10 weeks of age using Nissl. 8 pairs of males and

7 pairs of females were analyzed. Scale bar = 200 μm.

(F) Transcripts from the *Id1* associated gene cluster (Fig3E), *Ryr3* and *Lrrtm3* were

evaluated in *Grem1^{cKO}* and *Grem1$^{flox/flox}$* littermate control cortex at p10 using real

time qRT-PCR. *Ryr3* and *Lrrtm3* levels were normalized to *Gapdh*. n=4-6 biological

replicates. Columns, mean; bars, SD. t-test. *p<0.05, ****p<0.0001

1134

Fig.S8 Memory is not impaired in *Grem1^{cKO}* mice compared to *Grem1^{flox/flox}*

littermate controls

(A,B) Y maze test. Behavior was compared between age and sex matched *Grem1^{cKO}*

mice and *Grem1^{flox/flox}* littermate controls at 7-10 weeks of age. Male, n=12 control

and n=10 *Grem1^{cKO}*, Female, n=14 control and n=13 *Grem1^{cKO}*. t-test. (A)

Cumulative duration spent in new arm of Y maze. (B) The number of entries to new

arm of the Y maze. Columns, mean; bars, SEM.

A

B GO_GLUTAMATE_SECRETION

C 14.5dpc (24hours)   20.5dpc (7days)
TBR1 TdTomato DAPI

D

E