Interplay between medial nuclear stalling and lateral cellular flow underlies cochlear duct spiral morphogenesis

Mamoru Ishii[1], Tomoko Tateya[2], Michiyuki Matsuda[1,3], Tsuyoshi Hirashima*[3,4]

1. Laboratory of Bioimaging and Cell Signaling, Graduate School of Biostudies, Kyoto University
2. Department of Speech and Hearing Sciences and Disorders, Faculty of Health and Medical Sciences, Kyoto University of Advanced Science
3. Department of Pathology and Biology of Diseases, Graduate School of Medicine, Kyoto University
4. Japan Science and Technology Agency, PRESTO

Keywords: Differential growth, FRET imaging, Interkinetic nuclear migration, MAPK/ERK, Mathematical model, Morphogenesis, Spiral pattern formation

*Corresponding author: Tsuyoshi Hirashima, Ph.D

Graduate School of Biostudies, Kyoto University

Yoshida-Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan

Phone: +81-75-753-9450, Fax: +81-75-753-4655

hirashima.tsuyoshi2m@kyoto-u.ac.jp

Abstract

A notable example of spiral architecture in organs is the mammalian cochlear duct, where the

duct morphology is critical for hearing function. Molecular genetics has revealed the necessary

signaling molecules for the formation of spirals in organs, but it remains unclear how cellular

dynamics generate bending and coiling of the cochlear duct during development. Here we show

two modes of multicellular dynamics underlying the morphogenetic process by combining

deep tissue live-cell imaging, Förster resonance energy transfer (FRET)-based quantitation,

and mathematical modeling. First, surgical separation of the cochlear duct revealed that

bending forces reside primarily in the medial side of the duct. In the medial pseudostratified

epithelium, we found that nuclei stall at the luminal side during interkinetic nuclear migration,

which would cause the extension of the luminal side, thereby bending the duct. Second, long-

term organ-scale FRET imaging of extracellular signal-regulated kinase (ERK) activity showed

that helical ERK activation waves propagate from the duct tip concomitant with the reverse

multicellular flow in the lateral side of the duct, resulting in advection-based duct elongation.

We propose an interplay of distinct multicellular behaviors underpinning spiral morphogenesis

in the developing cochlear duct.

Introduction

Spiral shapes are a widely occurring motif in many varied biological tissues and organisms, including shells, horns, and plants, but how spiral shapes form has remained unclear (Thompson, 1942). The general principle of spiral formation is differences in growth rate between the outer and inner tissue of the extending organ, with the growth rate of the outer tissue being faster than that of the inner one, which has been theoretically and experimentally demonstrated in shells and plants (Johnson et al., 2019; Raup and Michelson, 1965; Smyth, 2016; Wada and Matsumoto, 2018). The cellular processes causing this differential tissue growth are different in each organ (Johnson et al., 2019; Saffer et al., 2017), and so identifying the organ-specific mechanisms underlying these differential tissue growth rates is crucial to understanding the developmental process of spiral morphogenesis.

An example of a spiral organ is the mammalian cochlear duct, which is a tonotopically organized auditory organ in the inner ear (Fig. 1A). During murine development, the cochlear duct, composed of epithelial cells, elongates, bends, and coils to form a spiral. The molecular basis for the morphogenesis of the cochlear duct has been the subject of several previous studies. Gene knockout studies have clarified that the elongation of the cochlear duct requires sonic hedgehog (SHH) signaling from the cochleovestibular ganglion in the conical central axis of the cochlea (Bok et al., 2013; Liu et al., 2010; Tateya et al., 2013), fibroblast growth factor (FGF) signaling of epithelial cells (Pauley et al., 2003; Pirvola et al., 2000; Urness et al., 2018, 2015), and non-canonical Wnt–planar cell polarity (PCP) signaling of prosensory cells (Mao et al., 2011; Montcouquiol and Kelley, 2020; Qian et al., 2007; Saburi et al., 2008; Wang et al., 2005). Deletion of *Shh* expression leads to a shortening of the cochlear duct and a significant decrease in cell proliferation exclusively at the base region (Bok et al., 2013). In *Fgf10* null mutant mice, the cochlear duct is remarkably shorter, but cell proliferation is unaffected (Urness et al., 2015), suggesting that cell proliferation and other cellular processes regulate ductal outgrowth. From embryonic day (E) 14.5 onwards, the mediolateral active migration of prosensory cells, during which these cells intercalate radially with their neighbors (known as convergent extension), contributes to longitudinal duct extension in a Wnt–PCP pathway-dependent manner (Chen et al., 2002; Cohen et al., 2019; Driver et al., 2017; Yamamoto et al., 2009). This cell intercalation drives ductal elongation; however, it cannot explain the asymmetrical morphogenetic mode underlying the duct bending before E14.5. Although the underlying signaling pathways are well characterized, the physical cellular mechanisms underlying spiral morphogenesis of the cochlear duct remain elusive. In the present study, we

aimed to identify the multicellular dynamics giving rise to the bending and coiling of the
developing cochlear duct using a combination of live cell-imaging, FRET quantitation, and
mathematical modeling.

Results

The medial epithelial layer causes cochlear duct bending

Mediolateral asymmetry characterizes the spiral form of the cochlear duct (Fig. 1A). We

therefore first examined the force balance between the medial and the lateral tissue by

surgically separating the cochlear duct along the lateral side of the hair cells (Fig. 1B); the

medial side included Kölliker's organ and the prosensory domain, and the lateral side included

the outer sulcus after separation. This manipulation led to the release of the mechanical stress

affected mutually between the medial and the lateral tissue, enabling us to inspect the internal

stresses of local tissues. The separated medial tissue curled much more after separation than

before, while the lateral tissue was relatively uncurled (Fig. 1C and Mov. 1), indicating that

active bending force are applied from the medial side, but not from the lateral side.

We next examined the morphology of the developing cochlear duct from E12.5 to E14.5

by staining an epithelial marker, E-cadherin, and performing organ-scale 3D imaging. During

this developmental period, the cochlear duct elongates and coils without changes to the

mediolateral width (Fig. 1D). The curvature and thickness of the epithelial layer are

significantly larger on the medial side than on the lateral side in a horizontal section of the roof-

floor axis (Fig. 1D). Hereafter, we refer to the epithelial layers on the medial and lateral sides

as the medial and lateral epithelial layers (MEL and LEL), respectively (Fig. 1E).

Morphological quantification revealed that the curvature and thickness of the MEL are highest

at a point more than 100 μm away from the tip of the duct along the MEL (Fig. 1F and S1). In

contrast, the curvature and thickness of the LEL remain constant (Fig. 1F and S1). These

observations prompted us to explore the structure and dynamics of the MEL at single cell

resolution.

Zooming in the epithelial layer clarified that most of the nuclei in the LEL align within a

few nucleus diameters of each other (~20 μm) (Fig. 1G, left). However, nuclei in the MEL

distribute broadly across a ~50 μm range, and mitotic cell rounding only occurs on the luminal

side of the MEL (Fig. 1G, right). We also found that nuclei that stained positive for the M phase

marker phospho-histone H3 (pHH3) were localized only on the luminal side of the epithelial

layers, while nuclei in the S phase cells labeled with a short pulse of ethynyl deoxyuridine
(EdU) were distributed predominantly on the basal side (Fig. 1H). Mosaic cell labeling
revealed that single cell bridges extended between the luminal and basal edges of the MEL via
long protrusions (Fig. 1I), clearly indicating that the MEL is a proliferative pseudostratified
epithelial tissue.

Figure 1 The pseudostratified medial epithelial layer curves in the cochlear duct
(A) Schematic diagrams showing the tissue axis and labels of the cochlear duct. (B)
Immunofluorescence image of anti-E-cadherin (green) and anti-Myosin VIIA, a marker for
sensory hair cells (magenta), with nuclear counterstaining using DAPI (blue) in the cochlea at
E17.5 during tissue separation, with cuts represented by dashed lines. N=3 was confirmed.
Scale bar, 300 μm. (C) Images of the cochlear duct before and after tissue separation.
Stereomicroscope images (upper) and the corresponding illustrations of the lateral (blue, left)
and medial (magenta, right) layers. N=3 was confirmed. Scale bar, 300 μm. (D)
Immunofluorescence images of anti-E-cadherin staining in the murine developing cochlea. The
lower rows are magnified images of the dotted squares in the upper rows. Yellow dotted lines
represent the edges of the epithelial layer. Scale bar, 200 μm. (E) Schematic diagram showing

regions used for morphological quantification. (F) Curvature and thickness as a function of the
arc length from the tip along the lateral epithelial layer (LEL, left) and the medial epithelial
layer (MEL, right) at E12.5. Mean ± standard deviation (s.d.) N=3. (G) Images of Lifeact-
EGFP (white) and H2B-mCherry (blue) showing cell shape and nuclear position in the LEL
and MEL. Yellow dotted lines represent the luminal and basal edges. Scale bar, 30 μm. (H)
Fluorescence labeling images of anti-pHH3 (green) and EdU (magenta) with nuclear
counterstaining using DAPI (blue) in the LEL and MEL. Yellow dotted lines represent the
luminal and basal edges. Scale bar, 30 μm. (I) Fluorescence image of GFP transfected cells
labelling single cells (green) and anti-E-cadherin (magenta) with nuclear counterstaining using
DAPI (blue) in the MEL (left) and the corresponding 3D rendered image (right). Yellow dotted
lines on the left represent the luminal and basal edges. Scale bar, 10 μm.
151
152

Luminal nuclear stalling promotes MEL bending

We next focused on the cellular dynamics in the MEL with a two-photon microscope. For live

imaging of the MEL, the outer cartilaginous shell that will develop into the bony labyrinth was

completely removed to expose the cochlear duct prior to *ex vivo* culture. We used a reporter

mouse line ubiquitously expressing fluorescence proteins localized in the cytoplasm, allowing

us to recognize the nuclei as the regions without fluorescence, as well as the luminal and basal

edges of MEL in the pseudostratified epithelium. We observed differences in the nuclear

movement between the flat region proximal to the tip of the apex and the curved region distal

to the tip of apex (Fig. 2A, Mov. 2). The nuclei moved to the luminal edge of the epithelial

layer before cell division, and after cell division the daughter nuclei returned to the basal side

in the flat region (Fig. 2B). In the curved region, however, the cell nuclei remained at the

luminal side even after cell division (Fig. 2C). To make this difference clearer, we tracked the

nuclei and quantified their distance from the luminal edge at 3 hours after nuclear division.

After classifying the tissue into two categories – flat and curved – based on the half-maximal

curvature (Fig. 2D), fewer nuclei move to the basal side in the curved region than in the flat

region (Fig. 2D'), suggesting that there are two different modes of luminal-basal nuclear

movement in the MEL, depending on the distance from the duct tip. In addition, we examined

two angles of cell division orientation, φ and θ, in the 3D polar coordinates from the live

imaging data, with φ and θ representing the zenith and azimuth angle, respectively (Fig. 2E).

The φ angle was between 0° to 45°, indicating that cell division mostly occurs in parallel to the

luminal surface of the MEL (Fig. 2F). The distribution of θ shows that most cells divide along

the apex-base axis rather than along the roof-floor axis (Fig. 2G), suggesting that the orientation

of cell division contributes to the apex-base local expansion of the luminal side of the MEL.

Luminal-basal nuclear movement in the flat region exhibits the behavior known as
interkinetic nuclear migration (IKNM), which has been observed in various pseudostratified
epithelia (Grosse et al., 2011; Kosodo et al., 2011; Meyer et al., 2011; Norden et al., 2009;
Sauer, 1935). During this process, nuclei at the basal side move close to the luminal surface
prior to mitosis, and the daughter nuclei move back to the basal side, which results in even
growth within the epithelial layer (Fig. 2H, upper). This mode of nuclear migration would not
necessarily result in bending of the epithelial tissue. In contrast, lack of basalward movement
during IKNM, which we refer to as 'luminal nuclear stalling' hereafter, results in asymmetric
one-way flux of nuclei from the basal side to the luminal surface of the layer, which results in
local cell crowding and expansion due to the division occurring at the luminal side (Fig. 2H,
lower). This asymmetrical IKNM, along with oriented cell division, may contribute to
differential growth between the luminal and basal sides of the MEL, and thus cause the physical
bending of the MEL.
To test this possibility, we built a cell-based mechanical model and examined whether
nuclear movements affect the curvature of an epithelial layer. Single epithelial cells in the layer
were represented as polygons with vertices consisting of the luminal and basal edges, which
could deform according to the cell cycle-dependent nuclear position, based on mechanical
interactions between neighboring cells (Fig. 2I). We then introduced a parameter γ
controlling the degree of basalward movement after IKNM – the nucleus moves to the basal
edge when $\gamma = 1$ and stays at the luminal side when $\gamma = 0$. Our mathematical simulation
demonstrated that the curvature of the epithelial layer monotonically decreases as a function of
γ (Fig. S2A), which supports our observation that luminal nuclear stalling mechanically
contributes to bending of the epithelial layer. We used an *in silico* experiment to predict that
cell cycle arrest at the S-phase entry would decrease the curvature of the MEL (Fig. 2J, 2K,
and Mov. 3). This prediction was validated by treating explant cochlea with the DNA synthesis
inhibitor Mitomycin C. After 1 day of treatment with 10 μM Mitomycin C, EdU-positive S
phase cells were no longer detected in the cochlea (Fig. S2B). Under these conditions, the
amount of curvature along the MEL was significantly decreased in the curved region (Fig. 2L
and 2M), which is consistent with the model prediction. These results corroborate our
hypothesis that luminal nuclear stalling after luminal mitosis promotes MEL bending.
206
207

Figure 2 Live imaging and mathematical modelling demonstrate that luminal nuclear stalling promotes MEL bending via luminal-basal differential growth

(A) Image of a section of an E14.5 cochlea in cytoplasmic reporter mice (CFP channel of ERK-FERT mice). The MEL was classified into flat (green) and curved (magenta) regions and the boundary between the flat and curved regions is indicated by a dotted line. The circle indicates the duct tip. Scale bar, 50 µm. (B, C) Kymographic images of the flat region (B) and the curved region (C). Dotted lines denote a change in position of manually marked nuclei over time. Scale bars, 20 µm. (D, D') For 16 nuclear divisions that could be tracked, the distance from the luminal edge 3 hours after nuclear division was plotted over the arc length from the tip (D') together with the MEL curvature (D). At half-maximal curvature (D), the samples in (D') were classified into two groups, i.e., flat (n=14) and curved (n=18). Mann–Whitney U test, p=0.00038. (E) Nuclear division orientation is represented by two angles, φ and θ, in the polar coordinate O'. (F, G) Angle distribution of φ and θ. n=16. Rayleigh test, p=0.00096 for (F) and p=0.15 for (G). (H) Schematics for the expected morphology depending on the mode of nuclear movement. IKNM would lead to the flat (upper) and asymmetric IKNM would lead to the curved (lower). (I) Virtual epithelial layer in the model simulation. The colored circles denote nuclei with the cell cycle state represented in the bottom. (J, K) Model prediction in the cell cycle arrest before S phase entry in the growing virtual MEL. Representative images (J) and the curvature over the arc length (K). Mean ± s.d. N=10. Mann–Whitney U test, p<0.001. Scale bar, 50 µm. (L, M) The effect of mitomycin C treatment. Representative images of resultant cochlear duct immunostaining of anti-E-cadherin (J) and the curvature over the arc length (M).

Mean ± s.d. N=3. Mann–Whitney U test, p=0.0017. Scale bar, 50 µm.

Spatial heterogeneity of cell proliferation suggests cellular inflow to the lateral side of the

apex to realize cochlear bending

One plausible mechanism for achieving the spiral morphology of the cochlear duct is MEL-

driven differential growth, but another possibility is that cell proliferation might occur at a

faster rate on the lateral side than on the medial side. To address this, we examined the spatial

distribution of proliferating cells within 30 min of labeling of EdU on the roof and floor side

of the cochlear duct (Fig. 3A and 3B). For quantification, the image domain of the cochlear

duct was divided into interrogation regions and the averaged fluorescence intensity of labelled

EdU was measured within each region (Fig. 3C). On the floor side, EdU-positive cells were

more abundant in the medial side than in the lateral side around the apex (arrows, Fig. 3B and

3D, left). However, on the roof side of the cochlear duct, EdU-positive cells were distributed

evenly, with slightly fewer cells at the base than at the apex and without a significant bias along

the mediolateral axis (Fig. 3B and 3D, right). These observations negate the possibility that cell

proliferation rates on the lateral side of the cochlear duct could drive mediolateral differential

tissue growth and cause duct bending. The spatial map of EdU intensity shows that cell

proliferation rates were higher in the floor-base region (arrowheads, Fig. 3B and 3D, left).

Supposing that cell proliferation is the main driver of local tissue growth, the higher volumetric

growth observed in the medial side than in the lateral side would contribute to the duct bending

inward at the lateral side, which contradicts the innate cochlear morphogenesis. We thus

hypothesized that the cells in the lateral side of the growing apex may be supplied by

proliferation 'hot spot' in the basal region of the cochlear duct, which could resolve the

observed mismatch between tissue growth rates in the medial and lateral sides of the duct.

Figure 3 Cell proliferation mapping suggests cellular inflow to the lateral side of the cochlear duct apex to realize cochlear bending

(A) Schematic of tissue axis of the cochlear duct. (B) Maximum projection images of stained anti-E-cadherin (white) and EdU (magenta) in the roof and floor region of cochlear duct at E12.5. Representative images showing EdU signals in the cochlear duct are shown. Arrows indicate the region of EdU signal gradient from the medial to the lateral side of the duct. Arrowheads indicate the base region where the EdU intensity is concentrated. Scale bar, 100 μm. (C) Regional mapping from irregular grids in the cochlear duct to regular grids. (D) Heat map of EdU intensity in the roof and floor region of the duct. Arrows and arrowheads are the same as in (B). N=3 was confirmed.

Retrograde helical ERK activation waves drive base-to-apex multicellular flow

How are cells supplied from the base of the cochlear duct to the lateral side? Earlier studies reported that FGF signaling is critical for cochlear duct outgrowth (Pirvola et al., 2000; Urness et al., 2015). We therefore focused on ERK MAP kinase, a downstream kinase in the FGF signaling pathway, and used a reporter mouse line that ubiquitously expresses a Förster resonance energy transfer (FRET)-based biosensor for ERK activity in the cytosol (Harvey et al., 2008; Komatsu et al., 2018, 2011). 3D FRET imaging using two-photon microscopy revealed that ERK is preferentially activated in the lateral-roof side of the cochlear duct, including the outer sulcus and stria vascularis (Fig. 4A, 4A', and Mov. 4), which is consistent with the previously reported distribution of *Fgfr2* expression (Urness et al., 2015).

For continuous observation during ductal outgrowth, we established an explant culture method in which the capsule above the duct tip was partially removed, allowing 3D organ-scale long-term imaging of ERK activity. Surprisingly, the time-lapse images of the cochlea

dissected at E12.5 revealed that ERK activation propagates intercellularly as oscillatory waves from the apex to the base of the floor side (Fig. 4B and Mov. 5), while ERK is constitutively activated around the ductal tip of the roof side (Fig. S3A and S3B). We next quantified multicellular tissue flow by particle image velocimetry (PIV) at the supra-cellular (4–5 cell length) scale, and found that cells coherently move as clusters of ~100 µm diameter from the base to the apex of the floor side, again as oscillatory waves, and similarly to the ERK activation waves of the floor side (Fig. 4C and Mov. 5). In the roof side, the multicellular tissue flows directly toward the direction of elongation around the duct tip (Fig. S3C). Kymography of ERK activity along the apex-base line of the lateral-floor side (Fig. 4D) shows oscillatory retrograde ERK activity waves (Fig. 4E and Fig. S3D), which proceed at a speed of 0.42 ± 0.078 µm min^{-1} (mean ± standard deviation) in space-fixed coordinates (Fig. 4F).

Since ERK activation can be induced by cell extension during collective cell migration (Hino et al., 2020), we further calculated the ERK activity rate, i.e., the time derivative of ERK activity, and the extension-shrinkage rate, i.e., the local tissue strain rate along the apex-base line, by using time-series data of ERK activity and PIV speed, respectively (Fig. S3E). We found that both the ERK activity rate and the extension-shrinkage rate oscillate across the time course (Figure 4G). Moreover, cross-correlation analysis revealed that the local tissue deformation, represented by the extension-shrinkage rate, precedes ERK activity changes by 24 min on average (Fig. 4H). This is consistent with a regime of mechano-chemical coupling for collective cell migration: extension-triggered ERK activation promotes cell contraction, pulling the neighboring cells, which eventually evokes another round of ERK activation in the neighboring cells (Boocock et al., 2020; Hino et al., 2020). The observed lag time between the ERK activity rate and the extension-shrinkage rate can be reproduced using a minimal mathematical model of the mechano-chemical coupling, but not by an uncoupling model under which the ERK activation waves regulate the cell deformation unidirectionally, supporting the plausibility of mechano-chemical coupling rather than an uncoupling regime (Fig. S4). The role played by ERK was confirmed by treating the cochlear duct with an inhibitor of the upstream kinase MEK, PD0325901. Treatment with the MEK inhibitor at 1 µM resulted in a significant decrease of the tissue flow speed (Fig. 4I) as well as ERK inactivation (Fig. S3F), corroborating that ERK activation waves contribute to the base-to-apex multicellular tissue flow.

Finally, we extended the analysis to 3D dynamics of ERK activity and cell movement in the developing cochlear duct. Surface rendering of the ERK activity map in the cytosol indicated that ERK activity peaks shift from the apex-roof to the base-floor in the lateral side

318   of the cochlear duct (Fig. 4J and Mov. 6). Concomitantly with helical ERK activity waves,

319   coherent cell movements can be observed from the base-floor to the apex-roof in the opposite

320   direction to the ERK waves (Mov. 6). This observation strongly suggests that ERK-mediated

321   helical collective cell movement could drive 3D duct coiling underlying the spiral

322   morphogenesis of the cochlear duct (Fig. 4K).

Figure 4 Retrograde helical ERK activation waves drive base-to-apex multicellular flow
(A) 3D ERK activity map in the cochlear duct at E12.5. (A') Cross section view (medial-lateral
and roof-floor plane) of (A). Orange dotted line indicates the floor plane shown in (B) and (C).
Scale bar, 50 µm. (B) Time-lapse snapshots of ERK activity maps in the floor plane. Time
indicates the elapsed time of live imaging. Yellow arrowheads indicate the ERK activity peak.
Scale bar, 100 µm. (C) Time-lapse snapshots of tissue flow speed obtained by PIV in the floor
plane. Scale bar, 100 µm. (D) Schematic diagram showing the axis, the apex-base line for
kymography, and regions of interests (ROIs). (E) Representative kymograph of ERK activity.
The horizontal axis indicates the position on the apex-base line shown in (D) and the vertical
axis indicates the elapsed time of live imaging. Dotted lines represent oscillatory waves from
the apex to the base. Scale bar, 100 µm. (F) ERK wave speed with mean and s.d. n=5 from

N=3. (G) Time-series ERK activity rate and extension-shrinkage rate in representative 3
different ROIs. (H) Cross-correlation between the extension-shrinkage rate and ERK activity
rate. n=12. Mean ± s.d. (I) Tissue flow speed before and after MEK inhibitor PD0325901
treatment at 1 μM. n=285 from N=3. Two-sample t-test, p<0.001. (J) Time-lapse snapshots of
surface rendered ERK activity maps in the cochlear duct at E12.5. The green corners
correspond to the green corner on the images shown in (B) and viewed from the left-bottom
corner of (B). Circles indicate the position of ERK activity peaks and the connecting dotted
lines indicate a trace of the peak shift. The time scale is the same as in (B). (K) Schematics for
the ERK activity waves and cell flow.

Discussion

Previous genetic studies have revealed the molecular basis of cochlear duct elongation during

development, but have been unable to explain the physical mechanisms by which the duct

bends because of its severe phenotype (Bok et al., 2013; Groves and Fekete, 2012; Urness et

al., 2018, 2015). Motivated by these earlier studies, we have visualized the cochlear duct

development by two-photon microscopy and found that nuclei moving from the basal side stall

at the luminal side after cell division, which gives rise to bending via differential tissue growth

between the luminal and basal sides of MEL. We also elucidated that coherent cellular flow

occurs from the base to the apex exclusively in the lateral wall of the growing cochlear duct,

which is accompanied by retrograde ERK activation waves.   Thus, our long-term deep tissue

imaging has illuminated unprecedented dynamics of cells and kinase activity underpinning the

bending of developing cochlear duct.

We showed that the mode of luminal-basal nuclear migration switches at a position in the

continuous pseudostratified epithelium (Fig. 2D'), which controls the curvature of the MEL

(Fig. 2J-M). To our knowledge, we have provided the first experimental evidence that nuclei

stall at the luminal side of the pseudostratified epithelium during IKNM in normal development

(Norden, 2017). The luminal nuclear stalling results in luminal expansion, to some extent,

because of physical space occupation even in the curved pseudostratified epithelium, while it

remains unclear whether the luminal nuclear stalling results from the MEL curvature, i.e., the

convexity of the luminal side. Therefore, we propose luminal expansion driven by luminal

nuclear stalling as a mechanism for sculpting curves in pseudostratified epithelium, in addition

to the actomyosin-based basal shrinkage that has been previously reported in zebrafish

neuroepithelium (Sidhaye and Norden, 2017; Yanakieva et al., 2019). One important remaining

question is what causes the space-dependent mode transition of nuclear movement in the

continuous tissue. It has been previously shown that knock-down of cell-surface TAG-1

(transient axonal glycoprotein-1) impedes basalward nuclear movement during IKNM in the

mouse ventricular zone, which ultimately leads to overcrowding of neural progenitor cells in

the luminal side, and a severe cortical dysplasia(Okamoto et al., 2013). It has been proposed

that cytoskeletal machinery, including actomyosin and microtubules, regulates basalward

nuclear movement (Kosodo et al., 2011; Norden, 2017; Norden et al., 2009). This provides an

important basis for the identification of the molecules responsible for the nuclear behavior in

the MEL at the supra-cellular level, which needs to be clarified in the future. In addition,

unknown factors providing these molecules with positional information should be explored to

understand the mode transition of nuclear movement at the tissue level.

In the present study, the establishment of long-term imaging techniques and biosensors

for protein kinase activity has led to the discovery of unexpected spatiotemporal patterns of

cell movement and ERK activity in the developing cochlear duct. Previously, we and others

have observed intercellular ERK activation waves in the epithelium, such as migrating Madin-

Darby canine kidney (MDCK) cells (Aoki et al., 2017; Hino et al., 2020), developing

drosophila tracheal placode (Ogura et al., 2018), and wounded murine skin (Hiratsuka et al.,

2015). We have also proposed an ERK-mediated mechanochemical feedback system, in which

cell extension activates ERK followed by ERK-triggered cell contraction (Boocock et al., 2020;

Hino et al., 2020), which can explain the symphony between cell movement and ERK activity.

The ERK activation wave speed in the developing cochlear duct was 0.42 µm min^{-1} (Fig.

4F), which is significantly slower than in MDCK cells and wounded mouse epidermis, where

it proceeds at 2.5 µm min^{-1} and 1.4 µm min^{-1}, respectively (Aoki et al., 2017; Hino et al., 2020;

Hiratsuka et al., 2015). Interestingly, when normalized to the cell lengths of the developing

cochlear duct (4 µm), MDCK cells (20 µm), and the basal cells of the mouse epidermis (10

µm), the wave speed of the developing murine cochlear duct, which is 6 cells hr^{-1}, is

comparable with that of MDCK cells, which is 7 cells hr^{-1}, and that of wounded adult murine

skin, which is 8 cells hr^{-1}. Our findings couple multicellular flow and ERK activation waves in

the cochlea, corroborating the findings of earlier studies (Boocock et al., 2020; Hino et al.,

2020), and supporting the existence of a general regulatory mechanism for collective cell

migration during tissue morphogenesis.

Our 3D time-lapse imaging revealed coherent helical cell flow from the base-floor to the

apex-roof of the lateral side of the cochlear duct. Cell flow analysis revealed that the rate of

base-to-apex cell flow (0.24 µm min^{-1}, Fig. S3E) exceeds that of the duct elongation speed

(0.13 µm min^{-1}, Mov. 5). Thus, the cell flow rate may be sufficient to compensate for the lateral

tissue growth. We speculate that this ERK-mediated cell advection originating from the

heterogeneity of cell proliferation causes consistent mediolateral differential growth at the

tissue scale, and results in cochlear bending. In support of this, knockout of *Shh* causes a

significant decrease in the number of proliferating cells at the base of the cochlear duct, and

causes the shortening of the cochlear duct (Bok et al., 2013). As SHH is secreted mainly from

the spiral ganglions located in the central axis of the cochlea (Bok et al., 2013), further

investigations on the interaction between the cochlear duct and these ganglions will provide a

better understanding of cochlear morphogenesis.

Overall, we visualized multicellular behavior underlying the bending of the cochlear duct

during development using deep tissue live imaging. This contributes to a better understanding

of symmetry breaking in tissue morphogenesis during development and in generation of inner

ear organoids (Koehler et al., 2017, 2013). The live imaging technique used in the present study

forms the basis for further analysis of the interplay between morphogenesis and cell fate

decisions during cochlear development (Cohen et al., 2019; Tateya et al., 2019).

Acknowledgements

This work was supported by the JSPS KAKENHI 17KT0107 and 19H00993, by the JST PRESTO JPMJPR1949 and CREST JPMJCR1654, and by the Medical Research Support Center of Kyoto University. We would like to thank Akane Kusumi for technical assistance, and Edouard Hannezo, Naoya Hino, and Yoshiko Takahashi for fruitful discussion.

Author contributions

Conceptualization, T.T., T.H.; Methodology, M.I., T.T.. T.H.; Software, M.I., T.H.; Validation, M.I., T.H.; Formal analysis, M.I., T.H.; Investigation, M.I., T.H.; Resources, M.I., M.M., T.H.; Data curation, M.I., T.H.; Writing - original draft, M.I., T.H.; Writing - review & editing, T.T., M.M., T.H.; Visualization, T.H.; Supervision, M.M., T.H.; Project administration, T.H.; Funding acquisition, M.M., T.H.

Competing interests

The authors declare no competing interests.

References

Abe T, Kiyonari H, Shioi G, Inoue KI, Nakao K, Aizawa S, Fujimori T. 2011. Establishment

of conditional reporter mouse lines at ROSA26 locus for live cell imaging. *Genesis*

49:579–590. doi:10.1002/dvg.20753

Aoki K, Kamioka Y, Matsuda M. 2013. Fluorescence resonance energy transfer imaging of

cell signaling from in vitro to in vivo: basis of biosensor construction, live imaging, and

image processing. *Dev Growth Differ* **55**:515–22. doi:10.1111/dgd.12039

Aoki K, Kondo Y, Naoki H, Hiratsuka T, Itoh RE, Matsuda M. 2017. Propagating Wave of

ERK Activation Orients Collective Cell Migration. *Dev Cell* **43**:305-317.e5.

doi:10.1016/j.devcel.2017.10.016

Bok J, Zenczak C, Hwang CH, Wu DK. 2013. Auditory ganglion source of Sonic hedgehog

regulates timing of cell cycle exit and differentiation of mammalian cochlear hair cells.

Proc Natl Acad Sci U S A. doi:10.1073/pnas.1222341110

Boocock D, Hino N, Ruzickova N, Hirashima T, Hannezo E. 2020. Theory of mechano-

chemical patterning and optimal migration in cell monolayers. *bioRxiv*.

doi:https://doi.org/10.1101/2020.05.15.096479

Chen P, Johnson JE, Zoghbi HY, Segil N. 2002. The role of Math1 in inner ear development:

Uncoupling the establishment of the sensory primordium from hair cell fate

determination. *Development*.

Cohen R, Amir-Zilberstein L, Hersch M, Woland S, Taiber S, Matsuzaki F, Bergmann S,

Avraham KB, Sprinzak D. 2019. Shear forces drive precise patterning of hair cells in the

mammalian inner ear. *bioRxiv*. doi:10.1101/707422

Driver EC, Northrop A, Kelley MW. 2017. Cell migration, intercalation and growth regulate

mammalian cochlear extension. *Dev*. doi:10.1242/dev.151761

Fletcher AG, Osterfield M, Baker RE, Shvartsman SY. 2014. Vertex models of epithelial

morphogenesis. *Biophys J* **106**:2291–2304. doi:10.1016/j.bpj.2013.11.4498

Grosse AS, Pressprich MF, Curley LB, Hamilton KL, Margolis B, Hildebrand JD, Gumucio

DL. 2011. Cell dynamics in fetal intestinal epithelium: Implications for intestinal growth

and morphogenesis. *Development*. doi:10.1242/dev.065789

Groves AK, Fekete DM. 2012. Shaping sound in space: The regulation of inner ear

patterning. *Development*. doi:10.1242/dev.067074

Harvey CD, Ehrhardt AG, Cellurale C, Zhong H, Yasuda R, Davis RJ, Svoboda K. 2008. A

genetically encoded fluorescent sensor of ERK activity. *Proc Natl Acad Sci U S A*.

doi:10.1073/pnas.0804598105

Hino N, Rossetti L, Marín-Llauradó A, Aoki K, Trepat X, Matsuda M, Hirashima T. 2020.

ERK-mediated mechanochemical waves direct collective cell polarization. *Dev Cell* **in**

press. doi:10.1101/2019.12.25.888552

Hirashima T, Adachi T. 2019. Polarized cellular mechano-response system for maintaining

radial size in developing epithelial tubes. *Development* **146**:dev181206.

doi:10.1242/dev.181206

Hirashima T, Adachi T. 2015. Procedures for the Quantification of Whole-Tissue

Immunofluorescence Images Obtained at Single-Cell Resolution during Murine Tubular

Organ Development. *PLoS One* **10**:e0135343. doi:10.1371/journal.pone.0135343

Hiratsuka T, Fujita Y, Naoki H, Aoki K, Kamioka Y, Matsuda M. 2015. Intercellular

propagation of extracellular signal-regulated kinase activation revealed by in vivo

imaging of mouse skin. *Elife* **2015**. doi:10.7554/eLife.05178

Johnson AB, Fogel NS, Lambert JD. 2019. Growth and morphogenesis of the gastropod

shell. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1816089116

Koehler KR, Mikosz AM, Molosh AI, Patel D, Hashino E. 2013. Generation of inner ear

sensory epithelia from pluripotent stem cells in 3D culture. *Nature*.

doi:10.1038/nature12298

Koehler KR, Nie J, Longworth-Mills E, Liu XP, Lee J, Holt JR, Hashino E. 2017. Generation

of inner ear organoids containing functional hair cells from human pluripotent stem

cells. *Nat Biotechnol*. doi:10.1038/nbt.3840

Komatsu N, Aoki K, Yamada M, Yukinaga H, Fujita Y, Kamioka Y, Matsuda M. 2011.

Development of an optimized backbone of FRET biosensors for kinases and GTPases.

Mol Biol Cell **22**:4647–56. doi:10.1091/mbc.E11-01-0072

Komatsu N, Terai K, Imanishi A, Kamioka Y, Sumiyama K, Jin T, Okada Y, Nagai T,

Matsuda M. 2018. A platform of BRET-FRET hybrid biosensors for optogenetics,

chemical screening, and in vivo imaging. *Sci Rep*. doi:10.1038/s41598-018-27174-x

Kosodo Y, Suetsugu T, Suda M, Mimori-Kiyosue Y, Toida K, Baba SA, Kimura A,

Matsuzaki F. 2011. Regulation of interkinetic nuclear migration by cell cycle-coupled

active and passive mechanisms in the developing brain. *EMBO J*.

doi:10.1038/emboj.2011.81

Liu Z, Owen T, Zhang L, Zuo J. 2010. Dynamic expression pattern of sonic hedgehog in

developing cochlear spiral ganglion neurons. *Dev Dyn*. doi:10.1002/dvdy.22302

Mao Y, Mulvaney J, Zakaria S, Yu T, Morgan KM, Allen S, Basson MA, Francis-West P,

Irvine KD. 2011. Characterization of a Dchs1 mutant mouse reveals requirements for

Dchs1-Fat4 signaling during mammalian development. *Development*.

doi:10.1242/dev.057166

Mao Y, Tournier AL, Hoppe A, Kester L, Thompson BJ, Tapon N. 2013. Differential

proliferation rates generate patterns of mechanical tension that orient tissue growth.

EMBO J **32**:2790–2803. doi:10.1038/emboj.2013.197

Meyer EJ, Ikmi A, Gibson MC. 2011. Interkinetic nuclear migration is a broadly conserved

feature of cell division in pseudostratified epithelia. *Curr Biol*.

doi:10.1016/j.cub.2011.02.002

Montcouquiol M, Kelley MW. 2020. Development and patterning of the cochlea: From

convergent extension to planar polarity. *Cold Spring Harb Perspect Med*.

doi:10.1101/cshperspect.a033266

Nagai T, Honda H. 2001. A dynamic cell model for the formation of epithelial tissues. *Philos

Mag Part B* **81**:699–719. doi:10.1080/13642810108205772

Norden C. 2017. Pseudostratified epithelia - cell biology, diversity and roles in organ

formation at a glance. *J Cell Sci*. doi:10.1242/jcs.192997

Norden C, Young S, Link BA, Harris WA. 2009. Actomyosin Is the Main Driver of

Interkinetic Nuclear Migration in the Retina. *Cell*. doi:10.1016/j.cell.2009.06.032

Ogura Y, Wen F-L, Sami MM, Shibata T, Hayashi S. 2018. A Switch-like Activation Relay

of EGFR-ERK Signaling Regulates a Wave of Cellular Contractility for Epithelial

Invagination. *Dev Cell* **46**:162-172.e5. doi:10.1016/j.devcel.2018.06.004

Okamoto M, Namba T, Shinoda T, Kondo T, Watanabe T, Inoue Y, Takeuchi K, Enomoto Y,

Ota K, Oda K, Wada Y, Sagou K, Saito K, Sakakibara A, Kawaguchi A, Nakajima K,

Adachi T, Fujimori T, Ueda M, Hayashi S, Kaibuchi K, Miyata T. 2013. TAG-1-

assisted progenitor elongation streamlines nuclear migration to optimize subapical

crowding. *Nat Neurosci*. doi:10.1038/nn.3525

Okuda S, Inoue Y, Eiraku M, Adachi T, Sasai Y. 2014. Vertex dynamics simulations of

viscosity-dependent deformation during tissue morphogenesis. *Biomech Model

Mechanobiol* **14**:413–425. doi:10.1007/s10237-014-0613-5

Pauley S, Wright TJ, Pirvola U, Ornitz D, Beisel K, Fritzsch B. 2003. Expression and

function of FGF10 in mammalian inner ear development. *Dev Dyn*.

doi:10.1002/dvdy.10297

Pirvola U, Spencer-Dene B, Xing-Qun L, Kettunen P, Thesleff I, Fritzsch B, Dickson C,

Ylikoski J. 2000. FGF/FGFR-2(IIIb) Signaling Is Essential for Inner Ear

542   Morphogenesis. *J Neurosci* **20**:6125–6134. doi:10.1523/JNEUROSCI.20-16-

543   06125.2000

544 Qian D, Jones C, Rzadzinska A, Mark S, Zhang X, Steel KP, Dai X, Chen P. 2007. Wnt5a

545   functions in planar cell polarity regulation in mice. *Dev Biol*.

546   doi:10.1016/j.ydbio.2007.03.011

547 Raup DM, Michelson A. 1965. Theoretical morphology of the coiled shell. *Science (80-)*.

548   doi:10.1126/science.147.3663.1294

549 Riedl J, Flynn KC, Raducanu A, Gärtner F, Beck G, Bösl M, Bradke F, Massberg S, Aszodi

550   A, Sixt M, Wedlich-Söldner R. 2010. Lifeact mice for studying F-actin dynamics. *Nat*

551   *Methods*. doi:10.1038/nmeth0310-168

552 Saburi S, Hester I, Fischer E, Pontoglio M, Eremina V, Gessler M, Quaggin SE, Harrison R,

553   Mount R, McNeill H. 2008. Loss of Fat4 disrupts PCP signaling and oriented cell

554   division and leads to cystic kidney disease. *Nat Genet*. doi:10.1038/ng.179

555 Saffer AM, Carpita NC, Irish VF. 2017. Rhamnose-Containing Cell Wall Polymers Suppress

556   Helical Plant Growth Independently of Microtubule Orientation. *Curr Biol*.

557   doi:10.1016/j.cub.2017.06.032

558 Sauer FC. 1935. Mitosis in the neural tube. *J Comp Neurol* **62**:377–405.

559   doi:10.1002/cne.900620207

560 Serra-Picamal X, Conte V, Vincent R, Anon E, Tambe DT, Bazellieres E, Butler JP,

561   Fredberg JJ, Trepat X. 2012. Mechanical waves during tissue expansion. *Nat Phys*

562   **8**:628–634. doi:10.1038/nphys2355

563 Sidhaye J, Norden C. 2017. Concerted action of neuroepithelial basal shrinkage and active

564   epithelial migration ensures efficient optic cup morphogenesis. *Elife*.

565   doi:10.7554/eLife.22689

566 Smyth DR. 2016. Helical growth in plant organs: Mechanisms and significance. *Dev*.

567   doi:10.1242/dev.134064

568 Tateya T, Imayoshi I, Tateya I, Hamaguchi K, Torii H, Ito J, Kageyama R. 2013. Hedgehog

569   signaling regulates prosensory cell properties during the basal-to-apical wave of hair cell

570   differentiation in the mammalian cochlea. *Dev*. doi:10.1242/dev.095398

571 Tateya T, Sakamoto S, Ishidate F, Hirashima T, Imayoshi I, Kageyama R. 2019. Three-

572   dimensional live imaging of Atoh1 reveals the dynamics of hair cell induction and

573   organization in the developing cochlea. *Dev*. doi:10.1242/dev.177881

574 Thompson DW. 1942. On Growth and Form, Revised ed. ed, On growth and form. Dover

575   Publications. doi:10.2307/2019330

Urness LD, Wang X, Doan H, Shumway N, Noyes CA, Gutierrez-Magana E, Lu R, Mansour
SL. 2018. Spatial and temporal inhibition of FGFR2b ligands reveals continuous
requirements and novel targets in mouse inner ear morphogenesis. *Dev.*
doi:10.1242/dev.170142
Urness LD, Wang X, Shibata S, Ohyama T, Mansour SL. 2015. Fgf10 is required for
specification of non-sensory regions of the cochlear epithelium. *Dev Biol.*
doi:10.1016/j.ydbio.2015.01.015
Wada H, Matsumoto D. 2018. Twisting Growth in Plant Roots In: Gitmann A, Grill J,
editors. Plant Biomechanics; From Structure to Function at Multiple Scales. Springer
International Publishing. pp. pp127-140.
Wang J, Mark S, Zhang X, Qian D, Yoo SJ, Radde-Gallwitz K, Zhang Y, Lin X, Collazo A,
Wynshaw-Boris A, Chen P. 2005. Regulation of polarized extension and planar cell
polarity in the cochlea by the vertebrate PCP pathway. *Nat Genet.* doi:10.1038/ng1622
Yamamoto N, Okano T, Ma X, Adelstein RS, Kelley MW. 2009. Myosin II regulates
extension, growth and patterning in the mammalian cochlear duct. *Development.*
doi:10.1242/dev.030718
Yanakieva I, Erzberger A, Matejčić M, Modes CD, Norden C. 2019. Cell and tissue
morphology determine actin-dependent nuclear migration mechanisms in neuroepithelia.
J Cell Biol. doi:10.1083/jcb.201901077
595
596

Materials and Methods

(1) Experiments

Animals

For FRET imaging, we used the transgenic mice that ubiquitously express an ERK biosensor with a long flexible linker (hyBRET-ERK-NLS) reported elsewhere(Harvey et al., 2008; Komatsu et al., 2018, 2011) . For simultaneous imaging of F-actin and nuclei, we crossed Lifeact-EGFP (Riedl et al., 2010) and R26-H2B-mCherry(Abe et al., 2011). Lifeact-EGFP mice were generously provided by Takashi Hiiragi from EMBL Heidelberg and R26-H2B-mCherry mice were provided from RIKEN Large (CDB0204K). Otherwise, we used ICR mice purchased from Japan SLC, Inc. We designated the midnight preceding the plug as embryonic day 0.0 (E0.0), and all mice were sacrificed by cervical dislocation to minimize suffering. All the animal experiments were approved by the local ethical committee for animal experimentation (MedKyo 19090 and 20081) and were performed in compliance with the guide for the care and use of laboratory animals at Kyoto University.

Antibodies

The following primary and secondary antibodies were used for immunofluorescence: anti-E-cadherin rat antibody (Cell Signaling Technology, #3195, 1:100 dilution), anti-Myosin-VIIa rabbit polyclonal antibody (Proteus BioSciences Inc., #25-6790, 1:200 dilution), anti-Histone H3 (phospho S28) rat polyclonal antibody (Abcam, #ab10543, 1:200 dilution), Alexa Fluor 546-conjugated goat anti-rat IgG (H+L) antibody (Thermo Fisher Scientific, #A11081, 1:1000 dilution), Alexa Fluor 647-conjugated goat anti-rabbit IgG (H+L) antibody (Abcam, #ab150079, or Thermo Fisher Scientific, #A21247, 1:1000 dilution).

Small-molecule inhibitors

The following chemicals were used: Mitomycin C (Nacalai Tesque, #20898-21) and PD0325901 (FUJIFILM Wako Pure Chemical Corporation, #162-25291).

Whole-tissue staining and imaging

The cochleae were gently freed from the capsule and the staining and clearing were performed according to an earlier study (Hirashima and Adachi, 2015). Briefly, the samples were fixed with 4% PFA in PBS overnight at 4°C and then blocked by incubation in 10% normal goat serum (Abcam, #ab156046) diluted in 0.1% Triton X-100/PBS (PBT) for 3 h at 37°C. The

samples were treated with primary antibodies overnight at 4°C, washed in 0.1% PBT, and

subsequently treated with secondary antibodies conjugated to either Alexa Fluor 546 or Alexa

Fluor 647 overnight at 4°C. For counter staining of nucleus, we mixed Hoechst 33342 (5 µg/ml,

Dojindo Molecular Technologies, #H342-10) or DAPI (Dojindo Molecular Technologies,

#D523-10, 1:200 dilution) . The samples were mounted with 10 µL of 1% agarose gel onto a

glass-based dish (Greiner Bio-One, #627871) for stable imaging. Then, the samples were

immersed with the BABB (benzyl-alcohol and benzyl-benzoate, 1:2, #04520-96 and # 04601-

65, Nacalai Tesque) solution or CUBIC-R+ (Tokyo Chemical Industry Co., # T3741) solution

for optical clearing. Images were acquired using the confocal laser scanning platform Leica

TCS SP8 equipped with the hybrid detector Leica HyD with the ×40 objective lens (NA = 1.3,

WD = 240 µm, HC PL APO CS2, Leica) and the Olympus FluoView FV1000 with the ×30

objective lens (NA = 1.05, WD = 0.8 mm, UPLSAPO30XS, Olympus).

EdU assay

For EdU incorporation to embryos, 200 µL of 5 mg/mL EdU in PBS was intraperitoneally

injected to pregnant mice 30 min prior to dissection. For the incorporation to dissected cochleae,

10 µM of EdU was treated into the samples 1 hour prior to the chemical fixation. Before EdU

detection, whole-tissue immunofluorescence of E-cadherin and counter nuclei staining with

DAPI were performed. EdU was detected using the Click-iT® EdU Imaging Kits (Thermo

Fisher Scientific, #C10340). The samples were optically cleared with CUBIC-R+ and images

were acquired by confocal microscopy as described above.

Mosaic cell labeling

DNA solutions of the pCAG-GFP vectors (0.7 µg/µL) in PBS with 0.1% Fast Green FCF

(Sigma-Aldrich, #F7252-5G) were injected into the lumen of an inner ear dissected from E13.5

embryos through a fine glass capillary tube under a stereo microscope (SZ61, Olympus). After

injection of DNA solution, the cochlea was sandwiched by a pair of tweezer type electrodes

(Nepa Gene Co., #CUY650P5) and the DNA was electrotransferred using the NEPA21 Super

Electroporator (Nepa Gene Co.). Three 5 msec square pulses (175 V, +) with a 10% decay rate

at intervals of 50 msec were applied as poring pulses, followed by three 50 msec square pulses

(10 V, +/–) with a 40% decay rate at intervals of 50 msec as transfer pulses.

Surgical separation

We dissected cochleae from embryos at E17.5 and manually separated a cochlear duct into the

medial and lateral side. The images in Movie 3 were acquired using the stereomicroscope

(SZX16, Olympus) with a cooled color CCD camera (DP73, Olympus).

Explant cultures

We cultured the dissected cochleae without removing the capsule otherwise noted. The

cochleae were mounted on a the 35 mm glass based dish (Iwaki, #3910-035) with 1 μL of

growth factor reduced Matrigel (Corning, #356231), and filled with 2 mL of a culture medium

including FluoroBrite DMEM Media (Thermo Fischer Scientific, #A1896701) with 1%

GlutaMAX (Thermo Fischer Scientific, #35050061) and 1% N2 Supplement with Transferrin

(Holo)(FUJIFILM Wako Pure Chemical Corporation, #141-08941). The samples were

incubated at 37°C under 5% CO_2.

Live imaging for explants

We used two different sample preparations for live imaging under ex vivo culture condition.

For imaging of MEL, we completely removed the capsule and put the uncovered cochlear duct

attached with spiral ganglion onto the glass-bottom dish whose apex side faces the glass. For

long-term organ-scale imaging, we partially cut off the capsule adjacent to the apex tip of

cochlear duct using tweezers carefully, and the semicircular canals were removed. The isolated

cochlea was put onto the dish as described above. The former method allows us to take images

MEL at single cell resolution but failed to achieve elongation and bending of cochlear duct. In

contrast, the latter method recapitulates the cochlear duct morphogenesis but was unable to

detect fluorescence signals in the medial side. For microscopy, we used an incubator-integrated

multiphoton fluorescence microscope system (LCV-MPE, Olympus) with a ×25 water-

immersion lens (NA=1.05, WD=2 mm, XLPLN25XWMP2, Olympus) and an inverted

microscope (FV1200MPE-IX83, Olympus) with a ×30 silicone-immersion lens (NA=1.05,

WD=0.8 mm, UPLSAPO30XS, Olympus). The excitation wavelengths were set to 840 nm,

930 nm, and 1040 nm each for CFP, Lifeact-EGFP, and R26-H2B-mCherry (InSight DeepSee,

Spectra-Physics). Imaging conditions for Lifeact-EGFP and R26-H2B-mCherry were as

follows – scan size: 800×800 pixels, scan speed: 4.0 μsec/pixel, IR cut filter: RDM690

(Olympus), dichroic mirrors: DM505 and DM570 (Olympus), and emission filters: BA495-

540 for EGFP and BA575-630 for mCherry (Olympus). Imaging conditions for the FRET

biosensor were as follows – scan size: 800 × 800 pixels, scan speed: 10 μsec/pixel, IR cut filter:

RDM690 (Olympus), dichroic mirrors: DM505 and DM570 (Olympus), and emission filters:

BA460-500 for CFP and BA520-560 for FRET detection (Olympus).

(2) Quantification and Analysis

FRET image analysis

703 Image processing for FRET measurement was described elsewhere (Aoki et al., 2013). Briefly,
704 the median filter of 3×3 window was processed to remove shot noises, and background signal
705 was subtracted each in FRET channel and CFP channel. Then, the ratio of FRET intensity to
706 the CFP intensity was calculated by a custom-made MATLAB (MathWorks) script.

Measurement of layer curvature and thickness

For 2D measurement of curvature and thickness, we first performed whole-mount
immunofluorescence of E-cadherin to visualize the cochlear epithelium and acquired z-stack
images by confocal microscopy as described above. Next, we manually traced the apical and
basal side of epithelial cells on the middle horizontal section of the roof-floor axis. The
extracted epithelial layer was named as the medial/lateral epithelial layer according to the side
based on a given apex tip point. Then, the curve of the epithelial layer was obtained by the
iterative skeletonization, and discrete points (x_i, y_i) were sampled along the curves at regular
intervals of 15 μm. Finally, fitting the discrete points with a cubic spline function, the function
S_i at an interval $[x_i, x_{i+1}]$ is denoted as

$$S_i(x) = a_i(x - x_i)^3 + b_i(x - x_i)^2 + c_i(x - x_i) + d_i.$$

Due to a definition of curvature $\kappa(x) = S''(1 + S'^2)^{-3/2}$, the curvature from the spline
function was calculated as

$$\kappa_i(x) = \frac{6a(x - x_i) + 2b}{(1 + \{3a(x - x_i)^2 + 2b(x - x_i) + c\}^2)^{3/2}}.$$

The curve of the convex/concave to the duct lumen was assigned as positive/negative in κ. We
defined the layer thickness as a linear length connecting to the luminal and basal edge, which
is vertical to the curve of the epithelial layer at sampling points.

Quantification of nuclear migration

We manually measured the center position of a daughter cell and its luminal and basal edge.

Based on the cross point where the luminal-basal line and center line of medial epithelial layer

are intersected, we calculated the arc length from the duct tip to the nuclear position using a

custom built MATLAB code.

Angle measurement of cell division orientation

First, we performed two-photon live imaging using FV1200MPE-IX83 as described above,

and obtained volumetric images with a depth interval of 3 μm. In each mitotic event, we then

manually measured position (x, y, z) of the two daughter cells 3 hours later after the mitosis in

the fixed coordinate system O of the 3D image. Also, we manually acquired the position of the

luminal and basal edge of the mother cell to define a position (x', y', z') in a local coordinate

system, O′, the origin of which is the middle point of the two daughter nuclei. Each orthogonal

basis of the O′ coordinate was defined as follows: the apex-base axis in the MEL (x′), the

surface normal of the MEL (y′), i.e., luminal-basal axis in the MEL, and roof-floor axis (z′)

orthogonal to both x′ and y′ according to the right-handed system. With this local coordinate

system, we finally calculated two angles φ and θ in the local sphere coordinate resulting from

coordinate transformation from the O into the O′ system.

EdU intensity mapping

First, we separated 8-bit image stacks into two, roof and floor, based on z position at the middle

point, and performed maximum intensity projection onto xy plane both for staining images for

E-cadherin and EdU. Next, we binarized immunostaining signals for E-cadherin using Otsu's

method with morphological operations and detected periphery of the cochlear duct with the

MATLAB function 'bwperim'. With three reference points given manually, i.e., a) the duct tip,

b) the end point of MEL and c) that of LEL, the MEL and LEL curve was each defined by

connecting points between a) and b) and those between a) and c) along the duct periphery.

Then, we marked points to make 20 bins at a constant distance each along the MEL curve and

LEL curve. By connecting the marked point of MEL and that of LEL indexed by an order from

the duct tip and dividing the lines into 10, we partitioned the cochlear duct into small regions

for the measurement. Finally, we measured the averaged intensity of EdU signal within each

region and normalized by 255.

Tissue flow and ERK activity

To calculate velocity fields of cells in cochleae, we performed particle image velocimetry

(PIV)-based image processing using a free code MatPIV (a GNU public license software

distributed by Prof. Kristian Sveen in University of Oslo) was applied to time-lapse images of

the CFP channel. Velocity fields at time T was computed by displacement between T and $T+\Delta t$.

Δt was set as the sampling rate, 12 min. The size of the interrogation window was set to 40

pixels, approximately 25 μm, corresponding to 4-5 cell diameter, and the window overlap was

set to 50%. The obtained velocity data were then smoothened via median filtering to eliminate

peaky noises. We then obtained the 'tissue flow speed for the elongation' from PIV velocity

vector projected onto the apex-base line depicted in Figure 4D, and calculated the spatial

derivative of the tissue flow speed for the elongation between two adjacent interrogation

windows on the apex-base line according to the definition of a diagonal component of the strain

rate. This quantity was smoothened using the MATLAB function 'smooth' to eliminate high

frequency components and defined as the extension-shrinkage rate. As for the ERK activity,

we set thresholds in CFP images using Otsu's method within each interrogation window to

extract cytoplasmic region and calculated the mean FRET/CFP ratio in the binarized region.

The ERK activity rate was calculated as the time derivative of the ERK activity. Cross-

correlation analysis was performed using the MATLAB function 'xcorr'.

Statistical analysis

The number of cells or region of interests analyzed (n) and the number of biological replicates

(N) are indicated in the figure legends. No particular statistical method was used to

predetermine the sample size. A minimum of N=3 independent experiments were performed

based on previous studies in the field. No inclusion/exclusion criteria were used and all

analyzed samples were included in the analysis. No randomization was performed. Statistical

tests, sample sizes, test statistics, and P-values were described in the main text. P-values of less

than 0.05 were considered to be statistically significant in two-tailed tests, and were classified

as 4 categories; * ($P<0.05$), ** ($P<0.01$), *** ($P<0.001$), and n.s. (not significant, i.e., $P \geq$

0.05).

Software

For digital image processing, we used MATLAB (MathWorks) and Image J (National Institute

of Health). For graphics, we used MATLAB (MathWorks), Imaris (Bitplane) and ImageJ

(National Institute of Health). For statistical analysis, we used MATLAB (MathWorks).

(3) Mathematical model

A-1. A mechanical model for MEL morphogenesis

We modeled multicellular dynamics within an epithelial layer using a vertex dynamics model

(VDM) (Fletcher et al., 2014; Hirashima and Adachi, 2019; Nagai and Honda, 2001). Here we

focused on two-dimensional section on apex-base and roof-floor axes as shown in Fig. 2A. In

general, the 2D VDM model represents a single cell as a polygon, of which vertices are

elementary points that constitute the cell shape, and a group of cells is regarded as a set of

polygons shared by neighboring cells – we put four vertices, two of which are regarded as an

luminal edge and the other two of which are as a basal edge.

In the VDM, the dynamics of position of vertex i, r_i, obey the equation of motion based

on the principle of least potential energy U as follows:

$$\eta(\dot{r}_i - v_i) = -\nabla_i U,$$ Eq. 1

where η is a viscosity coefficient. v_i is a local velocity of vertex i, defined as $\langle v_i \rangle = \langle \dot{r}_{j_i} \rangle_{j_i}$,

where j_i is an index of cells contacting to vertex i, r_{j_i} is a centroid of cell j_i, and $\langle \, \rangle_{j_i}$

denotes averaging in j_i (Hirashima and Adachi, 2019; Mao et al., 2013; Okuda et al., 2014).

For a potential energy as a minimum expression, we defined as

$$U = \sum_j \left\{ \frac{k_a}{2}\left(a_j - a_j^*\right)^2 + \frac{k_b}{2}\left(b_j - b_j^*\right)^2 + \frac{k_l}{2}\left(l_j - l_j^*\right)^2 + \frac{k_A}{2}\left(A_j - A_j^*\right)^2 \right\} + \sum_i \frac{k_\theta}{2}\theta_i^{\,2}.$$

818   Eq. 2

The first to third terms each represent a regulation of cell edge length in luminal/apical, basal,

and lateral side with controlling parameters (k_a, k_b, k_l), current length (a_j, b_j, l_j), and the

target length (a_j^*, b_j^*, l_j^*). Target edge length of the luminal/apical and that of basal side are a

function of nuclear position within the epithelial layer, described later. The fourth term

represents cell area preservation with its coefficient k_A, current cell area A_j, and the target cell

area A_j^*. The fifth term represents bending energy of the luminal/apical side and that of basal

side attributed to each vertex; k_θ denotes the bending rigidity and θ_i is an angle of

luminal/apical edge or that of basal edge at a vertex i. Although the cell shape is relatively

flexible in a pseudostratified epithelial tissue, this model framework is valid as several

biological features can be incorporated, such as the nuclear position-dependent edge regulation

and cell size preservation.

Each cell has a unique cell cycle, length of which τ_{div} is assumed to be equally assigned

to all cells. We assume that a period of each cell cycle phase (G1, S, G2, and M) is partitioned

833 as 11:8:4:1, and a cell cycle status or timer τ_j, origin of which is defined at the end of cell

834 division, i.e., a boundary of M-G1, is provided according to the distribution of cell cycle phase.

835 We model that the cell timer τ_j in the cycle determines the distance from the luminal side of

836 a cell, i.e., nuclear position d_j $(0 \le d_j \le 1)$, with a parameter γ as follows:

838
$$d_j = \begin{cases} 2\gamma\tau_j/\tau_{div} & (0 < \tau_j \le 0.5\tau_{div}) \\ (-\tau_j/\tau_{div} + 0.95)\gamma/0.45 & (0.5\tau_{div} < \tau_j \le 0.95\tau_{div}) \\ 0 & (0.95\tau_{div} < \tau_j \le \tau_{div}). \end{cases} \qquad \text{Eq. 3}$$

840 This means that nuclei move toward the basal side of cell during G1-S phase with a degree γ

841 and get close to the luminal side before entering M phase. Of note, γ controls the degree of

842 basalward movement as described in the main text. Once a cell divides, τ_j in one of daughter

843 cells reset to zero and one in the other is set to a value stochastically chosen from 0 to $0.1\tau_{div}$

844 with a uniform distribution to avoid perfect synchronization between neighboring cells.

845  Occupied cell area is dominant in the medial epithelial layer and the cell position along

846 the luminal-basal axis should regulate the length of luminal edge and that of basal edge. Thus,

847 we define the target length as the simplest linear function of the nuclear position with two

848 parameters ξ_{max} and ξ_{min} as follows:

850 $a_j^* = (\xi_{min} - \xi_{max})d_j + \xi_{max},$

851 $b_j^* = (\xi_{max} - \xi_{min})d_j + \xi_{min}.$         Eq. 4

854 **A-2. Numerical simulation**

855 The ordinary differential equations were numerically solved by the forward Euler method with

856 time step 0.01. The code was generated with MATLAB (MathWorks). Regarding initial

857 condition, 20 rectangular cells, width/height of each which is 2.5/50 μm, are arrayed along a

858 horizontal line. Standard parameter set is $\eta = 1$, $k_A = 0.01$, $a_j^* = 125$ [μm^2], $k_a = 1$,

859 $k_b = 1$, $k_l = 1$, $k_\theta = 30$, $l_j^* = 50$ [μm], $\tau_{div} = 432$, $\gamma = 0.9$, $\xi_{max} = 5$ [μm], $\xi_{min} =$

860 1 [μm], otherwise noted. a_j^*, l_j^*, γ, ξ_{max}, and ξ_{min} were determined from obtained images,

861 and other parameters were determined empirically with some observations; at first, η, k_a, k_b,

862 and k_l were set to 1 and then k_A, τ_{div}, and k_θ were determined to reproduce the MEL

863 curvature observed in the curvature experimentally. Note that relative values rather than

absolute ones are critical for the dynamics.

A-3. Detailed setting in virtual experiments for cell cycle arrest

For an investigation towards mimicking the developmental process, we introduced luminal

stalling region, defined with an arclength along the epithelial layer L originated from the

boundary of apex side. Assuming once the cells in the luminal stalling region undergo division,

the daughter cells' timer τ_j do not count up, meaning that those nuclei stay at the luminal side

without reentering the cell cycle. This is due to experimental observations in which daughter

cells do not undergo mitosis exclusively in the curved region within a short time scale, at least

our observation window. For the cells in the non-luminal stalling region, τ_{div} was set

stochastically chosen from a uniform distribution from 216 to 864 each after the cell division

to incorporate variability and asynchronicity in nuclear dynamics. To recapitulate the

mitomycin C treatment in simulation, the cell cycle was arrested if the cell timer τ_j was ranged

in $0.25\tau_{div}$ to $0.8\tau_{div}$, corresponding to the mid G1 phase to the end of S phase, when the

simulation time exceeded 700. For the control case, we evaluated the curvature of virtual MEL

when the cell number reached at 100. For the case of mitomycin C treatment, we evaluated

when the simulation time reached at 1300 because the averaged simulation time in the control

case was 1298 (n=10).

B-1. Modeling oscillatory ERK activation waves and cell flows

We built a minimal 1D mechano-chemical coupling model for the collective cell migration

based on our previous studies (Boocock et al., 2020; Hino et al., 2020). Cells, each indexed as

$j=1,...,N$, are represented as a chain of springs, whose junctions including the boundaries are

labeled as $i=1,...,N+1$, with elastic constant k and each cell generates contractile force at the

rear side of the cell to move to the front with a force F. *i.e.*, the cell contractile force with $j=N$

($F_{j=N}$) is regarded as the force at the junction $i=N$ ($F_{i=N}$). Because the epithelial cells adhere to

neighboring cells and thus transmit the elastic force with viscous frictions $\boldsymbol{\eta_c}$, the dynamics of

cell collectives is represented as

$$\boldsymbol{\eta_c \dot{x}_i = k(\varepsilon_j - \varepsilon_{j+1}) + F_{j=i}},$$

$$\boldsymbol{\varepsilon_j = (x_{i+1} - x_i)/L - 1}, \quad\quad \text{for } i = 1,...,N \text{ and } j = 1,...,N \quad\quad \text{Eq. 5}$$

where $\boldsymbol{\varepsilon}$ is a cell strain and L is a typical cell length, i.e., 5 μm. At the front edge of the cells,

i.e., *i=N+1*, a self-propelling force F_{tip} is generated, reflecting an elongation of the duct tip.

Since the cells respond to stretching as activating the ERK, a coupling between the cell

kinematics and the ERK activity is formulated as

$$\eta_E \dot{E}_j = \tanh(\alpha \varepsilon_j) - E_j, \qquad \text{Eq. 6}$$

where α denotes a sensitivity parameter and $\boldsymbol{\eta_E}$ denotes a timescale of the dynamics. Then,

the ERK activity is converted to the self-contractile force represented as dynamics of the

$$\eta_F \dot{F}_j = \lambda E_j - F_j, \qquad \text{Eq. 7}$$

where $\boldsymbol{\lambda}$ denotes a controlling parameter of amplitude, and $\boldsymbol{\eta_F}$ denotes a timescale.

As for the uncoupling regime, ERK activity was given as the following traveling waves

instead of Eq. 6:

$$E(X) = \left(\sin\left(\frac{\pi X}{w} + vt\right) + 1\right) \times 0.5, \qquad \text{Eq. 8}$$

where w is the characteristic length of ERK activation and v is the ERK activation speed.

The parameters w was set as 84 μm, i.e., the wavelength of the ERK activity is 168 μm, from

Figure 4E and v was set as 0.42 μm min^{-1} from Figure 4F.

B-2. Numerical simulation

The ordinary differential equations were numerically solved by the forward Euler method with

time step 0.01 using the MATLAB. The number of cells N was set as 1000 and the one

boundary $i=1$ was fixed and the other $i=N+1$ was the moving boundary condition. Biologically

plausible parameter set was determined as $\boldsymbol{\eta_c}$ = 40 [nN min μm^{-1}], $\boldsymbol{k}$ = 20 [nN], F_{tip} = 6 [nN],

$\boldsymbol{\alpha}$ = 3, $\boldsymbol{\eta_E}$ = 30 [min], $\boldsymbol{\eta_F}$ = 10 [min], and $\boldsymbol{\lambda}$ = 9 [nN] according to the present study and a

previous study (Serra-Picamal et al., 2012).

Supplementary figure legends

Figure S1 Morphological quantification of epithelial layer

Curvature and thickness as a function of the arc length from the tip along the lateral epithelial layer (LEL, left) and the medial epithelial layer (MEL, right) at E13.5 to E15.5. Mean ± s.d. N=3.

Figure S2 Luminal-basal nuclear movement in the MEL

(A) Numerical investigation of the mathematical model. The curvature is plotted over γ. Mean ± s.d. N=10. (B) Treatment of cochlea with 10 µM mitomycin C for 1 day. Maximum intensity projection of immunostained images for the anti-E-cadherin (white) and EdU labelling (magenta). Scale bar, 100 µm.

Figure S3 ERK activity waves and cell flows

(A) Roof plane (orange dotted line) in the cross section view of 3D ERK activity map in the cochlea at E12.5. Scale bar, 50 µm. (B) Time-lapse snapshots of ERK activity map in the roof plane. Time indicates the elapsed time of live imaging. Scale bar, 100 µm. (C) Time-lapse snapshots of tissue flow speed obtained by the PIV in the roof plane. The multicellular flows direct toward the elongation direction around the duct tip despite winding in the region away from the tip in the roof side. Scale bar, 100 µm. (D) Kymograph of ERK activity for the two samples. Horizontal axis indicates the position on the apex-base line and vertical axis indicates the elapsed time of live imaging. Dotted lines represent the oscillatory wave trains from the apex to the base. Scale bar, 100 µm. (E) Time-series data of ERK activity and the tissue flow speed for elongation in the three different ROIs. (F) ERK activity before and after the MEK inhibitor PD0325901 treatment at 1 µM. Scale bar, 20 µm.

Figure S4 Mathematical model for ERK activation waves and cell flows

(A, A') Schematics for two regimes. ERK activity and cell deformation is reciprocally regulated in the mechano-chemical coupling regimes (A), but is unidirectionally regulated from the ERK activity to the cell deformation without closed feedbacks in the uncoupling regime (A'). (B, B') Kymograph of the ERK activity. In the mechano-chemical coupling regime, the ERK activity propagation from the apex to the base can be generated (B), even without given ERK wave as shown in the uncoupling regime (B'). Scale bar, 100 µm. (C, C') Time-series of ERK activity rate and extension-shrinkage rate in the coupling (C) and the uncoupling regime (C'). (D, D')

964   Cross-correlation between the extension-shrinkage rate and the ERK activity rate in the

965   coupling (D) and the uncoupling regime (D'). The lag time in (D) is -28 min, compatible with

966   the experiments, while the lag time in (D') is -2 min, almost no lag time.

Movie legends

Movie S1 Real-time movie showing surgical separation of the cochlear duct at E17.5, related to Figure 1.

After separation, the medial and lateral tissue are placed on the left and the right, respectively. Note that the medial tissue remains curled even after being straightened.

Movie S2 Time-lapse movie of MEL in the cochlear duct at E14.5, related to Figure 2.

Representative nuclei are labelled when those reach at the luminal surface. Green and magenta each denote the nucleus in the flat region and in the curved region. Time interval is 5 min.

Movie S3 Model simulation for cell cycle arrest, related to Figure 2.

Color in the epithelial layer indicates the curvature and that in the circles indicate the cell cycle state. Interkinetic nuclear migration is stopped at a timing of cell cycle arrest shown in the right panel.

Movie S4 3D ERK activity map of the cochlear at E12.5, related to Figure 4.

Color indicates the ERK activity level shown in Figure 4A. Views from roof side at initial, and ones from apex side at last.

Movie S5 Time-lapse movie of the ERK activity and multicellular tissue flow, related to Figure 4.

Color in the left panel indicates the ERK activity level shown in Figure 4B and that in the right panel indicates the PIV speed shown in Figure 4C. Views at the floor plane indicated in Figure 4A'. Time intervals is 12 min.

Movie S6 3D time-lapse imaging of ERK activity in the lateral side of the cochlear duct at E12.5, related to Figure 4.

Color indicate the ERK activity level shown in Figure 4J. Views from the lower left corner of Figure 4B. Green arrow indicates the ERK activation peak.

Figure 1

A Inner ear / Cochlear duct
Base / Apex / Roof / Floor / Lateral / Medial

B E-CAD MYO7A DAPI / Lateral / Medial / 300 μm

C Before / After / 300 μm / 300 μm / Lateral / Medial / Lateral / Medial

D E12.5 / E13.5 / E14.5 / E-CAD / 200 μm

E Lateral / Base / Medial / Apex / Curvature / Thickness / LEL: Lateral epithelial layer / Tip / MEL: Medial epithelial layer

F LEL / MEL / Curvature [μm⁻¹] / Thickness [μm] / Arc length from tip [μm]

G LEL / MEL / Luminal / Basal / 30 μm / Lifeact-EGFP H2B-mCherry

H LEL / MEL / Luminal / Basal / 30 μm / pHH3 EdU DAPI

I MEL / GFP E-CAD DAPI / Apical / Basal / 10 μm

Figure 2

Figure 3

Figure 4