POPULATION GENOMICS REVEALS MOLECULAR DETERMINANTS OF SPECIALIZATION TO TOMATO IN THE

POLYPHAGOUS FUNGAL PATHOGEN *BOTRYTIS CINEREA* IN FRANCE

Alex Mercier[1,2], Adeline Simon[1], Nicolas Lapalu[1], Tatiana Giraud[3], Marc Bardin[4], Anne-Sophie

Walker[1†], Muriel Viaud[1†] and Pierre Gladieux[5*†]

[1] Université Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, 78850 Thiverval-Grignon, France

[2] Université Paris-Saclay, Orsay, France

[3] Ecologie Systématique Evolution, CNRS, Université Paris-Saclay, AgroParisTech, 91400 Orsay, France

[4] UR0407 Pathologie Végétale, INRAE, 84143 Montfavet, France

[5] PHIM Plant Health Institute, Univ Montpellier, INRAE, CIRAD, Institut Agro, IRD, Montpellier, France

[*] Corresponding author: pierre.gladieux@inrae.fr

[†] contributed equally to this study

Abstract

Many fungal plant pathogens encompass multiple populations specialized on different plant species. Understanding the factors underlying pathogen adaptation to their hosts is a major challenge of evolutionary microbiology, and it should help preventing the emergence of new specialized pathogens on novel hosts. Previous studies have shown that French populations of the grey mould pathogen *Botrytis cinerea* parasitizing tomato and grapevine are differentiated from each other, and have higher aggressiveness on their host-of-origin than on other hosts, indicating some degree of host specialization in this polyphagous pathogen. Here, we aimed at identifying the genomic features underlying the specialization of *B. cinerea* populations to tomato and grapevine. Based on whole genome sequences of 32 isolates, we confirmed the subdivision of *B. cinerea* pathogens into two genetic clusters on grapevine and another, single cluster on tomato. Levels of genetic variation in the different clusters were similar, suggesting that the tomato-specific cluster has not recently emerged following a bottleneck. Using genome scans for selective sweeps and divergent selection, tests of positive selection based on polymorphism and divergence at synonymous and non-synonymous sites and analyses of presence/absence variation, we identified several candidate genes that represent possible determinants of host specialization in the tomato-associated population. This work deepens our understanding of the genomic changes underlying the specialization of fungal pathogen populations.

Keywords: Host specialization; grey mould; gene content variation; selective sweeps; positive selection

Introduction

Many fungal plant pathogens encompass multiple lineages, host races or *formae speciales* specialized on different plant species. Understanding the proximate (*i.e.* molecular) and ultimate (*i.e.* eco-evolutionary) factors underlying adaptation to hosts is a major goal for evolutionary microbiology, because emerging diseases are often caused by the appearance and spread of new pathogen populations specialized onto new hosts (Fisher *et al.* 2012; Stukenbrock & McDonald 2008). Evolutionary theory predicts that pathogen specialization should facilitate the emergence of new populations onto novel hosts, because specialization restricts encounters of potential mates within hosts and reduces the survival of offspring due to maladaptation of immigrants and hybrid offspring, thereby reducing gene flow between ancestral and emerging populations (Giraud *et al.* 2010; Nosil *et al.* 2005). The role of specialization as a barrier to gene flow is expected to be strong for pathogens mating within or onto their hosts because, for individuals evolving the ability to infect novel hosts, mating automatically becomes assortative with respect to host use, and reproductive isolation arises as a direct consequence of adaptive divergence (Gladieux *et al.* 2011; Servedio *et al.* 2011). Evolutionary theory also predicts that specialization, and the associated emergence of new populations, could be facilitated by the molecular basis of plant-pathogen interactions, because compatibility is often determined by a limited number of genes in the host and the pathogen, and selection is more efficient when it acts on a smaller number of genes (Giraud *et al.* 2010; Schulze-Lefert & Panstruga 2011). However, despite the apparent ubiquity of specialized fungal pathogens and the negative impact that specialized populations can have on food security and ecosystem health, the genomic features involved in host specialization

remain largely unknown. Acquiring knowledge about the genomic features underlying

pathogen specialization can provide key insights into the mechanisms of specialization.

The ascomycete *Botrytis cinerea* is often presented as a textbook example of a

polyphagous plant pathogen, parasitizing more than 1400 host plant species belonging to

580 genera (Elad *et al.* 2016). Previous population genetic studies reveal that *B. cinerea* is

not strictly speaking a generalist pathogen, and that a more appropriate qualifier is

"polyspecialist", i.e. a set of populations specialized to different host species. Indeed, with

the exception of California (Ma and Michailides, 2005; Caseys et al, 2020; Soltis et al, 2019;

Atwell et al, 2015), studies conducted in multiple regions of the world reveal that

populations were structured (reviewed in Walker 2016), and recognize the host as the factor

with the highest explanatory power for population structure in *B. cinerea*, ahead of

geography. In France, our previous work revealed population subdivision in *B. cinerea,* with

genetic differentiation between field populations infecting tomato (*Solanum lycopersicum*)

and grapevine (*Vitis vinifera*), respectively (Walker *et al.* 2015). This population structure in

B. cinerea was shown to be stable in time and was observed in multiple regions in France.

Furthermore, this structure was later associated with differences in performance on the two

hosts, with pathogens isolated from tomato being more aggressive on tomato than

pathogens isolated from grapevine, and reciprocally (Mercier *et al.* 2019). Altogether, these

data were consistent with a certain degree of specialization of *B. cinerea* populations onto

these two host plants.

Here, we aimed to identify the molecular basis of host specialization in the *B. cinerea*

tomato- and grapevine-associated populations, by addressing the following questions: (1)

Can we confirm the genetic subdivision between *B. cinerea* populations from tomato and

grapevine using genomic data, and what is the degree of divergence between them? (2) Can

we identify genes with footprints of positive selection and/or divergent selection in the

genomes of populations specialized to different hosts, and what are their predicted

functions? (3) Is there variation in gene content between *B. cinerea* populations associated

with tomato and grapevine? To address these questions, we used a set of *B. cinerea* isolates

collected on tomato and grapevine in different regions of France. We Illumina-sequenced

their genomes and identified single nucleotide polymorphisms by mapping sequencing reads

against a high-quality reference genome (van Kan *et al.* 2017). Because some tests of

selection can be biased by population subdivision, while other tests are based on patterns of

population differentiation, we first analyzed the population structure of *B. cinerea* collected

on tomato and grapevine. To detect genes potentially involved in the specialization of *B.*

cinerea to tomato, we searched for signatures of positive selection, by scanning genomes in

the tomato population for selective sweeps, and by estimating the direction and intensity of

selection using McDonald-Kreitman tests on coding sequences. Furthermore, we

investigated signatures of divergent selection using genomic differentiation between

populations, and we characterized variations in the presence/absence of predicted genes

between populations collected on tomato and grapevine using *de novo* genome assemblies,

gene prediction, and orthology analysis.

Materials and methods

Sample collection

Botrytis cinerea samples were selected in a collection of isolates (i.e. single-spored mycelial colonies) originating from three regions of France (Champagne, Occitanie and Provence) and previously characterized using analyses of population structure based on microsatellite markers and pathogenicity tests (Walker *et al.* 2015; Mercier *et al.*, 2019). For each region and each host, we randomly selected three to nine isolates with high membership proportions ($q>0.9$) in the cluster matching their host of origin in a previous analysis of population structure based on microsatellite genotyping (Table 1; Mercier *et al.* 2019). Collection sites were 15 to 133 km apart within regions, and 204 to 722 km apart between regions (Supplementary Figure S1). The set of 32 isolates originated from the following hosts: (i) tomato (*Solanum lycopersicum*; fruits; 13 isolates), (ii) grapevine (*Vitis vinifera*; berries; 16 isolates), (iii) bramble (*Rubus fruticosus*; berries; two isolates) and (iv) hydrangea (*Hydrangea macrophylla*; flower buds; one isolate). Samples from tomato originated from plastic tunnels with sides opened (Occitanie region) or glasshouses (Provence and Champagne regions). Mycelia were cultured on malt-yeast-agar (MYA; 20 g.L^{-1} malt extract, 5 g.L^{-1} yeast extract, 15 g.L^{-1} agar) at 23°C under continuous light until conidiation, and stored as conidial suspensions in glycerol 20% at -80°C until use.

Pathogenicity tests on tomato plants

Isolates of *B. cinerea* collected from tomato or grape were cultivated on MYA medium in a growth chamber (21°C, 14 hours light) for 14 days. Conidia were then washed with sterile distilled water. The conidial suspension was filtered through a 30 µm mesh sterile filter to

remove mycelium fragments. The conidial concentration was determined with a

haemacytometer and adjusted to 10^6 spores/mL. Seeds of tomato var. Clodano (Syngenta)

were sown in compost and transplanted after one week in individual pots. Plants were

grown in a glasshouse for 6 to 8 weeks where they received a standard commercial nutrient

solution once or twice a day, depending on needs. Plants had at least eight fully expanded

leaves when inoculated with a conidial suspension. Each isolate was inoculated on five

plants, from each of which two leaves were removed, leaving 5-10 mm petiole stubs on the

stems. Each pruning wound was inoculated with 10 μl of conidial suspension at 10^6

conidia/mL. Inoculated plants were incubated in a growth chamber in conditions conducive

to disease development (21°C, 16h-photoperiod, 162 $μmol.s^{-1}.m^{-2}$, relative humidity > 80%).

Due to a growth chamber area that did not allow all the isolates to be tested at the same

time, nine series of pathogenicity tests were conducted with 10-12 isolates each, together

with the BC1 reference isolate collected in 1989 in a tomato glasshouse in Brittany

(Decognet *et al.* 2009). For each isolate, two to four independent repetitions of the

pathogenicity test were performed. Lesion sizes (in mm) were assessed daily between the 4[th]

and the 7[th] day post-infection and the Area Under the Disease Progress Curve (AUDPC;

(Simko & Piepho 2012) was computed to take into account the kinetics of disease

development for each isolate. To compare the aggressiveness of isolates, an aggressiveness

index (AI), relative to the reference isolate BC1, was computed as follows: $AI_{isolate} =$

$100 \times (AUDPC_{isolate}/AUDPC_{BC1})$, where $AUDPC_{isolate}$ was the average AUDPC for a given

isolate and $AUDPC_{BC1}$ is the average AUDPC for the reference isolate BC1. Using the AI index

calibrating the AUDPC of a given isolate with that of the reference isolate BC1 in the same

test allows comparing isolates aggressiveness while taking into account the variability

occurring among assays (e.g. plant physiological state; (Leyronas *et al.* 2018). Because of

data non-normality, data were analysed using non-parametric tests. Three statistical tests

were carried out with STATISTICA: (1) we used a non-parametric analysis of variance (Kruskal-

Wallis test) to assess differences among isolates in terms of aggressiveness on grape and

tomato, considering the average values for each of the independent pathogenicity tests as

replications; (2) we used the Mann-Whitney U-test to compare the aggressiveness of isolates

from different hosts of origin (tomato vs. grape), considering the independent repetitions of

the pathogenicity test as blocks and the 13 isolates from tomato and 16 isolates from

grapevine as replicates; (3) we used a Kruskal-Wallis test to compare isolates from three

different clusters (see *Results* section) in terms of aggressiveness on tomato plants.

DNA preparation and sequencing

Isolates were cultivated for 48 h on MYA + cellophane medium at 23 °C in the dark and then

ground using a mortar and pestle in liquid nitrogen. DNA was extracted using a standard

sarkosyl procedure (Dellaporta *et al.* 1983). Paired-end libraries were prepared and

sequenced (2 x 100 nucleotides) on a HiSeq4000 Illumina platform at Integragen (Evry,

France). Sequencing coverage ranged from 58 to 305 X. Genomic data were deposited at SRA

under accession number PRJNA624742. Read quality was checked using FASTQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/).

SNP calling and filtering

SNPs were detected with the same workflow as described in (Zhong *et al.* 2017). Sequencing

reads were preprocessed with TRIMMOMATIC v0.36 (Bolger *et al.* 2014). Preprocessed reads

were mapped onto the *B. cinerea* B05.10 reference genome (van Kan *et al.* 2017) using BWA

v0.7.15 (Li & Durbin 2009). Aligned reads were filtered based on quality using SAMTOOLS v1.3

(Li *et al.* 2009) and PICARD TOOLS (http://broadinstitute.github.io/picard/) to remove

secondary alignments, reads with a mapping quality <30 and paired reads not at the

expected distance. SNP calling was performed with FREEBAYES v1.1 (Garrison & Marth 2012).

Further filtering was carried out using script VCFFiltering.py

(https://urgi.versailles.inra.fr/download/gandalf/VCFtools-1.2.tar.gz), following (Li 2014).

We kept only biallelic SNPs supported by more than 90% of aligned reads, detected outside

low-complexity regions or transposable elements (as identified in the reference isolate

B05.10: https://doi.org/10.15454/TFYH9N; Porquier *et al.* 2016) and with coverage lower

than twice the standard deviation from the mean depth coverage. The VCF file is available

on Zenodo (doi: 10.5281/zenodo.4293375).

Population structure and demographic history

We performed a principal component analysis based on biallelic SNPs using the python

library SCIKIT-ALLEL 1.3.2 (https://github.com/cggh/scikit-allel). We used the sNMF program to

infer individual ancestry coefficients in *K* ancestral populations. This program is optimized for

the analysis of large datasets and it estimates individual admixture coefficients based on

sparse non-negative matrix factorization, without assuming Hardy-Weinberg equilibrium

(Frichot et al., 2014). We used SPLITSTREE 4 (Huson & Bryant 2005) to visualize relationships

between genotypes in a phylogenetic network, with reticulations representing the

conflicting phylogenetic signals caused by recombination or incomplete lineage sorting. The

position of the root was determined using a *B. fabae* isolate as the outgroup. *Botrytis fabae*

is one of the closest known relatives of *B. cinerea* (Amselem *et al.* 2011; Walker 2016).

Summary statistics of genomic variation (segregating sites S, nucleotide diversity π,

Watterson's θ, Tajima's D) were estimated using EGGLIB 3.0 (https://egglib.org/) on 10kb

windows, excluding sites with more than 50% missing data and removing windows with

lseff<1000 (lseff is the number of analysed sites) or nseff<3 (nseff is the average number of

exploitable samples). Site frequency spectra were estimated using DADI 1.7.0 (GUTENKUNST *ET*

AL. 2009). Allelic richness and private allele richness were estimated with ADZE 1.0 (Szpiech

et al. 2008), using a generalized rarefaction approach to account for differences in sample

size among populations.

We employed the f3 statistic (Reich *et al.* 2009) to test for admixture based on shared

genetic drift, as implemented in the POPSTATS python script (Skoglund *et al.* 2015;

https://github.com/pontussk/popstats/). The *f3* statistic is used to test for admixture among

three populations Px, P1, P2. In the no-admixture case, the *f3* statistic measures the branch

length between Px and the internal node of the unrooted population tree (Px;P1,P2), and it

is therefore expected to be greater than zero. In the case Px has a mixed ancestry from P1

and P2, or populations closely related to them, the *f3* statistic is expected to be negative.

Significance was assessed by block jackknife by treating each chromosome as a block and

weighting each block by the number of SNPs. The standard error of the test statistic was

used to define a Z-score.

Genome assembly and gene prediction

Gene content was determined using two independent approaches. In the first approach, we used the read mapping coverage of the genes previously predicted in the reference genome B05.10 (van Kan *et al.* 2017). Read count per site for each gene and for each isolate was computed using SAMTOOLS BEDCOV, and normalized using EDGER (McCarthy *et al.* 2012; Robinson *et al.* 2010). Genes showing significant differences in mapping coverage across populations were identified with EDGER (adjusted p-value <0.05 and more than two-fold change). Putative gene duplication events (higher mapping coverage) and missing B05.10 genes (lack of mapping reads) were visually inspected in their genomic context using a genome browser for validation.

The second approach was based on genome assembly and *de novo* gene prediction. Illumina paired-reads were assembled using a combination of VELVET (Zerbino & Birney 2008), SOAPDENOVO and SOAPGAPCLOSER (Luo *et al.* 2012), as follows: (1) reads were trimmed at the first N, (2) contigs were generated with several k-mer values using SOAPDENOVO, (3) several VELVET assemblies were built using several k-mer values and as the input the trimmed reads and all SOAPDENOVO contigs considered as "long reads ", (4) the assembly that maximizes the criterion (N50*size of the assembly) was selected, (5) SOAPGAPCLOSER was run on the selected assembly, and (6) Contigs completely included in other longer contigs were deleted. Genomic regions mapping to transposable elements previously identified in *B. cinerea* (https://doi.org/10.15454/TFYH9N; Porquier *et al.* 2016) were masked with REPEATMASKER (http://www.repeatmasker.org) prior to gene prediction. Genes were predicted using the FGENESH *ab initio* gene-finder (Solovyev *et al.* 2006;

http://www.softberry.com/berry.phtml), the program previously used to annotate the

reference genome (Amselem *et al.* 2011), and for which *Botrytis*-specific gene-finding

parameters were thus available. Completeness of the assembly and gene prediction were

evaluated with BUSCO using the Ascomycota gene set (Seppey *et al.* 2019). We then used

ORTHOFINDER (Emms & Kelly 2015) in order to define groups of orthologous sequences

(hereafter "groups of orthologs") based on sequences of predicted genes translated into

protein sequences. To reduce the impact of incomplete gene prediction (*e.g.* truncated

genes in small contigs), groups of orthologs were then manually checked for the

presence/absence of the protein-encoding genes in the genomes using TBLASTN (Johnson *et*

al. 2008). Proteins were functionally annotated using INTERPROSCAN (Jones *et al.* 2014), and

SIGNALP (Almagro Armenteros *et al.* 2019) was used to predict secretion signal peptides.

Prediction of transmembrane helices in proteins was performed using TMHMM 2.0 (Krogh *et*

al. 2001; Sonnhammer *et al.*). As we detected traces of bacterial contamination in the

genome of the SI13 isolate, this genome was not included in orthology analysis. However,

we have kept this genome for analyzes based on polymorphism, because bacterial reads

cannot map to the reference genome.

Tests of positive selection based on polymorphism and divergence at synonymous and non-

synonymous sites

We estimated the intensity and direction of selective pressures exerted on genes in

populations using the McDonald-Kreitman test based on polymorphism and divergence at

synonymous and non-synonymous sites (McDonald & Kreitman 1991). This test is based on

the number of nucleotide polymorphisms and substitutions in gene sequences, and assumes

that synonymous mutations are neutral. Pseudo-sequences for coding sequences of all

genes in the reference genome were generated using the VCF file and the reference

sequence. Synonymous and non-synonymous divergence was computed with CODEML (model

0; Yang 1997, 2007), using *B. fabae* as the outgroup. Synonymous and non-synonymous

polymorphism was computed using EGGLIB 3 (https://egglib.org/), filtering out sites with

more than 80% missing data and excluding filtered alignments with less than 10 codons or

four sequences. Samples Sl1, Sl2, Sl3, Sl13, Vv3, Vv5, Vv2, Vv4 and Vv6 were excluded

because they introduced missing data that prevented calculations of synonymous and non-

synonymous divergence. The neutrality index (NI), defined as $(P_N/D_N)/(P_S/D_S)$ (Stoletzki &

Eyre-Walker 2011), was computed for every gene, with P_N and D_N the numbers of non-

synonymous polymorphisms and substitutions, respectively, and P_S and D_S the numbers of

synonymous polymorphisms and substitutions, respectively. Pseudocounts of one were

added to each cell of the McDonald-Kreitman tables to ensure the NI was always defined

(*i.e.* no division by zero).

Linkage disequilibrium and recombination

We used POPLDDECAY (Zhang *et al.* 2019a) to measure linkage disequilibrium (r^2) as a function

of the distance between pairs of SNPs. POPLDDECAY was configured with a maximum distance

between SNPs of 300 kbp, a minor allele frequency of 0.005 and a maximum ratio of

heterozygous allele of 0.88. Recombination rates were estimated for each chromosome with

PAIRWISE in LDHAT version 2.2 (Auton & McVean 2007). Sites with missing data were excluded.

279
Tests of positive selection based on linkage disequilibrium and the site frequency spectrum

We searched for signatures of selective sweeps along genomes using three different

softwares, each implementing a different approach. The SWEED 3.0 software (Pavlidis *et al.*

2013) implements a composite likelihood ratio (*CLR*) test based on the SWEEPFINDER algorithm

(Nielsen *et al.* 2005), which uses the site frequency spectrum (SFS) of a locus to compute the

ratio of the likelihood of a hard selective sweep at a given position to the likelihood of a null

model hypothesis without selection. The CLR statistic was computed for each chromosomes

of each population using a grid size of 50 or 200 (grid size is the number of positions where

the likelihood is calculated), but only results for a grid size of 200 are presented because the

CLR profiles were highly similar between the two settings. Input files included both

"unfolded" SNPs (i.e. SNPs for which ancestral and derived states can be determined using

the allelic state of the outgroup) and "folded" SNPs (i.e. SNPs for which the outgroup had

missing data). Only biallelic sites with sample size greater than or equal to five were

included. The nS_L method implemented in the NSL software (Ferrer-Admetlla *et al.* 2014)

detects hard and soft selective sweeps based on haplotype homozygosity. The nSL statistic

was computed for each chromosome of each population, including only biallelic sites with

sample size greater than or equal to five. Nine isolates (Sl1, Sl2, Sl3, Sl13, Vv3, Vv5, Vv2, Vv4,

Vv6) were excluded to reduce the proportion of missing data. The *hapFLK* method (Fariello

et al. 2013) implemented in the HAPFLK software is based on the original *FLK* method by

(Bonhomme *et al.* 2010), which detects signatures of selection from differentiation between

populations. This metric was used to test the null hypothesis of neutrality by contrasting

allele frequencies at a given locus in different populations. *hapFLK* extends the FLK method

to account for the haplotype structure in the sample, and the method is robust to the effects

of bottlenecks and migration. HAPFLK takes the number of cluster of haplotypes as a

parameter (K). To determine the number of clusters of haplotypes K, we ran FASTPHASE v1.4

(Scheet & Stephens 2006) and R package IMPUTEQ (Khvorykh & Khrunin 2020) on

polymorphism data for the largest chromosome BCIN01. For each population, we used

IMPUTEQ to generate five datasets with 10% of polymorphic positions masked, and for each

masked dataset we used FASTPHASE for imputing masked positions assuming clusters of

K=2,3...10 haplotypes and the following parameters: -T10 -C25 -H-1 -n -Z. We estimated

error using Estimate-Errors function in IMPUTEQ and the number of clusters that minimizes

the error was selected as the optimum. The HapFLK metric was computed individually on

each chromosome using as number of clusters K = 5, and nfit = 2. Accessory chromosomes

were not included in these analyses as they showed presence/absence polymorphism (see

Results section).

Results

Whole-genome sequencing, population structure and demographic history

Previous work using microsatellite data revealed differentiation between populations of *B.*

cinerea collected on tomato and grapevine (Mercier *et al.* 2019; Walker *et al.* 2015). To

investigate the genetic basis of specialization of the tomato- and grapevine-infecting

populations, we randomly selected for genome sequencing 32 isolates collected on tomato

(13 isolates), grapevine (16 isolates), *Rubus* (two isolates) and *Hydrangea* (one isolate) (Table

1). Isolates from *Rubus* and *Hydrangea* were previously shown to belong to generalist

populations (*i.e.* assigned to clusters found on all sampled hosts; Mercier *et al.* 2019). One

isolate of the sister species *B. fabae* was also sequenced and used as the outgroup.

Information about the sequenced isolates is summarized in Table 1. Alignment of sequencing

reads to the B05.10 reference genome (van Kan *et al.* 2017) followed by SNP calling

identified 249,084 high-quality SNPs.

In clustering analyses based on sparse nonnegative matrix factorization algorithms, as

implemented in the sNMF program (Frichot et al., 2014), the model with *K*=4 clusters was

identified as the best model based on cross-entropy (Supplementary Figure S2) and models

with K>4 did not identify well-delimited and biologically relevant clusters (Fig. 1A). At K=4,

one cluster was associated with tomato (hereafter referred to as "T" cluster), two clusters

were associated with grapevine (hereafter referred to as "G1" for the largest, and "G2" for

the smallest), and one cluster was formed by the isolates Rf1 and Hm1 from bramble (*R.*

fruticosus) and hydrangea (*H. macrophylla*) (Figure 1A). The isolate Rf2 collected on wild

blackberry displayed ancestry in multiple clusters, and the reference isolate B05.10 had

ancestry in the G2 and T clusters. No pattern of geographical subdivision was observed,

consistent with previous findings (Walker *et al.* 2015; Mercier *et al.,* 2019). The neighbor-net

network inferred with SPLITSTREE revealed three main groups, corresponding to the three

main clusters identified with sNMF, with cluster G2 (isolates Vv8, Vv9, Vv11 and Vv15) more

closely related to T than to G1 (Figure 1B). Reticulations in the neighbor-net network and

patterns of membership at K=2 and K=3 in the analysis with sNMF indicated that cluster G2

shared recent ancestry with clusters G1 and T. However, tests for admixture using the f3

statistic (Reich *et al.* 2009; Skoglund *et al.* 2015) did not support a scenario in which G2

derived from admixture between the two other clusters (Supplementary Table S1). The

principal component analysis corroborated the results of clustering analyses and the

neighbor-net network (Figure 1C). The first principal component separated isolates Hm1 and

Rf1 from the rest of the dataset. The second principal component individualized isolate Rf2,

as well as the three clusters T, G1 and G2. The third and fourth principal components

individualized cluster G2 and isolate Sl11, respectively. Together, analyses of population

subdivision revealed three clearly defined populations (two on grapevine and one on

tomato) and we therefore focused on these populations to identify the genes underlying

differences in host specialization.

On average across core chromosomes BCIN01 to BCIN16, nucleotide diversity π and

Watterson's θ were comparable in the three populations (from $\pi=0.0018$/bp in G1 to π

$=0.0030$/bp in G2; from $\theta=0.0017$/bp in G1 to $\theta=0.0027$/bp in G2; Table 2; Supplementary

Table S2), although all comparisons were statistically significant except between populations

T and G2 for θ (two-tailed Wilcoxon signed-rank test, P-value>0.05). Allelic richness was

slightly higher in G2 than in T (AR=1.104 vs AR=1.094), and lower in G1 (AR=1.078). Private

allele richness was higher in G1 (PAR=0.085), than in T (PAR=0.073) and G2 (PAR=0.071)

(Table 2).

The site frequency spectra estimated in populations G1 and T were U-shaped, indicating

an excess of high-frequency derived alleles (Figure 2; population G2 was excluded because of

too small a sample size), consistent with ongoing episodes of positive selection, mis-

assignment of ancestral alleles or gene flow (Marchi & Excoffier 2020). Estimates of Tajima's

D were positive but close to zero in clusters T and G1 (T: D=0.309; G1: D=0.167), indicating a

slight deficit of low frequency variants (Table 2; Supplementary Table 2), consistent with

balancing selection or population contraction. In cluster G2, the estimated Tajima's D was

D=1.470 but the estimate was likely upwardly biased by the small sample size, because

Watterson's θ is underestimated when sample size is small. The distance at which linkage

disequilibrium (LD) decayed to 50% of its maximum was an order of magnitude longer in G2

(LDdecay50: 16,800bp) than in T (LDdecay50: 8600bp) and G1 (LDdecay50: 3100bp)

(Supplementary Figure S3). The recombination rate was higher in G2 (ρ=0.2047/bp) than in

G1 and T (G1: ρ=0.0178/bp; T: ρ=0.0108/bp).

The accessory chromosome BCIN18 showed no polymorphism in all three population and

the accessory chromosome BCIN17 showed no polymorphism in population G2

(Supplementary Table S2). However, it should be noted that coverage analysis also revealed

that these accessory chromosomes, which contain a reduced number of genes (23 and 19

respectively, van Kan *et al.* 2017), were distributed irrespectively of the host of origin:

BCIN17 was found present in all but three isolates (Sl9, Sl10 and Vv15), while BCIN18 was

only present in five isolates (Sl3, Sl5, Sl9, Vv9 and Vv11; Supplementary Table S3). In

population T, the accessory chromosome BCIN17 displayed a relatively high and positive

value of Tajima's D (D=1.37; Supplementary Table S2) and approximately twice as much

nucleotide diversity as in core chromosomes (π=0.0058/bp; Supplementary Table S2). In

population G1, accessory chromosome BCIN17 displayed a negative value of Tajima's D (D=-

0.845) and two orders of magnitude less nucleotide diversity than core chromosomes

(π=5.7e-5/bp; Supplementary Table S2). The differences in Tajima's D estimates for BCIN17

reflect the existence of two divergent haplotypes in T, but not in G1 (not shown).

Isolates from the T population are more aggressive on tomato plants.

To test whether isolates collected on tomato are more aggressive on their host of origin,

compared to isolates collected on grapevine, pathogenicity assays were performed on whole

tomato plants in controlled conditions. To assess differences among isolates in terms of

aggressiveness on grape and tomato, we used a non-parametric analysis of variance,

considering the average values for each of the independent pathogenicity tests as

replications. We found a significant isolate effect (Kruskal Wallis test, H (28, N= 87)

=69.89221, p < 0.0001), consistent with the wide range of aggressiveness levels observed for

the 29 isolates.

To compare the aggressiveness on tomato of isolates from different hosts of origin

(tomato vs. grape), we tested for differences in the distribution of the aggressiveness index

between isolates originating from the two types of hosts, considering the independent

repetitions of the pathogenicity test as blocks and the 13 isolates from tomato and 16

isolates from grapevine as replicates. We observed a significant effect of the host of origin

(Figure 3, Mann-Whitney U test, p < 0.0001) and T isolates collected on tomato were on

average 2.7 times more aggressive on tomato plants than isolates collected on grapevine.

To compare the aggressiveness on tomato of isolates from three different clusters (T, 13

isolates; G1, 12 isolates; G2, 4 isolates), we tested whether the aggressiveness index of

isolates from different clusters originate from the same distribution. A Kruskall-Wallis test

rejected the null hypothesis that all clusters display the same median of the aggressiveness

index (H (2, N= 87) =19.80083, p < 0.0001). The T population was significantly different from

the G1 population but not from G2 (Kruskall-Wallis test, p< 0.0001 and p = 0.37,

respectively). G2 and G1 were not significantly different (Kruskall-Wallis test, p = 0.57).

Gene content slightly differs between tomato- and grapevine-associated populations

As variation in gene content can be involved in adaptation to novel hosts (Cummings *et al.*

2004; Inoue *et al.* 2017; Langridge *et al.* 2015), we sought to identify genes specific to the

tomato (T) and grapevine (G1 and G2) populations. We first explored the mapping coverage

418 of genes previously identified in the B05.10 reference to identify sets of genes that were

419 missing or showing duplication events (Supplementary Table S4). Five B05.10 genes were

420 identified as missing in the T population, including four consecutive genes in the

421 subtelomeric region of chromosome BCIN02 that could correspond to a secondary

422 metabolism gene cluster. Among these four genes, one is coding for a NRPS-like enzyme

423 similar to the protein MelA of *Aspergillus terreus* involved in the biosynthesis of an α-keto

424 acid dimer (Geib et al. 2016), two other genes encode putative biosynthetic enzymes (FAD-

425 binding and enoyl reductase domains). Mapping coverage of B05.10 genes also suggested

426 some possible duplication events in a subtelomeric region of chromosome BCIN08, with T

427 isolates showing approximately three times as many reads as the G1 and G2 isolates for the

428 four consecutive genes Bcin08g00060 to Bcin08g00090. This suggested that the

429 corresponding region of at least 25 kb would be in three copies in the genomes of the T

430 isolates. Among the four duplicated genes, two encode carbohydrate-active enzymes

431 (CAZymes) known as plant cell wall degrading enzymes (PCWDEs) as they act on pectin

432 (glycoside hydrolase GH28) and hemicellulose or pectin side chains (GH43).

433  We also analysed the variation in gene content using a different approach that makes no

434 use of reference genome B05.10. We built *de novo* assemblies of the genomes of T, G1 and

435 G2 isolates. The genome assembly size of the 28 isolates ranged from 41 Mb to 42.5 Mb

436 (Supplementary Table S5), which was slightly smaller than the genome assembly size of the

437 B05.10 reference isolate (42.6 Mb). We then predicted genes *ab initio* using FGENESH. The

438 number of predicted genes ranged from 11,109 to 11,311 among genomes. To compare

439 gene content in the T, G1 and G2 populations, we used ORTHOFINDER to identify 12,319

440 groups of orthologous sequences (i.e., orthogroups). The number of groups of orthogroups

shared by pairs of isolates within populations was higher that between populations

(Supplementary Table S6). By looking for orthologous groups that were present in at least

75% of the genomes of a focal population and missing in other populations, we identified 21

G1-specific genes, a single G2-specific gene, five genes specific to the G1 and G2 populations

(those already detected with the first approach described above), two genes missing

specifically in the G1 population, and a single gene specific to the T population

(Supplementary Table S7). This latter gene was a GH71 glycoside hydrolase (OG0011490, an

α-1,3-glucanase; (Lombard *et al.* 2014) acting on fungal cell wall. Among the genes specific

to G1, we found a GH10 glycoside hydrolase (in OG0011469; (Lombard *et al.* 2014) acting on

plant cell wall (*i.e.* hemicellulose). The proteins encoded by the other G1-specific genes had

no functional prediction though four of them shared a domain typical of metalloenzymes

(IPR11249) with putative peptidase activities and three other ones showed a versatile

protein-protein interaction motif involved in many functions (IPR011333). Two proteins with

secretion signal peptides were also found specific to G1 (OG0011305 and OG0011366), with

OG0011366 having a predicted function of interferon alpha-inducible protein-like (Rosebeck

& Leaman 2008) and a predicted transmembrane helix.

Together, these analyses revealed that the magnitude of gene content variation is limited

between *B. cinerea* populations, which emphasizes the need to investigate differences in

allelic content at shared genes for elucidating the genomic basis of host specialization.

McDonald-Kreitman tests of positive selection identify genes related to virulence

We investigated differences in the direction and intensity of natural selection driving the

evolution of gene sequences in the two populations with the greatest difference in terms of

quantitative pathogenicity on grape and tomato, which are also the two populations with

the largest sample size (G1 and T). More specifically, we searched for genes with signatures

of positive selection in both populations that also show high sequence divergence between

populations, or genes with signatures of positive selection in one population, but not in the

other population. The direction and intensity of selection was estimated using neutrality

indexes computed for each individual gene in each population based on McDonald-Kreitman

tables of polymorphisms and substitutions at synonymous and non-synonymous sites, using

B. fabae as the outgroup. The neutrality index is expected to be below one for genes under

positive selection (due to an excess of non-synonymous substitutions) and above one for

genes under negative selection (due to a deficit of non-synonymous polymorphisms). To

identify genes potentially involved in host specialization, we first selected genes showing low

values of the neutrality index in both populations (log [neutrality index] ≤-0.5), and high

values of the inter-population dN/dS ratio (dN/dS in the top 5% percentile; Supplementary

Figure S4). This analysis identified five genes: Bcin02p04900, Bcin07p02650, Bcin09p06530,

Bcin14p01690, Bcin09p02190 (Supplementary Table S8). Two genes (Bcin07p02650,

Bcin14p01690) code for glycosyl hydrolases of the GH5 family, a family that includes

enzymes acting on plant cell walls and enzymes act on fungal cell walls. One of the two

genes (Bcin14p01690), has a cellulose binding domain which strongly suggests a role as

PCWDE. Two genes (Bcin02p04900, Bcin09p02190) are involved in basic cell functions (a

DNA nuclease and a protein involved in ribosome biogenesis). The last gene (Bcin09p06530)

has no known domain.

We also selected genes showing low values of the neutrality index in one population (log

[neutrality index] ≤-0.5), and high values in the other (log [neutrality index]>=0)

(Supplementary Figure S4). This analysis identified 392 genes in G1 and 428 genes in T

(Supplementary Table S8). Functional enrichment analyses revealed contrasting results in

the G1 and T populations (Supplementary Table S8). In the G1 population, we identified a

significant two-fold enrichment in transporter-encoding genes among the 392 genes with

signatures of positive selection. These 28 transporters included many candidates with

putative roles in nutrition such as the transport of sugars and amino-acids (five genes of

each). Fifteen of them were proteins of the major facilitator superfamily (MFS). The MFS

transporter-encoding genes included five putative sugar transporters and ten unknown

transporters that could have roles in various processes, including obtaining nutrients from

the host, efflux of fungi-toxic compounds or the export of fungal phytotoxins (Hartmann *et*

al. 2018; Maruthachalam *et al.* 2011; Perlin *et al.* 2014). Finally, five of the 392 genes

encoded ATPase transporters including the BcPrm1 P-type Ca2+/Mn2+-ATPase that

mediates cell-wall integrity and virulence in *B. cinerea* (Plaza *et al.* 2015).

In the T population, the list of 428 genes with signatures of positive selection showed a

significant 2.5-fold enrichment in genes encoding for proteins involved in oxidative stress

response (eight genes). These genes encode enzymes that are able to detoxify reactive

oxygen species *i.e.* glutathione-S-transferases (BcGST1, 9 and 24), the superoxide dismutase

BcSOD1, and two peroxidases (BcPRX8 and BcCCP1). In addition, the list of 428 genes also

included *BcatrO*, which encodes the transporter BcAtrO involved in the resistance to H_2O_2

(Pane *et al.* 2008).

A 2.5-fold enrichment was also observed for the genes coding for CAZYmes acting as

PCWDEs (ten genes) especially for those involved in the modification of hemicellulose (five

genes; Espino *et al.* 2010 ; Supplementary Table S8) such as the xylanase BcXyn11A that is

required for full virulence on tomato (Brito *et al.* 2006).

Selective sweeps in regions encompassing genes encoding enzymes involved in carbohydrates

metabolism

To identify genomic regions with signatures of selective sweeps, we conducted three

different genome scans, using different features of the data: i) hapFLK, which detects hard

and soft sweeps based on patterns of differentiation between clusters of haplotypes

between populations; ii) nS_L, which detects hard and soft sweeps based on the distribution

of fragment length between mutations and the distribution of the number of segregating

sites between pairs of chromosomes; iii) and SWEED's CLR, which detects hard sweeps based

on the site frequency spectrum. The *CLR* and nS_L metrics are population-specific and were

computed for each population independently, while the *hapFLK* metric is F_{ST}-based and was

thus computed for populations G1 and T (Supplementary figure S5). We identified candidate

SNPs located in (hard or soft) selective sweeps as the SNPs that were in the top 5% of the

hapFLK statistic, but also in the top 5% of either the nSL or CLR statistic in a least one

population (Figure 4). In total, this approach identified 4,667 SNPs of which 1,300 were

localized in coding sequences, 256 in introns, 830 in untranslated transcribed regions, and

465 less than 1500bp upstream of coding sequences. These 2,851 SNPs corresponded to 351

genes, of which 15 were identified by SNPs in the top 5% of selective sweep metrics in both

populations, 200 by SNPs in the top 5% of selective sweep metrics in population T, and 175

by SNPs in the top 5% of selective sweep metrics in population G1 (Supplementary table S9).

Candidate genes in the selective sweep regions of each population included genes coding

for proteins with functions consistent with a role in infection, such as transporters, CAZymes,

putative effectors and two genes that are confirmed virulence factors. Indeed, one region

identified in the G1 population contains the gene encoding the Pectin Methyl Esterase

BcPme1 required for full virulence of *B. cinerea* on several host including grapevine (Valette-

Collet *et al.* 2003), and another region contains the gene encoding BcCgf1, a small secreted

protein that is essential for infection structure development (Zhang *et al.* 2020).

Nevertheless, it is unlikely that all genes in selective sweep regions have been direct targets

of positive selection, most of them being possible hitch-hikers. This could be the reason why

no significant functional enrichment was detected among these genes.

Finally, comparison of the lists of genes identified in the selective sweep regions and

those with signatures of positive selection according to McDonald-Kreitman tests identified

nine genes in common for the T population and four genes in common for the G1 population

(Supplementary table S9). This suggests that these genes were subjected to both recurrent

positive selection for amino-acid changes and to recent positive selection in populations of

B. cinerea. In addition, we can hypothesize that these genes may be the actual targets of

positive selection, and that surrounding candidate genes could be only hitch-hikers.

Functional annotations of the genes that show both recurrent positive selection for amino-

acid changes and recent positive selection signals further indicated a cutinase-encoding gene

(Bcin01g09430) in the T population and a sugar transporter encoding gene (Bcin16g00530) in

the G1 population, and also pointed out various other functions.

Discussion

Genetic differentiation in B. cinerea *between populations associated with grapevine and*

tomato

Genetic structure associated with the host of origin in *B. cinerea* has been extensively

investigated (reviewed in Walker 2016). Most of the studies based on sufficient sampling

size (n>100) found significant population differentiation between populations of *B. cinerea*

from different hosts (e.g. in Chile, Tunisia, Hungary, United Kingdom; Walker 2016). One

noticeable case is the population of *B. cinerea* collected from various hosts in California (Ma

& Michailides 2005), for which whole-genome sequencing data did not detect any host-

associated population structure despite differences in pathogenicity against different hosts

(including tomato) in cross-infectivity assays (Atwell *et al.* 2015; Caseys *et al.* 2020; Soltis *et*

al. 2019). In France, previous research concluded that *B. cinerea* populations were

differentiated according to some of their hosts, including tomato, grapevine and, to a lesser

extent, bramble (Fournier & Giraud 2008; Walker *et al.* 2015). In two recent studies

comparing the aggressiveness of isolates coming from diverse hosts, the disease severity

caused by isolates from tomato was significantly greater than the severity caused by isolates

from grape or other crops, thus indicating that *B. cinerea* populations parasitizing tomato

were specialized to this host (Bardin *et al.* 2018; Mercier *et al.* 2019). Here we show that

populations parasitizing tomato and grapevine are subdivided into three populations, two

being associated with grapevine (G1 and G2) and one with tomato (T). This pattern of

population genetic structure differs from previous findings (Walker et al., 2015), as three

populations parasitizing grapevine had previously been detected, but studies differ in terms

of sampling and genotyping schemes (thousands of SNPs vs 8 SSR markers, and an order of

magnitude of difference in the size of sample sets). The clear pattern of population

subdivision found in our study also stands in sharp contrast with the lack of host- or

geography-associated population subdivision across various hosts, including tomato and

grape, in California (Atwell *et al.* 2015; Atwell *et al.* 2018). This difference in population

structure between France and California indicates that the factors leading to host-specific

differentiation between *B. cinerea* pathogens from grape and tomato do not operate

everywhere. The differences in LD decay (up to an order of magnitude longer in our study

compared to Californian *B. cinerea*) and nucleotide diversity (half as much in our study

compared to Californian *B. cinerea*) also suggest that the demographic history and

population biology of the pathogen is contrasted between the two regions.

Multiple factors can contribute to reduce gene flow between populations parasitizing

grapevine and tomato. A first possible factor limiting gene flow is adaptation to host. Mating

in *B. cinerea* occurs on the host after infection, between individuals that were thus

sufficiently adapted to infect the same host, which induces assortative mating with respect

to host use and reduce opportunities for inter-population crosses (Giraud 2006; Giraud *et al.*

2010; Giraud *et al.* 2008). Another factor possibly limiting gene flow between populations

infecting tomato and grape is habitat isolation (*i.e.* reduced encounters caused by mating in

different habitats). Tomatoes are grown in nurseries before being dispatched to the fields,

tunnels or greenhouses, and this may generate habitat isolation if sexual reproduction in the

pathogen occurs in nurseries for the tomato-infecting population of *B. cinerea*. Such habitat

isolation may contribute to promote adaptation to tomato, by preventing the immigration of

alleles which are favorable for infection of non-tomato hosts but not favorable for infection

of tomato. Differences in the timing of epidemics are unlikely to contribute to this habitat

599 isolation, as the period of infection of greenhouse tomatoes runs from late to early winter,

600 which includes the period of infection of grape. The same goes for the location of epidemics,

601 since the sites studied were chosen because the two types of crops are grown nearby. A final

602 possibility to explain the lack of gene flow between *B. cinerea* populations from grape and

603 tomato is that the frequency of sexual reproduction might be lower in populations infecting

604 greenhouse tomatoes. Higher winter temperatures and the removal of plant residues in the

605 greenhouse represent conditions that are less conducive to sexual reproduction. However,

606 our estimates of LD decay and recombination rates are not consistent with a substantially

607 lower frequency of sexual reproduction in the population associated with tomato, compared

608 to the population associated with grape.

609  The differences in population structure observed between *B. cinerea* populations from

610 France and California could be due to different cultivation practices. In California, differences

611 in pathogenicity between tomato and other hosts did not lead to genome-wide

612 differentiation, indicating that gene flow occurs between hosts. In France, on the contrary,

613 differences in virulence are associated to genome-wide differentiation, indicating restriction

614 of gene flow. These differences in structure may be explained by the cultivation of tomatoes

615 in open fields in California, which favors dispersal to other crops, while French tomatoes are

616 generally grown in plastic tunnels or greenhouses.

618 *Widespread signatures of selection along genomes*

619 We identified little variation in the gene content among T, G1 and G2 populations, with one

620 gene specific to T, 22 genes specific to G1 and five genes shared between G1 and G2 but not

621 T, suggesting that gene gain or loss is not the main process of adaptation to tomato. In

parallel to our analysis of presence/absence variation, our genome scans for positive

selection pinpointed several genomic regions which may harbour determinants of ecological

differentiation between the population specialized to tomato and the population parasitizing

grapevine. In order to cover multiple time scales and different signatures of positive

selection, we used a variety of analytical approaches. The McDonald-Kreitman test focuses

on genes and detects repeated episodes of selective sweeps fixing non-synonymous

substitutions, thus generating a higher ratio of amino acid divergence to polymorphism (D_n /

P_n), relative to the ratio of silent divergence to polymorphism (D_s / P_s), than expected under

neutrality. The values of nucleotide diversity and Tajima's D measured in the T population

specialized in tomatoes were very close to the values measured for the two other

populations, which is not consistent with a very recent origin of this population and justifies

the use of the McDonald-Kreitman test. Genome scans for selective sweeps detect more

recent events, and by nature these methods can also detect genes that are not directly the

target of selection, but may have hitch-hiked due to physical linkage with sites under

positive selection. However, the LD decay values measured for the T population remain

moderate (9kb), and we used a combination of different selective sweep metrics to

substantially shorten the list of candidate genes, which should reduce the impact of genetic

hitch-hiking on our list of genes under recent positive selection. Despite using a more

conservative approach, we identified more selective sweeps than in the generalist

Sclerotinia sclerotiorum fungus (Derbyshire *et al.* 2019), or in the Californian generalist

population of *B. cinerea* (Soltis *et al.* 2019).

Genes under positive selection

We identified a number of genes showing signatures of positive selection using the

approaches discussed above and highlighting potential candidates for their role in host

specialization. Functional annotation of the *B. cinerea* genome and previous experimental

studies provided lists of genes involved in host-pathogen interaction and in other

developmental processes (Amselem *et al.* 2011; Nakajima & Akutsu 2014; Rodriguez-

Moreno *et al.* 2018). We used these published lists of genes to investigate whether some

specific biological processes were subjected to positive selection in the different

populations. Our data revealed that the five genes showing the strongest signatures of

selection in both the T and G1 populations, but also showing high sequence divergence

between the two populations, included two genes encoding for CAZymes with glycoside

hydrolase activity, which are potential PCWDEs. We also found that the 428 genes showing

the strongest signatures of selection in the T population with McDonald-Kreitman tests were

enriched both in genes coding for PCWDEs and in genes coding for enzymes involved in the

oxidative stress response. Notably, ten genes encoding secreted CAZymes targeting

compounds of the plant cell wall, *i.e.* cellulose and pectin (Amselem et al., 2011; Lombard *et*

al. 2014), were found under positive selection. One of these ten genes encodes the xylanase

BcXyn11A that has previously been shown to be a virulence factor on tomato (Brito *et al.*

2006). Another one encodes a cutinase (Bcin01g09430) that was further detected in a

selective sweep region suggesting, recurrent and recent positive selection events. Additional

PCWDEs were found in other genomic regions identified as selective sweeps. Finally, our

comparative genomic analysis suggested that a subtelomeric region that contains two

PCWDEs acting on pectin and/or hemicellulose is duplicated in the T population.

Necrotrophic species have important repertoires of CAZymes especially those corresponding

to PCWDEs which are known to act as major virulence factors in fungi (Zhao *et al.* 2013;

Rodriguez-Moreno *et al.* 2018). The genome of the reference isolate of *B. cinerea* (B05.10)

revealed 118 PCWDEs (Amselem et al., 2011) and our data suggest that, within this

repertoire, some cellulases and pectinases may be of particular importance for the

degradation of tomato cell wall. SNPs within a pectinesterase gene were also associated with

virulence on tomato in a previous genome-wide association study (Soltis *et al.* 2019).

In addition to PCWDEs, the single gene that was present in the T population but missing in

the G1 and G2 populations encoded a CAZyme acting on the fungal cell wall, an α-1,3-

glucanase classified as a member of the GH71 family. In the fungal cell wall, α-1,3-glucan is a

major component that encloses the α-(1,3)-glucan-chitin fibrillar core. Because of its

external localization and specific composition, α-1,3-glucan of pathogenic fungi plays a major

role in infection-related morphology and host recognition (Beauvais *et al.* 2013; King *et al.*

2017). A dozen of genes of *B. cinerea* encode for enzymes of the GH71 family (Amselem et

al., 2011). The T-specific GH71 CAZyme might therefore have been retained in the T

population as a mean to specifically facilitate infection of tomato by modification of the

fungal cell wall resulting in adaptive morphological changes or impairment of host

recognition.

As mentioned above, the McDonald-Kreitman tests also indicated that eight genes coding

for enzymes that detoxify reactive oxygen species showed signatures of positive selection in

the T population. During infection, *B. cinerea* encounters an oxidative burst, an early host

response that results in the death of plant cells. This mechanism is used in turn by *B. cinerea*

to achieve full virulence but this also implies that the fungus has to resist to this toxic

environment. The fungal oxidative stress response system includes detoxifying enzymes such

as superoxide dismutases (SODs) that convert O_2 into the less toxic H_2O_2, as well as

catalases, peroxidases and peroxiredoxins that convert H_2O_2. Additional non-enzymatic

mechanisms include the oxidation of compounds such as glutathione (Heller & Tudzynski

2011). In addition to the genes encoding SOD (BcSOD3), peroxidases (BcCCP1),

peroxiredoxins (BcPRX8) and other detoxifying enzymes sur as glutathione-S-transferases

(BcGST1, 9 and 24), a gene encoding the transporter BcAtrO also showed a signature of

positive selection in the T population. Inactivation of this gene previously suggested that it

allows the efflux of H_2O_2 and resistance this reactive oxygen species (Pane *et al.* 2008).

Altogether, our data suggest that the oxidative burst occurring in the *B. cinerea*/tomato

interaction is particularly challenging for the fungus.

Concluding remarks

We identified a population of *B. cinerea* specialized to tomato, which diverged from a

grapevine-associated population. Genome scans for selective sweeps and McDonald-

Kreitman tests revealed widespread signatures of positive selection that identified genes

that may contribute to the pathogen's adaptation to its tomato host. Candidate genes for

specialization to tomato were significantly enriched in those encoding cellulases, pectinases

and enzymes involved in the oxidative stress response, suggesting that the ability to degrade

the host cell wall and to cope with the oxidative burst are two key process in the *B.*

cinerea/tomato interaction. Our work sets the stage for future studies aiming to elucidate

the phenotypic and fitness effects of the candidate genes for specialization of *B. cinerea* to

tomato, for instance by knocking-out or replacing candidate genes for host specialization.

Acknowledgements

AM was supported by a grant from the Doctoral School « Sciences du Végétal », Université Paris-Saclay. We are grateful to INRAE-LIPM Bioinformatics Platform (Jérôme Gouzy, Sébastien Carrère, Erika Sallet, Ludovic Legrand) and Genotoul Bioinformatics Platforms Toulouse Occitanie (Bioinfo Genotoul, doi: 10.15454/1.5572369328961167E12) for providing computational support and resources. This work was supported by a grant overseen by the French National Research Agency (ANR) as part of the "Investissements d'Avenir" Programme (LabEx BASC; ANR-11-LABX-0034) and by the INRAE department SPE. The BIOGER unit also benefits from the support of "Saclay Plant Science-SPS" (ANR-17-EUR-0007).

References

Almagro Armenteros JJ, Tsirigos KD, Sonderby CK, *et al.* (2019) SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nature biotechnology* **37**, 420-423.

Amselem J, Cuomo CA, van Kan JA, *et al.* (2011) Genomic analysis of the necrotrophic fungal pathogens *Sclerotinia sclerotiorum* and *Botrytis cinerea*. *PLoS genetics* **7**, e1002230.

Atwell S, Corwin J, Soltis N, *et al.* (2015) Whole genome resequencing of Botrytis cinerea isolates identifies high levels of standing diversity. *Frontiers in microbiology* **6**, 996.

Atwell S, Corwin JA, Soltis N, *et al.* (2018) Resequencing and association mapping of the generalist pathogen Botrytis cinerea. *bioRxiv*, 489799.

Auton A, McVean G (2007) Recombination rate estimation in the presence of hotspots. *Genome research* **17**, 1219-1227.

Bardin M, Leyronas C, Troulet C, Morris CE (2018) Striking similarities between Botrytis cinerea from non-agricultural and from agricultural habitats. *Frontiers in plant science* **9**, 1820.

Beauvais A, Bozza S, Kniemeyer O, *et al.* (2013) Deletion of the α-(1,3)-glucan synthase genes induces a restructuring of the conidial cell wall responsible for the avirulence of *Aspergillus fumigatus*. *PLoS pathogens* **9**, e1003716.

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120.

Bonhomme M, Chevalet C, Servin B, *et al.* (2010) Detecting selection in population trees: the Lewontin and Krakauer test extended. *Genetics* **186**, 241-262.

Brito N, Espino JJ, González C (2006) The endo-β-1, 4-xylanase Xyn11A is required for virulence in Botrytis cinerea. *Molecular Plant-Microbe Interactions* **19**, 25-32.

Caseys C, Shi G, Soltis N, *et al.* (2020) Quantitative interactions drive *Botrytis cinerea* disease outcome across the plant kingdom. *bioRxiv*, 507491.

Cummings CA, Brinig MM, Lepp PW, van de Pas S, Relman DA (2004) Bordetella species are distinguished by patterns of substantial gene loss and host adaptation. *Journal of bacteriology* **186**, 1484-1492.

Decognet V, Bardin M, Trottin-Caudal Y, Nicot PC (2009) Rapid change in the genetic diversity of *Botrytis cinerea* populations after the introduction of strains in a tomato glasshouse. *Phytopathology* **99**, 185-193.

Dellaporta SL, Wood J, Hicks JB (1983) A plant DNA minipreparation: Version II. *Plant Molecular Biology Reporter* **1**, 19-21.

Derbyshire MC, Denton-Giles M, Hane JK, *et al.* (2019) A whole genome scan of SNP data suggests a lack of abundant hard selective sweeps in the genome of the broad host range plant pathogenic fungus Sclerotinia sclerotiorum. *PloS one* **14**, e0214201.

Elad Y, Pertot I, Prado AMC, Stewart A (2016) Plant hosts of Botrytis spp. In: *Botrytis–the fungus, the pathogen*
and its management in agricultural systems, pp. 413-486. Springer.

Emms DM, Kelly S (2015) OrthoFinder: solving fundamental biases in whole genome comparisons dramatically
improves orthogroup inference accuracy. *Genome biology* **16**, 157.

Espino JJ, Gutierrez-Sanchez G, Brito N, *et al.* (2010) The *Botrytis cinerea* early secretome. *Proteomics* **10**, 3020-
3034.

Fariello MI, Boitard S, Naya H, SanCristobal M, Servin B (2013) Detecting signatures of selection through
haplotype differentiation among hierarchically structured populations. *Genetics* **193**, 929-941.

Ferrer-Admetlla A, Liang M, Korneliussen T, Nielsen R (2014) On detecting incomplete soft or hard selective
sweeps using haplotype structure. *Molecular biology and evolution* **31**, 1275-1291.

Fisher MC, Henk DIA, Briggs CJ, *et al.* (2012) Emerging fungal threats to animal, plant and ecosystem health.
Nature **484**, 186.

Fournier E, Giraud T (2008) Sympatric genetic differentiation of a generalist pathogenic fungus, *Botrytis cinerea*,
on two different host plants, grapevine and bramble. *Journal of Evolutionary Biology* **21**, 122-132.

Garrison E, Marth G (2012) Haplotype-based variant detection from short-read sequencing. *arXiv:1207.3907 [q-*
bio].

Geib E, Markus Gressler M, Iuliia Viediernikova I, Falk Hillmann F, Ilse D. Jacobsen ID, Sandor Nietzsche S,
Christian Hertweck C, Matthias Brock M (2016) A Non-canonical Melanin Biosynthesis Pathway
Protects Aspergillus terreus Conidia from Environmental Stress. *Cell Chemical Biology* **23**, 587-597.

Giraud T (2006) Speciation: Selection against migrant pathogens: the immigrant inviability barrier in pathogens.
Heredity **97**, 316-318.

Giraud T, Gladieux P, Gavrilets S (2010) Linking the emergence of fungal plant diseases with ecological
speciation. *Trends in ecology & evolution* **25**, 387-395.

Giraud T, Refrégier G, Le Gac M, de Vienne DM, Hood ME (2008) Speciation in fungi. *Fungal genetics and*
biology : FG & B **45**, 791-802.

Gladieux P, Guerin F, Giraud T, *et al.* (2011) Emergence of novel fungal pathogens by ecological speciation:
importance of the reduced viability of immigrants. *Molecular ecology* **20**, 4521-4532.

Gladieux P, Ravel S, Rieux A, *et al.* (2018) Coexistence of multiple endemic and pandemic lineages of the rice
blast pathogen. *MBio* **9**.

Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD (2009) Inferring the joint demographic history
of multiple populations from multidimensional SNP frequency data. *PLoS genetics* **5**, e1000695.

Hartmann FE, McDonald BA, Croll D (2018) Genome-wide evidence for divergent selection between
populations of a major agricultural pathogen. *Molecular ecology* **27**, 2725-2741.

Heller J, Tudzynski P (2011) Reactive oxygen species in phytopathogenic fungi: signaling, development, and
disease. *Annual review of phytopathology* **49**, 369-390.

Huson DH, Bryant D (2005) Estimating phylogenetic trees and networks using SplitsTree 4. *Manuscript in*
preparation, software available from www. splitstree. org.

Inoue Y, Vy TTP, Yoshida K, *et al.* (2017) Evolution of the wheat blast fungus through functional losses in a host
specificity determinant. *Science* **357**, 80-83.

Johnson M, Zaretskaya I, Raytselis Y, *et al.* (2008) NCBI BLAST: a better web interface. *Nucleic acids research* **36**,
W5-W9.

Jones P, Binns D, Chang HY, *et al.* (2014) InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**, 1236-1240.

Khvorykh GV, Khrunin AV (2020) imputeqc: an R package for assessing imputation quality of genotypes and
optimizing imputation parameters. *BMC bioinformatics* **21**, 1-14.

King R, Urban M, Lauder RP, *et al.* (2017) A conserved fungal glycosyltransferase facilitates pathogenesis of
plants by enabling hyphal growth on solid surfaces. *PLoS pathogens* **13**, e1006672.

Krogh A, Larsson B, Von Heijne G, Sonnhammer ELL (2001) Predicting transmembrane protein topology with a
hidden Markov model: application to complete genomes. *Journal of molecular biology* **305**, 567-580.

Langridge GC, Fookes M, Connor TR, *et al.* (2015) Patterns of genome evolution that have accompanied host
adaptation in *Salmonella*. *Proceedings of the National Academy of Sciences of the United States of*
America **112**, 863-868.

Leyronas C, Bardin M, Berthier K, *et al.* (2018) Assessing the phenotypic and genotypic diversity of *Sclerotinia*
sclerotiorum in France. *European journal of plant pathology* **152**, 933-944.

Li H (2014) Toward better understanding of artifacts in variant calling from high-coverage samples.
Bioinformatics (Oxford, England) **30**, 2843-2851.

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*
25, 1754-1760.

Li H, Handsaker B, Wysoker A, *et al.* (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078-2079.

Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B (2014) The Carbohydrate-Active enZymes
database (CAZy) in 2013. *Nucleic acids research* **42**, D490-495.

Luo R, Liu B, Xie Y, *et al.* (2012) SOAPdenovo2: an empirically improved memory-efficient short-read de novo
assembler. *GigaScience* **1**, 18.

Ma Z, Michailides TJ (2005) Genetic structure of Botrytis cinerea populations from different host plants in
California. *Plant disease* **89**, 1083-1089.

Marchi N, Excoffier L (2020) Gene flow as a simple cause for an excess of high-frequency-derived alleles.
Evolutionary applications **13**, 2254-2263.

Maruthachalam K, Klosterman SJ, Kang S, Hayes RJ, Subbarao KV (2011) Identification of pathogenicity-related
genes in the vascular wilt fungus *Verticillium dahliae* by *Agrobacterium tumefaciens*-mediated T-DNA
insertional mutagenesis. *Molecular Biotechnology* **49**, 209-221.

McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq experiments
with respect to biological variation. *Nucleic acids research* **40**, 4288-4297.

McDonald JH, Kreitman M (1991) Adaptive protein evolution at the Adh locus in *Drosophila*. *Nature* **351**, 652-
654.

Mercier A, Carpentier F, Duplaix C*, et al.* (2019) The polyphagous plant pathogenic fungus *Botrytis cinerea*
encompasses host-specialized and generalist populations. *Environmental microbiology*.

Nakajima M, Akutsu K (2014) Virulence factors of *Botrytis cinerea*. *Journal of General Plant Pathology* **80**, 15-23.

Nielsen R, Beaumont MA (2009) Statistical inferences in phylogeography. *Molecular ecology* **18**, 1034-1047.

Nielsen R, Williamson S, Kim Y*, et al.* (2005) Genomic scans for selective sweeps using SNP data. *Genome*
research **15**, 1566-1575.

Nosil P, Vines TH, Funk DJ (2005) Perspective: reproductive isolation caused by natural selection against
immigrants from divergent habitats. *Evolution; international journal of organic evolution* **59**, 705-719.

Pane C, Rekab D, Firrao G, Ruocco M, Scala F (2008) A NOVEL GENE CODING FOR AN ABC TRANSPORTER IN
BOTRYTIS CINEREA (BOTRYOTINIA FUCKELIANA) IS INVOLVED IN RESISTANCE TO H$^+$ O$^+$. *Journal of*
Plant Pathology, 453-462.

Pavlidis P, Zivkovic D, Stamatakis A, Alachiotis N (2013) SweeD: likelihood-based detection of selective sweeps
in thousands of genomes. *Molecular biology and evolution* **30**, 2224-2234.

Perlin MH, Andrews J, Toh SS (2014) Essential letters in the fungal alphabet: ABC and MFS transporters and
their roles in survival and pathogenicity. *Advances in genetics* **85**, 201-253.

Plaza V, Lagües Y, Carvajal M*, et al.* (2015) bcpmr1 encodes a P-type Ca2+/Mn2+-ATPase mediating cell-wall
integrity and virulence in the phytopathogen Botrytis cinerea. *Fungal Genetics and Biology* **76**, 36-46.

Porquier A, Morgant G, Moraga J*, et al.* (2016) The botrydial biosynthetic gene cluster of *Botrytis cinerea*
displays a bipartite genomic structure and is positively regulated by the putative Zn(II)2Cys6
transcription factor BcBot6. *Fungal genetics and biology : FG & B* **96**, 33-46.

Reich D, Thangaraj K, Patterson N, Price AL, Singh L (2009) Reconstructing Indian population history. *Nature*
461, 489-494.

Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. *Bioinformatics (Oxford, England)* **26**, 139-140.

Rodriguez-Moreno L, Ebert MK, Bolton MD, Thomma BPHJ (2018) Tools of the crook- infection strategies of
fungal plant pathogens. *The Plant journal : for cell and molecular biology* **93**, 664-674.

Rosebeck S, Leaman DW (2008) Mitochondrial localization and pro-apoptotic effects of the interferon-inducible
protein ISG12a. *Apoptosis : an international journal on programmed cell death* **13**, 562-572.

Scheet P, Stephens M (2006) A fast and flexible statistical model for large-scale population genotype data:
applications to inferring missing genotypes and haplotypic phase. *The American Journal of Human*
Genetics **78**, 629-644.

Schulze-Lefert P, Panstruga R (2011) A molecular evolutionary concept connecting nonhost resistance,
pathogen host range, and pathogen speciation. *Trends in plant science* **16**, 117-125.

Seppey M, Manni M, Zdobnov EM (2019) BUSCO: assessing genome assembly and annotation completeness.
In: *Gene Prediction*, pp. 227-245. Springer.

Servedio MR, Van Doorn GS, Kopp M, Frame AM, Nosil P (2011) Magic traits in speciation: 'magic' but not rare?
Trends in ecology & evolution **26**, 389-397.

Simko I, Piepho H-P (2012) The area under the disease progress stairs: calculation, advantage, and application.
Phytopathology **102**, 381-389.

Skoglund P, Mallick S, Bortolini MC, *et al.* (2015) Genetic evidence for two founding populations of the
Americas. *Nature* **525**, 104-108.

Solovyev V, Kosarev P, Seledsov I, Vorobyev D (2006) Automatic annotation of eukaryotic genes, pseudogenes
and promoters. *Genome biology* **7**, S10.11-12.

Soltis NE, Atwell S, Shi G, *et al.* (2019) Interactions of tomato and Botrytis cinerea genetic diversity: Parsing the
contributions of host differentiation, domestication, and pathogen variation. *The Plant Cell* **31**, 502-
519.

Sonnhammer ELL, Von Heijne G, Krogh A A hidden Markov model for predicting transmembrane helices in
protein sequences **6**, 175-182.

Stoletzki N, Eyre-Walker A (2011) Estimation of the neutrality index. *Molecular biology and evolution* **28**, 63-70.

Stukenbrock EH, McDonald BA (2008) The origins of plant pathogens in agro-ecosystems. *Annual review of*
phytopathology **46**, 75-100.

Szpiech ZA, Jakobsson M, Rosenberg NA (2008) ADZE: a rarefaction approach for counting alleles private to
combinations of populations. *Bioinformatics (Oxford, England)* **24**, 2498-2504.

Valette-Collet O, Cimerman A, Reignault P, Levis C, Boccara M (2003) Disruption of Botrytis cinerea pectin
methylesterase gene Bcpme1 reduces virulence on several host plants. *Molecular Plant-Microbe*
Interactions **16**, 360-367.

van Kan JA, Stassen JH, Mosbach A, *et al.* (2017) A gapless genome sequence of the fungus *Botrytis cinerea*.
Molecular plant pathology **18**, 75-89.

Walker A (2016) Diversity within and between species of *Botrytis*. In: *Botrytis - The Fungus, the Pathogen and*
its Management in Agricultural Systems (eds. Fillinger S, Elad Y), pp. 91-125.

Walker A, Gladieux P, Decognet V, *et al.* (2015) Population structure and temporal maintenance of the
multihost fungal pathogen *Botrytis cinerea*: causes and implications for disease management.
Environmental microbiology **17**, 1261-1274.

Yang Z (1997) PAML: a program package for phylogenetic analysis by maximum likelihood. *Computer*
applications in the biosciences **13**, 555-556.

Yang Z (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Molecular biology and evolution* **24**, 1586-
1591.

Zerbino DR, Birney E (2008) Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome*
research **18**, 821-829.

Zhang C, Dong SS, Xu JY, He WM, Yang TL (2019a) PopLDdecay: a fast and effective tool for linkage
disequilibrium decay analysis based on variant call format files. *Bioinformatics (Oxford, England)* **35**,
1786-1788.

Zhang C, Dong SS, Xu JY, He WM, Yang TL (2019b) PopLDdecay: a fast and effective tool for linkage
disequilibrium decay analysis based on variant call format files. *Bioinformatics* **35**, 1786-1788.
Zhang MZ, Sun CH, Liu Y, *et al.* (2020) Transcriptome analysis and functional validation reveal a novel gene,
BcCGF1, that enhances fungal virulence by promoting infection-related development and host
penetration. *Molecular plant pathology* **21**, 834-853.
Zhao Z, Liu H, Wang C, Xu JR (2013) Comparative analysis of fungal genomes reveals different plant cell wall
degrading capacity in fungi. *BMC genomics* **14**, 274.
Zhong Z, Marcel TC, Hartmann FE, *et al.* (2017) A small secreted protein in *Zymoseptoria tritici* is responsible for
avirulence on wheat cultivars carrying the Stb6 resistance gene. *The New phytologist* **214**, 619-631.
915

Table 1. *Botrytis* isolates included in the study. Isolates from grapevine berries and tomatoes were collected in three regions of France between September 2005 and June 2007 for Champagne and Provence isolates (Walker *et al.* (2015), and between May and June 2009 for Occitanie isolates. Tomatoes in Occitanie were grown in plastic tunnels with sides open, whereas those from the Champagne and Provence areas were grown in glass-greenhouses. B05.10 is used as the reference isolate for genomic analysis (van Kan et al. 2016)

Species	Isolate ID	Other ID	Host, cultivar	Region, city
B. cinerea	Vv1	VC636	*Vitis vinifera,* Pinot noir	Champagne, Hautvilliers
	Vv2	VC095	*Vitis vinifera,* Pinot noir	Champagne, Vandières
	Vv3	VC621	*Vitis vinifera,* Pinot noir	Champagne, Hautvilliers
	Vv4	VC671	*Vitis vinifera,* Pinot noir	Champagne, Hautvilliers
	Vv5	VC224	*Vitis vinifera,* Pinot meunier	Champagne, Courteron
	Vv6	VC624	*Vitis vinifera,* Pinot noir	Champagne, Hautvilliers
	Vv7	ACBER342	*Vitis vinifera,* Grenache	Provence, Berre
	Vv8	ACBER356	*Vitis vinifera,* Grenache	Provence, Berre
	Vv9	ACBER358	*Vitis vinifera,* Grenache	Provence, Berre
	Vv10	ACSAR333	*Vitis vinifera,* Grenache	Provence, Sarrians
	Vv11	ACSAR334	*Vitis vinifera,* Grenache	Provence, Sarrians
	Vv12	ACSAR335	*Vitis vinifera,* Grenache	Provence, Sarrians
	Vv13	ACSAR342	*Vitis vinifera,* Grenache	Provence, Sarrians
	Vv14	ACSAR354	*Vitis vinifera,* Grenache	Provence, Sarrians
	Vv15	ACSAR357	*Vitis vinifera,* Grenache	Provence, Sarrians
	Vv16	VC610	*Vitis vinifera,* Pinot noir	Champagne, Hautvilliers
	Sl1	VA714	*Solanum lycopersicum,* Moneymaker	Champagne, Foissy-sur-Vanne
	Sl2	VC800	*Solanum lycopersicum,* Moneymaker	Champagne, Courceroy
	Sl3	VC806	*Solanum lycopersicum,* Moneymaker	Champagne, Courceroy
	Sl4	AABER19	*Solanum lycopersicum,* Alison	Provence, Berre
	Sl5	ACBER304	*Solanum lycopersicum,* Alison	Provence, Berre
	Sl6	ACPIE306	*Solanum lycopersicum,* Hipop	Provence, Pierrelatte
	Sl7	ADPIE463	*Solanum lycopersicum,* Hipop	Provence, Pierrelatte
	Sl8	ADPIE475	*Solanum lycopersicum,* Hipop	Provence, Pierrelatte
	Sl9	65_TT8	*Solanum lycopersicum,* Brenda	Occitanie, Alenya
	Sl10	5_TT8	*Solanum lycopersicum,* Brenda	Occitanie, Alenya
	Sl11	13_TT8	*Solanum lycopersicum,* Brenda	Occitanie, Alenya

	Sl12	9_TT8	*Solanum lycopersicum*, Brenda	Occitanie, Alenya
	Sl13	66_TT8	*Solanum lycopersicum*, Brenda	Occitanie, Alenya
	Rf1	VC902	*Rubus fruticosus*, wild	Champagne, Foissy-sur-Vanne
	Rf2	VC399	*Rubus fruticosus*, wild	Champagne, Courteron
	Hm1	MSN-Bot 2556	*Hydrangea macrophylla*, Leuchtfeuer	Anjou, Angers
	B05.10	-	-	-
B. fabae	Bfab	MSN-Bot 2220	*Vicia faba*	Region of Tunis (Tunisia)

Table 2. Summary statistics of genomic variation in three clusters of *Botrytis cinerea*

Cluster	S	π	θ	AR	PAR	D	H	LD50	ρ
T	37763	0.0027	0.0026	1.094 (0.0002)	0.073 (0.0002)	0.309	0.451	8600	0.0108
G1	26769	0.0018	0.0017	1.078 (0.0002)	0.085 (0.0002)	0.167	0.230	3100	0.0178
G2	14307	0.0030	0.0027	1.104 (0.0003)	0.071 (0.0002)	1.470	0.140	16800	0.2047

S, number of segregating sites; π, nucleotide diversity per site; θ, Watterson's estimate of the population mutation parameter per site; ρ, population recombination parameter per site; AR, allelic richness (standard error of the mean); PAR, private allele richness (standard error of the mean); D, Tajima's D; H, Fay and Wu's standardised H; LD50, distance (in bp) at which linkage disequilibrium reaches half of its maximum value. Only core chromosomes BCIN01 to BCIN16 were included in calculations. Per site estimates of π, ρ and θ were computed by summing across chromosomes and dividing by number of sites covered. Tajima's D and Fay and Wu's standardised H were computed across 10kb windows, and averaged. AR and PAR were computed using a generalized rarefaction approach and a standardized sample size of two haploid genomes.

Figure Captions

Figure 1. Population subdivision inferred based on SNPs identified in 32 isolates of *Botrytis cinerea* collected on tomato (red, Sl prefix, *Solanum lycopersicum*), grape (green, Vv prefix, *Vitis vinifera*), bramble (black, Rf prefix, *Rubus fruticosus*) and hydrangea (black, Hm prefix, *Hydrangea macrophylla*). (A) Ancestry proportions in K clusters, as estimated with the sNMF program. Each multilocus genotype is represented by a vertical bar divided into K segments, indicating membership in K clusters. (B) Neighbor-net phylogenetic network estimated with SPLITSTREE, with one isolate of *B. fabae* (Bfab) used as the outgroup. Reticulations indicate phylogenetic conflicts caused by recombination or incomplete lineage sorting. (C) Principal component analysis showing first four principal components PC1, PC2, PC3 and PC4. Isolate B05.10 in (A) and (B) is the reference genome for *B. cinerea* (van Kan et al. 2016)

Figure 2. Site frequency spectra estimated in populations T2 and Vv1 based on 85,415 SNPs using the python package DADI, projecting sample sizes to seven haploid genomes. Projection consists in averaging over all possible re-samplings of the larger sample size data, thus biallelic positions with data for less than seven individuals are not included in calculations and not counted as SNPs.

Figure 3. Boxplots representing the extent of aggressiveness (% relative to the reference isolate BC1) on tomato of the *Botrytis cinerea* isolates collected on grapevine (n=16) and tomato (n=13). For each boxplot, mean (crosses), median (horizontal lines), values of the aggressiveness index (circles), 25-75% quartiles, and maximum and minimum values are represented.

Figure 4. Genome scans for selective sweeps in the T and G1 populations of *Botrytis cinerea*, parasitizing tomato and grapevine, respectively. (A) and (C): Composite likelihood ratio

(CLR) estimated using the SᴡᴇᴇD software (Nielsen *et al.* 2005; Pavlidis *et al.*, 2013) in T and G1, respectively. (B) and (D) Number of segregating sites by length, nS_L, estimated using the ɴSL software (Ferrer-Admetlla *et al.* 2014) in T and G1, respectively. (E) hapFLK statistic estimated using the HᴀᴘFLK software (Bonhomme *et al.* 2010; Fariello *et al.* 2013). Horizontal dashed black lines represent the top 5%. Vertical dashed grey lines represent the boundaries of the 16 chromosomes and the chromosome names are set along the x axis. SNPs in chromosomes with names ending with odd numbers are represented in dark colors, while SNPs in chromosomes with names ending with even numbers are represented in light colors. Black dots represent SNPs belonging to the following set: [(top 5% CLR population G1) ∪ (top 5% CLR population T) ∪ (top 5% nSL population G1) ∪ (top 5% nSL population T)] ∩ [top 5% hapFLK].