Recognition of high-risk HPV E6 oncoproteins by 14-3-3 proteins studied by
interactomics and crystallography

Gergo Gogl[1]*, Kristina V. Tugaeva[2]*, Pascal Eberling[1], Camille Kostmann[1], Gilles Trave[1†],
Nikolai N. Sluchanko[2†]

1 Equipe Labellisee Ligue 2015, Department of Integrated Structural Biology, Institut de
Genetique et de Biologie Moleculaire et Cellulaire (IGBMC), INSERM U1258/CNRS UMR
7104/Universite de Strasbourg, 1 rue Laurent Fries, BP 10142, F-67404 Illkirch, France

2 A.N. Bach Institute of Biochemistry, Federal Research Center of Biotechnology of the
Russian Academy of Sciences, 119071, Moscow, Russia

Email addresses of the authors:

goglg@igbmc.fr

kri94_08@mail.ru

eberling@igbmc.fr

kostmanc@igbmc.fr

*Contributed equally: G.G. and K.V.T.

†Corresponding authors: G.T.: traveg@igbmc.fr ; N.N.S.: nikolai.sluchanko@mail.ru

Abstract

In tumors induced by high-risk mucosal human papillomaviruses (hrm-HPVs), HPV E6 oncoproteins inhibit apoptotic processes and sustain cell proliferation. E6 from all hrm-HPVs harbor a C-terminal short PDZ domain-binding motif (PBM), whose phosphorylation down-regulates PDZ binding but triggers E6 binding to 14-3-3 proteins. Here we classify PBMs of E6 proteins depending on their principle ability to be phosphorylated and subsequently acquire a 14-3-3-binding motif III consensus, (pS/pT)XX-COOH. Systematic competitive fluorescence polarization measurements show that the PBMs from four selected E6 oncoproteins bind all seven human 14-3-3 isoforms with distinct, wide-ranging affinities, obeying remarkable trends assigned to 14-3-3 isoform specificity and small E6 sequence variations. We crystallized the hrm-HPV18 E6 PBM bound to 14-3-3ζ, revealing a 14-3-3-motif III complex at 1.9 Å resolution. Using fluorescence polarization and crystallography, we also demonstrate that fusicoccin, a molecule that reinforces many known 14-3-3 complexes, destabilizes the 14-3-3-E6 interaction, indicating the druggability of that complex.

Keywords:

Protein chimera; protein-protein interactions; phosphopeptides; phosphorylation; crystal structure; binding affinity

INTRODUCTION

Papillomaviruses (PVs) are a large group of small oncogenic DNA viruses infecting various vertebrates including humans (human papillomaviruses, HPVs). PVs target mucosal or cutaneous epithelium cells and induce their proliferation, to amplify their circular double-stranded DNA genome and package it into virion particles ready for a new infectious cycle. 225 HPV types currently annotated in the Papillomavirus Episteme (PaVE) database (https://pave.niaid.nih.gov/) are classified into five phylogenetically distinct genera (alpha, beta, gamma, mu and nu), of which alpha HPVs display almost exclusively mucosal tropism [1]. While most HPV only generate benign proliferative events such as warts, lesions or condylomas, a subset of about 20 HPV types belonging to two alpha species have been associated to cancers and are therefore epidemiologically categorized as "high-risk" [2, 3, 4]. HPV16 and 18 are the most prevalent, causing up to 80% of squamous cervical carcinomas [5], whereas HPV16 is also highly prevalent in HPV-positive tumors of the oropharynx and the anus [6].

The E6 protein is one of the two main HPV oncoproteins expressed at the early stage of infection. In HPV-transformed cells, E6 participates in counteracting apoptosis, altering differentiation pathways, polarity and adhesion properties, and thereby sustains cell proliferation [1, 7]. Inhibition of E6 in HPV-positive cell lines results in the cell growth arrest and induces apoptosis or rapid senescence [8, 9, 10, 11].

E6 has been found to interact with numerous distinct cellular target proteins involved in a variety of cellular functions [12]. Importantly, E6 proteins from distinct HPV species recognize distinct subsets of the full panel of potential E6 targets [13]. This likely contributes to the particular biological traits of each HPV type in terms of tropism, viral cycle, or pathogenicity.

E6 proteins of most mammalian PVs comprise two zinc-binding domains (E6N and E6C) separated by a linker helix [1] (Fig. 1A), which altogether form a charged-hydrophobic pocket that recognizes LxxLL motifs interspaced with acidic residues, found in a variety of host cellular proteins [12]. Hrm-HPV E6 recruits simultaneously the LxxLL motif of ubiquitin ligase E6-Associated Protein (E6AP) and, via a distinct interface, the central "core" domain of tumor suppressor p53 [14]. The resulting E6-E6AP-p53 trimer drives the ubiquitinylation and subsequent proteasomal degradation of p53 [15]. In HPV-transformed cells, this leads to low levels of p53 protein that probably contribute to the tumorigenic phenotype. It is supposed that E6 similarly stimulates the E6AP-mediated degradation of other host proteins [15, 16].

In addition to their LxxLL-binding pocket and p53-binding surface, hrm-HPV E6 proteins harbor a C-terminal PDZ domain-binding motif (PBM) [17] (Fig. 1A), which remarkably is absent in most low-risk non-cancer-causing HPV E6 proteins [18]. The intrinsically disordered PBM enables the E6 interaction with a variety of host PDZ domain-containing proteins involved in the regulation of cell polarity, adhesion, differentiation or survival [19]. These interactions of E6 with PDZ proteins are potentially modulated by PBM phosphorylation. Several E6 PBMs are phosphorylatable *in vitro* at variable levels of specificity and efficiency by basophilic kinases, such as PKA [18, 20, 21], PKB/Akt [18, 21], CamKII [21], or Chk1, and by the glutamine-directed ATM/ATR kinases [22]. E6 PBMs from high-risk mucosal HPV types 16, 18, 33, 35 (Fig. 1B) are good substrates for PKA *in vitro* [18, 21, 23]. In addition, E6 of HPV16 and HPV18 (hereinafter called 16E6 and 18E6) are metabolically labeled by ^{32}P upon transient transfection *in vivo*, and forskolin promotes such labeling [23]. Phosphorylation of several HPV-E6 proteins at the PBM was confirmed by using phospho-specific antibodies raised against the C-terminal PBM of 18E6 [18, 21]. It was also shown that the level of 18E6 phosphorylation remains low in the normal cell cycle, until the activation of stress-response kinases and those involved in the DNA damage response (see in particular [22]).

While phosphorylation of HPV-E6 PBMs generally reduces their binding to PDZ domains [24], it turns several of them into candidate binding sites for 14-3-3 proteins [18, 21]. Despite being encoded by seven distinct genes, human 14-3-3 proteins share highly similar sequences and biochemical properties. Hence they are commonly referred to as "isoforms", individually named β, γ, ε, ζ, η, σ, and τ (beta, gamma, epsilon, zeta, eta, sigma and tau) [25], not to be confused with their splice variants. 14-3-3 proteins form homo- and heterodimers [26] characterized by their ability to bind phosphopeptides [27, 28]. 14-3-3 are involved in the regulation of multiple cellular processes including apoptosis, cell division and signal transduction [29]. Phosphorylated 14-3-3-binding sequences usually correspond to internal motifs I RSX(pS/pT)X(P/G) and II RXY/FX(pS/pT)X(P/G) [28] and to the C-terminal motif III (pS/pT)X$_{0-2}$-COOH [30, 31], where pS/pT denotes phosphorylated serine or threonine and X denotes any amino acid. The regulation by 14-3-3 binding typically protects 14-3-3 targets from dephosphorylation and degradation, affects their activity, intracellular localization, and interactions with other proteins by occlusion of the phosphorylated segments [32]. Given the involvement of 14-3-3 in the development of various pathological conditions associated with the deregulation of the corresponding protein-protein interactions, 14-3-3/phosphotarget complexes are considered with increasing attention [33].

Phosphorylated 16E6, 18E6, and 31E6 have been previously shown to bind to 14-3-3 proteins, albeit with different efficiencies [18, 21, 24]. In contrast, the binding of unphosphorylated or phosphomimetic E6 PBMs to 14-3-3 proteins was not detectable [21, 24].

This suggests that the phosphorylation-dependent interaction with 14-3-3 proteins is a general feature of various E6 PBMs.

Here we classified PBMs of hrm-HPV E6 proteins depending on their principle ability to be phosphorylated and subsequently acquire a 14-3-3-binding motif III consensus, (pS/pT)XX-COOH. We performed systematic quantitative binding assays of four phosphorylatable E6 PBMs with all seven human 14-3-3 isoform homodimers and solved a high-resolution crystal structure of the HPV18 E6 PBM complex with 14-3-3ζ. This combination of approaches has allowed us to rationalize differences in the affinities of tested PBMs and fully confirm this by mutated peptides. Finally, we used binding assays and crystallography to analyze the impact of natural toxin fusicoccin on the 14-3-3-E6 PBM interaction, which proved that the 14-3-3-E6 complexes are nominally druggable by small molecules.

RESULTS

PDZ-binding motifs of E6 proteins can be classified according to their phosphorylation and 14-3-3-binding propensity

Among all 225 HPV type E6 proteins curated in the PaVE database (https://pave.niaid.nih.gov/), 31 E6 proteins from the α-genera HPV, having predominantly mucosal tropism, possess a C-terminal PDZ-binding motif (PBM) that corresponds to the class 1 PBM [X(S/T)X(L/V/I/C)-COOH, where X is any amino acid residue [34, 35]]. It has previously been shown that the E6 PBMs are susceptible to phosphorylation by protein kinases on the conserved Ser/Thr residue at the antepenultimate C-terminal position, which contributes to the class 1 PBM consensus [18, 21, 23]. This phosphosite is preceded by arginine residues in most of the HPV-E6 PBM sequences with recognizable patterns of the two common basophilic kinase substrate consensus motifs of R(X/R)X(S/T) and RXRXX(S/T) [36, 37]. Additional Arg residues, e.g. at position -6 are most probably less involved in kinase recognition and more important for binding to other partners. For instance, Arg -6 of 18E6 is important for PDZ recognition but its mutation does not affect its phosphorylation efficiency by PKA, while mutation of Arg -5 and -4 blocks 18E6 PBM phosphorylation [38]. Considering the basic consensus motifs, these 31 E6 PBMs can be readily divided into three subgroups: Subgroups 1 and 2 having sequences prone to phosphorylation by the basophilic kinases and the third, orphan subgroup with a less predictable phosphorylation propensity (Fig. 1C). Accordingly, the PBM of HPV66 E6, which in our classification falls into the orphan group, is normally recalcitrant to phosphorylation by PKA *in cellulo* but becomes prone to it upon introduction of Arg residues at its positions -5 and -4 [38]. In addition, more than half of the identified HPV-E6 PBM sequences harbor candidate phosphorylation sites of DNA damage response ATM/ATR kinases with a consensus of (S/T)Q [39].

Therefore, the regulation of subgroup 1 and 2 E6 PBMs by phosphorylation is likely a rather general phenomenon. In line with earlier observations [21, 24, 40], the phospho-PBM sequences from subgroups 1 and 2 can ideally match the C-terminal 14-3-3-binding motif III [30] (Fig. 1B and C). It is therefore likely that phosphorylation of the C-terminal PBMs from HPV-E6 types belonging to subgroups 1 and 2 by different kinases triggers their recognition by 14-3-3 proteins [18, 21, 22, 24].

Fig. 1. Features of the structure and the C-terminal PDZ-binding motif of E6 proteins. **A.** The domain organization (top) and three dimensional structure (bottom) of HPV16 E6 (16E6) showing the location of the phosphorylatable PDZ-binding motif (PBM). N-terminal (E6N) and C-terminal (E6C) domains are color coded, structural Zn atoms are in black, PBM is in cyan, phosphorylatable Thr -2 is in red. **B.** E6 PBMs from high-risk mucosal HPV types overlap with the 14-3-3-binding motif III [30]. The positions are numbered above, according to conventional PBM numbering. **C.** Classification of 31 PBM-containing HPV-E6 proteins based on the correspondence of their C-terminal PBMs to the consensus motifs phosphorylatable by basophilic kinases (subgroups 1 and 2). The third, orphan subgroup comprises PBMs of HPV-E6 proteins whose phosphorylation is less certain. Bold font indicates the E6 PBMs used in this work. Note that many PBMs overlap with recognition motifs (S/TQ) for phosphorylation by DNA damage response kinases ATM/ATR (TQ and SQ sites highlighted). Below are shown Weblogo diagrams [41] for the PBMs within each of the three subgroups (positions are numbered according to the PBM convention).

E6 PBMs from four distinct hrm-HPV types display wide-ranging affinities to the seven 14-3-3 proteins with parallel binding profiles

Four phospho-PBMs from E6 proteins of HPV types 16, 18, 33 and 35 belonging to subgroups 1 and 2 (as defined in Fig. 1C) were analyzed for their interaction with the all seven human 14-3-3 homodimers. For comparison, we also measured two non-viral phospho-PBMs originating from protein kinase RSK1 [24]. We used a competitive fluorescence polarization assay that measures the displacement of a fluorescent tracer 14-3-3-binding peptide in the presence of an increasing amount of a competitor molecule (all binding curves are shown in Supplementary Fig. 1A).

All phospho-PBMs (hereinafter called p16E6, p18E6, p33E6, p35E6, RSK1_-1P, and RSK1_-2P) detectably bound to 14-3-3 proteins, in sharp contrast to their unphosphorylated counterparts. The interactions between E6 phospho-PBMs and 14-3-3 proteins spanned very wide affinity ranges, from just below 1 µM (p33E6–14-3-3γ) to above 300 µM (Fig. 2A and Supplementary Fig. 1). Such large binding affinity differences are noteworthy since the four E6 PBM sequences are very similar (Fig. 1B), and all 14-3-3 isoforms share highly conserved phosphopeptide-binding grooves.

Remarkably, the six phospho-PBMs showed consistent hierarchy in their relative binding preferences towards any of the seven 14-3-3 isoforms, albeit with a systematic overall shift in affinity from one peptide to another. The seven 14-3-3 isoforms clustered as four groups of decreasing affinity, in a conserved order from the strongest to the weakest phospho-PBM binder: gamma, eta, zeta/tau/beta and epsilon/sigma (γ, η, ζ/τ/β, and ε/σ) (Fig. 2B). These conserved relative affinity shifts can be quantified by calculating, for two distinct 14-3-3 isoforms, their differences of free energy of binding ($\Delta\Delta G$) towards each individual phosphopeptide, then calculating the average difference ($\Delta\Delta G_{av}$) with its standard deviation (Fig. 2C). Between the strongest and the weakest binders (isoforms γ and σ, respectively) the average phosphopeptide-binding energy difference is $\Delta\Delta G_{av}$ = -5.1 ± 1.3 kJ/mol, roughly corresponding to a 11-fold K_D ratio.

Noteworthy enough, another study [42] also measured the binding of seven human 14-3-3 isoforms to the completely unrelated cystic fibrosis transmembrane conductance regulator (CFTR) phosphopeptide CFTR-R7, representing an internal 14-3-3-binding motif. A roughly comparable hierarchy of affinities was observed, also showing a 11-fold K_D ratio between the strongest binder (gamma isoform, K_D = 1 µM) and the weakest binder (epsilon isoform, K_D = 11 µM); both values being remarkably close to those obtained herein for the strongest-binding E6 phospho-PBM, p33E6. Similar trends were also detected upon interaction of the Leucine-Rich Repeat Kinase 2 (LRRK2) internal motif phosphopeptides with 14-3-3 isoforms, when gamma and eta were the strongest and epsilon and sigma were the weakest binders [43]. The K_D ratio for the gamma and sigma isoforms averaged for the three of LRRK2 phosphopeptides (~20) [43] is remarkably similar to the 11-fold difference described here for the E6 PBM interaction with the strongest and weakest human 14-3-3 isoforms. Therefore, our data support the idea that specificity of 14-3-3 isoforms for protein partners follows a general trend.

The seven 14-3-3 isoforms also showed consistent profiles in their relative binding preferences towards any of the four E6 phospho-PBMs. For each 14-3-3 isoform, the four phospho-PBMs systematically rank the same way from the strongest to the weakest binder: p33E6, p18E6, p16E6 and p35E6 (Fig. 2C). The average 14-3-3 binding free

energy difference between p33E6 and p35E6 was $\Delta\Delta G_{av}$ = -10.9 ± 0.7 kJ/mol, roughly corresponding to a 100-fold K_D ratio.

A

	K_D (µM) ± std						
	14-3-3γ	14-3-3η	14-3-3ζ	14-3-3τ	14-3-3β	14-3-3ε	14-3-3σ
pHPV33E6	0.94 ± 0.25	1.98 ± 0.15	2.72 ± 0.39	3.52 ± 0.28	4.05 ± 0.65	10.3 ± 2.39	6.77 ± 1.13
pHPV18E6	11.1 ± 0.6	23.6 ± 6.2	22.4 ± 1.3	37.8 ± 6.0	42.2 ± 6.3	101 ± 20	139 ± 24
pHPV16E6	37.2 ± 3.0	80.5 ± 16.9	73.6 ± 19.1	144 ± 28	159 ± 45	>300	>300
pHPV35E6	125 ± 6	163 ± 19	191 ± 18	233 ± 14	>300	>300	>300
pHPV35E6$_{T-6R}$	71.2 ± 6.7	85.0 ± 20.6	139 ± 21	194 ± 37	179 ± 29	270 ± 20	>300
pHPV35E6$_{E-1A}$	9.31 ± 0.72	12.4 ± 1.5	17.6 ± 3.6	21.5 ± 1.2	20.2 ± 1.4	31.9 ± 4.2	44.1 ± 13.3
pHPV35E6$_{T-6R, E-1A}$	3.27 ± 0.16	4.77 ± 0.53	7.32 ± 0.42	5.84 ± 0.43	7.06 ± 0.51	13.0 ± 2.0	18.9 ± 1.9
RSK1_-1P	0.31 ± 0.03	0.46 ± 0.06	0.96 ± 0.10	0.78 ± 0.13	0.74 ± 0.04	1.58 ± 0.21	1.71 ± 0.23
RSK1_-2P	0.20 ± 0.01	0.66 ± 0.05	1.54 ± 0.09	0.45 ± 0.08	0.96 ± 0.03	1.22 ± 0.16	3.57 ± 0.32

B

C

$\Delta\Delta G_{av}$ = -0.71±1.19   -1.50±0.63   -0.34±0.66   -0.14±1.39   -0.87±0.78   -1.35±0.82 (kJ/mol)

14-3-3σ < 14-3-3ε < 14-3-3β < 14-3-3τ < 14-3-3ζ < 14-3-3η < 14-3-3γ

$\Delta\Delta G_{av}$ =   -2.06±0.67   -3.09±0.15   -5.98±0.65   (kJ/mol)

HPV35E6 < HPV16E6 < HPV18E6 < HPV33E6

Fig. 2. Summary of affinity measurements carried out in this study. **A**. Affinities of four selected HPV-E6 phospho-PBMs, p35E6 mutants and RSK1 phosphopeptides towards the seven human 14-3-3 isoforms as determined by fluorescence polarization using FITC-labeled HSPB6 phosphopeptide as a tracer. Apparent K_D values determined from competitive FP experiments are presented. **B**. The heatmap representation of the data on panel A showing the affinity trends in the interaction profiles between 14-3-3 isoforms and four HPV-E6 phospho-PBMs from weakest (white) to strongest (red). **C**. Averaged ΔΔG values between 14-3-3 isoforms or E6 phospho-PBM pairs, calculated based on their observed order of binding affinities (from weakest to strongest). Individual K_D values from Supplementary Fig. 1 were first converted into ΔG values (at T=295 K; excluding cases when K_D > 300 µM) and average ΔΔG values (ΔΔG$_{av}$) were calculated between the indicated motifs/isoforms.

Atomic structure reveals the 14-3-3ζ / 18E6 PBM interface

To get structural insight into the 14-3-3ζ interactions with 18E6 PBM, we designed, as previously reported [44, 45], a protein-peptide chimera, which we co-expressed with PKA and purified from *E. coli* cells by using an engineered cleavable His$_6$ tag (Supplementary Fig. 2A-C). The 14-3-3ζ core was modified to block phosphorylation of the semiconserved Ser58 located in the dimer interface [46] and to facilitate crystallization [47] and was tethered to

the 18E6 phosphorylatable peptide RRRETQV-COOH. This approach was justified
because of the disordered nature of the E6 PBMs [48, 49].

The 14-3-3ζ-18E6 PBM chimera was stoichiometrically phosphorylated by co-expressed
PKA, which gave a distinct downward shift on native PAGE (Supplementary Fig. 2D), and
readily formed diffraction-quality crystals (all statistics is presented in Table 1). Structure
determination was carried out at a 1.9 Å resolution with a single 14-3-3 dimer in the
asymmetric unit (Supplementary Fig. 3A). The 18E6 phosphopeptide of this dimer was
bound *in-trans* in the amphipathic 14-3-3 grooves of the crystallographic symmetry
neighbor molecule, resulting in swapped phosphopeptides (Supplementary Fig. 3A). Such
arrangement, previously observed for 14-3-3σ chimeras with other phosphopeptides [44, 50]
provided us with high-resolution insight into the 14-3-3/18E6 PBM interaction (Fig. 3A-C).
All residues of the 18E6 phosphopeptides could be traced including the conformation of the
C-terminal carboxyl-group. The phosphopeptide is stabilized by multiple polar interactions
involving those between phospho-Thr156 (position -2) and conserved residues of the basic
pocket in the amphipathic groove of 14-3-3 (Arg56, Arg129, Lys49 and Tyr128, 14-3-3ζ
numbering), the phosphopeptide backbone interactions to Asn224 and Asn173, and a
remarkable polar contact between the carboxyl-group of the peptide and the side chains of
Lys120 and one of the two alternative conformations of Ser45 of 14-3-3ζ (Fig. 3C). The
carboxyl-group orientation is further stabilized by *in-cis* H-bond with the side chain of the
preceding residue Gln157 (position -1) (Fig. 3C). The side chain of the C-terminal Val158
(position 0) of the peptide makes a hydrophobic contact with Val46 of 14-3-3ζ. A second
remarkable hydrophobic contact is observed between the side chain of Val176 of 14-3-3ζ
and the methyl-group of phospho-Thr156 (position -2), which hints at a more stable binding
of pThr than would be in the case of pSer.

Importantly, the conformation of 18E6 phosphopeptide bound to 14-3-3ζ within the chimera
is practically identical (RMSD = 0.17 Å upon superimposition of the six core Cα atoms of
the peptides) to the 14-3-3σ-bound conformation of a synthetic 16E6 phosphopeptide
reported very recently at a much lower resolution (Fig. 3D) [24]. In both cases, the carboxyl-
group gives a polar contact to the Lys120 side chain and the hydrophobic side chain of the
last peptide residue (Val158 in 18E6 and Leu151 in 16E6) faces the side chain of Val46 of
14-3-3. This validates the chimeric approach [44] as a strategy to obtain high-resolution
crystal structures of 14-3-3/phosphopeptide complexes. Arg residues that are likely to be
involved in kinase recognition also contribute to 14-3-3 binding. The Arg residue occupying
position -4 forms an *in-cis* interaction with the phosphoryl group, and the Arg residue
occupying position -5 forms in both 16E6 and 18E6 a π-stacking interaction with Arg60 of
14-3-3 (Fig. 3D). Most of interface contacts of the 18E6/14-3-3ζ and 16E6/14-3-3σ
complexes are similar, suggesting that the two current crystal structures can also serve as
templates to build accurate homology models of other 14-3-3/E6 complexes and, more
generally, any motif III complexes with a pS/pTXX-COOH consensus.

However, some noteworthy differences appear in a subset of the crystallographic
conformers of 14-3-3/16E6 and 14-3-3/18E6 complexes. On the one hand, in 1 of the 4
conformers observed in the asymmetric unit of the 14-3-3σ/16E6 crystal, the side chains of
Arg -7 (Gln in 18E6) and Glu -3 form an additional *in-cis* salt bridge (Fig. 3D). On the other
hand, Arg -6 of 18E6 (Thr in 16E6) mediates a bipartite interaction with 14-3-3 in most of
the observed conformers. It simultaneously interacts with the carbonyl of Asp223 and
participates in a water-mediated interaction with Asn224 (Fig. 3C and D).

Fig. 3. Molecular interface between 14-3-3ζ and phosphorylated 18E6 PBM at a 1.9Å resolution. **A.** An overall view on the 14-3-3ζ dimer (subunits are in tints of grey) with two bound 18E6 phosphopeptides (cyan sticks). **B.** A magnified view on one of the amphipathic grooves of 14-3-3ζ showing the conformation of the 18E6 phosphopeptide and the corresponding $2F_o$-F_c electron density maps contoured at 1σ. Positions are numbered according to the PBM convention. **C.** Polar contacts (dashed lines) and hydrophobic interactions (semitransparent spheres) stabilizing the bound 18E6 peptide conformation. **D.** An overlay of the two 14-3-3 bound phosphopeptides from 16E6 (6TWZ.pdb) and 18E6 (this work) showing the similarity of the conformation. # denotes the C-terminus (-COOH). w – the water molecule, π – π-stacking interaction. Important positions are numbered according to the PBM convention. **E.** Averaged ΔΔG values between 14-3-3 isoforms or 35E6 phospho-PBM pairs, calculated based on their observed order of binding affinities (from weakest to strongest). Individual K_D values from Supplementary Fig. 1 were first converted into ΔG values (at T=295 K; excluding cases when K_D > 300 µM) and average ΔΔG values ($ΔΔG_{av}$) were calculated between the indicated motifs/isoforms.

Rescue of the weakest E6-14-3-3 interaction by rational design

We hypothesized that the remarkable affinity differences of the four studied E6 phospho-PBMs ($\Delta\Delta G_{av}$ = -10.9 ± 0.7 kJ/mol between the weakest and the strongest 14-3-3-binder) might result from two opposed mechanisms: interface peptide-14-3-3 contacts favoring complex formation versus intra-peptide contacts within the unbound peptide disfavoring it. On the one hand, position -6 is an Arg in the two strongest 14-3-3-binders (18E6 and 33E6) versus a Thr in the weakest ones (35E6 and 16E6). Accordingly, the crystal structures have shown that Arg -6 can mediate more interactions with the generic 14-3-3 interface (Fig. 3D). On the other hand, all E6 phospho-PBMs have a delicate charge distribution, with an acidic C-terminus (that is also involved in PDZ-domain binding) and a basic N-terminal segment (that is also involved in kinase recognition). These local charged segments may form transient *in-cis* interactions within the unbound phosphopeptide, so-called "charge clamps" [51], potentially unfavorable for 14-3-3 binding. We speculated that Glu -1 in p35E6, the weakest 14-3-3 binder, might participate in such a charge clamp, thereby disfavoring the binding.

To test these mechanisms, we synthesized three variants of p35E6. The first variant contained a T-6R substitution, which in principle could allow a more stable bound conformation, but may also stabilize charge-clamps in the free form of the motif. The second variant contained an E-1A substitution, which in principle could destabilize *in-cis* charge-clamps. A third variant contained both substitutions. All substitutions turned out to reinforce the binding affinities of 35E6 without altering the apparent preferences of the different 14-3-3 isoforms (Fig. 2A and Fig. 3E). Taken individually, T-6R moderately increased binding ($\Delta\Delta G_{av}$ = -1.1 ± 0.5 kJ/mol, 1.5-fold K_D ratio), while E-1A strongly reinforced it ($\Delta\Delta G_{av}$ = -5.1 ± 0.2 kJ/mol, 11-fold K_D ratio). When combined, the two substitutions synergistically increased binding ($\Delta\Delta G_{av}$ = -8.7 ± 0.4 kJ/mol, 35-fold K_D ratio), thereby turning p35E6 from the weakest 14-3-3 binder into the second strongest one, just below p33E6.

Fusicoccin partially destabilizes the 14-3-3/E6 PBM interaction

Fusicoccin (FSC) is commonly used as a stabilizer of 14-3-3 complexes, when its binding in the distinct pocket in the 14-3-3/phosphopeptide interface is allowed by phosphopeptide side chains of the amino acids in downstream positions relative to the phospho-residue [42, 52, 53, 54]. Binding of FSC is known to stabilize the 14-3-3 interaction with few internal motif I phosphopeptides (e.g., Gab2 peptide, 5EXA.pdb [55] and CFTR peptide, 5D3F.pdb [42]) and numerous peptides with the short motif III consensus, pS/pT**X**-COOH [31, 56, 57]. Most of the internal motif phosphopeptides, when bound to 14-3-3, apparently leave insufficient space for FSC and therefore hinder formation of the ternary complexes with FSC. Many motif III peptides with a consensus pS/pT**X**-COOH enable synergistic binding of the phosphopeptide and FSC, reinforcing the 14-3-3/phosphopeptide complex [31, 52, 53, 57, 58, 59]. At the same time, very little is known about the effect of FSC on binding of the longer motif III peptides to 14-3-3, such as in the case of E6 PBMs.

We investigated whether binding of phosphorylated E6 PBM to 14-3-3 might be affected by this small molecule and analyzed its effect on the affinity of the E6 PBMs to 14-3-3. We used FP experiments to measure equilibrium binding affinity constants of complexes

between the four HPV-E6 phosphopeptides and 14-3-3 isoforms ζ and γ, in the presence of 100 μM FSC (Supplementary Fig. 1B and Fig. 4A). The addition of FSC consistently decreased by 1.5 to 2 fold the affinities of all interactions ($\Delta\Delta G_{av}$ = -1.3 ± 0.5 and -1.8 ± 0.4 kJ/mol for ζ and γ, respectively) without altering the apparent preferences of the different peptides (Fig. 4A and B). Therefore, fusicoccin moderately destabilized all studied 14-3-3/E6 interactions, in contrast to its much broadly reported, stabilizing effect on 14-3-3 complexes [31, 42, 52]. Nevertheless, the binding of E6 peptides and FSC was not mutually exclusive.

A

	K_D (μM) ± std			
	14-3-3ζ	**14-3-3ζ +FSC**	**14-3-3γ**	**14-3-3γ +FSC**
pHPV33E6	2.72 ± 0.39	3.94 ± 0.40	0.94 ± 0.25	2.15 ± 0.26
pHPV18E6	22.4 ± 1.3	37.3 ± 5.2	11.1 ± 0.6	19.3 ± 2.0
pHPV16E6	73.6 ± 19.1	161 ± 23	37.2 ± 3.0	91.7 ± 13.2
pHPV35E6	191 ± 18	294 ± 15	125 ± 6	228 ± 61

B

$\Delta\Delta G_{av}$ = -1.28±0.45 (kJ/mol)
14-3-3ζ + FSC < 14-3-3ζ

$\Delta\Delta G_{av}$ = -1.77±0.42 (kJ/mol)
14-3-3γ + FSC < 14-3-3γ

C 14-3-3ζ dimer

FSC
FSC
18E6 phosphopeptides
...RRRE**pT**^156^QV#

D
L216 D223
D213 I217 N224
R152
E155
FSC K120 Q157 pT156
R153
N42 N173 V176
D124 R154
S45 #V158 E180
E131 R60
V46 Y128 R56
K49

E
Fusicoccin A (FSC)

F 14-3-3ζ
4 Å
B-factors:
18.1–102.4Å²
FSC
3.5 Å

Fig. 4. The destabilizing effect of FSC on the 14-3-3ζ/18E6 PBM interaction. **A**. Affinities of four selected HPV-E6 phospho-PBMs towards human 14-3-3ζ and 14-3-3γ in the absence and in the presence of FSC as determined by fluorescence polarization using FITC-labeled HSPB6 phosphopeptide as a tracer. Apparent K_D values determined from competitive FP experiments are presented. The binding curves are shown in Supplementary Fig. 1. **B**. Averaged ΔΔG values between 14-3-3-E6 phospho-PBM pairs in the absence or in the presence of FSC, calculated based on their observed order of binding affinities (from

weakest to strongest). Individual K_D values from Supplementary Fig. 1 were first converted
into ΔG values (at T=295 K; excluding cases when K_D > 300 μM) and average ΔΔG values
(ΔΔG_{av}) were calculated between the indicated motifs/isoforms. **C**. An overall view on the
ternary complex between 14-3-3ζ (subunits are shown by surface using two tints of grey),
18E6 phosphopeptide (cyan sticks) and FSC (pink sticks). FSC was soaked into the 14-3-
3ζ-18E6 chimera crystals. $2F_o$-F_c electron density maps contoured at 1σ show are shown
for the phosphopeptide and FSC only. **D**. A closeup view showing polar contacts (dashed
lines) and hydrophobic interactions (semitransparent spheres) positioning the 18E6
phosphopeptide (cyan sticks) and FSC (thin pink sticks) in the amphipathic groove of the
14-3-3ζ subunit (semitransparent light grey ribbon). $2F_o$-F_c electron density maps
contoured at 1σ are shown for the peptide and FSC. # denotes the C-terminus (-COOH).
The GGGG linker is omitted for clarity. **E**. Chemical formula of FSC showing the positions
of the functional groups discussed in the text. **F**. The effect of FSC binding. Conformational
changes upon FSC binding are shown by red arrows reflecting the closure of the C-
terminal α-helix of 14-3-3 and the rotation and translocation of the C-terminal residues in
the 18E6 peptide, accompanied by a significant rise of the local B-factors (are shown as a
gradient from blue to red as indicated). The amplitudes of the conformational changes in
14-3-3 and 18E6 peptide are indicated in Å by dashed arrows.

Structural basis for the destabilizing effect of fusicoccin

To get structural insights into the peculiar destabilizing effect of FSC on the 14-3-3/E6 PBM
interaction, we soaked the 14-3-3ζ/18E6 PBM chimera crystals with FSC and solved the
crystal structure of the 14-3-3ζ/18E6 PBM/FSC ternary complex (Fig. 4C-F and Table 1).
FSC soaking did not disrupt the crystal lattice of the 14-3-3ζ/18E6 PBM chimera crystals
(Supplementary Fig. 3) and the overall assembly (Fig. 4C and D); however, significant local
structural rearrangements could be observed. FSC (Fig. 4E) binding results in a ~4 Å
closure of the last α-helix of 14-3-3ζ (Fig. 4F), similar to what has already been described
for other 14-3-3 complexes containing FSC [58]. Crystal lattice obtained by FSC soaking
preserved the characteristic phosphopeptide swap stabilizing the contacts with the
neighboring chimera molecules, and FSC was bound in each 14-3-3ζ subunit, side-to-side
with the C-terminal end of the 18E6 PBM (Fig. 4C and D, Supplementary Fig. 3). All
residues and atoms within the analyzed area could be observed in the electron density
thanks to the high resolution (Table 1).

FSC occupies its well-defined cavity where it is positioned by hydrophobic interactions with
Phe117, Ile166, Ile217 and Leu216, polar contacts with residues Asn42 and Asp213, and a
remarkable H-bond involving its 3-metoxy oxygen and the side chain of Lys120 of 14-3-3ζ
(Fig. 4D). The latter contact breaks the Lys120 interaction with the carboxyl-group of the
18E6 PBM formed in the absence of FSC, displacing the carboxyl to another position,
where it establishes a new contact with the 12-hydroxy group of FSC (Fig. 4D and E). 14-3-
3ζ Lys49 also switches its position, and loses a contact to the backbone carbonyl of
pThr156 (Fig. 3C) to establish instead a contact with the 12-hydroxy group of FSC. The
side chain of the C-terminal Val158 of the 18E6 PBM shifts 3.5 Å towards the phosphate
moiety of Thr156, breaking the hydrophobic contact with Val46 of 14-3-3ζ and significantly
dispersing the local electron density (Fig. 4F). As a result, while most of the peptide
conformation remained unchanged, the B-factors of the last 18E6 PBM residue in the
refined FSC-bound structure increased significantly (Fig. 4F). Therefore, FSC binding

causes an apparent strain in the conformation of the C-terminus of 18E6 PBM (Figs 4 and
5). In addition, the presence of FSC significantly reshuffles the water network that
surrounds the E6 carboxy-terminal extremity, which directly contacts the bound FSC (Fig.
5). The simultaneous binding of the 18E6 PBM and FSC in the amphipathic groove of 14-3-
3 and the mild destabilizing influence of FSC indicate that this ternary complex can be used
as a starting point to design both stabilizers and inhibitors of 14-3-3/E6 interactions.

Fig. 5. Comparison of the water-mediated polar contacts formed in the 14-3-3ζ/18E6
interface in the absence (**A**) or in the presence of fusicoccin (FSC) (**B**). The main residues
involved in the interactions are shown by sticks with color coding: 14-3-3 residues are
shown in light grey, 18E6 residues are in cyan, phospho-group of Thr156 is shown by
orange sticks. FSC is shown by thin magenta sticks, water molecules affected by FSC
binding are shown by lime green, those similar in two structures are black. The C-terminal
18E6 residue (V158) is denoted by #. Note the significant redistribution of water molecules
upon FSC binding.

DISCUSSION

E6 oncoproteins of all high-risk alpha HPV types contain a conserved C-terminal PDZ-
Binding Motif which can turn into a potential 14-3-3-binding motif upon phosphorylation of

the conserved Thr/Ser residue at position -2 (Fig. 1). We demonstrated here that the
phosphorylated forms of four selected alpha HPV-E6 PBMs detectably bound to all seven
human 14-3-3 isoforms. Unexpectedly, the binding affinities of these 28 distinct 14-3-3/E6
complexes showed large variations, spanning a 10-fold K_D range for different 14-3-3
isoforms binding to a given PBM, and a 100-fold K_D range for different E6 phospho-PBMs
binding to a given 14-3-3 isoform. Furthermore, affinity variations followed trends that were
all but random. The 14-3-3 binding profiles of all E6 phosphopeptides turned out to be
remarkably parallel, and so were the E6-binding profiles of all 14-3-3 isoforms (Fig. 2B and
C).

When considering the present data together with previously published literature, 14-3-3
interactions with phosphorylated motifs from different origins have a strikingly wide affinity
range, spanning from low nanomolar to low millimolar detectable dissociation constants
(Fig. 6). Despite variable interaction modes (e.g. monovalent vs. divalent), the 14-3-3
family-wide affinity trends are surprisingly parallel, with 14-3-3γ and 14-3-3η consistently
being the strongest binders and 14-3-3σ and 14-3-3ε being the weakest binders.
Independently of the nature of the target motif, the average maximal K_D ratio between the
strongest-binding and the weakest-binding 14-3-3 is around 10-fold in our present work
and also in the literature (Fig. 6), Conversely, phosphopeptides, even very similar in
sequence, can sample much wider ranges of binding affinities for a given 14-3-3 isoform.
For instance, for 14-3-3γ, the K_D ratio between the strongest and the weakest binding
phosphopeptide is almost 625-fold in the present work, and 39,000-fold when taking into
account other reports [42, 43, 60, 61] (Fig. 6). However, the discussed weak interactions do not
necessarily lead to the absence of complex formation *in cellulo*, as 14-3-3 proteins are
among the most abundant proteins in human cells [62, 63]. Indeed, according to Protein
Abundance Database [63], 14-3-3ε is the 48[th] most abundant human protein (2479 ppm) and
14-3-3ζ is the 72[nd] (1680 ppm), whereas all seven 14-3-3 isoforms are within the 820 most
abundant proteins (i.e., the top 4.1%) out of 19949 proteins in the integrated whole human
body dataset. Given such high abundance, strong 14-3-3 interactions might lead to
extremely tight complex formation, while weak interactions might lead to transient, more
dynamic 14-3-3 complexes.

The observed strong and parallel affinity variations are noteworthy if one considers the high
degree of conservation of the amphipathic grooves of the seven 14-3-3 isoforms and of the
amphipathic groove-binding residues of the four E6 PBMs. Indeed, affinity differences may
stem not only from structural particularities of the bound complexes but also from intrinsic
properties of the unbound partners. Conformations unfavorable for complex formation may
exist in various proportions for each 14-3-3 isoform and each E6 phosphopeptide, thereby
influencing at various degrees the formation of the complexes and their resulting binding
affinities. On the one hand, full-length 14-3-3 proteins feature the flexible C-terminal tails
that are the most variable elements among isoforms; at least for some isoforms they have
been shown to sample conformations also occupying the amphipathic 14-3-3 grooves and
thereby regulating their affinities for target peptides [64, 65]. Conformational dynamics of the
14-3-3 dimers along the open-closed state trajectory [26, 44] may also differ among 14-3-3
isoforms, which can probably affect the effectiveness of adopting the peptide-bound
conformation. On the other hand, it has been shown that a phosphoryl-group in a
disordered segment can form local "charge clamps" *in-cis* with a neighboring Arg/Lys
residue, thereby decreasing its apparent availability for binding partners [51]. The four E6
PBMs studied herein display distinctive charged, polar or non-polar residues at several

positions (-1; -6; -7; -8; -9) that may influence their explorable conformational spaces. By replacing the Glu -1 by Ala in the weakest 14-3-3 binder, p35E6, we obtained a significantly stronger 14-3-3 binder, suggesting that we successfully disrupted a charge clamp that is unfavorable to binding of the p35E6 motif. Nonetheless, fine differences of contact networks at the interfaces of E6/14-3-3 complexes may also influence affinity. In this line, comparison of the present p18E6/14-3-3ζ structure to the previously solved p16E6/14-3-3σ structure [24] suggested that an Arg residue at position -6 established favorable interface contacts (Fig. 3C and D). Accordingly, replacing Thr -6 of p35E6 with Arg -6 as found in p18E6 and p33E6, effectively reinforced the binding event. Finally, combining both Glu→Ala and Thr→Arg substitutions at positions -1 and -6, respectively, resulted in a p35E6 variant that bound 14-3-3 proteins almost as strongly as the strongest studied E6 binder, p33E6 (Fig. 2A and Fig. 3E). Interestingly, the RSK1 phospho-PBMs (RRVRKLPSTpT**L**-COOH and RRVRKLPSpT**T**L-COOH), featuring non-charged residues Thr and Leu in the downstream position relative to the phosphothreonine, and therefore being unable to form the equivalent charge-clamp that is suggested in the case of p35E6 (SKPTRREpT**E**V-COOH), show remarkably high affinities to all 14-3-3 isoforms *per se* (Fig. 2A and Supplementary Fig. 1). Our data prove that charge clamps can indeed strongly contribute to the binding preferences of different phosphopeptide motifs. This principle needs to be further examined for all kinds of phosphorylation-regulated interactions as charge-clamp formation might be an often ignored biochemical property of practically all phosphorylation sites.

Fig. 6. Compilation of 14-3-3 family-wide affinity measurements of exemplary 14-3-3-binding motifs. A. Affinity maps of 14-3-3 interactions based on experimentally determined dissociation constants against the 14-3-3ome, as obtained in the current work and in refs [42, 43, 60, 61]. B. When normalized to the strongest partner-binder 14-3-3γ, all studied peptides are following very similar affinity trends between the different 14-3-3 isoforms. Between the strongest and the weakest partner-binding 14-3-3 isoform, a 12-fold dissociation constant decrease was found on average. C. When normalizing to the generally strongest studied 14-3-3-binding motif, the isoform-specific affinity variations for each peptide are compensated to clearly show the average differences between different binding motifs. Between the strongest and the weakest analyzed 14-3-3-binding motif, a 34,000-fold K_D decrease was found on average for all 14-3-3 isoforms. Only counting the very similar HPV-E6 peptides sequence-wise, this affinity difference can still be larger than 100-fold. Therefore, the motif-to-motif affinity differences are expected to be much higher than the 14-3-3 isoform-to-isoform affinity differences. Binding motifs that are analyzed in other studies are highlighted with a grey background [42, 43, 60, 61]. The color scale is either based on affinity values or in K_D ratios. √ denotes affinities weaker than the limit of quantitation of the fluorescence polarization assay.

When phosphorylated, all the E6 PBMs studied here ideally match motif III consensus recognized by 14-3-3 with two C-terminal residues downstream of the phosphorylated threonine (pS/pT**XX**-COOH) (Fig. 1B). Structures of motif III complexes with 14-3-3 have been less documented than the internal motif I/II complexes. Moreover, the pS/pT**XX**-COOH subtype of motif III sites, studied here, is underrepresented in the Protein Data Bank, where mostly motif III sites with only one residue after a phospho-residue have been co-crystallized with 14-3-3 (Supplementary Table 1). To the best of our knowledge, the only two reported structures with pS/pT**XX**-COOH consensus peptides are those of 14-3-3γ with the influenza virus protein NS1 (4O46.pdb) and our recently reported 14-3-3σ complex with the synthetic papillomavirus 16E6 phosphopeptide (6TWZ.pdb [24]), however, both of them have rather low resolution (2.8-2.9 Å) (Supplementary Table 1). Together with the henipavirus protein W binding to 14-3-3σ via its C-terminal motif III peptide RRMpS**N**-COOH [66], the adeno-associated virus protein Rep68 binding to 14-3-3 proteins using its C-terminal motif III peptide RGHpS**L**-COOH [67] and the hepatitis B virus protein X binding to 14-3-3ζ via an internal motif I peptide RPLpSGP [68], these examples illustrate that viral proteins commonly use their elements, efficiently mimicking the host 14-3-3-binding peptides, to hijack the cellular functions controlled by 14-3-3 proteins. One of the possible mechanisms is the 14-3-3-mediated stabilization of viral proteins to evade dephosphorylation and degradation, which may prolong the half-life and increase chances for successful replication and further infections [21, 68].

The phospho-PBM conformation revealed by our crystal structures, ideally congruous to the amphipathic groove of 14-3-3, leaves vacant the cavity that is known to be druggable and occupied by a fungal toxin called fusicoccin (FSC) [33, 54, 56]. Benefitting from the well-diffracting crystals of the 14-3-3ζ-18E6 PBM chimera, we proceeded with their soaking with FSC, which resulted in a 1.85-Å crystal structure of the ternary 14-3-3ζ/18E6 PBM/FSC complex. The direct comparison of the 14-3-3ζ/18E6 complex in the absence and in the presence of FSC revealed a significant conformational change in the last α-helix of 14-3-3ζ, a rearrangement of bound water molecules and a remarkable strain in the C-terminal part of the 18E6 PBM peptide (Figs 4 and 5). This strain forced the translocation of the carboxyl-group and increased the displacement factors for the last C-terminal residue, suggesting destabilization of the 14-3-3ζ/18E6 PBM complex by FSC. This partial destabilization, and thus the less-documented inhibitory action of FSC, was confirmed by *in vitro* binding assays using fluorescence polarization (Fig. 4A and B). Noteworthily, the structural data on the effect of FSC on the C-terminal end of the E6 PBM is very consistent with the preservation of the affinity differences for various 14-3-3/phospho-PBMs complexes in the presence of FSC (Fig. 4A). This supports the notion that those affinity differences are dictated by positions beyond the last C-terminal residue, including those involved in formation of charge clamps.

We could find only one earlier reported example when FSC decreased the binding affinity of a 14-3-3 to a motif III peptide (the interleukin 9 receptor alpha chain (IL-R9α) peptide, RSWpT**F**-COOH [31]), but in that case no experimental structural information was available. In addition, it was reported that binding of the shortest motif III peptide from the cyclin-dependent kinase inhibitor (p27[Kip1]) RRQpT-COOH to 14-3-3 is not affected by FSC, most likely due to the absence of direct contacts between FSC and the phosphopeptide, which both bind in the amphipathic groove of 14-3-3 independently [31]. The structural basis for the inhibitory action of FSC has very recently been reported for several internal motif I 14-3-3-

binding peptides [58]. Therefore, complementing the range of reports on the stabilizing effect of FSC [31, 42, 54, 55, 56], our results provide the first structural evidence that FSC can be a negative regulator of 14-3-3 interactions with typical motif III peptides.

Thus, our structural and *in vitro* binding data with FSC confirm the druggability of the 14-3-3-E6 interaction and suggest that appropriate modification and optimization of the small molecule may provide promising opportunities for selective modulation of viral protein-14-3-3 interactions in the future.

METHODS

Cloning, protein expression and purification and peptide synthesis

Previously described chimeras contained the C-terminally truncated human 14-3-3σ (Uniprot ID P31947; residues 1-231, 14-3-3σΔC) bearing on its N-terminus a His$_6$-tag cleavable by 3C protease and phosphorylatable peptides tethered to the 14-3-3σ C-terminus by a GSGS linker [44]. The novel chimera was designed taking into account the following modifications. First, it contained the C-terminally truncated human 14-3-3ζ sequence (Uniprot ID P63104; residues 1-229, 14-3-3ζΔC) connected to the PKA-phosphorylatable 18E6 heptapeptide around Thr156. Second, the 14-3-3ζ core was modified to block Ser58 phosphorylation (S58A) [69, 70]. Third, to improve crystallizability, the 14-3-3ζ sequence was mutated by introducing the ^{73}EKK^{75}→AAA and ^{157}KKE^{159}→AAA amino acid replacements in the highest-scoring clusters 1 and 2 predicted by the surface entropy reduction approach [47, 71]. Finally, the linker was changed to GGGG to exclude its unspecific phosphorylation (Supplementary Fig. 2A).

cDNA of the 14-3-3ζ-18E6 chimera was codon-optimized for expression in *Escherichia coli* and synthesized by IDT Technologies (Coralville, Iowa, USA). The 14-3-3ζΔC gene was flanked by *NdeI* and *AgeI* restriction endonuclease sites to enable alteration of the 14-3-3 or E6 PBM peptide sequences. The entire 14-3-3ζ-GGGG-18E6 PBM construct was inserted into a pET28-his-3C vector [72] using *NdeI* and *XhoI* restriction endonuclease sites. The resulting vector was amplified in DH5α cells and verified using DNA sequencing in Evrogen (Moscow, Russia, www.evrogen.ru).

The assembled vector (Kanamycin resistance) was transformed into chemically competent *E. coli* BL21(DE3) cells for expression either in the absence or in the presence of the His$_6$-tagged catalytically active subunit of mouse PKA [72]. Protein expression was induced by the addition of isopropyl-β-thiogalactoside (IPTG) to a final concentration of 0.5 mM and continued for 16 h at 25 °C. The overexpressed protein was purified using subtractive immobilized metal-affinity chromatography (IMAC) and gel-filtration essentially as described earlier for 14-3-3σ chimeras [44] (Supplementary Fig. 2B and C). The purified phosphorylated 14-3-3ζ-18E6 chimera revealed the characteristic downward shift on native PAGE compared to the unphosphorylated counterpart (Supplementary Fig. 2D). Given the absence of PKA phosphorylation sites in the modified 14-3-3ζ core and the linker, this strongly indicated 18E6 phosphorylation by co-expressed PKA. The chimera was fully soluble and stable at concentrations above 20 mg/ml required for crystallization. Protein concentration was determined at 280 nm on a Nanophotometer NP80 (Implen, Germany) using extinction coefficient equal to 0.93 (mg/ml)$^{-1}$ cm^{-1}.

For affinity measurements, full-length human 14-3-3 constructs with a rigid N-terminal MBP fusion were used. The coding sequences of the full-length 14-3-3 epsilon, gamma and zeta

were received from Prof. Lawrence Banks. cDNAs encoding other full-length 14-3-3
isoforms β, τ, η and σ were obtained as codon-optimized for *E.coli* expression synthetic
genes from IDT Technologies (Coralville, Iowa, USA). All 14-3-3 isoforms were fused via a
three-alanine linker to the C-terminus of a mutant MBP carrying the following amino acid
substitutions: D83A, K84A, K240A, E360A, K363A and D364A, as previously described [49].
All resulting clones were verified by sequencing. The MBP-fused proteins were expressed
in *E.coli* BL21 with IPTG induction. Proteins were affinity purified on an amylose column
and were further purified by ion-exchange chromatography (HiTrap Q HP, GE Healthcare).
Protein concentrations were determined by UV spectroscopy. The double-purified samples
were supplemented with glycerol and TCEP before aliquoting and freezing in liquid
nitrogen.

HPV peptides (35E6: biotin-ttds-SKPTRRETEV; 16E6: biotin-ttds-SSRTRRETQL; 18E6:
biotin-ttds-RLQRRRETQV; 33E6: biotin-ttds-SRSRRRETAL; p35E6: biotin-ttds-
SKPTRREpTEV; p35E6 E-1A: biotin-ttds-SKPTRREpTAV; p35E6 T-6R: biotin-ttds-
SKPRRREpTEV; p35E6 E-1A T-6R: biotin-ttds-SKPRRREpTAV; p16E6: biotin-ttds-
SSRTRREpTQL; p18E6: biotin-ttds-RLQRRREpTQV; p33E6: biotin-ttds-SRSRRREpTAL)
and RSK1 peptides (RSK1_-1P: biotin-ttds-RRVRKLPSTpTL and RSK1_-2P: biotin-ttds-
RRVRKLPSpTTL) were chemically synthesized in-house on an ABI 443A synthesizer with
Fmoc strategy. The fluorescently labeled HSPB6 (WLRRApSAPLPGLK) peptide (fpB6)
was prepared by FITC labeling of the chemically synthesized peptide as described
previously [24].

Fluorescence polarization (FP) assay

Fluorescence polarization was measured with a PHERAstar (BMG Labtech, Offenburg,
Germany) microplate reader by using 485 ± 20 nm and 528 ± 20 nm band-pass filters (for
excitation and emission, respectively). In direct FP measurements, a dilution series of the
14-3-3 protein was prepared in 96-well plates (96 well skirted pcr plate, 4ti-0740, 4titude,
Wotton, UK) in a 20 mM HEPES pH 7.5 buffer containing 150 mM NaCl, 0.5 mM TCEP,
0.01% Tween 20, 50 nM fluorescently-labeled fpB6 peptide and 100 µM fusicoccin (FSC),
if indicated. The volume of the dilution series was 40 µl, which was later divided into three
technical replicates of 10 µl upon transferring to 384-well micro-plates (low binding
microplate, 384 well, E18063G5, Greiner Bio-One, Kremsmünster, Austria). In total,
polarization of the probe was measured at 8 different protein concentrations (whereas one
contained no protein and corresponded to the free peptide). In competitive FP
measurements, the same buffer was supplemented with the protein to achieve a complex
formation of 60-80%, based on the titration. Then, this mixture was used for creating a
dilution series of the unlabeled competitor (i.e. the studied peptides) and the measurement
was carried out identically as in the direct experiment. Analysis of FP experiments were
carried out using ProFit, an in-house developed, Python-based fitting program [73]. The
dissociation constant of the direct and competitive FP experiment was obtained by fitting
the measured data with quadratic and competitive equation, respectively [73, 74]. ΔG values
were calculated using the formula $\Delta G = -RT \cdot \ln(K_D)$, at 295 K. $\Delta\Delta G_{av}$ values were obtained
by calculating the average and the standard deviation of all obtained individual ΔΔG values
(between different motifs or different proteins), excluding cases when $K_D > 300$ µM.

Crystallization and structure determination

Crystallization conditions were screened using commercially available and in-house developed kits (Qiagen, Hampton Research, Emerald Biosystems) by the sitting-drop vapor-diffusion method in 96-well MRC 2-drop plates (SWISSCI, Neuheim, Switzerland), using a Mosquito robot (TTP Labtech, Cambridge, UK) at 4 °C. The optimized condition of the crystals consisted of 19% polyethylene glycol 4000, 0.1M cacodylate buffered at pH 5.5. For soaking, crystals were transferred to a mother-liquor solution containing (saturated, partially precipitated) 5 mM fusicoccin and crystals were harvested after an 18h incubation period. All crystals were flash-cooled in a cryoprotectant solution containing 20% glycerol and stored in liquid nitrogen.

X-ray diffraction data were collected at the Synchrotron Swiss Light Source (SLS) (Switzerland) on the X06DA (PXIII) beamline and processed with the program XDS [75]. The crystal structure was solved by molecular replacement with a high-resolution crystal structure of 14-3-3ζ (PDB ID 2O02) using Phaser [76] and structure refinement was carried out with PHENIX [77]. TLS refinement was applied during the refinement. The crystallographic parameters and the statistics of data collection and refinement are shown in Table 1.

Acknowledgments

We would like to thank Prof. Lawrence Banks for the shared plasmids, Prof. Alexey Babakov for the provided fusicoccin preparations and Dr. Yaroslav Faletrov for fusicoccin identity verification. N.N.S. is grateful to the Russian Science Foundation for the grant no. 19-74-10031. We thank the support of the Swiss Light Source synchrotron (P. Scherrer Institute, Villigen, Switzerland) and the help of the beam-scientist at the PXIII beamline. The work was supported by the Ligue contre le cancer (équipe labellisée 2015 to G.T.), the French Infrastructure for Integrated Structural Biology (FRISBI), and Instruct-ERIC. G.G. was supported by the Post-doctorants en France program of the Fondation ARC.

Author contributions

G.G. purified proteins, carried out FP experiments, performed crystallographic studies, analyzed the data and edited the paper. K.V.T. cloned, purified and characterized proteins. C.K. cloned and purified proteins. P.E. synthesized the peptides. G.T. performed data analysis and data interpretation and edited the paper. N.N.S. contributed to protein purification and crystallographic experiments, supervised the research, analyzed the data and wrote the paper.

Conflict of interests

The authors declare no conflict of interest.

Data availability

The refined model and the structure factor amplitudes have been deposited in the PDB with the accession codes 6ZFD and 6ZFG. All other data supporting the findings of this study are available from the corresponding authors upon reasonable request.

References

1.   Suarez I, Trave G. Structural Insights in Multifunctional Papillomavirus Oncoproteins. *Viruses* **10**, (2018).

774
775   2.   Munoz N*, et al.* Epidemiologic classification of human papillomavirus types associated with
776   cervical cancer. *N Engl J Med* **348**, 518-527 (2003).
777
778   3.   Bosch FX*, et al.* Epidemiology and natural history of human papillomavirus infections and
779   type-specific implications in cervical neoplasia. *Vaccine* **26 Suppl 10**, K1-16 (2008).
780
781   4.   Bouvard V*, et al.* A review of human carcinogens--Part B: biological agents. *The Lancet
782   Oncology* **10**, 321-322 (2009).
783
784   5.   Basukala O, Sarabia-Vega V, Banks L. Human papillomavirus oncoproteins and post-
785   translational modifications: generating multifunctional hubs for overriding cellular
786   homeostasis. *Biological chemistry* **401**, 585-599 (2020).
787
788   6.   McBride AA. Oncogenic human papillomaviruses. *Philos Trans R Soc Lond B Biol Sci* **372**,
789   (2017).
790
791   7.   Poirson J*, et al.* Mapping the interactome of HPV E6 and E7 oncoproteins with the ubiquitin-
792   proteasome system. *FEBS J* **284**, 3171-3201 (2017).
793
794   8.   Celegato M*, et al.* A novel small-molecule inhibitor of the human papillomavirus E6-p53
795   interaction that reactivates p53 function and blocks cancer cells growth. *Cancer Lett* **470**,
796   115-125 (2020).
797
798   9.   Kolluru S, Momoh R, Lin L, Mallareddy JR, Krstenansky JL. Identification of potential
799   binding pocket on viral oncoprotein HPV16 E6: a promising anti-cancer target for small
800   molecule drug discovery. *BMC molecular and cell biology* **20**, 30 (2019).
801
802   10.   Zanier K*, et al.* The E6AP binding pocket of the HPV16 E6 oncoprotein provides a docking
803   site for a small inhibitory peptide unrelated to E6AP, indicating druggability of E6. *PLoS
804   One* **9**, e112514 (2014).
805
806   11.   Ramirez J*, et al.* Targeting the Two Oncogenic Functional Sites of the HPV E6 Oncoprotein
807   with a High-Affinity Bivalent Ligand. *Angew Chem Int Ed Engl* **54**, 7958-7962 (2015).
808
809   12.   Vande Pol SB, Klingelhutz AJ. Papillomavirus E6 oncoproteins. *Virology* **445**, 115-137
810   (2013).
811
812   13.   White EA, Kramer RE, Tan MJ, Hayes SD, Harper JW, Howley PM. Comprehensive
813   analysis of host cellular interactions with human papillomavirus E6 proteins identifies new
814   E6 binding partners and reflects viral diversity. *J Virol* **86**, 13174-13186 (2012).
815
816   14.   Martinez-Zapien D*, et al.* Structure of the E6/E6AP/p53 complex required for HPV-
817   mediated degradation of p53. *Nature* **529**, 541-545 (2016).
818
819   15.   Masuda Y*, et al.* Stepwise multipolyubiquitination of p53 by the E6AP-E6 ubiquitin ligase
820   complex. *J Biol Chem* **294**, 14860-14875 (2019).
821

822   16.   Vats A, Thatte J, Banks L. Identification of E6AP-independent degradation targets of HPV
823   E6. *The Journal of general virology* **100**, 1674-1679 (2019).

825   17.   Songyang Z*, et al.* Recognition of unique carboxyl-terminal motifs by distinct PDZ domains.
826   *Science* **275**, 73-77 (1997).

828   18.   Boon SS, Tomaic V, Thomas M, Roberts S, Banks L. Cancer-causing human papillomavirus
829   E6 proteins display major differences in the phospho-regulation of their PDZ interactions. *J
830   Virol* **89**, 1579-1586 (2015).

832   19.   Ganti K*, et al.* The Human Papillomavirus E6 PDZ Binding Motif: From Life Cycle to
833   Malignancy. *Viruses* **7**, 3530-3551 (2015).

835   20.   Delury CP*, et al.* The role of protein kinase A regulation of the E6 PDZ-binding domain
836   during the differentiation-dependent life cycle of human papillomavirus type 18. *J Virol* **87**,
837   9463-9472 (2013).

839   21.   Boon SS, Banks L. High-risk human papillomavirus E6 oncoproteins interact with 14-3-
840   3zeta in a PDZ binding motif-dependent manner. *J Virol* **87**, 1586-1595 (2013).

842   22.   Thatte J, Massimi P, Thomas M, Boon SS, Banks L. The Human Papillomavirus E6 PDZ
843   Binding Motif Links DNA Damage Response Signaling to E6 Inhibition of p53
844   Transcriptional Activity. *J Virol* **92**, (2018).

846   23.   Kuhne C, Gardiol D, Guarnaccia C, Amenitsch H, Banks L. Differential regulation of human
847   papillomavirus E6 by protein kinase A: conditional degradation of human discs large protein
848   by oncogenic E6. *Oncogene* **19**, 5884-5891 (2000).

850   24.   Gogl G*, et al.* Dual Specificity PDZ- and 14-3-3-Binding Motifs: A Structural and
851   Interactomics Study. *Structure* **28**, 747-759 e743 (2020).

853   25.   Aitken A. 14-3-3 proteins: a historic overview. *Semin Canc Biol* **16**, 162-172 (2006).

855   26.   Yang X*, et al.* Structural basis for protein-protein interactions in the 14-3-3 protein family.
856   *Proc Natl Acad Sci U S A* **103**, 17237-17242 (2006).

858   27.   Muslin AJ, Tanner JW, Allen PM, Shaw AS. Interaction of 14-3-3 with signaling proteins is
859   mediated by the recognition of phosphoserine. *Cell* **84**, 889-897 (1996).

861   28.   Yaffe MB*, et al.* The structural basis for 14-3-3:phosphopeptide binding specificity. *Cell* **91**,
862   961-971 (1997).

864   29.   Mackintosh C. Dynamic interactions between 14-3-3 proteins and phosphoproteins regulate
865   diverse cellular processes. *Biochem J* **381**, 329-342 (2004).

867   30.   Ganguly S, Weller J, Ho A, Chemineau P, Malpaux B, Klein D. Melatonin synthesis: 14-3-
868   3-dependent activation and inhibition of arylalkylamine N-acetyltransferase mediated by
869   phosphoserine-205. *Proc Natl Acad Sci U S A* **102**, 1222-1227 (2005).

31.   Paiardini A*, et al.* The phytotoxin fusicoccin differently regulates 14-3-3 proteins association
to mode III targets. *IUBMB Life* **66**, 52-62 (2014).

32.   Obsil T, Obsilova V. Structural basis of 14-3-3 protein functions. *Semin Cell Dev Biol* **22**,
663-672 (2011).

33.   Stevers LM*, et al.* Modulators of 14-3-3 Protein-Protein Interactions. *J Med Chem* **61**, 3755-
3778 (2018).

34.   Luck K, Charbonnier S, Trave G. The emerging contribution of sequence context to the
specificity of protein interactions mediated by PDZ domains. *FEBS Lett* **586**, 2648-2661
(2012).

35.   Puntervoll P*, et al.* ELM server: A new resource for investigating short functional sites in
modular eukaryotic proteins. *Nucleic Acids Res* **31**, 3625-3630 (2003).

36.   Miller CJ, Turk BE. Homing in: Mechanisms of Substrate Targeting by Protein Kinases.
Trends Biochem Sci **43**, 380-394 (2018).

37.   Ben-Shimon A, Niv MY. Deciphering the Arginine-binding preferences at the substrate-
binding groove of Ser/Thr kinases by computational surface mapping. *PLoS Comput Biol* **7**,
e1002288 (2011).

38.   Sarabia-Vega V, Banks L. Acquisition of a phospho-acceptor site enhances HPV E6 PDZ-
binding motif functional promiscuity. *The Journal of general virology*, (2019).

39.   Abraham RT. Cell cycle checkpoint signaling through the ATM and ATR kinases. *Genes*
Dev **15**, 2177-2196 (2001).

40.   Espejo AB*, et al.* PRMT5 C-terminal Phosphorylation Modulates a 14-3-3/PDZ Interaction
Switch. *J Biol Chem* **292**, 2255-2265 (2017).

41.   Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: a sequence logo generator.
Genome research **14**, 1188-1190 (2004).

42.   Stevers LM*, et al.* Characterization and small-molecule stabilization of the multisite tandem
binding between 14-3-3 and the R domain of CFTR. *Proc Natl Acad Sci U S A* **113**, E1152-
1161 (2016).

43.   Manschwetus JT*, et al.* Binding of the Human 14-3-3 Isoforms to Distinct Sites in the
Leucine-Rich Repeat Kinase 2. *Front Neurosci* **14**, 302 (2020).

44.   Sluchanko NN, Tugaeva KV, Greive SJ, Antson AA. Chimeric 14-3-3 proteins for
unraveling interactions with intrinsically disordered partners. *Sci Rep* **7**, 12014 (2017).

45.   Tugaeva KV, Remeeva A, Gushchin I, Cooley RB, Sluchanko NN. Design, expression,
purification and crystallization of human 14-3-3zeta protein chimera with phosphopeptide
from proapoptotic protein BAD. *Protein Expr Purif*, 105707 (2020).

46.   Woodcock JM, Murphy J, Stomski FC, Berndt MC, Lopez AF. The dimeric versus
921   monomeric status of 14-3-3zeta is controlled by phosphorylation of Ser58 at the dimer
922   interface. *J Biol Chem* **278**, 36323-36327 (2003).

924   47.   Goldschmidt L, Cooper DR, Derewenda ZS, Eisenberg D. Toward rational protein
925   crystallization: A Web server for the design of crystallizable protein variants. *Protein Sci* **16**,
926   1569-1576 (2007).

928   48.   Nomine Y*, et al.* Structural and functional analysis of E6 oncoprotein: insights in the
929   molecular pathways of human papillomavirus-mediated pathogenesis. *Mol Cell* **21**, 665-678
930   (2006).

932   49.   Zanier K*, et al.* Structural basis for hijacking of cellular LxxLL motifs by papillomavirus E6
933   oncoproteins. *Science* **339**, 694-698 (2013).

935   50.   Tugaeva KV, Titterington J, Sotnikov DV, Maksimov EG, Antson AA, Sluchanko NN.
936   Molecular basis for the recognition of steroidogenic acute regulatory protein by the 14-3-3
937   protein family. *FEBS J*, 10.1111/febs.15474 (2020).

939   51.   Gogl G*, et al.* Dynamic control of RSK complexes by phosphoswitch-based regulation.
940   *FEBS J* **285**, 46-71 (2018).

942   52.   Camoni L, Visconti S, Aducci P. The phytotoxin fusicoccin, a selective stabilizer of 14-3-3
943   interactions? *IUBMB Life* **65**, 513-517 (2013).

945   53.   De Vries-van Leeuwen IJ*, et al.* Interaction of 14-3-3 proteins with the estrogen receptor
946   alpha F domain provides a drug target interface. *Proc Natl Acad Sci U S A* **110**, 8894-8899
947   (2013).

949   54.   Wurtele M, Jelich-Ottmann C, Wittinghofer A, Oecking C. Structural view of a fungal toxin
950   acting on a 14-3-3 regulatory complex. *EMBO J* **22**, 987-994 (2003).

952   55.   Bier D*, et al.* Small-Molecule Stabilization of the 14-3-3/Gab2 Protein-Protein Interaction
953   (PPI) Interface. *ChemMedChem* **11**, 911-918 (2016).

955   56.   Sengupta A, Liriano J, Miller BG, Frederich JH. Analysis of Interactions Stabilized by
956   Fusicoccin A Reveals an Expanded Suite of Potential 14-3-3 Binding Partners. *ACS Chem
957   Biol* **15**, 305-310 (2020).

959   57.   Ohkanda J*, et al.* Structural Effects of Fusicoccin upon Upregulation of 14-3-3-
960   Phospholigand Interaction and Cytotoxic Activity. *Chemistry* **24**, 16066-16071 (2018).

962   58.   Kaplan A*, et al.* Polypharmacological Perturbation of the 14-3-3 Adaptor Protein
963   Interactome Stimulates Neurite Outgrowth. *Cell chemical biology* **27**, 657-667 e656 (2020).

965   59.   Saponaro A*, et al.* Fusicoccin Activates KAT1 Channels by Stabilizing Their Interaction
966   with 14-3-3 Proteins. *Plant Cell* **29**, 2570-2580 (2017).
967

968   60.   Kilisch M, Lytovchenko O, Arakel EC, Bertinetti D, Schwappach B. A dual phosphorylation
969   switch controls 14-3-3-dependent cell surface expression of TASK-1. *J Cell Sci* **129**, 831-
970   842 (2016).
971

61.   Centorrino F, Ballone A, Wolter M, Ottmann C. Biophysical and structural insight into the
USP8/14-3-3 interaction. *FEBS Lett* **592**, 1211-1220 (2018).

62.   Boston PF, Jackson P, Thompson RJ. Human 14-3-3 protein: radioimmunoassay, tissue
distribution, and cerebrospinal fluid levels in patients with neurological disorders. *J
Neurochem* **38**, 1475-1482 (1982).

63.   Wang M, Herrmann CJ, Simonovic M, Szklarczyk D, von Mering C. Version 4.0 of PaxDb:
Protein abundance data, integrated across model organisms, tissues, and cell-lines.
Proteomics **15**, 3163-3168 (2015).

64.   Silhan J, *et al.* 14-3-3 protein C-terminal stretch occupies ligand binding groove and is
displaced by phosphopeptide binding. *J Biol Chem* **279**, 49113-49119 (2004).

65.   Truong A, Masters S, Yang H, Fu H. Role of the 14-3-3 C-terminal loop in ligand
interaction. *Proteins* **49**, 321-325 (2002).

66.   Edwards MR, *et al.* Henipavirus W Proteins Interact with 14-3-3 To Modulate Host Gene
Expression. *J Virol* **94**, (2020).

67.   Han SI, *et al.* Rep68 protein of adeno-associated virus type 2 interacts with 14-3-3 proteins
depending on phosphorylation at serine 535. *Virology* **320**, 144-155 (2004).

68.   Tang Y, *et al.* 14-3-3zeta binds to hepatitis B virus protein X and maintains its protein
stability in hepatocellular carcinoma cells. *Cancer medicine* **7**, 5543-5553 (2018).

69.   Gu Y-M, Jin Y-H, Choi J-K, Baek K-H, Yeo C-Y, Lee K-Y. Protein kinase A
phosphorylates and regulates dimerization of 14-3-3 epsilon. *FEBS Lett* **580**, 305-310
(2006).

1001

1002   70.   Sluchanko NN, Uversky VN. Hidden disorder propensity of the N-terminal segment of
1003   universal adapter protein 14-3-3 is manifested in its monomeric form: Novel insights into
1004   protein dimerization and multifunctionality. *Biochim Biophys Acta* **1854**, 492-504 (2015).

1005

1006   71.   Goldschmidt L, Cooper DR, Derewenda ZS, Eisenberg D. SERp Server. (ed^(eds) (2007).

1007

1008   72.   Tugaeva KV, Tsvetkov PO, Sluchanko NN. Bacterial co-expression of human Tau protein
1009   with protein kinase A and 14-3-3 for studies of 14-3-3/phospho-Tau interaction. *PLoS One*
1010   **12**, e0178933 (2017).

1011

1012   73.   Simon MA, *et al.* High-throughput competitive fluorescence polarization assay reveals
1013   functional redundancy in the S100 protein family. *FEBS J* **287**, 2834-2846 (2020).

1014

1015   74.   Roehrl MH, Wang JY, Wagner G. A general framework for development and data analysis
1016   of competitive high-throughput screens for small-molecule inhibitors of protein-protein
1017   interactions by fluorescence polarization. *Biochemistry* **43**, 16056-16066 (2004).
1018
1019   75.   Kabsch W. Xds. *Acta Crystallogr D Biol Crystallogr* **66**, 125-132 (2010).
1020
1021   76.   McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser
1022   crystallographic software. *J Appl Crystallogr* **40**, 658-674 (2007).
1023
1024   77.   Adams PD*, et al.* PHENIX: a comprehensive Python-based system for macromolecular
1025   structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
1026
1027
1028

1029 **TABLES**

1030 **Table 1. Crystallographic statistics.**

	14-3-3ζ-18E6 chimera	14-3-3ζ-18E6 chimera + FSC
Data collection		
Wavelength	1.00	1.00
Resolution range	**39.26 - 1.9** (1.95 - 1.9)	**38.05 - 1.85** (1.9 - 1.85)
Space group	P 21 21 21	P 21 21 21
Unit cell (a, b, c, α, β, γ)	72.35, 78.53, 90.3, 90, 90, 90	73.23, 76.1, 88.95, 90, 90, 90
Total reflections	547783 (37726)	557315 (41375)
Unique reflections	41285 (2986)	41927 (3038)
Multiplicity	13.3 (12.6)	13.3 (13.6)
Completeness (%)	100 (100)	97.1 (96.5)
Mean I/sigma(I)	13.19 (1.39)	12.36 (1.40)
R-meas	16.6 (205)	16.3 (216)
CC1/2	99.9 (54.7)	99.8 (58.4)
Refinement		
R-work	0.1764	0.1908
R-free	0.2071	0.2192
Number of non-hydrogen atoms	4468	4589
macromolecules	4034	4004
ligands	24	102
solvent	410	483
Protein residues	481	482
RMS(bonds)	0.006	0.006
RMS(angles)	0.83	0.71
Ramachandran favored (%)	99.36	98.51
Ramachandran allowed (%)	0.64	1.49
Ramachandran outliers (%)	0	0
Rotamer outliers (%)	1.65	2.13
Clashscore	6.68	3.17
Average B-factor	33.39	29.6
macromolecules	32.42	28.49
ligands	62.93	32.67
solvent	41.2	38.11
Number of TLS groups	15	11
PDB ID	**6ZFD**	**6ZFG**

1031

1032

1033