Structural and Genetic Determinants of Convergence in the *Drosophila* tRNA Structure-Function Map

`Running title:` Convergence in *Drosophila* tRNAs

Version 0.2

Julie Baker Phillips[1,2] and David H. Ardell[1,3,#]

[1] Quantitative and Systems Biology Program, University of California, Merced, CA 95343

[2] Department of Biology, Cumberland University, 1 Cumberland Square, Lebanon, TN 37087

[3] Department of Molecular and Cell Biology, University of California, Merced, CA 95343

August 5, 2020

Abstract

The evolution of tRNA multi-gene families remains poorly understood, exhibiting unusual phenomena such as functional conversions of tRNA genes through anticodon shift substitutions. We improved FlyBase tRNA gene annotations from twelve *Drosophila* species, incorporating previously identified orthologies to compare substitution rates across tRNA bodies at single-site resolution. All rapidly evolving sites fell within the same metal-ion-binding pocket at the interface of the two major stacked helical domains. We applied our tRNA Structure-Function Mapper (tSFM) method independently to each *Drosophila* species and one outgroup species *Musca domestica*. We found that, although predicted tRNA structure-function maps are generally highly conserved in flies, one tRNA Class-Informative Feature (CIF) within the rapidly-evolving ion-binding pocket — Cytosine 17 (C17), ancestrally informative for lysylation identity — independently gained asparaginylation identity and substituted in

[#]Contact: dardell@ucmerced.edu

parallel across tRNA$^{\text{Asn}}$ paralogs at least once, possibly at least three times, in evolu-

tion of the genus. In *D. melanogaster*, most tRNA$^{\text{Lys}}$ and tRNA$^{\text{Asn}}$ genes are co-arrayed

in nearby heterologous gene clusters, suggesting that heterologous gene conversion

as well as structural similarities of the closely related asparaginyl-tRNA synthetase

(AsnRS) and lysyl-tRNA synthetase (LysRS) proteins may have played a role in these

changes. A previously identified Asn-to-Lys anticodon shift substitution in *D. ananas-*

sae may have arisen to compensate for the convergent and parallel gains of C17 in

tRNA$^{\text{Asn}}$ paralogs of the *D. ananassae* lineage. Our results underscore the functional

and evolutionary relevance of our tRNA structure-function map predictions and il-

luminate multiple genomic and structural factors contributing to rapid, parallel and

compensatory evolution of tRNA multi-gene families.

Keywords— class-informative feature; ion-binding pocket; parallel substitutions; convergent evolution; structure-

function map

Introduction

Transfer RNAs (tRNAs) were the first family of RNAs to be directly sequenced (Holley 1965) and

the first to be solved by X-ray crystallography (Holley et al. 1965). Historically, algorithms to esti-

mate phylogeny and substitution patterns were tested on tRNA genes (Cedergren et al. 1981; Eigen

et al. 1989). Early in the genome era, it was reported that tRNA genes can evolve to switch their func-

tional (aminoacylation) identities through anticodon shift substitutions (Saks et al. 1998). However,

the small sizes and high similarities of tRNA genes pose obstacles to inferring their orthology, which

is needed to better understand the evolutionary processes underlying functional turnover of tRNA

genes. An important step forward came from the "micro-syntenic" approach to infer tRNA gene or-

thology using flanking sequences, first applied in *Drosophila* (Rogers, Bergman, et al. 2010) and later to

other eukaryotes (Rogers and Griffiths-Jones 2014). These studies revealed that functional turnover

of tRNA genes through anticodon shift substitutions is more frequent and widespread than previ-

ously known. However, Rogers and Griffiths-Jones (2014) were unable to discern whether anticodon

shift substitutions occur more often within or between the two conserved and ancient superfamilies

of aminoacyl-tRNA synthetases (aaRSs), called Classes I and II (Eriani et al. 1990). The two super-

families may be further divided into three subclasses each (Cusack 1997), all of which pre-date the

divergence of bacteria, archaea and eukarotes as exemplified by the consistency with which aaRSs

may be used to root the statistical Tree of Life (Brown and Doolittle 1995).

More recent advances in orthology estimation for tRNAs exploit positional homology and the organization of tRNA genes as repeated elements in tandem gene arrays, and revealed both the great extent of functional turnover and important role of gene conversion in tRNA evolution Velandia-Huerto et al. 2016. It has also been shown that initiator tRNAs, the distinct functional class of tRNA initiating protein synthesis at start codons, may evolve through anticodon shift substitutions and functional conversion from elongator tRNAs (Bhattacharyya and Varshney 2016). The advances in ortholog estimation for tRNA genes have made it possible for the first time to undertake a detailed analysis of substitution rates and patterns in tRNA genes, which is one of two requirements to understand how tRNA genes evolve to switch functions. The second requirement is a means to predict the functional significance of tRNA sequence features outside the anticodon — what we call the "tRNA structure-function map."

In earlier work, we developed an approach to predict the tRNA structure-function map from genome sequences alone, based on structure-conditioned functional information statistics (Freyhult et al. 2006). Our algorithm recovered known tRNA identity elements and predicted new ones subsequently verified as functionally important in tRNA biogenesis and aminoacylation (Bailly et al. 2006; Sethi et al. 2009). Later, in a study of alphaproteobacterial evolution, we reported that the features predicted by our algorithm, which we call tRNA Class-Informative Features (CIFs), are phylogenetically informative and robust to compositional convergence and horizontal gene transfer (Amrine et al. 2014). We developed this approach further into a machine learning "phyloclassifier" to robustly predict the cyanobacterial progenitor of plastids (Lawrence et al. 2019). More recently we have shown that that tRNA structure-function maps, including Class-Informative Base-Pairs (CIBPs) and Class-Informative Mis-Pairs (CIMPs), are highly conserved both within trypanosomes and betweeen trypanosomes and humans, and also that tRNA CIFs can be used to predict functional differences between homologous enzymes from trypanosomes and humans (Kelly et al. 2020). We also showed evidence that tRNA gene clusters are conserved in trypanosomes and show clear evidence of duplication, deletion, and rearrangements (Kelly et al. 2020).

By combining data from multiple ortholog sets of microRNA genes, the *Drosophila* Twelve Genomes Consortium yielded one of the first analyses comparing evolutionary rates over different structurally-defined categories of sites in a non-coding RNA family (Drosophila 12 Genomes Consortium 2007). In the present work, we extend this approach, exploiting the conservation and high conformity (Wolfson et al. 2001) of tRNA genes to estimate and compare evolutionary rates across different structural components and at individual single- and paired-sites in tRNA genes from the *Drosophila* Twelve Genomes. We developed a strategy to optimize the selection of species to maximize the length of concatenated

alignments across tRNA ortholog sets. Analyzing our optimized alignment using MrBayes (Huelsen-
beck and Ronquist 2001; Ronquist et al. 2012), we discovered that one of several metal ion-binding
pockets in *Drosophila* tRNAs has evolved rapidly, and that this rapid evolution is directly related to
repeated convergent conversion of a tRNA CIF from one tRNA family to another, in parallel over
multiple tRNA genes on different chromosomes. We were able to identify multiple structural and
genomic factors that are likely to have contributed to this convergent evolution. Our results suggest
for the first time that anticodon shift substitutions may play a compensatory role in evolution of the
tRNA structure-function map.

Results

D. simulans Contains a Non-Canonical Initiator tRNA Gene Created by an Anti-codon Shift Substitution

We built a custom database of tRNA genes for the *Drosophila* twelve genomes based on FlyBase re-
lease 2008-07 (Tweedie et al. 2009) downloaded on October 19, 2011, which contained a total of 3494
tRNA genes. On this set, we integrated orthology annotations from Rogers, Bergman, et al. (2010),
COVE scores from tRNAscan-SE 1.3.1 (Lowe and Eddy 1997) and initiator tRNA gene predictions from
TFAM v1.3 (Tåquist et al. 2007) as described in Methods below. In addition to refolding FlyBase tRNA
gene models in tRNAscan-SE, we also refolded them with ARAGORN v1.2.34 (Laslett and Canback
2004) as an additional source of functional and structural predictions. Our initiator tRNA gene pre-
dictions were consistent with experimentally based annotations in *D. melanogaster* and across species, in
that if TFAM annotated any gene in any ortholog set from Rogers, Bergman, et al. (2010) and Rogers
and Griffiths-Jones (2014) as an initiator tRNA gene, then every other gene in that ortholog set would
also be independently annotated by TFAM as an initiator tRNA gene. We annotated between five and
seven initiator tRNA genes per *Drosophila* species we examined, for a total of 70 annotated initiator
tRNA genes across all species, of which 66 belonged to the ortholog sets from Rogers, Bergman, et al.
(2010). Our tRNA gene annotations are provided in Supplementary Data.

We found that one alloacceptor anticodon shift substitution (which converts the functional identity
of a tRNA from one amino acid to another), which was previously reported and validated as $tRNA_{CAU}^{Met}$
$\rightarrow tRNA_{CGU}^{Thr}$ (or in their notation, CAT:M $\rightarrow$ CGT:T) in *Drosophila simulans* (Rogers, Bergman, et al. 2010;
Rogers and Griffiths-Jones 2014), in fact represents the evolution of a non-canonical initiator tRNA
gene in *D. simulans* (non-canonical in the sense that its predicted tRNA product does not contain the

Table 1. Statistics on alignments of concatenated tRNA ortholog sets from subsets of *Drosophila* species

Species Subset	Species	Ortho. Sets[a]	Sites (gap-free)	Var. Sites (Partitions)[b]	P.I. Sites (Partitions)[c]
101111110000[d]	7	147	10878 (10598)	2729 (68)	27 (15)
110011100111[e]	8	85	6290 (6134)	1589 (67)	19(15)
111111110111[f]	11	83	6142 (5991)	1554 (67)	20 (15)
111111111111	12	47	3478 (3389)	896 (58)	16 (12)

[a] Number of tRNA ortholog sets from Rogers, Bergman, et al. (2010)
[b] Number of variable gap-free sites (or variable site-partitions excluding gap-free sites, of 74) over concatenated alignment
[c] Number of parsimoniously informative gap-free sites (or partitions excluding gap-free sites, of 74) over concatenated alignment
[d] Sophophora excluding *D. simulans* and *D. willistoni*
[e] All species excluding *D. sechellia D. yakuba*, *D. persimilis* and *D. willistoni*
[f] All species excluding *D. willistoni*

methionine anticodon CAU). This non-canonical initiator tRNA gene in *D. simulans* with FlyBase gene

ID gn0256165 and transcript ID tr0296323, evolved as part of Ortholog Set 183 from Rogers, Bergman,

et al. (2010). This ortholog set contains initiator tRNA orthologs in all eight species of Sophophora

excluding *D. willistoni*. The *D. simulans* ortholog carries only a single A $\rightarrow$ G anticodon shift substitution

at Sprinzl position 35 (Sprinzl coordinates are a standardized coordinate system for the consensus

universal secondary structure of tRNAs as well as conserved, more sub-function-specific structures

like the long variable arms of tRNA$^{\text{Leu}}$ and tRNA$^{\text{Ser}}$ (Sprinzl, Horn, et al. 1998)). The non-canonical

initiator tRNA gene in *D. simulans* is also identical in sequence to all other initiator tRNA genes we

annotated in *D. simulans*, except for the previously noted anticodon shift substitution at Sprinzl position

35.

Optimizing Species Exclusion to Maximize Gap-Free Alignment Length of Concatenated tRNA Ortholog Sets

We next prepared to undertake a global analysis of site-specific substitution rates and patterns in

Drosophila tRNA genes. To do this we analyzed a total of 753 orthologous sets from Rogers, Bergman,

et al. (2010) encompassing 3218 unique FlyBase tRNA transcript IDs, a subset of our reannotated Fly-

Base set. We found the representation of species to be uneven across the Rogers *et al.* ortholog sets.

Only 47 sets, about 6%, contained orthologs from all 12 *Drosophila* species (Table 1). One species in

particular, *D. willistoni*, was represented in very few sets. We proceeded to remove this species from

our subsequent substitution rate analyses. We also removed ortholog sets that contained isoacceptor

or alloacceptor anticodon shift substitutions, uncertain functional annotations, or predicted pseudo-

genes. We wished to integrate data from the largest number of ortholog sets that we could in order

to compare substitution rates across sites. However, we also wished to avoid having a large number

139	of missing or unidentified orthologs in the concatenated alignments that we made from the Rogers
140	*et al.* ortholog sets, so as to reduce the influence of missing data associated with large blocks of gap
141	characters on our results. We therefore computed statistics over automatically generated alignments
142	corresponding to 562 species sets containing seven or more of the eleven *Drosophila* species excluding
143	*D. willistoni*, in order to find the species set that resulted in concatenated alignments of greatest gap-free
144	length. To make these concatenated alignments for each species set, first we structurally aligned all
145	tRNA genes as described in Methods, and then we extracted aligned tRNA gene sequences for every
146	ortholog set that containing at least all species in the defined species set. We then concatenated these
147	extracted ortholog set alignments together and mapped all sites in this concatenated alignment into
148	site-partitions corresponding to each Sprinzl coordinate.

We found that a single species subset, labelled as "101111110000" in Table 1 and representing Sophophora excluding *D. simulans* and *D. willistoni* yielded the longest gap-free alignment concatenating the most ortholog sets, the greatest numbers of variable and parsimoniously informative sites, and the greatest average number of pairwise differences per site among all alignments we examined. We proceeded to focus on this alignment for downstream analysis as well as two others for the sake of comparison: a randomly picked one with a complementary pattern of species exclusion ("110011100111"), and one with nearly all species included, namely all eleven species except *D. willistoni* ("111111110111").

Substitution Patterns in *Drosophila* tRNAs

To compute site-specific substitution rates from our concatenated alignments of tRNA ortholog sets, we compared substitution rates across different site-partitions of our concatenated alignments corresponding either to tRNA secondary structural elements or to individual sites defined by Sprinzl coordinates (Sprinzl and Vassilenko 2005), using MrBayes 3.2.1 (Huelsenbeck and Ronquist 2001; Ronquist et al. 2012), aggregating over different functional classes of tRNA gene orthologs and using the fixed known species tree (Drosophila 12 Genomes Consortium 2007). We fitted unpaired sites and loops with the General Time Reversible (GTR) model (Lanave et al. 1984; Tavaré 1986; Rodríguez et al. 1990) with invariant sites (GTR+I), and paired-sites and stems to Double(GTR)+I (Ronquist et al. 2012). We obtained very similar results using other models including the Hasegawa-Kishino-Yano (HKY) model (Hasegawa et al. 1985) and using Gamma-distributed site-rate variation.

In our coarse-grained analysis of substitution rates by secondary structural elements we observed that, regardless of which alignments or models we used, the largest substitution rates occur in the D-loop and acceptor stem followed closely by the T-loop (Figure 1). Considering the relative constraint

Figure 1. Credible intervals, interquartile ranges and medians of substitution rates by secondary structural elements in *Drosophila* tRNAs as calculated in MrBayes 3.2.1 using the GTR+I/Double(GTR+I) models for loops/stems and the partitioned concatenated alignments indicated by bitsets as defined in Table 1.

that acceptor stems are expected to have from identity-driven interactions with proteins (Giegé, Sissler,

et al. 1998), the acceptor stem shows a surprisingly high rate of evolution. Furthermore, regardless

of which alignments or models we used, in our analysis of substitution rates by individual sites, we

found that three sites corresponding to Sprinzl coordinates 16, 17, and 60 showed markedly higher

rates of substitution than any other site except site 45 (Figure 2).

Sprinzl coordinates 16, 17 and 60 are among eight sites that form an ion-binding pocket between

the D- and T-arms (Sprinzl coordinates 15, 16, 17, 18, 19, 20, 59, and 60), at the interface of the two

tRNA helical domains (Behlen et al. 1990), as shown in Figure 3. This pocket was first described as site

1 in the original orthorhombic crystal form (Holbrook et al. 1977) and site 3 in the monoclinic struc-

Figure 2. Credible intervals, interquartile ranges and medians of substitution rates by individual site/Sprinzl coordinate in *Drosophila* tRNA genes as calculated in MrBayes 3.2.1 using the GTR+I models for individual sites and the partitioned concatenated alignments indicated by bit-sets as defined in Table 1.

Figure 3. Rapidly-evolving sites in Drosophila tRNA are part of a conserved, structurally plastic ion-binding pocket near the interface of the two tRNA helical domains. This image visualizes PDB structure accession ID 1EHZ (Shi and Moore 2000) and was rendered in VMD 1.9.2 (Humphrey et al. 1996).

ture (Jack et al. 1977). This ion-binding pocket was shown to potentially bind different metal ions including magnesium, cobalt, manganese, and lead (Jack et al. 1977; Holbrook et al. 1977; Shi and Moore 2000; Behlen et al. 1990). More recent structural work improving the resolution of the tRNA structure better resolved the structure of Sprinzl site 16, which is a fairly conserved uracil among eukaryotes (Marck and Grosjean, 2002), most likely post-translationally modified to dihydrouridine and demonstrated that this site is sensitive to ionic conditions (Shi and Moore 2000). The hypergeometric probability that three sites of elevated substitution rate randomly occur within the eight sites that form the ion-binding pocket out of 74 possible sites is $\frac{\binom{8}{3}\binom{66}{0}}{\binom{74}{3}} = 8.6 \times 10^{-4}$.

In the alignment with the most species (111111110111), we identified that 45 orthology groups spanning 14 functional classes carry substitutions in sites 16, 17 and 60 of this ion-binding pocket, as shown in Table S1 of Supplementary Materials. One set showed signs of covariance between sites 16 and 17, and another between sites 17 and 60.

Chromosomal gene location was significantly associated with substitutions in pocket sites 16, 17

and 60 ($X^2 = 56.3266$, $N = 692$, $p = 6.96 \times 10^{-11}$), and extended pocket sites ($X^2 = 87.4858$,

$N = 726$, $p < 2.2 \times 10^{-16}$, Supplementary Materials, Table S2. Substitutions in sites 16, 17 or 60

of the pocket were not associated with alloacceptor class switches overall ($X^2 = 20.9616$, $N = 672$,

$p = 0.52311$), nor were substitutions in the extended pocket ($X^2 = 23.2573$, $N = 703$, $p = 0.3873$.

Ortholog sets containing anticodon shift substitutions more often exhibited substitutions in pocket

sites 16,17, and 60 ($X^2 = 8.8797$, $N = 713$, $p = 0.002884$), but not extended pocket sites ($X^2 =$

2.7304, $N = 696$, $p = 0.09846$).

We provide estimates of all six transition and transversion rates for *Drosophila* tRNA genes in Sup-

plemetary Materials Table S3.

Class-Informative Base-Pairs and Base-Mispairs in *Drosophila melanogaster*

We used a recent reimplementation of our previously published algorithm to estimate the structure-

conditioned functional information statistics of Freyhult et al. (2006) in Python and extended it to

compute the information for all sixteen possible base-pair or base mis-pair features in the consensus

secondary structure of tRNAs, given a structural alignment and structural and functional annotations

of tRNA gene data. The software, called tSFM v1.0 for "tRNA Structure-Function Mapper," uses a

permutation-based approach to calculate the significance of the tRNA CIFs that it predicts, and cor-

rects for multiple comparisons by computing various measures of the Family-Wise Error Rate (FWER)

or False Discovery Rate, including that of Benjamini and Hochberg (Benjamini and Hochberg 1995)

and is freely available for download on GitHub. All code and data to reproduce results are available

in the Supplementary Materials.

We filtered our reannotated FlyBase tRNA gene dataset for tRNA genes from *Drosophila melanogaster*

with tRNAscan-SE 1.3.1 COVE scores of at least 50 bits and removed tRNA genes of indeterminate

function and selenocysteine tRNA genes, leaving 288 tRNA genes remaining. We then computed

tRNA Class-Informative Base Pairs and Base Mis-Pairs from these data, with only those features with

a Benjamini-Hochberg False Discovery Rate of 5% shown in Fig. 4. A great deal of functional informa-

tion appears to reside in paired features. Several base-mispairs are functionally informative, including

U13:U22 in the D-arm associated to both tRNA[Pro] and tRNA[Val] (contained in 32 tRNA genes of 32 to-

tal), U31:U39 in the C-arm contained in all six tRNA[Met] genes, C50:U64 in the T-arm contained in

nine of 11 tRNA[Ile] genes, G13:A22 in the D-arm contained in all 42 tRNA[Ser] and tRNA[Leu] genes, and

a G4:G69 mis-pair in the acceptor stem associated with nine of 16 tRNA[Thr] genes. We show a plot of

the significance of paired-site CIFs as a function of CIF information in Supplementary Figure S1. A

tRNA Class-Informative Base-Pairs and Mis-Pairs for *D. melanogaster*

from 288 tRNA genes with COVE scores ≥ 50 bits and Benjamini-Hochberg FDR < 5%

(5':3')-paired Sprinzl coordinates in tRNA acceptor (A), D, coding (C), and T arms

Figure 4. Class-Informative Base-Pairs and Base Mis-Pairs from 288 reannotated tRNA genes in *Drosophila melanogaster* with COVE scores of at least 50 bits as computed in tSFM v1.0 with a Benjamini-Hochberg False Discovery Rate of 5% (CIFs not meeting this significance threshhold are not shown). The total height of a stack of letters quantifies information potentially gained about the functional type of a tRNA by a tRNA-binding protein if it specifically recognizes the paired features indicated. The letters within each stack symbolize functional types of tRNAs, wherein IUPAC one-letter amino acid codes represent elongator tRNA aminoacylation identities and "X" symbolizes initiator tRNAs. The relative heights of letters within each stack quantifies the over-representation of tRNA functional types carrying that feature relative to the background frequency determined by the frequencies of genes of various functional types (as calculated through normalized log-odds).

224   text-file containing all statistics for paired-site and single-site CIFs in *D. melanogaster* is provided in the

225   Supplementary Code and Data.

Conservation of Class-Informative Features in Drosophila tRNAs

The wealth of information we have on *Drosophila* evolution provides an unprecedented opportunity to understand how tRNA Class-Informative Features evolve. To examine the conservation of tRNA CIFs in *Drosophila*, we computed tRNA CIFs in each species independently in the same way as we did for *D. melanogaster*, except for that we treated each site as an unpaired site. In order to polarize evolutionary changes in tRNA CIFs we additionally analyzed tRNA gene annotations and CIFs in the outgroup *Musca domestica*, which diverged from *Drosophila* between 20–80 million years ago (Wiegmann et al. 2003). As shown in Figure 5, tRNA CIFs are very highly conserved in *Drosophila* and flies generally. Broadly speaking, our predictions are very nearly identical in all fly genomes we analyzed, with one visibly clear exception standing out: C17, which occurs within the metal ion-binding pocket previously identified as having elevated substitution rates.

As shown in Fig. 6, CIF C17 is ancestrally associated with tRNA[Lys] in flies but it became associated with all or nearly all tRNA[Asn] paralogs in at least three phylogenetically distinct *Drosophila* lineages, one of which, leading to (*D. ananassae*), also underwent the Asn-to-Lys anticodon shift previously reported in Rogers, Bergman, et al. (2010). To underscore the validity and biological significance of our CIF evolution results, we emphasize that all of the 138 tRNA[Asn] genes analyzed in Fig. 6 have COVE scores of 50 bits are greater — in fact, all but two of them have scores above 70 bits, while all but six have scores above 80 bits. In addition, 86 of 111 FlyBase tRNA[Asn] genes belong to the Rogers *et al.* ortholog sets. For comparison's sake, ten of 111 FlyBase records for tRNA[Asn] genes have COVE scores below 50 bits, of which two belong to the Rogers *et al.* ortholog sets. We emphasize that the Rogers *et al.* ortholog sets have twice undergone rigorous peer review. Looking more closely at sequence variation, we found that all of the tRNA[Asn] genes that have undergone substitutions to C17 mostly have zero additional substitutions and look otherwise like other tRNA[Asn] genes rather than tRNA[Lys] genes. Of the 29 genes that substituted to C17 that belong to the Rogers *et al.* ortholog sets, 15 are considered "core genes" in the sense of Rogers *et al.*. All of them were annotated as tRNA[Asn] genes by TFAM.

Separate logo figures predicting tRNA CIFs in each species that we analyzed are provided in the Supplementary Materials. In Figure S2, also in Supplementary Materials, we show that as expected the relative average information of single-site tRNA CIFs is inversely related to its relative compositional frequency in *D. melanogaster*. The relative frequency of bases in *D. melanogaster* tRNAs decreases from G to C to U to A, while relative average information increases correspondingly.

Figure 5. Conservation of tRNA single-site CIFs in *Drosophila* with *Musca domestica* as an outgroup. Tree topology from Drosophila 12 Genomes Consortium (2007), with divergence dates from Tamura et al. (2003) and Hennig et al. (1981).

Discussion

257   In an analysis of tRNA gene ortholog sets from flies, primates and other groups, Rogers and Griffiths-

258   Jones (2014) reported no evidence of preferential anticodon shift substitutions within or between func-

259   tional classes of tRNAs charged by either of the two aminoacyl-tRNA synthetase superfamilies, Class

260   I or Class II. However, they did report that an Asn-to-Lys alloacceptor anticodon shift substitution is

Figure 6. Evolution of functional association of CIF C17 in flies. Cytosine single-site function logos are from Fig. 5, the tree topology is from Drosophila 12 Genomes Consortium (2007), and divergence dates are from Tamura et al. (2003) and Hennig et al. (1981). The three clades in which C17 became associated with tRNA^{Asn} genes are indicated by purple boxes on the tree, one of which (*D. ananassae*), also underwent the Asn-to-Lys anticodon shift previously reported in Rogers, Bergman, et al. (2010) as indicated in top purple box at figure left. The stacked bar graph at figure right shows the frequencies of U17 or C17 across all tRNA^{Asn} genes in each genome showing that C17 substituted in parallel across all tRNA^{Asn} genes in these three lineages.

261  one of only two types of anticodon shift substitutions that occurred more than once out of a total of 30

262  in their dataset, once in *Drosophila* and twice independently in primates. In this work, we show that in

263  flies, the previously reported Asn-to-Lys anticodon shift substitution occurs together with co-option

264  of CIF C17 from tRNA^{Lys} to tRNA^{Asn} in one lineage, that this co-option of CIF C17 from tRNA^{Lys} to

265  tRNA^{Asn} occurred at least three times independently in *Drosophila* evolution, and that when it occurred,

266  it was associated with parallel substitutions across all or most tRNA^{Asn} genes. We note that AsnRS

267  and LysRS are both Class IIb synthetases, suggesting that perhaps co-option of tRNA CIFs occurs more

268  frequently within aaRS sub-classes, perhaps because aaRSs of the same sub-class have more similar

269  binding interfaces on tRNAs Giegé, Touzé, et al. 2008. In our recently published theory for the evo-

270  lution of aaRS-tRNA interaction networks, tRNA mutations can influence interactions with multiple

271  aaRSs when they occur inside shared interfaces (Collins-Hed and Ardell 2019). The pattern we re-

272  port here is consistent with an interpretation that the Asn-to-Lys anticodon shift substitution evolved

273  to compensate the co-option of C17 from Lys to Asn in the *D. ananassae* lineage.

274    Comparing the genomic locations of tRNA^{Lys} and tRNA^{Asn} genes from the FlyBase data and vi-

sualizing them in the UCSC Genome Browser (Kent et al. 2002) for *Drosophila melanogaster*, we found

that eight of ten tRNA^{Asn} genes co-occur in a heterologous gene array along with eight tRNA^{Lys} genes

in both direct and inverted orientations within approximately 45 Kb on Chromosome 2R, far away

from two additional tRNA^{Asn} gene singletons, one on 2R and another on 3R. While a renewed tRNA

gene reannotation effort with long-read assembly data is needed to better understand the causes of

the changes we have reported here, the location data in *D. melanogaster* is consistent with a scenario in

which gene conversion by illegitimate recombination may have played a role in the convergent and

parallel evolution of the tRNA structure-function map in *Drosophila* that we describe here. Further-

more, if the organization of tRNA^{Asn} genes in *D. melanogaster* are broadly representative across species,

a simple scenario of concerted gene evolution (Nei and Rooney 2005) is insufficient to explain the par-

allel substitutions of C17 in tRNA^{Asn} genes that we observed. Rapid evolution of tRNA gene arrays

through rearrangements appears to be universal in eukaryotes (Velandia-Huerto et al. 2016) and at

least some prokaryotes (Tremblay-Savard et al. 2015).

In summary, interactions at least three different levels of biological structure appear to have con-

tributed to a specific and recurring pattern of rapid, parallel evolution and functional turnover of

tRNA genes in *Drosophila*: of physically interacting sites within tRNA tertiary structure, of co-clustered

tRNA genes within genomes, and of overlapping interfaces for structurally similar and closely-related

tRNA-binding proteins. Further work is needed to generalize the observations we have made and to

discern their causes.

Methods

Data, Ortholog Sets and Alignments

We obtained tRNA gene sequences and annotations for 12 species of *Drosophila* from FlyBase (2008_07 release, Mc-

Quilton et al. 2012) on October 16, 2011. For further functional re-annotation, we re-folded and re-annotated these

tRNA sequences using tRNAscan-SE 1.3.1 (Lowe and Eddy 1997) and ARAGORN 1.2.34 (Laslett and Canback

2004). The genome for *Musca domestica* (Scott et al. 2014) was downloaded from NBCI on October 8th, 2013. The

genome was annotated using predictions from tRNAscan-SE 1.3.1 (Lowe and Eddy 1997).

We identified Initiator tRNA genes using TFAM 1.3 (Tåquist et al. 2007). We aligned all tRNAs using Infernal

1.1 (Nawrocki et al. 2009) with the RFAM covariance model for tRNAs (RF00005) (Burge et al. 2012). We edited

alignments manually using Seaview 4.3.4 (Gouy et al. 2010) to produce a final alignment of length 74, and mapped

each site manually to Sprinzl coordinates (Sprinzl and Vassilenko 2005). We verified our coordinate mapping using

tRNAdb (Jühling et al. 2009). We retained Sprinzl coordinate 20A and removed the majority of the variable arm,

except Sprinzl coordinates 45 through 49, for subsequent analysis.

We downloaded ortholog sets of tRNA genes on October 18, 2011 from supplementary data available at `http:`

`//gbe.oxfordjournals.org/content/2/467/suppl/DC1` (Rogers, Bergman, et al. 2010). To optimize or-

tholog sets, we calculated lengths of pruned species trees using the Bio::TreeIO module in BioPerl 1.4.0 (Stajich,

Block, et al. 2002) and calculated numbers of variable and parsimoniously informative sites of alignments, as well

as average fraction of pairwise differences, using the Bio::PopGen modules (Stajich and Hahn 2005) in BioPerl 1.4.0.

Analysis of Substition Rates

All subsets of data were curated into concatenated alignments by previously published ortholog sets (Rogers,

Bergman, et al. 2010). Ortholog sets with anticodon shift substitutions, genes of indeterminate function, or pseu-

dogenes were removed from all analyses of substitution rates.

We estimated substitution rates with MrBayes 3.2.1 (Huelsenbeck and Ronquist 2001; Ronquist et al. 2012) using

the fixed known species tree (Drosophila 12 Genomes Consortium 2007). For all runs we constrained change to the

tree topology by setting rates of stochastic TBR and branch multipliers to zero probability. All Bayesian analyses

were run with two simultaneous chains for 4×10^6 iterations, monitoring convergence of split frequency standard

deviations and saved parameters every 500 iterations.

We computed evolutionary rates by assigning sites or site-pairs into corresponding data partitions. For struc-

tural data partitions, we considered nine structural tRNA components: acceptor stem (Sprinzl sites in *Drosophila*: 1

$- 7, 67 - 73$), D stem ($10 - 13, 23 - 26$), D loop ($14 - 22$), anticodon stem ($28 - 32, 40 - 44$), anticodon loop ($33 - 39$),

variable arm ($45 - 49$), T stem ($50 - 54, 62 - 66$), T loop ($55 - 61$), and "other sites" ($8, 9, 27, 74$). Sites in "other" are

not involved in base-pairs or considered part of loop structures in tRNA. We estimated rates using the General Time

Reversible (GTR) substitution model (Lanave et al. 1984; Tavaré 1986; Rodríguez et al. 1990) with Invariant sites, al-

lowing stationary state frequencies and all other substitution model parameters to be independent across partitions.

Alignment data and MrBayes initialization scripts in NEXUS format with corresponding alignments are provided

in supplementary data. The first 25% of parameter calculations were discarded as burn-in for statistical analysis. Pa-

rameter posterior probabilities were imported to R (R Core Team 2013). We used the coda package (Plummer et al.

2006) for Markov Chain Monte Carlo simulations diagnostics and the lattice package (Sarkar 2008) for multivariate

analysis. All data from MCMC simulations is presented with 95% Bayesian credible intervals.

Detection of Class-Informative Features

For analysis of tRNA CIFs, we removed genes with anticodon shift substitutions, genes of indeterminate function,

and pseudogenes and we further filtered genes to have tRNAscan-SE COVE scores (Lowe and Eddy 1997) of at least

50 bits. We identified Class-Informative features (CIFs) for all 12 species of *Drosophila* using tSFM v1.0, available

from `https://github.com/tlawrence3/tSFM`, using the Nemenman-Shafee-Bialek (NSB) Bayesian entropy

estimator (Nemenman et al. 2002) for features in two or more sequences and an exact estimator (Schneider et al.

1986) otherwise. To compute the significance of paired-site CIFs in *D. melanogaster*, we computed permutation *p*-

values (permuting functional class assignments over sequences that contain a CIF) for the total information of

paired-sites only, and then computed Benjamini-Hochberg False Discovery Rates (Benjamini and Hochberg 1995)

from those *p*-values.

Data Access

Acknowledgments

347 The authors thank Carolin Frank, Michael Cleary, Michael Colvin, Josh Phillips and David Pollock for helpful dis-

348 cussions. DHA was supported by a National Science Foundation grant (INSPIRE-1344279), a National Institutes of

349 Health grant (AI127582), and a Julius Kuhn Guest Professorship at Martin Luther University in Halle-Wittenberg,

350 United States. Computational research in this report was performed on the MERCED HPC cluster supported by

351 the National Science Foundation, United States (ACI-1429783).

References

Amrine KCH, Swingley WD, and Ardell DH. 2014. tRNA signatures reveal a polyphyletic origin of SAR11

strains among alphaproteobacteria. *PLoS computational biology*. **10**: e1003454.

Bailly M, Giannouli S, Blaise M, Stathopoulos C, Kern D, and Becker HD. 2006. A single tRNA base pair

mediates bacterial tRNA-dependent biosynthesis of asparagine. *Nucleic Acids Res*. **34**: 6083–6094.

Behlen LS, Sampson JR, DiRenzo AB, and Uhlenbeck OC. 1990. Lead-catalyzed cleavage of yeast tRNAPhe

mutants. *Biochemistry*. **29**: 2515–2523.

Benjamini Y and Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Ap-

proach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*. **57**: 289–300.

Bhattacharyya S and Varshney U. 2016. Evolution of initiator tRNAs and selection of methionine as the

initiating amino acid. *RNA Biology*. **13**: 810–819.

Brown JR and Doolittle WF. 1995. Root of the universal tree of life based on ancient aminoacyl-tRNA

synthetase gene duplications. *Proc Natl Acad Sci U S A*. **92**: 2441–2445.

Burge SW, Daub J, Eberhardt R, Tate J, Barquist L, Nawrocki EP, Eddy SR, Gardner PP, and Bateman A.

2012. Rfam 11.0: 10 years of RNA families. *Nucleic Acids Research*. **41**: D226–D232.

Cedergren RJ, Sankoff D, LaRue B, and Grosjean H. 1981. The evolving tRNA molecule. *CRC Critical Reviews*

in Biochemistry. **11**: 35–104.

Collins-Hed AI and Ardell DH. 2019. Match fitness landscapes for macromolecular interaction networks:

Selection for translational accuracy and rate can displace tRNA-binding interfaces of non-cognate

aminoacyl-tRNA synthetases. *Theor Popul Biol*.

Cusack S. 1997. Aminoacyl-tRNA synthetases. *Current Opinion in Structural Biology*. **7**: 881–889.

Drosophila 12 Genomes Consortium. 2007. Evolution of genes and genomes on the Drosophila phylogeny.

Nature. **450**: 203–18.

Eigen M, Lindemann BF, Tietze M, Winkler-Oswatitsch R, Dress A, and Haeseler A von. 1989. How old is

the genetic code? Statistical geometry of tRNA provides an answer. *Science*. **244**: 673–679.

Eriani G, Delarue M, Poch O, Gangloff J, and Moras D. 1990. Partition of tRNA synthetases into two classes

based on mutually exclusive sets of sequence motifs. *Nature*. **347**: 203–206.

Freyhult E, Moulton V, and Ardell DH. 2006. Visualizing bacterial tRNA identity determinants and an-

tideterminants using function logos and inverse function logos. *Nucleic Acids Research*. **34**: 905–916.

Giegé R, Sissler M, and Florentz C. 1998. Universal rules and idiosyncratic features in tRNA identity. *Nucleic*

Acids Res. **26**: 5017.

Giegé R, Touzé E, Lorber B, Théobald-Dietrich A, and Sauter C. 2008. Crystallogenesis Trends of Free and

Liganded Aminoacyl-tRNA Synthetases. *Crystal Growth & Design*. **8**: 4297–4306.

Gouy M, Guindon S, and Gascuel O. 2010. SeaView version 4: A multiplatform graphical user interface

for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution*. **27**: 221–224.

Hasegawa M, Kishino H, and Yano T. 1985. Dating of the human-ape splitting by a molecular clock of

mitochondrial DNA. *Journal of molecular evolution*. **22**: 160–174.

Hennig W et al. 1981. Insect phylogeny. In. John Wiley & Sons Ltd.

Holbrook SR, Sussman JL, Warrant RW, Church GM, and Kim SH. 1977. RNA-ligand interactions:(I) mag-

nesium binding sites in yeast tRNAPhe. *Nucleic Acids Research*. **4**: 2811–2820.

Holley RW. 1965. Structure of an alanine transfer ribonucleic acid. *Journal of the American Medical Association*.
194: 868–871.

Holley RW, Apgar J, Everett GA, Madison JT, Marquisee M, Merrill SH, Penswick JR, and Zamir A. 1965.
Structure of a Ribonucleic Acid. *Science*. **147**: 1462–1465.

Huelsenbeck JP and Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*.
17: 754–755.

Humphrey W, Dalke A, and Schulten K. 1996. VMD: visual molecular dynamics. *Journal of Molecular Graphics*.
14: 33–8– 27–8.

Jack A, Ladner JE, Rhodes D, Brown RS, and Klug A. 1977. A crystallographic study of metal-binding to
yeast phenylalanine transfer RNA. *Journal of Molecular Biology*. **111**: 315–328.

Jühling F, Mörl M, Hartmann RK, Sprinzl M, Stadler PF, and Pütz J. 2009. tRNAdb 2009: compilation of
tRNA sequences and tRNA genes. *Nucleic Acids Research*. **37**: D159–62.

Kelly P, Hadi-Nezhad F, Liu DY, Lawrence TJ, Linington RG, Ibba M, and Ardell DH. 2020. Targeting tRNA-
synthetase interactions towards novel therapeutic discovery against eukaryotic pathogens. *PLoS Negl*
Trop Dis. **14**: e0007983.

Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, and Haussler aD. 2002. The Human
Genome Browser at UCSC. *Genome Res.* **12**: 996–1006.

Lanave C, Preparata G, Saccone C, and Serio G. 1984. A new method for calculating evolutionary substi-
tution rates. *Journal of Molecular Evolution*. **20**: 86–93.

Laslett D and Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nu-
cleotide sequences. *Nucleic Acids Research*. **32**: 11–16.

Lawrence TJ, Amrine KC, Swingley WD, and Ardell DH. 2019. tRNA functional signatures classify plastids
as late-branching cyanobacteria. *BMC Evol. Biol.* **19**: 224.

Lowe TM and Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in
genomic sequence. *Nucleic Acids Research*. **25**: 955–964.

McQuilton P, St Pierre SE, Thurmond J, and FlyBase Consortium. 2012. FlyBase 101– the basics of navi-
gating FlyBase. *Nucleic Acids Research*. **40**: D706–14.

Nawrocki EP, Kolbe DL, and Eddy SR. 2009. Infernal 1.0: inference of RNA alignments. *Bioinformatics*. **25**:
1335–1337.

Nei M and Rooney AP. 2005. Concerted and Birth-and-Death Evolution of Multigene Families. *Annual*

Review of Genetics. **39**: 121–152.

Nemenman I, Shafee F, and Bialek W 2002. Entropy and inference, revisited. English (US). In: *Advances*

in Neural Information Processing Systems 14 - Proceedings of the 2001 Conference, NIPS 2001. Advances in Neural

Information Processing Systems. Neural information processing systems foundation.

Plummer M, Best N, Cowles K, and Vines K. 2006. CODA: Convergence Diagnosis and Output Analysis

for MCMC. *R News*. **6**: 7–11.

R Core Team 2013. R: A Language and Environment for Statistical Computing. R Foundation for Statistical

Computing. Vienna, Austria.

Rodríguez F, Oliver JL, Marín A, and Medina JR. 1990. The general stochastic model of nucleotide substi-

tution. *Journal of Theoretical Biology*. **142**: 485–501.

Rogers HH, Bergman CM, and Griffiths-Jones S. 2010. The evolution of tRNA genes in Drosophila. *Genome*

Biol Evol. **2**: 467–477.

Rogers HH and Griffiths-Jones S. 2014. tRNA anticodon shifts in eukaryotic genomes. *RNA*. **20**: 269–281.

Ronquist F, Teslenko M, Mark P van der, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, and

Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across

a large model space. *Systematic Biology*. **61**: 539–542.

Saks ME, Sampson JR, and Abelson J. 1998. Evolution of a Transfer RNA Gene Through a Point Mutation

in the Anticodon. *Science*. **279**: 1665–1670.

Sarkar D. 2008. Lattice: Multivariate Data Visualization with R. In. Springer, New York.

Schneider TD, Stormo GD, Gold L, and Ehrenfeucht A. 1986. Information content of binding sites on

nucleotide sequences. *Journal of Molecular Biology*. **188**: 415–431.

Scott JG et al. 2014. Genome of the house fly, Musca domestica L., a global vector of diseases with adap-

tations to a septic environment. *Genome Biol.* **15**: 466.

Sethi A, Eargle J, Black AA, and Luthey-Schulten Z. 2009. Dynamical networks in tRNA:protein complexes.

Proceedings of the National Academy of Sciences of the United States of America. **106**: 6620–6625.

Shi H and Moore PB. 2000. The crystal structure of yeast phenylalanine tRNA at 1.93 Å resolution: a classic

structure revisited. *RNA*. **6**: 1091–1105.

Sprinzl M, Horn C, Brown M, Ioudovitch A, and Steinberg S. 1998. Compilation of tRNA sequences and
sequences of tRNA genes. *Nucleic Acids Research*. **26**: 148–153.

Sprinzl M and Vassilenko KS. 2005. Compilation of tRNA sequences and sequences of tRNA genes. *Nucleic*
Acids Research. **33**: D139–40.

Stajich JE, Block D, et al. 2002. The Bioperl toolkit: Perl modules for the life sciences. *Genome Research*. **12**:
1611–1618.

Stajich JE and Hahn MW. 2005. Disentangling the effects of demography and selection in human history.
Molecular Biology and Evolution. **22**: 63–73.

Tamura K, Subramanian S, and Kumar S. 2003. Temporal patterns of fruit fly (*Drosophila*) evolution revealed
by mutation clocks. *Molecular Biology and Evolution*. **21**: 36–44.

Tåquist H, Cui Y, and Ardell DH. 2007. TFAM 1.0: an online tRNA function classifier. *Nucleic Acids Research*.
35: W350–3.

Tavaré S. 1986. Some probabilistic and statistical problems in the analysis of DNA sequences. *Lectures in*
Mathematics of Life Sciences. **17**: 57–86.

Tremblay-Savard O, Benzaid B, Lang BF, and El-Mabrouk N. 2015. Evolution of tRNA repertoires in Bacil-
lus inferred with OrthoAlign. *Mol Biol Evol*. **32**:

Tweedie S et al. 2009. FlyBase: enhancing Drosophila Gene Ontology annotations. *Nucleic Acids Res*. **37**:
D555–D559.

Velandia-Huerto CA, Berkemer SJ, Hoffmann A, Retzlaff N, Romero Marroquín LC, Hernández-Rosales
M, Stadler PF, and Bermúdez-Santana CI. 2016. Orthologs, turn-over, and remolding of tRNAs in
primates and fruit flies. *BMC Genomics*. **17**: 617.

Wiegmann BM, Yeates DK, Thorne JL, and Kishino H. 2003. Time Flies, a New Molecular Time-Scale for
Brachyceran Fly Evolution Without a Clock. *Systematic Biology*. **52**: 745–756.

Wolfson AD, LaRiviere FJ, Pleiss JA, Dale T, Asahara H, and Uhlenbeck OC 2001. tRNA Conformity. In:
The Ribosome. CSHL Press, pp. 185–193.