The *App^NL-G-F* mouse retina is a site for preclinical Alzheimer's disease diagnosis and research

Marjan Vandenabeele[1,2], Lien Veys[1,2], Sophie Lemmens[3], Xavier Hadoux[4], Géraldine Gelders[1,2], Luca

Masin[1,2], Lutgarde Serneels[2,5], Jan Theunis[6], Takashi Saito[7,8], Takaomi C. Saido[7], Murali Jayapala[9], Patrick

De Boever[6,10], Bart De Strooper[2,5], Ingeborg Stalmans[3], Peter van Wijngaarden[4], Lieve Moons[1,2], Lies De

Groef[1,2]*

[1]KU Leuven, Neural Circuit Development and Regeneration Research Group, Department of Biology,

Leuven, Belgium

[2]Leuven Brain Institute, Leuven, Belgium

[3]University Hospitals Leuven, Department of Ophthalmology, Leuven, Belgium; KU Leuven, Research

Group Ophthalmology, Department of Neurosciences, Leuven, Belgium

[4]Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne, Australia;

Ophthalmology, Department of Surgery, University of Melbourne, Parkville, Australia

[5]VIB Center for Brain & Disease Research, Leuven, Belgium; KU Leuven, Laboratory for the Research of

Neurodegenerative Diseases, Department of Neurosciences, Leuven, Belgium

[6]VITO (Flemish Institute for Technological Research), Health Unit, Mol, Belgium

[7]Laboratory for Proteolytic Neuroscience, RIKEN Center for Brain Science, Saitama, Japan

[8]Department of Neurocognitive Science, Institute of Brain Science, Nagoya City University Graduate

School of Medical Sciences, Japan

[9]Imec (Interuniversity Microelectronics Centre), Leuven, Belgium

[10] Hasselt University, Center of Environmental Sciences, Diepenbeek, Belgium; University of Antwerp,

Department of Biology, Wilrijk, Belgium

*Corresponding author:

Lies De Groef

Neural Circuit Development and Regeneration Research Group

Department of Biology

University of Leuven (KU Leuven)

Naamsestraat 61, box 2464

3000 Leuven

Belgium

Tel. + 32 (0)16 37 37 64

Abstract

In this study, we report the results of a comprehensive phenotyping of the retina of the App^{NL-G-F} mouse. We demonstrate that soluble Aβ accumulation is present in the retina of these mice early in life and progresses to Aβ plaque formation by midlife. This rising Aβ burden coincides with local microglia reactivity, astrogliosis, and abnormalities in retinal vein morphology. Electrophysiological recordings reveal signs of neuronal dysfunction yet no overt neurodegeneration was observed and visual performance outcomes were unaffected in the App^{NL-G-F} mouse. Furthermore, we show that hyperspectral imaging can be used to quantify retinal Aβ, underscoring its potential as a biomarker for AD diagnosis and monitoring. These findings suggest that the App^{NL-G-F} retina mimics the early, preclinical stages of AD, and, together with retinal imaging techniques, offers unique opportunities for drug discovery and fundamental research into preclinical AD.

I.   Introduction

Alzheimer's disease (AD) is the number one neurodegenerative disorder and cause of dementia. Accumulation of amyloid-beta (Aβ) plaques and hyperphosphorylated Tau in tangles are two hallmarks of AD, and are believed to lead to neurotoxic inflammation, neuronal dysfunction, and eventually neurodegeneration. Despite decades of research, the AD field is still struggling with finding techniques for adequate diagnosis and disease monitoring, rational treatment strategies and valid research models.

AD is supposed to start 20 years before the first cognitive symptoms appear [20, 39, 51], and it is nowadays believed that one should focus on this time window for diagnosis and future treatment [19, 47]. In the search for new biomarkers and animal models that allow evaluation of this preclinical phase of AD, increasing attention has gone to the retina. Indeed, given that the retina is an integral part of the central nervous system, cellular and molecular mechanisms underlying neurodegeneration are conserved, and pathological processes occurring in the retina may be an indicator of those processes ongoing in other parts of the central nervous system [3, 11, 26]. In addition, state-of-the-art technologies for ocular imaging (e.g. optical coherence tomography and confocal scanning laser ophthalmoscopy) allow these processes to be visualised at a resolution of at least an order of a magnitude higher than conventional brain imaging techniques, without the need for invasive, costly procedures or tracers, and in a reproducible and quantifiable manner [11].

One emerging technique in the field of retinal biomarkers for AD, is retinal hyperspectral imaging. This non-invasive imaging spectroscopy technique relies on the wavelength-dependent effect of Aβ on light scatter, to measure the reflectance spectrum of a locus over a range of wavelengths and convert this into a hyperspectral score of which the magnitude correlates to the amount of Aβ. The exact pathological correlate of this hyperspectral signature is still unclear, although previous *in vitro* studies and theoretical

modeling suggested that soluble $A\beta_{42}$ is being measured [29, 30]. The latter would be a particular

advantage, as converging evidence suggests that, in the preclinical phase of AD, soluble oligomeric Aβ

accumulates to a threshold that eventually leads to symptomatic AD. Rather than intermediates to

generation of Aβ plaques, these Aβ oligomers are nowadays regarded as being the most pathogenic form

of Aβ. Aβ oligomers initiate neuropathological processes such as neuroinflammation, gliosis, degradation

of synapses, impaired axonal transport, and oxidative damage, thereby leading to an enlarging pool of

neurons with limited functionality and ultimately driving the diseased brain to the tipping point where

actual neurodegeneration ensues [2, 7, 15, 52].

A multitude of studies has evaluated the retinal phenotype of AD mouse models, and revealed retinal

abnormalities that mirror observations in human AD subjects, including elevated Aβ levels, Aβ deposits

co-localizing with sites of apoptosis, tauopathy, neurodegeneration of ganglion and amacrine cells,

microgliosis and astrogliosis, abnormal electrophysiology, as well as tight junction attenuation and vessel

malformations [4, 10, 16, 24, 35]. However, results are often conflicting, and interpretation of their

biological relevance is obscured by the fact that these transgenic AD mice all rely on ectopic

overexpression of familial AD-associated genes (mutations). Here, we address this shortcoming with a

comprehensive retinal phenotyping study of the App^{NL-G-F} knock-in mouse, which is characterized by

physiological expression of humanized mouse APP, containing the Swedish, Beyreuther/Iberian and Arctic

mutations, in disease-relevant central nervous system regions and cell types [32, 40]. For this first-time

retinal phenotyping study of the App^{NL-G-F} mouse, we heavily relied on *in vivo* techniques to assess the

possibility of longitudinal disease monitoring via retinal imaging and electrophysiology, besides the gold

standard *post mortem* assays. Additionally, we evaluated the use of hyperspectral imaging to quantify

retinal Aβ burden, based on the hypothesis that it detects soluble $A\beta_{42}$ and therefore may offer a way to

study preclinical AD. Altogether, this approach represents a novel tool box, with the retina of the App^{NL-G-}

F mouse as a model and retinal imaging techniques as unique read-outs, for both fundamental and drug

discovery research in the preclinical phase of AD.

II.   Materials and methods

<u>Animals</u>

B6.129S5-*App*tm3.1Tcs (here referred to as App^{NL-G-F}) [32, 40], App^{NL-G-F} x CX_3CR-$1^{GFP/+}$ (generated via

crossing of App^{NL-G-F} and B6.129P2-*Cx3cr1*tm1Litt (The Jackson Laboratory)), B6;C3Tg

(APPswe,PSEN1dE9)85Dbo/Mmjax (here referred to as APP/PS1) mouse lines and corresponding wild type

(WT) controls were used. The App^{NL-G-F} mouse carries an *App* gene construct, which contains a humanized

Aβ region and includes the Swedish "NL", Iberian "F", and Arctic "G" mutations. The APP/PS1 mouse

contains transgenes for both APP and presenilin 1 (PSEN1 or PS1), under the control of the mouse prion

promotor, with chimeric mouse/human APP with the Swedish mutation, and human PS1 lacking exon 9.

The App^{NL-G-F} model was chosen as it avoids potential artifacts introduced by APP overexpression by using

a knock-in approach to express APP at endogenous levels and with appropriate cell-type and temporal

specificity. The APP/PS1 overexpression mouse was included in this study as a reference AD model, given

that its retinal phenotype is well-described, e.g. [9, 12, 22, 29, 33, 45, 54]. Mice were bred under standard

laboratory conditions. App^{NL-G-F} and WT female mice were sacrificed at 3, 6, 9, 12 and 18 months of age (±

1 month). Male mice were also included for the 18-month age group.

<u>Aβ extraction and ELISA detection</u>

For soluble Aβ extraction, two retinas from one mouse were pooled and homogenized in lysis buffer (50

mM NaCl, 0.4 % diethanolamine, and protease inhibitor cocktail (Roche)). Samples were spun down and

supernatant was centrifuged for 50 min at 100.000 x g. Supernatant containing the soluble Aβ fraction

was collected. From the pellet, insoluble Aβ was extracted with GuHCl (6 M GuHCl/50 mM Tris-HCl) by

sonication, incubation at 25°C for 60 min and centrifugation for 20 min at 170.000 x g. Aβ levels were

determined via ELISA Meso Scale Discovery 96-well plates coated with antibodies for $A\beta_{40}$ JRFcAβ40/28

and $A\beta_{42}$ JRFcAβ42/26. JRFAβN/25 (human specific N-terminal epitope) labeled with sulfo-TAG was used

as the detection antibody. Antibodies were provided by Dr. Marc Mercken (Janssen Pharmaceutica).

<u>Immunohistochemistry</u>

Following perfusion with 4% paraformaldehyde (PFA), eyes were enucleated and post-fixated in 4% PFA

for 1 hour. For cryosections (12 μm), eyes went through a sucrose gradient series for cryoprotection prior

to embedding in O.C.T. compound (Tissue-Tek, Sakura). For wholemount preparations, retinas were

flatmounted and post-fixated for one hour in 4% PFA. For Aβ immunostaining and hyperspectral imaging

of retinal wholemounts, no additional fixation was done.

Retinal cryosections were stained for glial acidic fibrillary protein (GFAP), S100 calcium binding protein B

(S100B), brain-specific homeobox/POU domain protein 3A (Brn3a), Ceh-10 Homeodomain-Containing

Homolog (Chx10) or oligomeric Aβ (A11). For GFAP, antigen retrieval was performed via proteinase K

treatment (5 min, 2 μg/ml, Qiagen). After blocking, sections were incubated overnight with anti-GFAP

(1:2000, #Z0334, DAKO), anti-S100B (1:600, #ab52642, Abcam), anti-Brn3a (1:750, #sc-31984, Santa Cruz),

anti-Chx10 (1:200, #X1180P, Sanbio) or A11 (1:500, #AHB0052, Invitrogen) primary antibodies. For GFAP,

S100B, Brn3a and Chx10, an Alexa-conjugated secondary antibody (Invitrogen) was used; for A11, a biotin-

conjugated antibody (Jackson ImmunoResearch) followed by tyramide signal amplification (Perkin Elmer)

was used. 4',6-diamidino-2-phenylindole (DAPI) was used as a nuclear counterstain.

For wholemount (immuno)staining, wholemounts were frozen for 15 min at -80°C, and incubated

overnight with primary antibodies for melanopsin (1:5000, #AB-N38, Advanced Targeting Systems),

ionized calcium-binding adapter molecule 1 (Iba1; 1:1000, #019-19741, Wako), choline acetyltransferase

(ChAT; 1:500, #AB144P, Millipore), platelet-derived growth factor receptor beta (PDGFRβ) (1:500,

#ab32570, Abcam), GFAP (1:1000, #Z0334, DAKO), RNA-binding protein with multiple splicing (RBPMS)

(1:250, #1830-RBPMS, PhosphoSolutions), protein kinase C alpha (PKCα) (1:100, #sc-10800, Santa Cruz),

green fluorescent protein (GFP; 1:500, #ab13970, Abcam) and/or Aβ (6E10, 1:500, #803001, BioLegend

and 82E1, 1:300, #10323, Immuno-Biological Laboratories), or with biotinylated isolectin B4 (1:50, #B-

1205, Vector Laboratories), followed by incubation with corresponding Alexa-conjugated secondary

antibodies or Alexa-conjugated streptavidin (Invitrogen). For Aβ staining, a three-minute step in 100%

formic acid was performed before freezing, as well as an extra blocking with goat anti-mouse Fab

fragments (1:50, Jackson ImmunoResearch) for 1 hour.

Microscopy and image analysis

Microscopy was performed either via confocal or multiphoton imaging (Olympus FV1000 or FV1000-M;

for Aβ, Iba1, Cx3cr1-GFP, ChAT and isolectin B4) or conventional fluorescence microscopy (Leica DM6; for

melanopsin, GFAP and S100B). For confocal or multiphoton imaging of retinal wholemounts, mosaic z-

stack images (step size 2-4 μm) were taken. Image analyses were performed using Fiji software [42], unless

stated otherwise.

(i)   *Microglia numbers and morphological parameters* were analyzed on maximum intensity

projection images of retinal wholemounts, as described previously [5].

(ii)   *Macroglia reactivity* was quantified on retinal sections by manually counting GFAP-

immunopositive fibers at the inner plexiform-inner nuclear layer border [43]. GFAP+ fibers

were counted over a length of 300 μm, on 20 images per mouse (two peripheral and two

central images per section; five sections per mouse, including the central section containing

the optic nerve head, and the sections located 240 and 480 μm anterior/posterior). For S100B,

the immunopositive area was measured in the nerve fiber/ganglion cell layer over a length of

300 μm, on 20 images per mouse (*cfr*. above) [18].

(iii)   *Retinal vasculature* was analyzed on retinal wholemounts, on maximum intensity projections

168   from the upper vasculature. Vessel density and branching complexity was analyzed with

169   AngioTool software [56] on a binarized image. Retinal vessel diameter was measured

170   manually at a distance of 300-400 μm from the optic nerve head; 5 measurements were

171   averaged per vessel.

172   *(iv)*   *RBPMS+ retinal ganglion cells* were counted on retinal wholemounts, using a validated

173   automated counting method [27].

174   *(v)*   *Melanopsin+ retinal ganglion cell* numbers were counted manually on wholemount images

175   [50].

176   *(vi)*   *ChAT+ cells* were counted in 3D, using the 'spots' module of Imaris software (version 7.4.2) on

177   z-stack pictures of the wholemounts.

Optical coherence tomography

Upon general anesthesia and pupil dilatation (0.5% tropicamide, Tropicol, Théa), optical coherence

tomography scans of the retina (1000 A-scans, 100 B-scans, 1.4 x 1.4 mm, Bioptigen Envisu R2200) were

acquired. Retinal layer thickness was measured using InVivoVue Diver 3.0.8 software (Bioptigen), at 16

locations equally spaced around the optic nerve head and averaged per mouse.

Optomotor response test

Visual performance was tested using the virtual reality optomotor test (Optomotry, Cerebral Mechanics),

as previously described [43]. Briefly, the mouse was place on a platform in the testing arena and, using

the staircase method, the maximal spatial frequency of moving black-white stripes for which the

optomotor reflex was still present, was determined.

 <u>Electroretinogram</u>

 Following overnight dark adaptation, electroretinograms (Celeris, Diagnosys) were recorded for

 anesthetized mice. Lens electrodes with integrated stimulators were placed on the cornea after pupil

 dilation (0.5% tropicamide, Tropicol, Théa and 15% phenylephrine, Théa). Eyes were alternately

 stimulated, and the electrode on the contralateral eye was used as a reference. Full-field flash

 electroretinogram was recorded at increasing flash intensities of 0.003, 0.01, 0.1, 1, 2.5 and 7.5 cd*s/m^2.

 For each intensity, 3 flashes were averaged. Inter-flash time increased from 5 s for the first two flash

 intensities, 10 s for the next two intensities to 30 and 45 s for the two last intensities. The scotopic

 threshold response was measured at $5*10^{-6}$ cd*s/m^2 and 50 flashes were averaged. Analysis was

 performed with Espion v6.59.9 software (Diagnosys), as previously described [43]. The scotopic threshold

 response amplitude was defined as the peak of the curve. Using a band pass filter (75-300 Hz), the

 oscillatory potentials on the rising part of the b-wave were retrieved and the area under the curve was

 calculated. The latency of the oscillatory potentials was defined for OP2 [43].

 <u>Hyperspectral imaging</u>

 A hyperspectral visible and near-infrared snapscan camera (150 bands, 470-900 nm, Imec; HSI Snapscan

 software (version 1.2.0)) was mounted on a Leica DM6 microscope with LED light source (400-700 nm).

 The snapscan camera acquires the hyperspectral image (2048 x 1536 pixels) by the movement of the

 linescan hyperspectral imaging sensor with on-chip filter technology on a miniaturized translation

 platform inside the camera body. Filters for each spectral band cover five adjacent pixel rows. The camera

 was used in stop-motion mode in steps of three pixel rows to reduce cross-talk between adjacent spectral

 bands and with HDR enabled (three frames) to reduce signal-to-noise ratio. Acquisition was set to cover

 only the 70 visible light spectral bands (471-691 nm). For hyperspectral imaging, retinal wholemounts

 were stained with DAPI, prior to mounting on a glass slide using a drop of saline and a cover slip. The

transmission spectra of central (1 mm from the optic nerve head) and peripheral (2 mm from the optic

nerve head) regions of each retinal quadrant were acquired using a 40x objective with the ganglion cell

layer nuclei used as the focal plane.

Spectral analysis was performed using a modification of published methods [13] (Supplementary fig. 1).

In brief, each frame of the recorded hyperspectral image was spatially resized to 20% of its original size to

smooth out acquisition noise. Each spectrum was normalized by the average spectral value of the three

longest wavelengths (685, 687 and 691 nm). Pixels with spectral values above 1.5 and below zero were

considered as outliers and masked out from further analysis. Spectral data from all mice were stacked in

a matrix and a corresponding class label (AD or wild type control [CO]) was given for each entry. The

between class covariance spectrum was calculated. This spectrum corresponds to the average spectral

difference between AD and CO groups. The hyperspectral score for a given spectral pixel was obtained by

calculating the dot product between its spectral data and the between class covariance spectrum. The

average spectral score for a given mouse was calculated using all pixels of all the HS images collected for

that retina.

Statistical analysis

Analyses were performed by a blinded observer. The employed statistical analyses and number of mice

are stipulated in the respective figure legend. Analyses were performed using Prism v.8.2.1 (GraphPad).

Differences were considered statistically significant for two-sided p-values < 0.05 (#, *p < 0.05; ##, **p <

0.01; ###, ***p < 0.001; ####, ****p < 0.0001; * for comparisons with baseline, # for comparisons

between genotype).

III.   Results

Amyloid accumulates in the retinas of App^{NL-G-F} mice, with Aβ plaque formation at old age

Retinal amyloid burden was studied in App^{NL-G-F} mice of various ages via ELISA. Soluble $Aβ_{42}$ was detected

at 3 months, with stable levels until 18 months, yet a steep increase at 24 months (p=0.0006) (Fig. 1a).

Insoluble $Aβ_{42}$, which is typically aggregated in amyloid plaques, was only clearly elevated at 12 months

and dramatically increased at 24 months (p<0.0001) (Fig. 1a). In contrast, and in line with the combined

effect of the Swedish mutation that increases total Aβ production and the Iberian mutation that increases

the $Aβ_{42}$/$Aβ_{40}$ ratio, a steady level of soluble $Aβ_{40}$ accumulation was present from 3 to 24 months, albeit

two orders of magnitude lower than $Aβ_{42}$ (Fig. 1b). Insoluble $Aβ_{40}$ (Fig. 1b) as well as soluble/insoluble $Aβ_{38}$

and $Aβ_{36}$ levels (data not shown) were below the detection limit of the assay. For comparison, in the

APP/PS1 overexpression mouse model, at 18 months of age (when retinal plaques are present [9, 12, 22,

33, 54]), we detected the accumulation of soluble $Aβ_{42}$ (comparable levels to App^{NL-G-F}), soluble $Aβ_{40}$

(higher levels than App^{NL-G-F}), and insoluble $Aβ_{42}$ (lower levels than App^{NL-G-F}) (Fig. 1c).

In accordance with these results, Aβ plaques were detected via 6E10 and 82E1 immunohistochemistry on

retinal wholemounts of App^{NL-G-F} mice of 18 months of age (Fig. 1d-i). Morphologically, these resembled

classical brain plaques, with a dense core and radiating fibrillary arms (Fig. 1k). Sizes typically ranged from

5 to 20 μm. Aβ plaque density was higher in the peripheral than in the central retina, while no overt

differences between the retinal quadrants were seen (Fig. 1j). Plaques mainly localized in the inner

plexiform layer (Fig. 1f), often closely associated with microglia (Fig. 1l & Fig. 4d-g), ChAT$^+$ strata with

cholinergic amacrine cell synapses (Fig. 1m), and PKCα$^+$ strata with bipolar cell terminals (Fig. 1n). In

addition, some were found around the blood vessels in the nerve fiber layer, in close association with

pericytes (Fig. 1o) and endothelial cells (Fig. 1p). In all these cases, the Aβ deposits were extracellularly.

Intracellular 6E10 staining was only seen in microglia (Fig. 1l), probably due to phagocytosis of Aβ plaques;

and in RBPMS[+] and melanopsin[+] ganglion cells (data not shown), likely reflecting endogenous APP

expression. Plaques did not associate with ganglion cells (RBPMS and melanopsin), bipolar (Chx10) and

amacrine (ChAT) cell somata, nor astrocytes and radial Müller glia (GFAP) (data not shown). Oligomeric

Aβ was visualized with an A11 immunostaining on retinal sections and was most abundant in all strata of

the inner plexiform layer – comprising amacrine and bipolar cell presynaptic and ganglion cell postsynaptic

terminals (Fig. 1q-u). Oligomeric Aβ was also observed extracellularly in the ganglion cell layer (Fig. 1t)

and intracellularly in Chx10[+] bipolar cells (Fig. 1u). A progressive accumulation over time – in 3- to 18- to

24-month-old mice –, increasingly affecting the inner plexiform layer and bipolar cells, was observed (Fig.

1q-s). No Aβ staining, with 6E10 nor A11, was observed in the outer retina.

These data show that, while soluble $Aβ_{42}$ (and $Aβ_{40}$) is present in the retinas of App^{NL-G-F} mice early in life,

retinal Aβ plaques only appear from 18 months of age onwards. Given this relatively slow progression of

amyloidopathy, this retinal mouse model may be used to study the conversion from early to late stages

of AD pathogenesis, as well as the effect of soluble Aβ on retinal structure and function. For the

characterization of the latter, in addition to longitudinal studies, we will largely focus on ages 3 and 18

months in the rest of this study, as these represent disease stages without and with Aβ plaques in the

retina, respectively.

<u>Hyperspectral imaging detects early retinal accumulation of Aβ in App^{NL-G-F} mice</u>

Aiming to provide a proof-of-concept for the use of hyperspectral imaging for the early detection of retinal

Aβ accumulation and for monitoring its progression, we next performed *ex vivo* hyperspectral imaging of

retinal wholemounts of 3- and 18-month-old App^{NL-G-F} mice and age-matched WT controls. For both ages,

the average spectra of the App^{NL-G-F} mice were clearly distinct from WT. In young mice, this separation was

most marked in the 475-500 nm range (Fig. 2a), while in old mice the spectral separation became even

more apparent, extending to 600 nm (Fig. 2b). A hyperspectral score was calculated for each mouse from

the *ex vivo* retinal reflectance data, using a modification of methods developed for *in vivo* hyperspectral

retinal imaging [13]. Hyperspectral scores were significantly higher in *App^{NL-G-F}* mice compared to age-

matched WT mice at both ages (p=0.0014 at 3 months, p=0.0002 at 18 months) (Fig. 2d-e). Furthermore,

the difference in the mean hyperspectral score of *App^{NL-G-F} versus* age-matched WT mice was also higher

at 18 months than at 3 months, in line with the progressive accumulation of Aβ seen in the ELISA.

Hyperspectral scores were calculated for each retinal quadrant in both central and peripheral locations

(Fig. 2j). In keeping with the immunohistochemistry findings, which demonstrated a predominately

peripheral distribution of Aβ, hyperspectral scores were higher in the peripheral retina than in the central

retina (Fig. 2g-h).

Remarkably, 18-month-old APP/PS1 mice had retinal hyperspectral signatures that closely mimics those

of 18-month-old *App^{NL-G-F}* mice (Fig. 2c, f, i). Given that these mice have similar levels of soluble Aβ$_{42}$, but

markedly different levels of soluble Aβ$_{40}$ and insoluble Aβ$_{42}$, it is possible that soluble Aβ$_{42}$ is most

contributory to the hyperspectral signature, as previously suggested by Vince and More [29].

<u>Neuronal dysfunction yet no degeneration in the inner retina of *App^{NL-G-F}* mice</u>

Many studies have shown neurotoxic effects of soluble Aβ, and that functional deficits may precede

neurodegeneration. Therefore, first electroretinography was performed with *App^{NL-G-F}* and WT mice at

different time points. Overall, no abnormalities were observed at early ages (Supplementary fig. 2). At 18

months, however, the latency of both the oscillatory potentials and the b-wave, a measure for amacrine

and bipolar cell function, respectively, showed significant changes (p<0.0001 for the effect of genotype)

in the *App^{NL-G-F}* mice relative to age-matched WT mice (Fig. 3d-g). Furthermore, also at 18 months of age,

the latency of the a-wave was affected at the lowest light intensity (p<0.0001), suggesting compromised

photoreceptor function (Fig. 3b-c).

Next, we assessed potential neurodegeneration in the *App*[NL-G-F] retina via a longitudinal *in vivo* optical

coherence tomography experiment. No differences in the overall thickness of the retina nor the nerve

fiber, ganglion cell, inner plexiform or inner nuclear layers were identified at any age (Fig. 3h). Of note, no

neurodegeneration was observed in the brains of the *App*[NL-G-F] mice either [40]. A small decrease in the

thickness of the outer nuclear layer was found in *App*[NL-G-F] mice, from 12 months (p=0.0016 for 12 months,

p<0.0001 for 18 months) of age onwards (Fig. 3i). Although these findings corroborate the

electroretinogram results, we are unsure how to interpret their biological relevance given that this

thinning is limited and occurs in the outer retina – where plaques nor oligomeric Aβ were seen. Loss or

dysfunction of the retinal pigment epithelium due to amyloidopathy in these support cells may underlie

this photoreceptor degeneration [6, 34], or, alternatively, the AD phenotype of the *App*[NL-G-F] mice may lead

to accelerated aging and therefore earlier age-related deterioration/loss of rods [21]. This should be

further investigated in follow-up studies.

Finally, based on the localization of plaques in the nerve fiber and inner plexiform layers, and in order to

confirm and complement the optical coherence tomography data, we next examined neuronal

subpopulations in the inner retinas of 18-month-old animals. We quantified: (i) ganglion cells, using the

pan-ganglion cell marker RBPMS (Fig. 3l); (ii) melanopsin[+] ganglion cells, given that this subtype appears

to be selectively vulnerable to Aβ pathology [31] (Fig. 3m); and (iii) ChAT[+] cholingeric amacrine cells (Fig.

3n). While a small loss of RBPMS[+] ganglion cells was observed (6.26 ± 2.28%, p=0.0158), melanopsin[+] and

ChAT[+] neuronal populations did not show any sign of degeneration in 18-month-old *App*[NL-G-F] mice. In line

with all of the above observations, and in keeping with the relatively minor loss of ganglion cells and neural

plasticity leading to reversible/inducible changes of retinal ganglion cell electrical activity during a critical

period of dysfunction preceding death [37], 18-month-old App^{NL-G-F} mice behaved normally in the

optomotor response test for visual acuity (Fig. 3o) and contrast sensitivity (data not shown).

Altogether, these results indicate that the majority of inner retinal neurons in the App^{NL-G-F} mouse survive

progressive amyloidosis up till 18 months. Nevertheless, although no cell death was observed at this age,

amacrine and bipolar cell function may be used as a sensitive read-out for AD pathology in the retina of

App^{NL-G-F} mice.

<u>Local microglia reactivity, astrogliosis and vascular changes in the retina of aged App^{NL-G-F} mice</u>

Because of the strong implication of microgliosis and astrogliosis in AD pathogenesis, and as neurotoxic

inflammation may precede neurodegeneration, we next examined the retinal glial cells. First, microgliosis

was investigated in entire Iba1-labeled retinal wholemounts of App^{NL-G-F} mice aged between 3 and 18

months. Overall quantification of microglial density (Fig. 4a), soma size (Fig. 4b) and roundness (data not

shown) did not reveal population-wide differences between the two genotypes at any age. Analysis of 18-

month-old App^{NL-G-F} x $CX_3CR-1^{GFP/+}$ mice retinas, with GFP-labeled microglia, confirmed these findings (data

not shown). However, a detailed investigation of these retinas, co-labeled for 6E10, revealed local

microglia reactivity around Aβ plaques, as was previously shown in the brains of these mice [40] and in

the retinas of other AD mouse models (3xTgAD, APP/PS1) [10, 36, 41]. Indeed, microglia clustered around

Aβ plaques, and these cells showed less ramified, thicker processes and larger somas (p=0.0112)

compared to microglia in regions without plaques (Fig. 4c-g) or in wild type $CX_3CR-1^{GFP/+}$ mice. Second,

Müller glia and astrocyte reactivity were assessed on transverse sections of the retina. Longitudinal

assessment of Müller glia reactivity, by counting the number of GFAP-positive radial fibers, revealed

higher reactivity in 18-month-old App^{NL-G-F} *versus* WT mice (p=0.0432) (Fig. 4h-j). Further substantiating

these findings, and in line with the brain phenotype [40], analysis of the S100B immunopositive area,

which is specific for astrocytes, showed astrogliosis in the retina of 18-month-old App^{NL-G-F} mice (p=0.0057)

(Fig. 4 k-m).

Looking at non-neuronal cell types, besides glial cells, also vasculature may undergo pathological

alterations in the context of AD. Changes in retinal blood vessel function and morphology, including

branching complexity, vessel density and vessel diameter, are being explored as potential early

biomarkers for AD [8, 38, 53, 55]. Given that a subset of plaques was found closely associated with the

retinal vasculature, we next assessed vessel morphology on retinal wholemounts stained with isolectin

B4 (Fig. 5). Vessel density (Fig. 5a) and branching complexity (Fig. 5b) were unaltered in App^{NL-G-F} *versus*

WT mice at 18 months of age. However, diameters of the retinal venules (Fig. 5c), yet not of the retinal

arterioles (Fig. 5d), were smaller in App^{NL-G-F} mice (p=0.0235). Of note, Shi *et al.* recently reported pericyte

loss accompanying retinal vascular amyloidosis, hence the observed reduction in diameter may be a

manifestation of this [44].

IV.   Discussion

In this study, we investigated the extent to which AD-associated disease processes are present in the

retina of the App^{NL-G-F} knock-in mouse, and whether they can be monitored via *in vivo* retinal imaging and

electrophysiology. First, we demonstrated that soluble Aβ (mostly Aβ$_{42}$) accumulation is present early in

life in these mice yet retinal Aβ plaques are only observable at 12 months of age and older. This rising Aβ

burden in the retina coincides with local microglia reactivity, astrogliosis, and abnormalities in retinal vein

morphology, and leads to compromised electrophysiological function of selected retinal neurons. No

overt neurodegeneration was seen, however, and visual performance outcomes were intact. Altogether,

these findings suggest that the App^{NL-G-F} retina resemble changes that occur in the brain in preclinical AD,

during which Aβ accumulation instigates the earliest detrimental effects on astrocytes, microglia and

vasculature, poses a stress on neuronal functionality but does not lead to overt functional impairment

and neuronal death yet. Second, the data presented in this study suggest that there is an identifiable pre-

plaque stage in AD in which only soluble forms of Aβ are present, followed by a phase in which soluble,

oligomeric forms of Aβ are likely in equilibrium with plaques. The *App^{NL-G-F}* retina and reported retinal

imaging techniques offer unique opportunities for further fundamental and preclinical research into this

sequence of biological events over the course of preclinical AD. Hence, the *App^{NL-G-F}* mouse is not only the

first AD model to accumulate Aβ without phenotypes related to APP overexpression and to more closely

model human pathophysiology, its retina also is the first model organ to study preclinical AD. Combined

with the available non-invasive, high-resolution imaging techniques and functional assessments of the

retina, this creates a toolbox for researchers to shed new light on Aβ proteostasis and novel disease-

modifying therapies focused thereon.

The rather mild phenotype is in stark contrast to observations made in the brains of *App^{NL-G-F}* mice. Indeed,

previous studies have shown that (sub)cortical Aβ plaques appear at 2-4 months of age, with astrogliosis

and microglial reactivity at 6-9 months, and cognitive impairment as early as 6 months [40]. The later

formation of Aβ plaques in the *App^{NL-G-F}* retina could be due to lower expression levels of Aβ compared to

the brain, as shown for other AD mouse models [1], or to different clearance rates. However, importantly,

it confers a unique strength to the *App^{NL-G-F}* retina as a model for AD research, as it creates a time window

to investigate the earliest phases of AD pathogenesis and is closer to the slow disease progression seen in

humans. A second asset of the *App^{NL-G-F}* retina is that, by diverging from many other available AD models,

its phenotype shows striking similarities to the neuropathological changes seen in retinas of human AD

patients. This includes the shape and size of the retinal plaques, which we found to be 5-10 times smaller

than their counterparts in the brain [22, 23]; the localization of these plaques to the inner retinal layers –

where endogenous APP expression has been observed [17, 25] – and associated with retinal blood vessels;

as well as the microglial reactivity, astrocytosis and changes in retinal vessel morphology. Of note, the

manifestations of retinal vascular amyloidosis are of particular interest in light of the recent findings by

Shi *et al.,* who reported pericyte loss accompanying retinal vascular amyloidosis in the human AD retina

[44]. With one exception [14], this phenotype is thus far unstudied in AD mouse models and our data

suggest that the *App*$^{NL-G-F}$ retina can be used to further explore this retinal manifestation of AD.

Furthermore, the *App*$^{NL-G-F}$ retina also appears to be a research model for in-depth investigation of the

synaptotoxic effects of Aβ, given the converging data on (soluble) Aβ accumulation in synapses in the

inner plexiform layer and electrophysiological abnormalities of amacrine and bipolar cells. Notably, an

explanation for the diverging and more severe phenotype of many other transgenic AD mouse models

may be sought in the fact that those models are far removed from the human disease state, with non-

endogenous promotors, overexpression of APP byproducts, PS1 mutation and/or abnormal APP and Aβ

expression levels. This study is pioneering in that it is the first-time retinal phenotyping study of the *App*$^{NL-}$

$^{G-F}$ knock-in mouse, which was created to overcome all of the aforementioned artefacts and currently still

is unique in its kind. Finally, we have also showcased here the unique potential of the retina for *in vivo*

monitoring of structural and functional changes in AD. Of the battery of tests used, hyperspectral imaging

and electroretinograms proved to be sensitive read-outs to assess AD disease status. Results from our

post mortem studies suggest that these may in future studies be complemented with confocal scanning

ophthalmoscopy for microgliosis and optical coherence tomography angiography or fundus imaging for

detailed retinal vascular assessments. Together, this creates a set of *in vivo* read-outs for preclinical

investigation that cannot be matched by any existing *in vivo* brain research methods. Notably, the fact

that identical techniques and outcome measures can be used in mice and patients, confers potential

translational value to this research.

A better understanding of the time course of the pathological processes underlying preclinical AD, is an

important prerequisite for the development of new therapies targeting this phase. Furthermore, retinal

imaging may provide an accurate and convenient indication of the efficacy of therapies. Currently, Aβ

status is evaluated via CSF assays or PET imaging, which is mainly indicative for the deposition of fibrillary

forms of Aβ. As such, these do not always provide information about oligomeric Aβ, which may be the

more relevant Aβ species to target [46, 49]. Our data substantiate that hyperspectral imaging is a highly

sensitive technique that can be used to quantify Aβ burden in the earliest stages of disease. This imaging

spectroscopy technique is based on the fact that soluble Aβ aggregates – most likely $Aβ_{42}$ – interfere with

traversing light, leading to Rayleigh light scattering and a characteristic hyperspectral signature, the

magnitude of which appear to be proportional to the amount of Aβ present [29, 30]. In the past year,

following promising *post mortem* and *in vivo* studies in both animal and human retinas [28–30], results of

the first clinical trial showing that *in vivo* retinal hyperspectral imaging can discriminate between Aβ PET-

positive cases and controls emerged [13]. This, together with our data, suggests that hyperspectral

imaging can be used to monitor retinal Aβ accumulation, with the potential to provide quantitative

measures of Aβ oligomers. As such, hyperspectral imaging in the App^{NL-G-F} retina may become a gateway

to improved preclinical studies of the new generation of Aβ therapeutics. Finally, on the long run, this

research may also lay the foundations for the rational use of this retinal imaging biomarker for AD

diagnosis, disease monitoring, patient stratification and even population-wide screenings.

V.   Conclusion

To conclude, decades of AD research have taught us that we should not try to unravel AD pathogenesis

from a simple, neuron-centric perspective, or by assuming a linear series of events leading from Aβ

accumulation to dementia. AD is to be understood as a complex pathological cascade, taking decades to

gradually evolve from a clinically silent phase to dementia, that is driven by the proteopathic stress

imposed by Aβ and Tau conformations, and feedback and feedforward responses of astrocytes, microglia,

and vascular cells [48]. Research based on this new framework requires novel approaches, including

mouse models that more closely mimic human pathophysiology and allow the study of the earliest,

preclinical phases of AD. In this research, we have shown that the retina of App^{NL-G-F} mice is a promising

model for preclinical AD, in which Aβ oligomers accumulate and pathological cellular responses are

elicited. Furthermore, we have added further evidence in support of the potential of hyperspectral

imaging to quantify retinal Aβ burden and to serve as an AD biomarker. Altogether, the App^{NL-G-F} retina

and retinal imaging techniques are unique tools for both fundamental and drug discovery research in the

preclinical phase of AD.

VI.   List of abbreviations

Alzheimer's disease (AD), amyloid-beta (Aβ), central nervous system (CNS), choline acetyltransferase

(ChAT), glial acidic fibrillary protein (GFAP), green fluorescent protein (GFP), hyperspectral imaging (HSI),

ionized calcium-binding adapter molecule 1 (Iba1), paraformaldehyde (PFA), RNA-binding protein with

multiple splicing (RBPMS), S100 calcium binding protein B (S100B), wild type (WT)

VII.   Declarations

<u>Ethics approval and consent to participate</u>

All animal experiments were performed according to the European directive 2010/63/EU and approved

by the KU Leuven institutional ethics committee for animal research.

<u>Consent for publication</u>

Not applicable

Availability of data and material

The data sets generated and analyzed during the current study are available from the corresponding

author upon reasonable request.

Competing interests

The authors declare the following competing interests: XH and PvW have filed an International (PCT)

Patent Application No PCT/AU2019/000003 relating to retinal hyperspectral imaging. They are co-

founders of Enlighten Imaging PTY LTD, a start-up company focused on developing novel retinal imaging

solutions for neurological and retinal diseases.

Funding

MV, LV, LuM and LDG are fellows of the Research Foundation Flanders (FWO). This research was

supported by the Alzheimer's Research Foundation (SAO-FRA), the European Union Horizon 2020

Research and Innovation Program (2014-2020) (HERALD project, granted by the ATTRACT consortium).

The Centre for Eye Research Australia receives Operational Infrastructure Support from the Victorian

Government. XH and PvW acknowledge funding support from the H & L Hecht Trust and the Yulgilbar

Alzheimer's Research Program.

Authors' contributions

MV designed the study, performed experiments, analyzed the data and wrote the manuscript; LV, SL, GG

performed experiments; XH, LS, JT, LuM analyzed the data; TS and TCS created the App^{NL-G-F} mice; SL, JT,

MJ and LDG implemented the hyperspectral imaging set-up; PDB, IS, BDS, PvW supervised the study and

reviewed the manuscript; LM supervised and designed the study, analyzed the data and wrote the

manuscript, LDG supervised and designed the study, performed experiments, analyzed the data and wrote

the manuscript.

<u>Acknowledgements</u>

We thank Véronique Brouwers and Evelien Herinckx for their assistance with mouse breeding and

experimentation; Wouter Charle, Toon Van Craenendonck and Arnout Standaert for advice on

hyperspectral imaging; Isabelle Etienne and Tine Van Bergen (Oxurion, Heverlee, Belgium) for help with

immunohistochemistry; RIKEN BioResource Center for creation and distribution of the *App^{NL-G-}*mice.

VIII.   Figures

Figure 1. Amyloid burden increases with age in the retina of *App^NL-G-F* mice. (a-b) Amyloid ELISA on retinal

lysates of *App^NL-G-F* mice shows accumulation of soluble $A\beta_{42}$ from 3 months of age, which rises up till 24

months. Levels of insoluble $A\beta_{42}$ only start to rise at 12 months. Accumulation of soluble, but not insoluble,

$A\beta_{40}$ is seen at all ages yet does not progress with age. One-way ANOVA with Dunnett's multiple

comparisons test ($F_{5,28}$=6.686, p=0.0006 for soluble $A\beta_{42}$; $F_{5,27}$=20.27, p<0.0001 for insoluble $A\beta_{42}$); n=5-6.

(c) Eighteen-month-old APP/PS1 mice have comparable soluble $A\beta_{42}$ levels yet lower levels of insoluble

$A\beta_{42}$ and higher soluble $A\beta_{40}$ than *App^NL-G-F* mice of the same age. n=5. Of note, human $A\beta$ levels are

undetectable in the retinas of WT mice and therefore these controls are not shown. (d-i) $A\beta$

immunostainings with antibodies 82E1 (d-f) and 6E10 (g-i) on retinal wholemounts of 18-month-old WT

(d, g) and *App^NL-G-F* mice (e, f, h, i) reveal the presence of $A\beta$ plaques in *App^NL-G-F* mice only. (f) Example of

a reconstruction of the z-plane of the image in panel e, with 82E1 and DAPI nuclear counterstaining,

showing that $A\beta$ plaques mainly localize to the inner plexiform layer. (j) Representative example of a

retinal wholemount on which 6E10-positive $A\beta$ plaques are depicted with dots to illustrate plaque

distribution in the retina. (k) $A\beta$ plaques in the cerebral cortex of an 18-month-old *App^NL-G-F* mouse. (l-p)

Double labeling for $A\beta$ (6E10) and retinal cell markers on retinal wholemounts of 18-month-old *App^NL-G-F*

mice. Single-tile images of the retinal flatmounts, with corresponding orthogonal projections of the z-

stack (') are shown. Dotted lines in m' and n' indicate the z depth for with the single-tile image is shown.

Dotted lines in o and p indicate the orientation of the orthogonal projections shown in o' and p'. (l) Double

labeling for $A\beta$ and Iba1 shows both extra- and intracellular $A\beta$ deposits in microglia, suggesting that

microglia surround and phagocytose plaques. (m-n) Most plaques are found in the inner plexiform layer,

where they align with the ChAT$^+$ strata in which the terminals of the cholinergic amacrine cells are

organized (m) and the PKCα^+ sublayer in which the terminals of the bipolar cells are organized (n). (o-p)

Plaques are also found in the uppermost layers of the retina, in association with the pericytes (m) and

endothelial cells (n) of the retinal blood vessels. (q-s) Immunostaining for oligomeric amyloid (A11) on

retinal cross-sections of 3-month- (3M) (q), 18-month- (18M) (r) and 24-month-old (24M) App^{NL-G-F} mice

(s) shows progressive accumulation of Aβ oligomers. (t, u) These Aβ oligomers are mostly found in the

inner plexiform layer, in association with ganglion cells (Brn3a$^+$) and displaced amacrine cells (Brn3a$^-$) in

the ganglion cell layer (t), and inside Chx10$^+$ bipolar cells (u). (v) Schematic overview of the retina and the

locations in which Aβ oligomers and plaques were found. Scalebars: 10 µm (m'-p'), 20 µm (d-i, k, n, q-u),

50 µm (p), 100 µm (l, m, o), 1 mm (j). Data are shown as mean $\pm$ SEM. GCL = ganglion cell layer, IPL = inner

plexiform layer, INL = inner nuclear layer, OPL = outer plexiform layer, ONL = outer nuclear layer, G =

ganglion cell, A = amacrine cell, B = bipolar cell, V = blood vessel, M = microglia, C = cone, R = rod.

Figure 2. Quantification of retinal Aβ burden via hyperspectral imaging reveals progressive retinal amyloidosis in *App^NL-G-F* mice. (a) Average spectra of 3-month-old *App^NL-G-F* and WT mice are different, with a clear separation in the short wavelength range. (b) Average spectra of 18-month-old *App^NL-G-F* and WT mice show significant spectral differences for all wavelengths up to 600 nm. (c) Average spectra of 18-month-old APP/PS1 and WT mice show significant spectral differences from ~500 to ~600 nm. Data are shown as mean ± 95% confidence intervals. (d-e) For both ages, comparison of hyperspectral (HS) scores

(obtained by compression of the spectral information to maximize between group separation) of *App*^{NL-G-}

F and WT mice shows a strong effect size and significant difference between genotypes. Unpaired two-

tailed t-tests (t_{14}=3.957,p=0.0014 at 3 months; t_{16}=4.919,p=0.0002 at 18 months); n=8-10. (f) HS scores of

18-month-old APP/PS1 and WT mice are significantly different. Unpaired two-tailed t-tests

(t_{10}=2.395,p=0.0376); n=4-8. (g-i) Heatmaps with a spatial representation of the HS score per image for

each mouse. Note that image fields 2, 4, 6 and 8, which correspond to the peripheral retinal locations,

have higher scores, indicative for a higher amyloid load. (j) Schematic representation of the retinal imaging

locations.

Figure 3. Neuronal dysfunction yet no degeneration in the retina of *App^{NL-G-F}* mice. (a) Representative

electroretinogram measurements, with increasing light intensities, in 18-month-old *App^{NL-G-F}* (green) and

WT (blue) mice. (b) The amplitude of the a-wave is unaffected in *App^{NL-G-F}* mice of 18 months, yet (c) the

latency time is decreased for the lowest light intensity. Two-way ANOVA with Sidak's multiple comparisons test ($F_{1,85}$ = 23.14 for the effect of genotype, p<0.0001); n=8-10. (d) Oscillatory potentials show a similar wave front (measured as the area under the curve, AUC), but (e) a reduced latency in 18-month-old App^{NL-G-F} mice, compared to WT. Two-way ANOVA with Sidak's multiple comparisons test ($F_{1,80}$ = 37.14 for the effect of genotype, p=0.0013, 0.0124, 0.0378 for 0.01, 0.1 and 1 cd*s/m^2 respectively); n=8-10. (f) Similarly, the amplitude of the b-wave is unaltered, but (g) the latency time is decreased in 18-month-old App^{NL-G-F} mice compared to WT mice. Two-way ANOVA with Sidak's multiple comparisons test ($F_{1,97}$ = 27.90 for the effect of genotype, p=0.0206 and 0.044 for 0.1 and 1 cd*s/m^2); n=8-10. Full electroretinogram data is given in Supplementary figure 2. (h) Analysis of the thickness of the retinal layers, as measured by optical coherence tomography, shows no differences between App^{NL-G-F} and WT mice from 2 to 18 months of age, (i) with the exception of a thinning of the outer nuclear layer. Two-way ANOVA with Sidak's multiple comparisons test ($F_{1,106}$ = 32.14 for the effect of genotype, p=0.0016 and p<0.0001 for 12 and 18 months, respectively); n=8-9. (j-k) Representative optical coherence tomography images of WT (j) and App^{NL-G-F} (k) retinas. Dotted lines delineate the outer nuclear layer. (l) Quantification of RBPMS$^+$ retinal ganglion cell density reveals ganglion cell loss in 18-month-old App^{NL-G-F} compared to WT mice. Unpaired two-tailed t-test (t_{14}=2.745,p=0.0158); n=8. (m-n) Cell counts of melanopsin$^+$ ganglion cells (m) and ChAT$^+$ neurons (n) show no differences in numbers between 18-month-old App^{NL-G-F} and WT mice. (o) A longitudinal study of visual acuity in App^{NL-G-F} mice from 3 till 18 months revealed a decrease with age but no genotype differences. Two-way ANOVA ($F_{10,180}$=21.11 for the effect of age, $F_{1,180}$=0.1149 for the effect of genotype); n=9. Scale bar: 25 µm. Data are depicted as mean ± SEM; RNFL = retinal nerve fiber layer, GCL = ganglion cell layer, IPL = inner plexiform layer, INL = inner nuclear layer, OPL = outer plexiform layer, ONL = outer nuclear layer, PL = photoreceptor layer.

Figure 4. Locally reactivated microglia and macrogliosis in the retinas of aged *App^NL-G-F* mice. (a-b)

Quantification of microglial density (a) and soma area (b) on Iba1-labeled retinal wholemounts of 3- to

18-month-old *App^NL-G-F* and WT mice reveals no differences between genotypes when analyzed for the

entire retina. An effect of aging was observed for microglia density (Two-way ANOVA, $F_{4,33}$ = 2.976,

p=0.0334); n=3-6. (c) Morphometric analysis of plaque-associated microglia *versus* microglia distant from

Aβ plaques in 18-month-old *App^NL-G-F* x *CX₃CR-1^GFP/+* retinas reveals that plaque-associated microglia have

a larger soma, which is indicative of their activation. Unpaired two-tailed t-test (t_{63}=2.614, p=0.0112);

n=25-40 cells from 6 mice. (d-g) Representative images of immunostaining for Aβ (6E10) on retinal

wholemounts of 18-month-old *App*$^{NL-G-F}$ x *CX₃CR-1*$^{GFP/+}$ (f-g) and *CX₃CR-1*$^{GFP/+}$ mice (d-e), illustrating that

green fluorescent microglia surrounding Aβ plaques display morphological alterations typical for reactive

microglia, with thicker and less ramified processes and a larger soma. (h) Counting the number of GFAP$^+$

radial fibers on retinal sections of *App*$^{NL-G-F}$ and WT mice shows that macrogliosis increases with age, and

significantly differs between genotypes at 18 months. Two-way ANOVA with Sidak's multiple comparisons

test ($F_{4,50}$=34.02 for the effect of age, $F_{1,50}$=4.411 for the effect of genotype, p=0.0432); n=5-8. (i-j)

Representative images of a GFAP immunostaining on retinal cross-sections of 18-month-old WT (i) and

App^{NL-G-F} (j) mice. (k) Quantification of the S100B immunopositive area on retinal sections of 18-month-

old *App*$^{NL-G-F}$ and WT mice shows astrogliosis in the *App*$^{NL-G-F}$ retina. Unpaired two-tailed t-test (t_{12}=3.358,

p=0.0057); n=7. (l-m) Representative images of the S100B immunostaining on WT (l) and *App*$^{NL-G-F}$ (m)

retinas. Scalebar (d-g): 20 µm, scalebar (i, j, l, m): 50 µm. Data are shown as mean ± SEM. GCL = ganglion

cell layer, IPL = inner plexiform layer, INL = inner nuclear layer, OPL = outer plexiform layer, ONL = outer

nuclear layer.

Figure 5. Retinal venules of *App^NL-G-F* mice have reduced diameters. (a-b) Retinal vasculature was analyzed at 3 and 18 months of age. Quantification of the density of the vascular network (a) and its branching complexity (b) shows no differences according to genotype. (c-d) (c) Retinal venular diameters are reduced in 18-month-old *App^NL-G-F* mice. Two-way ANOVA with Sidak's multiple comparison's test ($F_{1,25}$=3.807 for the effect of genotype, p=0.0235); n=7 for 3 to 6 months, n=7-8 for 18 months. (d) No differences are seen in retinal arteriolar diameters. (e-f) Representative images of isolectin B4-labeled retinal flatmounts of 18-month-old WT (e) and *App^NL-G-F* (f) mice. (e'-f') Magnification of the region of interest in which vessel analysis was performed. Arrows indicate retinal venules with a reduced diameter. Scalebar: 1000 μm. Data are depicted as mean ± SEM. A = artery, V = vein.

IX.   Additional files

Supplementary figure 1. Schematic overview of the hyperspectral imaging data analysis. Spectral data

from all mice were stacked in a matrix and a corresponding class label (AD or wild type control [CO]) was

given for each entry. The between class covariance spectrum was calculated. This spectrum corresponds

to the average spectral difference between AD and CO groups. The hyperspectral score for a given spectral

pixel was obtained by calculating the dot product between its spectral data and the between class

covariance spectrum. The average spectral score for a given mouse was calculated using all pixels of all

the HS images collected for that retina.

Supplementary figure 2. Electroretinogram measurements in AppNL-G-F mice. Electroretinogram

measurements were carried out in both AppNL-G-F and WT mice at 3,6,9,12 and 18 months of age. Data

are depicted as mean ± SEM; n=4-12 per group. (pdf file)

X.   References

1.   Alexandrov PN, Pogue A, Bhattacharjee S, Lukiw WJ (2011) Retinal amyloid peptides and

complement factor H in transgenic models of Alzheimer's disease. Neuroreport. doi:

10.1097/WNR.0b013e3283497334

2.   Cline EN, Bicca MA, Viola KL, Klein WL (2018) The Amyloid-β Oligomer Hypothesis: Beginning of

the Third Decade. J. Alzheimer's Dis. 64:S567–S610

3.   Cordeiro MF (2016) Eyeing the brain. Acta Neuropathol. 132:765–766

4.   Criscuolo C, Cerri E, Fabiani C, Capsoni S, Cattaneo A, Domenici L (2018) The retina as a window

to early dysfunctions of Alzheimer's disease following studies with a 5xFAD mouse model.

Neurobiol Aging 67:181–188. doi: 10.1016/j.neurobiolaging.2018.03.017

5.   Davis BM, Salinas-Navarro M, Cordeiro MF, Moons L, Groef L De (2017) Characterizing microglia

activation: A spatial statistics approach to maximize information extraction. Sci Rep 7:1–12. doi:

10.1038/s41598-017-01747-8

6.   Dong ZZ, Li J, Gan YF, Sun XR, Leng YX, Ge J (2018) Amyloid beta deposition related retinal

pigment epithelium cell impairment and subretinal microglia activation in aged APPswePS1

transgenic mice. Int J Ophthalmol 11:747–755. doi: 10.18240/ijo.2018.05.06

7.   Fantini J, Chahinian H, Yahi N (2020) Progress toward Alzheimer's disease treatment: Leveraging

the Achilles' heel of Aβ oligomers ? Protein Sci.

8.   Feke GT, Hyman BT, Stern RA, Pasquale LR (2015) Retinal blood flow in mild cognitive impairment

and Alzheimer's disease. Alzheimer's Dement Diagnosis, Assess Dis Monit. doi:

10.1016/j.dadm.2015.01.004

9.   Georgevsky D, Retsas S, Raoufi N, Shimoni O, Golzan SM (2019) A longitudinal assessment of

retinal function and structure in the APP/PS1 transgenic mouse model of Alzheimer's disease.

Transl Neurodegener. doi: 10.1186/s40035-019-0170-z

10. Grimaldi A, Brighi C, Peruzzi G, Ragozzino D, Bonanni V, Limatola C, Ruocco G, Di Angelantonio S

(2018) Inflammation, neurodegeneration and protein aggregation in the retina as ocular

biomarkers for Alzheimer's disease in the 3xTg-AD mouse model. Cell Death Dis 9:1–10. doi:

10.1038/s41419-018-0740-5

11. De Groef L, Cordeiro MF (2018) Is the Eye an Extension of the Brain in Central Nervous System

Disease? J Ocul Pharmacol Ther 34:129–133. doi: 10.1089/jop.2016.0180

12. Gupta VK, Chitranshi N, Gupta VB, Golzan M, Dheer Y, Wall R Vander, Georgevsky D, King AE,

Vickers JC, Chung R, Graham S (2016) Amyloid β accumulation and inner retinal degenerative

changes in Alzheimer's disease transgenic mouse. Neurosci Lett. doi:

10.1016/j.neulet.2016.04.059

13. Hadoux X, Hui F, Lim JKH, Masters CL, Pébay A, Chevalier S, Ha J, Loi S, Fowler CJ, Rowe C,

Villemagne VL, Taylor EN, Fluke C, Soucy JP, Lesage F, Sylvestre JP, Rosa-Neto P, Mathotaarachchi

S, Gauthier S, Nasreddine ZS, Arbour JD, Rhéaume MA, Beaulieu S, Dirani M, Nguyen CTO, Bui B

V., Williamson R, Crowston JG, van Wijngaarden P (2019) Non-invasive in vivo hyperspectral

imaging of the retina for potential biomarker use in Alzheimer's disease. Nat Commun 10:1–12.

doi: 10.1038/s41467-019-12242-1

14.   Harper DJ, Augustin M, Lichtenegger A, Gesperger J, Himmel T, Muck M, Merkle CW, Eugui P,

Kummer S, Woehrer A, Glösmann M, Baumann B (2020) Retinal analysis of a mouse model of

Alzheimer's disease with multicontrast optical coherence tomography. Neurophotonics 7:1. doi:

10.1117/1.nph.7.1.015006

15.   Harris SS, Wolf F, De Strooper B, Busche MA (2020) Tipping the Scales: Peptide-Dependent

Dysregulation of Neural Circuit Dynamics in Alzheimer's Disease. Neuron. doi:

10.1016/j.neuron.2020.06.005

16.   Hart NJ, Koronyo Y, Black KL, Koronyo-Hamaoui M (2016) Ocular indicators of Alzheimer's:

exploring disease in the retina. Acta Neuropathol. 132:767–787

17.   Ho T, Vessey KA, Cappai R, Dinet V, Mascarelli F, Ciccotosto GD, Fletcher EL (2012) Amyloid

precursor protein is required for normal function of the rod and cone pathways in the mouse

retina. PLoS One. doi: 10.1371/journal.pone.0029892

18.   Van Hove I, De Groef L, Boeckx B, Modave E, Tjing-Tjing H, Beets K, Etienne K, Van Bergen T,

Lambrechts D, Moons L, Feyen JH, Porcu M Single-cell transcriptome analysis of akimba retina

reveals cell type-specific insights into the pathobiology of diabetic retinopathy. Diabetologia

19.   Jack CR, Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, Holtzman DM, Jagust W,

Jessen F, Karlawish J, Liu E, Molinuevo JL, Montine T, Phelps C, Rankin KP, Rowe CC, Scheltens P,

Siemers E, Snyder HM, Sperling R, Elliott C, Masliah E, Ryan L, Silverberg N (2018) NIA-AA

Research Framework: Toward a biological definition of Alzheimer's disease. Alzheimer's Dement.

14:535–562

20.   Jack CR, Lowe VJ, Weigand SD, Wiste HJ, Senjem ML, Knopman DS, Shiung MM, Gunter JL, Boeve

BF, Kemp BJ, Weiner M, Petersen RC (2009) Serial PIB and MRI in normal, mild cognitive

impairment and Alzheimers disease: Implications for sequence of pathological events in

Alzheimers disease. Brain. doi: 10.1093/brain/awp062

21.   Kolesnikov A V., Fan J, Crouch RK, Kefalov VJ (2010) Age-related deterioration of rod vision in

mice. J Neurosci 30:11222–11231. doi: 10.1523/JNEUROSCI.4239-09.2010

22.   Koronyo-Hamaoui M, Koronyo Y, Ljubimov A V., Miller CA, Ko MHK, Black KL, Schwartz M, Farkas

DL (2011) Identification of amyloid plaques in retinas from Alzheimer's patients and noninvasive

in vivo optical imaging of retinal plaques in a mouse model. Neuroimage 54:S204. doi:

10.1016/j.neuroimage.2010.06.020

23.   Koronyo Y, Biggs D, Barron E, Boyer DS, Pearlman JA, Au WJ, Kile SJ, Blanco A, Fuchs DT, Ashfaq A,

Frautschy S, Cole GM, Miller CA, Hinton DR, Verdooner SR, Black KL, Koronyo-Hamaoui M (2017)

Retinal amyloid pathology and proof-of-concept imaging trial in Alzheimer's disease. JCI insight 2.

doi: 10.1172/jci.insight.93621

24.   Lim JKH, Li QX, He Z, Vingrys AJ, Wong VHY, Currier N, Mullen J, Bui B V., Nguyen CTO (2016) The

eye as a biomarker for Alzheimer's disease. Front. Neurosci. 10:536

25.   Loffler KU, Edward DP, Tso MOM (1995) Immunoreactivity against tau, amyloid precursor

protein, and beta-amyloid in the human retina. Investig Ophthalmol Vis Sci

26.   London A, Benhar I, Schwartz M (2013) The retina as a window to the brain - From eye research

to CNS disorders. Nat. Rev. Neurol. 9:44–53

27.   Masin L, Claes M, Bergmans S, Cools L, Andries L, Davis BM, Moons L, De Groef L (2020) RGCode:

a novel Retinal Ganglion Cell quantification tool based On DEep learning. Sci Rep (in press)

28.   More SS, Beach JM, McClelland C, Mokhtarzadeh A, Vince R (2019) In Vivo Assessment of Retinal

Biomarkers by Hyperspectral Imaging: Early Detection of Alzheimer's Disease. ACS Chem

Neurosci 10:4492–4501. doi: 10.1021/acschemneuro.9b00331

29. More SS, Beach JM, Vince R (2016) Early detection of amyloidopathy in Alzheimer's mice by

hyperspectral endoscopy. Investig Ophthalmol Vis Sci 57:3231–3238. doi: 10.1167/iovs.15-17406

30. More SS, Vince R (2015) Hyperspectral imaging signatures detect amyloidopathy in alzheimers

mouse retina well before onset of cognitive decline. ACS Chem Neurosci 6:306–315. doi:

10.1021/cn500242z

31. La Morgia C, Ross-Cisneros FN, Koronyo Y, Hannibal J, Gallassi R, Cantalupo G, Sambati L, Pan BX,

Tozer KR, Barboni P, Provini F, Avanzini P, Carbonelli M, Pelosi A, Chui H, Liguori R, Baruzzi A,

Koronyo-Hamaoui M, Sadun AA, Carelli V (2016) Melanopsin retinal ganglion cell loss in

Alzheimer disease. Ann Neurol 79:90–109. doi: 10.1002/ana.24548

32. Nilsson P, Saito T, Saido TC (2014) New mouse model of Alzheimer's. ACS Chem. Neurosci.

33. Ning A, Cui J, To E, Ashe KH, Matsubara J (2008) Amyloid-β deposits lead to retinal degeneration

in a mouse model of Alzheimer disease. Investig Ophthalmol Vis Sci. doi: 10.1167/iovs.08-1849

34. Park SW, Kim JH, Mook-Jung I, Kim KW, Park WJ, Park KH, Kim JH (2014) Intracellular amyloid

beta alters the tight junction of retinal pigment epithelium in 5XFAD mice. Neurobiol Aging

35:2013–2020. doi: 10.1016/j.neurobiolaging.2014.03.008

35. Parnell M, Guo L, Abdi M, Cordeiro MF (2012) Ocular manifestations of Alzheimer's disease in

animal models. Int. J. Alzheimers. Dis.

36. Perez SE, Lumayag S, Kovacs B, Mufson EJ, Xu S (2009) β-amyloid deposition and functional

impairment in the retina of the APPswe/PS1ΔE9 transgenic mouse model of Alzheimer's disease.

Investig Ophthalmol Vis Sci. doi: 10.1167/iovs.08-2384

37. Porciatti V, Ventura LM (2012) Retinal ganglion cell functional plasticity and optic neuropathy: A

comprehensive model. J. Neuro-Ophthalmology 32:354–358

38.   Querques G, Borrelli E, Sacconi R, De Vitis L, Leocani L, Santangelo R, Magnani G, Comi G,

730   Bandello F (2019) Functional and morphological changes of the retinal vessels in Alzheimer's

731   disease and mild cognitive impairment. Sci Rep. doi: 10.1038/s41598-018-37271-6

732   39.   Reiman EM, Quiroz YT, Fleisher AS, Chen K, Velez-Pardo C, Jimenez-Del-Rio M, Fagan AM, Shah

733   AR, Alvarez S, Arbelaez A, Giraldo M, Acosta-Baena N, Sperling RA, Dickerson B, Stern CE, Tirado

734   V, Munoz C, Reiman RA, Huentelman MJ, Alexander GE, Langbaum JBS, Kosik KS, Tariot PN,

735   Lopera F (2012) Brain imaging and fluid biomarker analysis in young adults at genetic risk for

736   autosomal dominant Alzheimer's disease in the presenilin 1 E280A kindred: A case-control study.

737   Lancet Neurol 11:1048–1056. doi: 10.1016/S1474-4422(12)70228-4

738   40.   Saito T, Matsuba Y, Mihira N, Takano J, Nilsson P, Itohara S, Iwata N, Saido TC (2014) Single App

739   knock-in mouse models of Alzheimer's disease. Nat Neurosci. doi: 10.1038/nn.3697

740   41.   Salobrar-García E, Rodrigues-Neves AC, Ramírez AI, de Hoz R, Fernández-Albarral JA, López-

741   Cuenca I, Ramírez JM, Ambrósio AF, Salazar JJ (2020) Microglial activation in the retina of a triple-

742   transgenic alzheimer's disease mouse model (3xTg-AD). Int J Mol Sci. doi: 10.3390/ijms21030816

743   42.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

744   Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A

745   (2012) Fiji: An open-source platform for biological-image analysis. Nat. Methods 9:676–682

746   43.   Sergeys J, Etienne I, Van Hove I, Lefevere E, Stalmans I, Feyen JHM, Moons L, Van Bergen T (2019)

747   Longitudinal in vivo characterization of the streptozotocin-induced diabetic mouse model: Focus

748   on early inner retinal responses. Investig Ophthalmol Vis Sci. doi: 10.1167/iovs.18-25372

749   44.   Shi H, Koronyo Y, Rentsendorj A, Regis GC, Sheyn J, Fuchs DT, Kramerov AA, Ljubimov A V.,

750   Dumitrascu OM, Rodriguez AR, Barron E, Hinton DR, Black KL, Miller CA, Mirzaei N, Koronyo-

751   Hamaoui M (2020) Identification of early pericyte loss and vascular amyloidosis in Alzheimer's

752   disease retina. Acta Neuropathol 139:813–836. doi: 10.1007/s00401-020-02134-w

45. Sidiqi A, Wahl D, Lee S, Ma D, To E, Cui J, To E, Beg MF, Sarunic M, Matsubara JA (2020) In vivo

Retinal Fluorescence Imaging With Curcumin in an Alzheimer Mouse Model. Front Neurosci

14:713. doi: 10.3389/fnins.2020.00713

46. Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM, Iwatsubo T, Jack CR, Kaye J,

Montine TJ, Park DC, Reiman EM, Rowe CC, Siemers E, Stern Y, Yaffe K, Carrillo MC, Thies B,

Morrison-Bogorad M, Wagster M V., Phelps CH (2011) Toward defining the preclinical stages of

Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's

Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's Dement

7:280–292. doi: 10.1016/j.jalz.2011.03.003

47. Sperling RA, Jack CR, Aisen PS (2011) Testing the right target and right drug at the right stage. Sci.

Transl. Med. 3:111cm33

48. De Strooper B, Karran E (2016) The Cellular Phase of Alzheimer's Disease. Cell 164:603–615

49. Tolar M, Abushakra S, Sabbagh M (2020) The path forward in Alzheimer's disease therapeutics:

Reevaluating the amyloid cascade hypothesis. Alzheimer's Dement. doi:

10.1016/j.jalz.2019.09.075

50. Valiente-Soriano FJ, García-Ayuso D, Ortín-Martínez A, Jiménez-López M, Galindo-Romero C,

Villegas-Pérez MP, Agudo-Barriuso M, Vugler AA, Vidal-Sanz M (2014) Distribution of melanopsin

positive neurons in pigmented and albino mice: evidence for melanopsin interneurons in the

mouse retina. Front Neuroanat. doi: 10.3389/fnana.2014.00131

51. Villemagne VL, Burnham S, Bourgeat P, Brown B, Ellis KA, Salvado O, Szoeke C, Macaulay SL,

Martins R, Maruff P, Ames D, Rowe CC, Masters CL (2013) Amyloid β deposition,

neurodegeneration, and cognitive decline in sporadic Alzheimer's disease: A prospective cohort

study. Lancet Neurol 12:357–367. doi: 10.1016/S1474-4422(13)70044-9

52. Wang ZX, Tan L, Liu J, Yu JT (2016) The Essential Role of Soluble Aβ Oligomers in Alzheimer's

Disease. Mol. Neurobiol. 53:1905–1924

53. Williams MA, McGowan AJ, Cardwell CR, Cheung CY, Craig D, Passmore P, Silvestri G, Maxwell AP,

McKay GJ (2015) Retinal microvascular network attenuation in Alzheimer's disease. Alzheimer's

Dement Diagnosis, Assess Dis Monit. doi: 10.1016/j.dadm.2015.04.001

54. Yang Y, Shiao C, Hemingway JF, Jorstad NL, Shalloway BR, Chang R, Keene CD (2013) Suppressed

Retinal Degeneration in Aged Wild Type and APPswe/PS1ΔE9 Mice by Bone Marrow

Transplantation. PLoS One. doi: 10.1371/journal.pone.0064246

55. Zabel P, Kaluzny JJ, Wilkosc-Debczynska M, Gebska-Toloczko M, Suwala K, Zabel K, Zaron A,

Kucharski R, Araszkiewicz A (2019) Comparison of retinal microvasculature in patients with

Alzheimer's disease and primary open-angle glaucoma by optical coherence tomography

angiography. Investig Ophthalmol Vis Sci. doi: 10.1167/IOVS.19-27028

56. Zudaire E, Gambardella L, Kurcz C, Vermeren S (2011) A Computational Tool for Quantitative

Analysis of Vascular Networks. PLoS One 6:e27385. doi: 10.1371/journal.pone.0027385