partR2: Partitioning R^2 in generalized linear mixed models

Martin A. Stoffel[1,2], Shinichi Nakagawa[3], Holger Schielzeth[1]

[1] Institute of Ecology and Evolution, Friedrich Schiller University Jena, Germany

[2] Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, UK

[3] Evolution & Ecology Research Centre and School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, Australia

ORCID: MAS 0000-0003-4030-3543, SN 0000-0002-7765-5182, HS 0000-0002-9124-2261

Corresponding authors

Martin A. Stoffel, martin.stoffel@ed.ac.uk

Holger Schielzeth, holger.schielzeth@uni-jena.de

31   **Abstract**

32   The coefficient of determination R^2 quantifies the amount of variance explained by regression

33   coefficients in a linear model. It can be seen as the fixed-effects complement to the repeatability R

34   (intra-class correlation) for the variance explained by random effects and thus as a tool for variance

35   decomposition. The R^2 of a model can be further partitioned into the variance explained by a

36   particular predictor or a combination of predictors using semi-partial (part) R^2 and structure

37   coefficients, but this is rarely done due to a lack of software implementing these statistics. Here, we

38   introduce `partR2`, an R package that quantifies part R^2 for fixed effect predictors based on

39   (generalized) linear mixed-effect model fits. The package iteratively removes predictors of interest

40   from the model and monitors the change in the variance of the linear predictor. The difference to

41   the full model gives a measure of the amount of variance explained uniquely by a particular predictor

42   or a set of predictors. `partR2` also estimates structure coefficients as the correlation between a

43   predictor and fitted values, which provide an estimate of the total contribution of a fixed effect to the

44   overall prediction, independent of other predictors. Structure coefficients can be converted to the

45   total variance explained by a predictor, here called 'inclusive' R^2, as the square of the structure

46   coefficients times total R^2. Furthermore, the package reports beta weights (standardized regression

47   coefficients). Finally, `partR2` implements parametric bootstrapping to quantify confidence intervals

48   for each estimate. We illustrate the use of `partR2` with real example datasets for Gaussian and

49   binomials GLMMs and discuss interactions, which pose a specific challenge for partitioning the

50   explained variance among predictors.

Introduction

Coefficients of determination R^2 are of interest in the study of ecology and evolution, because they quantify the amount of variation explained by a linear model (Edwards et al., 2008). By doing so, they go beyond significance testing in putting effects in perspective of the phenotypic variance. R^2 is expressed as a proportion of the total variance in the response, which represents a biologically relevant quantity if the total variation is representative for the total population (de Villemereuil et al., 2018). The total coefficient of determination in a generalised linear mixed model (GLMM) quantifies the variance explained by all fixed effects together (marginal R^2 *sensu* Nakagawa & Schielzeth, 2013, also known as the total correlation coefficient, Watanabe (1960)).

However, it is often of interest to attribute explained variation to individual predictors. Semi-partial coefficients of determination, also known as part R^2, decompose the variance of R^2 into components uniquely explained by individual predictors (Jaeger et al., 2017; Jaeger, Edwards & Gurka, 2019) or sets of predictors (Figure 1). The set of all predictors in the model yields the total proportion of variance explained by the fixed part of the model (total R^2). With correlations among predictors, it often happens that predictors in univariate regressions explain a large share of the variance, but do not show large part R^2 if other correlated predictors are included in the model. Note that part R^2 estimates the proportion of the variance in the response explained by a predictor *while* accounting for covariance between this predictor and the other predictors in the model, whereas the (arguably more familiar) partial R^2 estimates the proportion of the variance that is explained by a predictor of interest *after* accounting for the other predictors from the response as well as the predictor of interest. The difference is subtle, but important (see more below). Therefore, part R^2 represents 'variance accounted for' in relation to the total variance, but partial R^2 does not. Consequently, part R^2 will be conceptually easier to compare with (total) R^2.

Structure coefficients provide a valuable addition to part R^2 in the decomposition of the phenotypic variance (Nimon et al., 2008; Yeatts et al., 2017). Structure coefficients quantify the correlation between individual predictors and the linear predictor. Predictors that correlate well with a response, but are fitted with collinear predictors may show large structure coefficients as they are correlated to the predicted response, but low part R^2 as other predictors explain part of the same variance. Structure coefficients range from -1 to 1 with their absolute value expressing the correlation relative to a perfect correlation if a single predictor explains as much as the total fixed part of the model.

Structure coefficients are correlations and since the square of a correlation yields the variance

explained, we can use structure coefficients to estimate the total variance explained by a predictor

(Nimon et al., 2008). We call this the inclusive R^2 of a predictor and calculate it as the squared

structure coefficient, i.e. its contribution to the linear predictor independent of other predictors

(Nimon et al., 2008) times the proportion of variance explained by the linear predictor (which is the

'total' marginal R^2 of the model). As far as we are aware, inclusive R^2 has not been implemented

before, but it provides valuable insights into the structure of the variance explained (Figure 1).

Here, we introduce `partR2`, a versatile package for estimating part R^2, inclusive R^2, structure

coefficients and beta weights from mixed-effects models. The analysis is conceptually related to, but

not identical, to commonality analysis (Ray-Mukherjee et al., 2014; Seibold & McPhee, 1979; Zientek

& Thompson, 2006, Figure 1). We illustrate how to use `partR2` with real example datasets for

Gaussian and binomial GLMMs, discuss how to estimate part R^2 in the presence of interactions and

discuss some challenges and limitations.

Figure 1: Conceptual framework for the estimation of proportions of variance components in a mixed
model. The large grey circle symbolizes the variance in a response Y, the dark grey area on the top indicates
the share explained by random effects and the coloured ellipses symbolize variance in covariates with
intersections indicating jointly explained variances. `partR2` calculates total R^2, part R^2 for individual predictors
and sets of predictors as well as inclusive R^2. The package does not report partial R^2 and commonalities,
although they could be calculated from the `partR2` output.

Mathematical representation

Part R^2

A Gaussian mixed-effects model can be written as:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \sum \boldsymbol{\alpha}_k + \boldsymbol{\varepsilon} \qquad \text{(Eq. 1)}$$

Where $\mathbf{y}$ is a vector of response values (outcomes), $\mathbf{X}$ is the design matrix of fixed effects, $\boldsymbol{\beta}$ is a

vector of regression coefficients, $\sum \boldsymbol{\alpha}_k$ is the random part of the model that might contain multiple

(1 to k) random effects and $\boldsymbol{\varepsilon}$ is a vector or residual deviations. The linear predictor $\boldsymbol{\eta}$ represents

the vector of predicted values from the fixed part of the model as $\boldsymbol{\eta} = \mathbf{X}\boldsymbol{\beta}$.

Since we are interested in the proportion of the phenotypic variance explained, we symbolize

variance components by upper case Y and index by the source of variance (als in Figure 1). While

variances are frequently represented as V with the source of variance as an index, this leads to

ambiguity for V_X which might represent variance in y explained by x or variance in x itself, which is

why we use this alternative notation. The total variance in the response is $Y_{Total} = \mathrm{var}(y)$ and is

estimated from the raw data or from the model (see below). The variance of the residuals is

estimated by the model as $Y_R = \mathrm{var}(\varepsilon)$. The variance of the (sum of) random effects is estimated

by the model as $Y_{RE} = \mathrm{var}(\sum \boldsymbol{\alpha}_k)$ and the variance explained by fixed effects can be estimated as

the variance in the linear prector $Y_X = \mathrm{var}(\mathbf{X}\boldsymbol{\beta})$.

The coefficient of determination R^2 estimates the proportion of variance in the response that is

explained by fixed effects. The coefficient of determination R^2 for the total fixed part of the model

is thus:

$$R_x^2 = \frac{Y_X}{Y_X + Y_{RE} + Y_R} = \frac{Y_X}{Y_{Total}} \qquad \text{(Eq. 2)}$$

Note that the sum of the components in the denominator might deviate numerically from the total

outcome variance in the raw data. However, conceptually they are the same in that they represent

the population-level outcome variance. The variance in the outcome is an estimate from the

specific sample, while the sum of components of the mixed model represents a population-level

estimate given the data and the model.

A reduced model with a (set of) fixed effect predictors $x*$ removed but the same random effect

structure can be fitted as (now using the tilde to highlight the differences from Eq. 1):

$\mathbf{y} = \widetilde{\mathbf{X}}\widetilde{\boldsymbol{\beta}} + \sum \widetilde{\boldsymbol{\alpha}}_k + \widetilde{\boldsymbol{\varepsilon}}$ (Eq. 3)

with the variance in the linear predictor of the reduced model being $Y_{\widetilde{X}} = \mathrm{var}(\widetilde{\mathbf{X}}\widetilde{\boldsymbol{\beta}})$.

The variance uniquely explained by $x*$ is then the difference between the variance explained by

fixed effects in the full and the reduced model $Y_{X*} = Y_X - Y_{\widetilde{X}}$. Part R^2 sets this variance in

proportion to the total outcome variance:

$R^2_{X*} = \dfrac{Y_X - Y_{\widetilde{X}}}{Y_X + Y_{RE} + Y_R} = \dfrac{Y_X - Y_{\widetilde{X}}}{Y_{Total}}$ (Eq. 4)

The process of fitting a reduced model, estimation of Y_{X*} and estimation of R^2_{X*} can be repeated

for all predictors and combinations of predictors. At the limit for a model with all fixed effects

removed, $R^2_{x*} = R^2_X$.

Side-note on partial R^2

For completeness we note that the partial R^2 could be calculated as:

$R^2_{X*} = \dfrac{Y_X - Y_{\widetilde{X}}}{Y_{Total} - Y_{\widetilde{X}}}$ (Eq. 5)

However, this estimate does not put the explained variance in perspective of the total variance in

the response. It has the major disadvantage that the denominator depends on $Y_{\widetilde{X}}$. The same

effect in terms of Y_{x*} thus appears larger if the reduced model explains more variance (larger $Y_{\widetilde{X}}$).

Even in the case of independent additive predictors, the contributions of the different fixed effects

do not sum up to R^2_X, because of the change in the denominator that different Y_{x*} are compared

to. Finally, since we are interested in explaining phenotypic variation in some biological response

(the phenomenon to be explained), we think that part R^2 is the more relevant quantity, as it

represents the proportion of variance in the response uniquely explained by $x*$.

Inclusive R^2

Structure coefficients are the correlations between a particular predictor of interest $x*$ and the

linear predictor η. Structure coefficients are quantified from the full model as:

$SC_{x*} = \mathrm{cor}(\eta, x*)$ (Eq. 6)

The squared correlation between two variables *a* and *b* gives the variance explained for these

variables $\mathrm{cor}(a,b)^2 = R_a^2$. The squared structure correlations thus quantify the proportion of

variance in the linear predictor Y_X that is explained by a the predictor of interest $x*$. Since the

proportion of outcome variance explained by the linear predictor in the full model is R_x^2, the

inclusive variance explained by predictor $x*$ is:

$$IR_{x*}^2 = SC^2 \cdot R_x^2 \qquad\qquad\qquad \text{(Eq. 7)}$$

Inclusive R^2 as we define it here, complements part R^2 by giving additional insights. While part R^2

quantifies the variance uniquely explained by a predictor (or set of predictors), inclusive R^2

quantifies the total proportion of variance explained in the model, both uniquely and jointly with

other predictors. In the special case of a single predictor in a model $SC_{x*} = \mathrm{cor}(\eta, x*) = 1$, such

that $IR_{X*}^2 = R_x^2$.

Part R^2 in non-Gaussian models

For Gaussian models there is a single residual error term ε with variance $Y_R = \mathrm{var}(\varepsilon)$. For non-

Gaussian models, however, there is additional error that arises from the link function that translates

latent-level predictions to observed outcomes. This variance can be approximated for a variety of

link functions and error distributions (Nakagawa & Schielzeth, 2010; Nakagawa, Johnson &

Schielzeth, 2017). Our R package currently implements distribution-specific variances for Poisson

models with log and square root link functions and binomial models with logit and probit link

functions. For Poisson models and non-binary binomial models (proportion models), `partR2` also

fits an observational level random effect (if none is fitted already) to estimate variance due to

overdispersion (Harrison, 2014). Both the overdispersion variance, now denoted Y_R and the

distribution-specific variance Y_D are included in the denominator of the part R2 calculation:

$$R_{X*}^2 = \frac{Y_X - Y_{\widetilde{X}}}{Y_X + Y_{RE} + Y_D + Y_R} \qquad\qquad \text{(Eq. 8)}.$$

198
       Notably, there are other estimation methods for R^2 for non-Gaussian models or GLMM (Jaeger et
199
       al., 2017; Piepho, 2019). Currently, `partR2` only implments the method based on Nakagawa &
200
       Schielzeth (2013) and Nakagawa, Johnson & Schielzeth (2017).

Other implementations in R packages

There are a few R packages that calculate part R^2 for linear models (lm), for example `rockchalk::getDeltaRsquare` (Johnson & Grothendieck, 2019). Other packages calculate partial R^2 (not part R^2) such as `asbio::partial.R2` (Aho, 2020) and `rr2::R2` (Ives & Li, 2018) for lms and `rsq::rsq.partial` (Zhang, 2020) for linear models and generalized linear models (glm). Note that partial R^2 is different from part (semi-partial) R^2 (partial R^2 > part R^2), since it represents the unique variance explained by a particular predictor but after removing ('partialling out') the variance explained by the other predictors (Yeatts et al., 2017, Figure 1). The ppcor package calculates semi-partial and partial correlations, but does not work on fitted GLM or GLMM models (Kim, 2015). The package `yhat` features a suite of functions for full communality analyses in glms (Nimon, Oswald & Roberts, 2020). None of these packages estimates part R^2 for mixed-effects models that we focus on here.

Several packages estimate (marginal) R^2 as the variance explained by all fixed effects in linear mixed-effects models. This includes `performance::r2_nakagawa` (Lüdecke et al., 2020), `MuMIn::r.squaredGLMM` (Bartoń, 2019), and `rptR::rpt` (Stoffel, Nakagawa & Schielzeth, 2017). These packages do not allow to estimate part R^2. The only versatile package to estimate part R^2 from linear mixed-models is `r2glmm` (Jaeger, 2017). The function `r2glmm::r2beta` computes part R^2 from lmer, lme and glmmPQL model fits (also for linear models lm and glm) based on Wald statistics. However, it does neither support `lme4::glmer` for generalized linear model fits nor does it allow to estimate R^2 for combinations of predictors. Furthermore, it does not estimate structure coefficients, inclusive R^2 or part R^2 for multilevel factors as a unit.

Features of `partR2`

`partR2` takes a fitted (generalized) linear mixed-model (GLMM), from the popular mixed model package `lme4` (Bates et al., 2015) and estimates part R^2 by iteratively removing fixed effects (Nimon et al., 2008). The specific fixed effects of interest are specified by the `partvars` and/or by the `partbatch` argument. The package estimates part R^2 for all predictors specified in `partvars` individually and in all possible combinations (the maximum level of combinations can be set by the `max_level` argument). A custom specification of fixed effects of interest saves computation time as compared to an all-subset specification and is therefore required in `partR2`.

The central function `partR2` will work for Gaussian, Poisson and binomial GLMMs. Since the model

fit is done externally, there is no need to supply a family argument. For non-Gaussian GLMMs, the

package estimates link-scale R^2 *(sensu* Nakagawa & Schielzeth, 2013*)*. We implement parametric

bootstrapping to quantify sampling variance and thus uncertainty in the estimates. Parametric

bootstrapping works through repeated model fitting on simulated data based on fitted values

(Faraway, 2015). The number of bootstrap iterations is controlled by the `nboot` argument. We

recommend a low number of `nboot` for testing purposes and a large number (e.g. `nboot = 1000`)

for the final analysis.

The package returns an object of class `partR2` that contains elements for part R^2, inclusive R^2,

structure coefficients, beta weights (standardized regression slopes), bootstrapping iterations and

some other information. An extended summary, that includes inclusive R^2, structure coefficients and

beta weights can be viewed using the `summary` function. The `forestplot` function shows a

graphical representation of the variance explained by individual predictors and sets of predictors

along with their bootstrapping uncertainties. All computations can be parallelized across many cores

based on the `future` and `furrr` packages (Vaughan & Dancho, 2018; Bengtsson, 2020). An

extended vignette with details on the complete functionality accompanies the package.

Example with Gaussian data

We use an example dataset with hormone data collected from a population of captive guinea pigs

to illustrate the features of `partR2`. The dataset contains testosterone measurements of 31 male

guinea pigs, each measured at 5 time points (age between 120 and 240 days at 30-day intervals).

We analyze log-transformed testosterone titers and fit male identity as a random effect. As covariates

the dataset contains the time point of measurement and a rank index derived from behavioral

observations around the time of measurement (Mutwill et al., in prep.).

Rank and *Time* are correlated in the dataset (*r* = 0.40), since young individuals are typically low rank,

while older individuals tend to hold a high rank. *Time* might be fitted as a continuous predictor or as

a factor with five levels. Here we present the version of a factorial predictor to illustrate the estimation

of part R^2 for interactions terms. Hence, an interaction between time and rank will also be fitted.

First, the package needs to be loaded (after successful installation) in an R session (R Core Team,

2019). The package comes with the guinea pig dataset that also needs to be loaded using the data

function.

`library(partR2)`

```
264   data(GuineaPigs)
265
```

A single record contains missing values for testosterone measurements. Missing records can be problematic to handle in `partR2` and are better removed prior to the analysis. We also log-transform the response and convert *Time* to a factor and filter for the first three time points to simplify the output.

```
270
271   GuineaPigs <- subset(GuineaPigs,
272             !is.na(Testo) & !is.na(Rank) & (Time %in% c(1,3,5)))
273   GuineaPigs$TestoTrans <- log(GuineaPigs$Testo)
274   GuineaPigs$Time <- factor(GuineaPigs$Time)
275
```

We then fit a linear mixed effects model using `lmer` from the `lme4` package (Bates et al., 2015). Further exploration of the data and model checks are omitted here for simplicity, but are advisable in real data analysis.

```
279
280   library(lme4)
281   mod <- lmer(TestoTrans ~ Rank * Time + (1|MaleID), data=GuineaPigs)
282
```

The `partR2` analysis takes the `lmer` model fit (an `merMod` object) and a character vector `partvars` indicating the fixed effects to be evaluated. Interactions are specified with the colon syntax (see the package's vignette for further details).

```
286
287   res <- partR2(mod, partvars = c("Rank", "Time", "Rank:Time"),
288                 nboot=100)
289
```

The function returns a `partR2` object. The `print` function reports the part coefficients of determination and a more extensive summary can be viewed with the `summary` function which also shows inclusive R^2, structure coefficients and beta weights (standardized slopes) (Figure 2).

```
293
294   print(res)
```

```
295    summary(res, round_to = 2)

296

       R2 (marginal) and 95% CI for the full model:
        R2   CI_lower CI_upper ndf
        0.17 0.09     0.36      6

       ----------

       Part (semi-partial) R2:
        Predictor(s)         R2   CI_lower CI_upper ndf
        Model                0.17 0.09     0.36      6
        Rank                 0.00 0.00     0.18      6
        Time                 0.02 0.00     0.20      4
        Rank:Time            0.04 0.00     0.21      4
        Rank+Time            0.02 0.00     0.20      4
        Rank+Rank:Time       0.16 0.08     0.34      3
        Time+Rank:Time       0.04 0.00     0.22      2
        Rank+Time+Rank:Time  0.17 0.09     0.36      1

       ----------

       Inclusive R2 (SC^2 * R2):
        Predictor  IR2  CI_lower CI_upper
        Rank       0.13 0.03     0.26
        Time3      0.00 0.00     0.04
        Time5      0.00 0.00     0.04
        Rank:Time3 0.05 0.01     0.13
        Rank:Time5 0.01 0.00     0.07

       ----------

       Structure coefficients r(Yhat,x):
        Predictor  SC   CI_lower CI_upper
        Rank       0.87  0.56    0.94
        Time3      0.14 -0.18    0.43
        Time5      0.16 -0.26    0.48
        Rank:Time3 0.56  0.22    0.75
        Rank:Time5 0.28 -0.14    0.57

       ----------

       Beta weights (standardised estimates)
        Predictor  BW    CI_lower CI_upper
        Rank        0.50 -0.08    0.94
        Time3      -0.19 -0.53    0.14
        Time5       0.17 -0.20    0.55
        Rank:Time3  0.17 -0.36    0.83
        Rank:Time5 -0.36 -0.95    0.38

       ----------
297
```

298 **Figure 2:** Summary output for example data analysis with Gaussian data (guinea pig analysis).
299

300   The variances appear largely additive, since combinations of predictors explain about the sum of the

301   variance explained by individual predictors. The main components of the `partR2` object can be

302   accessed for further processing as `res$R2` for part R^2 (with point estimates and confidence intervals),

303   `res$SC` for structure coefficients, `res$IR2` for inclusive R^2 and `res$BW` for beta weights.

Dealing with interactions

306 Models with interaction are problematic, because the variance explained by a main factor can be

307 estimated in multiple ways (Figure 3) and because of internal parametrization of the model matrix.

Figure 3: Conceptual framework for dealing with interactions. An interaction is the product of two main

effects and thus often correlated with each of the main effects. The figure shows three options for estimating

the part R^2 for main effects that are involved in an interaction.

310
311
312
313

The model output above shows the number of parameters fitted in each model (Figure 2, each row

in the R2 part refers to a reduced model). In the `print` and `summary` output this is visible as a column

labelled 'ndf'. A close inspection shows that the removal of rank did not change the number of

parameters (6 for the full model, 6 for the model excluding rank). This is because the model matrix

is reparametrized in the reduced model and `lmer` will fit three terms for the interaction (here

`Time1:Rank, Time3:Rank, Time5:Rank`) rather than just two for the interaction in the full model.

Dummy coding of the factor can be usefully combined with centering of dummy coded variables

(Schielzeth, 2010) and gives more control over this re-parametrisation. It allows for example to

estimate the part R^2 for the average effect of *Rank* by constraining the average *Rank* effect to zero,

so that only the two contrasts are fitted (here `Time3:Rank, Time5:Rank`):

324
325 ```
GuineaPigs <- cbind(GuineaPigs, model.matrix(~ 0 + Time,
```

326  ```
     data=GuineaPigs))
```

327 ```
GuineaPigs$Time3 <- GuineaPigs$Time3 - mean(GuineaPigs$Time3)
```

328  ```
GuineaPigs$Time5 <- GuineaPigs$Time5 - mean(GuineaPigs$Time5)
```

The model can then be fitted with dummy predictors. Since the usual specification in `partR2` via

`partvars` would fit all possible combinations, including combinations of the different *Time* terms,

such a run can take a long time. However we are mostly interested in fitting and removing all dummy

predictors at a time. The package therefore features an additional argument `partbatch` to specify

a list of character vectors containing the sets of predictors that should always be kept together. In

the example, the list has two elements, a character vector for the dummy-coded main effects and a

character vector for the interaction terms. The analysis yields part R^2 for two batches of predictors as

well as *Rank* and their combinations.

338
339 ```
mod <- lmer(TestoTrans ~ (Time3 + Time5) * Rank + (1|MaleID),
```

340  ```
     data=GuineaPigs)
```

341 ```
batch <- c("Time3", "Time5")
```

342  ```
partR2(mod, partvars=c("Rank"), partbatch=list(Time=batch,
```

343 ```
     `Time:Rank`= paste0(batch, ":Rank")), nboot=100)
```

345  This, however, is only one way of dealing with interactions (Option A in Figure 3). It represents the

346  variance uniquely explained by main effects even in the presence of an interaction. Since interactions

347  are the products of main effects, interaction terms are typically correlated with main effects and the

348  part $R^2$ calculated above might not represent a biologically relevant quantity. There are two

349     alternative ways of how to deal with interactions. Both are possible in `partR2`, but since

350     requirements differ between applications, we do not implement one with priority.

351     One way to think about variance explained by main effects and their interactions is to pool the

352     variance explained by a main effect with the variance explained by interactions that the term is

353     involved in (Option B in Figure 3). In the guinea pig example, for instance, *Rank* might be considered

354     important either as a main effect or in interaction with time and we might want to estimate the total

355     effect of rank. This can be done for the guinea pig dataset by using `partbatch`:

356
```
357 mod <- lmer(Testo ~ Time * Rank + (1|MaleID), data=GuineaPigs)

358 partR2(mod, partbatch = list(Time=c("Time", "Time:Rank"),

359   Rank=c("Rank", "Time:Rank")), nboot=100)
```

361     A third, which we think usually preferable option is to prioritize main effects by assigning the

362     proportion of variance that is explained by a main effect together with the variance jointly explained

363     with its interaction to the main effect (Option C in Figure 3). This implies that part $R^2$ for a main effect

364     is estimated when its own interaction is excluded from the model (`mod1` and `part1` below). The

365     variance explained by the interaction is then estimated in a separate model (`mod2` and `part2` below).

366     We have implemented a helper function `mergeR2` that allows to merge two `partR2` runs.

367
```
368 mod1 <- lmer(Testo ~ Time * Rank + (1|MaleID), data=GuineaPigs)

369 part1 <- partR2(mod1, partvars = c("Time:Rank"), nboot=100)

370 mod2 <- lmer(Testo ~ Time + Rank + (1|MaleID), data=GuineaPigs)

371 part2 <- partR2(mod2, partvars = c("Time", "Rank"), nboot=100)

372 mergeR2(part1, part2)
```

374     All these results can be viewed by `print`, `summary` and plotted by `forestplot`. It is important to

375     bear in mind the differences in the interpretation as illustrated in Figure 3.

377     **An example with proportion data**

378     As an example for proportion data, we analyze a dataset on spatial variation in color morph ratios in

379     a color-polymorphic species of grasshopper. Individuals of this species occur either in a green or a

380   brown color variant and the dataset contains counts of brown and green individuals (separated for

381   females and males) from 42 sites sampled in the field (Dieker et al., 2018). Site identity will be fitted

382   as a random effect. As covariates the dataset contains a range of Bioclim variable that describe

383   various aspects of ecologically relevant climatic conditions (Karger et al., 2017). The aim is to identify

384   the climatic conditions that favour one or the other colour variant.

385   We first load the grasshopper dataset. We standardise all Bioclim variables using the `scale` function

386   and add an observation-level counter that will be used as an observation-level random effect (OLRE)

387   to account for overdispersion (Harrison, 2014).

389   ```
data(Grasshoppers)
```

390   ```
for (i in which(substr(colnames(Grasshoppers),1,3)=="Bio")){
```

391   ```
    Grasshoppers[,i] <- scale(Grasshoppers[,i])
```

392   ```
}
```

393   ```
Grasshoppers$OLRE <- 1:nrow(Grasshoppers)
```

We first fit a GLMM with binomial error structure and logit link using the `glmer` function from the

`lme4` package (Bates et al., 2015). A previous analysis has shown that the first principle component

of the Bioclim data explains a small, but significant part of variation in morph ratios (Dieker et al.,

2018). For illustration, we use the four Bioclim variables that show a loading of more than 0.30 on

the first principle component.

```
mod <- glmer(cbind(nGreen, nBrown) ~ Bio7 + Bio14 + Bio17 + Bio19 +
```

402   ```
    (1|SiteID) + (1|OLRE), data=Grasshoppers, family="binomial")
```

403   ```
res <- partR2(mod, partvars=c("Bio7", "Bio14", "Bio17", "Bio19"),
```

404   ```
    max_level = 1, nboot=100)
```

The `summary` output informs us (at the bottom) that there have been warnings in the bootstrapping

processes. This is not unusual since bootstrapping frequently generates data, for which one of the

parameters is estimated at the boundary (in particular if one of the variance components is very

small). The results can be visualised using the `forestplot` function (Figure 4). Plotting is based on

`ggplot2` (Wickham, 2016), and multiple forest plots can easily be assembled using the `patchwork`

package (Pedersen, 2020). Forest plots show the effect sizes graphically and can be set to either

show part $R^2$ when `type = "R2"` (the default), inclusive $R^2$ when `type = "IR2"`, structure

coefficients when `type = "SC"`, and beta weights (standardized model estimates) with `type =`

`"BW"`.

```
416  p1 <- forestplot(res, type = "R2")

417  p2 <- forestplot(res, type = "IR2")

418  p3 <- forestplot(res, type = "SC")

419  p4 <- forestplot(res, type = "BW")

420

421  library(patchwork)

422  (p1 + p2) / (p3 + p4) + plot_annotation(tag_levels = "A",

423      tag_prefix = "(", tag_suffix = ")")
```

Figure 4: Comparison of part R^2 for individual predictors (A), inclusive R^2 (B), structure coefficients (C) and beta weights (D) for an example dataset with proportion data from grasshoppers.

A comparison of part R^2, inclusive R^2, structure coefficients beta weights shows the different insights

that can be gained from these different summaries of the model fit (Figure 3). In this case, three of

the Bioclim variables (*Bio14*, *Bio17*, *Bio19*) are highly positively correlated (r ≥ 0.93), while a fourth

one (*Bio7*) is moderately negatively correlated to all three of them (r ≤ -0.63). Part R^2 are thus low,

because none of the parameters uniquely explains a large share of the variance. *Bio17* seems to be

the best predictor of morph ratios, with the largest (negative) beta weight, largest part R^2, largest

structure coefficients and largest inclusive R^2. Beta weights for the two positively correlated (but

slightly weaker) predictors, *Bio14* and *Bio19*, switch sign as is not unusual for collinear predictors.

This means that after accounting for the effect of *Bio17*, they contribute positively to prediction.

However, structure coefficients show that both variables load negatively on the linear predictor, as

does *Bio17*.

Challenges

Using transformation or functions in the formula argument can lead to issues with matching the terms

of the model with the `partvars` argument of `partR2`. It is therefore important that the names in

`partvars` match exactly the terms in the `merMod` object. However, any complications are easily

circumvented by implementing the transformations before fitting the model and storing them in the

data frame used in the analysis. It is also worth to be aware that unusual names may cause

complications and renaming can offer an easy solution.

We have repeatedly seen model outputs where the point estimate does not fall within the confidence

interval. This might seem like in the bug in the package, but in our experience usually indicates issues

with the data and/or the model. In fact, parametric bootstrapping can be seen as a limited form of

posterior predictive model checks (Gelman & Hill, 2007). If generating new data from the fitted

model (as done with parametric bootstrapping) results in data that are dissimilar to the original data,

then the model is probably not a good fit to the data.

Bootstrap iterations can sometimes yield slightly negative estimates of part R^2, in particular if the

variance explained by a predictor is low. These negative estimates happen in mixed-effects models,

because estimates of random-effect variance might change when a predictor is removed and this

can lead to a slight decrease in the residual variance, and hence a proportional increase in R^2 (see

also Rights & Sterba, 2019). By default, partR2 sets negative R^2 values to 0, but this can be changed

by setting `allow_neg_r2` to `TRUE`. It also happens that inclusive R^2 is estimated slightly lower than

part R^2 when the contribution of a particular predictor is very large. We consider both cases as

sampling error that should serve as a reminder that variance components are estimated with

relatively large uncertainly and minor differences should not be over-interpreted.

A warning needs to be added for the estimation of R^2 (and, in fact, also repeatability R) from small

datasets. In particular if the number of levels of random effect is low, variance components might be

slightly overestimated (Xu, 2003). This issue applies similarly to the variance explained by fixed

effects, in particular if the number of predictors is large relative to the number of data points.

Code and data availability

The `partR2` package can be downloaded from CRAN with `install.packages("partR2")` and

includes the data used in the examples. The development version can be obtained from GitHub

(https://github.com/mastoffel/partR2).

Author contributions

SN and HS conceived the idea. HS and MAS wrote the manuscript with contributions from SN. MAS

wrote the package and the vignette, supervised by HS. All authors contributed critically to the

manuscript, package and vignette and gave final approval for publication.

Acknowledgements

MAS and HS were supported by the German Research Foundation (DFG) as part of the SFB TRR 212

(NC³) (funding INST 215/543-1, 396782608). SN was supported by the ARC Discovery Project grant

(DP180100818). We thank Alexandra Mutwill for providing the guinea pig data.

References

Aho K. 2020.asbio: A Collection of Statistical Tools for Biologists. *Available at https://CRAN.R-*

project.org/package=asbio (accessed July 25, 2020).

Bartoń K. 2019.MuMIn: Multi-Model Inference. *Available at https://CRAN.R-*

project.org/package=MuMIn (accessed July 25, 2020).

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4.

Journal of Statistical Software 67:1–48.

Bengtsson H. 2020.future: Unified Parallel and Distributed Processing in R for Everyone. *Available*

at https://CRAN.R-project.org/package=future (accessed July 25, 2020).

Dieker P, Beckmann L, Teckentrup J, Schielzeth H. 2018. Spatial analyses of two colour

polymorphisms in an alpine grasshopper reveal a role of small-scale heterogeneity.

Ecology and Evolution 8:7273–7284. DOI: 10.1002/ece3.4156.

Edwards LJ, Muller KE, Wolfinger RD, Qaqish BF, Schabenberger O. 2008. An R2 statistic for fixed

effects in the linear mixed model. *Statistics in Medicine* 27:6137–6157. DOI:

10.1002/Sim.3429.

Faraway JJ. 2015. *Linear models with R*. Boca Raton: CRC Press, Taylor & Francis Group.

Gelman A, Hill J. 2007. *Data analysis using regression and multilevel/hierarchical models*.

Cambridge; New York: Cambridge University Press.

Harrison XA. 2014. Using observation-level random effects to model overdispersion in count data

in ecology and evolution. *PeerJ* 2:e616. DOI: 10.7717/peerj.616.

Ives A, Li D. 2018. rr2: An R package to calculate R2s for regression models. *Journal of Open*

Source Software 3:1028.

Jaeger B. 2017.r2glmm: Computes R Squared for Mixed (Multilevel) Models. *Available at*

https://CRAN.R-project.org/package=r2glmm (accessed July 25, 2020).

Jaeger BC, Edwards LJ, Das K, Sen PK. 2017. An R2 statistic for fixed effects in the generalized

linear mixed model. *Journal of Applied Statistics* 44:1086–1105. DOI:

10.1080/02664763.2016.1193725.

Jaeger BC, Edwards LJ, Gurka MJ. 2019. An R 2 statistic for covariance model selection in the

linear mixed model. *Journal of Applied Statistics* 46:164–184.

Johnson PE, Grothendieck G. 2019.rockchalk: Regression Estimation and Presentation. *Available at*

https://CRAN.R-project.org/package=rockchalk (accessed July 25, 2020).

Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP,

Kessler M. 2017. Climatologies at high resolution for the earth's land surface areas.

Scientific data 4:170122.

Kim S. 2015. ppcor: an R package for a fast calculation to semi-partial correlation coefficients.

Communications for statistical applications and methods 22:665.

Lüdecke D, Makowski D, Waggoner P, Patil I. 2020.performance: Assessment of Regression

Models Performance. *Available at https://CRAN.R-project.org/package=performance*

(accessed July 25, 2020).

Mutwill AM, Schielzeth H, Zimmermann TD, Richter H, Kaiser S, Sachser N. in prep. Endocrine

mechanisms underlying dominance rank acquisition in males: individuality meets plasticity

in a complex social environment.

Nakagawa S, Johnson PCD, Schielzeth H. 2017. The coefficient of determination R2 and intra-class

correlation coefficient from generalized linear mixed-effects models revised and

expanded. *Journal of the Royal Society Interface* 14:20170213. DOI:

10.1098/rsif.2017.0213.

Nakagawa S, Schielzeth H. 2010. Repeatability for Gaussian and non-Gaussian data: a practical

guide for biologists. *Biological Reviews* 85:935–956.

Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R2 from generalized

linear mixed-effects models. *Methods in Ecology and Evolution* 4:133–142. DOI:

10.1111/j.2041-210x.2012.00261.x.

Nimon K, Lewis M, Kane R, Haynes RM. 2008. An R package to compute commonality coefficients

in the multiple regression case: an introduction to the package and a practical example.

Behavior Research Methods 40:457–66. DOI: 10.3758/brm.40.2.457.

Nimon K, Oswald F, Roberts JK. 2020.yhat: Interpreting Regression Effects. *Available at*

https://CRAN.R-project.org/package=yhat (accessed July 25, 2020).

Pedersen TL. 2020.patchwork: The Composer of Plots. *Available at https://CRAN.R-*

project.org/package=patchwork (accessed July 25, 2020).

Piepho H-P. 2019. A coefficient of determination (R2) for generalized linear mixed models.

Biometrical Journal 61:860–872.

R Core Team. 2019. *R: A language and environment for statistical computing. R Foundation for*

Statistical Computing. Vienna, Austria.

Ray-Mukherjee J, Nimon K, Mukherjee S, Morris DW, Slotow R, Hamer M. 2014. Using commonality

analysis in multiple regressions: a tool to decompose regression effects in the face of

multicollinearity. *Methods in Ecology and Evolution* 5:320–328. DOI: Doi 10.1111/2041-

210x.12166.

Rights JD, Sterba SK. 2019. New recommendations on the Use of R-squared differences in

multilevel model comparisons. *Multivariate Behavioral Research*:1–32. DOI:

10.1080/00273171.2019.1660605.

Schielzeth H. 2010. Simple means to improve the interpretability of regression coefficients.

Methods in Ecology and Evolution 1:103–113. DOI: 10.1111/j.2041-210X.2010.00012.x.

Seibold DR, McPhee RD. 1979. Commonality analysis: a method for decomposing explained

variance in multiple regression analyses. *Human Communication Research* 5:355–365.

Stoffel MA, Nakagawa S, Schielzeth H. 2017. rptR: repeatability estimation and variance

decomposition by generalized linear mixed-effects models. *Methods in Ecology and*

Evolution 8:1639–1644. DOI: 10.1111/2041-210X.12797.

Vaughan D, Dancho M. 2018.furrr: Apply Mapping Functions in Parallel using Futures. *Available at*

https://CRAN.R-project.org/package=furrr (accessed July 25, 2020).

de Villemereuil P, Morrissey M, Nakagawa S, Schielzeth H. 2018. Fixed effect variance and the

estimation of repeatabilities and heritabilities: Issues and solutions. *Journal of Evolutionary*

Biology 31:621–632. DOI: 10.1111/jeb.13232.

Watanabe S. 1960. Information theoretical analysis of multivariate correlation. *IBM Journal of*

Research and Development 4:66–82. DOI: DOI 10.1147/rd.41.0066.

Xu RH. 2003. Measuring explained variation in linear mixed effects models. *Statistics in Medicine*

22:3527–3541. DOI: Doi 10.1002/Sim.1572.

Yeatts PE, Barton M, Henson RK, Martin SB. 2017. The use of structure coefficients to address

multicollinearity in sport and exercise science. *Measurement in Physical Education and*

Exercise Science 21:83–91.

Zhang D. 2020.rsq: R-Squared and Related Measures. *Available at https://CRAN.R-*

project.org/package=rsq (accessed July 25, 2020).

Zientek LR, Thompson B. 2006. Commonality analysis: Partitioning variance to facilitate better

understanding of data. *Journal of Early Intervention* 28:299–307.

576