Systematic investigation of imprinted gene expression and enrichment in the mouse brain explored at single-cell resolution

M. J. Higgs[1], M. J. Hill[2,3], R. M. John[4], and A. R. Isles[1,*]

1 Behavioural Genetics Group, MRC Centre for Neuropsychiatric, Genetics and Genomics, Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff, UK
2 Division of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Cardiff, UK
3 UK Dementia Research Institute, School of Medicine, Cardiff University, Cardiff, UK
4 School of Biosciences, Cardiff University, Cardiff, UK
* Corresponding author: A. R. Isles, Behavioural Genetics Group, MRC Centre for Neuropsychiatric Genetics and Genomics, Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff CF24 4HQ, UK.
E-mail: islesar1@cardiff.ac.uk

Key words: **Genomic Imprinting, Single Cell Genomics, Imprinted Gene Network, Neuroendocrine**

ABSTRACT

Background

Although a number of imprinted genes are known to be highly expressed in the brain, and in certain brain regions in particular, whether they are truly over-represented in the brain has never been formally tested. Using thirteen single-cell RNA sequencing datasets we systematically investigated imprinted gene over-representation at the organ, brain region, and cell-specific levels.

Results

We established that imprinted genes are indeed over-represented in the adult brain, and in neurons particularly compared to other brain cell-types. We then examined brain-wide datasets to test enrichment within distinct brain regions and neuron subpopulations and demonstrated over-representation of imprinted genes in the hypothalamus, ventral midbrain, pons and medulla. Finally, using datasets focusing on these regions of enrichment, we identified hypothalamic neuroendocrine populations and the monoaminergic hindbrain neurons as specific hotspots of imprinted gene expression.

Conclusions

These analyses provide the first robust assessment of the neural systems on which imprinted genes converge. Moreover, the unbiased approach, with each analysis informed by the findings of the previous level, permits highly informed inferences about the functions on which imprinted gene expression

converges. Our findings indicate the neuronal regulation of motivated behaviours such as feeding and

sleep, alongside the regulation of pituitary function as functional hotspots for imprinting, thus adding

statistical rigour to prior assumptions and providing testable predictions for novel neural and

behavioural phenotypes associated with specific genes and imprinted gene networks. In turn, this work

sheds further light on the potential evolutionary drivers of genomic imprinting in the brain.

BACKGROUND

Imprinted genes demonstrate a preferential or exclusively monoallelic expression from either the

maternal or paternal allele in an epigenetically predetermined manner (a parent-of-origin effect, POE),

and to date approximately 260 imprinted genes (demonstrating biased allelic expression and/or

associated with a parental-specific epigenetic mark) have been identified in the mouse (~230 in humans)

(9, 10). This epigenetic regulation makes genomic imprinting an evolutionary puzzle as many of these

genes are effectively haploid and thereby negate many of the benefits of diploidy (11). Studying the

patterns of expression and function of imprinted genes may therefore shed light on the drivers leading

to the evolution of genomic imprinting. For instance, functional characterisation of a number of

imprinted genes points to convergence on placental function (12), in line with the predictions of early

theoretical ideas (13). Outside of the placenta, the brain consistently emerges as an adult tissue with a

large number of expressed imprinted genes (14-16). However, given that it is estimated that ~80% of

all genes in the genome are expressed in the brain (17, 18), the question remains, *is* imprinted gene

expression actually enriched in the brain compared to other adult tissues? To date this has never been

formally tested.

A role for imprinted genes in the brain was initially suggested by (19) and neurological phenotypes

have been observed in early imprinted gene mouse models (20), as well as the behavioural deficits seen

in imprinting disorders such as Prader-Willi and Angelman syndromes (21, 22). Subsequent studies

have revealed diverse roles for imprinted genes in the brain. During development, several imprinted

genes are involved in the processes of neural differentiation, migration, axonal outgrowth and apoptosis

(23). In the adult brain, studies of mice carrying manipulations of individual imprinted genes have

suggested a wide range of behavioural roles including maternal care (24), feeding (25), social behaviour

(26, 27), learning/memory (28), cognition (29, 30), and more recently, sleep and circadian activity (31).

In addition to studies on individual imprinted genes, there are a limited number of studies that take a

systems level approach to characterizing the role of genomic imprinting in the brain. Early studies

examining developing and adult chimeras of normal and parthenogenetic/gynogenetic (Pg/Gg - two

maternal genomes) or androgenetic (Ag - two paternal genomes) cells indicated distinct regional

distribution for maternally (cortex and hippocampus) and paternally (hypothalamus) expressed genes

(20, 32). More recently, Gregg, Zhang (16) used the known imprinting status of 45 imprinted genes and

the Allen Brain Atlas to track dichotomous expression of imprinted genes across 118 brain regions to

identify brain-wide patterns of expression. Most imprinted genes were expressed in every brain region,

but detectable expression of the largest number of imprinted genes was found in regions of the

hypothalamus (medial preoptic area, arcuate nucleus), central amygdala, basal nuclei of the stria

terminalis and the monoaminergic nuclei, suggesting some form of specialisation. Although pioneering,

this study, and others identifying novel imprinted genes and/or mapping allelic expression in the brain

(14, 15, 33, 34), did not test whether the expression of these genes was especially enriched in given

brain regions but simply asked if they were expressed, at any level, or not.

Here we address the question of whether the brain and/or specific brain circuitry is a foci for genomic

imprinting by exploiting the rapidly expanding number of single-cell RNA sequencing (scRNA-seq)

datasets and systematically investigating imprinted gene enrichment and over-representation in the

murine brain. We performed this by a hierarchical sequence of data analysis, using datasets that allowed

a multi-organ comparison first, before proceeding to brain-specific and brain region-specific

comparisons with the outcome of each level informing the data selection for the next one, to identify a

consistent pattern of enrichment (Figure 1). We sought to provide a robust assessment of the neural

systems on which imprinted genes converge, statistically validating previous assumptions, identifying

neuronal domains that have received less emphasis in earlier studies, and providing testable predictions

for novel neural and behavioural phenotypes associated with specific genes and imprinted gene

networks.

RESULTS

Imprinted gene expression is enriched in the brain in a multi-organ analysis

The Mouse Cell Atlas (MCA) (35) and the *Tabula Muris* (TM) (36) are single cell compendiums

containing ~20 overlapping, but not identical, adult mouse organs. Key overlapping organs include the

bladder, brain, kidney, lung, limb muscle, and pancreas while organs included in only one dataset

include the ovary, testes, uterus, stomach within the MCA and the heart, fat, skin, trachea and diaphragm

within the TM. These compendiums create a snapshot of gene expression across adult tissues to assess

imprinted gene enrichment. Since this study focused on the adult body and brain, fetal tissues (including

the placenta) were not assessed.

Figure 1. The hierarchical set of datasets in this analysis. The datasets are sorted into multi-organ, whole brain and specific brain nuclei level analyses. The original publication and specific tissue/s analysed are provided for each analysis. White text in dark grey box indicates specifics to the analysis at that level – whether the analysis used the 'marker gene' Log2FC criteria or the relaxed Log2FC > 0 criterion, whether paternally and maternally expressed gene (PEG/MEG) analysis was carried out and whether the top tissue/cell-types and the average normalised expression were reported for imprinted genes.

96 An over-representation analysis (ORA) was performed on both datasets. All data were processed

97 according to the original published procedure, a list of upregulated genes was produced for each

98 tissue/identity group (vs. all other tissue/identity groups) and a one-sided Fisher's Exact test was

Table 1. Imprinted gene over-representation in MCA adult tissues (35). *Identity* – Tissue identities for the cells used in analysis; *Up Reg* – number of upregulated genes with $q \leq 0.05$ and Log2FC ≥ 1 (total number of genes in the dataset in brackets); *IG* – number of imprinted genes upregulated with $q \leq 0.05$ and Log2FC ≥ 1 (total number of IGs in the dataset in brackets); *ORA p* – *p* value from over representation analysis on groups with minimum 5% of total IGs; *ORA q* – Bonferroni corrected *p* value from ORA; *Mean FC IG* – mean fold change for upregulated imprinted genes; *Mean FC Rest* – mean fold change for all other upregulated genes; *Top Tissue IGs* – Sum of IGs with highest mean expression value for that identity group.

Tissue Identity	Up Reg (20,534)	IG (95)	ORA *p*	ORA *q*	Mean FC IG	Mean FC Rest	Top Tissue IGs

Tissue	Up Reg	IG	ORA p	ORA q	Mean FC IG	Mean FC Rest	Top Tissue IGs
Pancreas	2737	42	1.57E-13	**1.89E-12**	8.74	10.32	22
Brain	3401	34	4.43E-06	**5.31E-05**	8.76	125.00	19
Bladder	3183	29	0.000168	**0.002012**	4.45	8.51	8
Uterus	2567	22	0.002827	**0.033919**	4.66	8.46	7
Lung	1203	8	0.192705	1	3.82	151.41	4
Ovary	2219	13	0.223666	1	7.46	11.27	5
Kidney	1714	10	0.268425	1	13.76	182.89	5
Liver	1739	8	0.560145	1	4.55	80.51	3
Stomach	1821	7	0.748590	1	4.24	88.60	3
Thymus	1805	6	0.851579	1	2.78	6.76	2
Small Intestine	1719	5	0.908008	1	7.99	218.64	2
Testis	5212	14	0.995891	1	27.04	5058.36	10
Bone Marrow	1095	2	-	-	5.31	4.43	1
Mammary Gland Virgin	902	4	-	-	3.70	4.03	0
Muscle	1127	4	-	-	8.64	15.05	3
Peripheral Blood	1146	3	-	-	3.78	3.57	0
Prostate	369	0	-	-	0.00	478.10	0
Spleen	1501	1	-	-	4.90	4.77	1

performed using a custom list of imprinted genes (Supplemental Table S1) to identify tissues in which

imprinted genes were over-represented amongst the upregulated genes for that tissue. Each dataset in

this study was analysed independently which allowed us to look for convergent patterns of enrichment

between datasets of similar tissues/cell-types. Across only adult tissues, imprinted genes were

convergently over-represented in the pancreas, bladder and the brain in both datasets (Figure 2A). In

addition, in the MCA adult tissue dataset, there was a significant over-representation in the uterus (Table

1), and in the *Tabula Muris* analysis (Table 2), there was a significant over-representation in the muscle-

based tissues - diaphragm, trachea, and limb muscles. In addition to the ORA, to identify situations in

which imprinted genes were in fact enriched amongst the stronger markers of a tissue/cell-type, we

performed a Gene-Set Enrichment Analysis (GSEA) on tissues meeting minimum criteria (see

Methods), which assessed whether imprinted genes were enriched within the top ranked upregulated

genes for that tissue (ranked by Log2 Fold Change). No tissue at this level showed a significant GSEA

for imprinted genes. Mean normalised expression of imprinted genes across identity groups

(Supplemental Table S2) was the highest for Brain in the MCA and highest for Pancreas in the TM

(Brain (Non-Myeloid) was the fourth highest).

Table 2. Imprinted gene over-representation in *Tabula Muris* adult tissues (36). *GSEA p* – p value from Gene Set Enrichment Analysis for identity groups with 15+ IGs and Mean FC IG > Mean FC Rest; *GSEA q* – Bonferroni corrected p values from GSEA. All other column descriptions can be found in the legend of Table 1.

Tissue Identity	Up Reg (20,839)	IG (107)	ORA p	ORA q	Mean FC IG	Mean FC Rest	GSEA p	GSEA q	Top Tissue IGs
Diaphragm	416	19	3.66E-13	**4.75E-12**	6.49	4.83	0.1898	0.3796	4
Limb Muscle	761	24	6.32E-13	**8.22E-12**	9.02	5.09	0.0552	0.1104	8
Pancreas	4104	43	8.31E-07	**1.08E-05**	12.52	12.60	-	-	29
Trachea	1979	25	1.78E-05	**0.0002**	3.81	4.57	-	-	5
Brain (Non-Myeloid)	3081	31	0.0001	**0.0016**	12.16	14.17	-	-	14
Bladder	3338	31	0.0005	**0.0068**	3.30	5.30	-	-	16
Fat	1263	12	0.0286	0.3713	3.46	3.68	-	-	1

Heart	1108	10	0.0585	0.7601	2.87	5.14	-	-	0
Mammary Gland	1826	12	0.2264	1	3.52	5.24	-	-	3
Liver	1808	7	0.8307	1	6.19	54.93	-	-	3
Aorta	3515	14	0.8832	1	7.47	16.08	-	-	2
Tongue	4295	15	0.9696	1	4.15	7.16	-	-	8
Large Intestine	4758	11	0.9998	1	5.95	12.22	-	-	5
Brain (Myeloid)	1024	5	-	-	3.39	6.80	-	-	2
Kidney	584	3	-	-	24.94	22.90	-	-	1
Lung	914	2	-	-	2.73	5.41	-	-	0
Marrow	1957	5	-	-	7.65	5.25	-	-	4
Skin	1612	4	-	-	4.15	8.36	-	-	1
Spleen	625	1	-	-	4.63	4.28	-	-	0
Thymus	678	4	-	-	3.46	7.45	-	-	1

114 Given the interest in the different functions of maternally expressed genes (MEGs) and paternally

115 expressed genes (PEGs), we additionally ran the large-scale enrichment analyses (Levels 1 and 2) using

116 separate lists of PEGs and MEGs. At Level 1, MEGs and PEGs (Supplemental Table S3A, S3B, S4A

117 and S4B) revealed a similar pattern of enrichment in both datasets (Fig. 2). PEGs were over-represented

Figure 2. Level 1 multi-organ comparison summary graphics. (*A*) Venn diagram of upregulated imprinted genes in the brain in Mouse Cell Atlas and in the brain (non-myeloid) in the Tabula Muris. Imprinted genes are listed which show significant upregulation ($q \leq 0.05$ and Log2FC ≥ 1) in the tissues. Although these tissues are not identical, these were the two brain associated over-representations in the enrichment analysis. Parental-bias is indicated by colour (MEG - red, PEG - blue). From the 119 imprinted genes in the gene list, only 92 were common to both analyses (i.e., successfully sequenced and passed gene quality control filters). 34 imprinted genes were upregulated in the brain in the MCA and 31 genes in the *TM*. Genes in common from the two analyses are presented in bold and totalled in each section of the Venn Diagram, while genes found upregulated in one analysis but not available in the other analysis are included in small font and the number indicated in brackets. (*B*) Tissues with over-representation in MCA. Coloured tissues with bold labels were over-represented tissues using all imprinted genes, tissues with a blue circle behind were over-represented for PEGs alone, a red circle represent the same for MEGs, and a red/blue split circle were over-represented for both PEGs and MEGs. (*C*) Tissues with over-representation in *Tabula Muris*. See 2b description for details.

in the brain in both datasets (MCA - $q = 4.56 \times 10^{-6}$, *TM* - $q = 0.0005$) while MEGs were not. PEGs were

also over-represented in the diaphragm ($q = 0.0007$), limb muscle ($q = 0.0001$) and pancreas (MCA -

$q = 1.93 \times 10^{-5}$, *TM* - $q = 0.0002$), with a significant GSEA in the MCA pancreas ($p = 0.02$, Supplemental

Fig. S1). While MEGs were over-represented in the bladder (MCA - $q = 0.002$, *TM* - $q = 0.020$), the

pancreas (MCA - $q = 1.53 \times 10^{-7}$) and in the three muscular tissues of the *Tabula Muris* (diaphragm - q

$= 2.13 \times 10^{-8}$, limb muscle - $q = 2.43 \times 10^{-7}$, trachea - $q = 0.004$).

Imprinted gene expression is enriched in neurons and neuroendocrine cells of the brain

We next analysed cells from the whole mouse brain (Level 2), first analysing Ximerakis, Lipnick (6)

dataset, in which cells were grouped from the whole mouse brain (minus the hindbrain) into major cell

classes according to cell lineage. Imprinted genes were over-represented in neuroendocrine cells and

mature neurons (Table 3).

Neuroendocrine cells were defined as a heterogeneous cluster, containing peptidergic neurons and

neurosecretory cells expressing neuronal marker genes (e.g., *Syt1* and *Snap25*) alongside neuropeptide

genes (e.g., *Oxt*, *Avp*, *Gal*, *Agrp* and *Sst*) but distinguished by Ximerakis, Lipnick (6) by the unique

expression of *Baiap3* which plays an important role in the regulation of exocytosis in neuroendocrine

cells (37). GSEA additionally showed that the imprinted genes were enriched in the genes with the

highest fold change values for neuroendocrine cells only (Fig. 3). 26 imprinted genes had their highest

expression in the neuroendocrine cells and the mean normalised expression of imprinted genes was

almost twice as high for neuroendocrine cells as the next highest identity group (Supplemental Table

S2). The MEG/PEG analysis (Supplemental Table S5A and S5B) for this dataset found that PEGs were

over-represented in mature neurons ($q = 0.027$) and neuroendocrine cells ($q = 8.97 \times 10^{-6}$). MEGs were

also over-represented in neuroendocrine cells ($q = 0.047$) and uniquely over-represented in Arachnoid

barrier cells ($q = 0.014$). Only PEGs replicated the significant GSEA in neuroendocrine cells ($p = 4 \times 10^{-4}$

, Supplemental Fig. S2).

Table 3. Imprinted gene over-representation in neural lineage types (6). *Identity* – Cell lineage identities for the cells used in analysis. All other column descriptions can be found in the legend of Tables 1 & 2.

Cell Population Identity (Abbr.)	Up Reg (14,498)	IG (85)	ORA *p*	ORA *q*	Mean FC IG	Mean FC Rest	GSEA *p*	GSEA *q*	Top Cell IGs
Neuroendocrine cells (NendC)	3868	47	2.12E-08	**3.82E-07**	11.88	5.42	0.0017	**0.0051**	26

Mature Neurons (all types) (mNEUR)	2968	32	0.0002	**0.0035**	8.80	9.28	-	-	2
Arachnoid barrier cells (ABC)	2287	20	0.0396	0.7120	16.84	22.63	-	-	7
Tanycytes (TNC)	1279	12	0.0692	1	6.64	12.01	-	-	8
Vascular and leptomeningeal cells (VLMC)	1714	15	0.0724	1	15.06	13.03	0.0468	0.1404	4
Oligodendrocyte precursor cells (OPC)	1524	13	0.1067	1	3.03	7.17	-	-	1
Pericytes (PC)	1801	14	0.1649	1	8.20	8.22	-	-	2
Olfactory ensheathing glia (OEG)	1086	9	0.1848	1	7.95	26.03	-	-	1
Oligodendrocytes (OLG)	1183	9	0.2561	1	3.73	12.91	-	-	5
Choroid plexus epithelial cells (CPC)	2602	17	0.3524	1	7.43	19.34	-	-	5
Hemoglobin-expressing vascular cells (Hb_VC)	1798	11	0.4889	1	5.25	6.33	-	-	3
Vascular smooth muscle cells (VSMC)	3006	17	0.6093	1	8.94	6.71	0.1376	0.4128	5
Astrocyte-restricted precursors (ARP)	1445	8	0.6214	1	4.50	5.09	-	-	1
Neural stem cells (NSC)	1009	5	0.7138	1	4.00	4.09	-	-	0
Ependymocytes (EPC)	3233	17	0.7346	1	15.04	53.27	-	-	4
Endothelial cells (EC)	1455	7	0.7619	1	5.80	8.54	-	-	0
Hypendymal cells (HypEPC)	1525	6	0.8946	1	17.24	20.80	-	-	5
Neuronal-restricted precursor (NRP)	2339	10	0.8979	1	3.07	10.20	-	-	1
Astrocytes (ASC)	1384	4	-	-	2.22	6.04	-	-	0
Dendritic cells (DC)	1209	1	-	-	3.50	16.02	-	-	1
Immature Neurons (ImmN)	652	4	-	-	3.37	5.79	-	-	0
Macrophages (MAC)	1222	2	-	-	3.47	21.56	-	-	0
Microglia (MG)	1342	3	-	-	19.28	19.22	-	-	3
Monocytes (MNC)	947	2	-	-	16.49	19.13	-	-	1
Neutrophils (NEUT)	519	2	-	-	9.18	62.13	-	-	0

The second dataset at this level was Zeisel, Hochgerner (1) Mouse Brain Atlas (MBA) and it allowed a

much deeper investigation of nervous system enrichment with sequencing of the entire murine nervous

system and identifying cells by both brain region and cell type. Concordant with the previous findings,

primary analysis separating cells by lineage revealed over-representation of imprinted genes in neurons

only (Table 4). The overlap between the upregulated imprinted genes for the over-represented neural-

lineage cells from the Level 2 datasets are displayed in Figure 4. Additionally, PEGs alone demonstrated

no significant over-representations in cell lineage types while MEGs demonstrated over-representation

in vascular cells only ($q = 0.0004$) (Supplemental Table S6A and S6B).

Table 4. Imprinted gene over-representation in nervous system cell types (1). *Identity* – Cell identities for the cells used in analysis. All other column descriptions can be found in the legend of Table 1.

Cell Population Identity	Up Reg (19,547)	IG (108)	ORA *p*	ORA *q*	Mean FC IG	Mean FC Rest	Top Cell IGs
Neurons	5710	44	0.0081	**0.0487**	11.73	24.97	45
Vascular	2473	22	0.0171	0.1029	17.91	26.64	16
Oligos	1587	11	0.2701	1	4.64	11.48	12
Peripheral Glia	2820	16	0.5117	1	5.42	12.64	12
Ependymal	3683	20	0.5912	1	24.52	66.97	15
Immune	1564	7	0.7787	1	13.42	93.05	5
Astrocytes	1539	4	-	-	2.88	10.73	3

150   **The hypothalamus, ventral midbrain, pons and medulla are enriched for imprinted gene**

151   **expression**

152   After confirming neuron-specific enrichment of imprinted genes in the MBA dataset, further MBA

153   analysis was performed on cells classified as neurons and then grouped by brain/nervous system

Figure 3. GSEA and dot plots for imprinted genes upregulated in neuroendocrine cells in the Ximerakis, Lipnick (6) whole mouse brain dataset. (*A*) GSEA for imprinted genes upregulated in the neuroendocrine cells. In the analysis, genes are sorted by strength by which they mark this neuronal cluster (sorted by Log2FC values) indicated by the bar (bottom). The genes are arrayed left (strongest marker) to right and blue lines mark where imprinted genes fall on this array. The vertical axis indicates an accumulating weight, progressing from left to right and increasing or decreasing depending on whether the next gene is an imprinted gene or not. The *p*-value represents the probability of observing the maximum value of the score (red dashed line) if the imprinted genes are distributed randomly along the horizontal axis. The *q*-value for this analysis was significant at 0.0036. (*B*) Dot plot of imprinted genes upregulated in the 'Neuroendocrine cells' plotted across all identified cell types (Abbr. in Table 3). Imprinted genes were plotted in chromosomal order. Size of points represented absolute mean expression; colour represented the size of the Log2FC value for the cell identity group (e.g., neuroendocrine cells) vs. all other cells. Unique colour scales are used for MEGs (red/orange) and PEGs (blue).

154   regions. Significant over-representation was seen in neurons of the hypothalamus, ventral midbrain,

155   medulla, and pons (Table 5). The pons and medulla had the largest number, 45 and 44 respectively, of

156   imprinted genes upregulated (Figure 5A).

Table 5. Imprinted gene over-representation in nervous system region(1). *Identity* – Nervous system regional identities for the cells used in analysis genes. All other column descriptions can be found in the legend of Tables 1 & 2.

Brain Region Identity	Up Reg (18,335)	IG (105)	ORA *p*	ORA *q*	Mean FC IG	Mean FC Rest	GSEA *p*	GSEA *q*	Top Region IGs
Medulla	3147	45	8.38E-10	**1.26E-08**	4.79	4.01	0.1	0.2	15
Hypothalamus	1040	22	9.81E-08	**1.47E-06**	4.92	5.84	-	-	8
Pons	3581	44	1.62E-07	**2.43E-06**	4.20	3.91	0.1169	0.2338	22
Vent. Midbrain	1228	18	0.0002	**0.0034**	4.90	4.99	-	-	3
Vent. Striatum	689	8	0.0463	0.6941	3.92	4.92	-	-	0
Posterior Cortex	1090	9	0.1788	1	2.64	3.20	-	-	2
Enteric Nervous System	3885	26	0.2311	1	8.98	121.04	-	-	11
Sympathetic Nervous System	2804	18	0.3535	1	11.37	57.96	-	-	9
Anterior Cortex	979	6	0.5016	1	2.72	3.30	-	-	1
Dors. Midbrain	1045	6	0.5663	1	2.20	4.85	-	-	3
Thalamus	1441	8	0.6000	1	2.90	6.36	-	-	0
Hippocampus - CA1	1082	6	0.6008	1	3.01	4.02	-	-	2
Somatosensory Cortex	2121	11	0.6943	1	4.09	3.70	-	-	8
Dors. Striatum	1196	6	0.6974	1	4.03	5.43	-	-	2
Dorsal Root Ganglion	3607	16	0.9088	1	11.56	75.89	-	-	9
Middle Cortex	623	5	-	-	3.29	3.24	-	-	0
Spinal Cord	972	5	-	-	4.57	12.36	-	-	1
Amygdala	452	4	-	-	4.65	4.11	-	-	2
Dentate Gyrus	796	4	-	-	3.79	4.16	-	-	2
Hippocampus	631	4	-	-	4.86	3.82	-	-	2
Olfactory Bulb	445	4	-	-	4.02	8.27	-	-	2
Antero-Middle Cortex	646	3	-	-	4.59	4.31	-	-	1
Cerebellum	240	0	-	-	0.00	32.30	-	-	0

Figure 4. Venn diagram of upregulated imprinted genes in the mature neuronal cells in the whole brain datasets of Zeisel, Hochgerner (1) and Ximerakis, Lipnick (6). Imprinted genes are listed which show significant upregulation ($q \leq 0.05$ and Log2FC ≥ 1) in the cells. Although these cell types are not identical, these were all mature neural lineage cells with over-representations in the enrichment analysis. Parental-bias is indicated by colour (MEG - red, PEG - blue. From the 119 imprinted genes in the gene list, only 88 were common to both analyses (i.e., successfully sequenced and passed gene quality control filters). 45 imprinted genes were upregulated in neurons in the MBA, and in Ximerakis, Lipnick (6), 33 imprinted genes were upregulated in neurons and 48 genes in neuroendocrine cells. Genes in common from the two analyses are presented in bold and totalled in each section of the Venn Diagram, while genes found upregulated in one analysis but not available in the other analysis are included in small font and the number indicated in brackets.

Regional analysis for MEGs and PEGs separately (Supplemental Table S7A and S7B), revealed that

PEGs were over-represented in hypothalamus ($q = 6.53 \times 10^{-7}$), ventral midbrain ($q = 0.018$), the pons (q

$= 4.65 \times 10^{-5}$) and the medulla ($q = 4.10 \times 10^{-6}$); while MEGs were only over-represented in the medulla

($q = 0.002$) but had a significant GSEA for the pons ($q = 0.027$, Supplemental Fig. S3); see Figure 5B.

Neurons were then recategorized into unique subpopulations identified by marker genes (1) to uncover

the specific neural populations underlying the enrichment seen in the hypothalamus, pons and medulla,

and midbrain (Fig. 6; Supplemental Table S8). Each neural population was identified by its distinct

Figure 5. Level 2ii) brain region analysis summary figures. (*A*) Venn diagram of upregulated imprinted genes in the neurons of enriched nervous system regions from the Mouse Brain Atlas (1). Imprinted genes are listed which show significant upregulation ($q \leq 0.05$ and Log2FC ≥ 1) in the regions specified. The number of imprinted genes in each region of the Venn diagram are specified. Parental-bias of imprinted genes is indicated by colour (MEG - red, PEG - blue). (*B*) Brain regions enriched for imprinted gene expression via ORA or GSEA in the MBA (1). Regions over-represented for all imprinted genes are bolded. Regions over-represented for PEG expression alone are coloured blue while regions enriched for MEG expression alone are coloured red.

164 gene expression and suspected location within the brain (see http://mousebrain.org/ for an online

165 resource with detailed information on each cluster).

166 The hypothalamus was represented by a selection of inhibitory and peptidergic neurons. Inhibitory

167 neurons with over-representation of imprinted genes included: Deinh3 (notable genes *Lhx8, Gabrq*)

168 representing a Subthalamic Nucleus population, Deinh5 (*Nts*, *Dlk1*) and Teinh3 (*Gal, Irs4*)

169 representing Preoptic Area/ BNST populations, Deinh6 (*Agrp, Otp*) representing an Arcuate nucleus

170 population, and Deinh7 (*Avp, Rgs16, Nms*) and Deinh8 (*Six6, Nms, Vip*) representing Suprachiasmatic

171 nucleus populations. For peptidergic neurons, over-representation was seen in HYPEP3 (*Gpr101,*

172 *Tac1, Baiap3*) a ventromedial population, HYPEP2 (*Otp, Trh, Ucn3*) a ventromedial/paraventricular

173 population, HYPEPE1 (*Trh, Otp, Ngb*) a lateral hypothalamic population, HYPEP4 (*Oxt, Otp*) an

174 oxytocin magnocellular population of the paraventricular and supraoptic nuclei, and HYPEP6 (*Hcrt,*

175 *Pdyn, Trhr*) an orexin producing population of the dorsomedial/lateral hypothalamus.

176 The midbrain, medulla and pons were represented by a number of cell groups, with over-representation

177 seen in the medulla-based adrenergic (HBAR) and noradrenergic (HBNOR) groups and the

178 dopaminergic neurons of the midbrain in the Periaqueductal Gray (PAG) (MBDOP1) and the Ventral

Figure 6. Anatomical labelling of all the neural subpopulations with a significant over-representation of imprinted genes ($q \leq 0.05$ and Log2FC ≥ 1) in the Mouse Brain Atlas (1). The predicted brain nuclei localisation of the 32 neuronal subpopulations (out of 214 populations identified across the nervous system) specified in the MBA and enriched for imprinted genes. Brain regions that were not found to be enriched for imprinted genes are greyed out. The full Enrichment Analysis is available in Supplemental Table S8.

Tegmental Area (VTA)/Substantia Nigra (SNc) (MBDOP2). There were also several inhibitory

(MEINH, HBIN) and excitatory neuron (MEGLU, HBGLU) types spread across the nuclei from the

three regions (Fig. 6). The serotonergic populations of the raphe nuclei of these regions (HBSER) were

particularly prominent since the pons and medulla-based serotonin neuron populations (HBSER2,

HBSER4 and HBSER5) were the only neuron subpopulations out of the 214 total to have a significant

GSEA for imprinted genes after correction (Supplemental Fig. S4).

Additional regions of over-representation included TEINH1&2 representing neurons in the pallidum

and striatum and DEINH4 representing PVN neurons from the thalamus. In this comparison of 214

neuron populations, no neurons from areas such as the cortex, cerebellum or peripheral nervous system

were enriched, and neither were they over-represented in the previous regional analysis. Hence, further

analysis focused on those brain regions enriched in this whole brain level analysis.

190 Imprinted gene expression is over-represented in specific hypothalamic neuron subtypes

We next sought to investigate whether those regional neuron enrichments found within the whole brain

comparisons would be further clarified with enriched expression in specific neuronal subpopulations

within those regions. Namely within the hypothalamus (to identify neural populations enriched across

the whole hypothalamus and those enriched within specific hypothalamic nuclei) and whether imprinted

gene expression was enriched in specific dopaminergic and serotonergic populations, located in the

ventral midbrain and hindbrain, which were found as key subpopulations in the whole brain analysis.

Two datasets with single cell sequencing data for the adult hypothalamus existed (3, 4). Both clustered

their data into neuronal subpopulations allowing us to look for convergent imprinted enrichment across

major hypothalamic neuronal subtypes. Analysis revealed a clear neuronal bias in expression of

imprinted genes (Supplemental Table S9A and S10A). Within the Romanov, Zeisel (3) data, there was

a significant over-representation of imprinted genes in neurons ($q = 0.02$) and a similar observation was

seen in the Chen, Wu (4) data ($q = 0.001$), and both also demonstrated a significant GSEA in neurons

(Fig. 7A-D, Romanov, Zeisel (3) – $p = 0.011$, Chen, Wu (4) - $p = 0.022$).

Within the Chen, Wu (4) dataset, 4/33 hypothalamic neuronal subtypes had a significant over-

representation of imprinted genes (Supplemental Table S9B). The four subtypes were all GABAergic

Figure 7. GSEA and Dot plots for imprinted genes upregulated in neurons across the whole hypothalamus. (*A*) GSEA for imprinted genes upregulated in the 'Neuron' cell type in the whole hypothalamic dataset of Chen, Wu (4). See legend of Figure 3A for a description of how to interpret the plot. (*B*) Dot plot of imprinted genes upregulated in the 'Neuron' cell type plotted across all identified cell types in the Chen, Wu (4) whole hypothalamic dataset. See legend of Figure 3B for a description of how to interpret the plot. Abbr: OPC = Oligodendrocyte Precursor Cell, MG = Myelinating Oligodendrocyte, IMG = Immature Oligodendrocyte, Astro = Astrocyte, Epith = Epithelial, Macro = Macrophage, Tany = Tanycyte, Ependy = Ependymocyte, Micro = Microglia, POPC = Proliferating Oligodendrocyte Progenitor Cell. (*C*) GSEA for imprinted genes upregulated in 'neurons' in the whole hypothalamic dataset of Romanov, Zeisel (3). See legend of Figure 3A for a description of how to interpret the plot. (*D*) Dot plot of imprinted genes upregulated in 'neurons' plotted across all identified cell types in the Romanov, Zeisel (3)whole hypothalamic dataset. See legend of Figure 3B for a description of how to interpret the plot.

neurons, specifically: GABA13:*Slc18a2/Gal* (*q* = 0.0079), representing Galanin neurons (present in a

several hypothalamic regions); GABA17:*Slc6a3* (*q* = 0.0001) a dopaminergic neuron type with high

expression of *Th* and *Prlr*, which most likely corresponds to the TIDA neurons of the arcuate nucleus;

GABA8:*Vipr2* (*q* = 0.0071) with very high *Avp* and *Nms* expression suggesting these are neurons from

the SCN; and GABA15:*Agrp* (*q* = 0.034) feeding promoting neurons of the Arcuate Nucleus. Within

the Romanov, Zeisel (3) dataset, 3/62 subtypes had significant over-representation of imprinted gene

expression (Supplemental Table S10B): GABA14:*Agrp*/*Npy* (*q* = 0.013) which were the Arcuate

Nucleus feeding neurons also reported in Chen, Wu (4); a *Ghrh/Th* neuronal type (*q* = 0.032), again

likely corresponding to neurons from the arcuate nucleus and the top hit was GABA5:*Calcr/Lhx1* (*q* =

1.63×10^{-6}) but this was a poorly segregated population likely due to a deeper inner cluster heterogeneity.

This cluster was interesting since the imprinted genes *Calcr* and *Asb4* were amongst its most significant

marker genes, and it was notably the only cluster with high expression of all three of *Th*, *Slc6a3* and

Prlr. Romanov, Zeisel (3) did not identify any of their populations as the TIDA neurons, but the above

pattern of gene expression suggests that GABA5 may contain these neurons. Furthermore,

GABA17:*Slc6a3* from the Chen, Wu (4) dataset shared 21/40 upregulated genes of GABA5 (see

Supplemental Table S11 for full comparison);

Having consistently found well-known neurons from the arcuate nucleus (*Agrp, Ghrh*), and

suprachiasmatic nucleus (*Avp, Vip*) we sought to test imprinted gene enrichment within these

hypothalamic regions at a high resolution using datasets sequencing neurons purely from these

hypothalamic regions.

Arcuate nucleus (ARC) (2)

The first nuclei investigated was the ARC sequenced by Campbell, Macosko (2). Imprinted gene over-

representation was found in 8/24 arcuate neuron types (Supplemental Table S12). These included the

Agrp/*Sst* neuron type (with high expression of *Npy, q* = 0.003) and two *Pomc* neuron types

(*Pomc/Anxa2*, *q* = 0.004; *Pomc/Glipr1*, *q* = 0.03). Pomc expressing neurons are known to work as

feeding suppressants (38). Additional significant over-representation was found in the *Ghrh* neuron

type (*q* = 0.009), which was also enriched in *Gal* and *Th*. Finally, a highly significant over-

representation of imprinted genes was found in the *Th*/*Slc6a3* neuron type (*q* = 1.72 x 10^{-8}) identified

by the authors as one of the most likely candidates for the TIDA dopaminergic neuron population.

Marker genes for this identity group overlapped with the TIDA candidates from the previous two

datasets (e.g., *Slc6a3*, *Th*, *Lhx1*, *Calcr*). *Agrp* neurons, *Ghrh* neurons and these TIDA candidate neurons

were identified in both whole hypothalamic datasets and at the nuclei level.

Suprachiasmatic Nucleus (SCN) (5)

Analysis of the 10x chromium data of SCN neurons (Supplemental Table S13) revealed a significant

over-representation (*q* = 1.51 x 10^{-8}) and GSEA (*p* = 0.004, Supplemental Fig. S5) in the *Avp/Nms*

neuronal cluster (out of 5 neuronal clusters). This cluster shows the strongest expression for *Oxt, Avp*,

Avpr1a and *Prlr* and is one of the three neural group that Wen, Ma (5) found had robust circadian gene

expression, and the only subtype with notable phase differences in circadian gene expression in the

dorsal SCN. This cluster likely

corresponds to the GABA8

cluster found enriched in the

Chen, Wu (4) dataset. Figure 8

presents the overlapping

upregulated imprinted genes

from the convergently

upregulated neuron subtypes

in the hypothalamic

analysis of Level 3a and

3b.

Figure 8. Venn diagrams of upregulated imprinted genes in the neuronal subpopulations from level 3b that were also identified in level 2 and 3a. Imprinted gene overlap was contrasted for *Agrp/Sst* neuronal populations of the Arcuate Nucleus (1-4) and *Avp/Nms* neurons from the Suprachiasmatic Nucleus (1, 4, 5) Imprinted genes are listed which show significant upregulation (*q* ≤ 0.05 and Log2FC > 0) in the subpopulation. Parental-bias is indicated by colour (MEG - red, PEG - blue).

Imprinted gene expression is over-represented in monoaminergic nuclei of the mid- and hindbrain

In the MBA, Whole Hypothalamus and Arcuate Nucleus analyses, dopaminergic clusters were consistently enriched and, to explore this further, analysis of Hook, McClymont (39) data allowed comparison for dopamine neurons across the brain (specifically from the olfactory bulb, arcuate nucleus and midbrain) at two developmental timepoints (E15.5 and Post-natal day (P) 7). The arcuate nucleus P7 dopamine neurons emerged as the clearest over-represented subgroups (Supplemental Table S14). This included the *Th/Slc6a3/Prlr* neurons ($q = 1.15\times10^{-8}$) and the *Th/Ghrh/Gal* cluster ($q = 4.79\times10^{-5}$) the latter of which were referred to as 'neuroendocrine' cells by Hook, McClymont (39), and the former a mixture of arcuate nucleus populations with *Prlr* was one of the marker genes, suggesting this includes the TIDA neurons. Additionally, P7 midbrain neurons were the other group with significant over-representation (specifically from the PAG and VTA) as well as the neuroblasts at this time point.

Although no specific adult mouse midbrain datasets exist, ventral midbrain sequencing at E11.5 - E18.5 by La Manno, Gyllborg (40) allowed us to identify imprinted enrichment within the midbrain at a timepoint when the major neuronal populations are differentiating but still identifiable (Supplemental Table S15). As anticipated, we found significant over-representation in both mature (DA1; high *Th* and *Slc6a3,* $q = 0.0103$), and developing (DA0, $q = 0.0129$) dopaminergic neurons, as well as the serotonergic neurons ($q = 3.09\times10^{-7}$), likely from the midbrain raphe nuclei.

Raphe nuclei from the midbrain/hindbrain are key serotonergic regions of the brain. Analysis of all cell types in the Dorsal Raphe Nucleus (DRN) sequenced by Huang, Ochandarena (41) revealed a clear enrichment of imprinted genes in the neuronal populations of the DRN as compared to the non-neuronal cell populations of the DRN (Supplemental Table S16A). When compared to all other cell populations, significant ORA was seen for Dopaminergic ($q = 0.009$), Serotonergic ($q = 0.012$) and Peptidergic neurons ($q = 0.0008$), however, a significant GSEA was found for all five neuronal populations (Supplemental Fig. S6). When compared against each other (i.e., serotonergic upregulation vs. the other neurons), only the serotonergic neurons of the DRN ($q = 0.0019$) were found to have a significant over-representation of imprinted genes (Supplemental Table S16B). GSEA's were non-significant but the

mean fold change for imprinted genes was markedly higher in both serotoninergic (52% higher) and

dopaminergic neurons (68% higher). When contrasting neuronal subpopulations of the DRN, two of

the five serotonin subpopulations had significant over-representation of imprinted genes: *Hcrtr1/Asb4*

($q = 0.0014$) and *Prkcq/Trh* ($q = 0.007$) (Supplemental Table S16C). These clusters were identified by

Huang, Ochandarena (41) as the only clusters localised in the dorsal/lateral DRN and the serotonin

clusters enriched in *Trh*. Huang, Ochandarena (41) hypothesised that these were the serotonin neurons

that project to hypothalamic nuclei, and motor nuclei in the brainstem (as opposed to cortical/striatal

projection).

Imprinted gene expression is over-represented in lactotrophs and somatotrophs of the pituitary

gland

Following on from the enrichment seen above for imprinted gene expression in the dopaminergic

arcuate nucleus neurons

coordinating pituitary

gland output, we sought

to identify whether any

cells in the pituitary

would display matching

over-representation for

imprinted gene

expression. The

pituitary was not

sequenced as part of the

multi-organ or whole

brain datasets analysed

above and so two

independent datasets

were analysed that

Figure 9. *(A)* Pituitary cell types showing over-representation for imprinted gene expression in multiple pituitary datasets. Over-represented cell types are bold and not in greyscale. The hormone/s released from the endocrine cell types are also indicated. *(B)* Venn diagram of upregulated imprinted genes in the Somatotrophs and Lactotrophs in Cheung, George (7) and Ho, Hu (8). Imprinted genes are listed which show significant upregulation ($q \leq 0.05$ and Log2FC > 0) in the cell types. Parental bias is indicated by colour (MEG –red, PEG – blue). Genes in common from two analyses are presented in bold and totalled in each section of the Venn Diagram, while genes found upregulated in one analysis but not available in the others are included in small font and the number indicated in brackets.

specifically sequencing the mouse pituitary at single cell resolution. Ho, Hu (8) recently sequenced the

anterior pituitary gland of male and female C57BL/6 mice using two sequencing technologies, both

10X genomic and Drop-Seq. This identified a variety of cell types from the endocrine and non-

endocrine pituitary. We analysed data from both technologies and found that imprinted gene expression

was convergently over-represented in the Lactotrophs (prolactin secreting) and Somatotroph (growth

hormone secreting) cells (Supplemental Table S17A & 17B). In a second independent dataset

sequencing cells from male mouse pituitary glands (7), we found significant over-representation in the

Somatotropes and Thyrotrope (secreting thyroid stimulating hormone). Figure 9 demonstrates the

overlap in imprinted genes significantly expressed in Somatotropes and Lactotropes across the datasets

since these were the only cell-types to be over-represented in more than one dataset (Supplemental

Table S18). It is notable that the two cell types represented here directly match the two regulatory

neurons found over-represented in the arcuate nucleus of the hypothalamus.

DISCUSSION

Using publicly available single cell transcriptomics data, we apply an unbiased systems biology

approach to examine the enrichment of imprinted genes at the level of the brain in comparison to other

adult tissues, refining this analysis to specific brain regions and then to specific neuronal populations.

We confirm a significant over-representation in the brain, with imprinted genes over-represented in

neurons at every level tested and a marked enrichment in neuroendocrine cells lineages. Within-brain

analyses revealed that the hypothalamus and the monoaminergic system of the mid- and hindbrain were

foci for imprinted gene enrichment. While not all imprinted genes follow these patterns of expression,

these findings highlight collective gene expression which is non-random in nature. As such, these

analyses identify 'expression hotspots', which in turn suggest 'functional hotspots'. Specifically, our

results at the systems and cellular level highlight a major role for imprinted genes in the neuronal

regulation of pituitary function, feeding and sleep.

Some of the earliest studies of genomic imprinting identified the brain as a key area for imprinted gene

expression (20, 32). However, it is estimated that ~80% of the genome is expressed in the brain and

consequently, imprinted gene expression here may not be a purposeful phenomenon. Our current

analysis definitively show that imprinted genes were significantly over-represented in the brain as a

whole. This over-representation was found again with PEGs alone, but not MEGs. Within specific brain

regions, imprinted genes were over-represented in the hypothalamus, ventral midbrain, pons and

medulla. This confirms some previous findings from studies of Pg/Gg and Ag chimera studies (20, 32)

and summaries of imprinted gene expression (16). However, unlike these earlier studies, our analyses

do not simply ask if imprinted genes are expressed (at any level) or not, but robustly test whether this

expression is meaningful, and the expression of these genes are especially enriched in any given brain

region. Additionally, in the chimera studies, Pg/Gg cells with two maternal genomes preferentially

allocated to the developing adult cortex and hippocampus, and Ag cells with two paternal genomes

preferentially allocated to the developing hypothalamus and midbrain. Our analysis does not reproduce

this distinct pattern of MEG and PEG expression in the brain, and indeed we find no specific enrichment

of imprinted genes in cortex or hippocampus. Although the pattern of regional enrichment seen with all

imprinted genes is replicated when analysing PEGs alone, separate analysis of MEGs only shows over-

representation in the pons and medulla. This difference between our analysis of enrichment and the

Pg/Gg and Ag chimeras studies indicate that the distribution of Pg/Gg and Ag cells in the brain is not

driven by, or indeed reflective of adult PEG and MEG expression, but instead is probably determined

by expression of specific imprinted genes during brain development (42).

At the whole brain level, mature neurons and, in particular, neural-lineage neuroendocrine cells had

disproportionately higher numbers of imprinted genes expressed, and high levels of imprinted gene

expression. It is likely that this neural-lineage neuroendocrine population comprises members of the

key hypothalamic populations in which the expression of imprinted genes are enriched and, when

treated as their own cluster, demonstrate strong imprinted gene enrichment compared to other cell

lineages of the brain, even other mature neurons.

Within the hypothalamus, a selection of informative neuronal subpopulations were over-represented.

Strikingly, and suggestive of meaningful enrichment, we saw convergence across our different levels

of analysis with several key neuronal types identified in the whole hypothalamus and/or hypothalamic-

region-level analysis, already having been identified against the background of general imprinted gene

expression in the whole-brain-level analysis. These subpopulations are collectively associated with a

few fundamental motivated behaviours. We consistently saw enriched imprinted gene expression in *Agrp* expressing neurons when contrasting neurons across the whole brain, whole hypothalamus and within the arcuate nucleus. *Agrp* neurons from the arcuate nucleus are well known feeding promotors and a few imprinted genes have previously been associated with their function (*Asb4, Magel2, Snord116*) (43, 44) but never as an enriched population. Feeding was further linked with imprinting through enrichment seen in *Pomc+* neurons (45) as well as *Hcrt+* and *Gal+* neurons. Circadian processes are controlled principally by the Suprachiasmatic Nucleus and here we find strong imprinted gene enrichment in *Avp/Nms* expressing neurons (an active circadian population). These neurons were found enriched again when contrasting neurons across the whole brain, whole hypothalamus and within the SCN. This population is of interest given the growing appreciation of the role imprinted genes play in circadian processes and the SCN suggested by studies of individual imprinted genes (46). Pituitary endocrine regulation also emerged as a key function, considering the over-representation in the dopaminergic: *Th/Slc6a3/Prlr* neuron type (top hit in the arcuate nucleus and across dopaminergic neurons of the brain) and the *Th/Ghrh* subpopulation. These neuron populations can regulate prolactin (regulating lactation, stress, weight gain, parenting and more (47, 48)) and growth hormone (promoting growth and lipid/carbohydrate metabolism) release, respectively. Remarkably, we also found a matching enrichment in the lactotroph and somatotroph cells in the pituitary. A role for imprinted genes in pituitary function is well known (49, 50) with pituitary abnormalities associated with imprinted disorders such as PWS (51) and recent sequencing work showing imprinted genes are amongst the highest expressed transcripts in the mature and developing pituitary (52). Specific genes we found highly expressed here, such as Dlk1 and Nnat, have been shown to alter somatotroph phenotypes (53, 54). Finally, we saw enrichment in galanin expressing neuronal populations (found enriched when contrasting neurons across the whole brain, whole hypothalamus). Galanin neurons in the hypothalamus have a diverse set of functions including subpopulations for thermoregulation, feeding, reproduction, sleep and parenting behaviour (55, 56), contributing to this consistent picture of IGs associating with neurons key for motivated behaviour.

In this analysis the hypothalamus was a clear hot spot for imprinted gene expression, in line with the prevailing view of imprinted gene and hypothalamic function (50, 57). However, outside of the

hypothalamus other distinct hotspot emerged from our whole brain analysis including the

monoaminergic system of the midbrain/hindbrain. Analysing data from the dorsal raphe nucleus and

ventral midbrain revealed the dopaminergic and serotonergic neurons to be a foci of imprinted gene

expression within this region. These midbrain dopamine neurons were enriched when contrasted to

other dopamine neurons from the brain and the enriched serotonergic neurons were those that project

to the subcortical regions of the brain known to be associated with feeding and other motivated

behaviours (58), providing convergence with the functional hotspots seen in the hypothalamus.

Analyses of these kind are always bound by the available data and therefore there are notable limitations

and caveats to this study. The aim of this study was to generate information about 'hotspots' of

imprinted gene expression. This approach, and the use of over-representation analysis and GSEA,

therefore do not provide an exhaustive list of sites of expression. For example, oxytocin neurons of the

hypothalamus were not significantly enriched in any of the hypothalamic datasets despite a clear

oxytocin neuron phenotype reported in a handful of imprinted gene models (24, 59). This may be an

example of a functional effect occurring below the level of over-representation, or that imprinted genes

act during development and are not functionally enriched in adult oxytocin neurons, or simply that

compared to other hypothalamic neuronal populations, oxytocin neurons are not a 'hotspot' of imprinted

expression. Specific sequencing of oxytocinergic brain regions will be required to distinguish between

these possibilities. A second caveat is that, due to the nature of the datasets used, not all imprinted genes

were included, and our analysis was missing a significant subset of imprinted genes encoding small

RNAs or isoforms from the same transcription unit. A third caveat is that we did not assess parent-of-

origin expression for the 119 imprinted genes we included in the analysis. Previous expression profiling

of imprinted genes have also not measured the POEs (16, 60) but have restricted their gene selection to

genes with reliable imprinting status. Consequently, we only included the canonical imprinted genes

and genes with more than one demonstration of a POE when looking for enrichment. Furthermore, for

the vast majority of these genes, a brain-based POE effect has also already been reported (Supplemental

Table S1). Although this does not replace validating the imprinting status of all 119 in the tissues and

subregions examined, it does provide justification for looking at imprinted gene over-representation.

To resolve this issue, scRNA-seq using tissues derived from reciprocal F1 crosses between distinct

mouse lines will be key; for example, the recent work of (61) with cortical cell types provides an

example of the allelic specific single-cell expression measurements necessary to confirm the

enrichments found in this study.

By exploiting scRNA-seq data we have asked whether imprinted genes as a group are disproportionately

represented in the brain, in specific brain regions, and in certain neuronal cell-types. In the adult brain

imprinted genes were over-represented in neurons, and particularly the hypothalamic neuroendocrine

populations and the monoaminergic hindbrain neurons, with the serotonergic neurons demonstrating

the clearest signal. Interestingly, PEGs, but not MEGs recreate this signal at Levels 1 and 2 - most

notably only PEGs display the hypothalamic neuronal enrichment. By extension, these data also identify

behaviours that are foci for the action of imprinted genes. Although there are high profile examples of

individual imprinted genes expressed in the key brain regions we highlight and that have roles in feeding

(*Magel2*) (62) and sleep (*Snord116*) (63), our analyses indicate that imprinted genes *as a group* are

strongly linked to these behaviours and also identify other individual genes that should be explored in

these domains. Conversely, there are high-profile examples of imprinted genes involved in

hippocampus related learning and memory (*Ube3a*) (28), but we did not find enrichment for cell types

related to this brain function. The idea that imprinted genes converge on specific physiological or

behavioural processes is not unprecedented. Specialisation of function is predicted when considering

why genomic imprinting evolved at all (13, 64-66). Moreover, there is increasing evidence that the

imprinted genes themselves appear to be co-expressed in an imprinted gene network (IGN) and have

confirmed regulatory links between each other (67-69). The idea of an IGN or, at the very least, heavily

correlated and coordinated expression between imprinted genes adds further support to the idea that

imprinted genes work in concert to influence processes, rather than in isolation, and that perturbating

one may influence many others (70). Our findings add substance to these general ideas and highlight

the neuronal regulation of pituitary function, feeding and sleep as being key functional hotspots on

which imprinted genes converge which probably provides the best current basis for discerning

evolutionary drivers of genomic imprinting in the brain.

METHODS

Data Processing

13 unique datasets were analysed across the three levels of analysis (see Fig.1) and analyses were

conducted on each dataset independently. At each level of analysis, we aimed to be unbiased by using

all the datasets that fitted the scope of that level, but the availability of public scRNA-seq datasets was

limited, which prevented us from exploring all avenues (for example, a direct comparison of

enrichment between hypothalamic nuclei). All sequencing data were acquired through publicly

available resources and each dataset was filtered and normalised according to the original published

procedure. Supplemental Table S19 details the basic parameters of each dataset. Once processed, each

dataset was run through the same basic workflow (see below and Fig. 10), with minor adjustments

laid out for each dataset detailed in the Supplemental Methods.

Due to the high variability in sequencing technology, mouse strain, sex and age, and processing

pipeline, we have avoided doing analysis on combined datasets. Rather we chose to perform our

analyses independently for each dataset and look for convergent patterns of imprinted gene

enrichment between datasets on similar tissues/brain regions. As with any single-cell experiment, the

identification of upregulation or over-representation of genes in a cell-type depends heavily on which

other cells are included in the analysis to make up the 'background'. Analysing separate datasets (with

overlapping cell-types alongside distinct ones) and looking for convergent patterns of enrichment is

one way of counteracting this limitation.

Basic Workflow

Data were downloaded in the available form provided by the original authors (either raw or

processed) and, where necessary, were processed (filtered, batch-corrected and normalized) to match

the author's original procedure. Cell quality filters were specific to each dataset and summarised in

Supplemental Table S19. A consistent filter, to remove all genes expressed in fewer than 20 cells, was

applied to remove genes unlikely to play a functional role due to being sparsely expressed. Datasets of

the whole brain/hypothalamus were analysed both at the global cell level (neuronal and non-neuronal

cells) and neuron specific level (only neurons) with genes filtered for the ≥ 20 cell expression at each

level before subsequent analysis. Cell identities were supplied using the outcome of cell clustering

carried out by the original authors, so that each cell included in the analysis had a cell-type or tissue-

type identity. This was acquired as metadata supplied with the dataset or as a separate file primarily

from the same depository as the data but occasionally acquired from personal correspondence with the

authors. Cells were used from mice of both sexes when provided and all mice were aged 15 weeks or

younger across all datasets. Although our focus was the adult mouse brain, embryonic data were

included in some comparisons or when no alternatives were present. However, embryonic and post-

natal cells were never pooled to contribute to the same cell populations.

Positive differential expression

between identity groups were

carried out using one-sided

Wilcoxon rank-sum tests (assuming

the average expression of cells

within the current identity group is

'greater' than the average of cells

from all other groups). The test was

performed independently for each

gene and for each identity group

vs. all other groups. The large

number of *p* values were

corrected for multiple

comparisons using a horizontal

Benjamini-Hochberg correction,

Figure 10. Basic workflow schematic. Single Cell Expression Matrices were acquired through publicly available depositories. Data were processed according to the author's original specifications and all genes were required to be expressed in 20 or more cells. Cell population identities were acquired from the author's original clustering. Positive differential gene expression was calculated via Wilcoxon Rank-Sum Test. Upregulated genes were considered as those with $q \leq$ 0.05 and a Log2FC $\geq$ 1 for analysis levels 1 and 2, while this criterion was relaxed to Log2FC > 0 for level 3. Our imprinted gene list was used to filter upregulated genes and two different enrichment analyses were carried out, over-representation analysis via Fisher's Exact Test and Gene Set Enrichment Analysis via Liger algorithm (Subramanian, Tamayo (71), https://github.com/JEFworks/liger). Venn diagrams and dot plots were utilised for visualisation.

creating *q* values. Fold-change (FC) values, percentage expression within the identity group and

percentage expressed within the rest were also calculated. We considered genes to be significantly

positively differentially expressed (significantly upregulated) in a group compared to background

expression if it had a $q \leq 0.05$. In addition, for Level 1 and Level 2 analyses, the criteria for

upregulated genes included demonstrating a Log2FC value of 1 or larger (i.e., 2-fold-change or

larger). The datasets at these levels represented cells from a variety of organs, regions and cell-types,

and in line with this cellular diversity, the aim of these analyses was to look for distinctive

upregulation, akin to a marker gene. Once the analysis was restricted to cell subpopulations within a

specific region of the brain (i.e., Level 3), the additional criteria for upregulation was relaxed to

demonstrating just a positive Log2FC (i.e., the gene has a higher expression in this cell type than

background). This was mainly because we were not expecting imprinted genes to be 'markers' of

individual subpopulations at this level, but our aim was to identify enriched expression profiles for

them. This additionally ensures consistent criteria for enrichment within levels, allowing meaningful

comparison.

The same custom list of imprinted genes with reliable parent-of-origin effects (see below) was used

for all analyses, and all genes were included as long as the gene passed the 20-cell filter. The first

statistical analysis for enrichment was an Over-Representation Analysis (ORA) using a one-sided

Fisher's Exact Test ('fisher.test' function in R core package 'stats v3.6.2'). The aim was to assess

whether the number of imprinted genes considered to be upregulated as a proportion of the total

number of imprinted genes in the dataset (passing the 20-cell filter) was statistically higher than

would be expected by chance when compared to the total number of upregulated genes as a

proportion of the overall number of genes in the dataset (passing the 20-cell filter). To limit finding

over-represented identity groups with only a few upregulated imprinted genes, an identity group was

required to have ≥ 5 % of the total number of imprinted genes upregulated for ORA to be conducted.

Subsequent p-values for all eligible identity groups were corrected using a Bonferroni correction. This

provided a measure of whether imprinted genes are expressed above expectation (as opposed to the

expression pattern of any random gene selection) in particular identity groups.

Venn diagrams of the upregulated imprinted genes making up over-represented identity groups across

datasets (within a level) were also reported. Full lists of upregulated imprinted genes can be found in

the 'Upregulated_IGs.csv' file for each analysis in the Supplemental Data.

To further examine the presence of imprinted genes within tissues/cell types, and to provide a

different perspective to over-representation, we conducted a Gene-Set Enrichment Analysis (GSEA)

for imprinted genes amongst the upregulated genes of an identity group using a publicly available,

light-weight implementation of the GSEA algorithm (71) in R (https://github.com/JEFworks/liger).

This was done in a manner similar to Moffitt, Bambah-Mukku (72) since we were similarly using this

computational method to identify enrichment of our gene sets *within the upregulated genes* of the

different identity groups. Here, the GSEA was conducted for each individual identity group using

Log2FC values to rank the upregulated genes. The GSEA acts as a more conservative measure than

the ORA since it tests whether imprinted genes are enriched in the stronger markers of a group (the

genes with the highest fold change for a group vs. the rest) and hence whether the imprinted genes are

enriched in those genes with a high specificity to that tissue/cell type. To prevent significant results

being generated from just 2 or 3 genes, identity group to be analysed were selected as having a

minimum of 15 upregulated imprinted genes (i.e. the custom gene set) to measure enrichment for (a

value suggested by the GSEA user guide (https://www.gsea-

msigdb.org/gsea/doc/GSEAUserGuide Frame.html)) and to prevent significant results in which

imprinted genes cluster at the tail, identity groups were selected as having an average fold change of

the upregulated imprinted genes greater than the average fold change of the rest of the upregulated

genes for that group. Again, multiple p values generated from GSEA were corrected using a

Bonferroni correction. To further elucidate the genes responsible for significant GSEA's, dot plots of

the imprinted genes upregulated in that identity group were plotted across all identity groups with

absolute expression and Log2FC mapped to size and colour of the dots, respectively. Graphical

representations of significant GSEA's (post-correction) are included in the main text or as

supplemental figures, all other graphs, including additional dot plots not discussed in this study, can

be found in the repository (https://osf.io/jx7kr/) and Supplemental Data. If no cell populations met

these criteria, GSEA was not run and not included for that analysis.

For Level 1 and Level 2 analyses, we also carried out parent-of-origin specific analyses. The

imprinted gene list was divided into MEGs and PEGs and the analyses detailed above were run

separately for these two gene groups. For imprinted genes with known parent-of-origin variability

based on tissue type (*Igf2* and *Grb10*), the parent-of-origin characterisation of these genes was

changed accordingly. The absolute number of imprinted genes top-expressed in a tissue/cell-type were

also reported for analyses in Level 1 and Level 2 in the tables, since these analyses included a variety

of cell-types and tissues which may demonstrate meaningful clustering of the highest normalised

expression values. The mean normalised expression for all imprinted genes across the series of

identity groups in the datasets in Level 1 and Level 2 was also calculated alongside the mean

normalised expression for the rest of the genes (Supplemental Table S2).

All graphical representations and statistical analyses were conducted using R 3.6.2 (73) in RStudio

(74). Diagrams in Figures 1:2, 4:6 and 8:10 were created with BioRender.com.

Custom Imprinted Gene List

The gene list for the analysis was based on the list of murine imprinted genes recently published in

Tucci, Isles (10). Although the original list of imprinted genes was 260 genes long, only 163 genes were

identified in the most comprehensive of the datasets. We further refined this list to 119 imprinted genes

(Supplemental Table S1) which excluded the X-linked genes, consisting of mostly the canonical

protein-coding and long noncoding RNA imprinted genes, but the criteria for inclusion was those genes

with at least two independent demonstrations of their POE status. The only exceptions to multiple

independent demonstrations of a POE were four genes (*Bmf*, *B3gnt2*, *Ptk2*, *Gm16299*) identified by

(34) where a POE was validated for 16 brain regions and 7 adult tissues. For Level 2, the MEG/PEG

status of a gene was primarily based on reported allelic expression within the brain. Small non-coding

RNAs such as micro-RNAs (miRs) and small nucleolar RNAs (snoRNAs), which represent ~10% of

identified imprinted genes, were excluded from the analysis as their sequences were not

detected/subsumed by larger transcripts in the majority of the datasets. Another caveat with short-read

RNA-seq libraries is that much of the expression data for a given transcription unit cannot discriminate

differentially imprinted isoforms nor do some of the technologies (e.g., Smart-Seq2) possess stranded

libraries to distinguish antisense transcripts. For complex imprinting loci such as the *Gnas* locus, most

reads as result map to only *Gnas* and *Nespas* ignoring several overlapping and antisense genes.

DECLARATIONS

Ethics approval and consent to participate

Not applicable. All samples had been collected in the context of previous studies.

Consent for publication

Not applicable.

Competing interests

All authors declare no financial and non-financial competing interests.

Availability of data and materials

All sequencing data used in this project were acquired through publicly available resources

(Databases, Accession Numbers and Depositories for each dataset can be found in the Supplemental

Methods). All data generated in this experiment is provided as Supplemental Data and in the

following Open Science Framework repository (https://osf.io/jx7kr/). Custom R scripts to analyse

each dataset are provided as Supplemental Code and are available at https://github.com/MJHiggs/IG-

Single-Cell-Enrichment.

Funding

This work was supported by a Wellcome Trust PhD studentship (220090/Z/20/Z).

Authors' Contributions

MJHiggs performed bioinformatic analysis, with input from MJHill; MJHiggs., and ARI contributed

to project design, data interpretation, and wrote the manuscript, MJHiggs produced all Figures; all co-

authors reviewed and edited the manuscript.

Acknowledgements

We would like to thank all the research groups that carried out the single-cell RNA sequencing that

made this study possible and to particularly acknowledge Dr. L. Cheung, Dr. P. Hook, Dr. A. Jackson,

and Dr. S. Wen for help accessing the cell metadata for their associated studies.

604   REFERENCES
605
1.   Zeisel A, Hochgerner H, Lönnerberg P, Johnsson A, Memic F, Van Der Zwan J, et al. Molecular
architecture of the mouse nervous system. Cell. 2018;174(4):999-1014. e22.
2.   Campbell JN, Macosko EZ, Fenselau H, Pers TH, Lyubetskaya A, Tenen D, et al. A molecular census
of arcuate hypothalamus and median eminence cell types. Nature neuroscience. 2017;20(3):484-96.
3.   Romanov RA, Zeisel A, Bakker J, Girach F, Hellysaz A, Tomer R, et al. Molecular interrogation of
hypothalamic organization reveals distinct dopamine neuronal subtypes. Nature neuroscience. 2017;20(2):176-
88.
4.   Chen R, Wu X, Jiang L, Zhang Y. Single-cell RNA-seq reveals hypothalamic cell diversity. Cell
reports. 2017;18(13):3227-41.
5.   Wen S, Ma D, Zhao M, Xie L, Wu Q, Gou L, et al. Spatiotemporal single-cell analysis of gene
expression in the mouse suprachiasmatic nucleus. Nature neuroscience. 2020;23(3):456.
6.   Ximerakis M, Lipnick SL, Innes BT, Simmons SK, Adiconis X, Dionne D, et al. Single-cell
transcriptomic profiling of the aging mouse brain. Nature neuroscience. 2019;22(10):1696-708.
7.   Cheung LY, George AS, McGee SR, Daly AZ, Brinkmeier ML, Ellsworth BS, et al. Single-cell RNA
sequencing reveals novel markers of male pituitary stem cells and hormone-producing cell types.
Endocrinology. 2018;159(12):3910-24.

8.   Ho Y, Hu P, Peel MT, Chen S, Camara PG, Epstein DJ, et al. Single-cell transcriptomic analysis of
adult mouse pituitary reveals sexual dimorphism and physiologic demand-induced cellular plasticity. Protein &
cell. 2020;11(8):565-83.
9.   Ferguson-Smith AC. Genomic imprinting: the emergence of an epigenetic paradigm. Nature Reviews
Genetics. 2011;12(8):565-75.
10.   Tucci V, Isles AR, Kelsey G, Ferguson-Smith AC, Bartolomei MS, Benvenisty N, et al. Genomic
imprinting and physiological processes in mammals. Cell. 2019;176(5):952-65.
11.   Orr HA. Somatic mutation favors the evolution of diploidy. Genetics. 1995;139(3):1441-7.
12.   Peters J. The role of genomic imprinting in biology and disease: an expanding view. Nature Reviews
Genetics. 2014;15(8):517-30.
13.   Moore T, Haig D. Genomic imprinting in mammalian development: a parental tug-of-war. Trends in
genetics. 1991;7(2):45-9.
14.   Andergassen D, Dotter CP, Wenzel D, Sigl V, Bammer PC, Muckenhuber M, et al. Mapping the mouse
Allelome reveals tissue-specific regulation of allelic expression. Elife. 2017;6:e25125.
15.   Babak T, DeVeale B, Tsang EK, Zhou Y, Li X, Smith KS, et al. Genetic conflict reflected in tissue-
specific maps of genomic imprinting in human and mouse. Nature genetics. 2015;47(5):544-9.
16.   Gregg C, Zhang J, Weissbourd B, Luo S, Schroth GP, Haig D, et al. High-resolution analysis of parent-
of-origin allelic expression in the mouse brain. science. 2010;329(5992):643-8.
17.   Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, et al. Genome-wide atlas of gene
expression in the adult mouse brain. Nature. 2007;445(7124):168-76.
18.   Negi SK, Guda C. Global gene expression profiling of healthy human brain and its application in
studying neurological disorders. Scientific reports. 2017;7(1):1-12.
19.   Cattanach BM, Kirk M. Differential activity of maternally and paternally derived chromosome regions
in mice. Nature. 1985;315(6019):496-8.
20.   Keverne EB, Fundele R, Narasimha M, Barton SC, Surani MA. Genomic imprinting and the
differential roles of parental genomes in brain development. Developmental Brain Research. 1996;92(1):91-100.
21.   Angulo M, Butler M, Cataletto M. Prader-Willi syndrome: a review of clinical, genetic, and endocrine
findings. Journal of endocrinological investigation. 2015;38(12):1249-63.
22.   Nicholls RD, Knoll JH, Butler MG, Karam S, Lalande M. Genetic imprinting suggested by maternal
heterodisomy in non-deletion Prader-Willi syndrome. Nature. 1989;342(6247):281-5.
23.   Perez JD, Rubinstein ND, Dulac C. New Perspectives on Genomic Imprinting, an Essential and
Multifaceted Mode of Epigenetic Control in the Developing and Adult Brain. Annual Review of Neuroscience.
2016;39(1):347-84.
24.   Li L-L, Keverne E, Aparicio S, Ishino F, Barton S, Surani M. Regulation of maternal behavior and
offspring growth by paternally expressed Peg3. Science. 1999;284(5412):330-4.
25.   Davies JR, Humby T, Dwyer DM, Garfield AS, Furby H, Wilkinson LS, et al. Calorie seeking, but not
hedonic response, contributes to hyperphagia in a mouse model for Prader–Willi syndrome. European Journal of
Neuroscience. 2015;42(4):2105-13.
26.   McNamara GI, John RM, Isles AR. Territorial behavior and social stability in the mouse require
correct expression of imprinted Cdkn1c. Frontiers in behavioral neuroscience. 2018;12:28.
27.   Garfield AS, Cowley M, Smith FM, Moorwood K, Stewart-Cox JE, Gilroy K, et al. Distinct
physiological and behavioural functions for parental alleles of imprinted Grb10. Nature. 2011;469(7331):534-8.
28.   Jiang Y-h, Armstrong D, Albrecht U, Atkins CM, Noebels JL, Eichele G, et al. Mutation of the
Angelman ubiquitin ligase in mice causes increased cytoplasmic p53 and deficits of contextual learning and
long-term potentiation. Neuron. 1998;21(4):799-811.
29.   Dent CL, Humby T, Lewis K, Ward A, Fischer-Colbrie R, Wilkinson LS, et al. Impulsive choice in
mice lacking paternal expression of Grb10 suggests intragenomic conflict in behavior. Genetics.
2018;209(1):233-9.
30.   Relkovic D, Doe CM, Humby T, Johnstone KA, Resnick JL, Holland AJ, et al. Behavioural and
cognitive abnormalities in an imprinting centre deletion mouse model for Prader–Willi syndrome. European
journal of neuroscience. 2010;31(1):156-64.
31.   Lassi G, Ball ST, Maggi S, Colonna G, Nieus T, Cero C, et al. Loss of Gnas imprinting differentially
affects REM/NREM sleep and cognition in mice. PLoS Genet. 2012;8(5):e1002706.
32.   Allen ND, Logan K, Lally G, Drage DJ, Norris ML, Keverne EB. Distribution of parthenogenetic cells
in the mouse brain and their influence on brain development and behavior. Proceedings of the National
Academy of Sciences. 1995;92(23):10782-6.
33.   DeVeale B, Van Der Kooy D, Babak T. Critical evaluation of imprinted gene expression by RNA–Seq:
a new perspective. PLoS Genet. 2012;8(3):e1002600.
34.   Perez JD, Rubinstein ND, Fernandez DE, Santoro SW, Needleman LA, Ho-Shing O, et al. Quantitative
and functional interrogation of parent-of-origin allelic expression biases in the brain. Elife. 2015;4:e07860.

35.   Han X, Wang R, Zhou Y, Fei L, Sun H, Lai S, et al. Mapping the mouse cell atlas by microwell-seq. Cell. 2018;172(5):1091-107. e17.

36.   Schaum N, Karkanias J, Neff NF, May AP, Quake SR, Wyss-Coray T, et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris: The Tabula Muris Consortium. Nature. 2018;562(7727):367.

37.   Zhang X, Jiang S, Mitok KA, Li L, Attie AD, Martin TFJ. BAIAP3, a C2 domain–containing Munc13 protein, controls the fate of dense-core vesicles in neuroendocrine cells. Journal of Cell Biology. 2017;216(7):2151-66.

38.   Rau AR, Hentges ST. The relevance of AgRP neuron-derived GABA inputs to POMC neurons differs for spontaneous and evoked release. Journal of Neuroscience. 2017;37(31):7362-72.

39.   Hook PW, McClymont SA, Cannon GH, Law WD, Morton AJ, Goff LA, et al. Single-cell RNA-seq of mouse dopaminergic neurons informs candidate gene selection for sporadic Parkinson disease. The American Journal of Human Genetics. 2018;102(3):427-46.

40.   La Manno G, Gyllborg D, Codeluppi S, Nishimura K, Salto C, Zeisel A, et al. Molecular diversity of midbrain development in mouse, human, and stem cells. Cell. 2016;167(2):566-80. e19.

41.   Huang KW, Ochandarena NE, Philson AC, Hyun M, Birnbaum JE, Cicconet M, et al. Molecular and anatomical organization of the dorsal raphe nucleus. Elife. 2019;8:e46464.

42.   Davies W, Isles AR, Wilkinson LS. Imprinted gene expression in the brain. Neuroscience & Biobehavioral Reviews. 2005;29(3):421-30.

43.   Cassidy FC, Charalambous M. Genomic imprinting, growth and maternal–fetal interactions. Journal of Experimental Biology. 2018;221(Suppl 1).

44.   Vagena E, Crneta J, Engström P, He L, Yulyaningsih E, Korpel NL, et al. ASB4 modulates central melanocortinergic neurons and calcitonin signaling to control satiety and glucose homeostasis. Science Signaling. 2022;15(733):eabj8204.

45.   Aponte Y, Atasoy D, Sternson SM. AGRP neurons are sufficient to orchestrate feeding behavior rapidly and without training. Nature neuroscience. 2011;14(3):351.

46.   Tucci V. Genomic imprinting: a new epigenetic perspective of sleep regulation. PLoS genetics. 2016;12(5):e1006004.

47.   Grattan DR, Steyn FJ, Kokay IC, Anderson GM, Bunn SJ. Pregnancy-induced adaptation in the neuroendocrine control of prolactin secretion. Journal of neuroendocrinology. 2008;20(4):497-507.

48.   Grattan D, Kokay I. Prolactin: a pleiotropic neuroendocrine hormone. Journal of neuroendocrinology. 2008;20(6):752-63.

49.   Davies W, Lynn PM, Relkovic D, Wilkinson LS. Imprinted genes and neuroendocrine function. Frontiers in neuroendocrinology. 2008;29(3):413-27.

50.   Ivanova E, Kelsey G. Imprinted genes and hypothalamic function. Journal of molecular endocrinology. 2011;47(2):R67-R74.

51.   Miller JL, Goldstone AP, Couch JA, Shuster J, He G, Driscoll DJ, et al. Pituitary abnormalities in Prader–Willi syndrome and early onset morbid obesity. American Journal of Medical Genetics Part A. 2008;146(5):570-7.

52.   Scagliotti V, Costa Fernandes Esse R, Willis TL, Howard M, Carrus I, Lodge E, et al. Dynamic Expression of Imprinted Genes in the Developing and Postnatal Pituitary Gland. Genes. 2021;12(4):509.

53.   Charalambous M, Da Rocha ST, Radford EJ, Medina-Gomez G, Curran S, Pinnock SB, et al. DLK1/PREF1 regulates nutrient metabolism and protects from steatosis. Proceedings of the National Academy of Sciences. 2014;111(45):16088-93.

54.   Huerta-ocampo I, Slack R, Beechey C, Skinner J, Peters J, Christian H, editors. Overexpression of the imprinted gene Neuronatin represses normal pituitary differentiation. Endocrine Abstracts; 2004: Bioscientifica.

55.   Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG. Galanin neurons in the medial preoptic area govern parental behaviour. Nature. 2014;509(7500):325-30.

56.   Mechenthaler I. Galanin and the neuroendocrine axes. Cellular and molecular life sciences: CMLS. 2008;65(12):1826-35.

57.   Pulix M, Plagge A. Imprinted Genes and Hypothalamic Function. Developmental Neuroendocrinology: Springer; 2020. p. 265-94.

58.   Donovan MH, Tecott LH. Serotonin and the regulation of mammalian energy balance. Frontiers in neuroscience. 2013;7:36.

59.   Dombret C, Nguyen T, Schakman O, Michaud JL, Hardin-Pouzet H, Bertrand MJ, et al. Loss of Maged1 results in obesity, deficits of social interactions, impaired sexual behavior and severe alteration of mature oxytocin production in the hypothalamus. Human molecular genetics. 2012;21(21):4703-17.

60.   Steinhoff C, Paulsen M, Kielbasa S, Walter J, Vingron M. Expression profile and transcription factor binding site exploration of imprinted genes in human and mouse. BMC Genomics. 2009;10:144.

61.   Laukoter S, Pauler FM, Beattie R, Amberg N, Hansen AH, Streicher C, et al. Cell-type specificity of genomic imprinting in cerebral cortex. Neuron. 2020;107(6):1160-79. e9.

62.   Schaller F, Watrin F, Sturny R, Massacrier A, Szepetowski P, Muscatelli F. A single postnatal injection
of oxytocin rescues the lethal feeding behaviour in mouse newborns deficient for the imprinted Magel2 gene.
Human Molecular Genetics. 2010;19(24):4895-905.
63.   Lassi G, Priano L, Maggi S, Garcia-Garcia C, Balzani E, El-Assawy N, et al. Deletion of the
Snord116/SNORD116 alters sleep in mice and patients with Prader-Willi syndrome. Sleep. 2016;39(3):637-44.
64.   Keverne E. Significance of epigenetics for understanding brain development, brain evolution and
behaviour. Neuroscience. 2014;264:207-17.
65.   Keverne EB, Martel FL, Nevison CM. Primate brain evolution: genetic and functional considerations.
Proceedings of the Royal Society of London Series B: Biological Sciences. 1996;263(1371):689-96.
66.   Trivers R, Burt A. Kinship and genomic imprinting. Genomic imprinting: Springer; 1999. p. 1-21.
67.   Al Adhami H, Evano B, Le Digarcher A, Gueydan C, Dubois E, Parrinello H, et al. A systems-level
approach to parental genomic imprinting: the imprinted gene network includes extracellular matrix genes and
regulates cell cycle exit and differentiation. Genome research. 2015;25(3):353-67.
68.   Varrault A, Gueydan C, Delalbre A, Bellmann A, Houssami S, Aknin C, et al. Zac1 regulates an
imprinted gene network critically involved in the control of embryonic growth. Developmental cell.
2006;11(5):711-22.
69.   Gabory A, Ripoche M-A, Le Digarcher A, Watrin F, Ziyyat A, Forné T, et al. H19 acts as a trans
regulator of the imprinted gene network controlling growth in mice. Development. 2009;136(20):3413-21.
70.   Patten MM, Cowley M, Oakey RJ, Feil R. Regulatory links between imprinted genes: evolutionary
predictions and consequences. Proceedings of the Royal Society B: Biological Sciences.
2016;283(1824):20152760.
71.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.
Proceedings of the National Academy of Sciences. 2005;102(43):15545-50.
72.   Moffitt JR, Bambah-Mukku D, Eichhorn SW, Vaughn E, Shekhar K, Perez JD, et al. Molecular,
spatial, and functional single-cell profiling of the hypothalamic preoptic region. Science. 2018;362(6416).
73.   Team R. R: A language and environment for statistical computing. Vienna, Austria; 2013.
74.   Team R. RStudio: integrated development for R. RStudio, Inc, Boston, MA URL http://www rstudio
com. 2015;42:14.
771