Systematic investigation of imprinted gene expression and enrichment in the mouse brain explored at single-cell resolution

M. J. Higgs[1], M. J. Hill[2,3], R. M. John[4], and A. R. Isles[1,*]

1 Behavioural Genetics Group, MRC Centre for Neuropsychiatric, Genetics and Genomics, Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff, UK
2 Division of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Cardiff, UK
3 UK Dementia Research Institute, School of Medicine, Cardiff University, Cardiff, UK
4 School of Biosciences, Cardiff University, Cardiff, UK
* Corresponding author: A. R. Isles, Behavioural Genetics Group, MRC Centre for Neuropsychiatric Genetics and Genomics, Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff CF24 4HQ, UK.
E-mail: islesar1@cardiff.ac.uk

ABSTRACT

Background

Although a number of imprinted genes are known to be highly expressed in the brain, and in certain brain regions in particular, whether they are truly over-represented in the brain has never been formally tested. Using thirteen single-cell RNA sequencing datasets we systematically investigated imprinted gene over-representation at the organ, brain region, and cell-specific levels.

Results

We established that imprinted genes are indeed over-represented in the adult brain, and in neurons particularly compared to other brain cell-types. We then examined brain-wide datasets to test enrichment within distinct brain regions and neuron subpopulations and demonstrated over-representation of imprinted genes in the hypothalamus, ventral midbrain, pons and medulla. Finally, using datasets focusing on these regions of enrichment, we identified hypothalamic neuroendocrine populations and the monoaminergic hindbrain neurons as specific hotspots of imprinted gene expression.

Conclusions

These analyses provide the first robust assessment of the neural systems on which imprinted genes converge. Moreover, the unbiased approach, with each analysis informed by the findings of the previous level, permits highly informed inferences about the functions on which imprinted gene

expression converges. Our findings indicate the neuronal regulation of motivated behaviours such as

feeding and sleep, alongside the regulation of pituitary function as functional hotspots for imprinting,

thus adding statistical rigour to prior assumptions and providing testable predictions for novel neural

and behavioural phenotypes associated with specific genes and imprinted gene networks. In turn, this

work sheds further light on the potential evolutionary drivers of genomic imprinting in the brain.

BACKGROUND

Imprinted genes demonstrate a preferential or exclusively monoallelic expression from either the

maternal or paternal allele in an epigenetically predetermined manner (a parent-of-origin effect, POE).

To date approximately 260 imprinted genes, demonstrating biased allelic expression and/or associated

with a parental-specific epigenetic mark, have been identified in the mouse (~230 in humans) (9, 10).

This epigenetic regulation makes genomic imprinting an evolutionary puzzle as many of these genes

are effectively haploid and thereby negate many of the benefits of diploidy (11). Studying the patterns

of expression and function of imprinted genes may shed light on the drivers leading to the evolution

of genomic imprinting. For instance, characterisation of a number of imprinted genes points to

convergence on placental function (12), in line with the predictions of early theoretical ideas (13).

Outside of the placenta, the brain consistently emerges as an adult tissue with a large number of

expressed imprinted genes (14-16). However, given that it is estimated that ~80% of all genes in the

genome are expressed in the brain (17, 18), the question remains, *is* imprinted gene expression

actually enriched in the brain compared to other adult tissues? To date this has never been formally

tested.

A role for imprinted genes in the brain was initially suggested by (19), and neurological phenotypes

observed in, early imprinted gene mouse models (20). In addition, behavioural deficits were seen in

imprinting disorders such as Prader-Willi and Angelman syndromes (21, 22). Subsequent studies have

revealed diverse roles for imprinted genes in the brain. During development, several imprinted genes

are involved in the processes of neural differentiation, migration, axonal outgrowth and apoptosis

(23). In the adult brain, studies of mice carrying manipulations of individual imprinted genes have

suggested a wide range of behavioural roles including maternal care (24), feeding (25), social

behaviour (26, 27), learning/memory (28), cognition (29, 30), and more recently, sleep and circadian

activity (31).

In addition to studies on individual imprinted genes, there are a limited number of studies that take a

systems level approach to characterizing the role of genomic imprinting in the brain. Early studies

examining developing and adult chimeras of normal and parthenogenetic/gynogenetic (Pg/Gg - two

maternal genomes) or androgenetic (Ag - two paternal genomes) cells indicated distinct regional

distribution for maternally (cortex and hippocampus) and paternally (hypothalamus) expressed genes

(20, 32). More recently, Gregg, Zhang (16) used the known imprinting status of 45 imprinted genes

and the Allen Brain Atlas to track dichotomous expression of imprinted genes across 118 brain

regions to identify brain-wide patterns of expression. Most imprinted genes were expressed in every

brain region, but detectable expression of the largest number of imprinted genes was found in regions

of the hypothalamus (medial preoptic area, arcuate nucleus), central amygdala, basal nuclei of the

stria terminalis and the monoaminergic nuclei, suggesting some form of specialisation. Although

pioneering, this study, and others identifying novel imprinted genes and/or mapping allelic expression

in the brain (14, 15, 33, 34), did not test whether the expression of these genes was especially

enriched in given brain regions but simply asked if they were expressed, at any level, or not.

Here we address the question of whether the brain and/or specific brain circuitry is a foci for genomic

imprinting by exploiting the rapidly expanding number of single-cell RNA sequencing (scRNA-seq)

datasets and systematically investigating imprinted gene enrichment and over-representation in the

murine brain. We performed this by a hierarchical sequence of data analysis, using datasets that

allowed a multi-organ (Level 1) comparison first, before proceeding to brain-specific (Level 2) and

brain region-specific (Level 3) comparisons with the outcome of each level informing the data

selection for the next one, to identify a consistent pattern of enrichment (Figure 1). We sought to

provide a robust assessment of the neural systems on which imprinted genes converge, statistically

validating previous assumptions, identifying neuronal domains that have received less emphasis in

earlier studies, and providing testable predictions for novel neural and behavioural phenotypes

associated with specific genes and imprinted gene networks.

RESULTS

89   **Imprinted gene expression is enriched in the brain in a multi-organ analysis (Level 1 Analysis)**

90   The Mouse Cell Atlas (MCA) (35) and the *Tabula Muris* (TM) (36) are single cell compendiums

91   containing ~20 overlapping, but not identical, adult mouse organs. Key overlapping organs include

92   the bladder, brain, kidney, lung, limb muscle, and pancreas while organs included in only one dataset

93   include the ovary, testes, uterus, stomach within the MCA, and the heart, fat, skin, trachea and

94   diaphragm within the TM. These compendiums create a snapshot of gene expression across adult

95   tissues to assess imprinted gene enrichment. Since this study focused on the adult body and brain,

Figure 1. The hierarchical set of datasets in this analysis. The datasets are sorted into Level 1 (Multi-Organ) , Level 2 (Whole Brain) and Level 3 (Specific Brain Nuclei) analyses. The original publication and specific tissue/s analysed are provided for each analysis. White text in dark grey box indicates specifics to the analysis at that level – whether the analysis used the 'marker gene' Log2FC criteria or the relaxed Log2FC > 0 criterion, whether paternally and maternally expressed gene (PEG/MEG) analysis was carried out and whether the number of IGs with highest expression in a cell population and the average normalised expression were reported for imprinted genes.

96   fetal tissues (including the placenta) were not assessed.

97   An over-representation analysis (ORA) was performed on both datasets. All data were processed

98   according to the original published procedure, a list of upregulated genes was produced for each

99   tissue/identity group (vs. all other tissue/identity groups) and a one-sided Fisher's Exact test was

Table 1. Imprinted gene over-representation in MCA adult tissues (35). *Identity* – Tissue identities for the cells used in analysis; *Up Reg* – number of upregulated genes with $q \leq 0.05$ and Log2FC $\geq$ 1 (total number of genes in the dataset in brackets); *IG* – number of imprinted genes upregulated with $q \leq 0.05$ and Log2FC $\geq$ 1 (total number of IGs in the dataset in brackets); *ORA p* – *p* value from over representation analysis on groups with minimum 5% of total IGs; *ORA q* – Bonferroni corrected *p* value from ORA; *Mean FC IG* – mean fold change for upregulated imprinted genes; *Mean FC Rest* – mean fold change for all other upregulated genes; *No. IGs with highest expression* – Number of IGs with highest mean expression for cells from that identity group.

Tissue Identity	Up Reg (20,534)	IG (95)	ORA p	ORA q	Mean FC IG	Mean FC Rest	No. IGs with highest expression
Pancreas	2737	42	1.57E-13	**1.89E-12**	8.74	10.32	22
Brain	3401	34	4.43E-06	**5.31E-05**	8.76	125.00	19
Bladder	3183	29	0.000168	**0.002012**	4.45	8.51	8
Uterus	2567	22	0.002827	**0.033919**	4.66	8.46	7
Lung	1203	8	0.192705	1	3.82	151.41	4
Ovary	2219	13	0.223666	1	7.46	11.27	5
Kidney	1714	10	0.268425	1	13.76	182.89	5
Liver	1739	8	0.560145	1	4.55	80.51	3
Stomach	1821	7	0.748590	1	4.24	88.60	3
Thymus	1805	6	0.851579	1	2.78	6.76	2
Small Intestine	1719	5	0.908008	1	7.99	218.64	2
Testis	5212	14	0.995891	1	27.04	5058.36	10
Bone Marrow	1095	2	-	-	5.31	4.43	1
Mammary Gland Virgin	902	4	-	-	3.70	4.03	0
Muscle	1127	4	-	-	8.64	15.05	3
Peripheral Blood	1146	3	-	-	3.78	3.57	0
Prostate	369	0	-	-	0.00	478.10	0
Spleen	1501	1	-	-	4.90	4.77	1

performed using a custom list of imprinted genes (Supplemental Table S1) to identify tissues in which

imprinted genes were over-represented amongst the upregulated genes for that tissue. Each dataset in

this study was analysed independently which allowed us to look for convergent patterns of enrichment

between datasets of similar tissues/cell-types. Across only adult tissues, imprinted genes were

convergently over-represented in the pancreas, bladder and the brain in both datasets (Figure 2A). In

addition, in the MCA adult tissue dataset, there was a significant over-representation in the uterus

(Table 1), and in the *Tabula Muris* analysis (Table 2), there was a significant over-representation in

Table 2. Imprinted gene over-representation in *Tabula Muris* adult tissues (36). *GSEA p – p* value from Gene Set Enrichment Analysis for identity groups with 15+ IGs and Mean FC IG > Mean FC Rest; *GSEA q* – Bonferroni corrected *p* values from GSEA. All other column descriptions can be found in the legend of Table 1.

Tissue Identity	Up Reg (20,839)	IG (107)	ORA p	ORA q	Mean FC IG	Mean FC Rest	GSEA p	GSEA q	No. IGs with highest expression
Diaphragm	416	19	3.66E-13	**4.75E-12**	6.49	4.83	0.1898	0.3796	4
Limb Muscle	761	24	6.32E-13	**8.22E-12**	9.02	5.09	0.0552	0.1104	8
Pancreas	4104	43	8.31E-07	**1.08E-05**	12.52	12.60	-	-	29
Trachea	1979	25	1.78E-05	**0.0002**	3.81	4.57	-	-	5
Brain (Non-Myeloid)	3081	31	0.0001	**0.0016**	12.16	14.17	-	-	14
Bladder	3338	31	0.0005	**0.0068**	3.30	5.30	-	-	16
Fat	1263	12	0.0286	0.3713	3.46	3.68	-	-	1
Heart	1108	10	0.0585	0.7601	2.87	5.14	-	-	0
Mammary Gland	1826	12	0.2264	1	3.52	5.24	-	-	3
Liver	1808	7	0.8307	1	6.19	54.93	-	-	3
Aorta	3515	14	0.8832	1	7.47	16.08	-	-	2
Tongue	4295	15	0.9696	1	4.15	7.16	-	-	8
Large Intestine	4758	11	0.9998	1	5.95	12.22	-	-	5
Brain (Myeloid)	1024	5	-	-	3.39	6.80	-	-	2
Kidney	584	3	-	-	24.94	22.90	-	-	1
Lung	914	2	-	-	2.73	5.41	-	-	0
Marrow	1957	5	-	-	7.65	5.25	-	-	4
Skin	1612	4	-	-	4.15	8.36	-	-	1
Spleen	625	1	-	-	4.63	4.28	-	-	0
Thymus	678	4	-	-	3.46	7.45	-	-	1

the muscle-based tissues - diaphragm, trachea, and limb muscles. In addition to the ORA, to identify

situations in which imprinted genes were in fact enriched amongst the stronger markers of a

tissue/cell-type, we performed a Gene-Set Enrichment Analysis (GSEA) on tissues meeting minimum

criteria (see Methods), which assessed whether imprinted genes were enriched within the top ranked

upregulated genes for that tissue (ranked by Log2 Fold Change). No tissue at this level showed a

significant GSEA for imprinted genes. Mean normalised expression of imprinted genes across identity

groups (Supplemental Table S2) was the highest for Brain in the MCA and highest for Pancreas in the

TM (Brain (Non-Myeloid) was the fourth highest).

Given the interest in the different functions of maternally expressed genes (MEGs) and paternally

expressed genes (PEGs), we additionally ran the large-scale enrichment analyses (Levels 1 and 2)

using separate lists of PEGs and MEGs. At Level 1, MEGs and PEGs (Supplemental Table S3A, S3B,

S4A and S4B) revealed a similar pattern of enrichment in both datasets (Fig. 2). PEGs were over-

Figure 2. Level 1 multi-organ comparison summary graphics. (*A*) Venn diagram of upregulated imprinted genes in the brain in Mouse Cell Atlas and in the brain (non-myeloid) in the Tabula Muris. Imprinted genes are listed which show significant upregulation (*q* ≤ 0.05 and Log2FC ≥ 1) in the tissues. Although these tissues are not identical, these were the two brain associated over-representations in the enrichment analysis. Parental-bias is indicated by colour (MEG - red, PEG - blue). From the 119 imprinted genes in the gene list, only 92 were common to both analyses (i.e., successfully sequenced and passed gene quality control filters). 34 imprinted genes were upregulated in the brain in the MCA and 31 genes in the *TM*. Genes in common from the two analyses are presented in bold and totalled in each section of the Venn Diagram, while genes found upregulated in one analysis but not available in the other analysis are included in small font and the number indicated in brackets. (*B*) Tissues with over-representation in MCA. Coloured tissues with bold labels were over-represented tissues using all imprinted genes, tissues with a blue circle behind were over-represented for PEGs alone, a red circle represent the same for MEGs, and a red/blue split circle were over-represented for both PEGs and MEGs. (*C*)

represented in the brain in both datasets (MCA - q = 4.56x10^{-6}, *TM* - q = 0.0005) while MEGs were

not. PEGs were also over-represented in the diaphragm (q = 0.0007), limb muscle (q = 0.0001) and

pancreas (MCA - q =1.93x10^{-5}, *TM* - q = 0.0002), with a significant GSEA in the MCA pancreas (p =

0.02, Supplemental Fig. S1). While MEGs were over-represented in the bladder (MCA - q = 0.002,

TM - q = 0.020), the pancreas (MCA - q = 1.53x10^{-7}) and in the three muscular tissues of the *Tabula*

Muris (diaphragm - q = 2.13x10^{-8}, limb muscle - q = 2.43x10^{-7}, trachea - q = 0.004).

Imprinted gene expression is enriched in neurons and neuroendocrine cells of the brain (Level 2

Analysis)

We next analysed cells from the whole mouse brain (Level 2), firstly using the Ximerakis, Lipnick (6)

dataset, in which cells were grouped from the whole mouse brain (minus the hindbrain) into major

cell classes according to cell lineage. Imprinted genes were over-represented in neuroendocrine cells

and mature neurons (Table 3).

Neuroendocrine cells were defined as a heterogeneous cluster, containing peptidergic neurons and

neurosecretory cells expressing neuronal marker genes (e.g., *Syt1* and *Snap25*) alongside neuropeptide

genes (e.g., *Oxt*, *Avp*, *Gal*, *Agrp* and *Sst*) but distinguished by Ximerakis, Lipnick (6) by the unique

expression of *Baiap3* which plays an important role in the regulation of exocytosis in neuroendocrine

cells (37). GSEA additionally showed that the imprinted genes were enriched in the genes with the

highest fold change values for neuroendocrine cells only (Fig. 3). 26 imprinted genes had their highest

expression in the neuroendocrine cells and the mean normalised expression of imprinted genes was

almost twice as high for neuroendocrine cells as the next highest identity group (Supplemental Table

S2). The MEG/PEG analysis (Supplemental Table S5A and S5B) for this dataset found that PEGs

were over-represented in mature neurons (q = 0.027) and neuroendocrine cells (q = 8.97 x 10^{-6}).

MEGs were also over-represented in neuroendocrine cells (q = 0.047) and uniquely over-represented

in Arachnoid barrier cells (q = 0.014). Only PEGs replicated the significant GSEA in neuroendocrine

cells (p = 4x10^{-4}, Supplemental Fig. S2).

Table 3. Imprinted gene over-representation in neural lineage types (6). *Identity* – Cell lineage identities for the cells used in analysis. All other column descriptions can be found in the legend of Tables 1 & 2.

Cell Population Identity (Abbr.)	Up Reg (14,498)	IG (85)	ORA *p*	ORA *q*	Mean FC IG	Mean FC Rest	GSEA *p*	GSEA *q*	No. IGs with highest expression
Neuroendocrine cells (NendC)	3868	47	2.12E-08	**3.82E-07**	11.88	5.42	0.0017	**0.0051**	26
Mature Neurons (all types) (mNEUR)	2968	32	0.0002	**0.0035**	8.80	9.28	-	-	2
Arachnoid barrier cells (ABC)	2287	20	0.0396	0.7120	16.84	22.63	-	-	7
Tanycytes (TNC)	1279	12	0.0692	1	6.64	12.01	-	-	8
Vascular and leptomeningeal cells (VLMC)	1714	15	0.0724	1	15.06	13.03	0.0468	0.1404	4
Oligodendrocyte precursor cells (OPC)	1524	13	0.1067	1	3.03	7.17	-	-	1
Pericytes (PC)	1801	14	0.1649	1	8.20	8.22	-	-	2
Olfactory ensheathing glia (OEG)	1086	9	0.1848	1	7.95	26.03	-	-	1
Oligodendrocytes (OLG)	1183	9	0.2561	1	3.73	12.91	-	-	5
Choroid plexus epithelial cells (CPC)	2602	17	0.3524	1	7.43	19.34	-	-	5
Hemoglobin-expressing vascular cells (Hb_VC)	1798	11	0.4889	1	5.25	6.33	-	-	3
Vascular smooth muscle cells (VSMC)	3006	17	0.6093	1	8.94	6.71	0.1376	0.4128	5
Astrocyte-restricted precursors (ARP)	1445	8	0.6214	1	4.50	5.09	-	-	1
Neural stem cells (NSC)	1009	5	0.7138	1	4.00	4.09	-	-	0
Ependymocytes (EPC)	3233	17	0.7346	1	15.04	53.27	-	-	4
Endothelial cells (EC)	1455	7	0.7619	1	5.80	8.54	-	-	0
Hypendymal cells (HypEPC)	1525	6	0.8946	1	17.24	20.80	-	-	5
Neuronal-restricted precursor (NRP)	2339	10	0.8979	1	3.07	10.20	-	-	1
Astrocytes (ASC)	1384	4	-	-	2.22	6.04	-	-	0
Dendritic cells (DC)	1209	1	-	-	3.50	16.02	-	-	1
Immature Neurons (ImmN)	652	4	-	-	3.37	5.79	-	-	0
Macrophages (MAC)	1222	2	-	-	3.47	21.56	-	-	0
Microglia (MG)	1342	3	-	-	19.28	19.22	-	-	3
Monocytes (MNC)	947	2	-	-	16.49	19.13	-	-	1
Neutrophils (NEUT)	519	2	-	-	9.18	62.13	-	-	0

The second dataset at this level was Zeisel, Hochgerner (1) Mouse Brain Atlas (MBA) which allowed

a much deeper investigation of nervous system enrichment with sequencing of the entire murine

nervous system and identifying cells by both brain region and cell type. Concordant with the previous

findings, primary analysis separating cells by lineage revealed over-representation of imprinted genes

in neurons only (Table 4). The overlap between the upregulated imprinted genes for the over-

represented neural-lineage cells from the Level 2 datasets are displayed in Figure 4. Additionally,

Table 4. Imprinted gene over-representation in nervous system cell types (1). *Identity* – Cell identities for the cells used in analysis. All other column descriptions can be found in the legend of Table 1.

Cell Population Identity	Up Reg (19,547)	IG (109)	ORA *p*	ORA *q*	Mean FC IG	Mean FC Rest	No. IGs with highest expression
Neurons	5710	44	0.0081	**0.0487**	11.73	24.97	45
Vascular	2473	22	0.0171	0.1029	17.91	26.64	16
Oligos	1587	11	0.2701	1	4.64	11.48	12
Peripheral Glia	2820	16	0.5117	1	5.42	12.64	12
Ependymal	3683	20	0.5912	1	24.52	66.97	15
Immune	1564	7	0.7787	1	13.42	93.05	5
Astrocytes	1539	4	-	-	2.88	10.73	3

152   PEGs alone demonstrated

153   no significant over-representations in cell lineage types while MEGs demonstrated over-

154   representation in vascular cells only ($q = 0.0004$) (Supplemental Table S6A and S6B).

A

Enrichment Score (cumulative weight)

P-value < 0.0017

Log2 FC Values

edge value = 3

Sorted Genes (by descending Log2FC)

B

Imprinted Gene

Normalised Mean Expression

Log2FC vs Background (MEGs)

Log2FC vs Background (PEGs)

Cell Lineage Identity (Whole Brain – Ximerakis et al., 2019)

Figure 3. GSEA and dot plots for imprinted genes upregulated in neuroendocrine cells in the Ximerakis, Lipnick (6) whole mouse brain dataset. (*A*) GSEA for imprinted genes upregulated in the neuroendocrine cells. In the analysis, genes are sorted by strength by which they mark this neuronal cluster (sorted by Log2FC values) indicated by the bar (middle). Fold change values are displayed along the bottom of the graph. The genes are arrayed left (strongest marker) to right and blue lines mark where imprinted genes fall on this array. The vertical axis indicates an accumulating weight, progressing from left to right and increasing or decreasing depending on whether the next gene is an imprinted gene or not. The *p*-value represents the probability of observing the maximum value of the score (red dashed line) if the imprinted genes are distributed randomly along the horizontal axis. The *q*-value for this analysis was significant at 0.0036. (*B*) Dot plot of imprinted genes upregulated in the 'Neuroendocrine cells' plotted across all identified cell types (Abbr. in Table 3). Imprinted genes were plotted in chromosomal order. Size of points represented absolute mean expression; colour represented the size of the Log2FC value for the cell identity group (e.g., neuroendocrine cells) vs. all other cells. Unique colour scales are used for MEGs (red/orange) and PEGs (blue). Where a gene was not expressed in a cell type, this appears as a blank space in the plot

The hypothalamus, ventral midbrain, pons and medulla are enriched for imprinted gene

expression (Level 2 Analysis)

After confirming neuron-specific enrichment of imprinted genes in the MBA dataset, further MBA

analysis was performed on cells classified as neurons and then grouped by brain/nervous system

regions. Significant over-representation was seen in neurons of the hypothalamus, ventral midbrain,

medulla, and pons (Table 5). The pons and medulla had the largest number, 45 and 44 respectively, of

Figure 4. Venn diagram of upregulated imprinted genes in the mature neuronal cells in the whole brain datasets of Zeisel, Hochgerner (1) and Ximerakis, Lipnick (6). Imprinted genes are listed which show significant upregulation (*q* ≤ 0.05 and Log2FC ≥ 1) in the cells. Although these cell types are not identical, these were all mature neural lineage cells with over-representations in the enrichment analysis. Parental-bias is indicated by colour (MEG - red, PEG - blue. From the 119 imprinted genes in the gene list, only 88 were common to both analyses (i.e., successfully sequenced and passed gene quality control filters). 45 imprinted genes were upregulated in neurons in the MBA, and in Ximerakis, Lipnick (6), 33 imprinted genes were upregulated in neurons and 48 genes in neuroendocrine cells. Genes in common from the two analyses are presented in bold and totalled in each section of the Venn Diagram, while genes found upregulated in one analysis but not available in the other analysis are included in small font and the number indicated in brackets.

imprinted genes upregulated (Figure 5A).

Regional analysis for MEGs and PEGs separately (Supplemental Table S7A and S7B), revealed that

PEGs were over-represented in hypothalamus ($q = 6.53 \times 10^{-7}$), ventral midbrain ($q = 0.018$), the pons

($q = 4.65 \times 10^{-5}$) and the medulla ($q = 4.10 \times 10^{-6}$); while MEGs were only over-represented in the

medulla ($q = 0.002$) but had a significant GSEA for the pons ($q = 0.027$, Supplemental Fig. S3); see

Figure 5B.

Table 5. Imprinted gene over-representation in nervous system region(1). *Identity* – Nervous system regional identities

for the cells used in analysis genes. All other column descriptions can be found in the legend of Tables 1 & 2.

Brain Region Identity	Up Reg (18,335)	IG (106)	ORA p	ORA q	Mean FC IG	Mean FC Rest	GSEA p	GSEA q	No. IGs with highest expression
Medulla	3147	45	8.38E-10	**1.26E-08**	4.79	4.01	0.1	0.2	15
Hypothalamus	1040	22	9.81E-08	**1.47E-06**	4.92	5.84	-	-	8
Pons	3581	44	1.62E-07	**2.43E-06**	4.20	3.91	0.1169	0.2338	22
Vent. Midbrain	1228	18	0.0002	**0.0034**	4.90	4.99	-	-	3
Vent. Striatum	689	8	0.0463	0.6941	3.92	4.92	-	-	0
Posterior Cortex	1090	9	0.1788	1	2.64	3.20	-	-	2
Enteric Nervous System	3885	26	0.2311	1	8.98	121.04	-	-	11
Sympathetic Nervous System	2804	18	0.3535	1	11.37	57.96	-	-	9
Anterior Cortex	979	6	0.5016	1	2.72	3.30	-	-	1
Dors. Midbrain	1045	6	0.5663	1	2.20	4.85	-	-	3
Thalamus	1441	8	0.6000	1	2.90	6.36	-	-	0
Hippocampus - CA1	1082	6	0.6008	1	3.01	4.02	-	-	2
Somatosensory Cortex	2121	11	0.6943	1	4.09	3.70	-	-	8
Dors. Striatum	1196	6	0.6974	1	4.03	5.43	-	-	2
Dorsal Root Ganglion	3607	16	0.9088	1	11.56	75.89	-	-	9
Middle Cortex	623	5	-	-	3.29	3.24	-	-	0
Spinal Cord	972	5	-	-	4.57	12.36	-	-	1
Amygdala	452	4	-	-	4.65	4.11	-	-	2
Dentate Gyrus	796	4	-	-	3.79	4.16	-	-	2
Hippocampus	631	4	-	-	4.86	3.82	-	-	2
Olfactory Bulb	445	4	-	-	4.02	8.27	-	-	2
Antero-Middle Cortex	646	3	-	-	4.59	4.31	-	-	1
Cerebellum	240	0	-	-	0.00	32.30	-	-	0

167   Neurons were then recategorized into unique subpopulations identified by marker genes (1) to

Figure 5. Level 2 Brain Region Analysis summary figures. (*A*) Venn diagram of upregulated imprinted genes in the neurons of enriched nervous system regions from the Mouse Brain Atlas (1). Imprinted genes are listed which show significant upregulation (*q* ≤ 0.05 and Log2FC ≥ 1) in the regions specified. The number of imprinted genes in each region of the Venn diagram are specified. Parental-bias of imprinted genes is indicated by colour (MEG - red, PEG - blue). (*B*) Brain regions enriched for imprinted gene expression via ORA or GSEA in the MBA (1). Regions over-represented for all imprinted genes are bolded. Regions over-represented for PEG expression alone are coloured blue while regions enriched for MEG expression alone are coloured red.

uncover the specific neural populations underlying the enrichment seen in the hypothalamus, pons and

medulla, and midbrain (Fig. 6; Supplemental Table S8). Each neural population was identified by its

distinct gene expression and suspected location within the brain (see http://mousebrain.org/ for an

online resource with detailed information on each cluster).

The hypothalamus was represented by a selection of inhibitory and peptidergic neurons. Inhibitory

neurons with over-representation of imprinted genes included: a Subthalamic Nucleus population

(notable genes *Lhx8, Gabrq*), two Preoptic Area/ BNST populations (*Nts*, *Dlk1 / Gal, Irs4*)

representing, an Arcuate nucleus population (*Agrp, Otp*), and two Suprachiasmatic nucleus

populations (*Avp, Nms*, *Six6, Vip*). For peptidergic neurons, over-representation was seen in a

ventromedial population (*Gpr101, Tac1, Baiap3*), a ventromedial/paraventricular population (*Otp,*

Trh, Ucn3), a lateral hypothalamic population (*Trh, Otp, Ngb*), an oxytocin magnocellular population

of the paraventricular and supraoptic nuclei (*Oxt, Otp*), and an orexin producing population of the

dorsomedial/lateral hypothalamus (*Hcrt, Pdyn, Trhr*).

Figure 6. Anatomical labelling of all the neural subpopulations with a significant over-representation of imprinted genes ($q \leq 0.05$ and Log2FC ≥ 1) in the Mouse Brain Atlas (1). The predicted brain nuclei localisation of the 32 neuronal subpopulations (out of 214 populations identified across the nervous system) specified in the MBA and enriched for imprinted genes. Brain regions that were not found to be enriched for imprinted genes are greyed out. The full Enrichment

181   The midbrain, medulla and pons were represented by a number of cell groups, with over-

182   representation seen in the medulla-based adrenergic (HBAR) and noradrenergic (HBNOR) groups and

183   the dopaminergic neurons of the midbrain in the Periaqueductal Gray (PAG) (MBDOP1) and the

Figure 7. GSEA and Dot plots for imprinted genes upregulated in neurons across the whole hypothalamus. (*A*) GSEA for imprinted genes upregulated in the 'Neuron' cell type in the whole hypothalamic dataset of Chen, Wu (4). See legend of Figure 3A for a description of how to interpret the plot. (*B*) Dot plot of imprinted genes upregulated in the 'Neuron' cell type plotted across all identified cell types in the Chen, Wu (4) whole hypothalamic dataset. See legend of Figure 3B for a description of how to interpret the plot. Abbr: OPC = Oligodendrocyte Precursor Cell, MG = Myelinating Oligodendrocyte, IMG = Immature Oligodendrocyte, Astro = Astrocyte, Epith = Epithelial, Macro = Macrophage, Tany = Tanycyte, Ependy = Ependymocyte, Micro = Microglia, POPC = Proliferating Oligodendrocyte Progenitor Cell. (*C*) GSEA for imprinted genes upregulated in 'neurons' in the whole hypothalamic dataset of Romanov, Zeisel (3). See legend of Figure 3A for a description of how to interpret the plot. (*D*) Dot plot of imprinted genes upregulated in 'neurons' plotted across all identified cell types in the Romanov, Zeisel (3) whole hypothalamic dataset. See legend of Figure 3B for a description of how to interpret the plot.

inhibitory (MEINH, HBIN) and excitatory neuron (MEGLU, HBGLU) types spread across the nuclei

from the three regions (Fig. 6). The serotonergic populations of the raphe nuclei of these regions

(HBSER) were particularly prominent since the pons and medulla-based serotonin neuron populations

(HBSER2, HBSER4 and HBSER5) were the only neuron subpopulations out of the 214 total to have a

significant GSEA for imprinted genes after correction (Supplemental Fig. S4).

Additional regions of over-representation included neurons in the pallidum and striatum and PVN

neurons from the thalamus. In this comparison of 214 neuron populations, no neurons from areas such

as the cortex, cerebellum or peripheral nervous system were enriched, and neither were they over-

represented in the previous regional analysis. Hence, further analysis focused on those brain regions

enriched in this whole brain level analysis.

Imprinted gene expression is over-represented in specific hypothalamic neuron subtypes (Level 3A&3B Analysis)

We next sought to investigate whether those regional neuron enrichments found within the whole

brain comparisons would be further clarified with enriched expression in specific neuronal

subpopulations within those regions. Namely we sought to identify neural populations enriched across

the whole hypothalamus and those enriched within specific hypothalamic nuclei, and also whether

imprinted gene expression was enriched in the other key subpopulation identified, the ventral

midbrain and hindbrain dopaminergic and serotonergic populations. Two datasets with single cell

sequencing data for the adult hypothalamus existed (3, 4). Both clustered their data into neuronal

subpopulations allowing us to look for convergent imprinted enrichment across major hypothalamic

neuronal subtypes (Level 3A). Analysis revealed a clear neuronal bias in expression of imprinted

genes (Supplemental Table S9A and S10A). Within the Romanov, Zeisel (3) data, there was a

significant over-representation of imprinted genes in neurons ($q = 0.02$) and a similar observation was

seen in the Chen, Wu (4) data ($q = 0.001$), and both also demonstrated a significant GSEA in neurons

(Fig. 7A-D, Romanov, Zeisel (3) – $p = 0.011$, Chen, Wu (4) - $p = 0.022$).

Within the Chen, Wu (4) dataset, 4/33 hypothalamic neuronal subtypes had a significant over-

representation of imprinted genes (Supplemental Table S9B). The four subtypes were all GABAergic

neurons, specifically: Galanin neurons (*Slc18a2/Gal)* present in a several hypothalamic regions ($q =$

0.0079); a dopaminergic neuron type (*Slc6a3*) with high expression of *Th* and *Prlr* suspected to be the

TIDA neurons of the arcuate nucleus ($q = 0.0001$); SCN neurons (*Vipr2)* with very high *Avp* and *Nms*

expression ($q = 0.0071$); and *Agrp* feeding promoting neurons of the Arcuate Nucleus ($q = 0.034$).

Within the Romanov, Zeisel (3) dataset, 3/62 subtypes had significant over-representation of

imprinted gene expression (Supplemental Table S10B): A*grp/Npy* neurons ($q = 0.013$), the Arcuate

Nucleus feeding neurons also reported in Chen, Wu (4); a *Ghrh/Th* neuronal type ($q = 0.032$), again

likely corresponding to neurons from the arcuate nucleus and the top hit ($q = 1.63 \times 10^{-6}$) was a poorly

segregated population (*Calcr/Lhx1*), likely due to a deeper inner cluster heterogeneity. This cluster

was interesting since the imprinted genes *Calcr* and *Asb4* were amongst its most significant marker

genes, and it was notably the only cluster with high expression of all three of *Th*, *Slc6a3* and *Prlr*.

Romanov, Zeisel (3) did not identify any of their populations as the TIDA neurons, but the above

pattern of gene expression suggests that this cluster may contain these neurons. Furthermore, the

suspected TIDA neurons from the Chen, Wu (4) dataset shared 21/40 upregulated genes of this

unresolved cluster (see Supplemental Table S11 for full comparison);

Having consistently found well-known neurons from the arcuate nucleus (*Agrp, Ghrh*), and

suprachiasmatic nucleus (*Avp, Vip*) we sought to test imprinted gene enrichment within these

hypothalamic regions at a high resolution using datasets sequencing neurons purely from these

hypothalamic regions (Level 3B).

Arcuate nucleus (ARC) (2)

The first nuclei investigated was the ARC sequenced by Campbell, Macosko (2). Imprinted gene

over-representation was found in 8/24 arcuate neuron types (Supplemental Table S12). These

included the *Agrp/Sst* neuron type (with high expression of *Npy, q = 0.003*) and two *Pomc* neuron

types (*Pomc/Anxa2*, *q* = 0.004; *Pomc/Glipr1*, *q* = 0.03). Pomc expressing neurons are known to work

as feeding suppressants (38). Additional significant over-representation was found in the *Ghrh* neuron

type (*q* = 0.009), which was also enriched in *Gal* and *Th*. Finally, a highly significant over-

representation of imprinted genes was found in the *Th*/*Slc6a3* neuron type (*q* = 1.72 x 10^{-8}) identified

by the authors as one of the most likely candidates for the TIDA dopaminergic neuron population.

Marker genes for this identity group overlapped with the TIDA candidates from the previous two

datasets (e.g., *Slc6a3*, *Th*, *Lhx1*, *Calcr*). *Agrp* neurons, *Ghrh* neurons and these TIDA candidate

neurons were identified in both whole hypothalamic datasets and at the nuclei level.

Suprachiasmatic Nucleus (SCN) (5)

Analysis of the 10x chromium data of SCN neurons (Supplemental Table S13) revealed a significant

over-representation (*q* = 1.51 x 10^{-8}) and GSEA (*p* = 0.004, Supplemental Fig. S5) in the *Avp*/*Nms*

neuronal cluster (out of 5 neuronal clusters). This cluster shows the strongest expression for *Oxt, Avp*,

Avpr1a and *Prlr* and is one of the three neural group that Wen, Ma (5) found had robust circadian

gene expression, and the only subtype with notable phase differences in circadian gene expression in

the dorsal SCN. This cluster likely corresponds to the GABA8 cluster found enriched in the Chen, Wu

(4) dataset. Figure 8 presents the overlapping upregulated imprinted genes from the convergently

upregulated neuron subtypes

in the hypothalamic analysis

of Level 3a and 3b.

Imprinted gene expression

is over-represented in

monoaminergic nuclei of

the mid- and hindbrain

(Level 3C Analysis)

In the MBA, Whole

Hypothalamus and Arcuate

Nucleus analyses,

Figure 8. Venn diagrams of upregulated imprinted genes in the neuronal subpopulations from level 3b that were also identified in level 2 and 3a. Imprinted gene overlap was contrasted for *Agrp/Sst* neuronal populations of the Arcuate Nucleus (1-4) and *Avp/Nms* neurons from the Suprachiasmatic Nucleus (1, 4, 5) Imprinted genes are listed which show significant upregulation (*q* ≤ 0.05 and Log2FC > 0) in the subpopulation. Parental-bias is indicated by colour (MEG - red, PEG - blue).

dopaminergic clusters were consistently enriched and, to explore this further, analysis of Hook,

McClymont (39) data allowed comparison for dopamine neurons across the brain (specifically from

the olfactory bulb, arcuate nucleus and midbrain) at two developmental timepoints (E15.5 and Post-

natal day (P) 7). The arcuate nucleus P7 dopamine neurons emerged as the clearest over-represented

subgroups (Supplemental Table S14). This included the *Th*/*Slc6a3*/*Prlr* neurons ($q = 1.15 \times 10^{-8}$) and

the *Th/Ghrh/Gal* cluster ($q = 4.79 \times 10^{-5}$) the latter of which were referred to as 'neuroendocrine' cells

by Hook, McClymont (39), and the former a mixture of arcuate nucleus populations with *Prlr* was

one of the marker genes, suggesting this includes the TIDA neurons. Additionally, P7 midbrain

neurons were the other group with significant over-representation (specifically from the PAG and

VTA) as well as the neuroblasts at this time point.

Although no specific adult mouse midbrain datasets exist, ventral midbrain sequencing at E11.5 -

E18.5 by La Manno, Gyllborg (40) allowed us to identify imprinted enrichment within the midbrain at

a timepoint when the major neuronal populations are differentiating but still identifiable

(Supplemental Table S15). As anticipated, we found significant over-representation in both mature

(DA1; high *Th* and *Slc6a3,* $q = 0.0103$), and developing (DA0, $q = 0.0129$) dopaminergic neurons, as

well as the serotonergic neurons ($q = 3.09 \times 10^{-7}$), likely from the midbrain raphe nuclei.

Raphe nuclei from the midbrain/hindbrain are key serotonergic regions of the brain. Analysis of all

cell types in the Dorsal Raphe Nucleus (DRN) sequenced by Huang, Ochandarena (41) revealed a

clear enrichment of imprinted genes in the neuronal populations of the DRN as compared to the non-

neuronal cell populations of the DRN (Supplemental Table S16A). When compared to all other cell

populations, significant ORA was seen for Dopaminergic ($q = 0.009$), Serotonergic ($q = 0.012$) and

Peptidergic neurons ($q = 0.0008$), however, a significant GSEA was found for all five neuronal

populations (Supplemental Fig. S6). When compared against each other (i.e., serotonergic

upregulation vs. the other neurons), only the serotonergic neurons of the DRN ($q = 0.0019$) were

found to have a significant over- representation of imprinted genes (Supplemental Table S16B).

GSEA's were non-significant but the mean fold change for imprinted genes was markedly higher in

both serotoninergic (52% higher) and dopaminergic neurons (68% higher). When contrasting

neuronal subpopulations of the DRN, two of the five serotonin subpopulations had significant over-

representation of imprinted genes: *Hcrtr1/Asb4* (q = 0.0014) and *Prkcq/Trh* (q = 0.007)

(Supplemental Table S16C). These clusters were identified by Huang, Ochandarena (41) as the only

clusters localised in the dorsal/lateral DRN and the serotonin clusters enriched in *Trh*. Huang,

Ochandarena (41) hypothesised that these were the serotonin neurons that project to hypothalamic

nuclei, and motor nuclei in the brainstem (as opposed to cortical/striatal projection).

Imprinted gene expression is over-represented in lactotrophs and somatotrophs of the pituitary

gland (Level 3D Analysis)

Following on from the enrichment seen above for imprinted gene expression in the dopaminergic

arcuate nucleus neurons coordinating pituitary gland output, we sought to identify whether any cells

in the pituitary would display matching over-representation for imprinted gene expression (Level 3D).

The pituitary was not sequenced as part of the multi-organ or whole brain datasets analysed above and

so two independent datasets were analysed that specifically sequencing the mouse pituitary at single

cell resolution. Ho, Hu

(8) recently sequenced

the anterior pituitary

gland of male and

female C57BL/6 mice

using two sequencing

technologies, both 10X

genomic and Drop-Seq.

This identified a variety

of cell types from the

endocrine and non-

endocrine pituitary. We

analysed data from

both technologies and

found that imprinted

Figure 9. *(A)* Pituitary cell types showing over-representation for imprinted gene expression in multiple pituitary datasets. Over-represented cell types are bold and not in greyscale. The hormone/s released from the endocrine cell types are also indicated. *(B)* Venn diagram of upregulated imprinted genes in the Somatotrophs and Lactotrophs in Cheung, George (7) and Ho, Hu (8). Imprinted genes are listed which show significant upregulation ($q \leq 0.05$ and Log2FC > 0) in the cell types. Parental bias is indicated by colour (MEG –red, PEG – blue). Genes in common from two analyses are presented in bold and totalled in each section of the Venn Diagram, while genes found upregulated in one analysis but not available in the others are included in small font and the number indicated in brackets.

gene expression was convergently over-represented in the Lactotrophs (prolactin secreting) and Somatotroph (growth hormone secreting) cells (Supplemental Table S17A & 17B). In a second independent dataset sequencing cells from male mouse pituitary glands (7), we found significant over-representation in the Somatotropes and Thyrotrope (secreting thyroid stimulating hormone). Figure 9 demonstrates the overlap in imprinted genes significantly expressed in Somatotropes and Lactotropes across the datasets since these were the only cell-types to be over-represented in more than one dataset (Supplemental Table S18). It is notable that the two cell types represented here directly match the two regulatory neurons found over-represented in the arcuate nucleus of the hypothalamus.

DISCUSSION

Using publicly available single cell transcriptomics data, we apply an unbiased systems biology approach to examine the enrichment of imprinted genes at the level of the brain in comparison to other adult tissues, refining this analysis to specific brain regions and then to specific neuronal populations. We confirm a significant over-representation in the brain, specifically in neurons at every level tested, with a marked enrichment in neuroendocrine cells lineages. Within-brain analyses revealed that the hypothalamus and the monoaminergic system of the mid- and hindbrain were foci for imprinted gene enrichment. While not all imprinted genes follow these patterns of expression, these findings highlight collective gene expression which is non-random in nature. As such, these analyses identify 'expression hotspots', which in turn suggest 'functional hotspots'. Specifically, our results at the systems and cellular level highlight a major role for imprinted genes in the neuronal regulation of pituitary function, feeding and sleep.

Some of the earliest studies of genomic imprinting identified the brain as a key area for imprinted gene expression (20, 32). However, it is estimated that ~80% of the genome is expressed in the brain and consequently, imprinted gene expression here may not be a purposeful phenomenon. Our current analysis definitively show that imprinted genes were significantly over-represented in the brain as a whole. This over-representation was found again with PEGs alone, but not MEGs. Critically, over-representation was also found when limiting our analysis to genes that have been found to be imprinted in non-brain tissues (Supplemental Table S19). Excluding those genes imprinted in the

Within specific brain regions, imprinted genes were over-represented in the hypothalamus, ventral

midbrain, pons and medulla. This confirms some findings from studies of Pg/Gg and Ag chimera

studies (20, 32) and summaries of imprinted gene expression (16). However, unlike these earlier

studies, our analyses do not simply ask if imprinted genes are expressed (at any level) or not, but

robustly test whether this expression is meaningful, and the expression of these genes are especially

enriched in any given brain region. Additionally, in the chimera studies, Pg/Gg cells (two maternal

genomes) preferentially allocated to the developing adult cortex and hippocampus, and Ag cells (two

paternal genomes) preferentially allocated to the developing hypothalamus and midbrain. Our analysis

does not reproduce this distinct pattern of MEG and PEG expression in the brain, and indeed we find

no specific enrichment of imprinted genes in cortex or hippocampus. Although the pattern of regional

enrichment seen with all imprinted genes is replicated when analysing PEGs alone, separate analysis

of MEGs only shows over-representation in the pons and medulla. This difference between our

findings and the Pg/Gg and Ag chimeras studies could indicate that the distribution of Pg/Gg and Ag

cells in the brain is not driven by, or indeed reflective of adult PEG and MEG expression, but instead

is determined by expression of specific imprinted genes during brain development (42).

At the whole brain level, mature neurons and, in particular, neural-lineage neuroendocrine cells, had

disproportionately higher numbers of imprinted genes expressed, and high levels of imprinted gene

expression. It is likely that this neural-lineage neuroendocrine population comprises members of the

key hypothalamic populations in which the expression of imprinted genes are enriched and, when

treated as their own cluster, demonstrate strong imprinted gene enrichment compared to other cell

lineages of the brain, even other mature neurons.

Within the hypothalamus, a selection of informative neuronal subpopulations were over-represented.

Strikingly, and suggestive of meaningful enrichment, we saw convergence across our different levels

of analysis with several key neuronal types identified in the whole hypothalamus and/or

hypothalamic-region-level, already having been identified against the background of general

imprinted gene expression in the whole-brain-level analysis. These subpopulations are collectively

associated with a few fundamental motivated behaviours. We consistently saw enriched imprinted

gene expression in *Agrp* expressing neurons when contrasting neurons across the whole brain, whole

hypothalamus and within the arcuate nucleus. *Agrp* neurons from the arcuate nucleus are well known

feeding promotors and a few imprinted genes have previously been associated with their function

(*Asb4, Magel2, Snord116*) (43, 44) but never as an enriched population. Feeding was further linked

with imprinting through enrichment seen in *Pomc+* neurons (45) as well as *Hcrt+* and *Gal+* neurons.

Circadian processes are controlled principally by the Suprachiasmatic Nucleus and here we find

strong imprinted gene enrichment in *Avp/Nms* expressing neurons (an active circadian population).

Again, these were found to be enriched when contrasting neurons across the whole brain, whole

hypothalamus and within the SCN. This population is of interest given the growing appreciation of

the role imprinted genes play in circadian processes and the SCN suggested by studies of individual

imprinted genes (46). Pituitary endocrine regulation also emerged as a key function, considering the

over-representation in the dopaminergic: *Th/Slc6a3/Prlr* neuron type (top hit in the arcuate nucleus

and across dopaminergic neurons of the brain) and the *Th/Ghrh* subpopulation. These neuron

populations can regulate prolactin (regulating lactation, stress, weight gain, parenting and more (47,

48)) and growth hormone (promoting growth and lipid/carbohydrate metabolism) release,

respectively. Remarkably, we also found a matching enrichment in the lactotroph and somatotroph

cells in the pituitary. A role for imprinted genes in pituitary function is well known (49, 50), with

pituitary abnormalities associated with imprinted disorders such as PWS (51) and recent sequencing

work showing imprinted genes are amongst the highest expressed transcripts in the mature and

developing pituitary (52). Specific genes highly expressed here, such as Dlk1 and Nnat, have been

shown to alter somatotroph phenotypes (53, 54). Finally, we saw enrichment in galanin expressing

neuronal populations (found enriched when contrasting neurons across the whole brain, whole

hypothalamus). Galanin neurons in the hypothalamus have a diverse set of functions including

subpopulations for thermoregulation, feeding, reproduction, sleep and parenting behaviour (55, 56),

contributing to this consistent picture of IGs associating with neurons key for motivated behaviour.

In this analysis the hypothalamus was a clear hot spot for imprinted gene expression, in line with the

prevailing view of imprinted gene and hypothalamic function (50, 57). However, outside of the

hypothalamus other distinct hotspot emerged from our whole brain analysis including the

monoaminergic system of the midbrain/hindbrain. Analysing data from the dorsal raphe nucleus and

ventral midbrain revealed the dopaminergic and serotonergic neurons to be a foci of imprinted gene

expression within this region. These midbrain dopamine neurons were enriched when contrasted to

other dopamine neurons from the brain and the enriched serotonergic neurons were those that project

to the subcortical regions of the brain known to be associated with feeding and other motivated

behaviours (58), providing convergence with the functional hotspots seen in the hypothalamus.

Analyses of these kind are always bound by the available data and therefore there are notable

limitations and caveats to this study. The aim of this study was to generate information about

'hotspots' of imprinted gene expression. This approach, and the use of over-representation analysis

and GSEA, therefore do not provide an exhaustive list of sites of expression, and non-differentially

expressed genes could still be highly expressed genes despite not contributing to this analysis. An

example of a known site of expression for imprinted genes not found to be enriched in our analysis

was the oxytocin neurons of the hypothalamus, since a clear oxytocin neuron phenotype has been

reported in a handful of imprinted gene models (24, 59). This may be an example of a functional

effect occurring below the level of over-representation, or that imprinted genes act during

development and are not functionally enriched in adult oxytocin neurons, or simply that compared to

other hypothalamic neuronal populations, oxytocin neurons are not a 'hotspot' of imprinted

expression. Specific sequencing of oxytocinergic brain regions will be required to distinguish between

these possibilities. A second caveat is that, due to the nature of the datasets used, not all imprinted

genes were included, and our analysis was missing a significant subset of imprinted genes encoding

small RNAs or isoforms from the same transcription unit. A third caveat is that we did not assess

parent-of-origin expression for the 119 imprinted genes we included in the analysis. Previous

expression profiling of imprinted genes have also not measured the POEs (16, 60) but have restricted

their gene selection to genes with reliable imprinting status. Consequently, we only included the

canonical imprinted genes and genes with more than one demonstration of a POE when looking for

enrichment. Furthermore, for the vast majority of these genes, a brain-based POE effect has also

already been reported (Supplemental Table S1). Although this does not replace validating the

imprinting status of all 119 in the tissues and subregions examined, it does provide justification for

looking at imprinted gene over-representation. To resolve this issue, scRNA-seq using tissues derived

from reciprocal F1 crosses between distinct mouse lines will be key; for example, the recent work of

(61) with cortical cell types provides an example of the allelic specific single-cell expression

measurements necessary to confirm the enrichments found in this study.

By exploiting scRNA-seq data we have asked whether imprinted genes as a group are

disproportionately represented in the brain, in specific brain regions, and in certain neuronal cell-

types. In the adult brain imprinted genes were over-represented in neurons, and particularly the

hypothalamic neuroendocrine populations and the monoaminergic hindbrain neurons, with the

serotonergic neurons demonstrating the clearest signal. Interestingly, PEGs, but not MEGs, recreate

this signal at Levels 1 and 2 - most notably only PEGs display the hypothalamic neuronal enrichment.

By extension, these data also identify behaviours that are foci for the action of imprinted genes.

Although there are high profile examples of individual imprinted genes expressed in the key brain

regions we highlight and that have roles in feeding (*Magel2*) (62) and sleep (*Snord116*) (63), our

analyses indicate that imprinted genes *as a group* are strongly linked to these behaviours and also

identify other individual genes that should be explored in these domains. Conversely, there are high-

profile examples of imprinted genes involved in hippocampus related learning and memory (*Ube3a*)

(28), but we did not find enrichment for cell types related to this brain function. The idea that

imprinted genes converge on specific physiological or behavioural processes is not unprecedented.

Specialisation of function is predicted when considering why genomic imprinting evolved at all (13,

64-66). Moreover, there is increasing evidence that the imprinted genes themselves appear to be co-

expressed in an imprinted gene network (IGN) and have confirmed regulatory links between each

other (67-69). The idea of an IGN or, at the very least, heavily correlated and coordinated expression

between imprinted genes adds further support to the idea that imprinted genes work in concert, rather

than in isolation, to influence processes and that perturbating one may influence many others (70).

Our findings add substance to these general ideas and highlight the neuronal regulation of pituitary

function, feeding and sleep as being key functional hotspots on which imprinted genes converge

which probably provides the best current basis for discerning evolutionary drivers of genomic

imprinting in the brain.

METHODS

Data Processing

13 unique datasets were analysed across the three levels of analysis (see Fig.1) and analyses were

conducted on each dataset independently. At each level of analysis, we aimed to be unbiased by using

all the datasets that fitted the scope of that level, but the availability of public scRNA-seq datasets was

limited, which prevented us from exploring all avenues (for example, a direct comparison of

enrichment between hypothalamic nuclei). All sequencing data were acquired through publicly

available resources and each dataset was filtered and normalised according to the original published

procedure. Supplemental Table S20 details the basic parameters of each dataset. Once processed, each

dataset was run through the same basic workflow (see below and Fig. 10), with minor adjustments

laid out for each dataset detailed in the Supplemental Methods.

Due to the high variability in sequencing technology, mouse strain, sex and age, and processing

pipeline, we have avoided doing

analysis on combined datasets.

Rather we chose to perform our

analyses independently for each

dataset and look for convergent

patterns of imprinted gene

enrichment between datasets on

similar tissues/brain regions. As

with any single-cell experiment,

the identification of upregulation

or over-representation of genes

in a cell-type depends heavily

on which other cells are

Figure 10. Basic workflow schematic. Single Cell Expression Matrices were acquired through publicly available depositories. Data were processed according to the author's original specifications and all genes were required to be expressed in 20 or more cells. Cell population identities were acquired from the author's original clustering. Positive differential gene expression was calculated via Wilcoxon Rank-Sum Test. Upregulated genes were considered as those with $q \leq 0.05$ and a Log2FC ≥ 1 for analysis levels 1 and 2, while this criterion was relaxed to Log2FC > 0 for level 3. Our imprinted gene list was used to filter

included in the analysis to make up the 'background'. Analysing separate datasets (with overlapping

cell-types alongside distinct ones) and looking for convergent patterns of enrichment is one way of

counteracting this limitation.

Basic Workflow

Data were downloaded in the available form provided by the original authors (either raw or

processed) and, where necessary, were processed (filtered, batch-corrected and normalized) to match

the author's original procedure. Cell quality filters were specific to each dataset and summarised in

Supplemental Table S20. A consistent filter, to remove all genes expressed in fewer than 20 cells, was

applied to remove genes unlikely to play a functional role due to being sparsely expressed. Datasets of

the whole brain/hypothalamus were analysed both at the global cell level (neuronal and non-neuronal

cells) and neuron specific level (only neurons) with genes filtered for the ≥ 20 cell expression at each

level before subsequent analysis. Cell identities were supplied using the outcome of cell clustering

carried out by the original authors, so that each cell included in the analysis had a cell-type or tissue-

type identity. This was acquired as metadata supplied with the dataset or as a separate file primarily

from the same depository as the data but occasionally acquired from personal correspondence with the

authors. Cells were used from mice of both sexes when provided and all mice were aged 15 weeks or

younger across all datasets. Although our focus was the adult mouse brain, embryonic data were

included in some comparisons or when no alternatives were present. However, embryonic and post-

natal cells were never pooled to contribute to the same cell populations.

Positive differential expression between identity groups were carried out using one-sided Wilcoxon

rank-sum tests (assuming the average expression of cells within the current identity group is 'greater'

than the average of cells from all other groups). The test was performed independently for each gene

and for each identity group vs. all other groups. The large number of p values were corrected for

multiple comparisons using a horizontal Benjamini-Hochberg correction, creating q values. Fold-

change (FC) values, percentage expression within the identity group and percentage expressed within

the rest were also calculated. We considered genes to be significantly positively differentially

expressed (significantly upregulated) in a group compared to background expression if it had a $q \leq$

0.05. In addition, for Level 1 and Level 2 analyses, the criteria for upregulated genes included

demonstrating a Log2FC value of 1 or larger (i.e., 2-fold-change or larger). The datasets at these

levels represented cells from a variety of organs, regions and cell-types, and in line with this cellular

diversity, the aim of these analyses was to look for distinctive upregulation, akin to a marker gene.

Once the analysis was restricted to cell subpopulations within a specific region of the brain (i.e., Level

3), the additional criteria for upregulation was relaxed to demonstrating just a positive Log2FC (i.e.,

the gene has a higher expression in this cell type than background). This was mainly because we were

not expecting imprinted genes to be 'markers' of individual subpopulations at this level, but our aim

was to identify enriched expression profiles for them. This additionally ensures consistent criteria for

enrichment within levels, allowing meaningful comparison.

The same custom list of imprinted genes with reliable parent-of-origin effects (see below) was used

for all analyses, and all genes were included as long as the gene passed the 20-cell filter. The first

statistical analysis for enrichment was an Over-Representation Analysis (ORA) using a one-sided

Fisher's Exact Test ('fisher.test' function in R core package 'stats v3.6.2'). The aim was to assess

whether the number of imprinted genes considered to be upregulated as a proportion of the total

number of imprinted genes in the dataset (passing the 20-cell filter) was statistically higher than

would be expected by chance when compared to the total number of upregulated genes as a

proportion of the overall number of genes in the dataset (passing the 20-cell filter). To limit finding

over-represented identity groups with only a few upregulated imprinted genes, an identity group was

required to have ≥ 5 % of the total number of imprinted genes upregulated for ORA to be conducted.

Subsequent p-values for all eligible identity groups were corrected using a Bonferroni correction. This

provided a measure of whether imprinted genes are expressed above expectation (as opposed to the

expression pattern of any random gene selection) in particular identity groups.

Venn diagrams of the upregulated imprinted genes making up over-represented identity groups across

datasets (within a level) were also reported. Full lists of upregulated imprinted genes can be found in

the 'Upregulated_IGs.csv' file for each analysis in the Supplemental Data.

To further examine the presence of imprinted genes within tissues/cell types, and to provide a

different perspective to over-representation, we conducted a Gene-Set Enrichment Analysis (GSEA)

for imprinted genes amongst the upregulated genes of an identity group using a publicly available,

light-weight implementation of the GSEA algorithm (71) in R (https://github.com/JEFworks/liger).

This was done in a manner similar to Moffitt, Bambah-Mukku (72) since we were similarly using this

computational method to identify enrichment of our gene sets *within the upregulated genes* of the

different identity groups. Here, the GSEA was conducted for each individual identity group using

Log2FC values to rank the upregulated genes. The GSEA acts as a more conservative measure than

the ORA since it tests whether imprinted genes are enriched in the stronger markers of a group (the

genes with the highest fold change for a group vs. the rest) and hence whether the imprinted genes are

enriched in those genes with a high specificity to that tissue/cell type. To prevent significant results

being generated from just 2 or 3 genes, identity group to be analysed were selected as having a

minimum of 15 upregulated imprinted genes (i.e. the custom gene set) to measure enrichment for (a

value suggested by the GSEA user guide (https://www.gsea-

msigdb.org/gsea/doc/GSEAUserGuide Frame.html)) and to prevent significant results in which

imprinted genes cluster at the tail, identity groups were selected as having an average fold change of

the upregulated imprinted genes greater than the average fold change of the rest of the upregulated

genes for that group. Again, multiple p values generated from GSEA were corrected using a

Bonferroni correction. To further elucidate the genes responsible for significant GSEA's, dot plots of

the imprinted genes upregulated in that identity group were plotted across all identity groups with

absolute expression and Log2FC mapped to size and colour of the dots, respectively. Graphical

representations of significant GSEA's (post-correction) are included in the main text or as

supplemental figures, all other graphs, including additional dot plots not discussed in this study, can

be found in the repository (https://osf.io/jx7kr/) and Supplemental Data. If no cell populations met

these criteria, GSEA was not run and not included for that analysis.

For Level 1 and Level 2 analyses, we also carried out parent-of-origin specific analyses. The

imprinted gene list was divided into MEGs and PEGs and the analyses detailed above were run

separately for these two gene groups. For imprinted genes with known parent-of-origin variability

based on tissue type (*Igf2* and *Grb10*), the parent-of-origin characterisation of these genes was

changed accordingly. The absolute number of imprinted genes top-expressed in a tissue/cell-type were

also reported for analyses in Level 1 and Level 2 in the tables, since these analyses included a variety

of cell-types and tissues which may demonstrate meaningful clustering of the highest normalised

expression values. The mean normalised expression for all imprinted genes across the series of

identity groups in the datasets in Level 1 and Level 2 was also calculated alongside the mean

normalised expression for the rest of the genes (Supplemental Table S2).

All graphical representations and statistical analyses were conducted using R 3.6.2 (73) in RStudio

(74). Diagrams in Figures 1:2, 4:6 and 8:10 were created with BioRender.com.

Custom Imprinted Gene List

The gene list for the analysis was based on the list of murine imprinted genes recently published in

Tucci, Isles (10). Although the original list of imprinted genes was 260 genes long, only 163 genes

were identified in the most comprehensive of the datasets. We further refined this list to 119 imprinted

genes (Supplemental Table S1a) which excluded the X-linked genes, consisting of mostly the

canonical protein-coding and long noncoding RNA imprinted genes, but the criteria for inclusion was

those genes with at least two independent demonstrations of their POE status (See Supplemental

Table S1b for full list of 260 imprinted genes and reasons for gene exclusion). The only exceptions to

multiple independent demonstrations of a POE were four genes (*Bmf*, *B3gnt2*, *Ptk2*, *Gm16299*)

identified by (34) where a POE was assessed across 16 brain regions and 7 adult tissues within one

study. For Level 2, the MEG/PEG status of a gene was primarily based on reported allelic expression

within the brain. Small non-coding RNAs such as micro-RNAs (miRs) and small nucleolar RNAs

(snoRNAs), which represent ~10% of identified imprinted genes, were excluded from the analysis as

their sequences were not detected/subsumed by larger transcripts in the majority of the datasets.

Another caveat with short-read RNA-seq libraries is that much of the expression data for a given

transcription unit cannot discriminate differentially imprinted isoforms nor do some of the

technologies (e.g., Smart-Seq2) possess stranded libraries to distinguish antisense transcripts. For

complex imprinting loci such as the *Gnas* locus, most reads as result map to only *Gnas* and *Nespas*

ignoring several overlapping and antisense genes.

DECLARATIONS

Ethics approval and consent to participate

Not applicable. All samples had been collected in the context of previous studies.

Consent for publication

Not applicable.

Competing interests

All authors declare no financial and non☐financial competing interests.

Availability of data and materials

The datasets analysed during the current study were acquired from publicly available resources and are available in the following GEO repositories, Mouse Cell Atlas – GSE108097, *Tabula Muris* – GSE109774, Aging Mouse Brain – GSE129788, Hypothalamus (Chen) – GSE87544, Hypothalamus (Romanov) – GSE74672, Arcuate Nucleus – GSE93374, Suprachiasmatic Nucleus – GSE132608, Dopamine Neurons – GSE108020, Ventral Mid Brain – GSE76381, Dorsal Raphe Nucleus – GSE134163, Pituitary Gland (Ho) - GSE146619, Pituitary Gland (Cheung) - GSE120410 and the following SRA repository, Mouse Brain Atlas – SRP135960. The data generated in this experiment is provided as Supplemental Data and in an Open Science Framework repository entitled – "Imprinted Gene Enrichment at Single-Cell Resolution" (https://osf.io/jx7kr/). Custom R scripts to analyse each dataset are provided as Supplemental Code and are available at https://github.com/MJHiggs/IG-Single-Cell-Enrichment.

Funding

This work was supported by a Wellcome Trust PhD studentship (220090/Z/20/Z).

Authors' Contributions

MJHiggs performed bioinformatic analysis, with input from MJHill; MJHiggs., and ARI contributed to project design, data interpretation, and wrote the manuscript, MJHiggs produced all Figures; all co-authors reviewed and edited the manuscript.

Acknowledgements

617   We would like to thank all the research groups that carried out the single-cell RNA sequencing that

618   made this study possible and to particularly acknowledge Dr. L. Cheung, Dr. P. Hook, Dr. A. Jackson,

619   and Dr. S. Wen for help accessing the cell metadata for their associated studies.

REFERENCES

1.   Zeisel A, Hochgerner H, Lönnerberg P, Johnsson A, Memic F, Van Der Zwan J, et al. Molecular architecture of the mouse nervous system. Cell. 2018;174(4):999-1014. e22.

2.   Campbell JN, Macosko EZ, Fenselau H, Pers TH, Lyubetskaya A, Tenen D, et al. A molecular census of arcuate hypothalamus and median eminence cell types. Nature neuroscience. 2017;20(3):484-96.

3.   Romanov RA, Zeisel A, Bakker J, Girach F, Hellysaz A, Tomer R, et al. Molecular interrogation of hypothalamic organization reveals distinct dopamine neuronal subtypes. Nature neuroscience. 2017;20(2):176-88.

4.   Chen R, Wu X, Jiang L, Zhang Y. Single-cell RNA-seq reveals hypothalamic cell diversity. Cell reports. 2017;18(13):3227-41.

5.   Wen S, Ma D, Zhao M, Xie L, Wu Q, Gou L, et al. Spatiotemporal single-cell analysis of gene expression in the mouse suprachiasmatic nucleus. Nature neuroscience. 2020;23(3):456.

6.   Ximerakis M, Lipnick SL, Innes BT, Simmons SK, Adiconis X, Dionne D, et al. Single-cell transcriptomic profiling of the aging mouse brain. Nature neuroscience. 2019;22(10):1696-708.

7.   Cheung LY, George AS, McGee SR, Daly AZ, Brinkmeier ML, Ellsworth BS, et al. Single-cell RNA sequencing reveals novel markers of male pituitary stem cells and hormone-producing cell types. Endocrinology. 2018;159(12):3910-24.

8.   Ho Y, Hu P, Peel MT, Chen S, Camara PG, Epstein DJ, et al. Single-cell transcriptomic analysis of adult mouse pituitary reveals sexual dimorphism and physiologic demand-induced cellular plasticity. Protein & cell. 2020;11(8):565-83.

9.   Ferguson-Smith AC. Genomic imprinting: the emergence of an epigenetic paradigm. Nature Reviews Genetics. 2011;12(8):565-75.

10.   Tucci V, Isles AR, Kelsey G, Ferguson-Smith AC, Bartolomei MS, Benvenisty N, et al. Genomic imprinting and physiological processes in mammals. Cell. 2019;176(5):952-65.

11.   Orr HA. Somatic mutation favors the evolution of diploidy. Genetics. 1995;139(3):1441-7.

12.   Peters J. The role of genomic imprinting in biology and disease: an expanding view. Nature Reviews Genetics. 2014;15(8):517-30.

13.   Moore T, Haig D. Genomic imprinting in mammalian development: a parental tug-of-war. Trends in genetics. 1991;7(2):45-9.

14.   Andergassen D, Dotter CP, Wenzel D, Sigl V, Bammer PC, Muckenhuber M, et al. Mapping the mouse Allelome reveals tissue-specific regulation of allelic expression. Elife. 2017;6:e25125.

15.   Babak T, DeVeale B, Tsang EK, Zhou Y, Li X, Smith KS, et al. Genetic conflict reflected in tissue-specific maps of genomic imprinting in human and mouse. Nature genetics. 2015;47(5):544-9.

16.   Gregg C, Zhang J, Weissbourd B, Luo S, Schroth GP, Haig D, et al. High-resolution analysis of parent-of-origin allelic expression in the mouse brain. science. 2010;329(5992):643-8.

17.   Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, et al. Genome-wide atlas of gene expression in the adult mouse brain. Nature. 2007;445(7124):168-76.

18.   Negi SK, Guda C. Global gene expression profiling of healthy human brain and its application in studying neurological disorders. Scientific reports. 2017;7(1):1-12.

19.   Cattanach BM, Kirk M. Differential activity of maternally and paternally derived chromosome regions in mice. Nature. 1985;315(6019):496-8.

20.   Keverne EB, Fundele R, Narasimha M, Barton SC, Surani MA. Genomic imprinting and the differential roles of parental genomes in brain development. Developmental Brain Research. 1996;92(1):91-100.

21.   Angulo M, Butler M, Cataletto M. Prader-Willi syndrome: a review of clinical, genetic, and endocrine findings. Journal of endocrinological investigation. 2015;38(12):1249-63.

22.   Nicholls RD, Knoll JH, Butler MG, Karam S, Lalande M. Genetic imprinting suggested by maternal heterodisomy in non-deletion Prader-Willi syndrome. Nature. 1989;342(6247):281-5.

23.   Perez JD, Rubinstein ND, Dulac C. New Perspectives on Genomic Imprinting, an Essential and Multifaceted Mode of Epigenetic Control in the Developing and Adult Brain. Annual Review of Neuroscience. 2016;39(1):347-84.

24.   Li L-L, Keverne E, Aparicio S, Ishino F, Barton S, Surani M. Regulation of maternal behavior and offspring growth by paternally expressed Peg3. Science. 1999;284(5412):330-4.

25.   Davies JR, Humby T, Dwyer DM, Garfield AS, Furby H, Wilkinson LS, et al. Calorie seeking, but not
hedonic response, contributes to hyperphagia in a mouse model for Prader–Willi syndrome. European Journal of
Neuroscience. 2015;42(4):2105-13.
26.   McNamara GI, John RM, Isles AR. Territorial behavior and social stability in the mouse require
correct expression of imprinted Cdkn1c. Frontiers in behavioral neuroscience. 2018;12:28.
27.   Garfield AS, Cowley M, Smith FM, Moorwood K, Stewart-Cox JE, Gilroy K, et al. Distinct
physiological and behavioural functions for parental alleles of imprinted Grb10. Nature. 2011;469(7331):534-8.
28.   Jiang Y-h, Armstrong D, Albrecht U, Atkins CM, Noebels JL, Eichele G, et al. Mutation of the
Angelman ubiquitin ligase in mice causes increased cytoplasmic p53 and deficits of contextual learning and
long-term potentiation. Neuron. 1998;21(4):799-811.
29.   Dent CL, Humby T, Lewis K, Ward A, Fischer-Colbrie R, Wilkinson LS, et al. Impulsive choice in
mice lacking paternal expression of Grb10 suggests intragenomic conflict in behavior. Genetics.
2018;209(1):233-9.
30.   Relkovic D, Doe CM, Humby T, Johnstone KA, Resnick JL, Holland AJ, et al. Behavioural and
cognitive abnormalities in an imprinting centre deletion mouse model for Prader–Willi syndrome. European
journal of neuroscience. 2010;31(1):156-64.
31.   Lassi G, Ball ST, Maggi S, Colonna G, Nieus T, Cero C, et al. Loss of Gnas imprinting differentially
affects REM/NREM sleep and cognition in mice. PLoS Genet. 2012;8(5):e1002706.
32.   Allen ND, Logan K, Lally G, Drage DJ, Norris ML, Keverne EB. Distribution of parthenogenetic cells
in the mouse brain and their influence on brain development and behavior. Proceedings of the National
Academy of Sciences. 1995;92(23):10782-6.
33.   DeVeale B, Van Der Kooy D, Babak T. Critical evaluation of imprinted gene expression by RNA–Seq:
a new perspective. PLoS Genet. 2012;8(3):e1002600.
34.   Perez JD, Rubinstein ND, Fernandez DE, Santoro SW, Needleman LA, Ho-Shing O, et al. Quantitative
and functional interrogation of parent-of-origin allelic expression biases in the brain. Elife. 2015;4:e07860.
35.   Han X, Wang R, Zhou Y, Fei L, Sun H, Lai S, et al. Mapping the mouse cell atlas by microwell-seq.
Cell. 2018;172(5):1091-107. e17.
36.   Schaum N, Karkanias J, Neff NF, May AP, Quake SR, Wyss-Coray T, et al. Single-cell transcriptomics
of 20 mouse organs creates a Tabula Muris: The Tabula Muris Consortium. Nature. 2018;562(7727):367.
37.   Zhang X, Jiang S, Mitok KA, Li L, Attie AD, Martin TFJ. BAIAP3, a C2 domain–containing Munc13
protein, controls the fate of dense-core vesicles in neuroendocrine cells. Journal of Cell Biology.
2017;216(7):2151-66.
38.   Rau AR, Hentges ST. The relevance of AgRP neuron-derived GABA inputs to POMC neurons differs
for spontaneous and evoked release. Journal of Neuroscience. 2017;37(31):7362-72.
39.   Hook PW, McClymont SA, Cannon GH, Law WD, Morton AJ, Goff LA, et al. Single-cell RNA-seq of
mouse dopaminergic neurons informs candidate gene selection for sporadic Parkinson disease. The American
Journal of Human Genetics. 2018;102(3):427-46.
40.   La Manno G, Gyllborg D, Codeluppi S, Nishimura K, Salto C, Zeisel A, et al. Molecular diversity of
midbrain development in mouse, human, and stem cells. Cell. 2016;167(2):566-80. e19.
41.   Huang KW, Ochandarena NE, Philson AC, Hyun M, Birnbaum JE, Cicconet M, et al. Molecular and
anatomical organization of the dorsal raphe nucleus. Elife. 2019;8:e46464.
42.   Davies W, Isles AR, Wilkinson LS. Imprinted gene expression in the brain. Neuroscience &
Biobehavioral Reviews. 2005;29(3):421-30.
43.   Cassidy FC, Charalambous M. Genomic imprinting, growth and maternal–fetal interactions. Journal of
Experimental Biology. 2018;221(Suppl 1).
44.   Vagena E, Crneta J, Engström P, He L, Yulyaningsih E, Korpel NL, et al. ASB4 modulates central
melanocortinergic neurons and calcitonin signaling to control satiety and glucose homeostasis. Science
Signaling. 2022;15(733):eabj8204.
45.   Aponte Y, Atasoy D, Sternson SM. AGRP neurons are sufficient to orchestrate feeding behavior
rapidly and without training. Nature neuroscience. 2011;14(3):351.
46.   Tucci V. Genomic imprinting: a new epigenetic perspective of sleep regulation. PLoS genetics.
2016;12(5):e1006004.
47.   Grattan DR, Steyn FJ, Kokay IC, Anderson GM, Bunn SJ. Pregnancy□induced adaptation in the
neuroendocrine control of prolactin secretion. Journal of neuroendocrinology. 2008;20(4):497-507.
48.   Grattan D, Kokay I. Prolactin: a pleiotropic neuroendocrine hormone. Journal of neuroendocrinology.
2008;20(6):752-63.
49.   Davies W, Lynn PM, Relkovic D, Wilkinson LS. Imprinted genes and neuroendocrine function.
Frontiers in neuroendocrinology. 2008;29(3):413-27.
50.   Ivanova E, Kelsey G. Imprinted genes and hypothalamic function. Journal of molecular endocrinology.
2011;47(2):R67-R74.

51.   Miller JL, Goldstone AP, Couch JA, Shuster J, He G, Driscoll DJ, et al. Pituitary abnormalities in
Prader–Willi syndrome and early onset morbid obesity. American Journal of Medical Genetics Part A.
2008;146(5):570-7.
52.   Scagliotti V, Costa Fernandes Esse R, Willis TL, Howard M, Carrus I, Lodge E, et al. Dynamic
Expression of Imprinted Genes in the Developing and Postnatal Pituitary Gland. Genes. 2021;12(4):509.
53.   Charalambous M, Da Rocha ST, Radford EJ, Medina-Gomez G, Curran S, Pinnock SB, et al.
DLK1/PREF1 regulates nutrient metabolism and protects from steatosis. Proceedings of the National Academy
of Sciences. 2014;111(45):16088-93.
54.   Huerta-ocampo I, Slack R, Beechey C, Skinner J, Peters J, Christian H, editors. Overexpression of the
imprinted gene Neuronatin represses normal pituitary differentiation. Endocrine Abstracts; 2004: Bioscientifica.
55.   Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG. Galanin neurons in the medial preoptic
area govern parental behaviour. Nature. 2014;509(7500):325-30.
56.   Mechenthaler I. Galanin and the neuroendocrine axes. Cellular and molecular life sciences: CMLS.
2008;65(12):1826-35.
57.   Pulix M, Plagge A. Imprinted Genes and Hypothalamic Function. Developmental
Neuroendocrinology: Springer; 2020. p. 265-94.
58.   Donovan MH, Tecott LH. Serotonin and the regulation of mammalian energy balance. Frontiers in
neuroscience. 2013;7:36.
59.   Dombret C, Nguyen T, Schakman O, Michaud JL, Hardin-Pouzet H, Bertrand MJ, et al. Loss of
Maged1 results in obesity, deficits of social interactions, impaired sexual behavior and severe alteration of
mature oxytocin production in the hypothalamus. Human molecular genetics. 2012;21(21):4703-17.
60.   Steinhoff C, Paulsen M, Kielbasa S, Walter J, Vingron M. Expression profile and transcription factor
binding site exploration of imprinted genes in human and mouse. BMC Genomics. 2009;10:144.
61.   Laukoter S, Pauler FM, Beattie R, Amberg N, Hansen AH, Streicher C, et al. Cell-type specificity of
genomic imprinting in cerebral cortex. Neuron. 2020;107(6):1160-79. e9.
62.   Schaller F, Watrin F, Sturny R, Massacrier A, Szepetowski P, Muscatelli F. A single postnatal injection
of oxytocin rescues the lethal feeding behaviour in mouse newborns deficient for the imprinted Magel2 gene.
Human Molecular Genetics. 2010;19(24):4895-905.
63.   Lassi G, Priano L, Maggi S, Garcia-Garcia C, Balzani E, El-Assawy N, et al. Deletion of the
Snord116/SNORD116 alters sleep in mice and patients with Prader-Willi syndrome. Sleep. 2016;39(3):637-44.
64.   Keverne E. Significance of epigenetics for understanding brain development, brain evolution and
behaviour. Neuroscience. 2014;264:207-17.
65.   Keverne EB, Martel FL, Nevison CM. Primate brain evolution: genetic and functional considerations.
Proceedings of the Royal Society of London Series B: Biological Sciences. 1996;263(1371):689-96.
66.   Trivers R, Burt A. Kinship and genomic imprinting. Genomic imprinting: Springer; 1999. p. 1-21.
67.   Al Adhami H, Evano B, Le Digarcher A, Gueydan C, Dubois E, Parrinello H, et al. A systems-level
approach to parental genomic imprinting: the imprinted gene network includes extracellular matrix genes and
regulates cell cycle exit and differentiation. Genome research. 2015;25(3):353-67.
68.   Varrault A, Gueydan C, Delalbre A, Bellmann A, Houssami S, Aknin C, et al. Zac1 regulates an
imprinted gene network critically involved in the control of embryonic growth. Developmental cell.
2006;11(5):711-22.
69.   Gabory A, Ripoche M-A, Le Digarcher A, Watrin F, Ziyyat A, Forné T, et al. H19 acts as a trans
regulator of the imprinted gene network controlling growth in mice. Development. 2009;136(20):3413-21.
70.   Patten MM, Cowley M, Oakey RJ, Feil R. Regulatory links between imprinted genes: evolutionary
predictions and consequences. Proceedings of the Royal Society B: Biological Sciences.
2016;283(1824):20152760.
71.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.
Proceedings of the National Academy of Sciences. 2005;102(43):15545-50.
72.   Moffitt JR, Bambah-Mukku D, Eichhorn SW, Vaughn E, Shekhar K, Perez JD, et al. Molecular,
spatial, and functional single-cell profiling of the hypothalamic preoptic region. Science. 2018;362(6416).
73.   Team R. R: A language and environment for statistical computing. Vienna, Austria; 2013.
74.   Team R. RStudio: integrated development for R. RStudio, Inc, Boston, MA URL http://www rstudio
com. 2015;42:14.
787