Erythrocyte CD55 facilitates the internalization of *Plasmodium falciparum*

parasites

Bikash Shakya[1], Saurabh D. Patel[2], Yoshihiko Tani[3], and Elizabeth S. Egan[1*]

[1]Departments of Pediatrics and Microbiology & Immunology, Stanford University School

of Medicine, Stanford, CA 94305, USA

[2]Zuckerman Institute, Columbia University, New York City, NY 10027, USA

[3]Japanese Red Cross Osaka Blood Center, Osaka, Japan

*Correspondence: eegan@stanford.edu

Abstract

Invasion of human erythrocytes by the malaria parasite *Plasmodium falciparum* is a multi-

step process. Previously, a forward genetic screen for *P. falciparum* host factors

identified erythrocyte CD55 as essential for invasion, but its specific role and how it

interfaces with the other factors that mediate this complex process are unknown. Using

CRISPR-Cas9 editing, antibody-based inhibition, and live cell imaging, here we show

that CD55 is specifically required for parasite internalization. Pre-invasion kinetics,

erythrocyte deformability, and echinocytosis were not influenced by CD55, but entry was

inhibited when CD55 was blocked or absent. Visualization of parasites attached to CD55-

null erythrocytes point to a role for CD55 in progression of the moving junction. Our

findings demonstrate that CD55 acts after discharge of the parasite's rhoptry organelles,

and plays a unique role relative to all other invasion receptors. As the requirement for

CD55 is strain-transcendent, these results suggest that CD55 or its interacting partners

may hold potential as therapeutic targets for malaria.

Key Words

Malaria, CD55, erythrocyte, invasion, *Plasmodium falciparum*

Introduction

Malaria is caused by Apicomplexan parasites of the genus *Plasmodium*, of which

Plasmodium falciparum is responsible for the majority of severe disease cases in

humans. One of the world's major public health problems, malaria causes an estimated

216 million infections and ~445,000 deaths annually, primarily among young children and

pregnant women (WHO, 2018). *P. falciparum* has a complex life cycle involving stages in

the human and mosquito, but disease only occurs during the blood stage, when parasites

infect and replicate in human red blood cells (RBCs). As *P. falciparum* is an obligate

intracellular parasite, understanding the molecular determinants of its developmental

cycle within RBCs may lead to new therapies. For example, a number of *Plasmodium*

proteins that play key roles during erythrocyte invasion have shown promise as vaccine

candidates (Ord, Rodriguez, & Lobo, 2015; Sack, Kappe, & Sather, 2017; Salinas, Tang,

& Tolia, 2019). Since natural genetic variation in red cells can influence innate

susceptibility to malaria, host erythrocyte factors may also hold potential as therapeutic

targets (Taylor & Fairhurst, 2014). The identification and study of such factors, however,

has been severely limited by the intractability of mature RBCs, which lack a nucleus and

DNA.

P. falciparum invasion of erythrocytes involves a series of coordinated events that unfold

rapidly over the course of ~2 minutes. These events can be divided into three phases:

pre-invasion, active invasion, and echinocytosis (Gilson & Crabb, 2009; Weiss et al.,

2015). The process is initiated with the rupture of a "mother" parasite, termed a schizont,

which releases up to 32 daughter merozoites. During the pre-invasion phase, a free

merozoite makes initial contact with the red cell, stimulating shallow deformation of the

host cell plasma membrane. Next, the merozoite reorients so its apically-localized

organelles are abutting the cell surface. Reorientation is associated with significant

membrane deformation and involves interactions between *P. falciparum* ligands such as

the erythrocyte binding antigen (EBA) and reticulocyte binding-like homologues (Rh)

family proteins, and receptors on the red cell surface (Gilson & Crabb, 2009; Paul, Egan,

& Duraisingh, 2015; Riglar et al., 2011; Tham, Healer, & Cowman, 2012; Weiss et al.,

2015). Several ligand and receptor pairs have been shown to act at this stage, often in a

strain-specific manner (e.g. PfEBA-175 and GYPA; PfEBA-181 and GYPB PfEBA-140

and GYPC; and PfRh4 and CR1), but experimental data suggest their roles in apical

reorientation and host cell deformation are largely functionally redundant (Tham et al.,

2012).

The only receptor-ligand interaction known to be essential during the pre-invasion phase

involves basigin and PfRH5, which exists in a complex with PfRipr and CyRPA (Chen et

al., 2011; Crosnier et al., 2012; Dreyer et al., 2012; Reddy et al., 2015). Binding of the

PfRH5 complex to basigin is required for discharge of the rhoptry organelles into the

invaded cell and is associated with a calcium spike, potentially due to formation of a pore

at the erythrocyte surface (Volz et al., 2016; Weiss et al., 2015). Blocking the interaction

between PfRH5 and basigin with specific antibodies prevents invasion (Crosnier et al.,

2012).

Discharge of the rhoptry organelles heralds the start of active invasion. Among the

proteins injected from the rhoptries are those of the RON complex (RON 2, RON 4 and

RON5), which together form a receptor for binding by the PfAMA1 protein localized on

the merozoite surface (Alexander, Arastu-Kapur, Dubremetz, & Boothroyd, 2006;

Alexander, Mital, Ward, Bradley, & Boothroyd, 2005; Richard et al., 2010). The interaction

between PfAMA1 and the RON complex forms a moving junction between the parasite

and host cell that is believed to be entirely parasite-derived (Besteiro, Dubremetz, &

Lebrun, 2011; Besteiro, Michelin, Poncet, Dubremetz, & Lebrun, 2009; Harvey, Yap,

Gilson, Cowman, & Crabb, 2014; Koch & Baum, 2016). The moving junction provides an

anchoring point for the merozoite to actively invade using its own actinomyosin motor;

inhibiting the interaction between PfAMA1 and RON prevents invasion (Richard et al.,

2010; Srinivasan et al., 2011; Yap et al., 2014).

As invasion proceeds, a parasitophorous vacuole is formed from components of the host

cell membrane and rhoptries, yielding a protective niche for development of the new

daughter parasite. The third phase, echinocytosis, is a transient period of cell dehydration

and shrinkage observed after invasion; current evidence suggests it is triggered by

discharge of the rhoptry contents, prior to and independent of active invasion.

Echinocytosis is inhibited by reagents that prevent rhoptry discharge, such as antibodies

targeting basigin or PfRH5 (Weiss et al., 2015).

Most host factors known to play a role in *P. falciparum* invasion have been identified

based on their ability to bind to established invasion ligands or from studies of rare natural

mutants (Bei & Duraisingh, 2012). Given the inherent intractability of mature RBCs, which

lack a nucleus and DNA, the use of genetic approaches to identify and characterize

malaria host factors presents a logistical challenge (Egan, 2017). Recently, an shRNA-

based forward genetic screen using cultured red cells (cRBCs) derived ex-vivo from

primary human hematopoietic stem/progenitor cells (HSPCs) identified erythrocyte CD55

(aka DAF) as a critical host factor for *P. falciparum* invasion (Egan et al., 2015). A 70 kD

extracellular glycoprotein anchored to the red cell membrane by a

glycosylphosphatidylinositol (GPI) linkage, CD55 is broadly distributed in different tissues

and secretions, including blood cells (Cooling, 2015; Storry, Reid, & Yazer, 2010). On

erythrocytes, CD55 acts as a complement regulatory protein to prevent complement-

mediated damage. On epithelial cells, it has been shown to act as a receptor for Group B

coxsackie virus and Dr+ E. coli (Cooling, 2015; Coyne & Bergelson, 2006).

We have shown that *P. falciparum* invasion efficiency was reduced by ~50% in CD55-

knockdown cRBCs, and natural CD55-null erythrocytes from two rare donors with the

Inab phenotype were resistant to invasion (Egan et al., 2015). Importantly, the

requirement for CD55 was strain-transcendent, suggesting that it plays a conserved role

in *P. falciparum* invasion. However, the precise function of CD55 during invasion and how

it may interface with established ligands or receptors is unknown.

In this study, we investigated the functional role of CD55 during *P. falciparum* invasion

using CRISPR-Cas9 genome editing, antibody-based inhibition and live cell microscopy.

We found that CD55 plays a critical role during *P. falciparum* invasion of mature

erythrocytes, where it is specifically required for parasite internalization. As a host factor

that acts after discharge of the parasite's rhoptry contents, CD55 plays a unique role

relative to other receptors required for invasion, providing a crucial link between the

ligand-receptor interactions important for adhesion and deformation and effective

internalization.

Results

Generation of CD55-null red blood cells using CRISPR-Cas9

Previously, we observed that *P. falciparum* merozoites from several laboratory-adapted

strains and clinical isolates displayed impaired invasion into cryopreserved CD55-null

RBCs from two rare patients with the Inab phenotype (Egan et al., 2015). To study the

requirement for CD55 in an isogenic background, we sought to generate CD55-null

cRBCs from HSPCs using CRISPR-Cas9 genome editing by co-delivering single guide

RNAs (sgRNAs) and Cas9 in a ribonucleoprotein complex (RNP), a method that has been

used previously to generate null mutants in primary human CD34+ cells (Hendel et al.,

2015). We designed two chemically modified sgRNAs targeting the 5' end of *CD55*, and

delivered them individually or together into primary human CD34^{+} HSPCs via

nucleofection, in complex with recombinant Cas9 (Fig. 1 A). As a control, isogenic CD34^{+}

HSPCs from the same donor were nucleofected with Cas9 alone. After inducing the cells

to differentiate down the erythroid lineage for 18 days, we observed a high percentage of

CD55-null cRBCs in the RNP-transfected populations; a single sgRNA targeting exon 1

resulted in ~70% CD55-null cRBCs, while the combination of two sgRNAs targeting exons

1 and 2 increased the knockout efficiency to ~90% (Figure 1B). The cells proliferated ~

10,000-fold and the enucleation rate was >90% in both the CD55-null cells and wild-type

controls, demonstrating that the progenitors differentiated efficiently and that CD55 is not

required for this process (Figure S1). The absence of CD55 expression in the mutant

cRBCs was also confirmed using immunofluorescence assays (Figure S2).

Figure 1: CD55 is required for *P. falciparum* invasion (A) Schematic of *CD55* gene structure showing targeting sites of two single guide RNAs (sgRNAs). Vertical red lines indicate the positions of *CD55* exons. (B) Expression of CD55 on mutant (CD55-CRISPR) cRBCs generated with one sgRNA (left) or two sgRNAs (right), as compared to control (Cas9) cRBCs. (C) Invasion efficiency of *P. falciparum* 3D7 in Day 18 CD55-CRISPR cRBCs compared to isogenic controls (Cas9), relative to the invasion efficiency in peripheral blood erythrocytes (N=4 biological replicates; n=3 technical replicates; error bars indicate SEM; **p<0.01). (D) Time course of expression of CD71 on cRBCs harvested on different days of differentiation. Enucleated versus nucleated cells were gated using a nuclear dye. (E) Invasion efficiency of *P. falciparum* strain 3D7 in CD71-negative, CD55-CRISPR cRBCs compared to CD71-negative isogenic controls (Cas9). Invasion

efficiency is presented relative to the mean of Cas9 control. Three independent biological replicates are

shown; error bars represent SEM (n=2 or 3 technical replicates; **$p<0.01$, ***$p<0.005$). (F) Parasitemia in

population of CD55-CRISPR cRBCs, in which ~90% of the cells lack CD55 (CD55-negative), and the

remaining are CD55-positive, as quantified by immunofluorescence assays. Below are representative

images of CD55-CRISPR cRBCs infected with *P. falciparum*. Images are brightfield with fluorescence

overlaid. Blue, dapi. Red, anti-CD55-PE.

CD55 is required for *P. falciparum* invasion

To specifically assess the requirement for CD55 for *P. falciparum* invasion, we performed invasion assays using strain 3D7 in CD55-CRISPR cRBCs or in isogenic control cRBCs that had been differentiated for 18 days. While invasion into the CD55-CRISPR cRBCs was impaired relative to the control cells, it was only reduced by ~40% (Figure 1C). This subtle phenotype was reminiscent of the results for CD55-knockdown cRBCs, and differed from the strong invasion phenotype observed in erythrocytes from two CD55-null patients (Egan et al., 2015). Since cRBCs are less mature than erythrocytes from peripheral blood, we hypothesized that the relatively mild invasion phenotype observed in the CD55-null cRBCs may be explained by the overall maturation state of the ex-vivo cultures on day 18.

To begin to determine whether cell maturity modifies the requirement for CD55 during *P. falciparum* invasion, we assessed the surface expression of CD71 in the cRBCs, which is highly expressed on erythroblasts and young reticulocytes, but quickly disappears as reticulocytes mature into erythrocytes (J. Hu et al., 2013). Flow cytometry on day 17 demonstrated that approximately 50% of the enucleated cRBCs were CD71-positive and ~50% were CD71 negative, indicating a mix of cell maturity in the population (Figure 1D). This was further validated using a reticulocyte stain (Figure S3), which showed that day 17 cells were ~ 50% reticulocytes and 50% erythrocytes. A timecourse over 7 additional days of terminal differentiation revealed a progressive loss of CD71 expression, consistent with a change in the population structure to one dominated by erythrocytes rather than reticulocytes (Figure 1D).

To isolate a more homogeneous population of mature cRBCs for invasion assays, we

differentiated the cRBCs for 21-22 days and then used anti-CD71 antibody-immobilized

magnetic beads to enrich for CD71-negative cells (Figure S4). Using these highly mature

cells, invasion assays with *P. falciparum* strain 3D7 showed a ~75% reduction in invasion

into the CD55-CRISPR cRBCs as compared to isogenic control cRBCs in three biological

replicate experiments (Figure 1E). Since our CRISPR-Cas9 genome editing strategy

yielded a mixed population of ~90% CD55-null and 10% CD55-positive cells, we

suspected that the residual invasion observed in the "CD55-CRISPR" population was

attributable to wild-type cells. This was confirmed by immunofluorescence assays

showing that *P. falciparum* invasion was restricted to the cells expressing CD55: the

parasitemia in the minority, CD55-positive cells in the population was 22.5%, whereas

less than 0.5% of the CD55-null cRBCs were infected (Figure 1F). These results

demonstrate that CD55 is critical for *P. falciparum* invasion of fully differentiated cRBCs,

mimicking the prior observations with natural CD55-null erythrocytes, and validating

CD55 as an essential host factor for *P. falciparum* invasion. Based on previous

observations that the expression levels of many RBC surface proteins decline during

reticulocyte maturation (Chu et al., 2018; Malleret et al., 2013), we hypothesize that high

levels of other receptors important for invasion, such as basigin, may explain the reduced

requirement for CD55 for *P. falciparum* invasion of younger, CD71-positive reticulocytes.

Antibodies targeting CD55 inhibit *P. falciparum* invasion

To determine if *P. falciparum* invasion can be inhibited using antibodies targeting CD55,

we performed in vitro growth assays in the presence of three established anti-CD55

monoclonal antibodies individually and in combination. The antibodies each recognize a

different short consensus repeat (SCR) in the CD55 ectodomain: BRIC 230, BRIC 110

and BRIC 216 target SCR1, SCR2 and SCR3, respectively. While none of the antibodies

individually affected parasite growth, we observed a dose-dependent inhibition of parasite

growth over 72 hours in the presence of all three anti-CD55 monoclonal antibodies

combined, in comparison to an isotype control (Figure 2A). At the highest antibody

concentration tested (500µg/ml), the relative parasitemia was reduced by ~40% in the

presence of the combined monoclonals, compared to the isotype control.

Fig 2: Blocking CD55 with antibody inhibits growth of *P. falciparum* (A) *P. falciparum* strain 3D7 parasitemia after 72 hours of growth in RBCS in the presence of increasing concentrations of anti-CD55 monoclonal antibodies, relative to isotype control antibody (BRIC 170) at the same concentration. For the pooled antibodies, the indicated concentration was the total combined value, and there were equimolar amounts of each antibody. (N=3 biological replicates; n=2 technical replicates). Error bars indicate SEM.

(B) Parasitemia of *P. falciparum* strain 3D7 after 72 hours growth in non-enzyme treated RBCs with

increasing concentrations of polyclonal anti-CD55 IgG antibody, relative to that in isotype control antibody

at same concentration. (N=2 biological replicates; n=2 technical replicates). The highest antibody

concentration (10^0) was 400 ug/ml. (C) As in B, but with neuraminidase-treated RBCs. The highest antibody

concentration (10^1) was 40 ug/ml. (D) *P. falciparum* strain 3D7 parasitemia after 72 hours growth in 400

ug/ml Fab fragments generated from anti-CD55 polyclonal antibody or isogenic control. Error bars indicate

SEM. ***p<0.005. The starting parasitemias were 0.3% (Experiment 1) or 0.5% (Experiment 2).

To further study anti-CD55-mediated inhibition of *P. falciparum* invasion, we generated a

rabbit polyclonal antibody raised against the entire ectodomain of human erythrocyte

CD55. Flow cytometry analysis confirmed that the purified IgG antibody recognizes an

antigen on wild-type (WT) RBCs but not on CD55-null RBCs from an Inab donor

(Takahashi, 2008), confirming its specificity for CD55 (Figure S5). In *P. falciparum* growth

assays, we observed a dose-dependent inhibition of parasite growth in the presence of

anti-CD55 antibody relative to isotype control, with a ~40% reduction in relative

parasitemia at the highest concentration of antibody (400 µg/ml) (Figure 2B). This degree

of inhibition was very similar to that seen for the pooled monoclonal antibodies; together,

these findings suggest that blocking CD55 on the RBC can inhibit *P. falciparum* invasion.

We next tested the effect of anti-CD55 antibody on sialic acid-independent invasion

pathways by treating cells with neuraminidase (NM) to remove sialic acid. As has been

described previously, the growth of *P. falciparum* strain 3D7 was inhibited ~40% in NM-

treated RBCs as compared to untreated cells in the absence of antibody (Figure 2C),

reflecting some reliance of strain 3D7 on sialic acid for efficient invasion. In the presence

of increasing concentrations of anti-CD55 antibody, we observed a dose-dependent

inhibition of parasite growth in the NM-treated cells that is similar to untreated cells (~60%

versus ~40% at maximum concentration of antibody) (Figure 2B-C). The finding that

CD55 blockade inhibits *P. falciparum* regardless of the presence of membrane sialic acid

suggests that CD55 plays a role in both sialic acid-dependent and -independent invasion.

These results are consistent with previous findings showing that the requirement for CD55

is strain-transcendent, including strains that rely on sialic acid to various degrees (Egan

et al., 2015).

To confirm that the inhibitory effect of the polyclonal anti-CD55 antibody on *P. falciparum*

growth was not due to crosslinking, we tested the growth inhibitory activity of monovalent

anti-CD55 antibody fragments (Fab fragments). In the presence of 400 µg/ml of anti-CD55

Fab fragments, we observed ~40% reduction in relative parasitemia, recapitulating the

growth inhibitory activity observed with the bivalent anti-CD55 IgG antibody (Figure 2D).

These results further support the conclusion that CD55 is a critical host receptor for *P.*

falciparum.

Blocking CD55 inhibits *P. falciparum* internalization

To specifically determine the impact of CD55 blockade on *P. falciparum* invasion, we used

live cell imaging to visualize and quantify schizont rupture and merozoite invasion in real

time in the presence of the anti-CD55 polyclonal antibody or an isotype control (Figure

3A; Movies S1-S2). First, we quantified the efficiency of merozoite internalization in the

presence of anti-CD55 antibody versus isotype control. In the presence of the isotype

control antibody, of 345 *P. falciparum* merozoites that contacted an RBC, 53 invaded

successfully (15%) (Figure 3B). This frequency is similar to that described in previous live

microscopy studies analyzing the efficiency of 3D7 merozoite invasion in the absence of

antibodies (Volz et al., 2016; Weiss et al., 2015). In comparison, merozoite invasion was

significantly reduced in the presence of anti-CD55 antibody. Out of 312 merozoites that

made contact with an RBC in the presence of anti-CD55, only 21 successfully invaded

(6.7%) (Figure 3B). The number of invasion events per schizont rupture was similarly

reduced by half in the presence of anti-CD55 compared to control antibody (Figure 3C),

indicating that anti-CD55 antibody inhibits entry but does not prevent initial contact

between the merozoite and RBC.

Blocking CD55 has no impact on pre-invasion kinetics

Next, we examined the impact of CD55 blockade on the kinetics of the three main phases

of *P. falciparum* invasion: the pre-invasion time (period from initial contact to the onset of

internalization), the internalization time, and the time to echinocytosis (transient period of

cell dehydration that occurs after internalization) (Dvorak, Miller, Whitehouse, & Shiroishi,

1975; Weiss et al., 2015). For the subset of merozoites that ultimately invaded

successfully, there was no difference in the length of the pre-invasion time, internalization

time, or time to echinocytosis in the presence of anti-CD55 antibody compared isotype

control (Figure 3D-F). These results indicate that blocking CD55 does not impact pre-

invasion kinetics, at least not for the merozoites that manage to invade in the presence

of antibody.

Fig 3: Blocking CD55 with antibody inhibits invasion but not pre-invasion kinetics (A) Time-lapse images of invasion after initial merozoite contact. Arrowhead indicates invading merozoite. Time in seconds. (B) Percentage of merozoites that invaded an RBC after initial contact in presence of polyclonal anti-CD55 antibody (anti-CD55) or isotype control (IgG control). Bottom number is the total number of merozoites followed that made contact with the RBC, and top number is the subset that invaded. The data were acquired in three independent experiments, and the dots indicate the mean invasion efficiency from each experiment; *p=0.03. (C) Number of successful invasion events per schizont rupture (egress) in presence of anti-CD55 antibody or isotype control; *p=0.01. (D) Pre-invasion time (in seconds) for merozoites that

308 contact an RBC in the presence of anti-CD55 antibody or isotype control. (E) Internalization time (in

309 seconds) for merozoites in the presence of anti-CD55 antibody or isotype control antibody. (F) Time to

310 echinocytosis from the end of merozoite internalization in the presence of anti-CD55 antibody or isotype

311 control. Error bars indicate SEM; ns, not significant (B-F).

Merozoite-induced erythrocyte deformation is unaffected by CD55 blockade

During the pre-invasion period, parasite attachment to the RBC is associated with substantial deformation of the host cell membrane as the merozoite reorients apically (Dvorak et al., 1975; Gilson & Crabb, 2009; Paul et al., 2015). The degree of deformation is variable, and is mediated by interactions between parasite ligands released from the apical microneme organelles, such as the EBAs and Rhs, and receptors on the RBC membrane (Weiss et al., 2015). To investigate whether CD55 plays a role in RBC deformation, we used live microscopy to quantitate the efficiency and kinetics of merozoite-induced RBC deformation in the presence of anti-CD55 antibody versus isotype control. The efficiency of deformation was similar in the presence of the two antibodies: approximately 67% of merozoites deformed the RBC in the presence of anti-CD55 antibody, compared to ~60% deformation efficiency with the isotype control (Figure 4A). In addition, there was no significant difference in the duration of the deformation period in the presence of the two different antibodies (Figure 4B), suggesting that CD55 does not influence merozoite-induced deformation.

Fig 4: Merozoite-induced erythrocyte deformation is not affected by blocking CD55 (A) Efficiency of
RBC deformation by merozoites that make contact in presence of anti-CD55 or IgG control. The fractions
indicate the number of deformed RBCs out of the total merozoite-RBC contacts observed. The data were
acquired in three independent experiments, and the dots represent the mean for each experiment.; *p=0.02.
(B) Duration of erythrocyte deformation induced by attached merozoites that ultimately invaded in presence
of polyclonal anti-CD55 antibody (anti-CD55) or isotype control (IgG control). Time in seconds. (C-D)
Strength of merozoite-induced deformation in presence of anti-CD55 or isotype control for cases where
invasion was successful (C) or not successful (D); N/A, not applicable; ns, not significant, *p=0.0029. Error
bars indicate SEM (A-D).

We employed a four-point deformation scale to quantify the intensity of merozoite-induced

RBC deformation in the presence of antibody, where 0 denotes the absence of

deformation and 3 denotes the most extreme degree of deformation (Weiss et al., 2015)

(Movies S3-S5). Almost all merozoites that invaded successfully induced strong

deformation (scores of 2 or 3), regardless of which antibody was present (Figure 4C). In

comparison, merozoites that failed to invade had lower deformation scores, the majority

having scores of 0 or 1. For these merozoites, the distribution of scores skewed higher

in the presence of anti-CD55 antibody compared to the isotype control, perhaps reflecting

the inability of otherwise "fit" merozoites to complete invasion (Figure 4D). Taken

together, these findings demonstrate that blocking CD55 with antibody does not influence

the efficiency, duration, or strength of merozoite-induced RBC deformation, and instead

suggest that CD55 exerts its effect on invasion at a downstream step.

To further validate the conclusions of the antibody inhibition experiments, we took a

complementary approach and performed live cell imaging of *P. falciparum* invasion with

natural CD55-null erythrocytes from a rare donor with the Inab phenotype, where CD55

is absent due to a stop codon in exon 2(Takahashi, 2008). In traditional invasion assays,

we observed ~80% reduction in the efficiency of *P. falciparum* invasion into CD55-null

RBCs as compared to control RBCs after ~ 18 hours, as has been shown previously

(Egan et al., 2015) (Figure S6). When imaging invasion in real time, we did not observe

any successful invasion events into the CD55-null RBCs, out of 310 merozoites that made

contact (Figure 5A and Movies S6-S7). In comparison, there were 25 successful invasion

events into the control RBCs, out of 197 merozoites that made contact (~11%). Although

invasion into the CD55-null cells was clearly impaired, there was no difference in the

efficiency of merozoite-induced RBC deformation between the two genetic backgrounds

(Figure 5B). Moreover, we observed no significant difference in the distribution of

deformation scores between CD55 null and control RBCs for merozoites that did not

invade (Figure 5C). These results are consistent with the findings from the antibody

inhibition experiments, and suggest that CD55 influences invasion by acting after the

period of erythrocyte deformation.

Fig 5: Absence of CD55 prevents invasion but does not impact deformation. (A) Percentage of
merozoites that invaded wild-type (WT) or CD55-null RBCs after initial contact. Bottom number is the total

number of merozoites followed that made contact with the RBC, and top number is the subset that invaded.
The data were acquired in three independent experiments, and the dots indicate the mean invasion
efficiency from each experiment; *p=0.04. (B) Efficiency of WT or CD55-null RBC deformation upon
merozoite contact. Bottom number is the total number of RBCs contacted by a merozoite, and top number
is the subset that were deformed upon contact. The data were acquired in three independent experiments,
and the dots indicate the mean deformation efficiency for each experiment; ns, not significant. (C) Strength
of merozoite-induced deformation of WT or CD55-null RBCs among non-invading merozoites; ns, not
significant. Error bars indicate SEM (A-C).

CD55 acts downstream of rhoptry discharge

Since we observed that the blocking or deletion of CD55 impaired *P. falciparum* entry without altering pre-invasion kinetics or membrane deformation, we hypothesized that CD55 functions after the ligand-receptor interactions that mediate attachment and deformation. During the sequential steps of *P. falciparum* invasion, RBC deformation is followed by injection of the rhoptry organelle contents into the host cell cytoplasm, an event that requires an interaction between the parasite ligand PfRh5 and its RBC receptor, basigin (Weiss et al., 2015). The contents of the rhoptries in turn are believed to trigger echinocytosis. When the interaction between PfRH5 and basigin is blocked using inhibitory antibodies and rhoptry discharge is prevented, echinocytosis fails to occur (Volz et al., 2016; Weiss et al., 2015). To investigate whether CD55 is similarly required for release of the rhoptry contents, we quantified the incidence of RBC echinocytosis elicited by attached merozoites in the presence of anti-CD55 antibody versus isotype control using a cytochalasin-D (cyt-D) live microscopy assay (Figure 6A-B and Movie S8). Cyt-D is an inhibitor of actin polymerization that prevents merozoite internalization by inhibiting the actino-myosin motor, but does not impact attachment, rhoptry discharge, or echinocytosis (Miller, Aikawa, Johnson, & Shiroishi, 1979; Weiss et al., 2015). Upon treatment with cyt-D, ~60% of merozoites that attached to an RBC triggered echinocytosis in the presence of the isotype control antibody, and a similar rate of echinocytosis was observed in the presence of anti-CD55 antibody (Figure 6C). These results demonstrate that blocking CD55 does not significantly impact rhoptry discharge or echinocytosis, standing in distinct contrast to what has been observed for basigin and Rh5, where blocking their interaction with antibodies prevents echinocytosis (Weiss et al., 2015).

405   Together, these findings demonstrate that CD55 acts downstream of the interaction

406   between RH5 and basigin, facilitating a step of invasion that occurs after release of the

407   rhoptry organelles.

Fig 6: Blocking CD55 with antibody does not impact echinocytosis. (A) Cartoon illustrating echinocytosis elicited by an attached merozoite that has discharged its rhoptry contents in the presence of cyt-D, which prevents internalization. Reagents that block rhoptry discharge prevent echinocytosis. (B) Image showing an echinocyte with an attached merozoite. Scale bar=5um. (C) Echinocytosis efficiency in the presence of anti-CD55 antibody or isotype control. Bottom number is the total number of merozoite-RBC pairs observed to make contact, and top number is the subset in which echinocytosis occurred. The data are from eight independent experiments. Dots indicate the mean from each experiment and error bars represent the standard error of mean; ns, not significant.

CD55 may be required for progression of the moving junction

The moving junction that forms between the cell membranes of an invading merozoite and the erythrocyte during internalization involves interactions between PfAMA1, which derives from the micronemes, and the PfRON complex, which is expressed by the rhoptries. To investigate a possible role for CD55 in formation of the moving junction, we sought to visualize PfAMA1 and PfRON4 as merozoites attempted to invade CD55-null (Inab) or wild-type erythrocytes in the presence of cyt-D. Under these conditions, we observed a ~50% reduction in the efficiency of merozoite attachment to CD55-null erythrocytes as compared to WT controls by flow cytometry, as has been observed previously (Figure 7A) (Egan et al., 2015). Using confocal microscopy, we found that PfAMA1 and PfRON4 were frequently co-localized at the cellular interface for merozoites attached to WT control cells (~85%) (Figure 7B). In comparison, for merozoites attached to CD55-null erythrocytes, PfAMA1 and PfRON4 were co-localized at the cellular interface in only ~40% of cases (Figure 7B). For those merozoites in which PfAMA1 and PfRON4 were colocalized at the interface, we observed that approximately half of the WT control cells had an indentation at the point of merozoite contact, suggesting the merozoites were mid-invasion, whereas this was never observed for the CD55-null cells (Figure 7C). Together, these results support a model where CD55 is required for progression of the moving junction, and acts downstream of and distinctly from all other known host receptors for *P. falciparum* invasion.

Fig 7: Invasion fails to progress in the absence of CD55.

(A) Attachment efficiency of *P. falciparum* merozoites to CD55-null versus control RBCs in the presence of cyt-D, as measured by flow cytometry 90 min after the addition of synchronized schizonts (N=3, n=3). Error bars indicate SEM; *p<0.018. (B) Orientation of merozoites attached to WT versus CD55-null RBCs, indicated by the localization of AMA1, RON4, and a density at the merozoite's apical end as measured by confocal microscopy. Apical, AMA1 and RON4 co-localized at the cellular interface and merozoite density abutting RBC; Not apical, AMA1 and RON4 not co-localized at the cellular interface and merozoite density

not abutting RBC; N.D., indeterminate. The numbers above each bar indicate the subset of attached

merozoites in that orientation. (C) Representative confocal images showing merozoites attached to RBCs

with apical or non-apical orientation. Merozoite density at the apical end of the merozoite is indicated by an

arrow.   (D) Progression of invasion for merozoites apically attached to CD55-null or WT RBCs in the

presence of cyt-D. Panel on the right shows representative confocal images of a merozoite at the mid-

invasion stage.

Discussion

A comprehensive understanding of the molecular interactions required for *P. falciparum*

invasion has been limited by the absence of a robust genetic system to study red cells,

which are terminally differentiated and lack a nucleus and DNA. Previously, an RNAi-

based forward genetic screen in cultured red cells derived from HSPCs identified two new

candidate host factors for invasion: CD44 and CD55 (Egan et al., 2015). In this study, we

used a combination of genetics and inhibitory antibodies to determine the precise steps

of *P. falciparum* invasion during which CD55 functions. We have demonstrated that CD55

is specifically required for merozoite internalization, and plays a unique role relative to the

other host receptors known to act during invasion.

Prior work has shown that the efficiency of *P. falciparum* infection is reduced by ~

50% in CD55-deficient cRBCs where expression is downregulated using shRNAs. While

experiments using erythrocytes from two rare, CD55-null donors suggested that CD55 is

essential for *P. falciparum* invasion, this has never been demonstrated genetically. A

major roadblock to such experiments has been the technical challenges associated with

CRISPR-Cas9 genome editing in primary human hematopoietic stem cells, including

cytotoxicity from nucleic acids and low rates of transfection (Hendel et al., 2015). In this

study, we showed for the first time that fully mature, CD55-null cRBCs can be generated

efficiently from primary human HSPCs using CRISPR-Cas9 genome editing. Using

isogenic wild-type and mutant cells, we have demonstrated that CD55-null cRBCs are

resistant to *P. falciparum* invasion, confirming that CD55 is an essential host factor for *P.*

falciparum.

Here, we demonstrated the feasibility and benefits of generating truly mature erythrocytes

ex-vivo for the study of malaria host factors, as these cells closely mimic the target cell

for *P. falciparum* in the human bloodstream. Our approach involving the co-delivery of

two chemically modified sgRNAs together with Cas9 as a ribonucleoprotein complex has

been shown to be an effective strategy for CRISPR-based gene knockout in primary

HSPCs and T cells (Hendel et al., 2015). This method obviates the toxicity associated

with plasmid delivery, minimizes off-target activity, and improves sgRNA stability.

Combined with our ex-vivo erythropoiesis culture system, this method can efficiently

generate terminally differentiated, enucleated, CD71-low red cells with a high rate of gene

knock out that can be used to study host genetic determinants for *P. falciparum* in an

isogenic background.

We observed that the reliance of *P. falciparum* on CD55 for invasion increased

significantly as enucleated cRBCs matured into erythrocytes, likely reflecting the

substantial changes in protein abundance that occur during terminal maturation of human

red cells, including for proteins known to act as receptors, such as basigin and CR1

(Gautier et al., 2016; G. Hu et al.). The efficiency of *P. falciparum* invasion is further

influenced by the deformability of the red cell membrane (Tiffert et al., 2005), a biophysical

property that also changes as red cell progenitors proceed through enucleation and final

maturation (Giarratana et al., 2005). As clinical malaria is primarily a disease of mature,

peripheral blood erythrocytes, our results showing that terminal red cell maturation

modifies the requirement for CD55 during invasion highlights the potential drawbacks of

using cell lines to study host factors for *P. falciparum*. While considered erythroid in

nature, immortalized cell lines such as JK-1, EJ and BEL-A have low rates of enucleation

and terminal maturation (Kanjee et al., 2017; Satchwell et al., 2019; Scully et al., 2019),

suggesting their utility for genetic experiments on *P. falciparum* invasion may be limited

by incomplete phenotypes.

Antibodies targeting a variety of specific host receptors for *P. falciparum* have been

shown to have invasion inhibitory activity, including those against GYPA, CR1, and

basigin (Crosnier et al., 2012; Pasvol et al., 1989; Spadafora et al., 2010). While we found

that individual monoclonal antibodies against three distinct SCR domains of CD55 had

no discernable effect on *P. falciparum* growth, in combination they had potent dose-

dependent inhibitory activity, suggesting CD55's role in invasion is not restricted to a

specific SCR domain. Our demonstration that both the polyclonal anti-CD55 antibody and

anti-CD55 Fab fragments can inhibit *P. falciparum* growth aligns with the results from our

genetic studies, and corroborates the conclusion that CD55 is a critical host factor for *P.*

falciparum. In contrast to CR1, where antibodies are only inhibitory in the absence of red

cell sialic acid (Spadafora et al., 2010), anti-CD55 antibody blocked both sialic acid-

dependent and -independent invasion. This finding is consistent with prior studies

showing that the requirement for CD55 in *P. falciparum* invasion is strain-transcendent

(Egan et al., 2015), and implies a model where CD55 acts distinctly from CR1 and the

other alternative, strain-specific receptors.

Given the inherent limitations associated with studying invasion on a population scale in

longer term assays, live microscopy has been increasingly employed to visualize

individual *P. falciparum* invasion events in real time. Analysis of the morphology and

kinetics of discrete invasion steps in the context of blocking antibodies or soluble proteins

has contributed to a model describing the molecular events that occur during invasion

(Volz et al., 2016; Weiss et al., 2015). Our live microscopy experiments have added a

new dimension to this model by revealing that merozoite internalization is inhibited in the

presence of anti-CD55 antibody or with CD55-null erythrocytes, demonstrating for the first

time that CD55 is specifically required for parasite entry. Where and how does CD55 act

in relation to the *P. falciparum* ligands and erythrocyte receptors known to function during

invasion? We observed no impact of the anti-CD55 antibody on pre-invasion kinetics or

merozoite-induced erythrocyte deformability, findings that were validated using CD55-null

erythrocytes from a rare donor. These results suggest that CD55 functions distinctly from

the "alternative" receptors required for apical reorientation and red cell deformation (e.g.

glycophorins and CR1), as blocking interactions between these receptors and their

ligands strongly inhibits merozoite-induced deformability (Weiss et al., 2015).

Echinocytosis is a phenomenon of transient red cell shrinkage that commences soon after

merozoite internalization. Current evidence suggests that it is stimulated by changes in

the red cell cytoplasm that occur once an irreversibly-attached merozoite discharges its

rhoptry contents into the erythrocyte (Volz et al., 2016; Weiss et al., 2015). Echinocytosis

is inhibited by antibodies that block the interaction between PfRH5 and basigin, implying

that this interaction is necessary for rhoptry discharge (Weiss et al., 2015). These blocking

antibodies also inhibit Ca^{2+} flux into the erythrocyte, possibly reflecting formation of a pore

at the cellular interface when PfRH5 and basigin interact (Volz et al., 2016). The

glycophorins and CR1 act early in the pre-invasion phase during apical reorientation, and

blocking their interactions with ligands using genetics, enzyme treatments, or antibody

blockade also inhibits echinocytosis (Volz et al., 2016; Weiss et al., 2015). In contrast,

echinocytosis is not prevented by treatment with the actin polymerization inhibitor cyt-D

or reagents that block the interaction of PfAMA1 and the RON complex. Together, these

findings demonstrate that the stimulus for echinocytosis occurs after the interaction of

PfRH5 with basigin, but before establishment of the moving junction.

Our observation that anti-CD55 antibody had no effect the efficiency of merozoite-induced

echinocytosis demonstrates that CD55 acts distinctly from basigin and the other

established host receptors known to act during *P. falciparum* invasion. While both anti-

CD55 and anti-basigin inhibit invasion, echinocytosis occurs normally in the presence of

anti-CD55 but is blocked by anti-basigin antibody, likely due to inhibition of rhoptry

discharge (Volz et al., 2016). As blocking CD55 does not inhibit deformation nor

echinocytosis during invasion, our findings support a model in which CD55 acts after

rhoptry discharge, and specifically impacts merozoite internalization.

Merozoite internalization requires the formation of a moving junction between the cell

membranes of the invading merozoite and the erythrocyte that moves posteriorly down

the parasite as it invades. The current model of the moving junction involves interactions

between PfAMA1 expressed on the merozoite surface and the PfRON complex inserted

into the red cell membrane, independent of any host-encoded receptors (Alexander et al.,

2006; Alexander et al., 2005; Besteiro et al., 2009; Richard et al., 2010; Riglar et al., 2011;

Srinivasan et al., 2011). In addition to its key role in internalization, the moving junction

has also been proposed to be important for formation of the parasitophorous vacuole (PV)

(Yap et al., 2014).

Could CD55 function as a component of the moving junction during invasion? We

observed that PfAMA1 and PfRON4 were almost always co-localized at the cellular

interface for merozoites attempting to invade WT cells, whereas this was significantly less

common for merozoites attempting to invade CD55-null erythrocytes. For the minority of

merozoites that attached to CD55-null erythrocytes and had PfAMA1 co-localized with

RON at the cellular interface, none were observed to have progressed past early invasion,

unlike those attached to WT cells. Together, these findings support a model in which

CD55 is required for progression of the moving junction, either directly or indirectly. Future

studies involving higher resolution microscopy such as cryo-EM will be necessary to

better characterize how absence of CD55 may impact the moving junction or architecture

of the PV after invasion.

Our findings showing that the function of CD55 during invasion can be narrowed to the

internalization step are consistent with a model where CD55 acts distinctly from the other

known receptors for invasion, potentially by interacting with a specific parasite ligand. As

there is a precedent for CD55 on epithelial cells to act as a pathogen receptor that

transmits signals to the host cell (Coyne & Bergelson, 2006), it is tantalizing to

hypothesize that this molecule functions similarly in *P. falciparum* invasion of

erythrocytes. Ultimately, biochemical identification of parasite and erythrocyte interaction

partners of CD55 will yield important additional insights into the molecular function of

CD55 during merozoite internalization. Given the complexity of *P. falciparum* invasion and

the unique and essential role of CD55 relative to other established receptors, targeting its

activity or interaction partners in novel intervention strategies may enhance the

effectiveness of future therapies or vaccines for malaria.

Methods

***P. falciparum* culture, invasion assays and growth inhibition assays**

P. falciparum strain 3D7, a laboratory-adapted strain obtained from the Walter and Eliza

Hall Institute (Melbourne, Australia) was routinely grown in human erythrocytes (Stanford

Blood Center) at 2% hematocrit in complete RPMI-1640 supplemented with 25 mM

HEPES, 50 mg/L hypoxanthine, 2.42 mM sodium bicarbonate and 0.5% Albumax

(Invitrogen) at 37°C in 5% CO_2 and 1% O_2.

Parasite invasion assays were performed using synchronized late-stage schizont

parasites isolated using a MACS magnet (Miltenyi) and added at 1.0-1.5% initial

parasitemia to the cultured red blood cells (cRBCs) or peripheral red blood cells (pRBCs)

at 0.3% hematocrit. For some invasion assays, cRBCs were enriched for the CD71

negative population by using anti-CD71 antibody immobilized magnetic beads (Miltenyi),

as described below. Assays were performed in a volume of 100 μl per well in a 96 well

plate, or at 50 μl per well in Half Area 96-well plates (Corning) for the assays with CD71-

negative cRBCs. The ring stage parasitemia was determined after 18-24 hours by bright-

field microscopy of cytospin preparations stained with May-Grünwald and Giemsa. A

minimum of 1000 cells were counted for each technical replicate. For assays in which

there was no selection for CD71-negative cells, the invasion efficiency was determined

by normalizing the average ring stage parasitema in each genetic background to the

average ring stage parasitemia in control peripheral blood RBCs (pRBCs) for each

biological replicate. For assays that used CD71-negative cRBCs, the ring stage

parasitema for each genetic background was normalized to the mean for the control

cRBCs. Assays were performed at least 3 times using 2-3 technical replicates.

Antibody inhibition assays were performed using pRBCs in the presence of mouse

monoclonal antibodies obtained from IBGRL (BRIC 216, BRIC 230, BRIC 110, BRIC

170), polyclonal anti-CD55 antibody produced by New England Peptide, or isotype control

rabbit IgG antibody (Novus). All antibodies were dialyzed overnight in RPMI buffer prior

to use. Schizont stage parasites were added to untreated or neuraminidase treated

pRBCs at 0.5% hematocrit at an initial parasitemia of 0.5% in the presence of 0.05 to 500

µg/ml of the antibodies. For the neuraminidase treatment, the cells were incubated with

66.7mU/ml of neuraminidase from *Vibrio cholerae* (Sigma) at 37°C for 1 h with shaking

and washed 3 times in buffer before use in the growth inhibition assays. Some assays

were performed in presence of Fab fragments of anti-CD55 or isotype control IgG

antibodies at the concentration of 400 µg/ml. The Fab fragments were prepared as

described below. Assays were performed 2-3 times with 2-3 technical replicates in a

volume of 100 µl per well in 96 well plates. Parasitemias were determined on day 3 by

staining with SYBR Green 1 nucleic acid stain (Invitrogen) at 1:2000 dilution in PBS/0.3%

BSA for 20 minutes, followed by flow cytometry analysis on a MACSQuant flow cytometer

(Miltenyi).

Generation of Fab fragments

Polyclonal anti-CD55 antibody and control IgG were digested using the Pierce[TM] Fab

preparation kit (Thermo Fisher Scientific). The resulting Fab fragments were quantified

on a spectrophotometer, concentrated, and buffer exchanged with incomplete RPMI

using 30K Amicon Ultra 0.5 ml centrifugal filter (Millipore). For final quantification, the Fab

fragments were stained with Coomassie in a 10% SDS-PAGE gel along with known

concentrations of the undigested antibodies, and concentrations of the Fab fragments

were determined using ImageJ 1.50i(Schneider, Rasband, & Eliceiri, 2012).

Generation of cultured red blood cells from primary human CD34+ HSPCs

Bone marrow-derived primary human CD34+ HSPCs (Stem Cell Technologies) were

cultured in erythroid differentiation medium (cPIMDM) composed of Iscove Basal Medium

(IMDM) (Biochrom) supplemented with 4mM L-Glutamine (Sigma), 330 µg/ml holo-

transferrin (BBI Solutions), 10 µg/ml of recombinant human insulin (Sigma), 2 IU/ml

heparin (Affymetrix), 10^{-6} M hydrocortisone (Sigma), 100ng/ml SCF (R & D Systems), 5

ng/ml IL-3 (R & D Systems), 3 IU/ml Epo (Amgen) and 5% plasma (Octapharma) at 37°C

in 5% CO2 in air, as previously described (Egan et al., 2015; Giarratana et al., 2011). On

the 2^{nd} day of the culture, the cells were subjected to nucleofection with ribonucleoprotein

(RNP) complexes as described below. On day 5, the cells were washed and resuspended

in fresh cPIMDM to maintain concentration ~ 1×10^4/ml. Between day 8-13, the cells were

maintained in fresh cPIMDM devoid of IL3 and hydrocortisone at concentration of <5×10^5

cells/ml. On day 13, the cells were washed and plated at 7.5×10^5 - 1.0×10^6cells/ml in

cPIMDM without IL-3, hydrocortisone, or SCF. On day 15-16, the cells were co-cultured

at 1.0×10^6cells/ml concentration on a murine stromal cell layer (MS-5), as previously

described (Giarratana et al., 2005). The cells were harvested on days 18-22 for different

experiments. Growth and differentiation were monitored using hemocytometer-based

quantification and light microscopy of cytospin preparations stained with May-Grünwald

and Giemsa. To quantify the enucleation rate, cRBCs were incubated in Vybrant

DyeCycle violet (Life Technologies) (1:10,000) at 37°C for 30 min, followed by flow

cytometry analysis on a MACSQuant flow cytometer (Miltenyi).

CRISPR-Cas9 genetic modification of primary human CD34+ cells

Two sgRNAs targeting human *CD55* exons were designed using the Broad Institute's

GPP sgRNA design portal and synthesized as chemically modified sgRNAs by Synthego.

The sgRNA sequences were: GGGCCCCUACUCACCCCACA, which is predicted to

recognize a sequence in exon 1 of *CD55*, and CUGGGCAUUAGGUACAUCUG, which is

predicted to recognize a sequence in exon 2. Ribonucleoprotein (RNP) complexes

containing one or both sgRNAs were prepared by slowly adding 300 pmol of each sgRNA

to 150 pmol Cas9 protein in a 10 µl final volume with nuclease-free water and incubating

at room temperature for 10 min. On day 2 after thawing CD34+ cells, the RNP complexes

were added to 1×10^5 cells in 40 µl of P3 nucleofection buffer from the 4D-Nucleofector X

kit (Lonza). Half of the mixture was loaded to each well of a 16-well nucleofection cassette

and nucleofected using the using E0-100 program with the 4D-Nucleofector Lonza

Amaxa. After nucleofection, cells were transferred to 6 ml fresh cPIMDM and incubated

at 37°C in 5% CO2 in air.

Enrichment of CD71 negative cRBCs

Fully differentiated cRBCs were washed in degassed and chilled bead buffer (PBS + 0.5%

bovine serum albumin + 2mM EDTA), resuspended in the same buffer (45 µl for $6x10^6$

cells), and incubated the anti-CD71 antibody-immoblized beads (Miltenyi) (10µl beads for

$6x10^6$ cells) at 4°C for 15 min. The cells were washed in 1 ml of bead buffer and passed

through an LS magnetic column (Miltenyi) and washed 3 times with 2 ml of the buffer.

The flow through containing CD71- negative cells was collected, washed with cRPMI

medium and used for invasion assays.

Detection of cell surface proteins by flow cytometry

Expression of RBC surface proteins was measured in control or knockout cRBCs by flow

cytometry. $1x10^6$ cRBCs were washed 2 times with PBS/0.3% BSA and incubated with

primary monoclonal antibodies or fluorochrome conjugated antibodies at 4°C for 1 h.

Antibodies used: anti-CD55 (BRIC 216-PE, IBGRL; 1:50) and anti-CD71-PE (Miltenyi;

1:20). After incubation, the cells were washed 2 times in PBS/0.3% BSA, followed by flow

cytometry analysis on a MACSQuant flow cytometer (Miltenyi).

Immunofluorecence assays (IFA)

IFAs were performed as previously described, with some modifications (Tonkin et al.,

2004). For IFAs of CD55-CRISPR cRBCs infected with *P. falciparum*, cells were fixed in

4.0% paraformaldehyde and 0.0015% glutaraldehyde in PBS for 20 min at room

temperature and blocked for one hour in 3% BSA/PBS. The cells were incubated with

anti-CD55 antibody (BRIC 216-PE from IBGRL) at 1:50 concentration for 1 h at 4°C. Cells

were mounted Fluoromount-G with DAPI mounting medium (Electron Microscopy

Services) and the fluorescent images were captured with a 60X objective on a Keyence

BZ-X700 fluorescence microscope. For IFAs of merozoites attached to pRBCs, samples

were prepared as in the attachment assays (see below), and 60 ul of the samples were

fixed in the fixative containing 4.0% Paraformaldehyde and 0.015% Glutaraldehyde for

20 min, washed twice in PBS, and allowed to settle onto a Poly-L-Lysine coated coverslip

(Corning). The samples were then incubated in 0.1% Triton X-100/PBS for 10 min at

room temperature, washed in PBS, and incubated in 0.1mg/ml of NaBH4/PBS for 10 min.

at room temperature. Following a wash in PBS, the cells were blocked overnight in fresh

PBS/3.0% BSA at room temperature. Primary antibodies: mouse monoclonal Anti-

PfAMA1 1F9 (1:200) (Coley et al., 2001) and rabbit polyclonal anti-PfRON4 (1:200)

(Richard et al., 2010) were diluted in blocking buffer applied to cells for 2 hr at room

temperature. Following 3 washes, the cells were incubated in corresponding secondary

antibodies at 1:500 dilution: Alexa Fluor 555 (anti-rabbit) and Alexa Fluor 488 (anti-

mouse) for 1 h at room temperature. The cells were washed in PBS supplemented with

0.1ng/µl of DAPI (Thermofisher) and mounted in Vectashield anti-fade mounting medium.

Imaging was performed using a Zeiss LSM710 confocal microscope and the image

analysis was performed using Fiji/imageJ (Version 2.0.0-rc-69/1.52i). Apical orientation

was defined as PfAMA1 and PfRON4 co-localization at the cellular interface, and non-

apical orientation was defined as PfAMA1 and PfRON4 not co-localized at the cellular

interface. Merozoites were defined as being mid-invasion if AMA1 and RON4 were co-

localized at cellular interface and the merozoite appeared to protrude into the RBC,

creating an indentation.

Anti-CD55 polyclonal antibody generation

A CD55 cDNA (Asp 35-Ser 353) was cloned into a modified pTT5 vector (Raymond et

al., 2011), whose expression cassette consists of eGFP fused to a puromycin resistance

gene followed by a 2A skip peptide (Funston, Kallioinen, de Felipe, Ryan, & Iggo, 2008)

and a BiP signal peptide, and the resulting plasmid was verified by sequencing.   1.8L of

293-6E cells   grown in Freestyle 293 media (Invitrogen) to 0.8x10e6 cells/ml were

transfected with linear PEI (Polysciences)   with 397ug DNA at a 1:3 DNA:PEI

ratio.   TN1 tryptone (Organotechnie) was added to a final concentration of 0.5% (w/v)

one day after transfection, and cells were grown for an additional 5 days. Cells were spun

down and discarded, and the following reagents were added to the harvested media (final

concentration listed): 20mM Tris pH8, 350mM NaCl, 5mM Imidazole pH8, 0.2mM

NiCl2.   6ml GE IMAC sepharose 6 beads, and the sample was rocked at 4C for 1

hour. The sample was then poured into a glass column, and washed with 50ml of TBS

(10mM Tris pH8, 500mM NaCl) with imidazole at the following concentrations

sequentially: 5mM, 10mM, 50mM, and 250mM. Nickel was stripped from the column with

10ml 100mM EDTA.   The majority of the CD55-8xHis protein eluted in the 250mM

Imidazole fraction, with a smaller amount of equal purity in the 50mM imadazole

fraction. These were dialyzed separately into PBS pH 7.4 overnight at 4C, concentrated

to 10mg/ml using Amicon Ultra 15 with 30kDa cutoff (Millipore), snap frozen in liquid

nitrogen and stored at -80C until further use. 2.5 mg of CD55-8xHis was used for three

immunizations of two New Zealand white-SPF rabbits by New England Peptide.

Approximately 50ml of recovered antiserum was used for negative affinity purification with

an 8X HIS column, and subsequently purified with a protein A column. Approximately 15

ml of purified IgG at 1.209 mg/ml concentration was purified and stored at -80C

until further use.

Live cell imaging

Schizont stage *P. falciparum* strain 3D7 parasites at 4-5% parasitemia and 2% hematocrit

were tightly synchronized with 2 µM Compound 2 to prevent schizont rupture (Collins et

al., 2013). After ~4-6 hour incubation, they were washed three times and allowed to

recover in fresh cRPMI for 45 to 90 minutes at 37º C. After recovery, anti-CD55 antibody

or IgG isotype control was added to the cells at final antibody concentration of 400 µg/ml

and 1% hematocrit. In experiments involving previously cryopreserved CD55-null pRBCs

(Takahashi, 2008) or control WT pRBCs, late stage *P. falciparum* schizonts were isolated

using a MACS magnet, synchronized with Compound 2, and then mixed with the pRBCs

at 4-5% parasitemia and 1% hematocrit. The assays were loaded into a 20 mm diameter

Hybriwell sealing system (Grace Bio-labs) customized for live cell imaging. Rupture of the

parasite-infected red blood cells and subsequent invasion events were video recorded

using a 60X objective Keyence BZ-X700 live microscopy setup at 37°C supplied with 5%

CO_2 and 1% O_2 gaseous environment. For the live cell imaging of echinocytosis, 1µM

Cytochalasin D was added to the above mixtures. Kinetics and morphology of distinct

steps of invasion in the video were measured by using ImageJ (ImageJ 1.50i) as

described previously(Schneider et al., 2012; Weiss et al., 2015). Briefly, the following

events were quantified in each video: a) Contacts between merozoites and RBCs that

culminated in successful invasion; b) successful invasion that resulted echinocytosis; c)

contacts between merozoites and RBCs that could not proceed beyond initial

contact/deformation. d) RBCs contacted before successful invasion e) period of pre-

invasion (initial contact to end of deformation), f) during of internalization, g) time to the

onset of echinocytosis, h) efficiency and degree of merozoite-induced deformation of the

target RBC using a four-point deformation scale (0,1,2,3).

Merozoite attachment assays

Purified schizonts were added at 12.0% parasitemia to previously cryopreserved CD55-

null pRBCs (Takahashi, 2008) or control WT pRBCs at 1.0% hematocrit in the presence

of 1 µM Cytochalasin D, 50 U/ml Heparin or none in a final volume of 100 ul per well in a

96 well plate. The mixtures were incubated at 37°C for 90 min and the schizonts were

allowed to rupture. To quantify merozoite attachment by flow cytometry, 10 µl aliquots of

the samples were fixed in 2% Glutaraldehyde and 0.116M sucrose in PBS, washed in

PBS/0.3% BSA, and stained with SYBR Green 1 nucleic acid stain (Invitrogen). at 1:2000

dilution in PBS/0.3% BSA for 20 minutes, followed by flow cytometry analysis on a

MACSQuant flow cytometer (Miltenyi). The assays were performed 3 times with 3

technical replicates. The merozoite attachment rates to control or CD55-null RBCs were

calculated as the percent of RBCs with an attached merozoite, and were normalized for

background attachment in the presence of heparin. The merozoite attachment efficiency

was calculated for each genetic background in each biological replicate by normalizing to

the mean of the attachment rates in the three control experiments.

Data analysis

Stastistical analyses were performed and the graphs were generated using GraphPad Prism

8 Version 8.0.2 (159) for macOS.

795
796

Supplemental Information
Supplemental information includes six figures and eight movies.

Multimedia Files

Movie S1: *P. falciparum* 3D7 invasion in the presence of isotype control antibody.

Movie S2: *P. falciparum* 3D7 invasion in the presence of anti-CD55 antibody.

Movie S3: *P. falciparum* 3D7 merozoite-induced deformation with deformation score 1.

Movie S4: *P. falciparum* 3D7 merozoite-induced deformation with deformation score 2.

Movie S5: *P. falciparum* 3D7 merozoite-induced deformation with deformation score 3.

Movie S6: An example of a failed invasion despite successful merozoite-induced

deformation in CD55-null pRBC.

Movie S7: An example of a successful invasion of a wild-type pRBC.

Movie S8: Echinocytosis induced by attached *P. falciparum* merozoite to pRBCs in the

presence of Cytochalasin D. Black arrows: Attachment resulting echinocyte formation.

White arrows: Attachment resulting no echinocyte formation.

References

Alexander, D. L., Arastu-Kapur, S., Dubremetz, J. F., & Boothroyd, J. C. (2006). *Plasmodium falciparum* AMA1 binds a rhoptry neck protein homologous to TgRON4, a component of the moving junction in *Toxoplasma gondii*. *Eukaryot Cell, 5*(7), 1169-1173. doi:10.1128/EC.00040-06

Alexander, D. L., Mital, J., Ward, G. E., Bradley, P., & Boothroyd, J. C. (2005). Identification of the moving junction complex of *Toxoplasma gondii*: a collaboration between distinct secretory organelles. *PLoS Pathog, 1*(2), e17. doi:10.1371/journal.ppat.0010017

Bei, A. K., & Duraisingh, M. T. (2012). Functional analysis of erythrocyte determinants of *Plasmodium infection*. *Int J Parasitol, 42*(6), 575-582. doi:10.1016/j.ijpara.2012.03.006

Besteiro, S., Dubremetz, J. F., & Lebrun, M. (2011). The moving junction of apicomplexan parasites: a key structure for invasion. *Cell Microbiol, 13*(6), 797-805. doi:10.1111/j.1462-5822.2011.01597.x

Besteiro, S., Michelin, A., Poncet, J., Dubremetz, J. F., & Lebrun, M. (2009). Export of a *Toxoplasma gondii* rhoptry neck protein complex at the host cell membrane to form the moving junction during invasion. *PLoS Pathog, 5*(2), e1000309. doi:10.1371/journal.ppat.1000309

Chen, L., Lopaticki, S., Riglar, D. T., Dekiwadia, C., Uboldi, A. D., Tham, W. H., . . . Cowman, A. F. (2011). An EGF-like protein forms a complex with PfRh5 and is required for invasion of human erythrocytes by *Plasmodium falciparum*. *PLoS Pathog, 7*(9), e1002199. doi:10.1371/journal.ppat.1002199

Chu, T. T. T., Sinha, A., Malleret, B., Suwanarusk, R., Park, J. E., Naidu, R., . . . Chandramohanadas, R. (2018). Quantitative mass spectrometry of human reticulocytes reveal proteome-wide modifications during maturation. *Br J Haematol, 180*(1), 118-133. doi:10.1111/bjh.14976

Coley, A. M., Campanale, N. V., Casey, J. L., Hodder, A. N., Crewther, P. E., Anders, R. F., . . . Foley, M. (2001). Rapid and precise epitope mapping of monoclonal antibodies against *Plasmodium falciparum* AMA1 by combined phage display of fragments and random peptides. *Protein Eng, 14*(9), 691-698. doi:10.1093/protein/14.9.691

Collins, C. R., Hackett, F., Strath, M., Penzo, M., Withers-Martinez, C., Baker, D. A., & Blackman, M. J. (2013). Malaria parasite cGMP-dependent protein kinase regulates blood stage merozoite secretory organelle discharge and egress. *PLoS Pathog, 9*(5), e1003344. doi:10.1371/journal.ppat.1003344

858 Cooling, L. (2015). Blood Groups in Infection and Host Susceptibility. *Clin Microbiol Rev, 28*(3),
859   801-870. doi:10.1128/CMR.00109-14
860
861 Coyne, C. B., & Bergelson, J. M. (2006). Virus-induced Abl and Fyn kinase signals permit
862   coxsackievirus entry through epithelial tight junctions. *Cell, 124*(1), 119-131.
863   doi:10.1016/j.cell.2005.10.035
864
865 Crosnier, C., Bustamante, L. Y., Bartholdson, S. J., Bei, A. K., Theron, M., Uchikawa, M., . . .
866   Wright, G. J. (2012). Basigin is a receptor essential for erythrocyte invasion by
867   *Plasmodium falciparum*. *Nature, 480*(7378), 534-537. doi:nature10606 [pii]
868   10.1038/nature10606
869
870 Dreyer, A. M., Matile, H., Papastogiannidis, P., Kamber, J., Favuzza, P., Voss, T. S., . . . Pluschke,
871   G. (2012). Passive immunoprotection of *Plasmodium falciparum*-infected mice
872   designates the CyRPA as candidate malaria vaccine antigen. *J Immunol, 188*(12), 6225-
873   6237. doi:10.4049/jimmunol.1103177
874
875 Dvorak, J. A., Miller, L. H., Whitehouse, W. C., & Shiroishi, T. (1975). Invasion of erythrocytes by
876   malaria merozoites. *Science, 187*(4178), 748-750. doi:10.1126/science.803712
877
878 Egan, E. S. (2017). Beyond Hemoglobin: Screening for Malaria Host Factors. *Trends Genet*.
879   doi:10.1016/j.tig.2017.11.004
880
881 Egan, E. S., Jiang, R. H., Moechtar, M. A., Barteneva, N. S., Weekes, M. P., Nobre, L. V., . . .
882   Duraisingh, M. T. (2015). Malaria. A forward genetic screen identifies erythrocyte CD55
883   as essential for *Plasmodium falciparum* invasion. *Science, 348*(6235), 711-714.
884   doi:10.1126/science.aaa3526
885
886 Funston, G. M., Kallioinen, S. E., de Felipe, P., Ryan, M. D., & Iggo, R. D. (2008). Expression of
887   heterologous genes in oncolytic adenoviruses using picornaviral 2A sequences that
888   trigger ribosome skipping. *J Gen Virol, 89*(Pt 2), 389-396. doi:10.1099/vir.0.83444-0
889
890 Gautier, E. F., Ducamp, S., Leduc, M., Salnot, V., Guillonneau, F., Dussiot, M., . . . Mayeux, P.
891   (2016). Comprehensive Proteomic Analysis of Human Erythropoiesis. *Cell Rep, 16*(5),
892   1470-1484. doi:10.1016/j.celrep.2016.06.085
893
894 Giarratana, M. C., Kobari, L., Lapillonne, H., Chalmers, D., Kiger, L., Cynober, T., . . . Douay, L.
895   (2005). Ex vivo generation of fully mature human red blood cells from hematopoietic
896   stem cells. *Nat Biotechnol, 23*(1), 69-74. doi:nbt1047 [pii]10.1038/nbt1047
897
898 Giarratana, M. C., Rouard, H., Dumont, A., Kiger, L., Safeukui, I., Le Pennec, P. Y., . . . Douay, L.
899   (2011). Proof of principle for transfusion of in vitro-generated red blood cells. *Blood,
900   118*(19), 5071-5079. doi:blood-2011-06-362038 [pii] 10.1182/blood-2011-06-362038
901

Gilson, P. R., & Crabb, B. S. (2009). Morphology and kinetics of the three distinct phases of red
blood cell invasion by *Plasmodium falciparum* merozoites. *Int J Parasitol, 39*(1), 91-96.
doi:10.1016/j.ijpara.2008.09.007

Harvey, K. L., Yap, A., Gilson, P. R., Cowman, A. F., & Crabb, B. S. (2014). Insights and
controversies into the role of the key apicomplexan invasion ligand, Apical Membrane
Antigen 1. *Int J Parasitol, 44*(12), 853-857. doi:10.1016/j.ijpara.2014.08.001

Hendel, A., Bak, R. O., Clark, J. T., Kennedy, A. B., Ryan, D. E., Roy, S., . . . Porteus, M. H. (2015).
Chemically modified guide RNAs enhance CRISPR-Cas genome editing in human primary
cells. *Nat Biotechnol, 33*(9), 985-989. doi:10.1038/nbt.3290

Hu, G., Cabrera, A., Kono, M., Mok, S., Chaal, B. K., Haase, S., . . . Bozdech, Z. Transcriptional
profiling of growth perturbations of the human malaria parasite *Plasmodium falciparum*.
Nat Biotechnol, 28(1), 91-98. doi:nbt.1597 [pii] 10.1038/nbt.1597

Hu, J., Liu, J., Xue, F., Halverson, G., Reid, M., Guo, A., . . . An, X. (2013). Isolation and functional
characterization of human erythroblasts at distinct stages: implications for
understanding of normal and disordered erythropoiesis in vivo. *Blood, 121*(16), 3246-
3253. doi:10.1182/blood-2013-01-476390

Kanjee, U., Gruring, C., Chaand, M., Lin, K. M., Egan, E., Manzo, J., . . . Duraisingh, M. T. (2017).
CRISPR/Cas9 knockouts reveal genetic interaction between strain-transcendent
erythrocyte determinants of *Plasmodium falciparum* invasion. *Proc Natl Acad Sci U S A,
114*(44), E9356-E9365. doi:10.1073/pnas.1711310114

Koch, M., & Baum, J. (2016). The mechanics of malaria parasite invasion of the human
erythrocyte - towards a reassessment of the host cell contribution. *Cell Microbiol, 18*(3),
319-329. doi:10.1111/cmi.12557

Malleret, B., Xu, F., Mohandas, N., Suwanarusk, R., Chu, C., Leite, J. A., . . . Russell, B. (2013).
Significant biochemical, biophysical and metabolic diversity in circulating human cord
blood reticulocytes. *PLoS One, 8*(10), e76062. doi:10.1371/journal.pone.0076062

Miller, L. H., Aikawa, M., Johnson, J. G., & Shiroishi, T. (1979). Interaction between cytochalasin
B-treated malarial parasites and erythrocytes. Attachment and junction formation. *J Exp
Med, 149*(1), 172-184.

Ord, R. L., Rodriguez, M., & Lobo, C. A. (2015). Malaria invasion ligand RH5 and its prime
candidacy in blood-stage malaria vaccine design. *Hum Vaccin Immunother, 11*(6), 1465-
1473. doi:10.1080/21645515.2015.1026496
943

944 Pasvol, G., Chasis, J. A., Mohandas, N., Anstee, D. J., Tanner, M. J., & Merry, A. H. (1989).
945   Inhibition of malarial parasite invasion by monoclonal antibodies against glycophorin A
946   correlates with reduction in red cell membrane deformability. *Blood, 74*(5), 1836-1843.
947

948 Paul, A. S., Egan, E. S., & Duraisingh, M. T. (2015). Host-parasite interactions that guide red
949   blood cell invasion by malaria parasites. *Curr Opin Hematol, 22*(3), 220-226.
950   doi:10.1097/MOH.0000000000000135
951

952 Raymond, C., Tom, R., Perret, S., Moussouami, P., L'Abbe, D., St-Laurent, G., & Durocher, Y.
953   (2011). A simplified polyethylenimine-mediated transfection process for large-scale and
954   high-throughput applications. *Methods, 55*(1), 44-51. doi:10.1016/j.ymeth.2011.04.002
955

956 Reddy, K. S., Amlabu, E., Pandey, A. K., Mitra, P., Chauhan, V. S., & Gaur, D. (2015). Multiprotein
957   complex between the GPI-anchored CyRPA with PfRH5 and PfRipr is crucial for
958   *Plasmodium falciparum* erythrocyte invasion. *Proc Natl Acad Sci U S A, 112*(4), 1179-
959   1184. doi:10.1073/pnas.1415466112
960

961 Richard, D., MacRaild, C. A., Riglar, D. T., Chan, J. A., Foley, M., Baum, J., . . . Cowman, A. F.
962   (2010). Interaction between *Plasmodium falciparum* apical membrane antigen 1 and the
963   rhoptry neck protein complex defines a key step in the erythrocyte invasion process of
964   malaria parasites. *J Biol Chem, 285*(19), 14815-14822. doi:10.1074/jbc.M109.080770
965

966 Riglar, D. T., Richard, D., Wilson, D. W., Boyle, M. J., Dekiwadia, C., Turnbull, L., . . . Baum, J.
967   (2011). Super-resolution dissection of coordinated events during malaria parasite
968   invasion of the human erythrocyte. *Cell Host Microbe, 9*(1), 9-20.
969   doi:10.1016/j.chom.2010.12.003
970

971 Sack, B., Kappe, S. H., & Sather, D. N. (2017). Towards functional antibody-based vaccines to
972   prevent pre-erythrocytic malaria infection. *Expert Rev Vaccines, 16*(5), 403-414.
973   doi:10.1080/14760584.2017.1295853
974

975 Salinas, N. D., Tang, W. K., & Tolia, N. H. (2019). Blood-Stage Malaria Parasite Antigens:
976   Structure, Function, and Vaccine Potential. *J Mol Biol, 431*(21), 4259-4280.
977   doi:10.1016/j.jmb.2019.05.018
978

979 Satchwell, T. J., Wright, K. E., Haydn-Smith, K. L., Sanchez-Roman Teran, F., Moura, P. L.,
980   Hawksworth, J., . . . Baum, J. (2019). Genetic manipulation of cell line derived
981   reticulocytes enables dissection of host malaria invasion requirements. *Nat Commun,*
982   *10*(1), 3806. doi:10.1038/s41467-019-11790-w
983

984 Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of Image
985   Analysis. *Nature Methods, 9*(7), 671-675.
986

987   Scully, E. J., Shabani, E., Rangel, G. W., Gruring, C., Kanjee, U., Clark, M. A., . . . Duraisingh, M. T.
988   (2019). Generation of an immortalized erythroid progenitor cell line from peripheral
989   blood: A model system for the functional analysis of *Plasmodium* spp. invasion. *Am J*
990   *Hematol, 94*(9), 963-974. doi:10.1002/ajh.25543
991

992   Spadafora, C., Awandare, G. A., Kopydlowski, K. M., Czege, J., Moch, J. K., Finberg, R. W., . . .
993   Stoute, J. A. (2010). Complement receptor 1 is a sialic acid-independent erythrocyte
994   receptor of *Plasmodium falciparum*. *PLoS Pathog, 6*(6), e1000968.
995   doi:<u>10.1371/journal.ppat.1000968</u>
996

997   Srinivasan, P., Beatty, W. L., Diouf, A., Herrera, R., Ambroggio, X., Moch, J. K., . . . Miller, L. H.
998   (2011). Binding of *Plasmodium* merozoite proteins RON2 and AMA1 triggers
999   commitment to invasion. *Proc Natl Acad Sci U S A, 108*(32), 13275-13280.
1000   doi:10.1073/pnas.1110303108
1001

1002   Storry, J. R., Reid, M. E., & Yazer, M. H. (2010). The Cromer blood group system: a review.
1003   *Immunohematology, 26*(3), 109-118.
1004

1005   Takahashi, J. Y. S., Yamane S, Tanaka M, Takahashi H, et al. (2008). A case of Inab phenotype
1006   (IFC-) with anti-IFC. *Jpn J Transfusion and Cell Therapy, 54*(2), 258.
1007

1008   Taylor, S. M., & Fairhurst, R. M. (2014). Malaria parasites and red cell variants: when a house is
1009   not a home. *Curr Opin Hematol, 21*(3), 193-200. doi:10.1097/MOH.0000000000000039
1010

1011   Tham, W. H., Healer, J., & Cowman, A. F. (2012). Erythrocyte and reticulocyte binding-like
1012   proteins of *Plasmodium falciparum*. *Trends Parasitol, 28*(1), 23-30.
1013   doi:10.1016/j.pt.2011.10.002
1014

1015   Tiffert, T., Lew, V. L., Ginsburg, H., Krugliak, M., Croisille, L., & Mohandas, N. (2005). The
1016   hydration state of human red blood cells and their susceptibility to invasion by
1017   *Plasmodium falciparum*. *Blood, 105*(12), 4853-4860. doi:10.1182/blood-2004-12-4948
1018

1019   Tonkin, C. J., van Dooren, G. G., Spurck, T. P., Struck, N. S., Good, R. T., Handman, E., . . .
1020   McFadden, G. I. (2004). Localization of organellar proteins in Plasmodium falciparum
1021   using a novel set of transfection vectors and a new immunofluorescence fixation
1022   method. *Mol Biochem Parasitol, 137*(1), 13-21. doi:10.1016/j.molbiopara.2004.05.009
1023

1024   Volz, J. C., Yap, A., Sisquella, X., Thompson, J. K., Lim, N. T., Whitehead, L. W., . . . Cowman, A. F.
1025   (2016). Essential Role of the PfRh5/PfRipr/CyRPA Complex during *Plasmodium*
1026   *falciparum* Invasion of Erythrocytes. *Cell Host Microbe, 20*(1), 60-71.
1027   doi:10.1016/j.chom.2016.06.004
1028

1029   Weiss, G. E., Gilson, P. R., Taechalertpaisarn, T., Tham, W. H., de Jong, N. W., Harvey, K. L., . . .
1030   Crabb, B. S. (2015). Revealing the Sequence and Resulting Cellular Morphology of

1031   Receptor-Ligand Interactions during *Plasmodium falciparum* Invasion of Erythrocytes.

1032   *PLoS Pathog, 11*(2), e1004670. doi:10.1371/journal.ppat.1004670

1033

1034   WHO. (2018). *World Malaria Report 2018*. World Health Organization, Geneva.

1035

1036   Yap, A., Azevedo, M. F., Gilson, P. R., Weiss, G. E., O'Neill, M. T., Wilson, D. W., . . . Cowman, A.

1037   F. (2014). Conditional expression of apical membrane antigen 1 in *Plasmodium*

1038   *falciparum* shows it is required for erythrocyte invasion by merozoites. *Cell Microbiol,*

1039   *16*(5), 642-656. doi:10.1111/cmi.12287

1040