<u>Title:</u> Natural food intake patterns do not synchronize peripheral clocks

<u>Authors:</u> Xiaobin Xie[1,3], Ayaka Kukino[1], Haley E. Calcagno[1], Alec Berman[1], Joseph P. Garner[4],

Matthew P. Butler[1,2]

<u>Running Title:</u> Food is a weak zeitgeber

<u>Author Contributions:</u> Conceptualization, MPB; Methodology, MPB, JPG; Validation, MPB, XX,

HG; Investigation, MPB, XX, AK, HG, AB; Writing, MPB, XX; Supervision, MPB.

<u>Addresses:</u> [1]Oregon Institute of Occupational Health Sciences and [2]Department of Behavioral

Neuroscience, Oregon Health & Science University, Portland, OR. [3]Current address: Key

Laboratory of Protein Modification and Degradation, School of Basic Medical Sciences,

Guangzhou Medical University, Guangzhou 511436, China.. [4]Department of Comparative

Medicine, Stanford University, Stanford, CA.

<u>Corresponding Author:</u>

Matthew P. Butler

Oregon Institute of Occupational Health Sciences

Oregon Health & Science University

3181 SW Sam Jackson Park Road - L606

Portland, OR 97239

<u>butlema@ohsu.edu</u> | T: 503-418-4310

<u>Key words:</u> meal timing, peripheral clocks, liver, kidney, TRF, time restricted feeding

Abstract

Food is thought to synchronize circadian clocks in the body, but this is based on time-restricted

feeding (TRF) protocols. To test whether naturalistic feeding patterns are sufficient to phase-

shift and entrain peripheral tissues, we measured circadian rhythms of the liver, kidney, and

submandibular gland in *mPer2^Luc* mice under different feeding schedules. In ad lib feeding as

well as in a schedule designed to mimic the ad lib pattern, PER2::LUC bioluminescence peaked

during the night as expected. Surprisingly, shifting the scheduled feeding by 12h caused only

small advances (<3h). To isolate the effects of feeding from the light-dark cycle, clock phase

was then measured in mice acclimated to scheduled feeding and housed in constant darkness.

In these conditions, peripheral clock phases were better predicted by the rest-activity cycle than

the food schedule. Under natural feeding patterns, the master pacemaker in the brain sets the

phase of peripheral organs independent of feeding behavior.

Introduction

Healthy physiology is characterized by ~24h circadian rhythms in all tissues that are entrained

(synchronized) to cyclic environmental cues. Disruptions of the circadian system, such as by jet

lag and shift work, increase the risk of poor health including obesity, diabetes, and heart disease

in both humans and animal models (Karlsson et al., 2001; Turek et al., 2005; Suwazono et al.,

2006; Marcheva et al., 2010; Karatsoreos et al., 2011; Pan et al., 2011; Vimalananda et al.,

2015). Circadian disruptions cause adverse health outcomes via internal misalignment (in which

internal clocks become desynchronized from each other) and external misalignment (in which

clocks become desynchronized from external entraining cues) (Davidson et al., 2006; Scheer et

al., 2009; Kalsbeek et al., 2011; Karatsoreos et al., 2011; Buxton et al., 2012; LeGates et al.,

2012; Leproult et al., 2014; Mukherji et al., 2015b).

Internal clock alignment is maintained by the central pacemaker in the suprachiasmatic nucleus

(SCN) of the hypothalamus (Dibner et al., 2010a; Mohawk et al., 2012). The SCN is entrained

by light, and in turn, it coordinates the timing of peripheral clocks via several potential pathways,

one of which is thought to be the SCN's control of eating behavior. In studies of time-restricted

feeding (TRF), during which rodents are typically exposed to 4-12 h of food availability and 12-

20 h of fasting each day, the feeding schedule reliably entrains the circadian phase of peripheral

organ clocks, while the SCN remains entrained to the light-dark cycle (Damiola et al., 2000;

Stokkan et al., 2001). This has led to a general model of circadian entrainment in which the

SCN controls the phase of peripheral clocks via its control of feeding behavior.

TRF imposes long fasting intervals that do not typically occur when food is available ad libitum,

and it is not known whether natural eating patterns play a role in entraining peripheral clocks. To

test this, we measured peripheral organ rhythms in simulated natural feeding conditions in light-

dark cycles and in constant dim light (**Fig. 1, Supplementary Fig. S1**). We focused on the

phase of the liver and kidney, two tissues that are entrained by TRF, and the submandibular

gland, a tissue with a high amplitude oscillator that is insensitive to TRF and instead entrains to

the light-dark cycle (Damiola et al., 2000; Stokkan et al., 2001; Vujovic et al., 2008; Swamy et al.,

2018). Surprisingly, we found that simulated natural feeding had little effect on peripheral clocks:

peripheral oscillators generally remained entrained to either the light-dark cycle or to the

animal's own rest-activity cycle.

Results

<u>Experiment 1. Scheduled feeding in light-dark cycles.</u> We first tested whether simulated natural

food intake patterns were sufficient to shift locomotor activity or peripheral clocks in a light-dark

cycle (LD). After acclimating male mice to custom in-cage automatic feeders (**Supplementary**

Fig. S2), the average Ad-Lib intake pattern was calculated. Both meal size and inter-meal

interval varied across the light-dark cycle (**Fig 2A-C**), with 65% of food intake during the dark

phase. Mean meal size was 10.8 pellets (216 mg) (SD 6.2 pellets) and mean inter-meal interval

was 83 min (SD 62). To simulate natural intake patterns, inter-meal intervals were set to 90 min

Figure 1. Timeline for both experiments, showing time of baseline intake measures, food provision schedules (Ad Lib, Scheduled, Shifted, or TRF), the light-dark cycle (LD or DD), and points at which PER2::LUC imaging was conducted to measure the phase of peripheral organ circadian rhythms.

and the amount provided in each meal varied (**Fig. 2D,**

Supplementary Fig. S3). Note that because meals

were defined by at least 3 pellets, the inter-meal

intervals here are longer than the fasting intervals

(see Discussion). The group average profile was

applied to all mice but the amounts were adjusted to

each mouse's baseline intake. To ensure that mice

ate food when it was provided, the total food during

Scheduled conditions was reduced by 5% compared

to Ad Lib conditions. Body weight was maintained at

99-103% of baseline thereafter (**Supplementary Fig.**

S4). The scheduled food was later shifted 12 h

(Shifted-Food-LD). Finally, mice were switched to time

restricted feeding (TRF-LD), with eight equally sized

meals spread across the light portion of the

photocycle (**Fig. 2D**).

PER2::LUC bioluminescence rhythms from the liver,

kidney, and submandibular gland were assessed in

vivo at baseline and during each different feeding

schedule (**Fig. 3**). In the group-level analysis,

bioluminescence peaked in the late night in all tissues

in Ad Lib, Scheduled, and Shifted-Food conditions,

suggesting that shifting the food schedule did not

phase shift the peripheral clocks.

To capture individual variability, peak phase was

calculated for each mouse and tissue

(**Supplementary Figs. S5, S6**). In this analysis, there

was a significant effect of feeding condition on liver

phase (repeated measures ANOVA, $F_{3,24} = 44.5$,

p<.001) and kidney phase ($F_{3,24} = 19.6$, p<.001), but

there was no effect on the submandibular gland ($F_{3,24}$

= 0.39, p=.76) (**Fig. 3C**). Pairwise comparisons

showed that Scheduled meals did not change clock

Figure 2. A, B. Distributions of Ad Lib meal sizes in 20 mg pellets and inter-meal interval in minutes (n=10). Box plot shows the median, 25th, and 75th percentile. Mean is shown by the circle. C. Both meal size and inter-meal interval varied with time of day during Ad Lib conditions. The light-dark cycle is shown at the bottom, with Zeitgeber Time 0 (ZT0) and ZT12 defined as lights-on and lights-off, respectively. D. The mean pellet drop schedules are plotted for Scheduled, Shifted-Food (12h), and TRF conditions, all conducted in LD.

Figure 3. In vivo imaging and determination of activity phase. Every 4 h, mice were imaged after being lightly anesthetized and injected s.c. with luciferin. A. Ventral images from a representative mouse: note that peak PER2::LUC bioluminescence in the SMG occurs at ZT21 in all conditions, whereas peak bioluminescence in the liver occurs earlier only in the TRF light-fed condition. B. Mean bioluminescence is plotted as a function of time of day (n=9). Statistical results from a 2-way repeated measures ANOVA are shown (main and interaction effects of zeitgeber time and food condition). A significant interaction suggests a change in phase or amplitude of the rhythm. C. The phase of each tissue was calculated from the direction of the mean resultant vector and plotted (± SE). P value indicates the result of a 1 way repeated measures ANOVA (feeding condition as independent variable). Conditions that do not share a letter are significantly different (Tukey test, p<.05). Individuals are shown by gray traces.

phase relative to Ad Lib conditions (Tukey test, n.s.). When the Scheduled feeding was shifted

12 h (Shifted-Food), peripheral tissues continued to peak during the dark, with only small shifts

in liver (2.5±0.7 h) and kidney (2.8±1.1 h) (Tukey test, p<.05), and no advance in the

submandibular gland (**Fig. 3C,** 0.1±0.4 h, n.s.). The kidneys of one mouse and the liver of

another shifted robustly to the new food schedule. Without these, the group phase shifts were

less than 2 h. Conclusions based on circular statistics were similar, with the exception that

Shifted-Food schedule did not significantly advance the kidney clock (**Supplementary Fig. S5,**

Mardia-Watson-Wheeler test, p>.30). These data suggest that naturalistic food intake patterns

are not strong enough zeitgebers (time-setting cues) to phase shift the peripheral circadian

clock system.

The lack of food entrainment was surprising given previous reports from TRF treated mice

(Hatori et al., 2012; Tahara et al., 2012; Swamy et al., 2018), so mice were switched to a 12h

food availability cycle, with 8 equally sized meals presented during the 12 hours of the light

cycle (ZT0-12). After 2 weeks, this schedule was sufficient to significantly advance the liver and

kidney clocks relative to the first Scheduled condition by 6.5 and 6.0 h (**Fig. 3C,**

Supplementary Fig. S5). TRF had no effect on submandibular gland phase.

During different food conditions, mice remained
nocturnal as expected (Hatori et al., 2012; Bray et
al., 2013). The nocturnality index (nocturnal to
diurnal activity ratio) was significantly greater than 1
in all conditions, though there was a significant
reduction in this index during the last TRF-LD
condition (**Supplementary Figs. S7, S8**).

<u>Experiment 2. Scheduled feeding in constant</u>
<u>darkness</u>. In Experiment 1, shifting the simulated
natural food intake did not appreciably shift
locomotor activity or peripheral tissue clock phase.
But the zeitgeber strength necessary to shift clocks
may be much higher than to maintain entrainment,
for example as observed in responses to light
(Butler and Silver, 2011). Therefore, Experiment 2
was conducted on a new cohort of mice to
determine if simulated natural intake patterns were
sufficient to entrain the liver clock in the absence of
a light-dark cycle.

Mice exhibited a normal feeding pattern with 68% of
food consumed during the dark phase (**Fig. 4**). The
average intake profile for the second cohort of mice
was again applied using a 90 min inter-meal interval.
Body weight remained stable (**Fig. S9**), and
transitioning from Ad Lib to Scheduled feeding in LD
did not phase shift the circadian system (**Fig. S10**).

Without changing the Scheduled food condition, the
mice were released into constant dim red light (DD,
0.1 lux), and their activity rhythms began to free-run
according to their endogenous non-24h period. This
immediately established that simulated natural food
patterns do not entrain locomotor activity rhythms. In

Figure 4. A, B. Experiment 2 meal size and inter-meal interval distributions as in Fig. 2. C. Both meal size and inter-meal interval varied with time of day during ad lib conditions. D. The mean food rate was calculated for n=12 mice and applied to all mice, with mean level adjusted for individual intake at baseline. The peak in intake at the end of the night was not as pronounced as in Experiment 1. Here ZT12 and FT12 occur at 1700 PST. E. After free-running in darkness with scheduled feeding, mice were switched to TRF conditions with 12 h food availability with FT12 at 0500 PST.

DD, activity phase and peripheral organ phase were calculated periodically over 3 months, and

the PER2::LUC bioluminescence was analyzed as a function of the rest-activity cycle (circadian

time, with activity onset defined as CT12) or food time (FT12-24 representing the period of

greatest intake/provision and corresponding to the original 1700-0500 dark period) (**Fig. 5**).

PER2::LUC expression was synchronized to the rest-activity cycle: circadian time explained 36-

70% of the variance in the model (**Supplementary Table S1**). In contrast, peripheral rhythms

were desynchronized when plotted against the food intake cycle, and food time only explained

2-9% of variance.

Figure 5. PER2::LUC bioluminescence was measured *in vivo* 6 times/day (n=12 mice). This was done after 5 (circle), 7 (box), 9 (diamond), and 12 weeks (triangle) of housing in constant conditions. The same data are plotted against either the timing of food (left) or the timing of each animal's rest-activity cycle (right). Curves are the best fit cosines for each measurement point. R^2 values are from the full model that combined all four measurement days (**Supplementary Table S1**). The liver and kidneys are much better aligned with circadian rhythms of rest/activity, and thus with the brain's pacemaker, than with the schedule of food intake.

Over the four tests of phase during Scheduled-DD conditions, each tissue's phase was
calculated and plotted circularly as a function of food time or circadian time (**Supplementary**
Fig. S11). Scheduled food did not entrain peripheral organs. Instead, from test to test, the
phase in food time varied widely (Mardia-Watson-Wheeler test, W≥14.9, p≤.021). In contrast,
peripheral tissues were always in phase with circadian time with no test-to-test variability
(W≤10.7, p≥.099).

There was evidence for individual variability in sensitivity to food cues (**Supplementary Fig.**
S12). For each tissue, we tested whether circadian phase clustered better along food time or
circadian time by calculating the overall mean resultant vector from the four phases measured
during constant conditions (4 repeated measures of phase per tissue). Submandibular glands
did not entrain to food and always showed significant clustering in circadian time (**Fig. 6**, r≥.83,
p<.05 for all). In contrast, phase in some kidneys and livers was sensitive to food intake patterns.
One kidney and one liver (not from the same mouse) showed significant phase clustering in
food time, indicating food entrainment. Nevertheless, 8/12 livers and 11/12 kidneys showed
stronger clustering by circadian time (above the dotted line in **Fig. 6**); this was significant in 6/12
livers and 9/12 kidneys.

After 15 weeks in Scheduled-DD conditions, mice were switched to TRF with 12 h of feeding
and 12 h of fasting, to determine whether restricted feeding with a long daily fast could entrain
rhythms in DD. As in Experiment 1, mice were provided 8 equally sized meals presented during
the time of the original photophase (0500-1700 PST). Period shortened in the majority of mice,

Figure 6. Individual variability in liver and kidney entrainment to food. For each tissue in each mouse, a preferred phase was calculated and r calculated from the four measures completed in constant conditions. This was done both in food time and in circadian time. The two r statistics are plotted against each other. Points above the diagonal dotted line indicate that circadian time better predicts phase than food time. Points that fall in the grey shaded area have a significant preferred phase (r>0.82; p<0.05; Rayleigh test for n=4). Note: the liver and kidney with significant phase clustering by food time are not from the same mouse. Activity and tissue phases for the mouse with significant liver food time (empty circle on right) are shown in **Supplementary Fig. S12A**.

but locomotor activity entrained (period = 24h) in only two mice (**Supplementary Fig. S13**).

Though food schedules did not entrain locomotor activity in most mice, TRF did entrain rhythms

of the liver and kidney (**Supplementary Fig. S14**). When tissue phases were plotted by food

time, the liver and kidney phases strongly clustered around FT0 (Rayleigh test, p<.001), similar

to phases observed during Ad Lib conditions. Though the SMG displayed statistically significant

clustering, there was a wide range of phases. The opposite results were obtained when plotted

by circadian time: strong clustering for the SMG and weak clustering for the liver and kidney.

Therefore, TRF in DD effectively entrained kidney and liver but did not entrain either the SMG or

locomotor activity.

Discussion

We show here that natural feeding behavior has little resetting effect on peripheral clocks. In

Experiment 1, a 12-h shift of the simulated natural food provision schedule resulted in small

shifts in liver and kidney and no shift in the SMG. In Experiment 2, we removed the light-dark

cycle and tested whether the scheduled food would continue to entrain peripheral clocks.

Peripheral organ phase was nevertheless much better predicted by the rest-activity cycle; food

significantly entrained only 1 of 12 kidneys and 1 of 12 livers. As a positive control, TRF was an

effective zeitgeber in both experiments, unlike the simulated natural feeding. From these data,

we infer that under normal conditions of food intake, the SCN remains the dominant

synchronizer of peripheral clocks and that this does not require the intermediate step of feeding

behavior.

In TRF experiments, food reliably entrains peripheral organs (Damiola et al., 2000; Stokkan et

al., 2001). This has led to a model of peripheral entrainment in which the SCN entrains patterns

of behavior (activity and food intake), and these behaviors and their metabolic sequelae in turn

entrain peripheral organs. In this experiment of simulated natural feeding, however, the effects

of food intake rhythms on peripheral phase were small. This may be because natural intake

patterns of rodents do not feature long fasting intervals. The extended fast may be necessary to

entrain peripheral clocks via reduced circulating insulin (Mukherji et al., 2015a). The long fasts

are also important for the metabolic health benefits of TRF in rodents and humans (Hatori et al.,

2012; Chaix et al., 2014; Gill and Panda, 2015; Paoli et al., 2019; Stenvers et al., 2019).

Fasting intervals were longer during the day than the night, so by setting a constant inter-meal
interval (90 min), we may have removed a physiologically relevant fasting signal. The shortest
fast needed to entrain peripheral clocks is not known, but is likely between 4 and 8 h, given that
8h but not 4h fasts alter liver clock gene expression (Mukherji et al., 2015a). We therefore
analyzed the distribution of fasting intervals during Ad Lib conditions to determine how often >4h
fasts occur. True fasting intervals (no pellets dropped) are shorter than the inter-meal intervals,
since mice occasionally consumed 1-2 pellets and this was not counted as a meal. The
maximum average fasting intervals per bin in the two experiments were 93 min (ZT4.5-ZT6) and
91 min (ZT3-4.5), similar to the 90 min inter-meal interval imposed on the mice during
scheduled feeding and shorter than the peak mean inter-meal intervals (120–140 min, **Figs. 2,**
4). Of course, the averages may not be informative if individual mice still have long fasts each
day. The incidence of long fasts are shown in **Supplementary Tables S2-S3**. Over 80 mouse-
days under Ad Lib conditions in Experiment 1, only 13 intervals lasted longer than 4 hours
(0.7% of all fasting intervals). In Experiment 2, there were 53 such fasts in 120 mouse days
(2.3%). Based on TRF experiments, it would be the long fasts during the light phase that would
entrain normal night-time peaks of PER2::LUC in peripheral organs. These were not equally
distributed across mice. Though two mice in Experiment 2 had almost one long fast per day,
the other 20 mice had long fasts on half of their days or fewer and five never experienced a
light-phase fast >4 hours. These data match other reports that voluntary inter-meal intervals in
mice rarely exceed 5 h (Jensen et al., 2013). We therefore conclude that naturally occurring
fasting intervals during Ad Lib feeding are unlikely to entrain peripheral clocks.

Our results complement investigations of meal frequency effects on clocks. Kuroda et al. (2012)
show that both meal frequency and size affect the phase angle of entrainment in the liver,
kidney, and submandibular gland. Phase nevertheless generally tracked with the longest fast
each day (8-12h). In experiments of equally spaced meals to eliminate the rhythmic food signal,
peripheral rhythms remain entrained to the SCN in rats, mostly likely via adrenal hormone
signaling (Kalsbeek and Strubbe, 1998; Su et al., 2016). In mice, peripheral clock gene rhythms
are preserved but advanced ~2-4 h under the 6 meal/day schedule compared to ad libitum
feeding (Kuroda et al., 2012; Sen et al., 2017); the amount of advance seems to increase with
caloric restriction. These data establish that feeding rhythms are not necessary to entrain
peripheral clocks, but do not speak to the relative importance of meal-induced cues versus other
SCN-derived cues in ad lib fed mice. In turn, our results suggest that even in the presence of
naturally occurring food-derived metabolic signals, more proximate SCN-controlled cues

orchestrate the peripheral clock system. The SCN communicates time of day information via

neural, endocrine, and physiological pathways (Dibner et al., 2010b; Mohawk et al., 2012).

Potential entrainment mechanisms include glucocorticoids (Balsalobre et al., 2000; Valenzuela

et al., 2008; Pezuk et al., 2012; Cuesta et al., 2015), autonomic nervous system projections

(Ueyama et al., 1999; Cailotto et al., 2005), body temperature (Brown et al., 2002; Buhr et al.,

2010), and melatonin (von Gall et al., 2005).

Two results from the TRF conditions bear discussion: peripheral phase in Expt. 1 and the

differential entraining effect on clocks in Expt. 2. In Experiment. 1, the liver and kidney shifted

~6h, substantially less than the ~12h expected based on restricted feeding (Damiola et al.,

2000; Stokkan et al., 2001; Vollmers et al., 2009) and which we have observed when mice can

eat ad libitum during food availability periods (Swamy et al., 2018). A key difference may be the

use of equal-sized and -spaced meals: similar 4-7h shifts were reported in mice fed 3 meals in

either the light or dark {Kuroda, 2012 #2348}. Set meal sizes prevent the initial gorging

response to food availability that is only observed in light-fed mice (Bray et al., 2013). These

data suggest that both long fasts and a large refeeding response are necessary to fully reset

clocks in day-fed mice. In Experiment 2, locomotor activity rhythms continued to free-run in a

majority of mice despite the 12h feeding schedule during TRF-DD. This is consistent with some

reports (providing 2-6 h of food availability per day) (Abe et al., 1989; Mendoza et al., 2010;

Takasu et al., 2012) and differs from others (4-8 h of food/day) (Feillet et al., 2006; Sheward et

al., 2007). Others have shown an intermediate proportion of mice that entrain to food (44-67%)

(Marchant and Mistlberger, 1997; Refinetti, 2015). Both strain differences and duration of daily

food availability can contribute to this inconsistency (Castillo et al., 2004). The misalignment

between liver/kidney (entrained to food) and activity/submandibular gland (free-running and

synchronized to each other) further underscores that TRF has little effect on the central clock.

To what extent do our results shed light on human peripheral clock control? It is not yet known

whether meals entrain peripheral clocks in humans as they do in mice during TRF because

there are few methods to directly measure peripheral clock phase in humans. Studies with

shifted meal times have demonstrated that meals set the phase of glucose and other metabolite

rhythms, but the effects on clock phase in white blood cells is much smaller (Wehrens et al.,

2017; Skene et al., 2018) – indeed, clock gene phase shifts are similar to what we observed in

mice with physiologically relevant feeding pattern (Experiment 1). Though blood markers may

reflect an averaging of phase information from many tissues, the results suggest that meals are

weak zeitgebers for human clocks even as they drive daily rhythms in metabolic physiology. The

difference may lie in how mice and humans respond to fasting. Unlike in humans, an overnight

fast in a mouse can reduce body weight by 15%, induce a catabolic state, and engage

starvation protection mechanisms (Ayala et al., 2010; Jensen et al., 2013). Therefore, even

though TRF is a better match for human food intake patterns, ad lib-fed mice may better mimic

the gentler waxing and waning of circulating metabolic markers over the day in humans. It will

be of interest then to determine whether the SCN's control of peripheral rhythms via non-food

pathways is stronger in humans than in mice.

This experiment was enabled by home cage automated feeders that allowed precise

measurement of food intake and provision of a set pattern. The use of $mPer2^{Luc}$ mice and in

vivo imaging let us repeatedly sample circadian phase in multiple tissues in individuals,

revealing some inter-individual variability in zeitgeber sensitivity. Glucose, insulin, and other

potential resetting cues were not measured in this study, so although simulated natural food

patterns did not alter behavior and peripheral organ rhythms compared to ad lib intake, whether

and how the food schedule may have altered metabolic signaling remains unknown. Finally, the

scheduled food pattern was imposed by varying meal size, so the contribution of inter-meal

interval variability remains unknown.

These two experiments show that natural food intake patterns have little effect on the phase of

peripheral clocks. These data challenge a major model of circadian clocks in which peripheral

clocks are coordinated by the SCN via its control of food intake. Instead, our results suggest

that during normal feeding conditions, the SCN entrains peripheral clocks via feeding-

independent pathways.

Methods

<u>Animals and housing</u>. $mPer2^{Luc}$ mice on a C57Bl/6 background, bearing a knockin fusion protein

combining the clock gene *Period2* and a firefly luciferase reporter, were purchased from

Jackson Laboratories (B6.129S6-*Per2^{tm1Jt}*/J, Strain Code: 006852) (Yoo et al., 2004). Male mice

for this experiment were bred locally and maintained in a Thoren ventilated caging system in a

pathogen-free barrier facility on pelleted cellulose bedding (BioFresh Performance Bedding, ¼"

pelleted cellulose, Absorption Corp, Ferndale, WA), with food (LabDiet 5L0D) and water

available *ad libitum.* For experiments, adult mice were transferred to an animal enclosure

outside of the barrier facility, and housed individually in cages with a custom pellet feeder (Telos

Discovery Systems, West Lafayette, IN, **Supplementary Fig. S2**), in a single cabinet with light-

dark cycle control (Phenome Technologies, Lincolnshire, IL). The home cage feeder stands at

one end of the cage (Thoren model #1, 19.6 cm×30.9 cm×13.3 cm), and takes up 7.5 cm,

therefore limiting the living space length to 23.4 cm. Flat wire lids were used above the cage,

and to prevent the water bottle from interfering with activity recording (see below), the water

bottle was mounted at the end of cage, with the sipper tube extending through the watering

grommet (7.5 cm from the cage floor). Light was provided by green LEDs (525nm, full width at

half maximum (FWHM) 25nm, 125 lux) from 0500-1700 PST. To aid in husbandry and to

ameliorate potential dim-light induced effects on the circadian system during imaging

procedures, a dim red light was on continually in the cabinet (625nm, FWHM 25nm, 0.2 lux). All

procedures were approved by the Institutional Animal Care and Use Committee of Oregon

Health & Science University.

<u>Feeding</u>. At specified times, the feeders automatically dispensed 20 mg food pellets (Bio Serv.

Rodent Grain-Based Diet Sterile Product #S0163). The presence of the pellet and head entries

into the pellet trough (1.5cm × 1.5cm) were recorded by infrared beam breaks. During ad libitum

feeding, a new pellet was dropped each time the pellet in the trough is removed. A failsafe

mechanism drops another pellet 10 sec later if the first pellet is not detected in the trough. Each

dispensing event, head entry, and pellet removal is time stamped and logged by an Access

2010 database (Microsoft, Seattle, WA); analyses were then conducted on binned counts (10

min bins). Individual meals were identified in the record by feeding bouts of at least three pellets

(≥60 mg) with at least 20 min of fasting preceding and following the meal. Feeders were

checked periodically to ensure that all pellets were eaten prior to the next meal and that no

animals were hoarding food. Troughs were empty in 92% of 278 observations suggesting that

the dispensing schedule determined the intake schedule.

<u>Locomotor Activity.</u> Activity counts were measured by a passive infrared detector (Phidgets

1111_0 motion sensor) mounted 16.5 cm above the cage; data were collected as counts per 10

min bin. Activity was used to calculate the activity profile (mean activity per hour over 14 days)

and nocturnality index (dark phase activity divided by light phase activity over 14 days).

Additionally, circadian period and phase during constant conditions were calculated from

consecutive activity onsets over 7-10 days in ClockLab. Activity onset was defined as Circadian

Time 12 (CT12).

Protocol schedules. Experiment 1. The timeline of events is shown in **Fig. 1A**. At week 0, mice

(21-23 weeks of age, n=10) were transferred to the automatic feeding cages. Mice were

maintained in ad libitum (**Ad Lib**) feeding conditions with a 12:12 light:dark (**LD**) schedule for 4

weeks. From weeks 4-9, mice were exposed to Scheduled Feeding (**Scheduled-LD**) that

simulated the group's average feeding profile. At week 9, the scheduled feeding pattern was

shifted by 12 h (**Shifted-Food-LD**). Finally, at week 15, mice were exposed to time-restricted

feeding (**TRF-LD**) for 5 weeks, in which food was provided only during lights-on. Peripheral

PER2::LUC bioluminescence rhythms were measured once in each feeding condition. For one

mouse, a feeder failure introduced long fasts during the Scheduled and Shifted-Food conditions.

The mouse recovered but his data were excluded a priori from peripheral clock phase analyses,

leaving an analytic dataset of n=9. Experiment 2. Mice, aged 16-17 weeks (n=12), were

transferred to automatic feeding cages as above, and fed **Ad Lib** for 5 weeks (**Fig. 1B**).

Scheduled feeding was imposed at week 5, simulating the ad lib intake pattern for this new

cohort (**Scheduled-LD**). From week 8, the mice were housed in constant dim red light

(**Scheduled-DD**, 0.1 lux). Finally, they were shifted to TRF in DD (**TRF-DD**); food was available

during the initial time of lights-on (0500-1700 PST). As above, peripheral organ rhythms were

measured from PER2::LUC bioluminescence.

Imaging protocol. An in vivo imaging system (Stanford Photonics, Stanford, CA) was used to

measure PER2::LUC bioluminescence in the liver, submandibular gland, and kidney (Tahara et

al., 2012; Swamy et al., 2018); this system includes computer assisted capture of

bioluminescent images via an Electron Magnified (EM) CCD camera (ImageEM, Hamamatsu,

Japan, controlled by Piper software version 2.6.89.18, Stanford Photonics) connected to an

ONYX dark box (Stanford Photonics) in which mice can be maintained under isoflurane

anesthesia on a 37°C temperature controlled stage (mTCII micro-Temperature Controller, Cell

MicroControls, Norfolk, VA). Clock phase was determined from PER2::LUC bioluminescence

measured every 4 h (6 measures in one day). Mice were anesthetized with 2-4% isoflurane to

prevent movement during imaging. Prior to taking the first image, the fur was shaved from

surfaces close to the liver, submandibular gland, and kidneys. Mice were then injected s.c.

(15mg/kg) with D-luciferin potassium salt (Promega, Madison, WI) dissolved in sterile phosphate

buffered saline (30mg/10mL) and filtered (0.2μm). The dorsal and ventral surfaces of the mice

were imaged 8 and 10 min after luciferin injection, respectively. Bioluminescence was captured

by the camera in EM mode (sum of eight 125 ms exposures, gain 500). A brightfield reference

image was taken each time under dim red light (633 nm, FWHM 15 nm, 1.6 lux). After imaging,

the animals were immediately returned to their cages for recovery.

Bioluminescence was scored offline. Each image was opened in ImageJ (NIH), and the intensity

of the 24 bit greyscale image quantified using defined regions of interest, centered on the

brightest areas of the tissue (liver: 6 mm × 6 mm, both kidneys: 25 mm × 20 mm, submandibular

gland: 10 mm × 10 mm).

Statistics. Error is reported as SEM unless otherwise indicated, and all statistical tests were

evaluated as two-sided. Defining time. This experiment combined light cycles, food cycles, and

free-running cycles, so data were analyzed as a function of zeitgeber time, food time, and

circadian time, respectively (**Supplementary Fig. S1**). Zeitgeber time (ZT) is defined by the

light-dark cycle with lights-off at ZT12 and lights-on at ZT0. Food time (FT) is defined similarly,

with the major intake occurring from FT12 to FT0. For example, in Scheduled-LD conditions,

FT12 = ZT12. During TRF, FT12 is defined as the start of food availability. Lastly, Circadian

Time (CT) is specific to each animal's rest/activity cycle, and CT12 is defined as the time of

activity onset. Assessing rhythms. The six in vivo bioluminescence measurements per 24 h

were used to calculated a circadian profile and a peak phase. Because of variation in skin

pigmentation and fur shaving, the six measures were normalized as percent of mean within

mouse within organ within experimental time point (**Supplementary Figure S6**). *Circadian*

profile comparisons: In Experiment 1, circadian profiles of bioluminescence were analyzed by

repeated measures ANOVA with food condition and zeitgeber time as independent factors. In

Experiment 2, bioluminescence patterns were analyzed by cosinor analysis as a function of

circadian time or food time. A mouse's activity rhythm free-runs in constant conditions, so

circadian time drifted in and out of phase with the 24 h scheduled food time. The ability of

circadian time versus food time to predict peripheral organ phase was assessed by comparing

goodness of fit of the cosinor regression (R^2 and AICc). *Phase analysis*: Circadian phase of

individual organs was defined by the direction of the mean resultant vector (**Supplementary Fig.**

S6); this provided point estimates of phase for each tissue in each mouse. Circular statistics

were employed to determine whether there was a preferred phase in each condition (Rayleigh

test) and whether tissue phase differed between feeding conditions or between different weeks

into DD (Mardia-Watson-Wheeler test) (Batschelet, 1981). Individual differences. To analyze

individual differences, phase clustering within mouse and within organ was assessed by the

Rayleigh test (n=4 phase determinations per mouse per organ in DD). The mean resultant

vector lengths (r: r=1 when all points occur at the same phase, r=0 when points are uniformly

distributed around the clock) were plotted to illustrate the strength of phase clustering when

assessed by circadian time versus food time during DD (e.g., **Fig. 6**). <u>Other behaviors</u>.

Nocturnality was analyzed by ordinary regression on log-transformed activity with mouse as a

repeated measure, and are presented as median and 95% confidence interval. Groups are

more nocturnal if the 95% confidence interval does not include 1. Changes in circadian period

and phase were assessed by paired t-test.

Data availability

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

The authors thank Drs. Charles N. Allen, Steven A. Shea, Debra J. Skene, Andrew W. McHill, Mitchell S. Turker, Thijs J. Walbeek, and Brady Kalb for helpful discussions over the course of this work. This work was supported in part by grant NS102962, a New Investigator Grant from the Medical Research Foundation, and by the Oregon Institute of Occupational Health Sciences via funds from the State of Oregon (ORS 656.630).

References

References

Abe H, Kida M, Tsuji K, Mano T. 1989. Feeding cycles entrain circadian rhythms of locomotor
activity in CS mice but not in C57BL/6J mice. Physiol Behav **45**(2): 397-401.

Ayala JE, Samuel VT, Morton GJ, Obici S, Croniger CM, Shulman GI, Wasserman DH,
McGuinness OP. 2010. Standard operating procedures for describing and performing
metabolic tests of glucose homeostasis in mice. Dis Model Mech **3**(9-10): 525-534.

Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schutz G,
Schibler U. 2000. Resetting of circadian time in peripheral tissues by glucocorticoid
signaling. Science **289**(5488): 2344-2347.

Batschelet E. 1981. Circular Statistics in Biology. London, Academic Press.

Bray MS, Ratcliffe WF, Grenett MH, Brewer RA, Gamble KL, Young ME. 2013. Quantitative
analysis of light-phase restricted feeding reveals metabolic dyssynchrony in mice. Int J
Obes (Lond) **37**(6): 843-852.

Brown SA, Zumbrunn G, Fleury-Olela F, Preitner N, Schibler U. 2002. Rhythms of mammalian
body temperature can sustain peripheral circadian clocks. Curr Biol **12**(18): 1574-1583.

Buhr ED, Yoo SH, Takahashi JS. 2010. Temperature as a universal resetting cue for
mammalian circadian oscillators. Science **330**(6002): 379-385.

Butler MP and Silver R. 2011. Divergent photic thresholds in the non-image-forming visual
system: entrainment, masking and pupillary light reflex. Proceedings of the Royal
Society of London. Series B: Biological Sciences **278**(1706): 745-750.

Buxton OM, Cain SW, O'Connor SP, Porter JH, Duffy JF, Wang W, Czeisler CA, Shea SA. 2012.
Adverse metabolic consequences in humans of prolonged sleep restriction combined
with circadian disruption. Sci Transl Med **4**(129): 129ra143.

Cailotto C, La Fleur SE, Van Heijningen C, Wortel J, Kalsbeek A, Feenstra M, Pevet P, Buijs
RM. 2005. The suprachiasmatic nucleus controls the daily variation of plasma glucose
via the autonomic output to the liver: are the clock genes involved? Eur J Neurosci
22(10): 2531-2540.

Castillo MR, Hochstetler KJ, Tavernier RJ, Jr., Greene DM, Bult-Ito A. 2004. Entrainment of the
master circadian clock by scheduled feeding. Am J Physiol Regul Integr Comp Physiol
287(3): R551-555.

Chaix A, Zarrinpar A, Miu P, Panda S. 2014. Time-restricted feeding is a preventative and
therapeutic intervention against diverse nutritional challenges. Cell Metab **20**(6): 991-
1005.

Cuesta M, Cermakian N, Boivin DB. 2015. Glucocorticoids entrain molecular clock components
in human peripheral cells. FASEB J **29**(4): 1360-1370.

Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. 2000. Restricted
feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker
in the suprachiasmatic nucleus. Genes Dev **14**(23): 2950-2961.

Davidson AJ, Sellix MT, Daniel J, Yamazaki S, Menaker M, Block GD. 2006. Chronic jet-lag
increases mortality in aged mice. Curr Biol **16**(21): R914-916.

Dibner C, Schibler U, Albrecht U. 2010a. The mammalian circadian timing system: organization
and coordination of central and peripheral clocks. Annu Rev Physiol **72**: 517-549.

Dibner C, Schibler U, Albrecht U. 2010b. The mammalian circadian timing system: organization
and coordination of central and peripheral clocks. Annual review of physiology **72**: 517-
549.

Feillet CA, Ripperger JA, Magnone MC, Dulloo A, Albrecht U, Challet E. 2006. Lack of food
anticipation in Per2 mutant mice. Curr Biol **16**(20): 2016-2022.

Gill S and Panda S. 2015. A Smartphone App Reveals Erratic Diurnal Eating Patterns in
Humans that Can Be Modulated for Health Benefits. Cell Metab **22**(5): 789-798.

Hatori M, Vollmers C, Zarrinpar A, DiTacchio L, Bushong EA, Gill S, Leblanc M, Chaix A, Joens
M, Fitzpatrick JA, Ellisman MH, Panda S. 2012. Time-restricted feeding without reducing
caloric intake prevents metabolic diseases in mice fed a high-fat diet. Cell Metab **15**(6):
848-860.

Jensen TL, Kiersgaard MK, Sorensen DB, Mikkelsen LF. 2013. Fasting of mice: a review. Lab
Anim **47**(4): 225-240.

Kalsbeek A, Scheer FA, Perreau-Lenz S, La Fleur SE, Yi CX, Fliers E, Buijs RM. 2011.
Circadian disruption and SCN control of energy metabolism. FEBS Lett **585**(10): 1412-
1426.

Kalsbeek A and Strubbe JH. 1998. Circadian control of insulin secretion is independent of the
temporal distribution of feeding. Physiol Behav **63**(4): 553-558.

Karatsoreos IN, Bhagat S, Bloss EB, Morrison JH, McEwen BS. 2011. Disruption of circadian
clocks has ramifications for metabolism, brain, and behavior. Proc Natl Acad Sci U S A
108(4): 1657-1662.

Karlsson B, Knutsson A, Lindahl B. 2001. Is there an association between shift work and having
a metabolic syndrome? Results from a population based study of 27,485 people. Occup
Environ Med **58**(11): 747-752.

Kuroda H, Tahara Y, Saito K, Ohnishi N, Kubo Y, Seo Y, Otsuka M, Fuse Y, Ohura Y, Hirao A,
Shibata S. 2012. Meal frequency patterns determine the phase of mouse peripheral
circadian clocks. Sci Rep **2**: 711.

LeGates TA, Altimus CM, Wang H, Lee HK, Yang S, Zhao H, Kirkwood A, Weber ET, Hattar S.
2012. Aberrant light directly impairs mood and learning through melanopsin-expressing
neurons. Nature **491**(7425): 594-598.

Leproult R, Holmback U, Van Cauter E. 2014. Circadian misalignment augments markers of
insulin resistance and inflammation, independently of sleep loss. Diabetes **63**(6): 1860-
1869.

Marchant EG and Mistlberger RE. 1997. Anticipation and entrainment to feeding time in intact
and SCN-ablated C57BL/6j mice. Brain Res **765**(2): 273-282.

Marcheva B, Ramsey KM, Buhr ED, Kobayashi Y, Su H, Ko CH, Ivanova G, Omura C, Mo S,
Vitaterna MH, Lopez JP, Philipson LH, Bradfield CA, Crosby SD, JeBailey L, Wang X,
Takahashi JS, Bass J. 2010. Disruption of the clock components CLOCK and BMAL1
leads to hypoinsulinaemia and diabetes. Nature **466**(7306): 627-631.

Mendoza J, Albrecht U, Challet E. 2010. Behavioural food anticipation in clock genes deficient
mice: confirming old phenotypes, describing new phenotypes. Genes Brain Behav **9**(5):
467-477.

Mohawk JA, Green CB, Takahashi JS. 2012. Central and peripheral circadian clocks in
mammals. Annu Rev Neurosci **35**: 445-462.

Mukherji A, Kobiita A, Chambon P. 2015a. Shifting the feeding of mice to the rest phase creates
metabolic alterations, which, on their own, shift the peripheral circadian clocks by 12
hours. Proc Natl Acad Sci U S A **112**(48): E6683-6690.

Mukherji A, Kobiita A, Damara M, Misra N, Meziane H, Champy MF, Chambon P. 2015b.
Shifting eating to the circadian rest phase misaligns the peripheral clocks with the
master SCN clock and leads to a metabolic syndrome. Proc Natl Acad Sci U S A
112(48): E6691-6698.

Pan A, Schernhammer ES, Sun Q, Hu FB. 2011. Rotating night shift work and risk of type 2
diabetes: two prospective cohort studies in women. PLoS Med **8**(12): e1001141.

Paoli A, Tinsley G, Bianco A, Moro T. 2019. The Influence of Meal Frequency and Timing on
Health in Humans: The Role of Fasting. Nutrients **11**(4).

Pezuk P, Mohawk JA, Wang LA, Menaker M. 2012. Glucocorticoids as entraining signals for
peripheral circadian oscillators. Endocrinology **153**(10): 4775-4783.

Refinetti R. 2015. Comparison of light, food, and temperature as environmental synchronizers of
the circadian rhythm of activity in mice. J Physiol Sci **65**(4): 359-366.
Scheer FA, Hilton MF, Mantzoros CS, Shea SA. 2009. Adverse metabolic and cardiovascular
consequences of circadian misalignment. Proc Natl Acad Sci U S A **106**(11): 4453-4458.
Sen S, Raingard H, Dumont S, Kalsbeek A, Vuillez P, Challet E. 2017. Ultradian feeding in mice
not only affects the peripheral clock in the liver, but also the master clock in the brain.
Chronobiol Int **34**(1): 17-36.
Sheward WJ, Maywood ES, French KL, Horn JM, Hastings MH, Seckl JR, Holmes MC, Harmar
AJ. 2007. Entrainment to feeding but not to light: circadian phenotype of VPAC2
receptor-null mice. J Neurosci **27**(16): 4351-4358.
Skene DJ, Skornyakov E, Chowdhury NR, Gajula RP, Middleton B, Satterfield BC, Porter KI,
Van Dongen HPA, Gaddameedhi S. 2018. Separation of circadian- and behavior-driven
metabolite rhythms in humans provides a window on peripheral oscillators and
metabolism. Proc Natl Acad Sci U S A **115**(30): 7825-7830.
Stenvers DJ, Scheer F, Schrauwen P, la Fleur SE, Kalsbeek A. 2019. Circadian clocks and
insulin resistance. Nat Rev Endocrinol **15**(2): 75-89.
Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M. 2001. Entrainment of the circadian clock
in the liver by feeding. Science **291**(5503): 490-493.
Su Y, Cailotto C, Foppen E, Jansen R, Zhang Z, Buijs R, Fliers E, Kalsbeek A. 2016. The role of
feeding rhythm, adrenal hormones and neuronal inputs in synchronizing daily clock gene
rhythms in the liver. Mol Cell Endocrinol **422**: 125-131.
Suwazono Y, Sakata K, Okubo Y, Harada H, Oishi M, Kobayashi E, Uetani M, Kido T, Nogawa
K. 2006. Long-term longitudinal study on the relationship between alternating shift work
and the onset of diabetes mellitus in male Japanese workers. J Occup Environ Med
48(5): 455-461.
Swamy S, Xie X, Kukino A, Calcagno HE, Lasarev MR, Park JH, Butler MP. 2018. Circadian
disruption of food availability significantly reduces reproductive success in mice. Horm
Behav **105**: 177-184.
Tahara Y, Kuroda H, Saito K, Nakajima Y, Kubo Y, Ohnishi N, Seo Y, Otsuka M, Fuse Y, Ohura
Y, Komatsu T, Moriya Y, Okada S, Furutani N, Hirao A, Horikawa K, Kudo T, Shibata S.
2012. In vivo monitoring of peripheral circadian clocks in the mouse. Curr Biol **22**(11):
1029-1034.

Takasu NN, Kurosawa G, Tokuda IT, Mochizuki A, Todo T, Nakamura W. 2012. Circadian
regulation of food-anticipatory activity in molecular clock-deficient mice. PLoS One **7**(11):
e48892.
Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S,
Easton A, Jensen DR, Eckel RH, Takahashi JS, Bass J. 2005. Obesity and metabolic
syndrome in circadian *Clock* mutant mice. Science **308**(5724): 1043-1045.
Ueyama T, Krout KE, Nguyen XV, Karpitskiy V, Kollert A, Mettenleiter TC, Loewy AD. 1999.
Suprachiasmatic nucleus: a central autonomic clock. Nat Neurosci **2**(12): 1051-1053.
Valenzuela FJ, Torres-Farfan C, Richter HG, Mendez N, Campino C, Torrealba F, Valenzuela
GJ, Seron-Ferre M. 2008. Clock gene expression in adult primate suprachiasmatic
nuclei and adrenal: is the adrenal a peripheral clock responsive to melatonin?
Endocrinology **149**(4): 1454-1461.
Vimalananda VG, Palmer JR, Gerlovin H, Wise LA, Rosenzweig JL, Rosenberg L, Ruiz Narvaez
EA. 2015. Night-shift work and incident diabetes among African-American women.
Diabetologia.
Vollmers C, Gill S, DiTacchio L, Pulivarthy SR, Le HD, Panda S. 2009. Time of feeding and the
intrinsic circadian clock drive rhythms in hepatic gene expression. Proc Natl Acad Sci U
S A **106**(50): 21453-21458.
von Gall C, Weaver DR, Moek J, Jilg A, Stehle JH, Korf HW. 2005. Melatonin plays a crucial
role in the regulation of rhythmic clock gene expression in the mouse pars tuberalis. Ann
N Y Acad Sci **1040**: 508-511.
Vujovic N, Davidson AJ, Menaker M. 2008. Sympathetic input modulates, but does not
determine, phase of peripheral circadian oscillators. Am J Physiol Regul Integr Comp
Physiol **295**(1): R355-360.
Wehrens SMT, Christou S, Isherwood C, Middleton B, Gibbs MA, Archer SN, Skene DJ,
Johnston JD. 2017. Meal Timing Regulates the Human Circadian System. Curr Biol
27(12): 1768-1775.
Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Siepka SM, Hong HK, Oh
WJ, Yoo OJ, Menaker M, Takahashi JS. 2004. PERIOD2::LUCIFERASE real-time
reporting of circadian dynamics reveals persistent circadian oscillations in mouse
peripheral tissues. Proc Natl Acad Sci U S A **101**(15): 5339-5346.