1

From plants to ants: Fungal modification of leaf lipids for nutrition and communication in

the leaf-cutter ant fungal garden ecosystem

Authors
Lily Khadempour[1,†], Jennifer E. Kyle[2], Bobbie-Jo M. Webb-Robertson[2], Carrie D. Nicora[2], Francesca B. Smith[2], Richard D. Smith[2], Mary S. Lipton[3], Cameron R. Currie[1,#], Erin S. Baker[4,#], Kristin E. Burnum-Johnson[3,5,#]

Affiliations
[1]*Department of Bacteriology, University of Wisconsin-Madison, Madison, WI 53706, USA*

[2]*Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99352, USA*

[3]*The Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, WA, 99352, USA*

[4]*Department of Chemistry, North Carolina State University, Raleigh, NC 27607, USA*

†*Present address: Department of Earth and Environmental Sciences, Rutgers University-Newark, Newark, NJ 07102, USA*

[5]*Lead Contact*

[#]Co-corresponding authors:

Kristin E. Burnum-Johnson, Email: kristin.burnum-johnson@pnnl.gov
Erin S. Baker, Email: ebaker@ncsu.edu
Cameron Currie, Email: currie@bact.wisc.edu

Abstract

Lipids are essential to all living organisms, as an energy source, as an important cellular

structural component, and as a communication tool. In this study, we used global lipidomic

methods to evaluate the lipids in leaf-cutter ant fungal gardens. Leaf-cutter ants and their

coevolved fungal cultivar, *Leucoagaricus gongylophorus*, are a model mutualistic system. The

fungus enzymatically digests fresh plant material that the ants cut and deliver, converting energy

and nutrients from plants, and providing them to the ants through specialized hyphal swellings

called gongylidia. Using combined liquid chromatography, ion mobility spectrometry, and

tandem mass spectrometry we evaluated differences between the molecular speciation of lipids

in the leaf-cutter ant fungal garden ecosystem. This lipidomic study characterized leaves that are

fed to the gardens, gongylidia that are produced by the fungus to feed the ants, and spatially

resolved regions of the fungal garden through stages of leaf degradation. Lipids containing

alpha-linolenic acid (18:3) were enriched in leaves and the top of the gardens, but not dominant

in the middle or bottom regions. Gongylidia were dominated by lipids containing linoleic acid

(18:2). To evaluate the communicative potential of the lipids in fungal gardens we conducted a

behavioral experiment that showed *Atta* leaf-cutter ants responded differently to 18:3 and 18:2

fatty acids, with aggression towards 18:3 and attraction for 18:2. This work demonstrates the role

of lipids in both the transfer of energy and as an inter-kingdom communication tool in leaf-cutter

ant fungal gardens.

Importance

In this work we examined the role of lipids in the mutualism between leaf-cutter ants and fungus.

These ants cut fresh leaf material, which they provide to their fungal cultivar, that converts

energy and nutrients from the plants and provides it to the ants in specialized hyphal swellings

called gongylidia. This work constitutes the first example of a global lipidomics study of a

symbiotic system and provides insights as to how the fungus modifies plant lipids into a usable

source for the ants. Through a behavioral experiment, this work also demonstrates how lipids can

be used as an inter-kingdom communication tool, in this case an attractant, rather than as a

repellant, which is more often seen.

Introduction

60
Lipids are one of four fundamental components of organisms, the other three being

proteins, carbohydrates, and nucleic acids. Lipids serve many roles, and their characterization is

critical to understanding how organisms function individually, and in communities. Two roles

that we are particularly interested in are as a source of energy and their use in interspecies

communication. Lipids are widely used as a form of stored energy in organisms, and in diets

represent a dense source of energy, approximately double that of either proteins or carbohydrates

(1).

The role of lipids in communication is vital across biological scales. Lipids found in cell

membranes are important in cell-to-cell communication, either as a means to identify and

differentiate between cells of the same multicellular organism (2), conspecifics in unicellular

organisms (3), or cells that do not belong – pathogens (4), predators (5), or food (6). Lipids even

direct communication between host cells and virus, thereby impacting the favorability of

membrane fusion events (7). At a larger scale, lipids are used as signaling molecules in the form

of hormones (between cells of the same organism), pheromones (intraspecific communication),

and allelochemicals (interspecific communication).

Previous work on allelochemicals in symbiosis has focused largely in the realm of plant

defense against herbivory. When consumed by insect herbivores, some plants emit

allelochemicals that attract the herbivore's natural enemy, parasitic wasps (8). This system is a

fascinating example of communication in a multipartite symbiosis between a host, herbivore, and

indirect defensive symbiont. We expect that this type of communication would exist in beneficial

symbioses as well. Our work here explores the role of lipids, in both nutrition and

communication, in a model mutualistic system where fungi serve as an interface between ants

and plants.

Leaf-cutter ants are dominant herbivores that can consume as much as 17% of the leaf

biomass produced in Neotropical ecosystems (9). *Atta* leaf-cutter ant colonies contain millions of

workers that fall into specialized casts (10), contributing to the colony's success as ecosystem

engineers (11). Foraging leaf-cutter ants are often seen marching in conspicuous trails and

bringing pieces of leaves back to their underground nests that can be as large as 30 m^2 to 35

m^2 in area and several meters deep. Once to the nest, the ants use the leaves to feed their fungal

cultivar, *Leucoagaricus gongylophorus* (12), which is grown in their specialized fungal gardens.

By cultivating and feeding these fungal gardens, the ants are able to access nutrients in plant

biomass that would otherwise be unavailable (13-17).

Leaf-cutter ants and their microbial symbionts are paradigmatic in the field of insect

microbial symbiosis. Unlike other herbivores, the microbes in symbiosis with leaf-cutter ants can

be easily sampled because they exist as an external digestive system in the form of fungal

gardens. These fungal gardens exhibit vertical stratification, characterized by top, middle and

bottom regions. From top to bottom, there is a visual gradient of leaf degradation with leaves

starting as fresh, undigested leaves at the top, then processing through the initial stages of

degradation in the middle, and finally to more complete stages of degradation by the bottom

(18). To promote growth of the fungal garden, leaf-cutter ants perform different activities in the

various locations of the garden. Specifically, the ants triturate fresh plant material, incorporate it

into the top of the fungal garden, deposit tufts of fungal mycelium, and defecate with excrement

containing fungus-derived plant biomass-degrading enzymes to initiate lignocellulose

degradation (19, 20). As the plant material placed on the top of the fungal garden is digested, it

moves toward the bottom of the garden where it is removed from the fungal garden chamber and

deposited in a refuse dump by the ants (21), and by which time it has been depleted of most

nutrients. The gradual degradation of the plant material from top to bottom of the garden,

combined with specific ant activities, gives each region of the fungal garden a different visual

appearance and varied molecular properties (**Figure 1**). The middle of the fungal garden, where

the density of fungal hyphal growth is most pronounced, is also where the *L. gongylophorus*

produces gongylidia, or specialized hyphal swellings, that worker ants consume and feed to

larvae in the colony. While foraging workers obtain some nutrition from the plant sap they drink

when they cut the fresh leaf material, garden workers and larvae exclusively consume

gongylidia, where they obtain carbohydrates, polysaccharides, and energy-rich lipids (22, 23).

To elucidate the relationship between leaf-cutter ants and their fungal cultivar, earlier

studies have focused on understanding how the fungus breaks down plant biomass (13, 14). In

our previous metaproteomic analyses of the top, middle and bottom regions of the fungal garden,

we discovered that *L. gongylophorus* produced a majority of the lignocellulases found (13). The

studies also showed the resident bacterial community likely aids in this process by producing

amino acids and vitamins, thereby enabling the fungus to thrive (13, 14, 24). These results shed

great light into the symbiosis between the ant and the fungal cultivar, but to understand a more

complete picture other proteins were evaluated. Interestingly, the gongylidia (13, 20) showed an

enrichment in unique lipid-associated proteins when compared to different regions of the garden

(Data file S1), ultimately suggesting the gongylidia lipids would differ from those throughout the

fungal garden. A detailed analysis of the molecular speciation of lipids is needed to provide

biological insight into subpopulations and differential activities of complex samples, such as the

fungal garden ecosystem.

Here, we conduct the first global lipidomic study of the leaf-cutter ant-microbe symbiosis

using comprehensive liquid chromatography–tandem mass spectrometry and liquid

chromatography–ion mobility spectrometry-mass spectrometry. We examined spatiotemporal

changes in the molecular speciation of lipids across six heterogenous *Atta* leaf-cutter fungal

gardens through different stages of leaf degradation. First we evaluated the lipid content of

leaves the ants use to feed their cultivar to understand which ones existed initially. Then we

assessed the gongylidia, and the top, middle, and bottom regions of their fungal gardens at initial,

intermediate, and advanced stages of leaf degradation, respectively (**Figure 1**). The lipid content

of the leaf material was compared to the different regions of the fungal garden to understand

consumption of the leaf lipids and synthesis of the fungal garden consortium lipids through the

various regions. Additionally, the lipid content of the gongylidia was compared to the middle

region of the fungal garden to evaluate its specific properties versus the area where it was

harvested. Upon finding an enrichment of particular lipids in the leaves and fungal garden

components, we conducted a behavior experiment using seven *Atta cephalotes* colonies to

observe whether worker ants detected and responded to the lipids differently.

143
144   **Results**
145
We observed that the predominant lipids in the gongylidia and top and bottom of the

garden varied greatly. The top of the garden was most similar to the leaves and had predominant

lipid subclasses including monogalactosyldiacylglycerol (MGDG),

sulfoquinovosyldiacylglycerol (SQDG), and diacylglycerophosphoglycerol (PG), while the lipids

in the bottom of the gardens included diacylglycerophosphoethanolamines (PE) and

diacylglycerophosphoserine (PS). The gongylidia had dominant lipids including ceramides (Cer),

diacylglycerophosphoethanolamines (PE), and triacylglycerols (TG). Across all lipid subclasses

interesting trends were observed based on individual fatty acid composition; an evaluation of the

lipid subclasses and fatty acid composition differences is given below.

Lipid comparisons of leaves and fungal garden regions

To initiate the study, lipids present in the maple leaves fed to the fungal garden were evaluated

so they could be compared to the fungal garden regions and gongylidia. The lipids present in the

leaves were evaluated and leaf lipid categories included sphingolipids, glycerophospholipids, and

glycerolipids. These identifications were dominated by the PG subclass as well as galactolipids

(MGDG and SQDG) with only minor amounts of the other phospholipids

(diacylglycerophosphocholines (PC), PE, and diacylglycerophosphoinositols (PI)) (see Data file

S2 for all characterized lipids). Evaluation of the fungal garden samples showed that 274 lipids

were identified from the top, middle, and bottom regions of the six gardens (Data file S2 (a) &

(b)). Both the leaf lipids and garden lipids are depicted in **Figure 2** where the relative $\log_2$

expression level of each lipid is represented by a red-white-blue color scale with red denoting

high abundance and blue low abundance. Leaf lipids are included in the top row (highlighted

with a pink box), subsequent rows contain garden lipids, for the six top regions (highlighted with

a green bar), six middle regions (highlighted with a teal bar), and six bottom regions (highlighted

with an orange bar). In leaf samples, PGs and galactolipids were observed to be the most

abundant, which was distinct from the lipids detected in the top, middle, and bottom regions of

the garden, where TGs were found to be most abundant.

Garden lipids were also from sphingolipids, glycerophospholipids, and glycerolipids

categories and 13 subclasses. The glycerophospholipid and glycerolipid categories represented

97% of the identified lipids with 89 glycerophospholipids species and 177 glycerolipids species,

while the sphingolipid category was only a minor component with 8 species identified in the

garden (Data file S2(b)). Figure S1 and Data file S2(c) depict 110 lipids, including isomeric

species (25), with accurate quantitative values and no co-eluting lipid species detected. When the

top and bottom fungal garden regions were statistically compared to evaluate lipid changes and

leave degradation, lipid species were found to be statistically significant (p-value < 0.05) and a

trend was noted pertaining to chain length and degree of unsaturation. PE, PG, diacylglycerol

(DG), and TG species, were more unsaturated (i.e., contained more double bonds in the fatty acid

chains) in the top region of the garden, and PG and TG species had longer fatty acid chains.

An examination of the individual fatty acid composition of the lipids in the fungal garden

also illustrated interesting trends. The fatty acyl 16:0, 18:2, and 18:3 groups were found to

dominate the fatty acids profiles for the most abundant lipids identified in the phospholipid and

glycerolipid subclasses. **Figure 2** details the acyl chains for the 59 significantly enriched lipids

(p-value < 0.05), where those highlighted in green text (50 lipids) or orange text (9 lipids) are

significantly enriched in the top or bottom garden layer, respectively. For example,

PC(18:3_18:3), LPC(18:1)_A, PE(18:3_18:3), and PI(18:3_18:3) are highlighted with green text

in **Figure 2** since they greatly decreased from the top to bottom region of the fungal garden. Of

the 50 lipid species significantly elevated in the top layer, over half of them contained an 18:3

acyl chain. These 18:3 containing lipids significantly more abundant in the top layer include

MGDG, SQDG, monoacylglycerophosphocholine (LPC), PC, PG, DG, Cardiolipin (CL), PI,

monoacylglycerophosphoethanolamines (LPE), and PE species (signified with an * in **Figure 2**)

and TG(18:3/18:3/18:3) (Data file S2(b) denotes the 12 significant TGs, where no co-eluting

species were detected, increasing in the top layer). Many of these lipids also had a relatively high

abundance in the leaves, especially those containing two 18:3 fatty acyl groups (e.g., 18:3_18:3),

suggesting 18:3 containing lipids come from the leaves. Only nine lipids were found to

significantly increase in the bottom layer of the garden compared to the top and, all of these

lipids contained at least one 18:2 (signified with a [‡] in **Figure 2**), suggesting 18:3 is degraded

from the top to bottom regions of the garden by fungal garden microbial symbionts, while 18:2 is

synthesized by the fungal garden. Based on these results we hypothesized that 18:2 could be

enriched in the gongylidia.

Lipid changes between fungal garden and gongylidia

Next, the gongylidia were compared to the middle fungal garden where they are located

to assess similarities and differences. In the gongylidia, 263 lipids species were identified from

18 different subclasses (Data file S3 (a) & (b)). Figure S2 depicts 183 lipids, including isomeric

species, with accurate quantitative values and no co-eluting lipid species detected (Data file

S3(c)). From the statistical comparison of the gongylidia and surrounding middle garden region,

Figure 3 depicts the 116 lipid species found to be statistically significantly enriched, indicating

the gongylidia lipids are distinct from the surrounding middle region. Both the middle region

garden lipids and gongylidia lipids are depicted in **Figure 3** where the relative $\log_2$ abundance of

each lipid is represented by a red-white-blue color scale, red denotes high abundance and blue

denotes low abundance. Leaf lipids are included in the top row (highlighted with a pink box),

subsequent rows contain garden lipids, for the six middle regions (highlighted with a teal bar),

and six gongylidia samples (highlighted with a purple bar). The acyl chains for each lipid are

also detailed in **Figure 3**, where those highlighted in purple text or teal text are significantly

increasing in the gongylidia or surrounding middle garden region, respectively. Lipid categories

with high abundance in leaves from the initial garden study, MDGD, SQDG, and PG, in addition

to DGDG, showed low enrichment in the gongylidia samples; whereas, both detected

mannosylinositol phosphorylceramide (MIPC) lipids, MIPC(t18:0_24:0(2OH)) and

MIPC(t18:0_26:0(2OH)), and all five significant Cer species were enriched in the gongylidia.

PC, PE, phosphatidic acid (PA), PI, DG, and TG lipids showed differential enrichment

dependent on the lipid fatty acyl composition. All significantly changing 18:3 containing PC, PE,

PA, and DG species in the middle garden were depleted in the gongylidia (signified with an * in

Figure 3). Conversely, all significantly changing 18:2 containing PC, PE, PA, DG, and PI

species were enriched in the gongylidia (signified with a [‡] in **Figure 3**). Many TG species

contained both 18:3 and 18:2 as one of their three fatty acyl chains, yet ninety percent of all TG

species depleted in the gongylidia compared to the surrounding fungal garden contain at least

one 18:3 acyl chain and >84% of all TG species enriched in the gongylidia contain at least one

18:2 acyl chain. Additionally, lipid subclasses with high abundance in the leaves (MDGD,

SQDG, PG, and DGDG) showed relatively low concentrations in the gongylidia samples.

Behavior experiment

Our lipidomic analyses characterized a prevalence of lipid species containing linoleic

acid (18:2) in the gongylidia, the leaf-cutter ant's food source, and lipid species containing alpha-

linolenic acid (18:3) in leaf material. We performed a behavior experiment to test if the *Atta*

cephalotes workers had different responses to untreated paper discs and paper discs treated with

oleic acid (18:1), 18:2, and 18:3 (**Figure 4**; Figure S3, S4, & S5; Movie S1 & S2). Overall, the

ants spent significantly more time inspecting 18:2 (p=0.0012) and 18:1 (p=0.015) treated paper

discs than the control paper discs. Similarly, the ants picked up the discs infused with 18:2

(p=0.039) more frequently than the control discs, but during the experiment never picked up a

disc containing 18:3. However, the ants displayed aggressive behavior, including lunging and

biting the discs more frequently toward 18:3 compared to the control (p=0.014). The ants did not

exhibit aggressive behavior towards the discs with 18:1 or 18:2.

245
Discussion
Leaf-cutter ants, from the genera *Atta* and *Acromyrmex*, cultivate a fungus on fresh foliar

material, which converts plant biomass into new forms that are readily usable by the ants and

their larvae. This obligate mutualism is derived from within the ant subtribe Attina, which

represents a monophylogenetic lineage of ants that have associated with fungal cultivars for

approximately 55 million years. The leaf-cutter ants are the most derived group of the Attina and

diverged approximately 12 million years ago, coinciding with their association with *L.*

gongylophorus (26). Both organisms have undergone co-evolution. *Leucoagaricus*

gongylophorus has adapted to its association with the ants and has undergone gene reduction so

that it can no longer function as a saprotroph, like its closest free-living relatives (27). On the

other hand, the ant's digestive system has adapted to prevent digestion of fungus-produced plant

biomass degradation enzymes so that the ants can redistribute them through fecal fluid onto

freshly incorporated leaf material (19). In return, the fungus provides all of the nutrients to feed

the colony by producing carbohydrate, polysaccharide, and lipid rich gongylidia (22, 23).

In this study, characterization of both the lipid content of leaf material and lipids

synthesized by the fungus, allowed us to gain novel insight into the conversion of lipid nutrients

from the leaf material deposited on the top of the fungal garden to the enrichment of lipid

nutrients in the gongylidia for the ant to consume. We observed significant subclass trends

correlated to each region. There is an enrichment of lipids containing 18:3 in the leaves and the

top of the garden, and an enrichment of lipids containing 18:2 at the bottom of the garden and

gongylidia. Lipid subclasses that were significantly increased in the top of the garden and leaf

material included PG, MGDG, SQDG, and DGDG species, all of which are known to be

significant contributors to photosynthetic organelles (28). In addition, 18:3 has notable signaling

properties in plant tissues. When leaves are wounded, the polypeptide systemin is emitted from

the damaged cells into the apoplast, signaling the liberation of 18:3 from plant membrane lipids

into the cells. 18:3 begins the defense pathway by being converted to 12-oxophytodienoic acid

and then through beta-oxidation is converted into jasmonic acid (29). The main defense

mechanism in these leaves is jasmonic acid, and 18:3 is crucial to its synthesis.

The abundant lipids in the top of the fungal garden through to the gongylidia differ

greatly. This is an important result because while the ants ingest biomass-degrading enzymes

from the gongylidia and deposit them on top, the lipids are not being recycled in this manner. It

is clear then that the fungal garden metabolizes the lipids from the leaves and synthesizes new

lipids which are provided to the ants and larvae through the gongylidia, where they are ingested

and absorbed into their bodies. While this was assumed previously (22, 23), our study is the first

to actually track the molecular speciation and dynamic changes of these lipids through the fungal

garden.

We conducted the behavior experiment because we wanted to evaluate whether the ants

were able to detect the various lipids in the fungal gardens, and to observe any respective

differences in their behavior. We expected *Atta cephalotes* workers to be attracted to all three

tested lipids. Previous studies had shown that ants from the genera *Pogonomyrmex, Solenopsis,*

and *Atta* are attracted to 18:1, which is a lipid that dead ants emit from their bodies as a cue for

workers to move them out of the colony or toward the dump (30-32). We observed similar

attractive results with our ants, validating our methods. We expected that the workers would also

be attracted to 18:3, which is abundant in leaves they routinely cut and incorporate into their

gardens. However, the workers instead behaved aggressively toward 18:3 by lunging and biting

at the infused discs indicating that it is not this component of leaves that leaf-cutter ants are

attracted to. This ant response to the 18:3 lipids might be related to the lipid's role as a

constituent of the plant defense response lipid, jasmonic acid, which may be playing a role in

deterring herbivory by leaf-cutter ants. Finally, we also expected that the *Atta* workers would be

attracted to 18:2 since it has been shown attractive to another ant species (*Cataglyphis fortis*)

(33) and is abundant in the gongylidia they consume. The ants' attractive responses to 18:2 (both

disc pickup and time spent inspecting) differed most greatly from the controls, compared to the

two other lipids tested, which suggests that there is a strong attraction to this lipid. These

responses all relate back to the fungal garden metabolizing 18:3 containing lipids from leaves to

synthesize 18:2. 18:3 is repulsive to the ants, while 18:2 is highly attractive to them. One very

clear example of coevolution between the leaf-cutter ants and *L. gongylophrous* is the cultivar's

morphological and physiological adaptation in the production of gongylidia for ant consumption

(34). Our result, indicating that the gongylidia contain lipids that the leaf-cutter ants are attracted

to, provides a further example of the intricacies of coevolution through lipid communication in

this system. By restricting the enrichment of 18:2 lipids to the gongylidia, the fungus can focus

the ants' consumption to these specialized structures, thereby preventing damage to its hyphae.

Such communication from the fungus to the ants is vital for maintaining fungal fitness, and for

maintaining this mutualism.

The study of fungal allelochemicals that influence insect behavior is not new, but we

have taken a different approach in addressing this topic. Typically, behavior is observed between

an insect and a fungus, and then work is done to identify the allelochemicals involved in that

communication (35). Often, after identifying these allelochemicals, they are used as tools in pest

management (36). In the leaf-cutter ant literature as well, the approach has been largely from the

direction of behavior to mechanism, and it has been suggested that the cultivar uses volatile

semiochemicals to communicate with the ant colony (37) but attempts at identifying these have

not been conclusive (38, 39). Our approach was different. We were interested in identifying and

quantifying lipids in the fungal garden, and then based on proteomic and lipidomic data,

hypothesized that these lipids could be used as allelochemicals. Our behavioral experiment then

supported this and provides a new tract of study for understanding communication between leaf-

cutter ants and their cultivar.

As an important constitute of food, lipids act as a valuable source of energy, cellular

structural component, and as a communication tool. Thus, understanding what lipids are

consumed by leaf-cutter ants and how their cultivar functions to obtain, modify and deliver plant

lipids, further sheds light on this complex mutualistic relationship between these organisms. In a

broader context, it provides an important example of how microbes mediate the relationship

between herbivores and the plants that they consume, either directly, with microbes interfacing

with their hosts in the gut, or less directly, with the fungal garden serving as the ants' external

gut. Future lipidomic studies may help us to further understand how these lipids are used by the

ants in communication and symbiont maintenance, if and how they are incorporated into the

ants' bodies, and if specific lipids play a role as allelochemicals throughout the Attina and other

fungus farming insects, or if this phenomenon is limited to the leaf-cutter ants and their cultivar

L. gongylophorus.

Materials and Methods

Material collection

Six lab-maintained *Atta* leaf-cutter ant colonies were provided exclusively maple leaves

for the two weeks before sample collection. We collected leaf material as well as fungal garden

from the top, middle, and bottom of leaf-cutter ant fungal garden ecosystems. These layers are

differentiated based on color, texture and location in the garden (**Figure 1**). All of the six fungus

gardens that were sampled were maintained in the same size of container, 8 x 8 x 8 cm and so the

fungus garden that was sampled was from approximately the same distance from the edge for

each sample type.

Gongylidia are difficult to collect because of their small size and their sparseness in the

garden. To collect gongylidia, we first collected fungal garden in deep petri dishes and reduced

the number of workers. With fewer workers in the fungal garden, the gongylidia proliferate.

After several days, the gongylidia were picked off by needlepoint under a dissecting microscope

and they were placed in a microcentrifuge tube with water.

The leaf material used in this experiment was collected in the summer and kept frozen at

-20°C in vacuum sealed bags so that the ant colonies can be fed during the winter when fresh

leaves are not available.

Total lipid extraction

The collected material was lyophilized for untargeted lipidomics analysis. Lipids were

extracted using a modified Folch extraction (40). To break up the biological material,

approximately 25 mg of sample was bead beaten using a 3 mm tungsten carbide bead in 750 µl

of methanol for 2 min at a frequency of 30 Hz. The sample was then removed and transferred

into a 20 mL clean EPS glass vial with a Teflon lined cap. Another 750 µl of methanol was

added for a final volume of 1.5 mL methanol. Next, 3 mL of chloroform and 200 µl of water

were added to each sample. The samples were vortexed for 30 s, sonicated for 30 min, vortexed

again for 30 s, and then 0.925 µl of water was added to induce a phase separation. The samples

were incubated at 4ºC overnight and then the lower lipid layer was removed, dried down, and

stored at -20ºC at a concentration of 2 µg/µl in 2:1 chloroform/methanol until Mass

Spectrometry (MS) analysis.

Liquid chromatography (LC)-tandem mass spectrometry (MS/MS) and LC-ion mobility

spectrometry-mass spectrometry (LC-IMS-MS) analyses

All extracted lipids in this manuscript were analyzed by LC-MS/MS using a Waters

NanoAquity UPLC system interfaced with a Velos Orbitrap mass spectrometer (Thermo

Scientific, San Jose, CA) and an Agilent 6560 Ion Mobility QTOF MS system (Agilent, Santa

Clara, CA). The total lipid extracts (TLE) were reconstituted in methanol for a final abundance

of 0.4 µg TLE/µl. Seven µl of the TLE were injected onto Waters column (HSS T3 1.0 mm x

150 mm x 1.8 µm particle size). Lipids were separated over a 90 min gradient elution (mobile

phase A: ACN/H_2O (40:60) containing 10 mM ammonium acetate; mobile phase B: ACN/IPA

(10:90) containing 10 mM ammonium acetate) at a flow rate of 30 µl/min. Samples were

analyzed in both positive and negative ionization using HCD (higher-energy collision

dissociation) and CID (collision-induced dissociation) to obtain high coverage of the lipidome.

Velos Orbitrap MS abundance data is depicted in **Figure 2**, **Figure 3**, Data file S2, and Data file

S3. Figures and files apply LIPID MAPS Classification, Nomenclature and Shorthand Notation

for MS-derived Lipid Structures (41).

The LC-IMS-MS analyses were also performed in both positive and negative ion mode

and collected from 100-3200 m/z at a MS resolution of 40,000. The LC-IMS-MS data were

analyzed using in-house PNNL software for deisotoping and feature finding the

multidimensional LC, IMS and MS data (42).

Lipid targeted database and alignment

We used LIQUID software (43) for lipid targeted database alignment. This is achieved by

aligning all datasets (grouped by sample type and ionization mode) and matching unidentified

features to their identified counterparts using MZmine2 (44). Aligned features are manually

verified and peak apex intensity values are exported for statistical analysis.

Behavior experiment

An ant behavior experiment was conducted in order to determine if *A. cephalotes* ants

could detect various lipids found in the fungal garden, and to observe their corresponding

behavioral response to them. We tested the ants' response to linoleic acid (abundant in the

gongylidia, 18:2 fatty acids), alpha-linolenic acid (abundant in the leaves and top of garden, 18:3

fatty acids), and oleic acid (excreted by dead ants, 18:1 fatty acids). To test the response of the *A.*

cephalotes to these lipids, we used a method similar to those from López-Riquelme *et al.*(32).

All three of the lipids were dissolved in ethanol at 1 mg/mL concentration. We pipetted 5 μL of

the mixture onto 6 mm paper discs (Whatman 2017-006), that had been pre-labeled with pencil,

then waited several seconds for the ethanol to evaporate before introducing the discs to the *A.*

cephalotes colonies. Discs with only ethanol were used as a negative control. Each treatment

replicate was set up as a binary choice between the treatment oil, and the negative control. We

used seven *A. cephalotes* queenright colonies, which were all collected from La Selva Biological

Research Station in Costa Rica in April 2018 and were maintained at the University of

Wisconsin-Madison. The colonies were housed in 42.5 cm x 30.2 cm x 17.8 cm plastic

containers. Three treatment discs and three control discs were used for each assay and were

placed on a flat surface in the colony box. The ants were observed for five minutes from the time

the discs were placed in the colony. Behavior was coded using the BORIS program (45). We

recorded the number of times discs were picked up, characterized by ants using their mandibles

to lift the disc in the same manner that they lift leaf pieces when foraging. We recorded the

number and duration of ant inspections, characterized by ants approaching the discs while

waving antennae toward the discs, touching the disks with their antennae, or walking over the

discs. Finally, we recorded the number and duration of aggressive behaviors toward the discs,

characterized by lunging and biting, or moving slowly or stopping nearby with open mandibles

(46) (example behaviors can be seen in Movie S1 and Movie S2). It should be noted that if

several ants were simultaneously exhibiting the same behavior, this was not considered a

separate event. The event would be considered active as long as at least one ant was exhibiting

the behavior. We tested all three treatments with each colony.

Statistical Analysis

Lipidomics Statistical Analysis: All lipidomics datasets were first subjected to quality

control processing, which included an initial outlier identification step using the algorithm RMD-

PAV (47), which specifically performs a robust Principal Component Analysis (PCA) and

identifies outliers via distributional properties of the lipid measurements within each sample

(Average Pearson correlation within sample type group, median absolute deviation, skew and

kurtosis). From the robust PCA a robust Mahalanobis distance is computed and the statistical

score associated with each sample is computed from a chi-square distribution with 3 degrees of

freedom. Using RMD-PAV we found no outliers. Additionally, the data had very few missing

values and thus no lipid identifications were removed either as part of the quality control

processing. Quality control processing also evaluated if there was a garden effect, meaning that

the samples collected within a garden were more similar than samples collected from different

gardens. We computed a standard Pearson correlation values across all pairs of samples as our

metric of similarity and compared the within garden correlation values ($n1 = 18$) with the across

garden correlations ($n2 = 135$) using a two-sample t-test and did not observe a difference based

on garden (*p~0.894*); Pearson correlation of 0.879 and 0.876 for within and across samples,

respectively. The requirements of a two-sample t-test were tested for normality with a

Kolmogorov-Smirnov test (*p~0.587*) and equal variance with a two-sample F-test for equal

variances (*p~0.125*). This is confirmed in **Figure 2** with no obvious clustering outside sample

levels by garden.

A standard Analysis of Variance (ANOVA) was used to evaluate each lipid for a

statistical difference using the factor level (top, middle and bottom region of the garden) for the

first comparison with a Tukey's post-hoc test to compare the individual levels to one another

where each of the three levels contained 6 replicates. These 6 biological replicates provided

parallel measurements across 6 distinct fungal garden samples. **Figure 2** depicts the Tukey's

post-hoc garden top vs garden bottom test significant results at a p-value threshold of 0.05. To

test for differences between the middle and gongylidia for the second comparison (**Figure 3**) a

two-sample t-test was employed, again each factor contained 6 biological replicates.

For both the first comparison of the three (top, middle, and bottom) regions and the

second comparison of two (middle and gongylidia) regions the normality assumption was tested

for all lipids using a Kolmogorov-Smirnov test. For the first test ~3.6% of the lipids rejected the

null hypothesis that the data followed a normal distribution at a p-value threshold of 0.05 and

none reject the hypothesis for the second comparison. The 3.6% is within the expected

percentage to fail by random chance and none of these 3.6% lipids are significant after a

Bonferroni correction. For the first comparison of the three regions a Bartlett test of equal

variance was utilized, of where 16.2% failed the null hypothesis at a p-value threshold of 0.05.

450 However, none of these 16.2% lipids remained significant after a Bonferroni correction. A

451 similar result was found for the second dataset using a F-test for equal variances, 23.9%, of

452 which none remained after multiple test correction. These results did not warrant using non-

453 parametric statistics for the analysis. We did perform multiple corrections (Data file S3) with a

454 Bonferroni correction, which reduced the number of significant lipids by about three-quarters.

455 However, since we are interested in evaluating trends across lipid species due to class and fatty

456 acid composition, **Figures 2** and **3**, use a standard p-value threshold of 0.05. **Figure 3** lipids

457 important to our behavior study had Bonferroni corrected q-values below a threshold of 0.05

458 (i.e., lipids enriched in the gongylidia (TG(18:2/18:2/26:0), q-value 0.010; PE(18:2/18:2)_A, q-

459 value 0.003; PE(18:2/0:0)_B, q-value 0.027) and lipids enriched in the middle of the fungus

460 garden (TG(18:3/18:3/22:0), q-value 0.005; PI(18:3/18:3), q-value 0.032) (see Data file S3 for all

461 values).

462   <u>Statistical data included in **Figure 2** and **Figure 3**</u>: We evaluated if multivariate

463 differences in the lipid profiles between garden components were observable by unsupervised

464 learning, PCA. We conducted one PCA to compare the layers of the garden (**Fig. 2**) and another

465 for the comparison between gongylidia and the middle of the garden (**Fig. 3**). In both cases, we

466 used the function prcomp in the stats package of R with the scale option. To determine if the

467 distance between groups was significantly different than random, we used ANOSIM and

468 PERMANOVA, both in R (48). **Figure 2** contains the $\log_2$ intensity values for lipids

469 significantly changing (p-value < 0.05) between the top and bottom regions of the garden where

470 green denotes lipids relatively increased in the top of the garden and orange font denotes lipids

471 relatively increased in the bottom of the garden. **Figure 3** contains the $\log_2$ intensity values for

472 lipids significantly changing (p-value < 0.05) between the gongylidia and the middle region of

473 the fungus garden from where they were harvested where teal denotes lipids relatively increased

in the garden and purple font denotes lipids relatively increased in the gongylidia. Individual p-

values for each lipid can be found in Data File S2 and Data File S3.

<u>Behavior experiment Statistical Analysis</u>: For the behavior experiment, we did not

compare the various treatments to each other, but only to the control. Despite being the same age

and size, the colonies differed in terms of activity levels and number of ants, so we corrected for

this by comparing the response to the treatment disc to the control. For the duration

measurements we evaluated the difference in time between treatment and control, and for the

number of pickups and aggression events we used the proportion of times the event happened

between treatment discs and control discs. For the duration of time we conducted a one sample t-

test, where a mean different from zero indicated a significant difference between the treatment

and the control, and for the pickup and aggression events we performed a test of proportions,

where a proportion different from 0.5 indicated a significant difference (48).

486
487   **References**

488   1.   Merrill AL, Watt BK. 1973. Energy Value Of Foods - Basis and Derivation.
489   2.   Yeagle PL. 1989. Lipid regulation of cell membrane structure and function. FASEB J
490   3:1833-42.
491   3.   Waters CM, Bassler BL. 2005. Quorum sensing: Cell-to-cell communication in bacteria.
492   Annual Review of Cell and Developmental Biology 21:319-346.
493   4.   Lebeer S, Vanderleyden J, De Keersmaecker SCJ. 2010. Host interactions of probiotic
494   bacterial surface molecules: comparison with commensals and pathogens. Nature
495   Reviews Microbiology 8:171-184.
496   5.   Selander E, Kubanek J, Hamberg M, Andersson MX, Cervin G, Pavia H. 2015. Predator
497   lipids induce paralytic shellfish toxins in bloom-forming algae. PNAS112:6395-6400.
498   6.   Kearns DB, Shimkets LJ. 2001. Lipid chemotaxis and signal transduction in *Myxococcus*
499   *xanthus*. Trends in Microbiology 9:126-129.
500   7.   Ketter E, Randall G. 2019. Virus Impact on Lipids and Membranes. Annual Review of
501   Virology 6:319-340.
502   8.   Magalhaes DM, Da Silva ITFA, Borges M, Laumann RA, Blassioli-Moraes MC. 2019.
503   Anthonomus grandis aggregation pheromone induces cotton indirect defence and attracts
504   the parasitic wasp *Bracon vulgaris*. Journal of Experimental Botany 70:1891-1901.
505   9.   Costa AN, Vasconcelos HL, Vieira☐Neto EHM, Bruna EM. 2008. Do herbivores exert
506   top☐down effects in Neotropical savannas? Estimates of biomass consumption by
507   leaf☐cutter ants. Journal of Vegetation Science 19:849-854.
508   10.   Hölldobler B, Wilson EO. 2009. The superorganism: the beauty, elegance, and
509   strangeness of insect societies. WW Norton & Company, New York, NY.

11. Meyer ST, Leal IR, Tabarelli M, Wirth R. 2011. Ecosystem engineering by leaf-cutting
ants: nests of *Atta cephalotes* drastically alter forest structure and microclimate.
Ecological Entomology 36:14-24.
12. Holldobler B, Wilson EO. 2010. The Leafcutter Ants: Civilization by Instinct. W.W.
Norton and Company.
13. Aylward FO, Burnum-Johnson KE, Tringe SG, Teiling C, Tremmel DM, Moeller JA,
Scott JJ, Barry KW, Piehowski PD, Nicora CD, Malfatti SA, Monroe ME, Purvine SO,
Goodwin LA, Smith RD, Weinstock GM, Gerardo NM, Suen G, Lipton MS, Currie CR.
2013. *Leucoagaricus gongylophorus* produces diverse enzymes for the degradation of
recalcitrant plant polymers in leaf-cutter ant fungus gardens. Appl Environ Microbiol
79:3770-8.
14. Khadempour L, Burnum-Johnson KE, Baker ES, Nicora CD, Webb-Robertson BM,
White RA, 3rd, Monroe ME, Huang EL, Smith RD, Currie CR. 2016. The fungal cultivar
of leaf-cutter ants produces specific enzymes in response to different plant substrates.
Mol Ecol 25:5795-5805.
15. Suen G, Teiling C, Li L, Holt C, Abouheif E, Bornberg-Bauer E, Bouffard P, Caldera EJ,
Cash E, Cavanaugh A, Denas O, Elhaik E, Fave MJ, Gadau J, Gibson JD, Graur D,
Grubbs KJ, Hagen DE, Harkins TT, Helmkampf M, Hu H, Johnson BR, Kim J, Marsh
SE, Moeller JA, Munoz-Torres MC, Murphy MC, Naughton MC, Nigam S, Overson R,
Rajakumar R, Reese JT, Scott JJ, Smith CR, Tao S, Tsutsui ND, Viljakainen L, Wissler
L, Yandell MD, Zimmer F, Taylor J, Slater SC, Clifton SW, Warren WC, Elsik CG,
Smith CD, Weinstock GM, Gerardo NM, Currie CR. 2011. The genome sequence of the
leaf-cutter ant *Atta cephalotes* reveals insights into its obligate symbiotic lifestyle. PLoS
Genet 7:e1002007.
16. Nagamoto NS, Garcia MG, Forti LC, Verza SS, Noronha NC, Rodella RA. 2011.
Microscopic evidence supports the hypothesis of high cellulose degradation capacity by
the symbiotic fungus of leaf-cutting ants. Journal of Biological Research 16:308-312.
17. Suen G, Scott JJ, Aylward FO, Currie CR. 2011. The Microbiome of Leaf-Cutter Ant
Fungus Gardens, p 367-379. *In* de Bruijn FJ (ed), Handbook of Molecular Microbial
Ecology II doi:10.1002/9781118010549.ch36. John Wiley & Sons, Inc., Hoboken, NJ,
USA.
18. Moreira-Soto RD, Sanchez E, Currie CR, Pinto-Tomás AA. 2017. Ultrastructural and
microbial analyses of cellulose degradation in leaf-cutter ant colonies. Microbiology
163:1578-1589.
19. De Fine Licht HH, Schiott M, Rogowska-Wrzesinska A, Nygaard S, Roepstorff P,
Boomsma JJ. 2013. Laccase detoxification mediates the nutritional alliance between leaf-
cutting ants and fungus-garden symbionts. PNAS 110:583-7.
20. Aylward FO, Khadempour L, Tremmel DM, McDonald BR, Nicora CD, Wu S, Moore
RJ, Orton DJ, Monroe ME, Piehowski PD, Purvine SO, Smith RD, Lipton MS, Burnum-
Johnson KE, Currie CR. 2015. Enrichment and Broad Representation of Plant Biomass-
Degrading Enzymes in the Specialized Hyphal Swellings of *Leucoagaricus*
gongylophorus, the Fungal Symbiont of Leaf-Cutter Ants. PLoS One 10:e0134752.
21. Farji-Brener AG, Medina CA. 2000. The Importance of Where to Dump the Refuse: Seed
Banks and Fine Roots in Nests of the Leaf-Cutting Ants *Atta cephalotes* and *A.*
colombica. Biotropica 32:120-126.
22. Bass M, Cherrett JM. 1995. Fungal Hyphae as a Source of Nutrients for the Leaf-Cutting
Ant *Atta sexdens*. Physiological Entomology 20:1-6.

23.   Shik JZ, Rytter W, Arnan X, Michelsen A. 2018. Disentangling nutritional pathways
linking leafcutter ants and their co-evolved fungal symbionts using stable isotopes.
Ecology 99:1999-2009.
24.   Aylward FO, Burnum KE, Scott JJ, Suen G, Tringe SG, Adams SM, Barry KW, Nicora
CD, Piehowski PD, Purvine SO, Starrett GJ, Goodwin LA, Smith RD, Lipton MS, Currie
CR. 2012. Metagenomic and metaproteomic insights into bacterial communities in leaf-
cutter ant fungus gardens. ISME Journal 6:1688-1701.
25.   Kyle JE, Zhang X, Weitz KK, Monroe ME, Ibrahim YM, Moore RJ, Cha J, Sun X,
Lovelace ES, Wagoner J, Polyak SJ, Metz TO, Dey SK, Smith RD, Burnum-Johnson KE,
Baker ES. 2016. Uncovering biologically significant lipid isomers with liquid
chromatography, ion mobility spectrometry and mass spectrometry. Analyst 141:1649-
59.
26.   Schultz TR, Brady SG. 2008. Major evolutionary transitions in ant agriculture. PNAS
105:5435-40.
27.   Nygaard S, Hu H, Li C, Schiøtt M, Chen Z, Yang Z, Xie Q, Ma C, Deng Y, Dikow RB,
Rabeling C, Nash DR, Wcislo WT, Brady SG, Schultz TR, Zhang G, Boomsma JJ. 2016.
Reciprocal genomic evolution in the ant–fungus agricultural symbiosis. Nature
Communications 7:12233.
28.   Kobayashi K. 2016. Role of membrane glycerolipids in photosynthesis, thylakoid
biogenesis and chloroplast development. J Plant Res 129:565-580.
29.   Conconi A, Miquel M, Browse JA, Ryan CA. 1996. Intracellular Levels of Free
Linolenic and Linoleic Acids Increase in Tomato Leaves in Response to Wounding. Plant
Physiol 111:797-803.
30.   Gordon DM. 1983. Dependence of necrophoric response to oleic acid on social context in
the ant, *Pogonomyrmex badius*. J Chem Ecol 9:105-11.
31.   Wilson EO, Durlach NI, Roth LM. 1958. Chemical Releaser of Necrophoric Behavior in
Ants. Psyche: A Journal of Entomology 65:108-114.
32.   Lopez-Riquelme GO, Malo EA, Cruz-Lopez L, Fanjul-Moles ML. 2006. Antennal
olfactory sensitivity in response to task-related odours of three castes of the ant *Atta
mexicana* (Hymenoptera : Formicidae). Physiological Entomology 31:353-360.
33.   Buehlmann C, Graham P, Hansson BS, Knaden M. 2014. Desert ants locate food by
combining high sensitivity to food odors with extensive crosswind runs. Curr Biol
24:960-4.
34.   De Fine Licht HH, Boomsma JJ, Tunlid A. 2014. Symbiotic adaptations in the fungal
cultivar of leaf-cutting ants. Nat Commun 5:5675.
35.   Kecskeméti S, Szelényi MO, Erdei AL, Geösel A, Fail J, Molnár BP. 2020. Fungal
Volatiles as Olfactory Cues for Female Fungus Gnat, *Lycoriella ingenua* in the
Avoidance of Mycelia Colonized Compost. Journal of Chemical Ecology 46:917-926.
36.   Holighaus G, Rohlfs M. 2016. Fungal allelochemicals in insect pest management.
Applied Microbiology and Biotechnology 100:5681-5689.
37.   North RD, Jackson CW, Howse PE. 1999. Communication between the fungus garden
and workers of the leaf-cutting ant, *Atta sexdens rubropilosa*, regarding choice of
substrate for the fungus. Physiological Entomology 24:127-133.
38.   Sousa KKA, Catalani GC, Gianeti TMR, Camargo RS, Caldato N, Ramos VM, Forti LC.
2020. A Volatile Semiochemical Released by the Fungus Garden of Leaf-Cutting Ants.
Florida Entomologist 103:1-8, 8.
39.   Green PWC, Kooij PW. 2018. The role of chemical signalling in maintenance of the
fungus garden by leaf-cutting ants. Chemoecology 28:101-107.

40. Folch J, Lees M, Sloane Stanley GH. 1957. A simple method for the isolation and
purification of total lipides from animal tissues. J Biol Chem 226:497-509.
41. Liebisch G, Fahy E, Aoki J, Dennis EA, Durand T, Ejsing C, Fedorova M, Feussner I,
Griffiths WJ, Koefeler H, Merrill AH, Jr., Murphy RC, O'Donnell VB, Oskolkova OV,
Subramaniam S, Wakelam M, Spener F. 2020. Update on LIPID MAPS Classification,
Nomenclature and Shorthand Notation for MS-derived Lipid Structures. J Lipid Res
doi:10.1194/jlr.S120001025.
42. Crowell KL, Slysz GW, Baker ES, LaMarche BL, Monroe ME, Ibrahim YM, Payne SH,
Anderson GA, Smith RD. 2013. LC-IMS-MS Feature Finder: detecting multidimensional
liquid chromatography, ion mobility and mass spectrometry features in complex datasets.
Bioinformatics 29:2804-5.
43. Kyle JE, Crowell KL, Casey CP, Fujimoto GM, Kim S, Dautel SE, Smith RD, Payne SH,
Metz TO. 2017. LIQUID: an-open source software for identifying lipids in LC-MS/MS-
based lipidomics data. Bioinformatics 33:1744-1746.
44. Pluskal T, Castillo S, Villar-Briones A, Oresic M. 2010. MZmine 2: modular framework
for processing, visualizing, and analyzing mass spectrometry-based molecular profile
data. BMC Bioinformatics 11:395.
45. Friard O, Gamba M. 2016. BORIS: a free, versatile open-source event-logging software
for video/audio coding and live observations. Methods in Ecology and Evolution 7:1325-
1330.
46. Hernandez JV, Lopez H, Jaffe K. 2002. Nestmate recognition signals of the leaf-cutting
ant *Atta laevigata*. Journal of Insect Physiology 48:287-295.
47. Matzke MM, Waters KM, Metz TO, Jacobs JM, Sims AC, Baric RS, Pounds JG, Webb-
Robertson BJ. 2011. Improved quality control processing of peptide-centric LC-MS
proteomics data. Bioinformatics 27:2866-72.
48. Team RC. R: A language and environment for statistical computing. R Foundation for
Statistical Computing. Available at: http://wwwR-projectorg/ (Accessed: 23rd December
2013).

Acknowledgments

This work was funded by the U.S. Department of Energy (DOE), Office of Science, Office of

Biological and Environmental Research (BER), Early Career Research Program to KEBJ; Great

Lakes Bioenergy Research Center (DOE BER DE-FC02-07ER64494); National Institute of Food

and Agriculture, United States Department of Agriculture, under ID number 1003779; and

National Institutes of Health (NIH grant U19TW009872). This research used capabilities

developed by NIH National Institute of Child Health and Human Development (R21

HD084788), NIH National Institute of Environmental Health Sciences (R01 ES022190) and the

Pan-omics program (funded by DOE BER Genome Sciences Program). This work was

performed in the W. R. Wiley Environmental Molecular Sciences Laboratory (EMSL)

(grid.436923.9), a DOE Office of Science User Facility sponsored by the Office of Biological

and Environmental Research and located at Pacific Northwest National Laboratory (PNNL).

PNNL is a multiprogram national laboratory operated by Battelle for the Department of Energy

(DOE) under Contract DE-AC05-76RLO 1830. The funders had no role in study design, data

collection and interpretation, or the decision to submit the work for publication

649
Author contributions: Conceived of the project (KEBJ, ESB, CC, LK), conducted experiments

(LK, CN, KEBJ, ESB) performed statistical analysis (BJMWR, LK), analyzed data (LK, KEBJ,

ESB, JEK, FBS) designed sampling methodology (LK, KEBJ, CC), provided resources and

facilities (KEBJ, ESB, CC, MSL, RDS) and wrote the manuscript (LK, KEBJ, ESB).

Competing interests: There are no competing interests.

Data and materials availability: All data are available in the Supplementary Materials. Raw

lipidomic datasets are deposited on Mass Spectrometry Interactive Virtual Environment

(MassIVE). After the manuscript is published and has a PubMed ID, the data will be made

public and available on MassIVE; [https://massive.ucsd.edu] accession number MSV000084520.

659
FIGURE LEGENDS

Figure 1: A schematic diagram of the process by which lipids are processed through the

fungal garden. Fresh leaf material is first deposited on the top of the fungal garden and passes

through the middle and bottom of the garden. As the leaf material proceeds, lipids are extracted

by the fungus and modified. The final lipid profile in the gongylidia (the specialized fungal

structure that ants consume) differs from that which is in the leaf material and top of the fungal

garden. The gongylidia are characterized by lipids containing linoleic acid (18:2), whereas the

leaves are characterized by lipids containing alpha-linolenic acid (18:3). Lipid structures here are

only representative and other isomers are possible. Photo credits: Lily Khadempour, except

gongylidia by Don Parsons.

Figure 2: Heatmaps representing the relative log$_2$ expression level for lipids in the leaf food

source and the top, middle, and bottom strata of six leaf-cutter ant fungal gardens. Each

column represents the relative abundance of a unique lipid and the heatmaps are scaled by

column. Lipids significantly increased in the top are denoted with green font; lipids significantly

increasing in the bottom are denoted with orange font (p-values <0.05 were deemed significant);

_A and _B denote structural isomers. Diacylglycerophosphocholines (PC),

monoacylglycerophosphocholines (LPC), sulfoquinovosyldiacylglycerol (SQDG),

Monogalactosyldiacylglycerol (MGDG), Cardiolipin (CL), phosphatidic acid (PA), Hexosyl-

Ceramide (HexCer), Ceramide (Cer), Diacylglycerols (DG), diacylglycerophosphoglycerol (PG),

monoacylglycerophosphoglycerol (LPG), diacylglycerophosphoserine (PS),

diacylglycerophosphoethanolamines (PE), monoacylglycerophosphoethanolamines (LPE),

diacylglycerophosphoinositols (PI). Lipid abbreviations show the total number of acyl chain

carbons: total number of double bonds. ‡, 18:2 containing lipids significantly increased in the

bottom of the garden; *, 18:3 containing lipids significantly increased in the top of the garden.

Figure 3: Heatmaps representing the relative log$_2$ expression level for lipids in the leaf food

source and the gongylidia and surrounding fungal garden of six leaf-cutter ant fungal

garden ecosystems. Each column represents the relative abundance of a unique lipid and the

heatmaps are scaled by column. Lipids significantly enriched in the gongylidia compared to the

surrounding fungal garden are denoted with purple font; lipids significantly enriched in the

fungal garden are denoted with teal font (p-values <0.05 were deemed significant); _A and _B

denote structural isomers. Digalactosyldiacylglycerol (DGDG), sulfoquinovosyldiacylglycerol

(SQDG), Monogalactosyldiacylglycerol (MGDG), diacylglycerophosphoglycerol (PG),

monoacylglycerophosphoglycerol (LPG), Diacylglycerophosphocholines (PC),

monoacylglycerophosphocholines (LPC), mannosylinositol phosphorylceramide (MIPC),

Ceramide phosphoinositol (PI-Cer), diacylglycerophosphoinositols (PI), phosphatidic acid (PA),

lysophosphatidic acid (LPA), Hexosyl-Ceramide (HexCer), Ceramide (Cer),

diacylglycerophosphoethanolamines (PE), monoacylglycerophosphoethanolamines (LPE),

Triacylglycerols (TG), Diacylglycerols (DG). Lipid abbreviations show the total number of acyl

chain carbons: total number of double bonds. *, 18:3 containing lipids significantly decreased in

the gongylidia; ‡, 18:2 containing lipids significantly increased in the gongylidia.

Figure 4: **Results of the behavior experiment**. *Atta cephalotes* ants are attracted to linoleic acid

(18:2) and oleic acid (18:1) but show aggression toward alpha-linolenic acid (18:3). Each point

represents one experimental replicate, where the values for the control count or duration are

subtracted from the treatment count or duration. A one sample t-test was conducted for the

difference in duration and a one sample test of proportion was conducted for the number of

pickups and aggressions. Those whose mean is significantly different from zero are marked with

* $p < 0.05$ and ** $p < 0.01$. Box plot definitions: center line – median, upper and lower box

limits – upper and lower quartiles, whiskers – 1.5 x inter-quartile range, outliers – any points

outside the 1.5x inter-quartile range.

SUPPLEMENTARY MATERIALS LEGENDS

Figure S1: Heatmaps representing the relative $\log_2$ expression level for lipids in the leaf

food source and the top, middle, and bottom strata of six leaf-cutter ant fungal gardens.

Each column represents the relative abundance of a unique lipid and the heatmaps are scaled by

column. Lipids significantly increased in the top are denoted with green font; lipids significantly

increasing in the bottom are denoted with orange font (p-values <0.05 were deemed significant);

lipids not significantly changing across the top to the bottom of the garden are denoted with grey

font; _A and _B denote structural isomers. Diacylglycerophosphocholines (PC),

monoacylglycerophosphocholines (LPC), sulfoquinovosyldiacylglycerol (SQDG),

Monogalactosyldiacylglycerol (MGDG), Cardiolipin (CL), phosphatidic acid (PA), Hexosyl-

Ceramide (HexCer), Ceramide (Cer), Diacylglycerols (DG), diacylglycerophosphoglycerol (PG),

monoacylglycerophosphoglycerol (LPG), diacylglycerophosphoserine (PS),

diacylglycerophosphoethanolamines (PE), monoacylglycerophosphoethanolamines (LPE),

diacylglycerophosphoinositols (PI). Lipid abbreviations show the total number of acyl chain

carbons: total number of double bonds. ‡, 18:2 containing lipids significantly increased in the

bottom of the garden; *, 18:3 containing lipids significantly increased in the top of the garden.

Figure S2: Heatmaps representing the relative log$_2$ expression level for lipids in the leaf

food source and the gongylidia and surrounding fungal garden of six leaf-cutter ant fungal

garden ecosystems. Each column represents the relative abundance of a unique lipid and the

heatmaps are scaled by column. Lipids significantly enriched in the gongylidia compared to the

surrounding fungal garden are denoted with purple font; lipids significantly enriched in the

fungal garden are denoted with teal font (p-values <0.05 were deemed significant); lipids not

significantly changing between the gongylidia and the surrounding fungal garden are denoted

with grey font; _A and _B denote structural isomers. Digalactosyldiacylglycerol (DGDG),

sulfoquinovosyldiacylglycerol (SQDG), Monogalactosyldiacylglycerol (MGDG),

diacylglycerophosphoglycerol (PG), monoacylglycerophosphoglycerol (LPG),

Diacylglycerophosphocholines (PC), monoacylglycerophosphocholines (LPC), mannosylinositol

phosphorylceramide (MIPC), Ceramide phosphoinositol (PI-Cer), diacylglycerophosphoinositols

(PI), phosphatidic acid (PA), lysophosphatidic acid (LPA), Hexosyl-Ceramide (HexCer),

Ceramide (Cer), diacylglycerophosphoethanolamines (PE),

monoacylglycerophosphoethanolamines (LPE), Triacylglycerols (TG), Diacylglycerols (DG).

Lipid abbreviations show the total number of acyl chain carbons: total number of double bonds.

*, 18:3 containing lipids significantly decreased in the gongylidia; ‡, 18:2 containing lipids

significantly increased in the gongylidia.

Figure S3: *Atta* **ant behavior in response to alpha-linolenic acid (18:3) in comparison to**

control. Labels above panels represent the different colonies that were tested. Tracks represent

the times of event occurrences and their durations.

Figure S4: *Atta* **ant behavior in response to linoleic acid (18:2) in comparison to control**.

Labels above panels represent the different colonies that were tested. Tracks represent the times

of event occurrences and their durations.

Figure S5: *Atta* **ant behavior in response to oleic acid (18:1) in comparison to control**.

Labels above panels represent the different colonies that were tested. Tracks represent the times

of event occurrences and their durations.

Data file S1: Proteomic identification of lipid associated proteins. Quantities of protein

spectra related to lipid metabolism identified throughout the fungus garden, from the dataset

produced for Aylward et al. 2015.

Data file S2: Lipid comparisons of top, middle, and bottom fungal garden regions.

Lipidomic intensity data and statistical analysis results for comparisons across the top, middle,

and bottom fungal garden regions.

Data file S3: Lipid changes between fungal garden and gongylidia. Lipidomic intensity data

and statistical analysis results for comparisons across the middle fungal garden and gongylidia.

Movie S1: Aggressive behaviors. Discs labeled D are the negative control infused with only

ethanol. Discs labeled A are infused with ethanol and alpha-linolenic acid. In the first 34 s, ants

can be seen near or on both types of discs with their mandibles open, moving very slowly or

standing still. In the second part of the movie, an ant can be seen lunging and biting at the alpha-

linolenic acid infused discs. Movie can be viewed at https://youtu.be/fq6SVF-8v6M.

Movie S2: Interest/attractive behavior. Discs labeled D are the negative control infused with

only ethanol. Discs labeled B are infused with ethanol and linoleic acid. In the first 12 s of the

movie, ants can be seen approaching the linoleic acid discs, and waving their antennae. In the

second part of the movie, ants are inspecting the linoleic acid discs and one ant picks up a disc at

30 s. Movie can be viewed at https://youtu.be/unUL-ets3Vk.

Leaf

Top

Middle

Bottom

Lipids prevalent in leaf and top of fungus garden

SQDG(18:3_18:3)

DGDG(18:3_20:3)

PG(16:1_18:3)

MGDG(18:3_18:3)

Gongylidia

Ant

Lipids prevalent gongylidia

PC(18:2_22:0)

PI (18:2_18:2)

PE(16:1_18:2)

PC(16:1_18:2)

TG(16:0_16:1_18:2)

PE(18:2_18:2)

TG(18:2_18:2_26:0)