9 **ExBoX: a simple Boolean exclusion strategy to drive expression in**
10 **neurons**
16 Running title: Neuronal expression by Boolean Exclusion
17
18
19
20

Tyler Vahedi-Hunter*[1], Teresa Ubina*[1], Akshay Gupta[1], Wenny Wong[1], Will Agnew-
Svoboda[1], Vijayalakshmi Santhakumar[1], Martin M. Riccomagno[1]

[1] Neuroscience Graduate Program, Department of Molecular, Cell & Systems Biology,
University of California, Riverside, CA, USA

*Authors contributed equally to this work

Corresponding Author: Martin M. Riccomagno, martinmr@ucr.edu.

Keywords: Boolean logic, recombinases, combinatorial expression, intersectional

approaches, Cre, Flp, Adeno-Associated Virus.

Abstract

The advent of modern single-cell biology has greatly enhanced our

understanding of the striking diversity within cell populations that previously appeared

homogenous. This newly appreciated complexity has made intersectional genetic

approaches essential to understanding and probing cellular heterogeneity at the

functional level. Here we build on previous knowledge to develop a simple AAV-based

approach to define specific subpopulations of cells by Boolean exclusion logic (AND

NOT). This Expression by Boolean Exclusion (ExBoX) system encodes for a gene of

interest which is turned on by a particular recombinase (Cre or FlpO) and turned off by

another. ExBoX allows for the specific transcription of a gene of interest in cells

expressing only the activating recombinase, but not in cells expressing both. We show

the ability of the ExBoX system to tightly regulate expression of fluorescent reporters

both in vitro and in vivo, and further demonstrate the powerful adaptability of the system

by achieving expression of variety of virally-delivered coding sequences in the mouse

brain. This simple strategy will expand the molecular toolkit available for cell- and time-

specific gene expression in a variety of organisms and systems.

Abbreviations:

Hemagglutinin (HA)

Adeno-Associated Virus (AAV)

Double-floxed Inverse Orientation (DIO)

Dentate Gyrus (DG)

Gene Of Interest (GOI)

Introduction

Advancements in our understanding of the mechanisms underlying biological

processes have been greatly dependent on the development of new genetic tools. A big

breakthrough in mammalian genetics was the discovery and implementation of

homologous recombination to generate loss-of-function alleles in mice (Capecchi,

1989). The introduction of recombinases as genetically encoded tools, in combination

with conditionally targeted genetic alleles, made the control of genetic deletion in a

tissue-specific manner possible (Gu et al, 1994; Tsien et al, 1996). With the availability

of recombinase-dependent systems, region-specific gene knockouts and progenitor

tracing have now become routine experimental strategies in mouse genetics (Branda &

Dymecky, 2004).

Although cell diversity has always been appreciated in biology, single-cell

profiling has revolutionized the way we think about certain organs by uncovering cell

heterogeneity in populations that were once thought to display less complexity

(Darmanis et al, 2015). Our ability to appreciate cellular complexity is currently limited

by the available tools to label and manipulate cells with increased specificity. One way

to overcome this limitation is to simply expand the pool of tissue- and population-

specific recombinase lines. This requires the generation and characterization of

independent transgenic recombinase mouse lines for each newly discovered cell

population. A complementary approach that can take advantage of preexisting

transgenic lines is to incorporate genetically encoded intersectional approaches

(Awatramani et al, 2003). In comparison to conventional transgenic strategies that

select cellular targets based on expression of one particular promoter or driver,

intersectional approaches provide tighter specificity by selecting targets based on the

overlapping or sequential expression of multiple recombinases (Jensen & Dymecki,

2014). These intersectional strategies based on the combinatorial expression of two

recombinase systems (Cre/lox and Flp/FRT) were first used in the mouse to perform

fate mapping of previously elusive neural progenitors (Dymecki, Ray, & Kim 2010).

Intersectional approaches are not only able to label subpopulations with greater

specificity, but also make a variety of Boolean logic operations (AND, NOT, OR)

available (Daigle et al., 2019; Plummer et al, 2015). With multiple recombinase systems

driving the expression of reporter genes, targeting of subpopulations of cells can be

achieved through intersectional (expressing all drivers) and subtractive (lacking

expression of one or more drivers) strategies (Farago et al, 2006).

Viral vectors offer an alternative approach to driving conditional and

intersectional expression (Schnütgen et al, 2003; Atasoy et al, 2008; Gradinaru 2010;

Fenno et al, 2014). Strictly genetically encoded strategies have clear advantages over

surgically introduced viral vectors, like non-invasiveness and broad applicability in hard

to reach tissues during otherwise inaccessible embryonic stages (Jensen & Dymecki,

2014). However, viral expression systems can be extremely powerful given the ease of

generation, and the additional level of control one can gain by selecting the region and

time of injection (Zhang et al, 2007; Sternson et al, 2015). There are now multiple

versions of recombinase-dependent AAVs for recombinases like Cre, Flp, and DRE

(Atasoy et al, 2008; Saunders et al, 2012; Xue, Atallah, & Scanziani, 2014). More

recently, systems for intersectional expression driven by AAVs have also been

developed (Kakava-Georgiadou et al, 2019; Fenno et al, 2014). Combination of these

viral and genetically-encoded approaches will ultimately be essential to understand

whether newly discovered molecular diversity within cell populations amounts to any

notable phenotypic difference in terms of cellular structure and function.

In addition to being able to parse out functional diversity, there is an interest in

the field of developmental biology in being able to specifically turn genes on and off

during defined developmental windows or critical periods (Wiesel & Hubel,1963,

Kozorovitskiy et al, 2012). Intersectional approaches using Boolean negation/exclusion

could provide a solution to this problem: a gene of interest could be turned on by a

particular recombinase at a particular time point, while turned off later (AND NOT) by a

different recombinase (Table 1). Although a sophisticated system for AAV-driven

expression using exclusion logic has been generated elsewhere (Fenno et al, 2014), the

existing vectors have limitations in terms of being unnecessarily long and intricate in

design, somewhat restricting their applicability. In this study we describe the design and

characterization of an alternative expression system governed by Boolean exclusion

logic and driven by a single AAV. The newly developed vectors are simpler in design

and smaller in size, allowing for expression of longer genes of interest (GOI). As proof

of principle, we validated the system in neuronal populations, and also generated tools

for regulating neuronal activity in vivo.

Results

Design and construction of a combinatorial expression system using Boolean

exclusion: CreOn-FlpOff ExBox

To control expression and label subsets of cells with greater specificity using Cre

and FLP recombinases, we set out to develop a simple AAV-based expression system

governed by Boolean Exclusion logic (Expression by Boolean Exclusion or ExBoX). We

first generated a system in which expression of a coding sequence of interest is turned

ON by Cre recombinase and turned OFF by Flp recombinase: CreOn-FlpOff (Table 1).

The CreOn-FlpOff vector was designed to contain a cassette with a coding sequence

(CDS) of interest in-frame with a hemagglutinin tag (HA), P2A site, and EGFP reporter

inverted with respect to a promoter. The promoter chosen for these experiments was

that of human Synapsin I, which is a well characterized postmitotic neuronal promoter

(Glover et al., 2002). The inverted cassette is flanked by Frt sites, and double-lox sites

(lox2722 and loxP) in inverse orientation (DIO) (Schnütgen et al., 2003) (Figure 1A).

Upon Cre-mediated recombination, and HA-tagged coding sequence of interest and

EGFP get locked into the forward orientation, allowing for robust Cre-dependent

expression in neurons. Conversely, in the presence of Flp, the cassette is excised,

resulting in abolishment of CDS and EGFP reporter expression (Figure 1A). Thus, the

CDS is expressed only in the presence of Cre AND NOT FlpO (Table 1).

For proof of concept, we first generated a plasmid carrying only EGFP in the

reversible cassette of the CreOn-FlpOff system. The CreOn-FlpOff-EGFP plasmid was

co-transfected with Cre recombinase, FlpO recombinase, or both Cre and FlpO in a

neuroblastoma-derived cell line, Neuro2A (Figure 1B, left). A plasmid driving mCherry

under a constitutive promoter was co-transfected as transfection control. Upon

transfection of the CreOn-FlpOff-EGFP plasmid alone, no EGFP expression is found in

any cells. As expected, when CreOn-FlpOff-EGFP plasmid is co-transfected with Cre,

the EGFP cassette is inverted to the correct orientation and 93.9%±0.3 of transfected

cells express EGFP (Figure 1B). When CreOn-FlpOff-EGFP plasmid was co-transfected

with Flp, or co-transfected with Cre and Flp, no cells expressed EGFP, suggesting that

Flp can efficiently turn off reporter expression (Figure 1B). This confirms that expression

from CreOn-FlpOff ExBoX vectors can be successfully manipulated through Boolean

exclusion in N2A cells.

Design of a complementary system that is turned on by Flp and turned off by Cre:

FlpOn-CreOff ExBox

We next generated a complementary system that is turned on by Flp and off by

Cre, FlpOn-CreOff (Table 1). FlpOn-CreOff was designed using a similar internal logic,

where an expression cassette containing a CDS, an in-frame HA tag, P2A site, and

TdTomato reporter, are inverted with respect to the promoter. This expression cassette

is flanked by lox2722 sites, and a Flp controlled DIO switch (fDIO or fFLEX) (Xue et al,

2014) containing F14 and Frt sites (Figure 2A). In this system, Flp inverts the cassette,

resulting in CDS and TdTomato reporter expression, while Cre excises the cassette,

causing the abolishment of CDS and TdTomato reporter expression (Figure 2A). Thus

CDS expression should only occur when FlpO is present AND NOT Cre (Table 1).

As an initial proof-of-principle design, a FlpOn-CreOff vector driving simply a

TdTomato fluorescent reporter was generated. FlpOn-CreOff-TdTomato plasmid was

co-transfected with Cre, FlpO, or both Cre and FlpO. EGFP expressing plasmid was

used as transfection control. When the FlpOn-CreOff-TdTomato plasmid was co-

transfected with FlpO, approximately 85.8%±5.6 of transfected cells displayed

TdTomato expression (Figure 2B). When FlpOn-CreOff-TdTomato plasmid was co-

transfected with Cre alone, there were no TdTomato expressing cells. Finally, as

expected, co-transfection of the FlpOn-CreOff-TdTomato plasmid with both Cre and

FlpO resulted in no TdTomato reporter expression, confirming that Cre recombinase

can abolish expression of this constructs even in the presence of Flp (Figure 2B).

Therefore, expression of the construct occurs only in the presence of Flp AND NOT Cre

(Table 1).

Characterization of the ExBoX system in neurons

To test whether this newly developed system was able to drive expression in

neurons, we tested these constructs in primary cortical cultures. We used the same

plasmid combinations that were used in Neuro2A cells to perform ex-utero

electroporation prior to plating. Primary neuronal cultures were observed at 3 days in

vitro (DIV). Transfection of CreOn-FlpOff-EGFP plasmid alone resulted in no reporter

expression (Figure 3A). Notably, co-transfection of CreOn-FlpOff-EGFP plasmid with

Cre resulted in 100%±0.0 of transfected neurons expressing EGFP as expected (Figure

3A). When CreOn-FlpOff-EGFP plasmid was co-transfected with FlpO or with FlpO and

Cre, we found no EGFP reporter expression in any transfected cells (Figure 3A).

Similarly, transfection of FlpOn-CreOff-TdTomato plasmid alone resulted in no reporter

expression (Figure 3B). Co-transfection of FlpOn-CreOff-TdTomato plasmid with FlpO

resulted in 92.1%±2.0 of transfected cells expressing the TdTomato reporter, as

expected (Figure 3B). If FlpOn-CreOff-TdTomato plasmid is co-transfected with Cre

alone, or Cre and FlpO, it is expected that there will be no TdTomato expression since

Cre should excise the cassette. As predicted, no cells transfected in this combination

had any TdTomato expression (Figure 3B).

We then tested the neuronal expression of the ExBoX system *in vivo.* Initial

validation of CreOn-FlpOff was performed by *in utero* electroporation of CreOn-FlpOff-

EGFP plasmid with Cre, with FlpO, or with Cre and FlpO into the lateral ventricle in

E15.5 mice, followed by visualization 48 hours later. mCherry-expressing plasmids were

used as electroporation controls. When the CreOn-FlpOff-EGFP plasmid was

electroporated alone or with FlpO, we found no EGFP expression (Figure 4A). Tissue

co-transfected with CreOn-FlpOff-EGFP plasmid and Cre had robust expression of the

EGFP reporter, resulting in labeling of 97.7%±0.9 of transfected cells (Figure 4A). Co-

electroporation of CreOn-FlpOff-EGFP plasmid with Cre and FlpO resulted in negligible

EGFP expression (1.7%±0.9 of transfected cells, Figure 4A).

To validate FlpOn-CreOff, we electroporated the FlpOn-CreOff-TdTomato

plasmid alone and this resulted in no TdTomato expression. A plasmid driving EGFP

under the pCAG promoter was used as electroporation control. Co-electroporation of

FlpOn-CreOff-TdTomato plasmid with FlpO resulted in inversion of the cassette and

detectable expression of TdTomato in 86.9%±3.1 of transfected cells (Figure 4B). Co-

electroporation of FlpOn-CreOff-TdTomato plasmid with Cre showed no TdTomato

expression (Figure 4B). Finally, no TdTomato expression was detected when the

FlpOn-CreOff-TdTomato plasmid was co-electroporated with FlpO and Cre, indicating

successful AND NOT exclusion (Figure 4B). These data suggest that FlpOn-CreOff and

CreOn-FlpOff ExBoX work efficiently in neurons, *in vitro* and *in vivo*.

Validation of ExBoX AAVs *in vivo*

To generate tools that would be useful for manipulation of gene expression in the

postnatal brain, we packaged the FlpOn-CreOff construct into AAV9. To validate that

expression of the FlpOn-CreOff constructs is maintained in the virally-packaged form,

we performed stereotactic injections to deliver FlpOn-CreOff-TdTomato AAV into the

dentate gyrus (DG) of adult mice *in vivo*. FlpOn-CreOff-TdTomato AAV was co-injected

with either AAV-EF1α-FlpO-WPRE (AAV-EF1α-FlpO), or with AAV-EF1α-FlpO and

pENN.AAV.CamKII 0.4. Cre. SV40 (AAV-CamkII-Cre), and expression was observed

after 3 weeks (Figure 5). AAV expressing EGFP was co injected in WT and when Cre

was injected to visualize the injection site. Similar to our previous findings with FlpOn-

CreOff plasmids, robust TdTomato reporter expression was observed in DG neurons

co-infected with AAV-FlpOn-CreOff-TdTomato and AAV-EF1α-FlpO (Figure 5A, C),

verifying that expression of the construct in the presence of the 'ON' recombinase was

maintained in viral form. Furthermore, AAV-FlpOn-CreOff-TdTomato reporter

expression was inactivated by CamkII-Cre even in the presence of AAV-EF1α-FlpO

(Figure 5B, D; Figure S1). Thus, the FlpOn-CreOff construct works as expected when

packaged into AAV9.

We next packaged the complementary CreOn-FlpOff construct into AAV9. To

validate that expression of the CreOn-FlpOff construct is maintained in the virally-

packaged form, we performed stereotactic injections to deliver CreOn-FlpOff-EGFP

AAV into the dentate gyrus (DG) of adult mice *in vivo*. For validation for the viral CreOn-

FlpOff construct we incorporated a genetically-encoded Cre recombinase driven by a

transgene. The *Grik4-cre* mouse line was specifically chosen for providing robust Cre

expression in the hippocampus, including the DG (Supplemental Figure S2). AAV-

CreOn-FlpOff-EGFP was injected independently *Grik4-Cre* mice or Cre-negative

littermates (WT), or co-injected with AAV-EF1α-FlpO into *Grik4-Cre* animals (Figure 6).

AAV9 constitutively expressing TdTomato (pAAV-CAG-TdTomato) was co-injected in

WT animals and with the AAV-EF1α-FlpO injection to visualize the injection site. When

AAV-CreOn-FlpOff-EGFP was injected into WT (Cre-negative) animals, no EGFP

reporter expression was observed (Figure 6A, D, G). Conversely, injections into *Grik4-*

Cre positive animals demonstrated robust EGFP reporter expression (Figure 6B, E, H),

verifying the Cre-dependent activation of the virally-packaged construct. Furthermore, in

Grik4-Cre animals co-injection of AAV-EF1α-FlpO resulted in inactivation of EGFP

reporter expression (Figure 6C, F; Supplemental Figure S1). These results suggest that

the virally-packaged ExBoX constructs are successful at driving gene expression

through the expected AND NOT Boolean operation.

Generation of Activity-modulating CreOn-FlpOff Viruses for Neuroscience

To further validate the ExBoX system using multiple different cargos, we next

generated CreOn-FlpOff AAV constructs co-expressing EGFP and well-characterized

modulators of neuronal firing or synaptic release. To reduce activity, we generated a

CreOn-FlpOff virus that expresses Kir2.1 (AAV-CreOn-FlpOff-Kir2.1-EGFP), an inward-

rectifying K+ channel that has been demonstrated to increase rheobase current and

reduce resting membrane potential of neurons when overexpressed (Xue et al, 2014).

In parallel, we cloned a version of Kir2.1 carrying three point mutations that render the

channel in a non-functional state, referred to as Kir2.1Mut (AAV-CreOn-FlpOff-

Kir2.1Mut-EGFP) (Xue et al, 2014), which can be used as negative control for the Kir2.1

construct. We also designed a CreOn-FlpOff virus encoding mNaChBac (AAV-CreOn-

FlpOff-mNaChBac-EGFP). This bacterial Na^+ ion channel has been demonstrated to

increase firing of neurons when overexpressed by reducing action potential threshold

and rheobase (Xue et al, 2014). To serve as negative control for mNaChBac in neural

circuit experiments, we cloned a non-conducting mNaChBac mutant referred to as

mNaChBacMut into the CreOn-FlpOff vector (AAV-CreOn-FlpOff-mNaChBacMut-

EGFP) (Xue et al, 2014). Finally, we generated CreOn-FlpOff encoding tetanus toxin

(AAV-CreOn-FlpOff-TeTN-EGFP) to block synaptic transmission (Sweeney et al, 1995).

We validated reporter expression of these five viral constructs in adult mice using

stereotactic injection to deliver each independently or in combination with AAV-EF1α-

FlpO into the DG of adult Grik4-Cre mice or WT (Cre-negative) littermates (Figure 7). To

visualize the injection site pAAV-Syn-mCherry was co-injected in WT, or when AAV-

EF1α-FlpO was injected. As expected, injection of any of the viral constructs into Cre-

negative animals produced no EGFP reporter expression (Figure 7A, E, I, M, Q).

Conversely, each virus was able to produce robust EGFP reporter expression in DG

neurons when injected into Grik4-Cre positive animals (Figure 7B, F, J, N, R),

demonstrating expression of these viral constructs is Cre-dependent. Furthermore, in

Grik4-Cre mice injection of AAV-EF1α-FlpO prevented EGFP reporter expression

(Figure 7C, D, G, H, K, L, O, P, S, T; Supplemental Figure S3). Thus, despite the larger

cargo size, expression of these activity-modulating CreOn-FlpOff viral constructs was

similar to that of the CreOn-FlpOff-EGFP viral construct (Figure 6). Overall, our data

suggest that the ExBoX system can be used to efficiently control expression of multiple

genes of interest in a Cre- and Flp-dependent manner through AND NOT Boolean

operations.

Discussion

Design of a simpler, more space-efficient Boolean exclusion expression system

The ability to identify and characterize specific cell types and subpopulations has

been challenging in part because, until recently, cell targeting strategies have been

limited to populations defined by expression of a single marker or reporter. In order to

solve this issue and increase specificity of target selection, a number of intersectional

strategies have been developed (Atasoy et al, 2008; Fenno et al, 2014; Gradinaru et al,

2010; Jensen & Dymecki, 2014; Plummer et al, 2015; Schnütgen et al, 2003). A

previous study described the generation of an expression system using Boolean logical

operations, all governed by a single AAV vector (Fenno et al., 2014). This system,

called INTERSECT, incorporated a modular intron-containing system capable of driving

transcription using multiple Boolean logical operations such as AND, NOT, AND NOT,

XOR, and NAND (Fenno et al., 2014). While extremely elegant, the design for the AND

NOT constructs was constrained by the fact that it followed the same internal logic as

the AND INTRSECT constructs. As a result of this constraint the currently existing AND

NOT system has a couple of shortcomings. The first limitation is that this system adds

two introns that make the constructs longer by 194 base-pairs. This is not a serious

concern when expressing from a plasmid, but it becomes relevant when dealing with

viruses with limited packaging capacity like AAV (Dong et al., 1996). Another concern

with the use of introns is that they make the design and cloning of other coding

sequences into these vectors more difficult and cumbersome. In this study, we

generated an alternative system aimed at solving these constraints. Thus, by design,

the ExBoX system has two clear advantages over previous systems: it provides more

cargo room and has a simpler design, allowing for replacement of the CDS cassette in a

simple cloning step.

The ExBoX system allows for robust expression only in the presence of the

respective ON switch, which can be reliably inactivated in the presence of the OFF

switch (Table 1). This is true for both plasmid (Figures 1-4) and viral constructs (Figures

5-7) as validated by a battery of *in vitro* and *in vivo* experiments in embryonic and

postnatal tissues. Altogether, these experiments demonstrate that the ExBoX system is

effective in delivering strong recombinase-dependent expression in a variety of settings.

A Boolean system to control expression with temporal and cell-type specificity

Boolean logic AND-NOT operations made available by ExBoX can be used to

study previously inaccessible subpopulations. In situations where there is no suitable

specific driver for a subpopulation of interest, the ExBoX system can be used to drive

expression of a gene of interest or reporter in said subpopulation, provided there is a

recombinase driver available for a broader neuronal population, and one for the cells to

be excluded (Figure 8A). For example, the AND NOT logic of the ExBoX system could

be utilized to label or genetically manipulate 5HTR3a$^+$;VIP$^-$ interneurons in the cortex

(Tremblay et al., 2016). This could be done by turning on expression of a CreON-FlpOff

construct with *HTR3a-Cre*, while excluding inhibitory neurons expressing *VIP-FlpO*

(Figure 8A) (Che et al., 2018; He et al., 2017; Gerfen et al., 2013).

In addition to cell-specificity, these intersectional approaches can be used to

achieve temporal control. This is particularly important when studying developmental

events or degenerative diseases with defined critical periods. Using ExBoX one could

take advantage of the expression of a recombinase to provide cell-type specificity, and

a second recombinase to turn off expression at the desired time (Figure 8B). For

example, expression of a gene of interest in the DG can be turned ON by injecting

ExBoX AAVs at a desired developmental time point into *Grik4-Cre* mice, and later

turned OFF with Flp, thus creating a window of expression.

The tissue- and temporal-specificity conferred by the ExBoX system will make

developmental studies that were previously limited by the pre-existing tools possible.

With this in mind, we generated ExBoX vectors encoding for neuronal activity and

synaptic transmission modulators, which will allow for tight control of these functional

manipulations in time and space. These activity-modulating tools, although well

characterized in a variety of circuits (see Bando et al., 2016; Burrone et al., 2002; Johns

et al., 1999; Lin et al., 2010; Okada and Matsuda, 2008; Priya et al., 2018; Sim and

Antolin et al., 2013; Sweeney et al, 1995; Xue et al, 2014 for examples), will need to be

validated in each particular setting and/or system. To facilitate studies where the timing

of CDS expression is critical, we are currently generating AAV vectors driving

expression of a tamoxifen-dependent FlpO (data not shown) (Goodrich et al, 2018).

This AAV-FlpOERT2 could be delivered at the same time as the ExBoX AAVs and used

to turn off expression by orally providing tamoxifen at a later timepoint.

Expanding the toolkit to study gene function

Due to the ExBoX's simple design, the system can be easily modified to

incorporate alternative and/or additional recombinase systems. For example, inclusion

of the Dre-Rox recombinase system, in addition to Cre-lox and Flp-frt, would provide

even greater intersectional specificity (Plummer et al, 2015). In addition to Dre-ON and

OFF intersectional AND NOT systems, this could theoretically enable AND-NOT-NOT

Boolean logic operations, permitting an even greater expanse of possibilities. In

addition, by using other promoters in the AAV constructs, this system can be used to

drive expression in any cell-type or organ of interest. The broad applicability and

exquisite specificity of the system would make the use of these vectors particularly

useful for gene therapy applications.

In summary, we designed and generated a new set of viral tools to drive

expression using Boolean exclusion logic, ExBoX. ExBoX AAVs allow for tight spatial

and temporal control of transcription. These new tools are simple and can be easily

modified to express any desired gene of interest. Our design also increases the space

available for other coding sequences of interest, even if this increase in capacity is

somewhat small (194 bps). As proof of concept, we generated a variety of vectors

encoding for neuronal activity and synaptic transmission regulators that will be readily

available to the neuroscience community. Based on its simplicity, we believe the ExBoX

system will help universalize the use of combinatorial expression approaches.

Combining these newly developed tools with genetically encoded Cre and FlpO lines, in

addition to a growing variety of viral tools, will allow for cell-specific gene expression in a

variety of systems.

Materials and Methods

Animals. The ICR mouse strain was purchased from Taconic. The Grik-4-Cre

transgenic mouse strain was purchased from The Jackson Laboratory. Generation of

the Grik4-cre transgenic mouse line has been described previously (Nakazawa et al.,

2002). Mice were housed in a controlled environment maintained at approximately 22*C

on a 12-hour light/dark cycle. Mouse chow and water were provided ad libitum. The day

of vaginal plug observation was designated as embryonic day 0.5 (E0.5), and the day of

birth as postnatal day 0 (P0). All animal procedures presented were performed

according to the University of California Riverside's Institutional Animal Care and Use

Committee (IACUC) guidelines and approved by UC Riverside IACUC.

Viral constructs. The CreOn-FlpOff and FlpOn-CreOff ExBoX viral constructs were

designed using ApE (https://jorgensen.biology.utah.edu/wayned/ape/) and Vector

Builder (vectorbuilder.com) free software. The plasmids were synthesized and

assembled by Vector Builder. Plasmid DNA was purified using the Qiagen Maxi Prep kit

(Qiagen Cat#10023). AAVs were also packaged by Vector Builder into AAV2/9.

pENN.AAV.CamKII 0.4. Cre. SV40 (AAV5) (Addgene viral prep #105558-AAV5;

http://n2t.net/addgene:105558; RRID:Addgene_105558), pAAV-CAG-TdTomato (AAV5)

(Addgene viral prep#59462-AAV5; http://n2t.net/addgene:59462;

RRID:Addgene_59462) and pCAG-FLEX-EGFP-WPRE (AAV9) (Addgene viral prep

#51502-AAV9; http://n2t.net/addgene:51502; RRID:Addgene_51502) were obtained

from Addgene (Oh, S.W., Harris, J.A., Ng, L., et al., 2014). rAAV5/AAV-EF1α-FlpO-

WPRE and rAAV9/CAG-GFP were obtained from the UNC GTC Vector Core.

In utero electroporation. Pregnant ICR mice at E15.5 were anesthetized with

isoflurane and an abdominal incision was made to expose the uterus. Pups were

visualized through the uterine wall. Plasmids diluted in fast green and phosphate

buffered saline (PBS) (3 µg/µl) were injected through sharpened glass capillary needles

into the lateral ventricle. 5mm paddles were used to deliver five 40V pulses of 50ms

each with 950ms intervals across the hemispheres. Uterine horns were repositioned

inside the female and the abdominal cavity was filled with 5x penicillin/streptomycin

(pen/strep) in sterile PBS. Pups were collected at E17.5 via transcardial perfusion with

PBS and 4% paraformaldehyde (PFA) and fixed in 4% PFA for four hours at 4°C, rinsed

with PBS, and sectioned coronally at 150um on a vibrating microtome (VT100S; Leica).

Immunohistochemistry was performed as described (Polleux & Ghosh, 2002). Nuclei

were visualized with DAPI (10g/ml) and primary antibodies used were chicken anti-GFP

(1:500, Aves Labs Cat# GFP-1010) and rabbit anti-dsRed (1:500, Takara Bio Cat#

632475) with secondary antibodies goat anti-chicken 488 (1:1000, Thermo Fisher

Scientific Cat# A-11039) and goat anti-rabbit 546 (1:1000, Abcam Cat# ab60317).

Sections were mounted on glass microscope slides with Fluoro Gel (Electron

Microscopy Sciences Cat# 1798502) and imaged with a laser-scanning confocal

microscope (Leica SPEII).

***Ex utero* electroporation.** *Ex utero* electroporation was performed similarly as *in utero*

electroporation. Pregnant ICR mice at E14.5 were cervically dislocated, uterine horns

were removed, and pups were dissected out in Complete HBSS on ice. Plasmids were

injected bilaterally and 5mm paddles were used to deliver five 40V pulses of 50ms each

with 950 ms intervals across each hemisphere. After electroporation, cortices were

removed for primary cortical culture.

Primary Cortical Neuron Culture. Primary cortical neurons were isolated from E14.5

ICR mouse embryos as described (Kim & Magrané, 2011) using growth media as

described (Polleux and Ghosh 2010). 12mm circular coverslips were treated with 12M

HCL overnight and thoroughly neutralized with deionized water. Acid-treated coverslips

were stored in a glass petri dish with 70% to 100% Ethanol. The coverslips were fire-

sterilized prior to use and coated with laminin and poly-D-lysine (8.3ug/mL) in sterile

water in a 24-well sterile culture plate. *Ex utero* electroporation was performed prior to

dissection in Complete Hanks Balanced Salt Solution (HBSS, 2.5mM HEPES pH7.4,

2.5mM D-glucose, 1mM $CaCl_2$, 1mM $MgSO_4$, 2.5mM $NaHCO_3$). Neurons were plated at

a density of 2.6×10^5cells/cm^2 and maintained in Complete Neurobasal (Neurobasal

media, pen/strep, Glutamax, B27) in 5% CO_2 at 37°C. 50% of the media was

exchanged with fresh media 48 hours after plating. Neurons were collected after 72

hours and fixed in 4% PFA for 5 minutes, then rinsed with PBS 3x for 5 min. Cells were

incubated in blocking buffer (PBS, 1% goat or donkey serum, 0.1% Triton-X) for 30

minutes at RT, rinsed with PBS 1x, then incubated in primary antibodies (chicken anti-

GFP (1:1000, Aves Labs Cat# GFP-1010) and/or rabbit anti-dsRed (1:1000, Takara Bio

Cat# 632475) in antibody dilution buffer (PBS, 0.1% goat or donkey serum) at 4°C

overnight. The following day cells were rinsed in PBS 4x for 5min, and were incubated

in secondary antibodies (goat anti-chicken 488 (1:1000, Thermo Fisher Scientific Cat#

A-11039) and goat anti-rabbit 546 (1:1000, Abcam Cat# ab60317)) and DAPI (1ug/mL)

in antibody dilution buffer for 2 hours, then rinsed with PBS 4x for 10min. Coverslips

were mounted on glass slides with Fluoro Gel and imaged with a confocal microscope.

Neuro2A Cell Culture. Neuro2A (N2A) cells from ATCC were maintained in DMEM

with pen/strep and 10% fetal bovine serum and incubated in 5% CO_2 at 37°C. Cells

were plated on laminin and poly-D-lysine (8.3µg/ml) coated 12mm coverslips in a 24-

well plate at 7.8×10^4 cells/cm^2. Cells were transfected 24 hours later with Metafectene

PRO (Biontex Cat# T040-1.0) according to manufacturer's suggested protocol with a

1:4 DNA to Metafectene ratio. Cells were fixed 48 hours after transfection with 4% PFA

for 5 min, rinsed with PBS, stained with DAPI (1 µm/mL) for one hour, then rinsed again

with PBS 3x for 5min. Coverslips were mounted with Fluoro Gel onto a glass slide and

examined with confocal microscopy.

Stereotaxic Injections. Stereotaxic injections were performed as described previously

(Osten et al., 2007) in P30 mice. Mice were put under anesthesia for the duration of the

procedure using isoflurane. Injections for AAV-FlpOn-CreOff-TdTomato expression

experiments were performed in CD1 mice targeting the DG of the Hippocampus

(coordinates: A/P -1.70 mm; M/L ±1.90 mm; D/V, -1.70 mm). Right hemisphere

injections received a 400nl viral cocktail of AAV-FlpOn-CreOff-TdTomato (6.55×10^{12}

GC/mL), rAAV5/EF1α-FlpO-WPRE, pENN.AAV.CamKII 0.4. Cre. SV40, and pCAG-

FLEX-EGFP-WPRE in a 1:1:1:1 ratio, whereas the contralateral hemisphere received a

200nl viral cocktail containing FlpOn-CreOff-TdTomato (6.55×10^{12} GC/mL), EF1α-FlpO-

WPRE, in a 1:1 ratio. Titers for commercially available AAVs were as follows:

pENN.AAV.CamKII 0.4. Cre. SV40 (2.4×10^{13} GC/ml), pCAG-FLEX-EGFP-WPRE (3.3 x

10^{13} GC/ml), pAAV-CAG-TdTomato (1.2×10^{13} GC/ml), rAAV5/EF1α-FlpO-WPRE (2.6

x 10^{12} GC/ml), and rAAV9/CAG-GFP (2 x 10^{12} GC/ml). FlpO-negative controls were

injected with a viral cocktail of FlpOn-CreOff-TdTomato and pAAV-CAG-GFP. For viral

CreOn-FlpOff-EGFP construct experiments, injections were performed in *Grik4-Cre*

mice. Right hemisphere injections received 200nl of desired CreOn-FlpOff construct,

whereas the contralateral left hemisphere received a 500nl viral cocktail containing the

construct, rAAV5/EF1α-FlpO-WPRE, and pAAV-CAG-TdTomato in a 2:2:1 ratio

(respectively). Viral CreOn-FlpOff-EGFP constructs co-expressing following GOIs were

tested: Kir2.1 (3.34x10^{13} GC/ml), Kir2.1-mutant (4.6x10^{13} GC/ml), NaChBac (2.98x10^{12}

GC/ml), NaChBac-mutant (4.93x10^{13} GC/ml), and TeTN (2.52x10^{12} GC/ml). Three

weeks after injections, mice were perfused and fixed with 4% paraformaldehyde O/N at

4 °C, rinsed in PBS x1, and sectioned coronally on a vibratome (150 μm). Sections were

incubated in 4',6-diamidino-2-phenylindole (DAPI) and mounted onto slides with Fluoro

Gel fluorescence mounting medium (Electron Microscopy Sciences Cat# 1798502).

Expression was observed via confocal microscopy.

Image analysis. N2A and primary cortical neuron image analysis was performed with

FIJI (Schindelin et al., 2012). GFP+ and RFP+ cells were all manually counted. DAPI

counts were obtained by thresholding the DAPI channel, image was made binary, then

the watershed function was used and finally the analyze particles function to count

nuclei with a size between 20-2000 pixels^2 and 0.2-1.0 circularity. Transfection

efficiency was calculated as $\frac{\#RFP\ or\ \#GFP}{\#DAPI} \times 100$. Percentage of transfected cells in which

recombinase activity resulted in fluorescent reporter (RFP or GFP) expression were

calculated as $\frac{\#GFP\ or\ RFP}{\#\ transfected} \times 100$. Images obtained from *in utero* electroporations were

counted manually, avoiding the ventricular zone where it became impossible to

distinguish individual cells. Percentage of ON cells were calculated the same as in N2A

and primary cortical neurons. The N2A, primary cortical culture, and intrauterine

electroporation datasets were tested for normality with the Shapiro-Wilk test and QQ

plot. A One-way ANOVA with Tukey's multiple comparisons test was done for N2A

transfection and intrauterine electroporation data (Figures 1,2,4). Unpaired t-test was

performed on viral injection data (Figures S1 and S3). The data set for primary cortical

cell counts failed to meet normality and thus Fisher's exact test was performed (Figure

3). Image analysis for viral constructs was performed with ImageJ. A region of interest

in granule cell layer of the DG and defined by the site of injection was selected. Mean

pixel intensity of the viral reporter fluorophore in the ROI was measured. An uninjected

region of the hippocampus was simultaneously selected and measured for fluorescence

intensity, and defined as background fluorescence. Background fluorescence was then

subtracted from the mean pixel intensity of the ROI. The percentage of viral reporter

expression maintained when the OFF recombinase was co-injected was calculated as

$\frac{mean\ pixel\ intensity\ OFF}{mean\ pixel\ intensity\ ON} \times 100$ for each animal, and was averaged across samples (n=3) for

each viral construct.

Acknowledgements

We would like to thank Drs. Edward Zagha and Kevin Wright for critically reading the

manuscript and providing helpful comments. We would like to thank the University of

North Carolina Chapel Hill Gene Therapy Center Vector Core for packaging of viral

constructs. pENN.AAV.CamKII 0.4.Cre.SV40 was a gift from James M. Wilson, pCAG-

FLEX-EGFP-WPRE was a gift from Hongkui Zeng, and pAAV-CAG-TdTomato and

pAAV-CAG-GFP were a gift from Edward Boyden. AAV-EF1α-FlpO-WPRE was a gift

from Karl Deisseroth.

Competing interests
The authors declare no competing financial interests.

Funding sources
This work was supported by grants from the National Institutes of Health

(R21MH118640 and R01NS104026 to M.M.R.; R01NS069861 and R01NS097750 to

V.S) and a Hellman Foundation Fellowship to M.M.R.

References

(Bolded names = Co-first authorship)

Antón-bolaños, N., Sempere-ferràndez, A., Guillamón-Vivancos, T., Martini, F. J.,
Pérez-Saiz, L., Gezelius, H., Filipehuk, A., Valdeolmillos, M., López-Bendito, G. (2019).
Prenatal activity from thalamic neurons governs the emergence of functional cortical
maps in mice. *Science*, 364(6444), 987–990. https://doi.org/10.1126/science.aav7617

Atasoy, D., Aponte, Y., Su, H. H., & Sternson, S. M. (2008). A FLEX Switch Targets
Channelrhodopsin-2 to Multiple Cell Types for Imaging and Long-Range Circuit
Mapping. *Journal of Neuroscience, 28*(28), 7025–7030.
https://doi.org/10.1523/JNEUROSCI.1954-08.2008

Awatramani, R., Soriano, P., Rodriguez, C., Mai, J. J., & Dymecki, S. M. (2003). Cryptic
boundaries in roof plate and choroid plexus identified by intersectional gene activation,
Nat. Genet., 35(1), 70–75. https://doi.org/10.1038/ng1228

Bando, Y., Irie, K., Shimomura, T., Umeshima, H., Kushida, Y., Kengaku, M., Fujiyoshi,
Y., Hirano, T., Tagawa, Y. (2016). Control of Spontaneous Ca2+ Transients Is Critical
for Neuronal Maturation in the Developing Neocortex. Cerebral Cortex, 26(1), 106–117.
https://doi.org/10.1093/cercor/bhu180

Branda, C. S., & Dymecki, S. M. (2004). Talking about a Revolution: The Impact of Site-
Specific Recombinases on Genetic Analyses in Mice. *Developmental Cell, 6*, 7–28.

Burrone, J., O'Byrne, M., & Murthy, V. N. (2002). Multiple forms of synaptic plasticity
triggered by selective suppression of activity in individual neurons. *Nature*, 420(6914),
414–418. https://doi.org/10.1038/nature01242

Capecchi, M. (1989). Altering the genome by homologous recombination. *Science*,
244(4910), 1288–1292. https://doi.org/10.1126/science.2660260

Che, A., Babij, R., Iannone, A. F., Fetcho, R. N., Ferrer, M., Liston, C., Fishell, G., De
Marco Garcia, N. V. (2018). Layer I interneurons sharpen sensory maps during neonatal
development. *Neuron*, 99(1), 98–116.
https://doi.org/10.1016/j.neuron.2018.06.002.Layer

Daigle, T. L., Madisen, L., Hage, T. A., Valley, M. T., Knoblich, U., Larsen, R. S., … Li,
L. (2019). A suite of transgenic driver and reporter mouse lines with enhanced brain cell
type targeting and functionality. *Cell*, 174(2), 465–480.
https://doi.org/10.1016/j.cell.2018.06.035.A

Darmanis, S., Sloan, S. A., Zhang, Y., Enge, M., Caneda, C., Shuer, L. M., Hayden
Gephart, M. G., Barres, B. A., & Quake, S. R. (2015). A survey of human brain
transcriptome diversity at the single cell level. *Proceedings of the National Academy of*
Sciences, 112(23), 7285–7290. https://doi.org/10.1073/pnas.1507125112

Dong, J., Fan, P., & Frizzell, R. A. (1996). Quantitative analysis of Packaging Capacity
of Recombinant Adeno-Associated Virus. *Human Gene Therapy*, 7, 2101–2112.

Dymecki, S. M., Ray, R. S., & Kim, J. C. (2010). Mapping Cell Fate and Function Using
Recombinase-Based Intersectional Strategies. Guide to Techniques in Mouse
Development, Part B: Mouse Molecular Genetics. *Methods Enzymol.* (2nd ed., Vol.
477). https://doi.org/10.1016/S0076-6879(10)77011-7

Fenno, L. E., Mattis, J., Ramakrishnan, C., Hyun, M., Lee, S. Y., He, Tucciarone, J.,
Selimbeyoglu, A., Berndt, A., Grosenick, L., Zalocusky, K.A., Bernstein, H., Swanson,
H., Perry, C., Diester, I., Boyce, F.M., Bass, C., Neve, R., Huang, Z.J., & Deisseroth, K.
(2014). Targeting cells with single vectors using multiple-feature Boolean logic. *Nature
Methods, 11*, 763–772. https://doi.org/10.1038/nmeth.2996

Garcıa-Garcia, M. J., Eggenschwiler, J. T., Caspary, T., Alcorn, H. L., Wyler, M. R.,
Huangfu, D., Rakeman, A.S., Lee, J.D., Feinberg, E.H., Timmer, J.R., & Anderson, K. V.
(2005). Analysis of mouse embryonic patterning and morphogenesis by forward
genetics. *PNAS*, *5*(7), 5913–5919. www.pnas.org/cgi/doi/10.1073/pnas.0501071102

Gerfen, C. R., Paletzki, R., & Heintz, N. (2013). GENSAT BAC cre-recombinase driver
lines to study the functional organization of cerebral cortical and basal ganglia circuits.
Neuron, 80(6), 1368–1383. https://doi.org/10.1016/j.neuron.2013.10.016

Glover, C. P., Bienemann, A. S., Heywood, D. J., Cosgrave, A. S., & Uney, J. B. (2002).
Adenoviral-mediated, high-level, cell-specific transgene expression: a SYN1-WPRE
cassette mediates increased transgene expression with no loss of neuron specificity.
Molecular therapy: the journal of the American Society of Gene Therapy, *5*(5 Pt 1), 509–
516. https://doi.org/10.1006/mthe.2002.0588

Goodrich, M. M., Talhouk, R., Zhang, X., & Goodrich, D. W. (2018). An approach for
controlling the timing and order of engineered mutations in mice. *Genesis* (New York,
N.Y. : 2000), 56(8), e23243. https://doi.org/10.1002/dvg.23243

Gordon, J., & Ruddle, F. (1981). Integration and stable germ line transmission of genes
injected into mouse pronuclei. *Science*, 214(4526), 1244–1246.
https://doi.org/10.1126/science.6272397

Gradinaru, V., Zhang, F., Ramakrishnan, C., Mattis, J., Prakash, R., Diester, I.,
Goshen, I., Thompson, K. R., Deisseroth, K. (2010). Molecular and Cellular Approaches
for Diversifying and Extending Optogenetics. *Cell*, *141*(1), 154–165.
https://doi.org/10.1016/j.cell.2010.02.037

Gu, H., Marth, J. D., Orban, P. C., Mosmann, H., & Rajewsky, K. (1994). Deletion of a
DNA Polymerase β Gene Segment in T Cells Using Cell Type-Specific Gene Targeting.
Science, 265(5168), 103–106. https://doi.org/10.1126/science.8016642

He, M., Tucciarone, J., Lee, S., Nigro, M. J., Kim, Y., Levine, J. M., Kelly, S.M.,
Krugikov, I., Wu, P., Chen, Y., Gong, L., Hou, Y., Osten, P., Rudy, B., Huang, Z. J.

(2016). *Neuron*, 91(6), 1228–1243.
https://doi.org/10.1016/j.neuron.2016.08.021.Strategies

Hogan, B., & Williams, J. (1981). Integration of foreign genes into the mammalian germ
line: Genetic engineering enters a new era. *Nature*, 294(5836), 9–10.
https://doi.org/10.1038/294009a0

Jaenisch, R., & Mintz, B. (1974). Simian Virus 40 DNA Sequences in DNA of Healthy
Adult Mice Derived from Preimplantation Blastocysts Injected with Viral DNA.
Proceedings of the National Academy of Sciences, 71(4), 1250–1254.
https://doi.org/10.1073/pnas.71.4.1250

Jensen, P., & Dymecki, S. M. (2014). Essentials of Recombinase-Based Genetic Fate
Mapping in Mice. *Methods Mol Biol*, *1092*, 437–454. https://doi.org/10.1007/978-1-
60327-292-6

Johns, D. C., Marx, R., Mains, R. E., O'Rourke, B., & Marbán, E. (1999). Inducible
genetic suppression of neuronal excitability. *Journal of Neuroscience*, 19(5), 1691–
1697. https://doi.org/10.1523/jneurosci.19-05-01691.1999

Kakava-georgiadou, N., Zwartkruis, M. M., Bullich-vilarrubias, C., Adan, R. A. H., &
Marti, F. (2019). An Intersectional Approach to Target Neural Circuits With Cell- and
Projection-Type Specificity: Validation in the Mesolimbic Dopamine System. *Frontiers in*
Molecular Neuroscience, *12*(February), 1–9. https://doi.org/10.3389/fnmol.2019.00049

Kim H.J., Magrané J. (2011) Isolation and Culture of Neurons and Astrocytes from the
Mouse Brain Cortex. In: Manfredi G., Kawamata H. (eds) Neurodegeneration. Methods
in Molecular Biology (Methods and Protocols), vol 793. Humana Press, Totowa, NJ

Kozorovitskiy, Y., Saunders, A., Johnson, C. A., Lowell, B. B., Sabatini, B. L., Israel, B.,
& Medical, D. (2012). Recurrent network activity drives striatal synaptogenesis. *Nature*,
485(7400), 646–650. https://doi.org/10.1038/nature11052.

Kristianto, J., Johnson, M. G., Zastrow, R. K., Radcliff, A. B., & Blank, R. D. (2017).
Spontaneous recombinase activity of Cre-ERT2 in vivo. *Transgenic research*, 26(3),
411–417. https://doi.org/10.1007/s11248-017-0018-1

Lin, C. W., Sim, S., Ainsworth, A., Okada, M., Kelsch, W., & Lois, C. (2010). Genetically
Increased Cell-Intrinsic Excitability Enhances Neuronal Integration into Adult Brain
Circuits. *Neuron*, 65(1), 32–39. https://doi.org/10.1016/j.neuron.2009.12.001

Madisen, L., Garner, A. R., Carandini, M., Zeng, H., & Cheng, A. (2015). Transgenic
Mice for Intersectional Targeting of Neural Sensors and Effectors with High Specificity
and Performance. *Neuron, 85*, 942–958. https://doi.org/10.1016/j.neuron.2015.02.022

Merte, J., Wang, Q., Vander Kooi, C. W., Sarsfield, S., Leahy, D. J., Kolodkin, A. L., &
Ginty, D. D. (2010). A forward genetic screen in mice identifies Sema3A[K108N], which

binds to Neuropilin-1 but cannot signal. *Journal of Neuroscience, 30*(16), 5767-5775. https://doi.org/10.1523/JNEUROSCI.5061-09.2010

Nakazawa K; Quirk MC; Chitwood RA; Watanabe M; Yeckel MF; Sun LD; Kato A; Carr CA; Johnston D; Wilson MA; Tonegawa S. (2002). Requirement for hippocampal CA3 NMDA receptors in associative memory recall. Science 297(5579):211-8.

Nusslein-Volhard, C., & Wieschaus, E. (1980). Mutations affecting segment number and polarity in Drosophila. *Nature, 287*(October), 795–801. https://doi.org/10.1038/287795a0

Oh, S. W., Harris, J. A., Ng, L., Winslow, B., Cain, N., Mihalas, S., Wang, Q., Lau, C., Kuan, L., Henry, A. M., Mortrud, M. T., Ouellette, B., Nguyen, T., Sorensen, S. A., Slaughterbeck, C., Wakeman, W., Li, Y., Feng, D., … Anh Zeng, H. (2014). A mesoscale connectome of the mouse brain. Nature, 508(7495), 207–214. https://doi.org/10.1038/nature13186 Okada, M., & Matsuda, H. (2008). Chronic lentiviral expression of inwardly rectifying K+ channels (Kir2.1) reduces neuronal activity and downregulates voltage-gated potassium currents in hippocampus. *Neuroscience*, 156(2), 289–297. https://doi.org/10.1016/j.neuroscience.2008.07.038

Osten, P., Cetin, A., Komai, S., Eliava, M., & Seeburg, P. H. (2007). Stereotaxic gene delivery in the rodent brain. *Nature Protocols*, 1(6), 3166–3173.

Plummer, N. W., Evsyukova, I. Y., Robertson, S. D., Marchena, J. De, Tucker, C. J., & Jensen, P. (2015). Expanding the power of recombinase-based labeling to uncover cellular diversity. *Development, 142*, 4385–4393. https://doi.org/10.1242/dev.129981

Polleux, F., & Ghosh, A. (2002). The Slice Overlay Assay: A Versatile Tool to Study the Influence of Extracellular Signals on Neuronal Development. *Science's STKE*, 2002(136), pl9.

Priya, R., Paredes, M. F., Karayannis, T., Yusuf, N., Liu, X., Jaglin, X., Graef, I., Alvarez-Buylla A., Fishell, G. (2018). Activity Regulates Cell Death within Cortical Interneurons through a Calcineurin-Dependent Mechanism. *Cell Reports*, 22(7), 1695–1709. https://doi.org/10.1016/j.celrep.2018.01.007

Rubin, G., & Spradling, A. (1982). Genetic transformation of Drosophila with transposable element vectors. *Science, 218*(4570), 348–353. https://doi.org/10.1126/science.6289436

Sauer, B., & Mcdermott, J. (2004). DNA recombination with a heterospecific Cre homolog identified from comparison of the pac-c1 regions of P1-related phages. *Nucleic Acids Research, 32*(20), 1–10. https://doi.org/10.1093/nar/gkh941 1–10 doi:10.1093/nar/gkh941

Saunders, A., Johnson, C. A., Sabatini, B. L., & Miyamichi, K. (2012). Novel recombinant adeno-associated viruses for Cre activated and inactivated transgene

expression in neurons. *Frontiers in Neural Circuits*, *6*(47), 1–10.
https://doi.org/10.3389/fncir.2012.00047

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source
platform for biological-image analysis. *Nature Methods*, 9(7), 676–682.

Schnütgen, F., Doerflinger, N., Calléja, C., Wendling, O., Chambon, P., & Ghyselinck,
N. B. (2003). A directional strategy for monitoring Cre-mediated recombination at the
cellular level in the mouse. *Nature Biotechnology, 21*, 562–565.
https://doi.org/10.1038/nbt811

Sim, S., Antolin, S., Lin, C. W., Lin, Y. X., & Lois, C. (2013). Increased cell-intrinsic
excitability induces synaptic changes in new neurons in the adult dentate gyrus that
require Npas4. *Journal of Neuroscience*, 33(18), 7928–7940.
https://doi.org/10.1523/JNEUROSCI.1571-12.2013

Sternson, S. M., Atasoy, D., Betley, J. N., Henry, F. E., & Xu, S. (2016). An Emerging
Technology Framework for the Neurobiology of Appetite. *Cell Metabolism*, *23*(2), 234–
253. https://doi.org/10.1016/j.cmet.2015.12.002

Sweeney, S.T., Broadie, K., Keane, J., Niemann H., O'Kane, C.J. (1995). Targeted
expression of tetanus toxin light chain in Drosophila specifically eliminates synaptic
transmission and causes behavioral defects. *Neuron* 14, 341–351.

Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic Interneurons in the Neocortex:
From Cellular Properties to Circuits. *Neuron*, 91(2), 260–292.
https://doi.org/10.1016/j.neuron.2016.06.033

Tsien, J. Z., Chen, D. F., Gerber, D., Tom, C., Mercer, E. H., Anderson, D. J., Mayford,
M., Kandel, E. R., & Tonegawa, S. (1996). Subregion- and Cell Type–Restricted Gene
Knockout in Mouse Brain. *Cell*, 87(7), 1317–1326. https://doi.org/10.1016/S0092-
8674(00)81826-7

Wright, K. M., Lyon, K., Leung, H., Leahy, D. J., Ma, L., & David, D. (2013).
Dystroglycan organizes axon guidance cue localization and axonal pathfinding. *Neuron*,
76(5), 931–944. https://doi.org/10.1016/j.neuron.2012.10.009.

Xue, M., Atallah, B. V., & Scanziani, M. (2014). Equalizing excitation-inhibition ratios
across visual cortical neurons. *Nature*, *511*(7511), 596–600.
https://doi.org/10.1038/nature13321

Zhang, F., Aravanis, A.M., Adamantidis, A., Lecea, L., and Deisseroth, K. (2007).
Circuit-breakers: optical technologies for probing neural signals and systems. *Nature
Reviews Neuroscience* 8(August), 577–581. https://doi:10.1038/nr n2192

Figure Legends

Table 1. Conceptual basis for the Expression by Boolean Exclusion (AND NOT)

system using multiple recombinases. Expression of a gene of interest (GOI) can be

spatially and temporally manipulated through the use of multiple recombinases. In this

model, expression of a GOI depends on the presence of an ON recombinase, and is

prevented by the presence of an OFF recombinase. Thus, gene expression only occurs

in cells where the ON recombinase is present AND NOT the OFF recombinase.

Controlling where and when the ON and OFF recombinases are expressed provides a

high level of specificity for targeting cell populations.

Figure 1. Design and proof of concept for CreOn-FlpOff constructs. (A) Schematic

representation of the CreOn-FlpOff construct. It is composed of an inverted expression

cassette encoding for a coding sequence (CDS) fused to an HA tag, a P2A site, and

EGFP, which is flanked by Frt, lox2722, and loxP sites. Presence of Cre recombinase

results in inversion of the cassette and expression. Presence of FlpO results in removal

of the cassette. **(B)** Expression of fluorescent reporter proteins in Neuro2A cells

transfected with CreOn-FlpOff-EGFP construct (green) and pCAG-mCherry (red,

loading control). Co-transfection with pCAG-Cre plasmid causes inversion of the

cassette into the "ON" position and EGFP expression. Co-transfection of pCAG-FlpO

deletes the cassette and abolishes EGFP expression. (B, right panel) Quantification of

N2A cell transfections. Upon transfection with Cre, the CreOn-FlpOff-EGFP construct

results in 93.9%±0.3 of transfected cells expressing EGFP, while co-transfection with

FlpO results in abolishment of EGFP expression. N=3 independent experiments, error

bars ±SEM. ANOVA p<0.001; Tukey post-hoc test *** p<0.0001 vs. all other

transfections. Scale bar 30μm.

Figure 2. Design and proof of principle for FlpOn-CreOff constructs. (A) Schematic

representation of the FlpOn-CreOff construct. The inverted expression cassette is

flanked by lox2722, F14, and Frt sites. Presence of Flp recombinase results in inversion

of the cassette and expression. Presence of Cre results in removal of the cassette. **(B)**

Expression of fluorescent reporters in Neuro2A cells transfected with FlpOn-CreOff-

TdTomato construct (red) and pCAG-EGFP (green, transfection control) in Neuro2A

cells. Co-transfection with pCAG-FlpO causes inversion of the cassette into the "ON"

position and TdTomato expression. Transfection with pCAG-Cre removes the cassette

and abolishes TdTomato expression. Upon transfection with FlpO, the FlpOn-CreOff-

TdTomato construct results in detectable expression of reporter in 85.8%±5.6 of

transfected cells, while co-transfection with Cre results in abolishment of TdTomato

expression (right panel). N=3 independent experiments, error bars ±SEM. ANOVA

p<0.01; Tukey post-hoc test *** p<0.0001 vs. all other transfections. Scale bar 30μm.

Figure 3. Validation of CreOn-FlpOff in primary cortical neurons. (A) Primary

cortical neurons transfected with CreOn-FlpOff and Cre results in 100%±0.0 of

transfected cells expressing EGFP. Without Cre or in the presence of FlpO there is no

EGFP expression in any neurons. pCAG-mCherry was co-transfected as transfection

control. **(B)** Primary cortical neurons transfected with FlpOn-CreOff-TdTomato and FlpO

results in detectable TdTomato expression in 92.1%±2.0 of transfected cells (right).

pCAG-EGFP was co-transfected as transfection control. Without FlpO, or in the

presence of Cre, there is no TdTomato expression in any neurons. N=3, error bars

±SEM. Fisher's exact test ***$p<0.0001$. Scale bar 30μm.

Figure 4. *In vivo* validation of CreOn-FlpOff and FlpOn-CreOff vectors via *in utero*

electroporation (IUE). (A) Co-electroporation of CreOn-FlpOff-EGFP with mCherry

reporter, Cre + pCAG-mCherry, FlpO-mCherry, or Cre + FlpO-mCherry in e15.5 ICR

mouse embryos. Electroporation of CreOn-FlpOff-EGFP with Cre results in 97.7%±0.9

of transfected cells expressing EGFP (right). Electroporation with FlpO results in no

expression and electroporation with Cre and FlpO results in only 1.7%±0.9 of cells still

expressing EGFP. mCherry expression serves as electroporation control. **(B)** Co-

electroporation of FlpOn-CreOff-TdTomato with pCAG-IRES-EGFP reporter, Cre-IRES-

EGFP, FlpO, or Cre-IRES-EGFP + FlpO in e15.5 Swiss-Webster embryos. pCAG-IRES-

EGFP expression was used as electroporation control.Electroporation of FlpOn-CreOff-

TdTomato with FlpO results in 86.9%±3.1 of transfected cells expressing TdTomato

(right). Co-electroporation with Cre, or with Cre and FlpO, results in no cells expressing

TdTomato. N=3, error bars ±SEM. ANOVA $p<0.001$ for both sets of experiments; Tukey

post-hoc test: *** $p<0.0001$ vs. all other transfections. Scale bar 100μm.

Figure 5. *In-vivo* validation of FlpOn-CreOff AAV by stereotactic injection in

postnatal DG neurons. (A-I) Expression of viral FlpOn-CreOff-TdTomato construct

(red) in WT animals co-injected with pAAV-CAG-GFP **(A, D, G)**, with AAV-EF1α-FlpO-

WPRE **(B, E, H),** or AAV-EF1α-FlpO-WPRE + AAV-CamKII-Cre + AAV-CAG-FLEX-

EGFP **(C, F, I)**. EGFP expression is used to visualize the injection site in the absence of

TdTomato expression. Even in the presence of FlpO, Cre injection resulted in

inactivation of TdTomato reporter expression (**C**, **F**). DAPI (blue) was used to visualize

nuclei. Scale bar: 150 μm.

Figure 6. In-vivo validation of CreOn-FlpOff AAV in the dentate gyrus of postnatal

mice. (A-I) Expression of viral CreOn-FlpOff-EGFP construct (green) in WT animals (A,

D, G), in Grik4-Cre animals (B, E, H), and in Grik4-Cre mice co-injected with AAV-

EF1α-FlpO-WPRE (C, F, I). pAAV-CAG-TdTomato (red) was co-injected when

necessary to visualize the injection site. Even in the presence of Cre, FlpO injection

resulted in inactivation of reporter expression by the CreOn-FlpOff-EGFP virus (C, F).

EGFP labeling in the molecular layer of AAV-EF1α-FlpO injected animals (C,F) is from

axonal projections from the contralateral side. DAPI (blue) was used to visualize nuclei.

Scale bar: 150 μm.

Figure 7. *In-vivo* validation of CreOn-FlpOff AAV constructs to manipulate

neuronal activity. (A-T) Expression of fluorescent reporter by CreOn-FlpOff AAVs

(green) co-expressing (A-D) Kir2.1, (E-H) Kir2.1-Mutant, (I-L) NaChBac, (M-P)

NaChBac-mutant, and **(Q-T)** TETN in WT (A, E, I, M, Q) , *Grik4-Cre* animals **(B, F, J,**

N, R) and *Grik4-Cre* animals co-injected with AAV-EF1α-FlpO-WPRE (C, D, G, H, K, L,

O, P, S, T). Injections delivering AAV-EF1α-FlpO included pAAV-CAG-TdTomato (red)

in order to visualize infected cells. In the presence of Cre and FlpO, expression of the

CreOn-FlpOff virus was inactivated (C, D, G, H, K, L, O, P, S, T). EGFP labeling in the

molecular layer of TeTN/FlpO injected animals (T) is from axonal projections from the

contralateral side. Scale bar: 150 μm.

Figure 8. The ExBoX system can be utilized in multiple ways. (A) Expression of a

gene of interest (GOI) can be restricted to a specific subpopulation of cells. Only the

subpopulation of cells expressing the ON recombinase, but not the OFF recombinase,

will express the GOI. **(B)** ExBoX can also be used for temporal control of expression. In

this scheme, activation of the ON recombinase is used just prior to a critical period of

interest. Introduction or activation of the OFF recombinase at the end of this critical

period will turn off expression of the GOI. This serves to reduce confounding factors

during later experimental observation, as GOI expression is no longer present.

	No Recombinase	Cre Only	FlpO Only	Cre and FlpO
Cre = ON FlpO = OFF	○○○ ○○○ ○○○	🟢🟢🟢 🟢🟢🟢 🟢🟢🟢	○○○ ○○○ ○○○	○○○ ○○○ ○○○
FlpO = ON Cre = OFF	○○○ ○○○ ○○○	○○○ ○○○ ○○○	🔴🔴🔴 🔴🔴🔴 🔴🔴🔴	○○○ ○○○ ○○○

Table 1. Conceptual basis for the Expression by Boolean Exclusion (AND NOT) system using multiple recombinases.

A

B

Fig1

A

F14

Syn promoter ◁▶ Tdtomato P2A HA CDS ◀ WPRE Poly A

Frt

↓ FlpE

Syn promoter CDS HA P2A Tdtomato WPRE Poly A

Lox2722

↓ Cre

Syn promoter WPRE Poly A

B

FlpOn-CreOff-TdTomato

| | Vehicle | +Cre | +FlpO | +Cre +FlpO |

DAPI EGFP TdTomato

TdTomato

FlpOn-CreOff-TdTomato

***

% ON (RFP/GFP)

Vehicle   +Cre   +FlpO   +FlpO +Cre

Constructs

Fig2

Fig3

Fig4

AAV-FlpON-CreOff-TdTomato

Fig5

AAV-CreON-FlpOff-EGFP

-Cre | +Cre | +Cre +Flp

DAPI EGFP TdTomato

EGFP

TdTomato

Fig6

Fig7

A

Expression of Recombinases

Expression of GOI

No recombinase
+ON recombinase
+OFF Recombinase (± ON)

No Expression of GOI
Expression of GOI

B

Expression of GOI

Time

Critical Period

ON Recombinase

OFF Recombinase

Experimental observation

Fig8