Global drivers of obligate mycorrhizal symbionts diversification

Benoît Perez-Lamarque [1,2] *, Maarja Öpik [3], Odile Maliet [1], Ana C. Afonso Silva [1], Marc-André Selosse [2,4], Florent Martos [2], and Hélène Morlon [1]

[1] *Institut de biologie de l'École normale supérieure (IBENS), École normale supérieure, CNRS, INSERM, Université PSL, 46 rue d'Ulm, 75 005 Paris, France*

[2] *Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier 75 005 Paris, France*

[3] *University of Tartu, 40 Lai Street, 51 005 Tartu, Estonia*

[4] *Department of Plant Taxonomy and Nature Conservation, University of Gdansk, Wita Stwosza 59, 80-308 Gdansk, Poland*

* *corresponding author: benoit.perez@ens.psl.eu; ORCID: 0000-0001-7112-7197*

Abstract (200 words):

Arbuscular mycorrhizal fungi (AMF) are widespread microscopic fungi that provide mineral nutrients to most land plants by forming one of the oldest terrestrial symbioses. They have sometimes been referred to as an "evolutionary cul-de-sac" for their limited species diversity and their ecological niches restricted to plant-symbiotic life style. Here we use the largest global database of AMF to analyze their diversification dynamics in the past 500 million years (Myr) based on the small subunit (SSU) rRNA gene. We demonstrate that overall the SSU rRNA gene is variable enough to delineate AMF species and find that AMF have low diversification rates. After a diversification peak around 150 Myr ago, they experienced an important diversification slowdown in the last 100 Myr, likely related to a shrinking of their mycorrhizal niches. Our results identify patterns and drivers of diversification in a group of obligate symbionts of major ecological and evolutionary importance. They also highlight a striking example of a diversification slowdown that, instead of reflecting an adaptive radiation as typically assumed, may result from a limited ability to colonize new niches in an evolutionary cul-de-sac.

Introduction:

Arbuscular mycorrhizal fungi (AMF - subphylum Glomeromycotina) are obligate symbionts sometimes referred to as an "evolutionary cul-de-sac, albeit an enormously successful one" [1, 2]. They have limited morphological and species diversities, yet associate with the roots of >80% of land plants, where they provide mineral resources in exchange for photosynthates [3]. Present in most terrestrial ecosystems, AMF play key roles in plant protection, nutrient cycling, and ecosystem functions [4]. Fossil evidence and molecular phylogenies suggest that AMF contributed to the emergence of land plants [5–8] and coevolved with them for more than 400 million years (Myr)[8–10].

Despite the ecological ubiquity and evolutionary importance of AMF, large-scale patterns of their evolutionary history are poorly known. Studies on the diversification of AMF have been hampered by the difficulty of delineating species, quantifying global scale species richness, and building a robust phylogenetic tree for this group. Indeed, AMF are microscopic soil- and root-dwelling fungi that are poorly differentiated morphologically and difficult to cultivate. Although their classical taxonomy is mostly based on the characters of spores and root colonization [3, 11], AMF species delineation has greatly benefited from molecular data [12]. Experts have defined "virtual taxa" (VT) based on a minimal 97% similarity of a region of the 18S small subunit (SSU) rRNA gene and monophyly criteria [13, 14]. As for many other pragmatic species concepts, VT have rarely been tested for their biological relevance [15], and a consensual system of AMF classification is still lacking [16]. AMF are also poorly known genetically: the full SSU rRNA gene sequence is known in few species [17], other gene sequences in even fewer [10, 18], and complete genomes in very few [19].

The drivers of AMF diversification are unknown. A previous dated phylogenetic tree of VT found that many speciations occurred after the last major continental reconfiguration around 100 Myr ago [20], suggesting that AMF diversification is not linked to vicariant speciation during this geological event. Still, geographical speciation could play an important role in AMF diversification, as these organisms have spores that disperse

efficiently [21–23], which could result in frequent founder-event speciation [24]. Other

abiotic factors include habitat: tropical grasslands have, for example, been suggested as

diversification hotspots for AMF [25]. Besides abiotic factors, AMF are obligate symbionts

and, although relatively generalist [4, 26, 27], their evolutionary history could be largely

influenced by a diffuse coevolution with their host plants [10, 28, 29]. Over the last 400

Myr, land plants have experienced massive extinctions and radiations [29, 30], adaptations

to various ecosystems [31, 32], and associations with different soil microorganisms [33, 34].

All these factors could have influenced diversification dynamics in AMF.

Here, we reconstruct several thoroughly sampled phylogenetic trees of AMF,

considering several criteria of species delineations and uncertainty in phylogenetic

reconstructions. We combine this phylogenetic data with paleoenvironmental data and

data of current AMF geographic distributions, ecological traits, interaction with host

plants, and genetic diversity to investigate the global patterns and drivers of AMF

diversification in the last 500 Myr.

Material & methods:

Database choice:

We used AMF SSU rRNA gene sequences from MaarjAM database, the largest global database of AMF gene sequences [13]. Ideally, our analyses would be based on several independent genomic regions, however such data has yet to be acquired. In fungi, the usual barcode is the ITS region. However, for several reasons the SSU rRNA gene has been preferred over the ITS in AMF [35], and therefore ITS data on AMF are currently less common. In addition, we confirmed using the dataset of [35] that the ITS are very difficult to align, preventing us from reconstructing a robust phylogeny to investigate AMF evolutionary history (Supplementary Fig. 1).

Virtual taxa phylogenetic reconstruction:

We reconstructed several Bayesian phylogenetic trees of the 384 VT from the corresponding representative sequences available in the MaarjAM database [13] updated in June 2019 (Supplementary Methods 1). We used the full length (1,700 base pairs) SSU rRNA gene sequences from [17] to better align the VT sequences using MAFFT [36]. We selected the 520 base pair central variable region of the VT aligned sequences and performed a Bayesian phylogenetic reconstruction using BEAST2 [37]. We obtained a consensus VT tree and selected 12 trees equally spaced in 4 independent Bayesian chains to account for phylogenetic uncertainty in the subsequent diversification analyses, hereafter referred to as the VT replicate trees. We set the crown root age at 505 Myr [20], which is coherent with fossil data and previous dated molecular phylogenies [8, 10].

Delineation into Evolutionary Units (EUs):

We considered several ways to delineate AMF species based on the SSU rRNA gene.
In addition to the VT species proxy, we delineated AMF *de novo* into evolutionary units
(EUs) using 5 different thresholds of sequence similarity ranging from 97 to 99% and a
monophyly criterion. We gathered 36,411 AMF sequences of the SSU rRNA gene from
MaarjAM, mainly amplified by the primer pair NS31–AML2 (variable region) [38, 39]
(dataset 1, Supplementary Table 1), corresponding to 27,728 haplotypes. We first built a
phylogenetic tree of these haplotypes and then applied to this tree our own algorithm (R-
package RPANDA [40, 41]) that traverses the tree from the root to the tips, at every node
computes the average similarity of all sequences descending from the node, and collapses
the sequences into a single EU if their sequence dissimilarity is lower than a given
threshold (Supplementary Methods 2). Finally, we performed Bayesian phylogenetic
reconstructions of the EUs using BEAST2, following the procedure described in
Supplementary Methods 1.

Coalescent-based species delineation analyses:

Finally, we considered the Generalized Mixed Yule Coalescent method (GMYC) [42,
43], a species delineation approach that does not require specifying an arbitrary similarity
threshold. GMYC estimates the time *t* in a reconstructed calibrated tree that separates
species diversification (Yule process – before *t*) and intraspecific differentiation (coalescent
process – after *t*). GMYC is too computationally intensive to apply on the 36,411 SSU
sequences; we used it here on three smaller clades to investigate the ability of the SSU gene
to delineate AMF species despite its slow evolution [16], and as a way to evaluate the
biological relevance of the VT and various EUs delineations. We selected the following
AMF clades: the family Claroideoglomeraceae; the order Diversisporales; and an early-
diverging clade composed of the orders Archaeosporales and Paraglomerales. For each
clade, we reconstructed Bayesian phylogenetic trees of haplotypes following the procedure
described in Supplementary Methods 1. We then ran GMYC analyses (splits R-package
[44]) on each of these trees and evaluated the support of the GMYC model compared to a

null model in which all tips are assumed to be different species, using a likelihood ratio

test (LRT). If the LRT supports the GMYC model, different SSU haplotypes belong to the

same AMF species, *i.e.* the SSU rRNA gene has time to accumulate substitutions between

AMF speciation events.

Total diversity estimates:

We evaluated how thoroughly sampled our species-level AMF phylogenetic trees

are by estimating the total number of VT and EUs using rarefaction curves and the

Bayesian Diversity Estimation Software (BDES [45]) (Supplementary Methods 3).

Diversification analyses:

We estimated lineage-specific diversification rates using ClaDS, a Bayesian

diversification model that accounts for rate heterogeneity by modeling small rate shifts at

speciation events [46]. At each speciation event, the descending lineages inherit new

speciation rates sampled from a log-normal distribution with an expected value log[$\alpha \times \lambda$]

(where λ represents the parental speciation rate and α is a trend parameter) and a standard

deviation σ. We considered the model with constant turnover ε (*i.e.* constant ratio between

extinction and speciation rates; *ClaDS2*) and ran a newly developed ClaDS algorithm based

on data augmentation techniques which enables us to estimate mean rates through time

(https://github.com/OdileMaliet/ClaDS_Julia). We ran ClaDS2 with 3 independent chains,

checked their convergence using a Gelman-Rubin diagnostic criterion [47], and recorded

lineage-specific speciation rates. We also recorded the estimated hyperparameters (α, σ, ε)

and the value m=$\alpha \times \exp(\sigma^2/2)$, which indicates the general trend of the rate through time

[46].

In addition, we applied TreePar [48], another diversification approach that does not

consider rate variation across lineages, but models temporal shifts in diversification rates

affecting all lineages simultaneously. We searched for up to ten shifts in diversification rates at every 2-million-year interval in each phylogenetic tree. We estimated the number of temporal shifts in AMF diversification rates using maximum likelihood inferences and likelihood ratio tests. We also used CoMET, its equivalent piecewise-constant model in a Bayesian framework (TESS R-package [49, 50]). We chose the Bayesian priors according to maximum likelihood estimates from TreePar, disallowed mass extinction events, and ran the MCMC chains until convergence (minimum effective sample sizes of 500).

We also fitted a series of time-dependent and environment-dependent birth-death diversification models using RPANDA [40, 51] to test the influence of temperature, pCO_2, and land plant fossil diversity on AMF diversification. For the time-dependent models, we considered models with constant or exponential variation of speciation rates through time and null or constant extinction rates (*fit_bd* function). Models with constant rates correspond to the null hypothesis of clock-like speciation, whereas the exponential variation ensures positive rates and are an approximation of diversity dependence, a process often invoked during radiations [52]. For the environment-dependent models, we considered an exponential dependency of the speciation rates with the environmental variable (env), *i.e.* speciation rate=b*exp(a*env), where a and b are two parameters estimated by maximum likelihood (*fit_env* function). The exponential variation corresponds here to a simple linear regression of log-transformed rates. Environment curves were smoothed using the function *smooth.spline* (stats R-package). We used the corrected Akaike information criterion (AICc) to select the best-fit models, considering that a difference of 2 in AICc indicates that the model with the lowest AICc is better.

The influence of temperature was tested on the complete AMF phylogenetic trees, using estimates of past global temperature [53]. As these temporal analyses can be sensitive to the root age calibration, we replicated them using the youngest (437 Myr) and oldest (530 Myr) crown age estimates from [10]. The influence of pCO_2 [54] and of land plant fossil diversity was tested starting from 400 Myr ago, as these environmental data are not available for more ancient times. For these analyses we sliced the phylogenies at 400 and

200 Myr ago, and applied the diversification models to the sliced sub-trees larger than 50

tips. Estimates of land plant diversity were obtained using all available Embryophyta

fossils from the Paleobiology database (https://paleobiodb.org) and using the shareholder

quorum subsampling method (Supplementary Methods 4; [55]).

All diversification analyses were performed for each delineation on the consensus

and on the 12 replicate trees to account for phylogenetic uncertainty. We also considered

missing species by imputing sampling fractions, computed as the number of observed VT

or EUs divided by the corresponding BDES estimates of global AMF diversity estimated

from Sichel distributions (Supplementary Table 2). As such analyses depend on diversity

estimates, we replicated all diversification analyses using lower sampling fractions down

to 50%.

Testing for potential drivers of AMF diversification:

To further investigate the potential factors driving AMF diversification, we assessed

the relationship between lineage-specific estimates of present-day speciation rates and

characteristics of each AMF taxonomic unit, *i.e.* VT or EUs.

First, we characterized AMF niche width using a set of 10 abiotic and biotic variables

recorded in MaarjAM database for each AMF unit. In short, among a curated dataset

containing AMF sequences occurring only in natural ecosystems (dataset 2; [26]), for each

AMF unit, we reported the number of continents, ecosystems, climatic zones,

biogeographic realms, habitats, and biomes where it was sampled, as well as its number of

plant partners, their phylogenetic diversity, and its centrality in the plant-fungus bipartite

network, and performed a principal component analysis (PCA; Supplementary Methods

5). For AMF units represented by at least 10 sequences, we tested whether these PCA

coordinates reflecting AMF niche widths were correlated with the present-day speciation

rates using both linear mixed-models (not accounting for AMF phylogeny) or

MCMCglmm models [56]. For MCMCglmm, we assumed a Gaussian residual distribution, included the fungal phylogenetic tree as a random effect, and ran the MCMC chains for 1,300,000 iterations with a burn-in of 300,000 and a thinning interval of 500.

Next, we tested the relationship between speciation rates and geographic characteristics of AMF units. To test the effect of latitude, we associated each AMF unit with its set of latitudes and used similar MCMCglmm with an additional random effect corresponding to the AMF unit. To account for inhomogeneous sampling along the latitudinal gradient, we re-ran the model on jackknifed datasets (we re-sampled 1,000 interactions per slice of latitude of twenty degrees). Similarly, we tested the effect of climatic zone and habitat on the speciation rates.

Finally, to test the effect of dispersal capacity, we assessed the relationship between spore size and speciation rate for the few (n=32) VT that contain sequences of morphologically characterized AMF isolates [57]. We gathered measures of their average spore length [57] and tested their relationship with speciation rate by using a phylogenetic generalized least square regression (PGLS).

Estimating genetic diversity:

As a first attempt at connecting AMF macroevolutionary diversification to microevolutionary processes, we measured intraspecific genetic diversities across AMF units. For each AMF unit containing at least 10 sequences, we computed genetic diversity using Tajima's estimator [58]($\theta\pi$; Supplementary Methods 6). Using similar statistical tests as above, we investigated the correlation of AMF genetic diversity with speciation rate, niche width, geographic characteristics, and spore size. We tested the robustness of the results to the minimal number of sequences per AMF unit (10, 15 or 20) used to compute genetic diversity and to perform the PCA.

250   These statistical models were replicated on the different phylogenetic trees

251 (consensus or replicates) for each delineation and we reported p-values (P) corresponding

252 to two-sided tests.

Results:

AMF species delineations & phylogenetic reconstructions:

The EU97.5 and EU98 delineations (obtained using a threshold of 97.5% and 98% respectively) provided a number of AMF units (340 and 641) comparable to the 384 currently recognized VT, while the EU97 delineation had much less (182). Conversely, the EU98.5 and EU99 delineations yielded a much larger number of AMF units (1,190 and 2,647, respectively; Supplementary Tables 3 & 4) that was consistent with the number obtained using GMYC analyses (Supplementary Tables 4 & 5). This supports the idea that some VT might lump together several cryptic species [16](Supplementary Note 1), and that a 98.5 or 99% similarity threshold is more relevant for AMF species delineation. In addition, the GMYC indicated that the actual level of genetic variation within the SSU marker is overall sufficient to separate AMF species-like units among SSU haplotypes (GMYC LRT: P<0.05; Supplementary Fig. 2); on average, there are 10 SSU haplotypes for one AMF unit delineated using GMYC (Supplementary Table 5). Rarefaction curves as well as Bayesian and Chao2 estimates of diversity suggested that more than 90% of the total diversity of AMF is represented in our dataset regardless of the delineation threshold (Fig. 1b, Supplementary Tables 2, 5, & 6; Supplementary Note 2), which is consistent with the proportion of new AMF units detected in recent studies [59].

The reconstructed Bayesian phylogenetic trees based on VT and EU species delineations did not yield high support for the nodes separating the main AMF orders; yet, they had similar topologies and branching times of the internal nodes overall (Fig. 2, Supplementary Figs. 3). As expected, finer delineations resulted in an increase in the number of nodes close to the present (Supplementary Figs. 4). However, we observed a slowdown in the accumulation of new lineages close to the present in all lineage through time plots (LTTs), including those with the finest delineations (EU98.5 and EU99; Supplementary Fig. 5).

Temporal diversification dynamics:

AMF speciation rates ranged from 0.005 to 0.03 events per lineage per Myr (Fig. 2; Supplementary Fig. 6), and varied both within and among AMF orders, with Glomerales and Diversisporales having the highest present-day speciation rates (Supplementary Fig. 7). AMF experienced their most rapid diversification between 200 and 100 Myr ago according to estimates of diversification rates through time obtained with ClaDS (Fig. 2; Supplementary Fig. 8), and 150-50 Myr ago according to diversification models with piecewise constant rates (TreePar [48] and CoMET [50], Fig. 2; Supplementary Figs. 9 & 10).

The fast diversification of AMF around 150 Myr ago was followed by a slowdown in the recent past (Fig. 2; Supplementary Fig. 8), as suggested by the plateauing of the LTTs. A global decrease of the speciation rates through time was independently supported by ClaDS, TreePar, and CoMET analyses (Supplementary Figs. 11, 9 & 10), as well as time-dependent models in RPANDA [60](Supplementary Fig. 12). This slowdown was robust to all species delineations (Supplementary Figs. 8, 9, 10, & 12), the branching process prior (analyses not shown), phylogenetic uncertainty, and sampling fractions down to 50%, except in ClaDS analyses where the trend disappeared in some EU99 trees and for sampling fractions lower than 70% (Supplementary Figs. 13, 14, & 15).

We did not find a strong signal of extinction in our analyses: the turnover rate estimated from ClaDS was generally close to zero (Supplementary Fig. 11b), and models including extinctions were never selected in RPANDA (Supplementary Fig. 12). Similarly, the extinction rates estimated in piecewise-constant models were not significantly different from 0 (Supplementary Fig. 16).

AMF diversification drivers:

When fitting environment-dependent models of diversification, we found high support for temperature-dependent models compared to time-dependent models for all AMF delineations, sampling fractions, and crown ages (Fig. 3; Supplementary Figs. 17, 18, 19, 20, & 21), with the exception of some EU99 trees with a 50% sampling fraction (Supplementary Fig. 21). This signal of temperature dependency was not due to a temporal trend (Supplementary Note 3) nor to an artefact caused by rate heterogeneities (Supplementary Note 4). Evidence for temperature dependency, however, decreased in some clades closer to the present (see Methods), as small trees tend to be best fit by constant rather than environment-dependent models (Supplementary Fig. 22). We detected a significant positive dependency of the diversification rates on CO_2 concentrations in some sub-trees, but rarely found a significant effect of plant fossil diversity (Supplementary Fig. 22).

The PCA of AMF niche width characteristics had a first principal component (PC1) that indicated the propensity of each AMF unit (VT or EUs) to be vastly distributed among continents, ecosystems and/or associated with many plant species and lineages, whereas the second principal component (PC2) indicated the propensity of a given AMF unit to associate with few plant species on many continents (Fig. 4; Supplementary Fig. 23 & 24). Hence, PC1 reflects AMF niche width, whereas PC2 discriminates the width of the abiotic relatively to the biotic niche (Fig. 4a-b; Supplementary Fig. 25). We found a positive correlation between PC1 and lineage-specific speciation rates in the majority of the VT and EU99 trees (Fig. 4c-d; Supplementary Fig. 26a). However, these results were no longer significant when controlling for phylogenetic non-independence between AMF units (Supplementary Fig. 26b), likely because a single *Glomeraceae* clade, including the abundant and widespread morphospecies *Rhizophagus irregularis* and *R. clarus,* had both the highest speciation rates among AMF (Fig. 2a-b) and the largest niche widths (Supplementary Fig. 27).

We found no effect of latitude on speciation rates, regardless of the AMF delineation or the minimum number of sequences per AMF unit (MCMCglmm: $P>0.05$), and no effect of habitat or climatic zone either (Supplementary Fig. 28). Similarly, we recovered no significant correlation between spore size and speciation rate (Supplementary Fig. 29), nor between spore size and level of endemism (Supplementary Fig. 30).

Finally, Tajima's $\theta\pi$ estimator of AMF genetic diversity was significantly and positively correlated with niche width (PC1) for all AMF delineations and minimal number of sequences per AMF unit considered, and in particular with abiotic aspects of the niche (PC2) in many cases (Fig. 4e-h; Supplementary Fig. 26). Genetic diversity was not correlated with speciation rate (Supplementary Fig. 26), latitude, habitat, climatic zone (MCMCglmm: $P>0.05$), or spore size (PGLS: $P>0.05$).

Discussion

AMF species delineations, diversity, and phylogeny:

The species concept is difficult to apply in AMF, which are poorly differentiated morphologically and mainly characterized by environmental sequences [16]. In addition, their reproduction mode is not well known and they have unique nuclear dynamics in their spores and hyphae [61]. Our GMYC analyses suggest that biologically relevant AMF species-like units correspond to SSU rRNA haplotypes with a sequence similarity between 98.5 and 99%. With this criterion of species delineation, we estimate that there are between 1,300 and 2,900 AMF 'species'. These estimates are largely above the number of currently described morphospecies or VT (Supplementary Note 1) but remain low in comparison with other fungal groups, like the Agaricomycetes that include taxa forming ectomycorrhiza [62].

Species delineations and phylogenies constructed from a single gene and short sequences are limited, but in the current state of data acquisition, relatively short metabarcoding sequences provide for most microbial groups, including AMF, the only current possibility to analyze their diversification dynamics [20, 63, 64]. Here, our phylogenies did not resolve the branching of the AMF orders, with node supports similar to those of previous studies [12, 17, 20](Supplementary Note 5), confirming that additional genomic evidence is required to reach consensus. We considered this uncertainty in the phylogenetic reconstruction by repeating our analyses on a set of trees spanning the likely tree space. We hope that our study based on the SSU rRNA region alone will foster efforts to obtain more genetic data, including additional genes and genomic information, with the aim of reconstructing better supported, comprehensive phylogenies.

AMF diversify slowly:

We found speciation rates for AMF an order of magnitude lower than rates typically found for macro-eukaryotes [46, 65], like plants [29], or Agaricomycetes [62]. Low speciation rates in AMF may be linked to their (debated) asexual reproduction [66], to their occasional long-distance dispersal that homogenizes populations globally over evolutionary timescales [67], or to the fact that they are generalist obligate symbionts [68]. Regardless of the proximal cause, and contrary to Agaricomycetes for example, which present a large diversity of species, morphologies, and ecologies, the niche space exploited by AMF is limited to plant roots and the surrounding soil because of their obligate dependence on plants for more than 400 Myr [69, 70]. Thus, although AMF species delineation based on the SSU rRNA gene can be a poor predictor of their functional diversity, our analyses based on this gene has revealed that AMF, despite their ubiquity, have poorly diversified in the last 500 Myr compared with other groups.

We found little evidence for species extinction in AMF, including at mass extinction events. Although AMF are relatively widespread and generalists, and low extinction rates have been predicted before based on their ecology [2], these low extinction rate estimates could also come from the difficulty of estimating extinction from molecular phylogenies [71], one of the limitations of phylogeny-based diversification analyses (Supplementary Note 6).

AMF diversification through time:

The observed peak of AMF diversification detected between 200 and 100 Myr (or 150-50 Myr according to the models) was mainly linked to the fast diversification of the largest family Glomeraceae (Fig. 2). This peak was concomitant with the radiation of flowering plants [28], but also with a major continental reconfiguration, including the breakdown of Pangea and the formation of climatically contrasted landmasses [20]. This period was also

characterized by a warm climate potentially favorable to AMF diversification, such that disentangling the impact of these various factors on AMF diversification is not straightforward. Interestingly, a peak of diversification at this period was also found in the Agaricomycetes forming ectomycorrhiza [62].

This peak of diversification has been followed by a slowdown. Signals of diversification slowdowns sometimes result from methodological artifacts, including incorrect species delineations or under-sampling [72]. We carefully considered uncertainty in species delineations and under-sampling down to 50%. In addition, our GMYC analyses confirmed that the SSU rRNA gene evolves fast enough to delineate AMF species-like units; although some cryptic AMF species can have the same SSU sequence [12], our analyses support the overall existence of several SSU haplotypes per AMF unit. Slowdowns in diversification rates close to the present have often been interpreted as a progressive reduction of the number of available niches as species diversify and accumulate [72, 73]. In AMF, this potential effect of niche saturation could be exacerbated by a reduction of their niches linked to both repetitive breakdowns of their symbiosis with plants and climatic changes. Indeed, in the last 100 Myr, many plant lineages evolved alternative root symbioses or became non-symbiotic [5, 32, 33, 74]: approximately 20% of extant plants do not interact with AMF anymore [4]. Additionally, the cooling of the Earth during the Cenozoic reduced the surface of tropical regions [75, 76], which tend to be a reservoir of ecological niches for AMF [20, 32, 77].

The difficulty of reconstructing past symbiotic associations prevents direct testing of the hypothesis that the emergence of new root symbioses in plants led to a diversification slowdown in AMF. However, we tested the hypothesis that global temperature changes affected diversification rates and found a strong relationship (Fig. 3). Such associations between temperature and diversification rates have been observed before in eukaryotes and have several potential causes [78]. Two prevailing hypotheses are the evolutionary speed hypothesis, stipulating that high temperatures entail higher mutation rates and

faster speciation [79], and the productivity hypothesis, stating that resources and associated ecological niches are more numerous in warm and productive environments, especially when the tropics are large [80]. The latter hypothesis is particularly relevant for AMF, which have many host plant niches in the tropics and potentially less in temperate regions [81], where a higher proportion of plants are non-mycorrhizal [82] or ectomycorrhizal [32, 62]. Hence, the observed effect of past global temperatures could reflect the shrinkage of tropical areas and the associated decrease of the relative proportion of arbuscular mycorrhizal plants.

A few AMF clades displayed a significant support for diversification models with a positive dependency on CO_2 concentrations, which reinforces the idea that for the corresponding AMF, benefits retrieved from plants could have been amplified by high CO_2 concentrations and fostered diversification [83, 84]. Conversely, we found a limited effect of land plant fossil diversity, which indicates that variations in the tempo of AMF diversification did not systematically follow those of land plants. Still, the possible concordance of the peak of AMF diversification with the radiation of the Angiosperms is noteworthy, in particular in Glomeraceae that frequently interact with present-day Angiosperms [17]. The co-diversification with the plants might have been an important driver from the emergence of land plants until the Mesozoic [2, 10], but less so thereafter, when AMF diversification declined while some flowering plants radiated, including AMF-free groups such as the species-rich Orchidaceae, blurring co-diversification patterns (Supplementary Fig. 31)[30, 85].

AMF recent diversification:

Looking at the correlates of AMF present-day diversification rates, we found no effect of habitat or climatic zone, even though AMF are more frequent and diverse in the tropics (Supplementary Fig. 32)[20, 25, 81] and their speciation rates are positively correlated with global temperature. Further work, including a more thorough sampling of the distribution

of AMF species across latitudes and habitats, would be required to confirm these patterns and to distinguish whether speciation events are indeed no more frequent in the tropics or, if they are, whether long-distance dispersal redistributes the new lineages at different latitudes over evolutionary time scales [25]. Similarly, although the temporal changes in the availability of AMF niches likely influenced the diversification of the group, we found little support for AMF species with larger niche width having higher lineage-specific speciation rates (Fig. 4). We also note that there are important aspects of the niche that we do not (and yet cannot) account for in our characterization of AMF niche width: it is thought that some AMF species may mainly provide mineral nutrients extracted from the soil, whereas others may be more specialized in protecting plants from biotic or abiotic stresses [86] and such (inter- or intra-specific) functional variations may have evolutionary significance. Finally, although spore size is often inversely related to dispersal capacity [87], which can either promote diversification by favoring founder speciation events, or limit diversification by increasing gene flow, we found no significant correlation between spore size and diversification rates, which may be explained either by a weak or absent effect or by the low number of species for which this data is available. In addition, the absence of correlation between spore size and level of endemism suggests that even AMF with large spores experience long-distance dispersal [57, 88]. Thus, if large spores might limit dispersal at smaller (e.g. intra-continental) scales in AMF [22, 89], this does not seem to affect diversification.

In AMF, intraspecific variability is an important source of functional diversity [67, 90] and their genetic diversity may indicate the intraspecific variability on which selection can act, potentially leading to species diversification. Here, geographically widespread AMF species appear to be more genetically diverse, as previously suggested by population genomics [67], but do not necessarily speciate faster. Along with a decoupling between genetic diversity and lineage-specific speciation rate, this suggests that the accumulation of genetic diversity among distant subpopulations is not enough to spur AMF speciation.

Conclusion:

Our findings that AMF have low speciation rates, likely constrained by the availability of suitable niches, reinforce the vision of AMF as an "evolutionary cul-de-sac" [1]. We interpret the significant diversification slowdown in the past 100 Myr as the conjunction of the emergence of plant lineages not associated with AMF and the reduction of tropical areas induced by climate cooling, in the context of obligate dependence of AMF on plants. Diversification slowdowns have often been interpreted as the signal of adaptive radiations [72, 91], that is clades that experienced a rapid accumulation of morphological, ecological, and species diversity [92]. AMF provide a striking example of a clade with slow morphological, ecological and species diversification that features a pattern of diversification slowdown. In AMF, and potentially in many other species groups [78], such a slowdown likely reflects a reduction of the global availability of ecological niches.

References:

1. Malloch DM. The evolution of mycorrhizae. *Can J Plant Path* 1987; **9**: 398–402.

2. Morton JB. Species and clones of arbuscular mycorrhizal fungi (Glomales, Zygomycetes): their role in macro- and microevolutionary processes. *Mycotaxon (USA)* 1990; **37**: 493–515.

3. Smith SE, Read DJ. Mycorrhizal Symbiosis. *Mycorrhizal Symbiosis* . 2008. Elsevier.

4. van der Heijden MGAA, Martin FM, Selosse M-A, Sanders IR. Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytol* 2015; **205**: 1406–1423.

5. Selosse M-A, Le Tacon F. The land flora: a phototroph-fungus partnership? *Trends Ecol Evol* 1998; **13**: 15–20.

6. Field KJ, Pressel S, Duckett JG, Rimington WR, Bidartondo MI. Symbiotic options for the conquest of land. *Trends Ecol Evol* 2015; **30**: 477–486.

7. Feijen FAA, Vos RA, Nuytinck J, Merckx VSFT. Evolutionary dynamics of mycorrhizal symbiosis in land plant diversification. *Sci Rep* 2018; **8**: 10698.

8. Strullu-Derrien C, Selosse MA, Kenrick P, Martin FM. The origin and evolution of mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytol* 2018; **220**: 1012–1030.

9. Simon L, Bousquet J, Lévesque RC, Lalonde M. Origin and diversification of endomycorrhizal fungi and coincidence with vascular land plants. *Nature* 1993; **363**: 67–69.

10. Lutzoni F, Nowak MD, Alfaro ME, Reeb V, Miadlikowska J, Krug M, et al. Contemporaneous radiations of fungi and plants linked to symbiosis. *Nat Commun* 2018; **9**: 1–11.

11. Stürmer SL. A history of the taxonomy and systematics of arbuscular mycorrhizal fungi belonging to the phylum Glomeromycota. *Mycorrhiza* 2012; **22**: 247–258.

12. Krüger M, Krüger C, Walker C, Stockinger H, Schüßler A. Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from

phylum to species level. *New Phytol* 2012; **193**: 970–984.

13. Öpik M, Vanatoa A, Vanatoa E, Moora M, Davison J, Kalwij JM, et al. The online

database MaarjAM reveals global and ecosystemic distribution patterns in

arbuscular mycorrhizal fungi (Glomeromycota). *New Phytol* 2010; **188**: 223–241.

14. Öpik M, Davison J, Moora M, Zobel M. DNA-based detection and identification of

Glomeromycota: the virtual taxonomy of environmental sequences. *Botany* 2014; **92**:

135–147.

15. Powell JR, Monaghan MT, Öpik M, Rillig MC. Evolutionary criteria outperform

operational approaches in producing ecologically relevant fungal species

inventories. *Mol Ecol* 2011; **20**: 655–666.

16. Bruns TD, Corradi N, Redecker D, Taylor JW, Öpik M. Glomeromycotina: what is a

species and why should we care? *New Phytol* 2018; **220**: 963–967.

17. Rimington WR, Pressel S, Duckett JG, Field KJ, Read DJ, Bidartondo MI. Ancient

plants with ancient fungi: liverworts associate with early-diverging arbuscular

mycorrhizal fungi. *Proc R Soc B Biol Sci* 2018; **285**: 20181600.

18. James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, Cox CJ, et al.

Reconstructing the early evolution of Fungi using a six-gene phylogeny. *Nature*

2006; **443**: 818–822.

19. Venice F, Ghignone S, Salvioli di Fossalunga A, Amselem J, Novero M, Xianan X, et

al. At the nexus of three kingdoms: the genome of the mycorrhizal fungus

Gigaspora margarita provides insights into plant, endobacterial and fungal

interactions. *Environ Microbiol* 2020; **22**: 122–141.

20. Davison J, Moora M, Opik M, Adholeya A, Ainsaar L, Ba A, et al. Global

assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism.

Science (80-) 2015; **349**: 970–973.

21. Egan C, Li D-W, Klironomos J. Detection of arbuscular mycorrhizal fungal spores

in the air across different biomes and ecoregions. *Fungal Ecol* 2014; **12**: 26–31.

22. Bueno CG, Moora M. How do arbuscular mycorrhizal fungi travel? *New Phytol*

2019; **222**: 645–647.

23.   Correia M, Heleno R, da Silva LP, Costa JM, Rodríguez-Echeverría S. First evidence

for the joint dispersal of mycorrhizal fungi and plant diaspores by birds. *New Phytol*

2019; **222**: 1054–1060.

24.   Templeton AR. The reality and importance of founder speciation in evolution.

BioEssays . 2008. John Wiley & Sons, Ltd. , **30**: 470–479

25.   Pärtel M, Öpik M, Moora M, Tedersoo L, Szava-Kovats R, Rosendahl S, et al.

Historical biome distribution and recent human disturbance shape the diversity of

arbuscular mycorrhizal fungi. *New Phytol* 2017; **216**: 227–238.

26.   Perez-Lamarque B, Selosse M, Öpik M, Morlon H, Martos F. Cheating in arbuscular

mycorrhizal mutualism: a network and phylogenetic analysis of

mycoheterotrophy. *New Phytol* 2020; **226**: 1822–1835.

27.   Sanders IR. Preference, specificity and cheating in the arbuscular mycorrhizal

symbiosis. *Trends Plant Sci* 2003; **8**: 143–145.

28.   Sauquet H, Magallón S. Key questions and challenges in angiosperm

macroevolution. *New Phytol* 2018; **219**: 1170–1187.

29.   Zanne AE, Tank DC, Cornwell WK, Eastman JM, Smith SA, FitzJohn RG, et al.

Three keys to the radiation of angiosperms into freezing environments. *Nature*

2014; **506**: 89–92.

30.   Cleal CJ, Cascales-Miñana B. Composition and dynamics of the great Phanerozoic

Evolutionary Floras. *Lethaia* 2014; **47**: 469–484.

31.   Bredenkamp GJ, Spada F, Kazmierczak E. On the origin of northern and southern

hemisphere grasslands. *Plant Ecol* 2002; **163**: 209–229.

32.   Brundrett MC, Tedersoo L. Evolutionary history of mycorrhizal symbioses and

global host plant diversity. *New Phytol* 2018; **220**: 1108–1115.

33.   Werner GDA, Cornelissen JHC, Cornwell WK, Soudzilovskaia NA, Kattge J, West

SA, et al. Symbiont switching and alternative resource acquisition strategies drive

mutualism breakdown. *Proc Natl Acad Sci* 2018; **115**: 5229–5234.

34.   Werner GDA, Cornwell WK, Sprent JI, Kattge J, Kiers ET. A single evolutionary

innovation drives the deep evolution of symbiotic N2-fixation in angiosperms. *Nat*

Commun 2014; **5**: 4087.

35. Lekberg Y, Vasar M, Bullington LS, Sepp S-K, Antunes PM, Bunn R, et al. More

bang for the buck? Can arbuscular mycorrhizal fungal communities be

characterized adequately alongside other fungi using general fungal primers? *New*

Phytol 2018; **220**: 971–976.

36. Katoh K, Standley DM. MAFFT Multiple sequence alignment software version 7:

Improvements in performance and usability. *Mol Biol Evol* 2013; **30**: 772–780.

37. Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: a

software platform for Bayesian evolutionary analysis. *PLoS Comput Biol* 2014; **10**:

e1003537.

38. Simon L, Lalonde M, Bruns TD. Specific amplification of 18S fungal ribosomal

genes from vesicular-arbuscular endomycorrhizal fungi colonizing roots. *Appl*

Environ Microbiol 1992; **58**: 291–5.

39. Lee J, Lee S, Young JPW. Improved PCR primers for the detection and

identification of arbuscular mycorrhizal fungi. *FEMS Microbiol Ecol* 2008; **65**: 339–

349.

40. Morlon H, Lewitus E, Condamine FL, Manceau M, Clavel J, Drury J. RPANDA: An

R package for macroevolutionary analyses on phylogenetic trees. *Methods Ecol Evol*

2016; **7**: 589–597.

41. R Core Team. R: A language and environment for statistical computing. 2019. R

Foundation for Statistical Computing, Vienna, Austria.

42. Fujisawa T, Barraclough TG. Delimiting species using single-locus data and the

generalized mixed yule coalescent approach: A revised method and evaluation on

simulated data sets. *Syst Biol* 2013; **62**: 707–724.

43. Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, et al.

Sequence-based species delimitation for the DNA taxonomy of undescribed insects.

Syst Biol 2006; **55**: 595–609.

44. Ezard T, Fujisawa T, Barraclough TG. SPLITS: SPecies' LImits by Threshold

Statistics. 2009. , R package

45. Quince C, Curtis TP, Sloan WT. The rational exploration of microbial diversity.
ISME J 2008; **2**: 997–1006.

46. Maliet O, Hartig F, Morlon H. A model with many small shifts for estimating
species-specific diversification rates. *Nat Ecol Evol* 2019; **3**: 1086–1092.

47. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences.
Stat Sci 1992; **7**: 457–472.

48. Stadler T. Mammalian phylogeny reveals recent diversification rate shifts. *Proc Natl*
Acad Sci 2011; **108**: 6187–6192.

49. Höhna S, May MR, Moore BR. TESS: An R package for efficiently simulating
phylogenetic trees and performing Bayesian inference of lineage diversification
rates. *Bioinformatics* 2016; **32**: 789–791.

50. May MR, Höhna S, Moore BR. A Bayesian approach for detecting the impact of
mass-extinction events on molecular phylogenies when rates of lineage
diversification may vary. *Methods Ecol Evol* 2016; **7**: 947–959.

51. Condamine FL, Rolland J, Morlon H. Macroevolutionary perspectives to
environmental change. *Ecol Lett* 2013; **16**: 72–85.

52. Rabosky DL, Lovette IJ. Density-dependent diversification in North American
wood warblers. *Proc R Soc B Biol Sci* 2008; **275**: 2363–2371.

53. Royer DL, Berner RA, Montanez IP, Tabor NJ, Beerling DJ. CO2 as a primary driver
of Phanerozoic climate. *GSA Today* 2004; **14**: 4–10.

54. Foster GL, Royer DL, Lunt DJ. Future climate forcing potentially without precedent
in the last 420 million years. *Nat Commun* 2017; **8**: 14845.

55. Alroy J. Geographical, environmental and intrinsic biotic controls on Phanerozoic
marine diversification. *Palaeontology* 2010; **53**: 1211–1235.

56. Hadfield JD. MCMC methods for multi-response generalized linear mixed models:
The MCMCglmm R package. *J Stat Softw* 2010; **33**: 1–22.

57. Davison J, Moora M, Öpik M, Ainsaar L, Ducousso M, Hiiesalu I, et al. Microbial
island biogeography: isolation shapes the life history characteristics but not
diversity of root-symbiotic fungal communities. *ISME J* 2018; **12**: 2211–2224.

58.   Tajima F. Evolutionary relationship of DNA sequences in finite populations.
Genetics 1983; **105**: 437–460.

59.   Sepp S, Davison J, Jairus T, Vasar M, Moora M, Zobel M, et al. Non-random
association patterns in a plant–mycorrhizal fungal network reveal host–symbiont
specificity. *Mol Ecol* 2019; **28**: 365–378.

60.   Morlon H, Parsons TL, Plotkin JB. Reconciling molecular phylogenies with the
fossil record. *Proc Natl Acad Sci* 2011; **108**: 16327–16332.

61.   Kokkoris V, Stefani F, Dalpé Y, Dettman J, Corradi N. Nuclear dynamics in the
arbuscular mycorrhizal fungi. *Trends Plant Sci* 2020; **25**: 765–778.

62.   Varga T, Krizsán K, Földi C, Dima B, Sánchez-García M, Sánchez-Ramírez S, et al.
Megaphylogeny resolves global patterns of mushroom evolution. *Nat Ecol Evol*
2019; **3**: 668–678.

63.   Louca S, Shih PM, Pennell MW, Fischer WW, Parfrey LW, Doebeli M. Bacterial
diversification through geological time. *Nat Ecol Evol* 2018; **2**: 1458–1467.

64.   Lewitus E, Bittner L, Malviya S, Bowler C, Morlon H. Clade-specific diversification
dynamics of marine diatoms since the Jurassic. *Nat Ecol Evol* 2018; **2**: 1715–1723.

65.   Upham NS, Esselstyn JA, Jetz W. Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol* 2019;
17: e3000494.

66.   Yildirir G, Malar C M, Kokkoris V, Corradi N. Parasexual and sexual reproduction
in arbuscular mycorrhizal fungi: Room for both. *Trends Microbiol* 2020; **28**: 517–519.

67.   Savary R, Masclaux FG, Wyss T, Droh G, Cruz Corella J, Machado AP, et al. A
population genomics approach shows widespread geographical distribution of
cryptic genomic forms of the symbiotic fungus Rhizophagus irregularis. *ISME J*
2018; **12**: 17–30.

68.   Morlon H, Kemps BD, Plotkin JB, Brisson D. Explosive radiation of a bacterial
species group. *Evolution (N Y)* 2012; **66**: 2577–2586.

69.   Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, et al. Genome
of an arbuscular mycorrhizal fungus provides insight into the oldest plant

symbiosis. *Proc Natl Acad Sci U S A* 2013; **110**: 20117–22.

70.   Rich MK, Nouri E, Courty P-E, Reinhardt D. Diet of arbuscular mycorrhizal fungi:

Bread and butter? *Trends Plant Sci* 2017; **22**: 652–660.

71.   Rabosky DL. Challenges in the estimation of extinction from molecular

phylogenies: A response to Beaulieu and O'Meara. *Evolution (N Y)* 2016; **70**: 218–

228.

72.   Moen D, Morlon H. Why does diversification slow down? *Trends Ecol Evol* 2014; **29**:

190–197.

73.   Rabosky DL. Ecological limits and diversification rate: Alternative paradigms to

explain the variation in species richness among clades and regions. *Ecol Lett* 2009;

12: 735–743.

74.   Maherali H, Oberle B, Stevens PF, Cornwell WK, McGlinn DJ. Mutualism

persistence and abandonment during the evolution of the mycorrhizal symbiosis.

Am Nat 2016; **188**: E113–E125.

75.   S. Meseguer A, Condamine FL. Ancient tropical extinctions at high latitudes

contributed to the latitudinal diversity gradient. *Evolution (N Y)* 2020; **74**: 1966–

1987.

76.   Ziegler AM, Eshel G, McAllister Rees P, Rothfus TA, Rowley DB, Sunderlin D.

Tracing the tropics across land and sea: Permian to present. *Lethaia* 2003; **36**: 227–

254.

77.   Read DJ. Mycorrhizas in ecosystems. *Experientia* 1991; **47**: 376–391.

78.   Condamine FL, Rolland J, Morlon H. Assessing the causes of diversification

slowdowns: temperature-dependent and diversity-dependent models receive

equivalent support. *Ecol Lett* 2019; **22**: 1900–1912.

79.   Rohde K. Latitudinal gradients in species diversity: The search for the primary

cause. *Oikos* 1992; **65**: 514.

80.   Clarke A, Gaston KJ. Climate, energy and diversity. *Proc R Soc B Biol Sci* . 2006. The

Royal SocietyLondon. , **273**: 2257–2266

81.   Toussaint A, Bueno G, Davison J, Moora M, Tedersoo L, Zobel M, et al.

Asymmetric patterns of global diversity among plants and mycorrhizal fungi. *J Veg*
Sci 2020; **31**: 355–366.

82. Bueno CG, Moora M, Gerz M, Davison J, Öpik M, Pärtel M, et al. Plant mycorrhizal
status, but not type, shifts with latitude and elevation in Europe. *Glob Ecol Biogeogr*
2017; **26**: 690–699.

83. Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, et
al. Functional analysis of liverworts in dual symbiosis with Glomeromycota and
Mucoromycotina fungi under a simulated Palaeozoic CO2 decline. *ISME J* 2016; **10**:
1514–1526.

84. Humphreys CP, Franks PJ, Rees M, Bidartondo MI, Leake JR, Beerling DJ.
Mutualistic mycorrhiza-like symbiosis in the most ancient group of land plants. *Nat*
Commun 2010; **1**: 103.

85. Ramírez-Barahona S, Sauquet H, Magallón S. The delayed and geographically
heterogeneous diversification of flowering plant families. *Nat Ecol Evol* 2020; **4**:
1232–1238.

86. Chagnon P-L, Bradley RL, Maherali H, Klironomos JN. A trait-based framework to
understand life history of mycorrhizal fungi. *Trends Plant Sci* 2013; **18**: 484–491.

87. Nathan R, Schurr FM, Spiegel O, Steinitz O, Trakhtenbrot A, Tsoar A. Mechanisms
of long-distance seed dispersal. *Trends Ecol Evol* 2008; **23**: 638–647.

88. Kivlin SN. Global mycorrhizal fungal range sizes vary within and among
mycorrhizal guilds but are not correlated with dispersal traits. *J Biogeogr* 2020; **47**:
1994–2001.

89. Chaudhary VB, Nolimal S, Sosa-Hernández MA, Egan C, Kastens J. Trait-based
aerial dispersal of arbuscular mycorrhizal fungi. *New Phytol* 2020; **228**: 238–252.

90. Munkvold L, Kjøller R, Vestberg M, Rosendahl S, Jakobsen I. High functional
diversity within species of arbuscular mycorrhizal fungi. *New Phytol* 2004; **164**: 357–
364.

91. Harmon LJ, Schulte JA, Larson A, Losos JB. Tempo and mode of evolutionary
radiation in iguanian lizards. *Science (80-)* 2003; **301**: 961–964.

92.   Simpson GG. The major features of evolution. 1953. Columbia University Press,

745   New York.

746   93.   Janzen T, Etienne RS. Inferring the role of habitat dynamics in driving

747   diversification: evidence for a species pump in Lake Tanganyika cichlids. *bioRxiv*

748   2016; **11**: 1–18.

Acknowledgment:

The authors acknowledge C. Strullu-Derrien, M. Elias, D. de Vienne, A. Vogler, J.-Y. Dubuisson, C. Quince, S.-K. Sepp, and M. Chase for helpful discussions. They also thank L. Aristide, S. Lambert, J. Clavel, I. Quintero, I. Overcast, and G. Sommeria for comments on an early version of the manuscript and David Marsh for English editing. BPL acknowledges B. Robira, F. Foutel-Rodier, F. Duchenne, E. Faure, E. Kerdoncuff, R. Petrolli, and G. Collobert for useful discussions and C. Fruciano and E. Lewitus for providing codes. This work was supported by a doctoral fellowship from the École Normale Supérieure de Paris attributed to BPL and the École Doctorale FIRE – Programme Bettencourt. MÖ was supported by the European Regional Development Fund (Centre of Excellence EcolChange) and University of Tartu (PLTOM20903). Funding of the research of FM was from the Agence Nationale de la Recherche (ANR-19-CE02-0002). HM acknowledges support from the European Research Council (grant CoG-PANDA).

Author contributions:

All the authors designed the study. MÖ gathered the data and BPL performed the analyses. OM and ACAS provided some codes. BPL and HM wrote the first version of the manuscript and all authors contributed substantially to the revisions.

Competing Interests statement:

The authors declare that there is no conflict of interest.

Figure 1: Molecular-based species delineations of arbuscular mycorrhizal fungi (AMF) give consistent results and indicate a nearly complete sampling.

We compared the *virtual taxa* (VT) delineation from [13] with newly-developed automatic

delineations into *evolutionary units* (EUs) based on an average threshold of similarity and a

criterion of monophyly.

(a) The proportion of AMF units (VT or EUs) in each AMF family reveals constant

proportions across delineations, although Glomeraceae tend to be relatively less numerous

compared to the other AMF family in the VT delineation. The main AMF orders are

indicated on the right of the charts: Paraglomerales + Archaeosporales, Diversisporales,

and Glomerales (Glomeraceae + Claroideoglomeraceae).

(b) Rarefaction curves indicating the number of AMF units as a function of the percentage

of sampled AMF accession revealed that the AMF sampling in MaarjAM is close to

saturation for all delineations (VT or EUs). Rarefactions were performed 100 times every 5

percent and the median of the 100 replicates is represented here.

Figure 2: The diversification dynamic of arbuscular mycorrhizal fungi (AMF) varies significantly through time and between lineages.

(a-b): AMF consensus phylogenetic trees corresponding to the VT (a) and EU99 (b) species delineations. Branches are colored according to the lineage-specific speciation rates estimated by ClaDS using the BDES estimated sampling fraction: lineages with low and high speciation rates are represented in blue and red, respectively.
The main AMF clades are indicated with the following letters: P = Paraglomerales + Archaeosporales, D = Diversisporales, C = Claroideoglomeraceae, and G = Glomeraceae.

(c-d): Mean speciation rates through time estimated by ClaDS, for the VT (c) and EU99 (d) delineations and using the BDES estimated sampling fraction. The mean speciation rate corresponds to the maximum *a posteriori* (MAP) of the mean speciation rate across all fungal lineages back in time (including extinct and unsampled lineages). Orange and grey lines represent the 12 independent replicate trees and the consensus tree, respectively. Unlike most replicate trees, the EU99 consensus tree tends to present a limited diversification slowdown, which reinforces the idea that consensus trees can be a misleading representation [93].

(e-f): Net diversification rates (speciation rates minus extinction rates) through time estimated by TreePar, for the VT (c) and EU99 (d) delineations and using the BDES estimated sampling fraction. Orange and grey lines represent the 12 independent replicate trees and the consensus tree, respectively.

Figure 3: Temperature-dependent diversification models reveal that global temperature positively associates with the speciation rates of arbuscular mycorrhizal fungi (AMF) in the last 500 million years.

(a): Average global temperature in the last 500 million years (Myr) relative to the average temperature of the period 1960-1990. The smoothed orange line represents cubic splines with 33 degrees of freedom used to fit temperature-dependent models of AMF diversification with RPANDA. This default smoothing was estimated using the R function *smooth.spline*.

(b): AICc difference between the best-supported time-dependent model and the temperature-dependent model in RPANDA, for the VT (left) and EU99 (right) delineations, using the BDES estimated sampling fraction. An AICc difference greater than 2 indicates that there is significant support for the temperature-dependent model.

(c): Parameter estimations of the temperature-dependent models (speciation rate ~ exp(parameter * temperature)). A positive parameter value indicates a positive effect of temperature on speciation rates.

For both delineations, the boxplots represent the results obtained for the consensus tree and the 12 independent replicate trees. Boxplots indicate the median surrounded by the first and third quartiles, and whiskers extend to the extreme values but no further than 1.5 of the inter-quartile range. The horizontal dotted lines highlighted the values estimated for the consensus trees. Compared to the replicate trees, the consensus trees tend to present extreme values (stronger support for temperature-dependent model), which reinforces the idea that consensus trees can be a misleading representation [93].

Figure 4: Abiotic and biotic drivers of the species diversification and differentiation of arbuscular mycorrhizal fungi (AMF)

(a-b): Projection of 10 abiotic and biotic variables on the two principal coordinates according to the VT (a) or EU99 (b) delineations. Principal coordinate analysis (PCA) was performed for the AMF units represented by at least 10 sequences. Colors represent the contribution of the variable to the principal coordinates. The percentage for each principal coordinate (PC) indicates its amount of explained variance.

Tested variables were: the numbers of continents on which the AMF unit occurs (nb_continent), of realms (nb_realm), of ecosystems (nb_ecosystems), of habitats (nb_habitats), of biomes (nb_biomes), and climatic zones (nb_climatic) [13], as well as information about the associated plant species of each unit, such as the number of plant partners (nb_plants), the phylogenetic diversity of these plants (PD), and the betweenness and closeness measurement of each fungal unit in the plant-fungus interaction network (see Methods).

(c-d): Speciation rates as a function of the PC1 coordinates for each VT (c) or EU99 (d) unit. Only the AMF consensus tree is represented here (other replicate trees are presented in Supplementary Fig. 26).

(e-h): Genetic diversity (Tajima's $\theta\pi$ estimator) as a function of the PC1 (e-f) or PC2 (g-h) coordinates for each VT (e-g) or EU99 (f-h) unit. Only the AMF consensus tree is represented here (other replicate trees are presented in Supplementary Fig. 26). The grey lines indicate the statistically significant linear regression between the two variables inferred using MCMCglmm.