Increased influence of prior choices on perceptual decisions in autism

Helen Feigin[1]*, Shir Shalom-Sperber[1]*, Ditza A. Zachor[2,3] and Adam Zaidel[1+]

[1]Gonda Multidisciplinary Brain Research Center, Bar Ilan University, Ramat Gan 5290002, Israel

[2]The Autism Center/ALUT, Shamir Medical Center, Zerifin, 6093000, Israel;

[3]Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, 6139001, Israel;

*These authors contributed equally towards this study

[+]Corresponding author:
Adam Zaidel
Gonda Multidisciplinary Brain Research Center
Bar Ilan University
Ramat Gan, 5290002
Israel
Tel: +972-35317755
Email: adam.zaidel@biu.ac.il

ABSTRACT

Autism spectrum disorder (ASD) manifests sensory and perceptual atypicalities. Recent

theories suggest that these may reflect a reduced influence of prior information in ASD.

Although many studies have not found reduced long-term priors ASD, some have found

reduced adaptation to recent sensory stimuli. However, the effects of recent prior

stimuli and prior perceptual choices can counteract one-another. Here, we investigated

this using two different tasks (in two different cohorts): (i) visual location

discrimination, and (ii) multisensory (visual-vestibular) heading discrimination. We fit

that data using a logistic regression model to dissociate the specific effects of prior

stimuli and prior choices. In both tasks/cohorts perceptual decisions were biased toward

recent choices and away from prior stimuli. Notably, the 'attractive' effect of prior

choices was significantly larger in ASD, while there was no difference in the influence

of prior stimuli (in both tasks). These results challenge theories of reduced priors in

ASD, and rather suggest an increased consistency bias for perceptual decisions in ASD.

1 INTRODUCTION

Autism spectrum disorder (ASD) is primarily characterized by impairments in
social skills and by restricted and repetitive patterns of behavior (American Psychiatric
Association, 2013). Although increased or decreased responsiveness to sensory input is
also part of the current diagnostic criteria, the nature of atypical perceptual function in
ASD is not fully understood, and not part of the diagnostic criteria. Furthermore, the
connection between altered perceptual function and the core behavioral symptoms of
ASD is unclear.

Classic theories suggest that atypical perceptual function in ASD reflects a shift
in favor of processing local (or low-level) sensory details, over integration of those
details into a global (or high-level) percept. This may reflect (non-exclusively):
impaired global processing (Shah and Frith, 1983; Frith and Happe, 1994; Behrmann
et al., 2006; Booth and Happé, 2018), superior local processing (Mottron and Burack,
2001; Plaisted et al., 2003; Mottron et al., 2006), and/or a preference for detail (Plaisted
et al., 1999; Happe and Frith, 2006; Koldewyn et al., 2013). In recent years, it has been
proposed that the bias towards low-level processing in ASD, as well as other perceptual
and sensory atypicalities, such as hyper- or hypo-sensitivity to sensory stimuli (Rogers
and Ozonoff, 2005), may be explained by altered perceptual inference (Pellicano and
Burr, 2012; Friston et al., 2013; Lawson et al., 2014).

Perceptual inference has been broadly studied in typical brain function within
the Bayesian framework (Knill and Richards, 1996; Mamassian et al., 2002; Kersten
and Yuille, 2003; Knill and Pouget, 2004). The Bayesian approach provides a
principled (computational) account of perception as a process of statistical inference,
and posits that top-down expectations ('priors') are integrated together with bottom-up
sensory signals (which are inherently "noisy", i.e., variable) to improve perceptual
reliability. Although integrating 'priors' reduces sensory noise in favor of high-level
(global) perception, this might be at the expense of low-level sensory accuracy.
Accordingly, Pellicano and Burr (2012) proposed that individuals with ASD might use
priors to a lesser degree, resulting in superior local processing and altered sensory
responsiveness. However, relatively greater sensory (vs. prior) weighting might
actually be expected from Bayesian theory due to superior low-level sensory function
in ASD (Brock, 2012; Karvelis et al., 2018). Thus, whether or not priors are altered and
what aspects of perceptual inference may differ in ASD requires investigation.

Priors are complex – they reflect a compound (subjective) aggregate of an individual's beliefs about the probability of observing various events or stimuli in the world (Ma, 2019). This is influenced by context, statistics of the environment, and other factors, across broad and narrow timescales. For example, one's prior for hearing explosions might temporarily increase when watching an action movie (but return to normal when leaving the cinema). By contrast, the prior that objects fall to the ground (due to gravity) might be more long-term and constant. It is unclear therefore what aspects of priors might be affected in ASD. At minimum, these can be divided according to their recency: short-term (recent information such as context or immediately preceding events) and long-term (such as beliefs and expectations about the world that were learned throughout life).

Recent studies have found that long-term priors seem intact in ASD. The "light-from-above prior" (that light more frequently comes from above) was applied by individuals with ASD (Croydon et al., 2017), the influence of object size on perceptions of heaviness was similar in ASD (Buckingham et al., 2016), and a slow-speed prior was integrated in ASD (Noel et al., 2020) – like controls. These results indicate that sensory predictions based on statistical information gathered over long-term experience are indeed built and integrated in ASD. Also, an experiment that tested "Mooney" images (difficult to interpret pictures, comprising black and white patches) showed that prior exposure to the high-fidelity images helped individuals with ASD as much as controls (Van de Cruys et al., 2018), indicating that priors over a time-course of minutes are also used in ASD. Thus, an overall reduction of all priors in ASD does not seem likely.

By contrast, studies that found reduced adaptation to prior stimuli in ASD seem to provide evidence for a reduced impact of *recent* sensory information (in the range of seconds) on perception in ASD. For example: prolonged exposure to a specific facial identity, which typically biases subsequent perception away from that image, did so to a lesser degree in ASD (Pellicano et al., 2007). Also, adaptation to numerosity judgements (Turi et al., 2015) to loudness (Lawson et al., 2015) and to audio-visual asynchronies (Noel et al., 2017) was reduced in ASD. At face value, these results seem to support the notion of reduced use of recent prior information in ASD. However, 'adaptation' (like in these studies) most commonly refers to the phenomenon by which exposure to a stimulus repels subsequent perception away from that stimulus (often called a "negative" effect) – opposite to a Bayesian prior, which attracts perception

towards the more frequent stimuli (positive effect). Thus, while reduced adaptation in ASD may suggest a smaller impact of prior stimuli, this does not imply reduced priors. Also, adaptation is an umbrella term that includes a repertoire of effects (Kohn, 2007) – what aspects thereof are specifically altered in ASD requires investigation.

'Serial-dependence', a related topic of high recent interest, also studies the impact of recent sensory information (in the range of seconds) on subsequent perception. Specifically, it describes a positive effect by which perceptual decisions are biased toward recently experienced stimuli (Fischer and Whitney, 2014; Liberman et al., 2014; Taubert et al., 2016; Alexi et al., 2018; Liberman and Manassi, 2018) – opposite to the negative effect of adaptation. Here too initial evidence shows reduced serial-dependence in ASD (Molesworth et al., 2015; Lieder et al., 2019), suggesting a smaller impact of recent stimuli in ASD.

However, these studies of reduced serial-dependence and reduced adaptation in ASD only take into account the previous stimuli, and not the previous choices. This is important because there is mounting evidence that prior perceptual decisions (rather than sensory stimuli) elicit the positive effect of serial-dependence (Kaneko and Sakai, 2015; Talluri et al., 2018; Feigin et al., 2020). Thus the overall impact of recent sensory experience might reflect a composite effect, e.g., of repulsion away from the prior stimuli (adaptation) and attraction toward the prior choices (serial-dependence) which can, at least partially, cancel one-another (Fritsche et al., 2017; Pascucci et al., 2019). Therefore, understanding specifically which response (to recent prior stimuli or prior choices) is altered in ASD may change our understanding of these results profoundly.

The effect of prior choices is a factor that has been less considered so far in ASD. A recent study of high-level (executive) decision-making found greater consistency in the decisions of individuals with ASD, which may suggest an ASD tendency towards repetitive behavior also in decision-making (Wu et al., 2018). However, this study used a gambling task, which may be different from perceptual decision-making, and thus did not dissociate the serial effects of prior sensory from prior choice information, which needs to be tested together in a perceptual decision-making task. We hypothesized, based on the studies described above, that a positive effect of serial-dependence from prior choices may be enlarged in ASD, and that this may cancel out adaptation effects, which could appear as reduced adaptation in ASD if the effects of prior stimuli and choices are not accounted for separately.

In this study we aimed to dissociate how recent prior stimuli and choices (over a time-course of seconds) affect perceptual decisions in ASD. For this purpose, we tested the performance of individuals with ASD and controls in a visual location discrimination task designed to induce short-term biases over several trials (Feigin et al., 2020). In addition, we reanalyzed data from a previous study of multisensory (visual-vestibular) heading direction discrimination performed on a different group of participants (Zaidel et al., 2015). Both tasks comprised a two alternative forced choice (2AFC) regarding a single interval stimulus. For the location discrimination task, participants discriminated whether a visual stimulus (solid circle, presented on a screen) lay to the left or right of the screen center. For heading discrimination, participants discriminated whether a linear self-motion stimulus was leftward or rightward of straight ahead. We fit the data with a logistic regression model that separated between the effects of prior stimuli and prior choices.

Strikingly, in both studies (two different cohorts) we found a stronger effect of prior choices on perceptual decisions in ASD vs. controls. By contrast, individuals with ASD demonstrated a similar effect of adaptation to previous stimuli compared to controls (in both studies). These results challenge theories of reduced priors in ASD, and rather implicate an increased influence of prior choices in ASD. This may expose a novel aspect of repetitive behavior in ASD, reflected in perceptual decisions.

RESULTS

The main aim of this study was to investigate whether the influence of recent prior information on perceptual decision-making is altered in ASD. This was tested on two different datasets: (i) visual location discrimination (collected for this study), and (ii) multisensory visual-vestibular heading discrimination (reanalysis of previously published data, Zaidel et al., 2015). In both datasets, we found a stronger effect of recent prior information on perceptual decisions in ASD compared to controls. Moreover, we found that this effect was mediated by a greater influence of prior choices in ASD. Specifically, perceptual decisions were more likely to be the same as previous choices (an increased 'consistency' bias). Interestingly, no differences were found regarding the influence of recent sensory information.

Location discrimination

In the visual location discrimination task, the participants were required to report whether the visual stimulus (a filled circle) lay to the right or to the left of the screen center (2AFC). Stimulus locations were biased to the right or to the left for several 'prior' discriminations, followed by a single unbiased (i.e., balanced) test stimulus (Fig. 1; rightward and leftward biased trials were interleaved). We first tested the overall effect of these short-term priors by sorting the test stimuli according to their prior bias (right/left) and plotting separate psychometric functions for each prior type (per participant).

Figure 1. Location discrimination – event sequence within and across trials. All stimuli (filled circles) were discriminated according to location, i.e.: whether the circle lay to the left or to the right of the screen center (2AFC). Each trial comprised a series of stimuli and discriminations that began with a fixation point (at the screen center), followed by a series of 'prior' stimuli biased to the right (light green) or left (dark green), and then by a single (unbiased) 'test' stimulus. The black arrow marks the timecourse across trials, which interleaved trials with right and left prior types (light and dark green dashed boxes, respectively) pseudorandomly, each ending with a test stimulus (black dashed boxes). Prior circle locations were drawn from a normal distribution either biased to the right or left of the screen center (light and dark green probability distributions, P_R and P_L, respectively). Test circle locations followed an unbiased staircase procedure.

1   **Larger effect of recent priors in ASD for location discrimination**

2   Example psychometric plots for one participant with ASD and one control

3   participant (red and blue shades, left and right subplots of Figure 2A, respectively)

4   reveal that for both participants test stimulus discriminations were influenced by recent

5   priors. This is seen by a shift in the psychometric curves for rightward vs. leftward

6   biased priors (light vs. dark shades, respectively). Note that all psychometric curves are

7   plotted as a function of the test stimulus location (which was unbiased). Thus, any

8   differences between the light and dark shaded curves reflect different responses to the

9   same test stimuli (therefore shifts in the psychometric curves expose the effects of

10 recent priors).

Figure 2. Location discrimination – psychometric function shifts. (A) Psychometric curves for an example participant with ASD (left plot) and an example control participant (right plot) performing location discrimination. The psychometric curves represent the ratio of rightward choices as a function of stimulus location, sorted according to the prior type (left or right bias, represented by dark and light colors, respectively). The data (circles) were fit with cumulative Gaussian distribution functions (solid lines). Circle size reflects the number of trials for a given stimulus location and prior type. (B) PSE shifts (ΔPSE = PSE for left biased priors minus PSE for right biased prior). Red and blue bars represent ΔPSEs (mean ± SEM) for participants with ASD and controls, respectively. *$p < 0.05$.

11   To understand the direction of the shift – it is easiest to compare the

12 psychometric curves where they cross x = 0° (vertical dotted line). For both participants

13 the lighter shaded curve (right biased priors) crosses at a higher value vs. the darker

14 curve (left biased priors). This means that there is a higher probability for choosing

15 right following rightward biased priors (and a higher probability for choosing left

16 following leftward biased priors). Another way to view and to quantify this is by the

17 difference between the curves' PSEs (where they cross the horizontal dotted line y =

18 0.5). A larger PSE difference (ΔPSE, see Eq. 4 in the Methods) indicates a stronger

19 effect, and positive ΔPSE values (like these examples) means that the participant's

choices were biased in the direction of the recent priors. In these examples, the ΔPSE

was larger for the participant with ASD (difference between the red shaded curves)

versus the control participant (difference between the blue shaded curves).

At the group level, the ΔPSEs were highly significant for the ASD group (red

bar, Fig. 2B; $t_{17} = 5.2$, $p = 7 \cdot 10^{-5}$, t-test), and smaller, but still (albeit more marginally)

significant for the control group (blue bar, Fig. 2B; $t_{19} = 2.1$, $p = 0.048$; t-test). Notably,

the ΔPSEs were significantly larger for the ASD vs. control groups ($t_{36} = 2.1$, $p = 0.043$,

t-test). Hence, not only was a significant effect of recent priors seen also in ASD, it was

larger and more significant than that of controls. This result (and the results presented

below) does not result from a difference in response times between the groups, because

these were similar: mean $\pm$ $SD = 0.69 \pm 0.15$s for ASD and 0.77 ± 0.17s for controls;

$t_{36} = -1.5$, $p = 0.14$, t-test).

Figure 3. Perceptual decision model schematic. Perceptual choices were modeled using a logistic regression with four factors (providing their respective fitted beta coefficients): (i) the current stimulus ($\beta_{curr\_stimulus}$), (ii) previous stimuli ($\beta_{prev\_stimuli}$), (iii) previous choices ($\beta_{prev\_choices}$) and (iv) baseline bias (β_0). Black stems on the stimulus and choice axes depict the stimuli (s) and choices (c) for an example trial (indexed by time step), with a test stimulus at time t (black dashed box) and prior stimuli and choices n steps back (light green dashed boxes). The sum of the product of the factors with their respective beta coefficients (z) is passed through a logistic function to yield the probability of making a rightward choice in response to the current (test) stimulus at time t.

Enhanced influence of recent choices in ASD for location discrimination

The above (ΔPSE) analysis suggests that individuals with ASD have a larger influence of recent priors. However, it does not address the underlying reason – specifically, it does not dissociate the different influences of prior stimuli from prior choices, which may be different. Recent studies have shown that the influence of recent prior stimuli and choices might be in opposite directions – i.e., the effects or prior stimuli could be adaptive (reflected by a negative shift) whereas recent choices may lead to a consistency bias (reflected by a postive shift; Kaneko and Sakai, 2015; Talluri et al., 2018; Pascucci et al., 2019; Feigin et al., 2020). Thus, the effects of prior stimuli and choices could cancel each other out. Furthermore, a larger ΔPSE shift in ASD could result from a weaker stimulus adaptation effect, or a stronger influence of prior choices (or a combination thereof). Hence, conclusions cannot be drawn from the ΔPSE results alone.

To address this, we fit a logistic regression model (per participant) that predicted each participant's discriminations of test stimuli, based on four factors: the current stimulus, previous stimuli, previous choices and a general subjective baseline bias (Fig. 3). The fourth regression coefficient (baseline bias) was not part of the group comparison because this analysis aimed to investigate the difference in priors, and there was no expected difference in baseline bias (it was however, still needed for the model fits in order to better estimate the other parameters). We confirmed that the baseline coefficients (β_0) were indeed not significantly different between the groups ($t_{36} = -1.46$, $p = 0.15$). A comparison of the other three regression coefficients revealed a significant difference between the ASD and control groups (one-way MANOVA; $F_{3,34} = 3.314$, $p = 0.03$).

Further investigation into this group difference, using pairwise comparisons, found that the influence of previous choices ($\beta_{prev\_choices}$ coefficients) was significantly larger in ASD vs. controls (Fig. 4, center plot; $t_{36} = 2.5$, $p = 0.017$). Thus, while both groups were influenced by their previous choices in a 'positive' (attractive) manner ($\beta_{prev\_choices}$ coefficients were positive and significant for each group individually; $t_{17} = 9.9$, $p = 2 \cdot 10^{-8}$ and $t_{19} = 6.4$, $p = 4 \cdot 10^{-6}$ for ASD and controls, respectively) this effect was significantly increased in ASD. This not only negates hypotheses of reduced priors in ASD – it shows the opposite – an increased influence of prior choices in ASD.

Figure 4. Location discrimination – model parameters. Beta coefficients for model parameters: current stimulus, previous choices and previous stimuli (left, center and right plots, respectively). Red and blue bars represent the values (mean ± SEM) for participants with ASD and controls, respectively. *$p = 0.017$, n.s. – not significant.

By contrast, the influence of previous stimuli ($\beta_{prev\_stimuli}$ coefficients) was small (Fig. 4, right plot, note the difference in y-scale vs. the other plots; to enable comparison between choice and stimulus coefficients, stimulus values were normalized before model fitting to have RMS = 1, like choice). For both groups, the influence of previous stimuli tended to be negative, suggesting an adaptive effect. This was significant for the control group ($t_{19} = -2.5$, $p = 0.021$) and close to significant for the ASD group ($t_{17} = -1.99$, $p = 0.063$). There was no significant difference between groups (Fig. 4; $t_{36} = 0.01$, $p = 0.99$).

Lastly, as expected, the current stimulus coefficients ($\beta_{curr\_stimulus}$) were large, positive and significant for both groups (Fig. 4, left; $t_{17} = 9.2$, $p = 5 \cdot 10^{-8}$ and $t_{19} = 8.1$, $p = 1.5 \cdot 10^{-7}$ for the ASD and control groups respectively). Also, these did not differ significantly between groups ($t_{36} = -0.85$, $p = 0.4$). Hence, the only difference we found in ASD was an increased influence of previous choices.

Comparison to clinical scores

To test whether the increased influence of prior choices in ASD reflects clinical scores of repetitiveness and inflexibility, we correlated the $\beta_{prev\_choices}$ coefficients and ΔPSE values with the ADOS and ADI-R sub-scores (Table 1). There was a trend for a relationship between the overall effect of priors (ΔPSEs) and the ADI-Communication sub-scores, however, this did not survive correction for multiple (six) comparisons ($p = 0.023$ before correction). Correlations with the $\beta_{prev\_choices}$ coefficients were not significant ($p > 0.05$ before correction of multiple comparisons). A larger dataset in the future might provide more informative correlation results.

Heading discrimination

Following the location discrimination experiment, we aimed to test whether the results would generalize to a different task, and whether it could be replicated on a different cohort. For this, we reanalyzed data from a previously published experiment that tested self-motion perception and multisensory integration of visual and vestibular cues in ASD (Zaidel et al., 2015). The task in this experiment was heading discrimination – also a 2AFC: participants were required to discriminate whether their self-motion was to the right or left of straight-ahead. Accordingly, the data and analyses were similar. However, the heading discrimination experiment was not originally designed to test the effects of recent priors (and therefore there was no apriori biasing of previous stimuli). Hence, we defined priors here simply by the preceding trial.

Larger effect of recent priors in ASD for heading discrimination

To calculate ΔPSEs, we constructed psychometric curves conditioned on the previous trial. Based on our results from the location discrimination experiment (that recent choices drive the effect) the leftward or rightward prior type for each trial was defined according to the choice on the previous trial. We first present the ΔPSE results here, and in the following section (below) we present fits to the logistic regression model.

Psychometric curves for an example ASD participant and an example control participant (visual condition) are presented in Figure 5A (red and blue shaded curves, in the left and right plots, respectively). Here too, the psychometric curves were shifted, and in the same (positive) direction as the location discrimination results: namely, following rightward choices, test stimuli were more likely to be discriminated rightward, and following leftward choices, test stimuli were more likely to be discriminated leftward. Like above, in the examples here the shift for the ASD participant was larger (red colors) vs. the control participant (blue colors).

We calculated ΔPSEs for each participant, per condition: visual, vestibular and combined visual-vestibular cues (Fig. 5B, left, right and center plots, respectively). Since the data comprised several blocks with different levels of visual coherence, ΔPSEs were calculated per coherence (presented along the x-axis). Also in these data the ΔPSEs were consistently larger for the ASD group vs. controls (the red lines lie above the blue lines in Fig. 5B). This difference was statistically significant ($F_{1,705} =$

1 6.1, $p = 0.014$; two-way ANOVA). These results suggest that our findings of a greater

2 influence of prior choices in ASD generalize across more than one task and cue. But

3 once again, further analysis is required in order to dissociate the effects of recent stimuli

4 and choices.

Figure 5. Heading discrimination – psychometric function shifts. (A) Psychometric curves for an example participant with ASD (left plot) and an example control participant (right plot) performing heading discrimination (visual condition, with 50% coherence). The psychometric curves represent the ratio of rightward choices as a function of heading, sorted according to the prior type (left or right bias, represented by dark and light colors, respectively). The data (circles) were fit with cumulative Gaussian distribution functions (solid lines). Circle size reflects the number of trials for a given stimulus heading and prior type. (B) PSE shifts (ΔPSE = PSE for left biased priors minus PSE for right biased prior) for three stimulus conditions: visual, vestibular and combined cues (left, right and center plots, respectively). Red and blue lines represent ΔPSEs (mean ± SEM) as a function of visual coherence for participants with ASD and controls, respectively. Overall, PSE shifts are significantly higher in ASD than in controls ($p = 0.014$).

5 **Enhanced influence of recent choices in ASD for heading discrimination**

6   To separate the influence of previous stimuli from previous choices, we fit the

7 data using a logistic regression model, similar to location discrimination results

8 presented above (Fig. 3). But, here only one 'prior' was used ($n = 1$) because there was

no consistent bias (by design) beyond the previous trial. In line with the location

discrimination results, the heading discrimination model fits revealed a significant

difference between ASD and control groups in the weight given to the previous choice

($\beta_{prev\_choices}$; Fig. 6, center; $F_{1,698} = 12.81$, $p = 4 \cdot 10^{-4}$; mixed two-way ANOVA, with

condition as a second factor). The $\beta_{prev\_choices}$ coefficients were significantly positive for

the ASD group ($t_{215} = 4.69$, $p = 5 \cdot 10^{-6}$), indicating that perceptual choices were attracted

towards (i.e., more likely to be similar to) the previous choice. By contrast, the

$\beta_{prev\_choices}$ coefficients were not significant in the control group ($t_{501} = -1.49$, $p = 0.136$).

Thus, here too, individuals with ASD demonstrated a greater influence of prior choices.

Figure 6. Heading discrimination – model parameters. Beta coefficients for three model parameters: current stimulus, previous choice and previous stimulus (left, center and right plots, respectively). These were averaged over three stimulus conditions (vestibular, visual and combined) and five visual coherence levels. Red and blue bars represent the beta coefficients (mean ± SEM) for participants with ASD and controls, respectively. ***$p < 0.001$, n.s. – not significant.

The previous stimulus ($\beta_{prev\_stimuli}$) coefficients were significantly negative for

both the ASD and control groups (Fig. 6, right plot; $t_{222} = -8.4$, $p = 4 \cdot 10^{-15}$ and $t_{501} = -$

10.4, $p < 10^{-16}$, respectively). This indicates a significant (negative) effect of adaptation

to previous stimuli in the heading discrimination experiment. Accordingly, adaptation

seemed to be stronger in heading discrimination vs. location discrimination, which

showed a similar trend but with marginal statistics (possibly due to the duration of the

stimuli: 1s for heading stimuli and 0.25s for location stimuli). Notably, here too there

was no significant difference between the $\beta_{prev\_stimuli}$ coefficients for the ASD and

control groups ($t_{723} = -1.35$, $p = 0.177$). Thus, we found no evidence for reduced

adaptation in ASD. Rather, we found an increased (positive) influence of prior choices.

Since these effects can be in opposite directions, if they aren't dissociated, a greater

influence of prior choices could (mistakenly) be interpreted as reduced adaptation. For

example, in our vestibular data the overall (ΔPSE) shifts lie closer to zero for ASD vs.

controls (Figure 5B, right plot). However, this does not reflect reduced adaptation, but

rather a greater influence of prior choices in ASD.

Naturally, current stimulus coefficients ($\beta_{curr\_stimulus}$) were large, positive, and

significant in both ASD and control groups (Fig. 6, left; $t_{216} = 21.7$, $p < 10^{-16}$ and $t_{501} =$

35.4, $p < 10^{-16}$, respectively), indicating that, like in the location discrimination task,

the current stimulus was still the most crucial factor in the perceptual decision.

Therefore, the increased influence of prior choices in ASD does not simply reflect a

case of blind repetition.

Effect of recent priors in ASD ensues even when reported using different actions

In order to dissociate the effect of recent choices from motor repetition, an

additional location discrimination condition was run – the *Response Invariant*

condition. In this condition, different motor actions were used when responding to

'prior' vs. 'test' stimuli (identified by different colors, see *Methods* for more details).

Also in this condition ΔPSE values were significantly positive for both the control and

ASD groups individually ($t_{15} = 2.65$, $p = 0.018$ and $t_{15} = 2.68$, $p = 0.017$, respectively,

t-tests; Fig. 7A). However, there was no significant difference between the groups (t_{30}

$= 0.91$, $p = 0.93$, *t*-test; Fig. 7A).

Figure 7. Results for the *Response Invariant* location discrimination condition.
(A) PSE shifts (ΔPSE = PSE for left biased priors minus PSE for right biased prior).
Red and blue bars represent ΔPSEs (mean ± SEM) for participants with ASD and
controls, respectively. (B) Beta coefficients for three model parameters: current
stimulus, previous choices and previous stimuli (left, center and right plots,
respectively). Red and blue bars represent the beta coefficients (mean ± SEM) for
participants with ASD and controls, respectively. n.s. – not significant.

When comparing the logistic regression coefficients (like above for the primary condition) we found no significant difference between the groups (Fig. 7B; one-way MANOVA; $F_{3,28} = 1.18$, $p = 0.34$). For both the ASD and control groups there was a positive effect of previous choices ($t_{15} = 3.97$, $p = 0.001$ and $t_{15} = 6.73$, $p = 6.8 \cdot 10^{-6}$, respectively; Fig. 7B, center plot) while the effect of previous stimuli was small and not significant ($t_{15} = -1.24$, $p = 0.23$ and $t_{15} = 0.25$, $p = 0.8$, respectively; Fig. 7B, right plot). Similar to the primary condition, the current stimulus effect was large, positive and significant for both groups (Fig. 7B, left plot; $t_{15} = 16.26$, $p = 6.2 \cdot 10^{-11}$ and $t_{15} = 10.74$, $p = 1.9 \cdot 10^{-8}$, respectively). Here too, the baseline coefficients did not differ significantly between the groups ($t_{30} = 1.29$, $p = 0.2$). These results indicate that the effect of prior choices remained for both ASD and control groups, even when the prior and current perceptual decisions differed in terms of the the motor action used to report responses (and changes in stimulus color). Here too, response times did not differ between the two groups (mean $\pm$ SD = 1.18 ± 0.33s and 1.18 ± 0.38s, for ASD and controls respectively; $t_{30} = -0.48$, $p = 0.96$, t-test).

These results show that the positive effect of recent prior choices ensues even when context (stimulus color) and the motor actions used to report responses are altered – also for individuals with ASD. This strengthens the argument against a reduced effect of recent priors in ASD. Taking the results of the two location discrimination conditions (and heading discrimination) together, we conclude that individuals with ASD experience a positive bias from recent choices, like controls. When context is consistent (i.e. stimulus color is the same, and the same motor action is used to report choices) this effect is enlarged in ASD vs. controls. However, when context is changed (and perception-action coupling is broken) this effect is no longer enlarged in ASD (but is nonetheless still present).

DISCUSSION

In this study, we found that when making perceptual decisions, individuals with ASD are more greatly influenced (vs. controls) by their recent prior choices. For all participants (ASD and control) perceptual decisions were 'attracted' towards (i.e., more likely to be the same as) recent prior choices. However, this effect was significantly larger in ASD. This finding was replicated in two different tasks (visual localization

and visual/vestibular heading discrimination) performed by two different cohorts – one in Israel and one in the USA. These results are in line with a recent study that showed increased consistency in gambling choices in ASD (Wu et al., 2018). Here, we extend this to perceptual decision-making, and dissociated the effects of prior choices and prior stimulus information.

By contrast, the influence of recent prior sensory information was similar in ASD and controls (in both tasks). This reflected a 'repulsive' bias (away from the preceding stimuli) which was significant for both groups in the heading discrimination task (similar, but not significant in the location discrimination task). Notably, this effect (adaptation to prior stimuli) was in the opposite direction vs. the effect of prior choices. Opposing effects of prior sensory vs. prior choices on perceptual decisions has been shown in typically developed individuals (Kaneko and Sakai, 2015; Talluri et al., 2018; Pascucci et al., 2019; Feigin et al., 2020). However, this has not yet been studied in ASD. Importantly, these effects need to be dissociated in order to better understand suggestions of reduced adaptation in ASD (Pellicano et al., 2007; Lawson et al., 2015; Molesworth et al., 2015; Turi et al., 2015; Noel et al., 2017; Lieder et al., 2019)

When separating the effects of prior choices and prior stimuli, we did not find reduced adaptation to prior stimuli in ASD. Our results therefore suggest that observations of "reduced adaptation" in ASD might (at least partially) be explained by an *increased* positive (i.e., counteracting) effect of prior choices. However, our stimuli were of relatively short duration (250ms in the location discrimination task and 1s in the heading direction discrimination task). Longer stimuli might lead to greater adaptation (Pegors et al., 2015) which might better expose any differences. Thus, future work should test whether adaptation to longer duration stimuli (several seconds) is altered in ASD, while taking into account the effects of prior choices.

The finding of an increased influence of prior choices in ASD may suggest that repetitiveness and inflexibility symptoms might manifest also in perceptual decisions. However, we did not find a significant correlation between the beta coefficients for prior choices and ADOS and ADI-R sub-scores of repetitiveness and inflexibility. Accordingly, our findings might offer an additional measure of repetitive symptoms, which is perhaps not covered by these sub-scores (but this would need a larger data set to confirm). Although there was a trend between the overall effect of priors (ΔPSEs) and the ADI-Communication sub-scores, this did not survive correction for multiple

comparisons. Therefore, whether or not an increased influence of prior choices correlates with ASD symptoms warrants further investigation with a larger dataset in the future.

The *Response Invariant* condition tested the effect of prior choices when context was altered – choices were reported via different motor actions (signaled by different colored stimuli). Also here, individuals with ASD still showed a significant influence of prior choices. This challenges the theories of attenuated priors in ASD (Pellicano and Burr, 2012) and demonstrates that even when context is different, individuals with ASD nonetheless still demonstrate a robust influence of priors (specifically prior choices). However, in this condition, the effect was no longer larger than controls. The fact that an enlarged influence of prior choices was seen in ASD only in the primary condition (and in the heading discrimination condition) when motor responses to prior and test stimuli were the same, suggests that i may be mediated by altered perception-action coupling (Buckingham et al., 2016).

It has been proposed that individuals with high-functioning ASD have an increased ability and tendency to recognize repeating patterns in stimuli and to generate rules for prediction (Baron-Cohen, 2002, 2009). Our results suggest that an elaborate set of systematic, internal rules in ASD uses information from prior choices to a greater degree. Furthermore, consistency (in context and in the motor responses) might be an important factor for generating prediction rules in ASD.

In summary, we found here that individuals with ASD demonstrated an increased influence of recent prior choices on perceptual decisions (vs. controls), while the influence of recent stimulus history was unaltered. This was seen in two different tasks using different modalities (visual localization and visual/vestibular heading discrimination), tested in two different cohorts (Israel and USA). These results suggest that perceptual decisions in ASD are more repetitive and less flexible (i.e., more consistent with prior decisions).

METHODS

We studied serial-dependence of perceptual decisions in individuals with ASD vs. controls performing two different tasks: (i) visual location discrimination (designed for this study and performed in Israel), (ii) multisensory (visual-vestibular) heading

discrimination (reanalysis of previously published data, collected in the USA; Zaidel et al., 2015). We present below participant and task details of the former (visual location discrimination), and briefly summarize relevant details of the latter (heading discrimination), for which full details are available in the original publication.

Participants

The visual location discrimination task was performed by 23 children and adolescents with ASD and 27 age-matched controls (all male; age range: 8-17 years). This study was approved by the institutional Helsinki committee at The Shamir Medical Center and the internal review board at Bar-Ilan University. All participants (and one of their caregivers) signed informed consent. Participants with ASD were recruited trough The Autism Center/ALUT at The Shamir Medical Center in Israel, and were all high functioning and cooperative. The ASD group was selected from a population that underwent a comprehensive evaluation for (and diagnosis of) ASD at a tertiary autism center. The criteria for selection of the ASD participants included having verbal and nonverbal scores equal to or above 80 points on the Wechsler Intelligence Scale for Children – Fourth Edition (WISC-IV; Wechsler, 2010).

The behavioral assessment of ASD was based on two standardized tests, the Autism Diagnostic Interview-Revised ADI-R (Lord et al., 1994; Rutter et al., 2003b) and the Autism Diagnostic Observation Schedule (ADOS-II; Lord et al., 2012), and a clinical judgement based on the DSM-IV criteria (American Psychiatric Association, 2000). All participants (including controls) were screened at enrollment with the Social Communication Questionnaire (SCQ) current version (Rutter et al., 2003a).

To verify that the ASD and control groups were matched in terms of intellectual functioning, the control group was tested using two verbal (vocabulary, similarities) and two nonverbal (block-design, matrix-reasoning) WISC-IV subtests (these scores were available for the ASD group from previous assessment). Performance of the ASD and control groups for the four subtests did not differ significantly ($p > 0.12$ for all four pairwise comparisons). Therefore, there is no reason to believe that any task performance differences between the groups resulted from differences in IQ. All participants had normal or corrected-to-normal vision.

One of the participants from the control group and two participants from the ASD group did not complete the task, and were therefore excluded. The data of the

remaining participants were pre-screened to confirm task understanding and cooperation (see *Data Analysis* section below for details). This removed an additional five ASD participants and four control participants, leaving 18 participants in the ASD group and 20 participants in the control group for further analysis. Tables 1 and 2 present the ages, IQ and SCQ scores for the ASD and control groups, respectively, with clinical scores (for the ASD group) also presented in Table 1.

The heading discrimination experiment tested 14 adolescents with ASD and 22 age-matched controls (all male; ages 13–19). It was approved by the institutional review board at Baylor College of Medicine, where the study was performed. Further details regarding the cohort and task are available in the original publication (Zaidel et al., 2015).

Location discrimination task - primary condition

The location discrimination experiments were run in a dark, quiet room at the Gonda Multidisciplinary Brain Research Center, at Bar-Ilan University. Participants were seated comfortably in front of an LCD monitor (resolution 1920×1200 pixels, refresh rate 60 Hz) and their head position fixed by a chin-rest. The screen was positioned such that the center of the screen lay 50 cm directly in front of the participant's eyes. The stimuli were filled circles (6.36° visual angle diameter), generated and presented using PsychoPy software (Peirce, 2007) on a dark gray background. Circle locations were fixed in the vertical plane (halfway up the screen) and varied in the horizontal plane only (right/left). At the screen center the stimulus (s) was defined as $s = 0°$ such that circle locations positioned to the right and left of the screen center had positive ($s > 0°$) and negative ($s < 0°$) values, respectively. As the magnitude of s decreases (approaches zero), the stimulus becomes more difficult to discriminate.

The participants performed a two-alternative forced choice (2AFC) task in which they were required to discriminate the location of the circle stimulus (left or right relative to the screen's center) by pressing the corresponding arrow key (right or left arrow key, respectively) on a standard computer keyboard. They were instructed to make their choice as accurately and as quickly as possible. They were informed that task difficulty varies (according to the distance of the circle from the screen center) and instructed to make their best guess when in doubt.

The stimulus locations followed a structured and balanced paradigm that we have recently developed to induce and test short-term biases over several (prior) stimuli (Feigin et al., 2020) as follows: each trial comprised of a series of circle stimuli, starting with a random number of biased 'prior' circles, followed by a single (unbiased) 'test' stimulus. The number of prior circles (n) was drawn from a discrete normal distribution of natural numbers limited to the range 1-9 (with mean $\mu = 5$, and standard deviation $\sigma = 2$). On a given trial, the locations of the prior circles were drawn randomly from a normal distribution, biased either to the left ($\mu = -3.82$, $\sigma = 3.82°$) or to the right ($\mu = 3.82$, $\sigma = 3.82°$). Trials with leftward or rightward biased priors were interleaved. The priors on a given trial were biased in one direction (on average), such that they could still appear to the contralateral side, but with lower probability (~16%).

Test circle locations were balanced to the left/right, with the location sign randomly selected on each trial ($p = 0.5$ positive/negative). Test location magnitude (absolute distance from 0°) was set using a staircase procedure (Cornsweet, 1962). The starting location magnitude was $|s| = 6.36°$ and followed a "one-up, two-down" staircase rule: after a single incorrect response, the location magnitude $|s|$ increased (i.e., became easier), and after two successive correct responses, the magnitude decreased (i.e., became harder). This staircase rule converges to ~70.7% correct performance (Leek, 2001). Staircase step size followed logarithmic increments, where location magnitude $|s|$ was multiplied by 0.6 to increase task difficulty and divided by 0.6 to decrease task difficulty. Location magnitude was limited to a maximum of $|s| = 25.46°$, but this was never reached in practice. Separate staircase procedures were run for each prior type (rightward/leftward biased, each comprising 100 trials) and pseudo randomly interleaved.

Using this paradigm we were able to examine the short-term effects of the priors' bias on subsequent perceptional decisions, in a controlled and interleaved manner. The event sequence within a trial (and across trials) is presented in Figure 1. At the beginning of each trial, a fixation cross was presented at the center of the screen for 500 ms, with the intention of reducing possible carry-over effects from the previous trial. The fixation cross was not presented again during the rest of the trial in order not to interfere with the prior's effect being measured. Each circle (prior/test) was presented for 250 ms, and was preceded by a 250 ms delay (blank screen) such that the next circle appeared 250 ms after the participant's response. Testing both prior types (right and

left, 200 trials together) took ~20 minutes. A break was automatically given after 100 trials, and the experiment was resumed by the participant, when ready.

Location discrimination - *Response Invariant* condition

Thirty participants (16 ASD and 14 controls) from those who partook in the primary location discrimination experiment (described above) agreed to perform a second location discrimination task – the *Response Invariant* condition, right after the first (a break was given in between the two tasks). This condition was designed to dissociate the effect of previous choices from previous motor responses on subsequent perceptual decisions. The stimuli and procedures were the same as before, except: (i) prior circles were colored (either green or purple) while test circles remained white (all priors and test stimuli had equal brightness). (ii) The choices for prior (colored) circles were reported using a different set of keys from the test (white) circles. Participants were instructed to use the 'up' and 'down' arrow keys to report right and left choices for colored circles (counterbalanced across participants), and the 'right' and 'left' arrow keys for white circles (like before). The color was randomly chosen for each prior stimulus (50% green and 50% purple). Only distinction between colored vs. white circles was required (no distinction was needed between purple and green circles), but we still used two colors in order to be consistent with our previous study in which different colors were used because of other manipulations (Feigin et al., 2020).

Heading discrimination task

In order to investigate whether the effects of recent biased priors, revealed by our location discrimination task, generalize across other tasks, we reanalyzed data from a previous study of heading discrimination in ASD. Extended details of this experiment are found in the original publication (Zaidel et al., 2015). In brief: participants were seated on a motion platform, and viewed a screen (also mounted on the platform) through 3D glasses. They were presented with self-motion stimuli that were either vestibular-only (inertial motion of the motion platform, in darkness), visual-only (optic-flow simulating self-motion of the participant through a cloud of "stars", while the motion platform remained stationary), or combined vestibular and visual cues (inertial motion in conjunction with synchronized optic flow). The self-motion stimuli were primarily in a forward heading direction (single-interval), with slight deviations to the

right or left of straight ahead. The participants' task was to discriminate whether the heading direction was to the right or left of straight ahead (2AFC).

Visual cue reliability was varied by manipulating the motion coherence of the optic-flow pattern (the percentage of stars moving coherently). Vestibular reliability was fixed throughout the experiment. Five levels of visual coherence were tested, in separate blocks: 100%, 90%, 75%, 50%, and 0%. One coherence was tested per block. Visual, vestibular, and combined cue conditions were interleaved, each with their own staircase. For 0% coherence the visual only condition was not tested (only vestibular and combined cue conditions were tested) because 0% coherence represents complete visual noise.

This experiment was not originally designed to assess the effects of priors, and thus test stimuli were not preceded by a series of biased priors. So, for these data each trial was analyzed as a 'test' stimulus, conditioned on the previous trial (which served as its 'prior'). To analyze psychometric shifts due to priors (see below), the trials were sorted based on the choice of the previous trial. In the subsequent model analysis (see below), the effects of both prior stimuli and choices were accounted for.

Data analysis

Data analysis was similar for the location discrimination and heading discrimination data, and performed with custom software using Matlab (The MathWorks). Psychometric functions were defined by the proportion of rightward choices as a function of stimulus *s* (circle location or heading direction) with positive and negative values representing stimuli to the right and left, respectively, and $s = 0°$ marking straight ahead. Psychometric functions were calculated by fitting the data (per participant, condition) with the following function:

$$\varphi(s) = \lambda + (1 - 2\lambda)F(s; \mu, \sigma) \qquad \text{(Equation 1)},$$

where *F(s)* is a cumulative Gaussian distribution (with mean μ and standard deviation σ) and λ represents the lapse rate. The mean (μ) reflects the point of subjective equality (PSE; the stimulus level with equal probability for a right/left choice), and we define the psychophysical threshold by the standard deviation (σ). Smaller thresholds (smaller σ) reflect better performance.

The goodness-of-fit of the psychometric functions was evaluated using the Likelihood-ratio based *pseudo-R-squared* (Hosmer and Lemeshow, 1989; Menard,

2000), which was calculated by the proportional reduction in the deviance of the fitted

psychometric model (D_{fitted}) compared to that of the null model (D_{null}):

$$R^2{}_L = 1 - \frac{D_{fitted}}{D_{null}} \qquad \text{(Equation 2)}.$$

Only participants whose psychometric curves all produced $R^2{}_L > 0.5$ were included the

analysis. Psychometric functions and deviances were calculated using the *psignifit*

toolbox for Matlab (Version 4) (Schütt et al., 2016).

The aggregate effect of priors on subsequent discriminations was assessed by

calculating separate psychometric functions for the test stimuli preceded by left and

right biased priors and calculating the difference in PSE, as follows:

$$\Delta PSE = PSE_{left} - PSE_{right} \qquad \text{(Equation 3)},$$

where the sign of ΔPSE corresponds the direction of the priors' effect. Positive ΔPSE

values indicate an attractive effect (perceptual choices are more likely to be to the same

side as the prior bias) and negative values indicate a repulsive effect (perceptual choices

are more likely to be to the opposite side of the prior bias).

Perceptual decision model fit

While the measure of ΔPSE (Eq. 3) quantifies the aggregate influence of the

prior discriminations, it does not dissociate the effect of the prior stimuli from that of

the prior choices. Moreover, these can have opposing effects and therefore (at least

partially) cancel each other out. To address this, we fit a binomial logistic regression

model (per participant and condition) that predicts a participant's choices based on four

factors: (i) the current stimulus, (ii) previous stimuli, (iii) previous choices, and (iv) the

participant's baseline bias to choose right or left (see Figure 3 for a schematic of the

model). Specifically, the logistic regression model estimates the probability of making

a rightward choice (p_t) for the current stimulus (at time t) as follows:

$$p_t = \frac{1}{1+e^{-z_t}} \qquad \text{(Equation 4)},$$

where z_t is a linear combination of the factors:

$$z_t = \beta_0 + \beta_{curr\_stimulus} \cdot s_t + \beta_{prev\_stimuli} \cdot \frac{1}{n} \sum_{i=t-n}^{t-1} s_i + \beta_{prev\_choices} \cdot \frac{1}{n} \sum_{i=t-n}^{t-1} c_i$$

$$\text{(Equation 5)}.$$

Here β_0 represents the participant's baseline bias, s_t is the current stimulus (at time t) and $\beta_{curr\_stimulus}$ its fitted weight. The other two terms account for the effects of previous stimuli and choices. Namely, $\frac{1}{n}\sum_{i=t-n}^{t-1} s_i$ is the average stimulus from the n previous discriminations (and $\beta_{prev\_stimuli}$ its fitted weight) and $\frac{1}{n}\sum_{i=t-n}^{t-1} c_i$ is the rightward choice proportion from the n previous discriminations (and $\beta_{prev\_choices}$ its fitted weight). For the location discrimination data, responses to the test stimuli were fit using all the n priors (per trial). For the heading discrimination data only the previous discrimination was used as a prior ($n = 1$).

Choices were binary (right/left; coded by '1' and '-1', respectively), while stimulus values (circle locations or heading directions) were continuous (with negative and positive values for stimuli to the left/right, respectively). In order to allow for better comparison between the logistic regression weights, stimulus intensities were normalized (before model fitting) by the root-mean-square (RMS) of all the actual stimulus values presented in the task. That way, stimulus and choice factors were of comparable magnitude, both with RMS = 1.

AUTHOR CONTRIBUTIONS

HF collected and analyzed the location discrimination data. SS analyzed the heading discrimination data. DAZ recruited the ASD participants and conducted/supervised their comprehensive evaluation, and obtained Helsinki approval from the Shamir Medical Center. AZ conceived of and supervised the study. All authors contributed to writing the paper.

CONFLICT OF INTERESTS

The authors have no conflict of interest to report.

ACKNOWLEDGMENTS

We thank Mor Weinberger and Yarden Menashri-Sinai for help with location discrimination data collection, and Tamar Harpaz for administrative assistance. This work was supported by a grant from The Israeli Centers of Research Excellence (I-CORE, Center No. 51/11) to A.Z.

Table 1. ASD participant details

ASD participant number	Age	SCQ	IQ – WISC IV				ADOS			ADI-R		
			Block-Design	Matrix-Reasoning	Vocabulary	Similarities	Calibrated Severity Score	Social Affect	RRB	Reciprocal Social Interaction	Communication	RRB
1	8	17	7	9	10	9	9	7	7	13	11	5
2	9	17	11	9	9	8	9	10	10	27	14	6
3	10	21	13	16	13	13	8	8	8	10	13	1
4	10	14	12	13	10	16	9	9	9	11	8	5
5	10	11	9	7	7	7	9	9	9	12	12	2
6	11	21	11	10	7	10	8	7	7	13	15	10
7	11	15	7	11	8	7	8	8	8	11	11	5
8	11	19	12	13	10	16	7	7	7	17	13	5
9	11	22	12	9	11	10	7	10	10	23	19	7
10	12	13	15	13	12	10	9	8	8	11	8	3
11	12	29	10	12	7	8	7	7	7	20	19	4
12	13	22	8	7	11	8	10	10	10	14	9	9
13	13	12	7	10	8	10	10	9	9	23	11	4
14	14	19	15	9	15	17	10	9	9	20	14	8
15	14	14	14	9	12	15	7	5	5	11	7	5
16	15	18	7	9	10	9	6	6	6	8	4	5
17	16	30	13	9	11	12	8	6	6	13	13	8
18	17	22	9	11	8	10	10	10	10	11	5	6
Mean ± SD	12.1 ± 2.4	18.7 ± 5.3	10.7 ± 2.8	10.3 ± 2.3	9.9 ± 2.2	10.8 ± 3.2	8.4 ± 1.2	8.1 ± 1.6	8.1 ± 1.6	14.9 ± 5.4	11.4 ± 4.2	5.4 ± 2.3

SCQ scores are current from the time of the study. Other scores are from within 0–5 years of the study. RRB - Restricted and Repetitive Behaviors

Table 2. Control participant details

Control participant number	Age	SCQ	IQ – WISC IV			
			Block-Design	**Matrix-Reasoning**	**Vocabulary**	**Similarities**
1	8	5	14	10	18	12
2	9	1	14	10	15	10
3	9	1	13	13	17	14
4	9	1	13	14	14	15
5	10	1	11	10	10	12
6	11	1	9	15	15	8
7	11	3	10	8	9	10
8	12	10	6	11	10	9
9	12	1	12	7	8	9
10	12	1	9	8	11	9
11	13	3	14	12	13	18
12	13	0	5	8	8	9
13	13	8	14	17	11	16
14	14	3	10	15	8	13
15	15	0	10	14	12	13
16	15	5	12	16	9	7
17	15	7	13	16	12	15
18	16	2	14	14	8	11
19	16	3	8	8	9	9
20	16	2	11	10	5	11
Mean ± SD	12.5 ± 2.6	2.9 ± 2.8	11.1 ± 2.7	11.8 ± 3.2	11.1 ± 3.4	11.5 ± 3.0

IQ and SCQ scores are current from the time of the study.

REFERENCES

Alexi J, Cleary D, Dommisse K, Palermo R, Kloth N, Burr D, Bell J (2018) Past visual experiences weigh in on body size estimation. Sci Rep 8:1–8.

American Psychiatric Association (2000) Diagnostic and statistical manual of mental disorders (4th ed., text rev.). Washington, DC.

American Psychiatric Association (2013) The Diagnostic and Statistical Manual of Mental Disorders: DSM-5, 5th ed. Washington, DC: American Psychiatric Publishing.

Baron-Cohen S (2002) The extreme male brain theory of autism. Trends Cogn Sci 6:248–254.

Baron-Cohen S (2009) Autism: The empathizing-systemizing (E-S) theory. Ann N Y Acad Sci 1156:68–80.

Behrmann M, Avidan G, Leonard GL, Kimchi R, Luna B, Humphreys K, Minshew N (2006) Configural processing in autism and its relationship to face processing. Neuropsychologia 44:110–129 Available at: http://www.ncbi.nlm.nih.gov/pubmed/15907952.

Booth RDL, Happé FGE (2018) Evidence of Reduced Global Processing in Autism Spectrum Disorder. J Autism Dev Disord 48:1397–1408.

Brock J (2012) Alternative Bayesian accounts of autistic perception: Comment on Pellicano and Burr. Trends Cogn Sci 16:573–574.

Buckingham G, Michelakakis EE, Rajendran G (2016) The Influence of Prior Knowledge on Perception and Action: Relationships to Autistic Traits. J Autism Dev Disord 46:1716–1724.

Cornsweet TN (1962) The staircase-method in psychophysics. Am J Psychol 75:485–491.

Croydon A, Karaminis T, Neil L, Burr D, Pellicano E (2017) The light-from-above prior is intact in autistic children. J Exp Child Psychol 161:113–125.

Feigin H, Baror S, Bar M, Zaidel A (2020) Perceptual decisions are biased toward relevant prior choices. bioRxiv:858324.

Fischer J, Whitney D (2014) Serial dependence in visual perception. Nat Neurosci 17:738–743.

Friston KJ, Lawson R, Frith CD (2013) On hyperpriors and hypopriors: comment on Pellicano and Burr. Trends Cogn Sci 17:1 Available at: http://dx.doi.org/10.1016/j.tics.2012.11.003 [Accessed May 22, 2020].

Frith U, Happe F (1994) Autism: beyond "theory of mind." Cognition 50:115–132 Available at: http://www.ncbi.nlm.nih.gov/pubmed/8039356.

Fritsche M, Mostert P, de Lange FP (2017) Opposite Effects of Recent History on Perception and Decision. Curr Biol 27:590–595.

Happe F, Frith U (2006) The weak coherence account: detail-focused cognitive style in autism spectrum disorders. J Autism Dev Disord 36:5–25 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16450045.

Hosmer DW, Lemeshow S (1989) Applied Logistic Regression. In. New York: Wiley.

Kaneko Y, Sakai K (2015) Dissociation in decision bias mechanism between probabilistic information and previous decision. Front Hum Neurosci 9:1–11.

Karvelis P, Seitz AR, Lawrie SM, Seriès P (2018) Autistic traits, but not schizotypy, predict increased weighting of sensory information in bayesian visual integration. Elife 7.

Kersten D, Yuille A (2003) Bayesian models of object perception. Curr Opin Neurobiol 13:150–158 Available at: http://www.ncbi.nlm.nih.gov/pubmed/12744967.

Knill D, Richards W (1996) Perception as Bayesian inference. Available at: https://www.google.com/books?hl=en&lr=&id=_cTLCgAAQBAJ&oi=fnd&pg=PP1&dq=yuille+bulthoff+bayesian+decision+1996&ots=CfmIwIrFTU&sig=ikURWmJZ3nw4xnopMqZks858A9M [Accessed October 27, 2019].

Knill DC, Pouget A (2004) The Bayesian brain: the role of uncertainty in neural coding and computation. Trends Neurosci 27:712–719 Available at: http://www.ncbi.nlm.nih.gov/pubmed/15541511.

Kohn A (2007) Visual adaptation: Physiology, mechanisms, and functional benefits. J Neurophysiol 97:3155–3164.

Koldewyn K, Jiang Y V, Weigelt S, Kanwisher N (2013) Global/local processing in autism: not a disability, but a disinclination. J Autism Dev Disord 43:2329–2340 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23378063.

Lawson RP, Aylward J, White S, Rees G (2015) A striking reduction of simple loudness adaptation in autism. Sci Rep 5:1–7.

Lawson RP, Rees G, Friston KJ (2014) An aberrant precision account of autism. Front Hum Neurosci 8.

Leek MR (2001) Adaptive procedures in psychophysical research. Percept Psychophys 63:1279–1292.

Liberman A, Fischer J, Whitney D (2014) Serial dependence in the perception of faces. Curr Biol 24:2569–2574.

Liberman A, Manassi M (2018) Serial dependence promotes the stability of perceived emotional expression depending on face similarity. Attention, Perception, Psychophys 80:1461–1473.

Lieder I, Adam V, Frenkel O, Jaffe-Dax S, Sahani M, Ahissar M (2019) Perceptual bias reveals slow-updating in autism and fast-forgetting in dyslexia. Nat Neurosci 22:256–264 Available at: https://doi.org/10.1038/s41593-018-0308-9 [Accessed May 26, 2020].

Lord C, Rutter M, DiLavore PC, Risi S, Gotham K, Bishop S (2012) Autism diagnostic observation schedule, second edition (ADOS-2) manual (part I): Modules 1–4. Torrance, CA: Western Psychological Services.

Lord C, Rutter M, Le Couteur A (1994) Autism Diagnostic Interview-Revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. J Autism Dev Disord 24:659–685.

Ma WJ (2019) Bayesian Decision Models: A Primer. Neuron 104:164–175.

Mamassian P, Landy M, Maloney LT (2002) Bayesian Modelling of Visual Perception. In: Probabilistic Models of the Brain, pp 13–36 Available at: https://www.google.com/books?hl=en&lr=&id=mzBlvComcqwC&oi=fnd&pg=PA13&dq=Bayesian+modelling+of+visual+perception&ots=ro_6xaHxyf&sig=RPnB1Z-VkvPWD9fEnfqNAm-VVzY [Accessed May 22, 2020].

Menard S (2000) Coefficients of Determination for Multiple Logistic Regression Analysis. Am Stat 54:17–24.

Molesworth C, Chevallier C, Happé F, Hampton JA (2015) Children with autism do not show sequence effects with auditory stimuli. J Exp Psychol Gen 144:48–57.

Mottron L, Burack J (2001) Enhanced perceptual functioning in the development of autism. Available at: https://psycnet.apa.org/record/2001-01233-007 [Accessed May 22, 2020].

Mottron L, Dawson M, Soulieres I, Hubert B, Burack J (2006) Enhanced perceptual functioning in autism: an update, and eight principles of autistic perception. J Autism Dev Disord 36:27–43 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16453071.

Noel J-P, Lakshminarasimhan KJ, Park H, Angelaki DE (2020) Increased variability but intact integration during visual navigation in Autism Spectrum Disorder. Proc Natl Acad Sci 117:11158–11166 Available at: http://www.pnas.org/lookup/doi/10.1073/pnas.2000216117 [Accessed May 22, 2020].

Noel JP, De Niear MA, Stevenson R, Alais D, Wallace MT (2017) Atypical rapid audio-visual temporal recalibration in autism spectrum disorders. Autism Res 10:121–129.

Pascucci D, Mancuso G, Santandrea E, Libera C Della, Plomp G, Chelazzi L (2019) Laws of concatenated perception: Vision goes for novelty, decisions for perseverance. PLoS Biol 17:e3000144.

Pegors TK, Mattar MG, Bryan PB, Epstein RA (2015) Simultaneous perceptual and response biases on sequential face attractiveness judgments. J Exp Psychol Gen 144:664–673 Available at: https://pubmed.ncbi.nlm.nih.gov/25867223/ [Accessed July 26, 2020].

Peirce JW (2007) PsychoPy — Psychophysics software in Python. J Neurosci Methods 162:8–13.

Pellicano E, Burr D (2012) When the world becomes "too real": a Bayesian explanation of autistic perception. Trends Cogn Sci 16:504–510 Available at: http://www.ncbi.nlm.nih.gov/pubmed/22959875.

Pellicano E, Jeffery L, Burr D, Rhodes G (2007) Abnormal Adaptive Face-Coding Mechanisms in Children with Autism Spectrum Disorder. Curr Biol 17:1508–1512.

Plaisted K, Saksida L, Alcantara J, Weisblatt E (2003) Towards an understanding of the mechanisms of weak central coherence effects: experiments in visual configural learning and auditory perception. Philos Trans R Soc L B Biol Sci 358:375–386 Available at: http://www.ncbi.nlm.nih.gov/pubmed/12639334.

Plaisted K, Swettenham J, Rees L (1999) Children with Autism Show Local Precedence in a Divided Attention Task and Global Precedence in a Selective Attention Task. J Child Psychol Psychiatry 40:733–742.

Rogers SJ, Ozonoff S (2005) Annotation: what do we know about sensory dysfunction in autism? A critical review of the empirical evidence. J Child Psychol Psychiatry 46:1255–1268 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16313426 [Accessed May 22, 2020].

Rutter M, Bailey A, Lord C (2003a) The social communication questionnaire: Manual. Los Angeles, CA: Western Psychological Services.

Rutter M, Le Couteur A, Lord C (2003b) Autism diagnostic interview-revised. Los Angeles, CA: Western Psychological Services.

Schütt HH, Harmeling S, Macke JH, Wichmann FA (2016) Painfree and accurate Bayesian estimation of psychometric functions for (potentially) overdispersed data. Vision Res 122:105–123.

Shah A, Frith U (1983) An islet of ability in autistic children: a research note. J Child Psychol

Psychiatry 24:613–620 Available at: http://www.ncbi.nlm.nih.gov/pubmed/6630333.

Talluri BC, Urai AE, Tsetsos K, Usher M, Donner TH (2018) Confirmation Bias through
Selective Overweighting of Choice-Consistent Evidence. Curr Biol 28:3128–3135.

Taubert J, Burg E Van Der, Alais D (2016) Love at second sight : Sequential dependence of
facial attractiveness in an on-line dating paradigm. Sci Rep 6:2–6.

Turi M, Burr DC, Igliozzi R, Aagten-Murphy D, Muratori F, Pellicano E (2015) Children
with autism spectrum disorder show reduced adaptation to number. Proc Natl Acad Sci
U S A 112:7868–7872 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26056294.

Van de Cruys S, Vanmarcke S, Van de Put I, Wagemans J (2018) The Use of Prior
Knowledge for Perceptual Inference Is Preserved in ASD. Clin Psychol Sci 6:382–393
Available at: http://journals.sagepub.com/doi/10.1177/2167702617740955 [Accessed
May 26, 2020].

Wechsler D (2010) WISC-IV HEB. Wechsler Intelligence Scale for Children–Fourth Edition
(Hebrew Version). Jerusalem, Israel: Psychtech.

Wu HC, White S, Rees G, Burgess PW (2018) Executive function in high-functioning autism:
Decision-making consistency as a characteristic gambling behaviour. Cortex 107:21–36.

Zaidel A, Goin-Kochel RP, Angelaki DE (2015) Self-motion perception in autism is
compromised by visual noise but integrated optimally across multiple senses. Proc Natl
Acad Sci U S A 112:6461–6466 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/25941373.