Light environment influences mating behaviours during the early stages of divergence in tropical butterflies

Alexander E. Hausmann[1]*, Chi-Yun Kuo[1,2], Marília Freire[1,3], Nicol Rueda-M[4], Mauricio Linares[4], Carolina Pardo-Diaz[4], Camilo Salazar[4], Richard M. Merrill[1]*

[1] Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Grosshaderner Str. 2, 82152 Planegg-Martinsried, Germany

[2] Smithsonian Tropical Research Institute, Gamboa, 0843-03092, Panama

[3] Present address: Department of Evolutionary Neuroethology, Max Planck Institute for Chemical Ecology, Hans-Knöll-Straße 8, 07745 Jena, Germany

[4] Department of Biology, Faculty of Natural Sciences, Universidad del Rosario, Carrera 24 No 63C-69, Bogotá 111221, Colombia

*Corresponding authors: hausmann@bio.lmu.de and merrill@bio.lmu.de

ABSTRACT

Species divergence is facilitated when traits under divergent selection also act as mating cues. Fluctuations in sensory conditions can alter signal perception independently of adaptation to contrasting sensory environments, but how this fine scale variation affects behavioural isolation has received less attention, especially in terrestrial organisms. The warning patterns of *Heliconius* butterflies are under selection for aposematism and act as mating cues. Using computer vision, we extracted behavioural data from 1481 hours of video footage for 387 individuals. We show that the putative hybrid species *H. heurippa* and its close relative *H. timareta linaresi* differ in their response to divergent warning patterns, and that these differences are strengthened with increased local illuminance. Trials with live individuals reveal low-level assortative mating that are sufficiently explained by differences in visual attraction. Finally, results from hybrid butterflies are consistent with linkage between a major warning pattern gene and the corresponding behaviour, though the differences in behaviour we observe are unlikely to cause

rapid reproductive isolation as predicted under a model of hybrid trait speciation. Overall, our

results highlight that the role of ecological mating cues for behavioural isolation may depend on

the immediate sensory conditions during which they are displayed to conspecifics.

Keywords: *Heliconius* – Ecological Speciation – Sensory Environment – Magic traits – Hybrid

Trait Speciation – Computer vision

BACKGROUND

During ecological speciation, barriers to gene flow evolve as a result of ecologically-based

divergent selection [1]. These barriers are generally expected to build up gradually over time [2],

with premating isolation evolving early alongside ecological differences [1,3,4]. The evolution of

premating isolation is facilitated when traits under ecological selection act as mating cues

(sometimes known as 'magic traits' [5,6]), as this allows divergent natural selection acting on

ecological traits to be transferred to mating behaviours. However, the strength of behavioural

barriers (and consequently also the strength of genetic associations between ecological and

mating traits) may be influenced by the immediate, and perhaps rapidly changing, sensory

conditions, but this has received relatively little empirical attention (but see [7–11]). One reason

may be that robustly detecting these effects during the early stages of divergence, when they may

be most relevant, presents a substantial empirical challenge.

It is well established that the sensory environment can alter signal detection and

perception [12,13]. Colour perception of an object depends not only on its reflectance spectrum,

but also the illumination, available light spectrum and background colouration of the

environment, all of which may change within seconds and over very short distances [14]. This

can affect within-population preferences. For example, red colouration in male *Habronattus*

spiders is only an efficient mating signal if presented in broad-spectrum sunlight [15]. Beyond

vision, other sensory modalities can be also affected by the environment and lead to variation in

preferences within populations. Female swordtail fish are strongly attracted to the chemical

signals of conspecific males in clean water, but this preference is absent if signalling occurs in

polluted water [8]. Similarly, urban noise can disrupt the transmission of bird songs and interfere

with acoustic-based mate choice, unless singers show plasticity in song pitch [16].

The sensory environment may affect the evolution of reproductive isolation in two main

ways. First, adaptations to meet the ecological needs of different sensory environments can

influence female preferences, subsequently driving divergence in male signals, and leading to

reproductive isolation through sensory drive [17]. For example, in two closely related cichlid

fishes, *Pundamilia pundamilia* and *Pundamilia nyererei*, adaptation of the visual system to the

local environment [18,19] is associated with divergent female mate preference for male

colouration [18,20–24]. Second, prevailing environmental conditions may alter the efficacy of

signals used in mate choice [12], so that individual mating preferences may act as an important

reproductive barrier under some sensory conditions, but not others ('context-sensitive

preferences' [25]). If the strength of preference alleles depends on the sensory environment, then

this will influence their contribution to reproductive isolation. However, compared to the role of

sensory adaptation, the immediate influence of local sensory conditions on mating behaviours

and how this may relate to the evolution and maintenance of new species, has been less-studied.

By affecting context-sensitive preferences, the sensory conditions during signalling may

also influence the strength of genetic associations (*i.e.* linkage disequilibrium, LD) between

mating and ecological traits, which are typically required for speciation to proceed when gene

flow persists [26]. Specifically, when ecological traits act as mating cues, LD (between

ecological and preference loci) will arise as a natural consequence of mating preferences, but this

will be proportional to the strength of preference [27], which may be influenced by the sensory

environment. Regardless, unless preferences are very strong, more robust coupling between

mating and ecological components of reproductive isolation likely requires genetic architectures

that impede recombination between relevant loci (or one-allele mechanisms, see [28]). These

genetic architectures include physical linkage, which may be further strengthened by genomic

rearrangements like inversions, or – in the extreme – pleiotropy, where distinct traits are

controlled by the same allele [29]. To date, studies have reported physical linkage between

behavioural and ecological components of reproductive isolation for a handful of animal taxa,

including pea aphids [30], fish [31] and butterflies [32].

Heliconius butterflies are known for their bright warning patterns, which are often under

selection for Müllerian mimicry [33]. Closely related *Heliconius* species frequently – but not

always – have very different wing patterns, which additionally act as mating cues [34–40]. This

contributes to assortative mating because males almost invariably prefer wing patterns

resembling their own, and warning patterns in *Heliconius* are considered one of the best examples

of 'magic traits' in nature [6,34,41]. Variation in these warning pattern cues is largely controlled

by just four genes: *optix* [42], *cortex* [43], *WntA* [44] and *aristaless* [45]. The genetics of the

corresponding shifts in visual mate preference are less well known, though recent work has

implicated a handful of genes associated with neural signalling in close physical linkage to the

colour pattern gene *optix* [32,46].

There is substantial evidence that colour pattern alleles have introgressed between

otherwise separately evolving *Heliconius* lineages [47–53]. In particular, taxa within the

heurippa-timareta group, found in the eastern slopes of the South-American Andes, have

acquired red colour pattern elements from local races of the closely related species *H. melpomene*

via introgression of the alleles at the *optix* locus [48]. Adaptive introgression of colour pattern

alleles has frequently led to near perfect mimicry between local races of *H. timareta* and *H.*

melpomene; elsewhere, however, the resulting colour patterns are not shared with local

Heliconius species. In particular, *H. heurippa* has a red-yellow banded forewing pattern that is

not shared with other local *Heliconius* (Fig. 1A). Its close relative *H. timareta linaresi*, which is

assumed to represent the ancestral wing colour pattern of the *heurippa-timareta* group [54], only

displays a yellow band (Fig. 1A) and does not have an obvious co-mimic. These two populations

are geographically adjacent (Fig. 1A) and presumably share a contact zone (though it has not yet

been discovered). Despite their nominal status as separate species, *H. t. linaresi* and *H. heurippa*

likely represent an early stage of divergence; hybrids between *H. t. linaresi* and *H. heurippa* are

fully viable and fertile [55] and any post-mating isolation is likely limited to selection acting

against immigrant warning patterns.

The red patterns elements of *H. heurippa* are presumably maintained by strong frequency-

dependent selection imposed by predators [56]. However, their efficacy as intraspecific signals

may be influenced by how they are perceived by conspecifics under natural sensory conditions.

In this way, the evolution and maintenance of premating reproductive barriers may depend on

interactions between the signal, environment and receiver. *H. melpomene* are broadly separated

from *H. heurippa-timareta* (along with closely related *H. cydno*) taxa across a gradient of open to

closed forest and decreasing light intensity in a pattern of mosaic sympatry [57,58]. These

contrasting sensory environments are predicted to alter how colour patterns are perceived by

butterfly visual systems [59]. While *H. heurippa* and *H. t. linaresi* are not ecologically isolated in

this way, their forest habitats are highly heterogenous and *Heliconius* are known to settle with

their wings open in more brightly illuminated 'light patches' [60]. Whether changes in

illuminance affect male behaviours has not been investigated, but such an effect would suggest

that the efficacy of wing patterns as premating reproductive barriers may depend on fluctuations

in sensory conditions as females move through the environment.

To test whether warning patterns contribute to premating isolation between *H. heurippa*

and *H. t. linaresi*, we used a novel computer vision pipeline to extract behavioural data from

~1500h of video footage, alongside choice trails to determine levels of assortative mating. Our

resulting data, including ~17,000 behavioural interactions for 387 *H. heurippa*, *H. t. linaresi* and

hybrid males, allowed us to test (i) whether the two species show differences in visual mate

attraction towards con- and hetero-specific female patterns and (ii) whether differences in visual

attraction segregate with the colour patterns, consistent with physical linkage between the

behavioural and known colour pattern genes. Measuring illuminance in real-time at each female

pattern also allowed us to ask (iii) whether these behavioural differences are influenced by

fluctuating local conditions.

MATERIAL AND METHODS

(a) Butterfly collection, maintenance and crossing design. We maintained outbred butterfly

stocks, estabished from wild caught individuals, at the Universidad del Rosario insectaries in La

Vega, Colombia (4°59' N, 74°20' W, elevation 1257 m) between January 2016 and September

2019. *H. timareta linaresi* was collected from Guayabal (2°41'04"N, 74°53'17"W), and *H.*

heurippa was collected from three locations, Lejanías (03°34'0"N, 74°04'20.4"W), Buenavista

(4°10'30"N, 73°40'41"W) and Santa María (04°53'28.2"N, 73°15'11.4"W). We used these

stocks to generate F$_1$ hybrids (*H. t. linaresi* mother and *H. heurippa* father) and backcross hybrids

to each species (BL, backcross to *H. t. linaresi* and BH, backcross to *H. heurippa*). Details of

broods can be found in Table S1. All butterflies were supplied with ~10% sugar solution in

addition to *Lantana*, *Psiguria* and *Gurania spp.* as a source of pollen. Females were kept

individually, with *Passiflora* plants for oviposition, whereas experimental males were kept in

mixed groups. With the exception of a few individuals, eggs were collected every other day and

the caterpillars were raised until eclosion in plastic cups, and fed on fresh *Passiflora* leaves.

Figure 1: A) Locations at the Eastern slopes of the Colombian Andes where *H. heurippa* (red dots) and *H. timareta*

linaresi (blue dots) are known to occur. B) Video recording setup. The two static mounted females were presented

simultaneously to males and the area around each 'female' was surveyed with a camera. A light logger was

positioned at each station, recording illuminance every second. The position of each setup in the cage was changed

every hour. C) Computer vision pipeline for *post-hoc* motion detection. A difference image is formed between each

frame and its respective predecessor frame (identifying pixels with a change in value). Thresholding, blurring and

again thresholding pick out significant local changes in pixels ('signals'). Frames where 'signals' were detected (as

well as surrounding frames) then get extracted and combined into a new video file.

(b) Trials with mounted females. To test for differences in visual mating behaviours, we

assayed males in choice trials with dead mounted *H. heurippa* and *H. t. linaresi* females

presented simultaneously in a sun-exposed 4 x 4 x 2 m insectary. Virgin females were frozen on

the day of eclosion with their wings spread and kept at -20°C for at least one week. They were

then washed in hexane to remove residual cuticular hydrocarbons and other pheromones, and

mounted onto a small piece of wire. Over the course of the experiment, we used 21 *H. heurippa*

and 23 *H. t. linaresi* mounted females, and each day a randomly chosen pair of available females

was used for the experiment (*i.e.* one *H. heurippa* and one *H. t. linaresi* female). Mounted

females were individually attached to a horizontal wire (at ~70cm height above ground), itself

attached to a plastic pipe support buried in a bucket of cement at one of six locations within the

experimental cage. This location was changed every hour (hereafter 'trial'), and the new location

was chosen at random (female types were never in the same location twice during the same day).

Thus, our experiment included six different micro-environments with unique light conditions.

A *GoPro Hero 5 Black™* (GoPro, San Matteo, CA) camera, equipped with 128 GB

SanDisk™ Extreme SD card, was installed at each position, and about ~50cm above the

respective mounted female (Fig. 1B). Resolution was set to 1920 x 1080 pixels, at a frame rate of

60 fps and a shutter speed of 1/480. The field of view was set to *Narrow*. Colour was set to

GoPro and sharpness to *High*; ISO was set to 400; and any *auto*-setting, except for white balance,

was disabled. Each camera had a *Neewer™* UV filter attached, which reduced overexposure of

the yellow bands, and was constantly charged with an *Xlayer™ powerbank 210033*. At same

height as the mounted female and ~30cm away from it, we attached a *HOBO™ UA-002-64*

logger to measure illuminance [*lux*] every second during the recordings (Fig. 1B). Cameras and

light loggers were synced to the computer's system time using *GoPro Quik™* and *HOBOware™*

software, respectively, allowing us to precisely match the individual video frames and light

measurements.

Most of the virgin naive males matured in a separate communal cage before being

introduced into the experimental cage >4 days after eclosion, where they were tested in mixed

groups (median group size = 22, min = 7, max = 30). Males were numbered on the ventral side of

the wing and received a unique code of dots on the dorsal side of the wings, which allowed

subsequent identification from video footage. Each male was tested in the experimental cage over

the course of multiple days (median = 12 d, min = 1 d, max = 35 d). We recorded an average of

3.01 h (min = 0.20 h, max = 5.94 h) of material on each of the 246 recording days, with an

average starting time at 9.35AM (earliest = 7.48AM, latest = 2.23PM) and an average ending

time at 12.43PM (earliest = 9.34AM, latest = 4.23PM). Conducting this vast amount of

behavioural trials across different seasons and at different times allowed us to capture a variety of

light environments.

(c) Computer vision and video analyses. We used a custom motion-detection pipeline, which

post-hoc discarded video footage with no activity. The detection of frames with motion

('signals') was based on difference imaging and consecutive steps of blurring and thresholding,

as implemented in the *OpenCV* library available for C++ (Fig. 1C). Not all of the frames of male

motion sequences made it over the threshold, so we extracted the 'signal'-frames and the

surrounding frames one sec before and after a signal ('chunks'). Chunks were calculated in R

[61], and the extraction of the frames and final reduced videos were done with the *OpenCV*

library in C++. Code is accessible in our GitHub repository

(github.com/SpeciationBehaviour/visual_preference_heurippa_linaresi). Video material was

processed on a HP^{TM} ProDesk 600 G3 SFF Desktop computer (i7-7700 CPU @ 3.60GHz, 3600

MHz, 4 cores), faster than real time speed (runtime about 80% of video duration). All videos

were processed under the same threshold and blur settings. Remaining footage (with activity) was

curated manually at 66.6% speed using the *MPCHC^{TM} player*. We recorded three interaction

behaviours: 'approach' (the male is changing its direction of flight towards the mounted female,

resulting in a curve or circling motion), 'courtship initiation' (sustained hovering above mounted

female) and 'sitting' (male sits down on mounted female). All three types of interactions were

combined for subsequent analyses.

(d) Tetrad experiments with live females. We also performed 'tetrad' trials with virgin males

and females to test for assortative mating between *H. heurippa* and *H. t. linaresi*. For each trial,

sexually mature *H. heurippa* and *H. t. linaresi* males (one of each) were allowed to acclimatize

for 15 mins in a 2 x 4 x 2 m insectary, at which point *H. heurippa* and *H. t. linaresi* virgin

females (one of each) were introduced into the same cage. Once the first mating occurred, the

experiment was stopped and the combination of male and female recorded. Only trials which

ended in mating were included in subsequent analysis.

(e) Statistical analyses. We measured the illuminance at both mounted females for about 83% of

the behavioural data points (94% for *H. t. linaresi* and *H. heurippa* males, 73% for BL hybrids,

88% for all other hybrids tested). Illuminance refers to the intensity of light falling onto a surface

and is measured in *lux*. For subsequent statistical modelling, we used the logarithm to the base 10

to transform *lux* measures to a measure of perceived changes in light conditions at each mounted

female (log-illuminance). We additionally scaled (and centred) each set of log-illuminance

measurements, therefore making the choice of the base of the logarithm irrelevant for analyses.

All analyses were conducted in R, and raw data and an R Markdown for analysis can be

found in our GitHub repository

(https://github.com/SpeciationBehaviour/visual_preference_heurippa_linaresi). Posteriors will be

described with a 95% equal-tailed credible interval and the mean as a point estimate. We

analysed data from trials with mounted females using logistic regression with the *brms* package

[64], an interface to the Bayesian software Stan [65] and implemented in R [61]. Male

interactions directed towards the *H. t. linaresi* or *H. heurippa* mounted females were fitted as a

binary Bernoulli-distributed response variable (*i.e.* 0 and 1, respectively). Consequently,

estimates from the model can be understood as a proportion of interactions with the mounted *H.*

heurippa female, with higher values indicating stronger relative attraction to the *H. heurippa*

'female' and lower values indicating stronger relative attraction to the *H. t. linaresi* 'female'.

Models initially included all possible nested variations of the fully saturated model explaining

effects of 1) male type, 2) log-illuminance at the *H. heurippa* 'female', and 3) log-illuminance at

the *H. t. linaresi* 'female', as well as all possible interactions between 1-3 (resulting in 19

possible model structures). Individual ID and trial were fitted as random effects in all of the

models, correcting for overdispersion of the data. Including trial (and in many cases also

individual ID) as a random effect has often been disregarded in previous analyses (but see e.g.

[66]). However, we found strong correlation among individuals' behaviour during trials (see

supplementary R Markdown), likely due to the unique combination of males, mounted females

and their position in the experimental cage during a given trial. The latter may have also affected

sensitivity of our motion-detection software. We therefore considered it inevitable to correct for

this.

To test for 'species' differences, we initially fitted models for the 'pure' males, where

male type = '*H. heurippa*' or '*H. t. linaresi*'. We next tested for physical linkage between the loci

underlying variation in warning pattern and relative attraction. Because differences in wing

colour pattern between *H. heurippa* and *H. t. linaresi* are predominantly controlled by a single

locus whose alleles control the expression of the gene *optix*, which is responsible for the red

forewing bar seen in *H. heurippa,* segregation of the red bar in BL individuals allowed us to test

for physical linkage (see [32,37]). This was done by fitting models for the BL males, where male

type = 'red' (*Bb* genotype) or 'no red' (*bb* genotype). For the latter, we additionally fitted brood

ID as a random effect. To determine which fixed and interaction terms to retain [67], we

calculated the widely applicable information criterion (WAIC) and the leave-one-out information

criterion (LOOIC) for each model, using the *loo* package [68], and WAIC weights using the *brms*

package [64]. In all of our models, weakly informative priors (centred around the value for no

preference) were set for coefficients corresponding to the different male types, which gives small

prior probabilities for extreme values very close to 0 or 1. For all other coefficients, default (non-

informative) priors were applied. Posteriors of the estimated marginal means (EMMs) were

calculated using the *emmeans* package [69]. From this, we retrieved posteriors of contrasts and

calculated the posterior probability (PP) that the EMMs differ from each other. We additionally

fitted 'categorical illuminance' models to the 'pure' and the BL males, adopting the best fitting

model structure determined for each dataset previously. For the 'pure' and BL males separately,

we determined the median illuminance value at the *H. heurippa* 'female' (see Fig. S1) and the

median illuminance value at the *H. t. linaresi* 'female' when male responses occurred. Values

smaller than or equal to the median were considered 'poorly lit', and values greater than the

median were considered 'brightly lit'.

For data from the tetrad experiments, we fitted observed counts of each mating outcome

(type of male and female involved) as a Poisson-distributed response variable. We included the

specific male-female combination of the mating outcome as a fixed factor. Transforming the

resulting model estimates into proportions makes this effectively a multinomial model [70]. Non-

informative default priors were applied to all of the coefficients. PP were calculated with the

brms package [64]. We then compared the observed frequency of each mating combination to the

predicted frequencies based on our measurements of visual preference from the mounted female

experiment. Predictions were derived by multiplying the frequency that a male type was involved

in *any* combination of mating during the tetrad experiments by its respective EMMs from the

models fitted to the mounted female data. Predictions were based either on the overall EMM for

each type, or on the interaction term EMMs from the categorical model. Posterior distributions

for each prediction were calculated using the *binom* package under the default prior [71].

RESULTS

(a) Species comparisons: i) Divergent visual attraction behaviours in *H. timareta linaresi*

and *H. heurippa* males are dependent on the light environment.

Over the course of 1.5 years, we collected 1481 hours of video footage. Using our computer

vision pipeline, we reduced this to 66 hours that required human curation (*i.e.* 4.5% of the total

footage recorded), during which we recorded 16,995 individual behavioural 'interactions' from

387 individual males (on average 43.9 'interactions' per male). These data allowed us to

determine the effects of male type and illuminance on the relative visual attraction to the *H.*

heurippa mounted 'female' (hereafter 'preference'). The best fitting model for the pure *H.*

heurippa and *H. t. linaresi* males retained male type, log-illuminance at the *H. heurippa* 'female'

and their interaction (Table S2: model #1). Overall, our results suggest that the local light

environment can influence the strength of visual attraction.

Across the entire dataset, illuminance at the mounted *H. heurippa* female increased the

differences in preference between the male types, as evidenced by the interaction between male

type and log-illuminance at *H. heurippa* female (Fig. 2A, PP simple slope *H. heurippa* > simple

slope *H. t. linaresi* = 0.996). This was largely driven by an increase in log-illuminance at the *H.*

heurippa 'female' leading to a stronger conspecific preference in *H. heurippa* males (PP simple

slope > 0 = 0.993); there was limited support for an increase in log-illuminance at the *H.*

heurippa 'female' being associated with an increase in *H. t. linaresi* male preference for their

own phenotype (PP simple slope < 0 = 0.768). Overall, *H. heurippa* males showed a higher

proportion of interactions with the *H. heurippa* mounted females than the *H. t. linaresi* males.

Although supported by our Bayesian model with high credibility (posterior probability (PP) *H.*

heurippa relative visual attraction > *H. t. linaresi* relative visual attraction = 0.969), this

difference in preference was relatively subtle (0.07, CrI: 0.00 - 0.14) and characterised by

considerable within-population variation (Fig. 2B). Nevertheless, this effect nearly doubled when

the *H. heurippa* 'female' was in brighter light (0.13, CrI: 0.05 - 0.22; Fig 2D) and was absent

when the *H. heurippa* 'female' was poorly lit (0.01, CrI: -0.07 - 0.09; Fig 2C).

Figure 2: The effect of illuminance at the mounted *H. heurippa* female on relative attraction towards *H. heurippa* 'females' (*i.e.* the proportion of interactions with the mounted *H. heurippa* 'female' as opposed to the *H. t. linaresi* 'female'). **A**: Relative attraction towards *H. heurippa* 'females' of *H. t. linaresi* (blue) and *H. heurippa* males (red) under changing illuminance at the mounted *H. heurippa* female. Illuminance score on the x-axis was log-scaled and the coloured area around each regression line represents the 95% credible interval (CrI). Dashed vertical line represents the mean log-illuminance used as a cutoff to define the poorly and brightly lit conditions (below). Overall relative attraction towards *H. heurippa* 'females' for *H. t. linaresi* males (blue) and *H. heurippa* males (red); (**B**) across light environments; (**C**) for poorly lit *H. heurippa* 'female'; (**D**) brightly lit *H. heurippa* 'female'. Note that only preference data with the associated illuminance data were included in the underlying statistical models. B-D show the Gardner-Altman plots (see (Ho et al. 2019)), where the right-hand axis shows the difference between the two male types: horizontal lines project from the means of the posteriors for each male type (means and CrIs can be found in Table S3). Vertical thick line shows the 95% CrI for the posterior of the difference between the means, while the dot in its middle shows the mean of this posterior. In B-D, each point represents a single individual and its size is relative to the number of associated observations. The same individual males may appear in all three panels (unless they experienced only one of the light environments around the *H. heurippa* 'female' or illuminance data was not recorded during their responses, in which case they only appear in B; for individual trajectories, see Fig. S5AF). Custom swarmplot was used to distribute the dots horizontally.

(b) Species comparisons: ii) *H. heurippa* **males mate more often with conspecific females in**

choice trials.

To test for assortative mating we also conducted 'tetrad' mate choice trials between *H. heurippa*

and *H. t. linaresi*. During 89 tetrad trials with a total of 356 butterflies, we observed 50 con- and

39 heterospecific matings, supporting a degree of assortative mating (PP for positive assortative

mating = 0.88). This trend was driven by a higher proportion of conspecific matings involving an

H. heurippa male (0.405, CrI: 0.305 - 0.508) than heterospecific matings involving an *H.*

heurippa male (0.270, CrI: 0.183 - 0.366]) (PP = 0.94). *H. t. linaresi* males did not participate

more frequently in con- rather than heterospecific matings (PP = 0.43) (Fig. 3). In general, *H.*

heurippa males mated more often than *H. t. linaresi* males (60 vs. 29 times). Our results closely

match the predictions of the observed proportions from the mounted females experiment with

brightly lit *H. heurippa* 'female' (predictions indicated as horizontal purple bars in Fig. 3), and,

to some extent, those derived from all of the illuminance conditions combined (Table S4).

Figure 3: *H. heurippa* males show a preference for live, conspecific females in the tetrad experiments, but *H. t. linaresi* males do not. Dashed vertical line indicates expectation if neither species shows a preference to mate with its own and if both species are equally likely to engage into mating. Posterior distributions for each mating pair combination are displayed as a histogram. 95% CrI is coloured in red. Vertical orange line indicates the mean of the posterior distribution. Predictions based on the visual attraction data from the mounted females experiment with brightly lit *H. heurippa* 'female' are displayed with their 95% CrI as horizontal purple lines above the respective posterior distribution.

(c) Hybrid comparisons: Patterns of behaviour in backcross hybrids are consistent with linkage between colour and preference loci.

Estimates of preference suggest that first generation hybrid (F_1) males behave similarly as the *H. heurippa* males (Fig. S4D). This might suggest that *H. heurippa* alleles for attraction to red colour are dominant over the corresponding *H. t. linaresi* alleles; however, challenging this hypothesis, BH (backcross to *H. heurippa*) males seem to display a preference for the *H. t. linaresi* 'females' (Fig. S4E). For BL (backcross to *H. t. linaresi*) males, the fully saturated model (*i.e.* including all possible interactions) was the best fit (Table S5, model #1). Once again, the six best fitting models included the interaction between type (*i.e.* red and non-red wing pattern) and log-illuminance at the *H. heurippa* 'female' (91% of cumulative WAIC weight for an effect of this interaction term), suggesting that illuminance of the *H. heurippa* 'female' may increase existing differences in preference between the two male genotypes tested. BL males with a red band (*Bb*) were more likely to court/interact with the *H. heurippa* 'female' than BL males lacking the red band (*bb*) (Fig. S5). This difference was supported with moderately high probability by our Bayesian model (PP preference $Bb > bb = 0.897$). Although these represent small effects in absolute terms (0.03, CrI: -0.02 - 0.08]), this difference accounts for ~50% of the difference in means of the parental populations, consistent with linkage between colour and preference loci.

As for males from the parental populations, the difference in preference between males with the two backcross to BL genotypes (*i.e. Bb* vs. *bb*) was more pronounced when the *H.*

heurippa 'female' was in bright light (Fig. 4CD). There was only a small effect of the three-way

interaction with illuminance measures at the *H. t. linaresi* 'female' - the difference was slightly

higher when the *H. heurippa* mounted female was brightly lit (Fig. 4C, 0.11, CrI: 0.02 - 0.21), as

opposed to when both were brightly lit (Fig. 4D, 0.07, CrI: 0.00 - 0.13). A brightly lit *H. t.*

linaresi mounted female led to a slightly reversed pattern (*i.e. bb > Bb*), but only when the *H.*

heurippa mounted female was poorly lit (Fig. 4B, -0.03, CrI: -0.13 - 0.07). If both mounted

females where poorly lit, there was no difference in preference (Fig. 4A, 0.01, CrI: -0.05 - 0.08).

Across the entire dataset, illuminance at the mounted *H. heurippa* female increased the

differences in preference between the two BL genotypes (Fig. S2B, PP simple slope *Bb* > simple

slope *bb* = 0.999). While the slope for red backcross males is only slightly positive (PP = 0.859),

the slope for males without red is strongly negative (PP = 0.999). However, we observed a

stronger effect of illuminance at the *H. t. linaresi* 'female' than for the pure males. We found a

slightly negative slope for the interaction term with illuminance at the *H. t. linaresi* 'female' for

red BL males (Fig. S2D, PP = 0.722), but BL males without red showed a stronger preference for

the *H. heurippa* 'female' when the *H. t. linaresi* 'female' was bright (Fig. S2D, PP = 0.999).

Similar to the pure males, the three-way interaction indicates that a bright *H. heurippa* 'female'

may more strongly induce preference for own phenotype when the *H. t. linaresi* 'female' is less

bright (Fig. S2GH and signatures in raw data Fig. S3CDGH).

Figure 4: The effect of illuminance at the mounted *H. heurippa* female on relative attraction towards *H. heurippa* 'females' (i.e. the proportion of interactions with the mounted *H. heurippa* 'female' as opposed to the *H. t. linaresi* 'female') for BL hybrid males. Relative attraction towards *H. heurippa* 'females' for non-red (*bb*; blue dots) and red males (*Bb*; red dots): (**A**) with poorly lit *H. t. linaresi* and *H. heurippa* 'females'; (**B**) brightly lit *H. t. linaresi* and poorly lit *H. heurippa* 'females'; (**C**) poorly lit *H. t. linaresi* and brightly lit *H. heurippa* 'females'; and (**D**) brightly-lit *H. t. linaresi* and *H. heurippa* 'females'. Note that only preference data with the associated illuminance data were included in the underlying statistical model. Means and CrIs can be found in Table S3. Gardner-Altman plots are as in Figures 1B-D. For individual trajectories, see Fig. S4BC.

DISCUSSION

By collecting ~1500h of mate choice data, we have shown that the light environment can influence visual mating behaviours during the early stages of divergence in *Heliconius* butterflies. Although our data are characterised by considerable individual variation, we observed significant differences in the visual attraction behaviours of *H. heurippa* and *H. timareta linaresi* males, and this difference is stronger when the female patterns are more brightly lit. Experiments with live males and females revealed a degree of assortative mating, and the differences in visual attraction behaviours that we observe are sufficient to explain this. We have also shown that differences in visual attraction are associated with the presence or absence of the red forewing band in interspecific hybrids under bright light conditions, perhaps suggesting physical linkage between the ecologically relevant colour pattern gene and those for the corresponding behaviour.

Studies of speciation often focus on already diverged groups, which are maintained by multiple reproductive barriers [66,72], making it difficult to draw conclusions about the role of the individual barriers and their ecological context. At the other extreme, barriers acting at the early stages of divergence may be of small effect and, especially in the case of behavioural phenotypes, require very large datasets to draw robust conclusions, which may not always be feasible. Behavioural researchers increasingly use computer vision [73], and software for automated tracking and individual identification is widely available (e.g. [74,75]). However, applying these techniques to footage with heterogenous backgrounds, variable light environments and arenas larger than the camera's field of view remains a challenge. To overcome these limitations, we combined computer vision, allowing *post-hoc* motion detection, with subsequent human curation. This permitted a low-cost solution, without the need for complex equipment, and

importantly allowed us to capture frames before and after motion is detected from many hundreds

of hours of video footage.

This large volume of data revealed that increased illuminance at the red mounted *H.*

heurippa female increased the frequency of interactions between *H. heurippa* males with *H.*

heurippa females (relative to *H. t. linaresi* females). *H. heurippa* males were no more likely than

H. t. linaresi males to interact with the *H. heurippa* female when she was poorly lit (lower 50%

quantile of illuminance values; Fig. 2C), but were 1.3 times (*i.e.* 0.57/0.44, Table S3) more likely

to interact with the *H. heurippa* female in brighter light (upper 50% quantile of illuminance

values; Fig. 2D). Similar effects of the light environment have been observed in aquatic

organisms. For example, the attractiveness of red colour during courtship is dependent on the

spectrum of available light, influenced by water depth and turbidity, and recognition of the con-

specifics may fail entirely under certain light environments (e.g. [7,10,18,23,76]). However, far

fewer studies have tested in behavioural experiments how changing light environments influence

mating preferences that contribute to population divergence in terrestrial organisms (but see

[15,77,78]).

The exact mechanisms underlying the differences in behaviour we observe under

contrasting lighting conditions isn't immediately clear. Insects, including papilionid butterflies

[79], have frequently evolved colour constancy across light environments [80,81], and although

this may only partially succeed [79,80], we generally expect individuals to be able to distinguish

H. heurippa and *H. t. linaresi* patterns under different brightness conditions. Alternatively, male

attraction to female colour patterns might to some degree be 'wavelength-specific', where

triggering of behaviours depends greatly on an object's emission of specific wavelengths and on

the intensity of those wavelengths [81]. Under this scenario, a colour cue might only trigger a

behaviour when presented at intense illuminance including wavelengths from the relevant part of

the spectrum [81–83]. The differences in illuminance we measured may correlate with spectral

differences; under shaded conditions, the available light may be reflected from the vegetation or

from outside the path of direct sunlight, and consequently lack red wavelengths (but be rich in

greenish or bluish light, respectively) [14,15]. Indeed, previous work modelling *Heliconius* vision

suggests that red patterns are more conspicuous to *Heliconius* when presented in bright sunlight

[59] (though whether this affects *Heliconius* behaviours has not previously been tested).

Regardless of the proximate mechanism, our data indicate that the prevailing light

conditions can influence visual mating behaviours in *Heliconius* butterflies. Unlike many other

examples, which largely involve anthropogenic induced changes in aquatic environments

[7,9,76], the effects we see here potentially act across very short time and spatial scales. This

reflects the forest environment where the light environment can vary rapidly across both time and

space. As such, male response may depend less on the broader environment (perhaps influenced

by forest type), than on the movement of females across smaller spatial scales (for example

between dimly and brightly lit patches). *Heliconius* butterflies are known to bask in the sun with

their wings open, particularly during the morning hours [60]. This may primarily be for

thermoregulation, though other species are known to display in local environments where they

are most conspicuous, and *Heliconius* might follow a similar strategy. *Anolis* lizards, for

example, are known to occupy micro-habitats in which their dewlap colour is most conspicuous

[84,85]. Visual modelling suggests *Heliconius* red patterns are more conspicuous to avian

predators when presented in bright sunlight, suggesting it may also enhance aposematism [59].

Whatever the ultimate reason for aggregating in sun-exposed patches, our data suggest that these

behaviours could enhance the strength of divergent mating preferences as a by-product.

Whether the differences in visual attraction behaviours that we observe translate to

assortative mating, and therefore contribute to reproductive isolation, is an important question.

Previous studies testing for assortative mating between *Heliconius* species closely related to *H.*

heurippa and *H. t. linaresi* in tetrad experiments similar to ours report very low frequencies of

interspecific matings [34,86,87]. This was not the case in our experiments, where trials resulting

in interspecific mating between *H. heurippa* and *H. t. linaresi* were not uncommon. Nevertheless,

the large number of trials in our study allowed us to detect assortative mating, albeit at low

levels, though this was much stronger for the 60 trials in which *H. heurippa* males mated (Fig. 3).

Unfortunately, we do not have illuminance data for these experiments, and this would be difficult

to measure given the movement of the females. Considering that the live females may have been

basking in the sun during the trials, that the overall activity of *Heliconius* is highest when it is

sunny [88] and that mating in *Heliconius* seems to occur much more frequently on sunny than on

cloudy days (M. Linares, pers. obs.), it is likely that a large proportion of the trials with mating

outcomes in our experiments included sun-exposed females. When accounting for the more

frequent involvement of *H. heurippa* males in mating, the mating rates between *H. heurippa* and

H. t. linaresi in our tetrad experiments closely match estimates derived from our experiments

with dead mounted females in bright light. Although we cannot rule out a role for cues acting

across other sensory modalities, the differences in visual attraction alone are sufficient to explain

the levels of assortative mating we observe.

Considering our data for *H. heurippa* and *H. t. linaresi* hybrids, overall, it is difficult to

ascertain patterns of dominance for visual attraction from our data. This is perhaps not surprising

given that the differences in behaviour between populations are subtle and are shaped by

considerable variation. However, segregation of the *optix* alleles, which control the red pattern

elements in *Heliconius*, in the backcross to *H. t. linaresi* did allow us to test for linkage between

the warning pattern cue and the corresponding behaviour. Once again, we observed illuminance-

induced shifts in visual attraction, so that the two types of backcross to *H. t. linaresi* males (*i.e.*

red, *Bb* vs. non-red, *bb*) differed in behaviour, but only under higher illuminance conditions.

These results are consistent with physical linkage between behavioural loci and *optix*, as has been

shown elsewhere – but with much greater effects – between the closely related species *H. cydno*

and *H. melpomene* [32]. Physical linkage will help to maintain key genetic associations (*i.e.* LD)

between loci underlying ecologically relevant traits and those for premating isolation, like mating

preferences, facilitating speciation with gene flow in general [26,28], and hybrid trait speciation

in particular. Although the differences between backcross to *H. t. linaresi* genotypes account for

~50% of that observed in the two parental taxa, in absolute terms, the effects are unlikely to

permit sufficient power for a formal QTL study and caution should be exercised when

interpreting our results. Nevertheless, our results are consistent with a simple genetic mechanism

by which behavioural alleles were acquired alongside red colour pattern alleles through

introgression from *H. melpomene* into the '*heurippa*/*timareta*/*cydno*' lineage [89,90]. However, it

seems unlikely that the behavioural differences we observe here would rapidly lead to strong

reproductive isolation as predicted under a model of hybrid trait speciation [86,89].

In conclusion, the behavioural differences we report here for *H. heurippa* and *H. t.*

linaresi are similar in strength to those reported elsewhere for other *Heliconius* taxa at the early

stages of divergence [e.g. 38,66]. Alone these may represent only weak barriers to gene flow;

however, by augmenting divergent ecological selection acting on the warning pattern cue, they

may facilitate the accumulation of additional barriers as speciation proceeds. In addition, our

results suggest that the degree to which *Heliconius* warning pattern contribute to premating

isolation may depend on local illuminance, which can change rapidly in both time and space.

Without more stable differences in the sensory environment, perhaps facilitated by shifts in

habitat use, these fluctuations might constrain speciation. However, *Heliconius* are known to

aggregate and display their warning patterns in sun-exposed patches within the broader forest

environment, and our results suggest that this context would enhance premating isolation. Traits

predominantly shaped by ecology frequently act as mating cues, which by coupling divergent

natural selection to premating isolation can promote speciation with gene flow [6]; our results

highlight that this effect may depend on the immediate sensory conditions during which these

ecological mating cues are displayed to conspecifics.

ACKNOWLEDGEMENTS

We are very grateful to the Abondano-Almeida family for helping AEH and MF settle in

Colombia; Juan Sebastián Sánchez, Óscar Penagos, Isabel Leon and Lina Gabriela Melo for

assistance in the insectaries; Annika Neuhaus, Morgan Oberweiser, Saoirse McMahon and Lucas

Asis for help scoring videos; and Martin Küblbeck, Lucie Queste, Matteo Rossi, Daniel Shane

Wright and Stephen Montgomery for comments on the manuscript. Field collections and

insectary rearing were conducted under permit no. 530 issued by the Autoridad Nacional de

Licencias Ambientales de Colombia (ANLA). CS and CP were funded by Universidad del

Rosario (Grant Number: IV-FGD005). ML was partially funded by the Faculty of Natural

Sciences at Universidad del Rosario. AEH, MF and RMM were supported by an Emmy Noether

fellowship and research grant awarded to RMM by the Deutsche Forschungsgemeischaft (DFG)

(Grant Number: GZ: ME 4845/1-1).

REFERENCES

1. Nosil P. 2012 *Ecological speciation*. Oxford University Press. (doi:10.1093/acprof:osobl/9780199587100.001.0001)
2. Mallet J. 2008 Hybridization, ecological races and the nature of species: empirical evidence for the ease of speciation. *Philos. Trans. R. Soc. B.* **363**, 2971–2986. (doi:10.1098/rstb.2008.0081)
3. Coyne JA, Orr HA. 1997 "Patterns of speciation in Drosophila" revisited. *Evolution* **51**, 295–303. (doi:10.1111/j.1558-5646.1997.tb02412.x)
4. Grant PR, Grant BR. 1997 Genetics and the origin of bird species. *PNAS* **94**, 7768–7775. (doi:10.1073/pnas.94.15.7768)
5. Gavrilets S. 2004 *Fitness landscapes and the origin of species (MPB-41)*. Princeton University Press. (doi:10.2307/j.ctv39x541)
6. Servedio MR, Van Doorn GS, Kopp M, Frame AM, Nosil P. 2011 Magic traits in speciation:'magic'but not rare? *Trends Ecol. Evol.* **26**, 389–397. (doi:10.1016/j.tree.2011.04.005)
7. Seehausen O, Van Alphen JJM, Witte F. 1997 Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* **277**, 1808–1811. (doi:10.1126/science.277.5333.1808)
8. Fisher HS, Wong BBM, Rosenthal GG. 2006 Alteration of the chemical environment disrupts communication in a freshwater fish. *Proc. R. Soc. B* **273**, 1187–1193. (doi:10.1098/rspb.2005.3406)
9. Heuschele J, Mannerla M, Gienapp P, Candolin U. 2009 Environment-dependent use of mate choice cues in sticklebacks. *Behav. Ecol.* **20**, 1223–1227. (doi:10.1093/beheco/arp123)
10. Maan ME, Seehausen O, Van Alphen JJM. 2010 Female mating preferences and male coloration covary with water transparency in a Lake Victoria cichlid fish. *Biol. J. Linn. Soc.* **99**, 398–406. (doi:10.1111/j.1095-8312.2009.01368.x)
11. Johnson AM, Stanis S, Fuller RC. 2013 Diurnal lighting patterns and habitat alter opsin expression and colour preferences in a killifish. *Proc. R. Soc. B* **280**, 20130796. (doi:10.1098/rspb.2013.0796)
12. Endler JA. 2000 Evolutionary implications of the interaction between animal signals and the environment. In *Animal signals: signalling and signal design in animal communication* (ed RG Espmark Y, Amundsen T), pp. 11–46. Tapir Academic Press.
13. Price TD. 2017 Sensory drive, color, and color vision. *Am. Nat.* **190**, 157–170. (doi:10.1086/692535)
14. Endler JA. 1993 The color of light in forests and its implications. *Ecol. Monogr.* **63**, 1–27. (doi:10.2307/2937121)
15. Taylor LA, McGraw KJ. 2013 Male ornamental coloration improves courtship success in a jumping spider, but only in the sun. *Behav. Ecol.* **24**, 955–967. (doi:10.1093/beheco/art011)
16. Slabbekoorn H, Peet M. 2003 Birds sing at a higher pitch in urban noise. *Nature* **424**, 267. (doi:10.1038/424267a)
17. Boughman JW. 2002 How sensory drive can promote speciation. *Trends Ecol. Evol.* **17**, 571–577. (doi:10.1016/s0169-5347(02)02595-8)
18. Seehausen O *et al.* 2008 Speciation through sensory drive in cichlid fish. *Nature* **455**, 620. (doi:10.1038/nature07285)

19. Maan ME, Seehausen O, Groothuis TGG. 2017 Differential survival between visual
environments supports a role of divergent sensory drive in cichlid fish speciation. *Am. Nat.*
189, 78–85. (doi:10.1086/689605)
20. Maan ME, Hofker KD, van Alphen JJM, Seehausen O. 2006 Sensory drive in cichlid
speciation. *Am. Nat.* **167**, 947–954. (doi:10.1086/503532)
21. P. Haesler M, Seehausen O. 2005 Inheritance of female mating preference in a sympatric
sibling species pair of Lake Victoria cichlids: implications for speciation. *Proc. R. Soc. B*
272, 237–245. (doi:10.1098/rspb.2004.2946)
22. Selz OM, Pierotti MER, Maan ME, Schmid C, Seehausen O. 2014 Female preference for
male color is necessary and sufficient for assortative mating in 2 cichlid sister species.
Behav. Ecol. **25**, 612–626. (doi:10.1093/beheco/aru024)
23. Seehausen O, van Alphen JJM. 1998 The effect of male coloration on female mate choice
in closely related Lake Victoria cichlids (Haplochromis nyererei complex). *Behav. Ecol.*
Sociobiol. **42**, 1–8. (doi:10.1007/s002650050405)
24. Stelkens RB, Pierotti MER, Joyce DA, Smith AM, van der Sluijs I, Seehausen O. 2008
Disruptive sexual selection on male nuptial coloration in an experimental hybrid
population of cichlid fish. *Philos. Trans. R. Soc. B*. **363**, 2861–2870.
(doi:10.1098/rstb.2008.0049)
25. Rosenthal GG. 2017 *Mate choice: the evolution of sexual decision making from microbes*
to humans. Princeton University Press. (doi:10.2307/j.ctt1vwmhb0)
26. Kopp M *et al.* 2018 Mechanisms of assortative mating in speciation with gene flow:
connecting theory and empirical research. *Am. Nat.* **191**, 1–20. (doi:10.1086/694889)
27. Kirkpatrick M, Barton NH. 1997 The strength of indirect selection on female mating
preferences. *PNAS* **94**, 1282–1286. (doi:10.1073/pnas.94.4.1282)
28. Felsenstein J. 1981 Skepticism towards Santa Rosalia, or why are there so few kinds of
animals? *Evolution* **35**, 124–138. (doi:10.2307/2407946)
29. Smadja CM, Butlin RK. 2011 A framework for comparing processes of speciation in the
presence of gene flow. *Mol. Ecol.* **20**, 5123–5140. (doi:10.1111/j.1365-
294x.2011.05350.x)
30. Hawthorne DJ, Via S. 2001 Genetic linkage of ecological specialization and reproductive
isolation in pea aphids. *Nature* **412**, 904–907. (doi:10.1038/35091062)
31. Bay RA *et al.* 2017 Genetic coupling of female mate choice with polygenic ecological
divergence facilitates stickleback speciation. *Curr. Biol.* **27**, 3344–3349.
(doi:10.1016/j.cub.2017.09.037)
32. Merrill RM, Rastas P, Martin SH, Melo MC, Barker S, Davey J, McMillan WO, Jiggins
CD. 2019 Genetic dissection of assortative mating behavior. *PLoS Biol.* **17**.
(doi:10.1101/282301)
33. Merrill RM *et al.* 2015 The diversification of Heliconius butterflies: what have we learned
in 150 years? *J. Evol. Biol.* **28**, 1417–1438. (doi:10.1111/jeb.12672)
34. Jiggins CD, Naisbit RE, Coe RL, Mallet J. 2001 Reproductive isolation caused by colour
pattern mimicry. *Nature* **411**, 302–305. (doi:10.1038/35077075)
35. Jiggins CD, Estrada C, Rodrigues A. 2004 Mimicry and the evolution of premating
isolation in Heliconius melpomene Linnaeus. *J. Evol. Biol.* **17**, 680–691.
(doi:10.1111/j.1420-9101.2004.00675.x)
36. Kronforst MR, Young LG, Kapan DD, McNeely C, O'Neill RJ, Gilbert LE. 2006 Linkage
of butterfly mate preference and wing color preference cue at the genomic location of
wingless. *PNAS* **103**, 6575–6580. (doi:10.1073/pnas.0509685103)

37.   Merrill RM, Van Schooten B, Scott JA, Jiggins CD. 2011 Pervasive genetic associations
between traits causing reproductive isolation in Heliconius butterflies. *Proc. R. Soc. B* **278**,
511–518. (doi:10.1098/rspb.2010.1493)
38.   Merrill RM, Gompert Z, Dembeck LM, Kronforst MR, McMillan WO, Jiggins CD. 2011
Mate preference across the speciation continuum in a clade of mimetic butterflies.
Evolution **65**, 1489–1500. (doi:10.1111/j.1558-5646.2010.01216.x)
39.   Sánchez AP, Pardo-Diaz C, Enciso-Romero J, Muñoz A, Jiggins CD, Salazar C, Linares
M. 2015 An introgressed wing pattern acts as a mating cue. *Evolution* **69**, 1619–1629.
(doi:10.1111/evo.12679)
40.   Rosser N *et al.* 2019 Geographic contrasts between pre-and postzygotic barriers are
consistent with reinforcement in Heliconius butterflies. *Evolution* **73**, 1821–1838.
(doi:10.1111/evo.13804)
41.   Merrill RM, Wallbank RWR, Bull V, Salazar PCA, Mallet J, Stevens M, Jiggins CD. 2012
Disruptive ecological selection on a mating cue. *Proc. R. Soc. B* **279**, 4907–4913.
(doi:10.1098/rspb.2012.1968)
42.   Reed RD *et al.* 2011 Optix drives the repeated convergent evolution of butterfly wing
pattern mimicry. *Science* **333**, 1137–1141. (doi:10.1126/science.1208227)
43.   Nadeau NJ *et al.* 2016 The gene cortex controls mimicry and crypsis in butterflies and
moths. *Nature* **534**, 106–110. (doi:10.1038/nature17961)
44.   Martin A *et al.* 2012 Diversification of complex butterfly wing patterns by repeated
regulatory evolution of a Wnt ligand. *PNAS* **109**, 12632–12637.
(doi:10.1073/pnas.1204800109)
45.   Westerman EL *et al.* 2018 Aristaless controls butterfly wing color variation used in
mimicry and mate choice. *Curr. Biol.* **28**, 3469–3474. (doi:10.1016/j.cub.2018.08.051)
46.   Rossi M, Hausmann AE, Thurman TJ, Montgomery SH, Papa R, Jiggins CD, Mcmillan
WO, Merrill RM. 2020 Visual mate preference evolution during butterfly speciation is
linked to neural processing genes. *bioRxiv* (doi:10.1101/2020.03.22.002121)
47.   Heliconius Genome Consortium. 2012 Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. *Nature* **487**, 94. (doi:10.1038/nature11041)
48.   Pardo-Diaz C, Salazar C, Baxter SW, Merot C, Figueiredo-Ready W, Joron M, McMillan
WO, Jiggins CD. 2012 Adaptive introgression across species boundaries in Heliconius
butterflies. *PLoS Genet.* **8**. (doi:10.1371/journal.pgen.1002752)
49.   Kronforst MR, Hansen MEB, Crawford NG, Gallant JR, Zhang W, Kulathinal RJ, Kapan
DD, Mullen SP. 2013 Hybridization reveals the evolving genomic architecture of
speciation. *Cell Rep.* **5**, 666–677. (doi:10.1016/j.celrep.2013.09.042)
50.   Wallbank RWR *et al.* 2016 Evolutionary novelty in a butterfly wing pattern through
enhancer shuffling. *PLoS Biol.* **14**. (doi:10.1371/journal.pbio.1002353)
51.   Zhang W, Dasmahapatra KK, Mallet J, Moreira GRP, Kronforst MR. 2016 Genome-wide
introgression among distantly related Heliconius butterfly species. *Genome Biol.* **17**, 25.
(doi:10.1186/s13059-016-0889-0)
52.   Enciso-Romero J, Pardo-Díaz C, Martin SH, Arias CF, Linares M, McMillan WO, Jiggins
CD, Salazar C. 2017 Evolution of novel mimicry rings facilitated by adaptive introgression
in tropical butterflies. *Mol. Ecol.* **26**, 5160–5172. (doi:10.1111/mec.14277)
53.   Jay P, Whibley A, Frézal L, de Cara MÁR, Nowell RW, Mallet J, Dasmahapatra KK,
Joron M. 2018 Supergene evolution triggered by the introgression of a chromosomal
inversion. *Curr. Biol.* **28**, 1839–1845. (doi:10.1016/j.cub.2018.04.072)
54.   Arias CF, Giraldo N, McMillan WO, Lamas G, Jiggins CD, Salazar C. 2017 A new

subspecies in a Heliconius butterfly adaptive radiation (Lepidoptera: Nymphalidae). *Zool.*
J. Linn. Soc. **180**, 805–818. (doi:10.1093/zoolinnean/zlw010)

55.   Rodríguez-Monje SV, Linares M. 2010 Ruta alternativa de reconstrucción del patrón de
coloración de Heliconius heurippa. Uniandes. See
https://univdelosandes.on.worldcat.org/oclc/916327490.

56.   Mallet J, Barton NH. 1989 Strong natural selection in a warning-color hybrid zone.
Evolution **43**, 421–431. (doi:10.2307/2409217)

57.   Merot C, Salazar C, Merrill RM, Jiggins CD, Joron M. 2017 What shapes the continuum
of reproductive isolation? Lessons from Heliconius butterflies. *Proc. R. Soc. B* **284**,
20170335. (doi:10.1098/rspb.2017.0335)

58.   Estrada C, Jiggins CD. 2002 Patterns of pollen feeding and habitat preference among
Heliconius species. *Ecol. Entomol.* **27**, 448–456. (doi:10.1046/j.1365-2311.2002.00434.x)

59.   Dell'Aglio DD, Troscianko J, Stevens M, McMillan WO, Jiggins CD. 2019 The
conspicuousness of the toxic Heliconius butterflies across time and habitat. *BioRxiv* ,
662155. (doi:10.1101/662155)

60.   Jiggins CD. 2017 *The ecology and evolution of Heliconius butterflies*. Oxford University
Press. (doi:10.1093/acprof:oso/9780199566570.001.0001)

61.   R Core Team. 2019 R: A Language and Environment for Statistical Computing.

62.   Laughlin SB. 1994 Matching coding, circuits, cells, and molecules to signals: general
principles of retinal design in the fly's eye. *Prog. Retin. Eye Res.* **13**, 165–196.
(doi:10.1016/1350-9462(94)90009-4)

63.   Akre KL, Johnsen S. 2014 Psychophysics and the evolution of behavior. *Trends Ecol.*
Evol. **29**, 291–300. (doi:10.1016/j.tree.2014.03.007)

64.   Bürkner P-C. 2018 Advanced Bayesian Multilevel Modeling with the R Package brms.
The R Journal, 10 (1), 395--411. doi: 10.32614. (doi:10.32614/rj-2018-017)

65.   Carpenter B *et al.* 2017 Stan: A probabilistic programming language. *J. Stat. Softw.* **76**.
(doi:10.18637/jss.v076.i01)

66.   Chamberlain NL, Hill RI, Kapan DD, Gilbert LE, Kronforst MR. 2009 Polymorphic
butterfly reveals the missing link in ecological speciation. *Science* **326**, 847–850.
(doi:10.1126/science.1179141)

67.   Gelman A, Hwang J, Vehtari A. 2014 Understanding predictive information criteria for
Bayesian models. *Stat. Comput.* **24**, 997–1016. (doi:10.1007/s11222-013-9416-2)

68.   Vehtari A, Gabry J, Magnusson M, Yao Y, Gelman A. 2019 loo: Efficient leave-one-out
cross-validation and WAIC for Bayesian models. *R Packag. version 2.2.0*

69.   Lenth R. 2019 emmeans: Estimated Marginal Means, aka Least-Squares Means. *R Packag.*
version 1.4.6

70.   Royle JA, Dorazio RM. 2008 *Hierarchical modeling and inference in ecology: the*
analysis of data from populations, metapopulations and communities. Elsevier.
(doi:10.1016/B978-0-12-374097-7.X0001-4)

71.   Dorai-Raj S. 2014 binom: Binomial confidence intervals for several parameterizations. *R*
Packag. version 1.1-1

72.   Stroud JT, Losos JB. 2020 Bridging the process-pattern divide to understand the origins
and early stages of adaptive radiation: a review of approaches with insights from studies of
Anolis lizards. *J. Hered.* **111**, 33–42. (doi:10.1093/jhered/esz055)

73.   Egnor SER, Branson K. 2016 Computational analysis of behavior. *Annu. Rev. Neurosci.*
39, 217–236. (doi:10.1146/annurev-neuro-070815-013845)

74.   Graving JM, Chae D, Naik H, Li L, Koger B, Costelloe BR, Couzin ID. 2019

DeepPoseKit, a software toolkit for fast and robust animal pose estimation using deep
learning. *Elife* **8**, e47994. (doi:10.7554/elife.47994)

75.   Romero-Ferrero F, Bergomi MG, Hinz RC, Heras FJH, de Polavieja GG. 2019 Idtracker.
ai: tracking all individuals in small or large collectives of unmarked animals. *Nat. Methods*
16, 179–182. (doi:10.1038/s41592-018-0295-5)

76.   Engström-Öst J, Candolin U. 2007 Human-induced water turbidity alters selection on
sexual displays in sticklebacks. *Behav. Ecol.* **18**, 393–398. (doi:10.1093/beheco/arl097)

77.   Summers K, Symula R, Clough M, Cronin T. 1999 Visual mate choice in poison frogs.
Proc. R. Soc. B **266**, 2141–2145. (doi:10.1098/rspb.1999.0900)

78.   Théry M, Pincebourde S, Feer F. 2008 Dusk light environment optimizes visual perception
of conspecifics in a crepuscular horned beetle. *Behav. Ecol.* **19**, 627–634.
(doi:10.1093/beheco/arn024)

79.   Kinoshita M, Arikawa K. 2000 Colour constancy in the swallowtail butterfly Papilio
xuthus. *J. Exp. Biol.* **203**, 3521–3530.

80.   Chittka L, Faruq S, Skorupski P, Werner A. 2014 Colour constancy in insects. *J. Comp.*
Physiol. A **200**, 435–448. (doi:10.1007/s00359-014-0897-z)

81.   Song B-M, Lee C-H. 2018 Toward a mechanistic understanding of color vision in insects.
Front. Neural Circuits **12**, 16. (doi:10.3389/fncir.2018.00016)

82.   Maan ME, Cummings ME. 2009 Sexual dimorphism and directional sexual selection on
aposematic signals in a poison frog. *PNAS* **106**, 19072–19077.
(doi:10.1073/pnas.0903327106)

83.   Paris TM, Allan SA, Udell BJ, Stansly PA. 2017 Wavelength and polarization affect
phototaxis of the Asian citrus psyllid. *Insects* **8**, 88. (doi:10.3390/insects8030088)

84.   Fleishman LJ. 1992 The influence of the sensory system and the environment on motion
patterns in the visual displays of anoline lizards and other vertebrates. *Am. Nat.* **139**, S36--
S61. (doi:10.1086/285304)

85.   Leal M, Fleishman LJ. 2002 Evidence for habitat partitioning based on adaptation to
environmental light in a pair of sympatric lizard species. *Proc. R. Soc. B* **269**, 351–359.
(doi:10.1098/rspb.2001.1904)

86.   Mavárez J, Salazar CA, Bermingham E, Salcedo C, Jiggins CD, Linares M. 2006
Speciation by hybridization in Heliconius butterflies. *Nature* **441**, 868–871.
(doi:10.1038/nature04738)

87.   Kronforst MR, Young LG, Gilbert LE. 2007 Reinforcement of mate preference among
hybridizing Heliconius butterflies. *J. Evol. Biol.* **20**, 278–285. (doi:10.1111/j.1420-
9101.2006.01198.x)

88.   Crane J. 1955 Imaginal behavior of a Trinidad butterfly, Heliconius erato hydara
Hewitson, with special reference to the social use of color. *Zool.* **40**, 167–196.

89.   Jiggins CD, Salazar C, Linares M, Mavarez J. 2008 Hybrid trait speciation and Heliconius
butterflies. *Philos. Trans. R. Soc. B.* **363**, 3047–3054. (doi:10.1098/rstb.2008.0065)

90.   Melo MC, Salazar C, Jiggins CD, Linares M. 2009 Assortative mating preferences among
hybrids offers a route to hybrid speciation. *Evolution* **63**, 1660–1665. (doi:10.1111/j.1558-
5646.2009.00633.x)

Supplementary material: Light environment influences mating behaviours during the early

stages of divergence in tropical butterflies

Alexander E. Hausmann[1*], Chi-Yun Kuo[1,2], Marília Freire[1,3], Nicol Rueda-M[4], Mauricio

Linares[4], Carolina Pardo-Diaz[4], Camilo Salazar[4], Richard M. Merrill[1*]

[1] Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Grosshaderner Str. 2, 82152 Planegg-
Martinsried, Germany
[2] Smithsonian Tropical Research Institute, Gamboa, 0843-03092, Panama
[3] Present address: Department of Evolutionary Neuroethology, Max Planck Institute for Chemical Ecology, Hans-Knöll-Straße 8,
07745 Jena, Germany
[4] Department of Biology, Faculty of Natural Sciences, Universidad del Rosario, Carrera 24 No 63C-69, Bogotá 111221, Colombia

*Corresponding authors: hausmann@bio.lmu.de and merrill@bio.lmu.de

1) Supplementary Figures

Figure S1: 'Poorly lit', median and 'brightly lit' *H. heurippa* mounted female (left panel, from top to bottom), as
seen on processed videos (time stamp and trial ID are added by the motion-detection program onto the left top
corner). Median illuminance measurement around the *H. heurippa* mounted female (22'045 lux) was used as
threshold for the 'categorical' model to define poorly and brightly lit conditions. As representatives for poorly lit and
brightly lit conditions, we picked illuminance measures (specifically 6'200 and 132'268 lux) from the bottom and the
top quartile of the distribution. The right panel shows the *H. t. linaresi* mounted female at the same illuminance
values as a comparison. The presented frames were picked from different recording days. Each frame additionally
shows a conspecific male approaching/courting the mounted female and dots used to identify individuals are visible
on their wings. A change in hue of the red patch on *H. heurippa* seems to be visible under different illuminance
conditions, but it should not be forgotten that 1) human vision is vastly different from butterfly vision and 2) that this
is a representation produced by the GoPro camera (GoPro, San Matteo, CA), which is post-processing colour-
composition of frames. [*Figure follows on next page*]

Figure S2: All interaction terms involving male type from fully saturated models (*i.e.* Table S2, model #3 for *H.*
heurippa and *H. t. linaresi* males and Table S4, model #1 for backcross to *H. t. linaresi* males). **A+B**: Two-way
interactions between illuminance at the *H. heurippa* mounted female and male type. Proportion of interactions with
the *H. heurippa* female of *H. heurippa* (red) and *H. t. linaresi* (blue) males (A; resembles Fig. 2A, but comes from a
different underlying model); and differently coloured backcross to *H. t. linaresi* males (B, red males (red), males
without red (blue)). **C+D**: The same for the two-way interactions between illuminance at the *H. t. linaresi* mounted
female and male type **E-H**: Three-way interactions between male type and the two illuminance measures. Male type
is indicated on top of each graph. Colouration of each region in the graph depends on predicted preference for the *H.*
heurippa mounted female at the given light conditions (scale depicted on right). White contour lines show Kernel
densities weighted by preference score, which essentially gives the same information as the colour gradient ('peaks'
are where the landscape is most red). All illuminance-axes are logarithmically scaled. Dashed lines in G and H show
median illuminance measures at the respective axis, used to categorize conditions into 'poorly' and 'brightly' lit for
the categorical models.

 Figure S3: **A-D**: Raw data points of *H. t. linaresi*, *H. heurippa*, non-red and red backcross to *H. t. linaresi* males

(from top to bottom) in a two-dimensional illuminance space. Horizontal dimension shows illuminance at the

mounted *H. heurippa* female, vertical dimension shows illuminance at the mounted *H. t. linaresi* female. Blue dots

indicate responses of males to the mounted *H. t. linaresi* female and red dots indicate responses to the mounted *H.*

heurippa female. Kernel densities are shown for each set of data (blue lines for responses to *H. t. linaresi*, red lines

for responses to *H. heurippa*). **E-H**: the two-dimensional illuminance space was then divided into 15 squares for each

male type and local preferences within each square were calculated. Graphs directly relate to graphs A-D. The higher

the preference for the mounted *H. heurippa* female in a square, the more red the square; the higher the preference for

the *H. t. linaresi* female, the more blue (see scale on the right). Squares without response are white. These local

preferences have to be interpreted in combination with the local sample sizes (as shown in left column), since they

often rely on very few observations. All axes are logarithmically scaled.

Figure S4: Individual trajectories from categorical models. Visual preferences of pure males and hybrid males are

displayed by type. Within each type, data is split depending on illuminance around the *H. heurippa* mounted female,

into poorly and brightly lit. Whenever an individual appeared under both light conditions, it is connected by a

reaction norm line in this plot. Transparency of reaction norm lines within each panel is scaled by the maximum

number of observations available for one individual within this panel. For *H. t. linaresi*, *H. heurippa* and BL males,

posterior means for each EMM are calculated from the same underlying categorical models as used for Fig. 2 and

Fig. 4, respectively. For F_1 and BH males, a categorical model was fit including male type, the illuminance category

around the *H. heurippa* 'female' (determined again by the median measurement) and their interaction. EMMs were

then extracted using the same logic as for the other types. The difference between two groups (in this case between

poorly and brightly lit conditions within each type) is shown by the same Gardner-Altman plot type as used for Fig. 2

and Fig. 4. Small red and blue line located at the right side of each panel show estimators for *H. t. linaresi* and *H.*

heurippa males from Fig. 2B as a reference. [*Figure follows on next page*]

Figure S5: Relative attraction towards *H. heurippa* 'females' (i.e. the proportion of interactions with the mounted *H.*

heurippa 'female' as opposed to the *H. t. linaresi* 'female') for non-red (*bb*) BL males (blue dots) and red (*Bb*) BL

males (red dots). Data and estimators are across all light environments. Note that only preference data with

associated illuminance data were included in the underlying statistical models (a few of the raw data points displayed

here therefore were not considered by the underlying model). The difference between the two groups is shown with

the same logic as used for Fig. 2 and Fig. 4. Small red and blue line located at the right side of each panel show

estimators for *H. t. linaresi* and *H. heurippa* males from Fig. 2B as a reference.

2) Supplementary Tables

Table S1: Information on different experimental hybrid lines. Left column shows F_1 broods from which lines
originated. For the third line, we summarized two F_1 crosses and their subsequent crosses into one hybrid line. These
two F_1 crosses had the same *H. heurippa* father, but a different *H. t. linaresi* mother. Individual counts in brackets
show how many males of each brood were tested in the experiment. 32 other F_1 males from 7 broods were tested in
the experiment and are not mentioned here, as their brood was not involved in generating backcrosses (13
individuals/5 broods of those being from a cross between *H. t. linaresi* mother and *H. heurippa* father and 19
individuals/2 broods being from the reciprocal F_1 cross with *H. heurippa* mother and *H. t. linaresi* father). *H.*
heurippa is abbreviated as *heu* and *H. t. linaresi* as *lin* in the table.

F_1 brood	Backcross to *lin* with male F_1	Backcross to *lin* with female F_1	Backcross to *heu* with male F_1
mother: *lin*, father: *heu*	mother: *lin*, father: F_1	mother: F_1, father: *lin*	mother: *heu*, father: F_1
C18_002 (12 indiv.)	7 broods (45 indiv.)	2 broods (9 indiv.)	2 broods (20 indiv.)
C18_020 (6 indiv.)	3 broods (83 indiv.)	1 brood (15 indiv.)	1 brood (9 indiv.)
C18_034 & C18_036 (half-sib.) (13 indiv.)	4 broods (18 indiv.)	2 broods (10 indiv.)	3 broods (6 indiv.)
C19_014 (-)	1 brood (8 indiv.)	-	-

Table S2: 19 possible fixed-effect structures for the pure males of *H.* heurippa and *H. t.* linaresi (fixed effect 'type').

'I_H' and 'I_L' stand for log-illuminance measures at the *H.* heurippa and the *H. t.* linaresi mounted females,

respectively. A '*' sign in a model formula indicates that all involved variables as well as all possible interactions

between them are included. All models include trial and male ID as random effects. Each model structure has a

LOOIC and WAIC value, which mostly agree. Based on the WAIC value, differences between best model and other

models are calculated (ΔWAIC) as well as a model-weight score.

#	Fixed Effects Term	LOOIC	WAIC	ΔWAIC	Weight$_{WAIC}$
1	~ type * I_H	3773.40	3768.52	0.00	0.29
2	~ type * I_H + I_L	3774.63	3769.68	1.16	0.16
3	~ type * I_H * I_L	3775.20	3770.08	1.56	0.13
4	~ type * I_H + I_H * I_L	3775.99	3771.04	2.52	0.08
5	~ type * I_H + type * I_L	3776.53	3771.52	3.01	0.06
6	~ type * I_H + type * I_L + I_H * I_L	3776.62	3771.56	3.04	0.06
7	~ type	3777.82	3773.07	4.55	0.03
8	~ type + I_L	3778.10	3773.20	4.68	0.03
9	~ type + I_H	3778.11	3773.33	4.81	0.03
10	~ I_L	3778.50	3773.62	5.10	0.02
11	~ type * I_L	3778.83	3773.95	5.43	0.02
12	~ 1	3779.24	3774.44	5.92	0.02
13	~ I_H	3779.45	3774.58	6.06	0.01
14	~ type + I_H + I_L	3779.75	3774.87	6.35	0.01
15	~ I_H + I_L	3780.58	3775.58	7.06	0.01
16	~ type + I_H * I_L	3780.77	3775.80	7.28	0.01
17	~ type * I_L + I_H	3780.82	3775.99	7.47	0.01
18	~ type * I_L + I_H * I_L	3781.61	3776.60	8.09	0.01
19	~ I_H * I_L	3781.85	3776.78	8.26	0.00

915 **Table S3**: Posterior means and 95% equal-tailed credible intervals for male types from Fig. 2, Fig. S5 and Fig. 4.

916 Minus ("-") in the second column stands for a poorly lit mounted female, a plus ("+") for a brightly lit one.

Male type	Light environment	Mean	Lower	Upper
H. t. linaresi (left in Fig. 2B)	Across all	0.447	0.383	0.513
H. heurippa (right in Fig. 2B)	Across all	0.515	0.449	0.582
H. t. linaresi (left in Fig. 2C)	*H. heurippa* -	0.464	0.391	0.540
H. heurippa (right in Fig. 2C)	*H. heurippa* -	0.456	0.385	0.527
H. t. linaresi (left in Fig. 2D)	*H. heurippa* +	0.437	0.362	0.514
H. heurippa (right in Fig. 2D)	*H. heurippa* +	0.572	0.493	0.650
BL without red (left in Fig. S5)	Across all	0.461	0.414	0.508
BL with red (right in Fig. S5)	Across all	0.493	0.444	0.543
BL without red (left in Fig. 4A)	*H. heurippa* - / *H. t. linaresi* -	0.460	0.406	0.515
BL with red (right in Fig. 4A)	*H. heurippa* - / *H. t. linaresi* -	0.475	0.419	0.531
BL without red (left in Fig. 4B)	*H. heurippa* - / *H. t. linaresi* +	0.477	0.401	0.554
BL with red (right in Fig. 4B)	*H. heurippa* - / *H. t. linaresi* +	0.443	0.359	0.528
BL without red (left in Fig. 4C)	*H. heurippa* + / *H. t. linaresi* -	0.426	0.350	0.505
BL with red (right in Fig. 4C)	*H. heurippa* + / *H. t. linaresi* -	0.538	0.461	0.616
BL without red (left in Fig. 4D)	*H. heurippa* + / *H. t. linaresi* +	0.438	0.384	0.491
BL with red (right in Fig. 4D)	*H. heurippa* + / *H. t. linaresi* +	0.504	0.446	0.563

Table S4: Posterior means and 95% equal-tailed credible intervals for estimates of proportions of each mating
outcome from tetrad experiments. First row is based on empirically measured data, the other rows are stochastic
predictions based on mounted female data from all light environments, from a poorly lit *H. heurippa* mounted female
and a brightly lit one, respectively.

Estimates based on	*H. heurippa* ♀ *H. t. linaresi* ♂	*H. t. linaresi* ♀ *H. t. linaresi* ♂	*H. heurippa* ♀ *H. heurippa* ♂	*H. t. linaresi* ♀ *H. heurippa* ♂
tetrad experiment	0.168 [0.099; 0.253]	0.157 [0.09; 0.239]	0.405 [0.305; 0.508]	0.27 [0.183; 0.366]
mounted females: across all	0.149 [0.084; 0.23]	0.184 [0.111; 0.27]	0.349 [0.255; 0.45]	0.329 [0.236; 0.429]
mounted females: poorly lit *H. heurippa*	0.152 [0.086; 0.233]	0.181 [0.109; 0.266]	0.315 [0.224; 0.414]	0.363 [0.267; 0.465]
mounted females: brightly lit *H. heurippa*	0.146 [0.082; 0.226]	0.187 [0.114; 0.273]	0.387 [0.289; 0.489]	0.291 [0.203; 0.389]

Table S5: 19 possible fixed-effect structures for the BL males (fixed effect 'type' refers here to wing colour
phenotype of hybrids). 'I_H' and 'I_L' stand for log-illuminance measures at the *H. heurippa* and the *H. t. linaresi*
mounted females, respectively. A '*' sign in a model formula indicates that all involved variables as well as all
possible interactions between them are included. All models include trial, male ID and brood as random effects. Each
model structure has a LOOIC and WAIC value, which mostly agree. Based on the WAIC value, differences between
best model and other models are calculated (ΔWAIC) as well as a model-weight score.

#	Fixed Effects Term	LOOIC	WAIC	ΔWAIC	Weight$_{WAIC}$
1	~ type * I_H * I_L	6815.06	6811.69	0.00	0.38
2	~ type * I_H + type * I_L + I_H * I_L	6815.98	6812.60	0.91	0.24
3	~ type * I_H + type * I_L	6817.15	6813.84	2.15	0.13
4	~ type * I_H + I_H * I_L	6817.82	6814.43	2.74	0.10
5	~ type * I_H	6820.15	6816.86	5.17	0.03
6	~ type * I_H + I_L	6820.78	6817.48	5.79	0.02
7	~ type + I_H * I_L	6821.33	6817.96	6.27	0.02
8	~ I_H * I_L	6822.03	6818.67	6.98	0.01
9	~ type	6821.94	6818.68	6.99	0.01
10	~ type * I_L + I_H * I_L	6822.28	6818.95	7.27	0.01
11	~ 1	6822.46	6819.18	7.50	0.01
12	~ type + I_H	6822.96	6819.70	8.01	0.01
13	~ type + I_L	6823.18	6819.94	8.26	0.01
14	~ I_H	6823.50	6820.19	8.50	0.01
15	~ I_L	6823.80	6820.52	8.84	0.00
16	~ type + I_H + I_L	6824.36	6821.03	9.35	0.00
17	~ type * I_L	6825.21	6821.92	10.23	0.00
18	~ I_H + I_L	6825.20	6821.93	10.24	0.00
19	~ type * I_L + I_H	6825.55	6822.23	10.54	0.00

966
967