Genome-wide bioinformatic analyses predict key host and viral factors in SARS-CoV-2 pathogenesis

Mariana G. Ferrarini[1,§], Avantika Lal[2,§], Rita Rebollo[1], Andreas Gruber[3], Andrea Guarracino[4], Itziar Martinez Gonzalez[5], Taylor Floyd[6], Daniel Siqueira de Oliveira[7], Justin Shanklin[8], Ethan Beausoleil[8], Taneli Pusa[9], Brett E. Pickett[8,#] Vanessa Aguiar-Pulido[6,#]

1 University of Lyon, INSA-Lyon, INRA, BF2I, Villeurbanne, France
2 NVIDIA Corporation, Santa Clara, CA, USA
3 Oxford Big Data Institute, Nuffield Department of Medicine, University of Oxford, Oxford, UK
4 Centre for Molecular Bioinformatics, Department of Biology, University Of Rome Tor Vergata, Rome, Italy
5 Amsterdam UMC, Amsterdam, The Netherlands
6 Center for Neurogenetics, Weill Cornell Medicine, Cornell University, New York, NY, USA
7 Laboratoire de Biométrie et Biologie Evolutive, Université de Lyon; Université Lyon 1; CNRS; UMR 5558, Villeurbanne, France
8 Brigham Young University, Provo, UT, USA
9 Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Belvaux, Luxembourg

§: These authors contributed equally
#: Corresponding authors
Keywords: SARS-CoV-2, COVID-19, gene expression, RNA-seq, RNA-binding proteins, host-pathogen interaction, transcriptomics

Abstract

The novel betacoronavirus named Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) caused a worldwide pandemic (COVID-19) after initially emerging in Wuhan, China. Here we applied a novel, comprehensive bioinformatic strategy to public RNA sequencing and viral genome sequencing data, to better understand how SARS-CoV-2 interacts with human cells. To our knowledge, this is the first meta-analysis to predict host factors that play a specific role in SARS-CoV-2 pathogenesis, distinct from other respiratory viruses. We identified differentially expressed genes, isoforms and transposable element families specifically altered in SARS-CoV-2 infected cells. Well-known immunoregulators including *CSF2*, *IL-32*, *IL-6* and *SERPINA3* were differentially expressed, while immunoregulatory transposable element families were overexpressed. We predicted conserved interactions between the SARS-CoV-2 genome and human RNA-binding proteins such as hnRNPA1, PABPC1 and eIF4b, which may play important roles in the viral life cycle. We also detected four viral sequence variants in the spike, polymerase, and nonstructural proteins that correlate with severity of COVID-19. The host factors we identified likely represent important mechanisms in the disease profile of this pathogen, and could be targeted by prophylactics and/or therapeutics against SARS-CoV-2.

Graphical Abstract

Introduction

SARS-CoV-2 infects human cells by binding to the angiotensin-converting enzyme 2 (ACE2) receptor [82]. Recent studies have sought to understand the molecular interactions between SARS-CoV-2 and infected cells [24], some of which have quantified gene expression changes in patient samples or cultured lung-derived cells infected by this virus [10, 44, 80]. These studies are essential to understanding the mechanisms of pathogenesis and immune response which can facilitate the development of treatments for COVID-19 [34, 52, 85].

Viruses generally trigger a drastic host response during infection. A subset of these specific changes in gene regulation is associated with viral replication, and therefore can pinpoint potential drug targets. In addition, transposable element (TE) overexpression has been observed upon viral infection [48], and TEs have been actively implicated in gene regulatory networks related to immunity [15]. Moreover, SARS-CoV-2 is a virus with a positive-sense, single-stranded, monopartite RNA genome. Such viruses are known to co-opt host RNA-binding proteins (RBPs) for diverse processes including viral replication, translation, viral RNA stability, assembly of viral protein complexes, and regulation of viral protein activity [22, 43].

In this work we identified a signature of altered gene expression that is consistent across published datasets of SARS-CoV-2 infected human lung cells. We present extensive results from functional analyses (signaling pathway enrichment, biological functions, transcript isoform usage, metabolic flux prediction, and TE overexpression) performed upon the genes that are differentially expressed during SARS-CoV-2 infection [10]. We also predict specific interactions between the SARS-CoV-2 RNA genome and human RBPs that may be involved in viral replication, transcription or translation, and identify viral sequence variations that are significantly associated with increased pathogenesis in humans. Knowledge of these molecular and genetic mechanisms is important to understand the SARS-CoV-2 pathogenesis and to improve the future development of effective prophylactic and therapeutic treatments.

Results

We designed a comprehensive bioinformatics workflow to identify relevant host-pathogen interactions using a complementary set of computational analyses (Figure 1). First, we carried out an exhaustive analysis of differential gene expression in human lung cells infected by SARS-CoV-2 or other respiratory viruses, identifying gene, isoform- and pathway-level responses that specifically

characterize SARS-CoV-2 infection. Second, we predicted putative interactions between the ₃₁ SARS-CoV-2 RNA genome and human RBPs. Third, we identified a subset of these human RBPs ₃₂ which were also differentially expressed in response to SARS-CoV-2. Finally, we predicted four viral ₃₃ sequence variants that could play a role in disease severity. ₃₄

Figure 1. Overview of the bioinformatic workflow applied in this study. ₃₆

SARS-CoV-2 infection elicits a specific gene expression and pathway signature in human cells

We wanted to identify genes that were differentially expressed across multiple SARS-CoV-2 infected ₃₉ samples and not in samples infected with other respiratory viruses. As a primary dataset, we ₄₀ selected GSE147507 [10], which includes gene expression measurements from three cell lines derived ₄₁ from the human respiratory system (NHBE, A549, Calu-3) infected either with SARS-CoV-2, ₄₂ influenza A virus (IAV), respiratory syncytial virus (RSV), or human parainfluenza virus 3 (HPIV3), ₄₃ with different multiplicity of infection (MOI). We also analyzed an additional dataset GSE150316, ₄₄ which includes RNA-seq extracted from formalin fixed, paraffin embedded (FFPE) histological ₄₅ sections of lung biopsies from COVID-19 deceased patients and healthy individuals (see Figure 2A ₄₆ and Materials and Methods for further details). ₄₇

Hence, we retrieved 41 differentially expressed genes (DEGs) that showed significant and ₄₈ consistent expression changes in at least three datasets from cell lines infected with SARS-CoV-2, ₄₉ and that were not significantly affected in cell lines infected with other viruses within the same ₅₀ dataset (Supplementary Table 1A). To these, we added 23 genes that showed significant and ₅₁ consistent expression changes in two of four cell line datasets infected with SARS-CoV-2 and at least ₅₂ one lung biopsy sample from a SARS-CoV-2 patient. Results coming from FFPE sections were less ₅₃ consistent presumably due to the collection of biospecimens from different sites within the lung. ₅₄ Thus, the final set consisted of 64 DEGs: 48 up-regulated and 16 downregulated of which 38 had an ₅₅ absolute Log2FC > 1 in at least one dataset (relevant genes from this list are shown in Table 1). ₅₆

SERPINA3, an antichymotrypsin which was proposed as an interesting candidate for the ₅₇ inhibition of viral replication [13], was the only gene specifically upregulated in the four cell line ₅₈ datasets tested (Table 1). Other interesting up-regulated genes were the amidohydrolase *VNN2*, the ₅₉

pro-fibrotic gene *PDGFB*, the beta-interferon regulator *PRDM1* and the proinflammatory cytokines *CSF2* and *IL-32*. *FKBP5*, a known regulator of NF-kB activity, was among the consistently downregulated genes. We also generated additional lists of DEGs that met different filtering criteria (Supplementary Table 1B, see Supplementary File 1 for the complete DEG results for each dataset).

In order to better understand the underlying biological functions and molecular mechanisms associated with the observed DEGs, we performed a hypergeometric test to detect statistically significant overrepresented gene ontology (GO) terms among the DEGs having an absolute Log2FC > 1 in each dataset separately.

Table 1. Log2FC for selected genes that showed significant up-or down-regulation in SARS-CoV-2 infected samples (FDR-adjusted p-value < 0.05), and not in samples infected with the other viruses tested. Log2FC values are only provided for statistically significant samples.

| Gene | Cell Type and MOI | | | | Biopsies | |
| | A549 MOI 0.2 | A549 | Calu-3 | NHBE | Case 1 | Case 3 |
			MOI 2			
VNN2		6.18	0.42	6.13		
CSF2		3.56	7.30	2.70		
WNT7A		4.99	0.79	0.45		
PDZK1IP1		1.72	0.70	2.28		
SERPINA3	0.49	1.39	0.77	1.44		
RHCG		1.51	2.02	1.33	2.53	
IL32		1.64	1.23	1.21		
PDGFB		1.91	1.75	1.00		
ALDH1A3		1.09	1.32	0.39		
TLR2		1.63	0.89	0.84		
G0S2		0.66	3.79	0.83		
NRCAM		0.73	1.82	0.78		
SERPINB1		0.61	1.17	0.72		
PRDM1		0.82	3.49	0.59		
MT-TN		0.55	1.70	0.33		
ATF4		0.79	1.07	0.26		
BHLHE40		0.75	1.56	0.18		
PTPN12	0.48	0.97	1.23			
GPCPD1	0.36	0.94	1.69			
DUSP16	0.33	0.41	1.43			
FKBP5		-0.39		-0.36	-1.47	-2.14
DAP		-0.18	-0.61		-1.16	
FECH		-0.27	-0.36		-1.54	
MT-CYB		-0.30	-0.26		-3.68	
EIF4A1		-0.33	-0.63		-1.85	
POLE4	-0.23	-0.82	-1.24			
DDX39A	-0.23	-1.27	-0.54			
CENPP		-0.36	-0.40	-0.38		
TMEM50B		-0.48	-0.59	-0.53		
HPS1	-0.28	-0.31	-0.62			
SNX8	-0.30	-0.43	-0.56			

Consistent with the findings of Blanco-Melo et al. [10], GO enrichment analysis returned terms associated with immune system processes, Pi3K/AKT signaling pathway, response to cytokine, stress and virus, among others (see Supplementary File 2 for complete results). In addition, we report 285 GO terms common to at least two cell line datasets infected with SARS-CoV-2, and absent in the response to other viruses (Figure 2B, Supplementary Table 2A), including neutrophil and granulocyte activation, interleukin-1-mediated signaling pathway, proteolysis, and stress activated signaling cascades.

Figure 2. Overview of the RNA-seq based results specific to SARS-CoV-2 which were not detected in the other viral infections (IAV, HPIV3 and RSV). (A) Representation of the RNA-seq studies used in our analyses. (B) Non-redundant functional enrichment of DEGs. Here we report a subset of non-redundant reduced terms consistently enriched in more than one SARS-COV-2 cell line which were not detected in the other viruses' datasets. We added generic categories of immunity, metabolism, cellular processes and signaling/epigeneitcs to the GO terms as colored dots. (C) Top 20 differentially expressed isoforms (DEIs) in SARS-CoV-2 infected samples. Y-axis denotes the differential usage of isoforms (dIF) whereas x-axis represents the overall log2FC of the corresponding gene. Thus, DEIs also detected as DEGs by this analysis are depicted in blue. (D) The upper right diagram depicts different manners by which TE family overexpression might be detected. While TEs may indeed be autonomously expressed, the old age of most TEs detected points toward either being part of a gene (exonization or alternative promoter), or a result of pervasive transcription. We report the functional enrichment for neighboring genes of differentially expressed TEs (DETEs) specifically upregulated in SARS-CoV-2 Calu-3 and A549 cells (MOI 2). The same categories used in subfigure (B) were attributed to the GO terms reported here.

Next, we wanted to pinpoint intracellular signaling pathways that may be modulated specifically ₉₅ during SARS-CoV-2 infection. A robust signaling pathway impact analysis (SPIA) enabled us to ₉₆ identify 30 pathways, including many involved in the host immune response, that were significantly ₉₇ enriched among differentially expressed genes in at least one virus-infected cell line dataset ₉₈ (Supplementary Table 3). More importantly, we predicted four pathways to be specific to ₉₉ SARS-CoV-2 infection and observed that the significant pathways differed by cell type and ₁₀₀ multiplicity of infection. The significant results included only one term common to A549 (MOI 0.2) ₁₀₁ and Calu-3 cells (MOI 2), namely the interferon alpha/beta signaling. Additionally, we found the ₁₀₂ amoebiasis (A549 cells, MOI 0.2), the p75(NTR)-mediated and the trka receptor signaling pathways ₁₀₃ (A549 cells, MOI 2) as significantly impacted. ₁₀₄

We also used a classic hypergeometric method as a complementary approach to our SPIA ₁₀₅ pathway enrichment analysis. While there were generally higher numbers of significant results using ₁₀₆ this method, we observed that the vast majority of enriched terms (FDR < 0.05) described ₁₀₇ infections with various pathogens, innate immunity, metabolism, and cell cycle regulation ₁₀₈ (Supplementary Table 3). Interestingly, we were able to detect enriched KEGG pathways common to ₁₀₉ at least two SARS-CoV-2 infected cell types and absent from the other virus-infected datasets ₁₁₀ (Figure 2B, Supplementary Table 2B). These included pathways related to infection, cell cycle, ₁₁₁ endocytosis, signalling pathways, cancer and other diseases. ₁₁₂

SARS-CoV-2 infection results in altered lipid-related metabolic fluxes ₁₁₃

To integrate the gene expression changes with metabolic activity in response to virus infection, we ₁₁₄ projected the transcriptomic data onto the human metabolic network [75]. This analysis detected ₁₁₅ common decreased fluxes in inositol phosphate metabolism in both A549 and Calu-3 cells infected ₁₁₆ with SARS-CoV-2 at a MOI of 2 (Supplementary Table 4). The consensus solution (obtained taking ₁₁₇ into account the enumeration of all solutions) in A549 cells (MOI 2) also recovered decreased fluxes ₁₁₈ in several lipid pathways: fatty acid, cholesterol, sphingolipid, and glycerophospholipid. In addition, ₁₁₉ we detected an increased flux common to A549 and Calu-3 cell lines in reactive oxygen species (ROS) ₁₂₀ detoxification, in accordance with previous terms recovered from functional enrichment analyses. ₁₂₁

SARS-CoV-2 infection induced an isoform switch of genes associated ₁₂₂ with immunity and mRNA processing ₁₂₃

We wanted to analyze changes in transcript isoform expression and usage associated with ₁₂₄ SARS-CoV-2 infection, as well as to predict whether these changes might result in altered protein ₁₂₅ function. We identified isoforms experiencing a switch in usage greater than or equal to 30% in ₁₂₆ absolute value, and retrieved those with a Bonferroni-adjusted p-value less than 0.05. After ₁₂₇ calculating the difference in isoform usage (dIF) per gene (in each condition), we performed ₁₂₈ predictive functional consequence and alternative splicing analyses for all isoforms globally as well as ₁₂₉ at the individual gene level. ₁₃₀

We observed 3,569 differentially expressed isoforms (DEIs) across all samples (Supplementary ₁₃₁ Figure 1A, Supplementary Table 5A). Results indicate that isoforms from A549 cells infected with ₁₃₂ RSV, IAV and HPIV3 exhibited significant differences in biological events such as complete open ₁₃₃ reading frame (ORF) loss, shorter ORF length, intron retention gain and decreased sensitivity to ₁₃₄ nonsense mediated decay (Supplementary Figure 1B). These conditions also displayed various ₁₃₅ changes in splicing patterns, ranging from loss of exon skipping events, changes in usage of ₁₃₆ alternative transcription start and termination sites, and decreased alternative 5' and 3' splice sites ₁₃₇ (Supplementary Figure 1C). ₁₃₈

In contrast, isoforms from SARS-CoV-2 infected samples displayed no significant global changes ₁₃₉ in biological consequences or alternative splicing events between conditions (Supplementary Figures ₁₄₀ 1A and 1B respectively). Trends indicated transcripts in SARS-CoV-2 samples experienced decreases ₁₄₁ in ORF length, numbers of domains, coding capability, intron retention and nonsense mediated ₁₄₂ decay (Supplementary Figure 1A). These biological consequences may result from increased multiple ₁₄₃

exon skipping events and alternative transcription start sites via alternative 5' acceptor sites   144
(Supplementary Figure 1B). While not significant, these trends implicate that the SARS-CoV-2 virus   145
may globally trigger host cell machinery to generate shorter isoforms that, while not shuttled for   146
degradation, either do not produce functional proteins or produce alternative aberrant proteins not   147
utilized in non-SARS-CoV-2 tissue conditions.   148

Despite the lack of global biological consequence and splicing changes, individual isoforms from   149
SARS-CoV-2 infected samples experienced significant changes in gene expression and isoform usage   150
(Figure 2C). Top-expressing genes were associated with cellular processes such as immune response   151
and antiviral activity (*IFI44L*, *IL6*, *MX1*, *TRIM5*), transcription and mRNA processing (*DDX10*,   152
HNRNPA3F6, *JMJD7*, *ZNF487*, *ZNF599*) and cell cycle and survival (*BCL2L2-PABPN1*, *CDCA3*)   153
(Supplementary Table 5B). Similarly, significant genes from non-SARS-CoV-2 samples were   154
associated with processes such as immune cell development and response (*ADCY7*, *BATF2*, *C9orf72*,   155
ETS1, *GBP2*, *IFIT3*), transcription regulation and DNA repair (*ABHD14B*, *ATF3*, *IFI16*,   156
POLR2J2, *SMUG1*, *ZNF19*, *ZNF639*), mitochondrial function (*ATP5E*, *BCKDH8*, *TST*, *TXNRD2*),   157
and GTPase activity (*GBP2*, *RAP1GAP*, *RGS20*, *RHOBTB2*) (Supplementary Figure 1D,   158
Supplementary Table 5B).   159

Upon further inspection, we noticed that *IL-6*, a gene encoding a cytokine involved in acute and   160
chronic inflammatory responses, displayed 3 and 4-fold increases in expression in NHBE and A549   161
cells, respectively (infected with a MOI of 2) (Supplementary Figure 1B). To date, the Ensembl   162
Genome Reference Consortium has identified 9 *IL-6* isoforms in humans, with the traditional   163
transcript having 6 exons (*IL6-204*), 5 of which contain coding elements. NHBE cells expressed 4   164
known *IL-6* isoforms, while A549 cells expressed 1 unknown and 6 known isoforms. When evaluating   165
the actual isoforms used across conditions, NHBE cells used 3 out of 4 isoforms observed, while A549   166
cells used all 7 observed isoforms. Isoform usage is evaluated based on isoform fraction (IF), or the   167
percentage of an isoform found relative to all other identified isoforms associated with a specific gene.   168
For example, in the case of NHBE SARS-CoV-2 samples, the IF for the *IL6-201* isoform = 0.75,   169
IL6-204 = 0.05, *IL6-206* = 0.09, *IL6-209* = 0.06, and the sum of these IF values = 0.95, or 95%   170
usage of the *IL-6* gene. Both SARS-CoV-2 samples exhibited exclusive usage of non-canonical   171
isoform *IL6-201*, and inversely, mock samples almost exclusively utilized the *IL6-204* transcript. In   172
NHBE infected cells, isoform *IL6-201* experienced a significant increase in usage (dIF = 0.75) and   173
IL6-204 a significant decrease in usage (dIF = -0.95) when compared to mock conditions. Similarly,   174
isoform *IL6-201* in A549 infected cells experienced an increase in usage (dIF = 0.58), while uses of   175
all other isoforms remained non-significant in comparison to mock conditions.   176

Overexpression of TE families close to immune-related genes upon   177
SARS-CoV-2 infection   178

In order to estimate the expression of TE families and their possible roles in SARS-CoV-2 infection,   179
we mapped the RNA-seq reads against all annotated human TE families and detected DETEs   180
(Supplementary File 3). We found 68 common TE families upregulated in SARS-CoV-2 infected   181
A549 and Calu-3 cells (MOI 2). From this list, we excluded all TE families detected in A549 cells   182
infected with the other viruses. This allowed us to identify 16 families that were specifically   183
upregulated in Calu-3 and A549 cells infected with SARS-CoV-2 and not in the other viral infections.   184

The 16 families identified were MER77B, MamRep4096, MLT2C2, PABL_A, Charlie9, MER34A,   185
L1MEg1, LTR13A, L1MB5, MER11C, MER41B, LTR79, THE1D-int, MLT1I, MLT1F1,   186
MamRep137. Most of the TE families uncovered are ancient elements, incapable of transposing, or   187
harboring intrinsic regulatory sequences [36, 55, 68]. Eleven of the 16 TE families specifically   188
upregulated in SARS-COV-2 infected cells are long terminal repeat (LTR) elements, and include well   189
known TE immune regulators. For instance, the MER41B (primate specific TE family) is known to   190
contribute to Interferon gamma inducible binding sites (bound by STAT1 and/or IRF1) [14, 64].   191
Other LTR elements are also enriched in STAT1 binding sites (MLT1L) [14], or have been shown to   192
act as cellular gene enhancers (LTR13A [16, 31]).   193

Given the propensity for the TE families detected to impact nearby gene expression, we further ²⁹⁴ investigated the functional enrichment of genes near upregulated TE families (+- 5kb upstream, 1kb ¹⁹⁵ downstream). We detected GO functional enrichment of several immunity-related terms (e.g. MHC ¹⁹⁶ protein complex, antigen processing, regulation of dendritic cell differentiation, T-cell tolerance ¹⁹⁷ induction), metabolism related terms (such as regulation of phospholipid catabolic process), and ¹⁹⁸ more interestingly a specific human phenotype term called "Progressive pulmonary function ¹⁹⁹ impairment" (Figure 2D). Even though we did not limit our search only to neighboring genes which ²⁰⁰ were also DE, we found several similar (and very specific) enriched terms in both analyses, for ²⁰¹ instance related to immune response, endosomes, endoplasmic reticulum, vitamin (cofactor) ²⁰² metabolism, among others. This result supports the idea that some responses during infection could ²⁰³ be related to TE-mediated transcriptional regulation. Finally, when we searched for enriched terms ²⁰⁴ related to each one of the 16 families separately, we also detected immunity related enriched terms ²⁰⁵ such as regulation of interleukins, antigen processing, TGFB receptor binding and temperature ²⁰⁶ homeostasis (Supplementary File 4). It is important to note that given the old age of some of the ²⁰⁷ TEs detected, overexpression might be associated with pervasive transcription, or inclusion of TE ²⁰⁸ copies within unspliced introns (see upper box in Figure 2D). ²⁰⁹

The SARS-CoV-2 genome is enriched in binding motifs for 40 human ²¹⁰ RBPs, most of them conserved across SARS-CoV-2 genome isolates ²¹¹

Our next aim was to predict whether any host RNA binding proteins interact with the viral genome. ²¹² To do so, we first filtered the AtTRACT database [23] to obtain a list of 102 human RBPs and 205 ²¹³ associated Position Weight Matrices (PWMs) describing the sequence binding preferences of these ²¹⁴ proteins. We then scanned the SARS-CoV-2 reference genome sequence to identify potential binding ²¹⁵ sites for these proteins. Figure 3 illustrates our analysis pipeline. ²¹⁶

We identified 99 human RBPs with 11,897 potential binding sites in the SARS-CoV-2 ²¹⁷ positive-sense genome. Since the SARS-CoV-2 genome produces negative-sense intermediates as part ²¹⁸ of the replication process [35], we also scanned the negative-sense molecule, where we found 11,333 ²¹⁹ potential binding sites for 96 RBPs (Supplementary Table 6). ²²⁰

To find RBPs whose binding sites occur in the SARS-CoV-2 genome more often than expected by ²²¹ chance, we repeatedly scrambled the genome sequence to create 1,000 simulated genome sequences ²²² with an identical nucleotide composition to the SARS-CoV-2 genome sequence (30% A, 18% C, 20% ²²³ G, 32% T). We used these 1,000 simulated genomes to determine a background distribution of the ²²⁴ number of binding sites found for a specific RBP. This allowed us to pinpoint RBPs with ²²⁵ significantly more or significantly fewer binding sites in the actual SARS-CoV-2 genome than ²²⁶ expected based on the background distribution (two-tailed z-test, FDR-corrected P < 0.01). To ²²⁷ retrieve RBPs whose motifs were enriched in specific genomic regions, we also repeated this analysis ²²⁸ independently for the SARS-CoV-2 5'UTR, 3'UTR, intergenic regions, and for the sequence from the ²²⁹ negative sense molecule. Motifs for 40 human RBPs were found to be enriched in at least one of the ²³⁰ tested genomic regions, while motifs for 23 human RBPs were found to be depleted in at least one of ²³¹ the tested regions (Supplementary Table 7). ²³²

We next examined whether any of the 6,936 putative binding sites for these 40 enriched RBPs ²³³ were conserved across SARS-CoV-2 isolates. We found that 6,581 putative binding sites, ²³⁴ representing 34 RBPs, were conserved across more than 95% of SARS-CoV-2 genome sequences in ²³⁵ the GISAID database (≥ 26,213 out of 27,592 genomes). However, this is of limited significance as ²³⁶ RBP binding sites in coding regions are likely to be conserved due to evolutionary pressure on ²³⁷ protein sequences rather than RBP binding ability. We therefore repeated this analysis focusing only ²³⁸ on putative RBP binding sites in the SARS-CoV-2 UTRs and intergenic regions. We found 124 ²³⁹ putative RBP binding sites for 21 enriched RBPs in the UTRs and intergenic regions. Of these, 50 ²⁴⁰ putative RBP binding sites for 17 RBPs were conserved in >95% of the available genome sequences; ²⁴¹ 6 in the 5'UTR, 5 in the 3'UTR, and 39 in intergenic regions (Supplementary Table 8). ²⁴²

Subsequently, we interrogated publicly available data to validate the putative SARS-CoV-2 / ²⁴³

RBP interactions (Supplementary Table 9). According to GTEx data [25], 39 of the 40 enriched
RBPs and all 23 of the depleted RBPs were expressed in human lung tissue. Further, 31 of 40
enriched RBPs and 22 of 23 depleted RBPs were co-expressed with the ACE2 and TMPRSS2
receptors in single-cell RNA-seq data from human lung cells (GSE122960; [25,62]), indicating that
they are present in cells that are susceptible to SARS-CoV-2 infection. We next checked whether any
of these RBPs are known to interact with SARS-CoV-2 proteins and found that human poly-A
binding proteins C1 and C4 (PABPC1 and PABPC4) bind to the viral N protein [24]. Thus, it is
conceivable that these RBPs interact with both the SARS-CoV-2 RNA and proteins. Finally, we
combined these results with our analysis of differential gene expression to identify SARS-CoV-2
interacting RBPs that also show expression changes upon infection. The results of this analysis are
summarized for selected RBPs in Table 2.

Figure 3. Workflow and selected results for analysis of potential binding sites for human RNA-binding
proteins in the SARS-CoV-2 genome.

Motif enrichment in SARS-CoV-2 differs from related coronaviruses

We repeated the above analysis to calculate the enrichment and depletion of RBP-binding motifs in the genomes of two related coronaviruses: the SARS-CoV virus (Supplementary Table 10) that caused the SARS outbreak in 2002-2003, and RaTG13 (Supplementary Table 11), a bat coronavirus with a genome that is 96% identical with that of SARS-CoV-2 [4, 84].

We found that the pattern of enrichment and depletion of RBP binding motifs in SARS-CoV-2 is different from that of the other two viruses. Specifically, the SARS-CoV-2 genome is uniquely enriched for binding sites of CELF5 in its 5'UTR, PPIE on its 3'UTR, and ELAVL1 in the viral negative-sense RNA molecule. These three proteins are involved in RNA metabolism and are important for RNA stability (ELAVL1, CELF5) and processing (PPIE). Despite the high sequence identity between the two genomes, the single binding site for CELF5 on the SARS-CoV-2 5'UTR is conserved in 97% of available SARS-CoV-2 genome sequences but absent in the 5'UTR of RaTG13.

Table 2. Selected conserved human RBPs predicted to interact with the SARS-CoV-2 genome along with experimental information.

RPB	DE Analysis*1			Experimental evidence in human datasets				RBP binding site prediction	
	A549 LogFC	Calu-3 LogFC	SARS-CoV-2 Specific DEG	GTEx Lung Tissue (TPM)	scRNA*2	PPI Map*3	Interaction with viral RNA*4	Conserved*5	Region
HNRNPA1		-0.32	☐	331.336	✓		✓	✓	3'UTR
HNRNPA2B1	-1.08	-0.29	☐	539.829	✓		✓	✓	
PABPC1	0.72	0.44	✓	448.025	✓	N	✓	✓	
PABPC4	0.30	-0.28	☐	103.082	✓	N	✓	✓	
PPIE		-0.27	☐	13.827	✓		☐	✓	
CELF5	0.56		☐	0.079	✓		☐	✓	5'UTR
FMR1		0.75	☐	21.435	✓		☐	✓	
RBM24	0.34		☐	1.412	☐		☐	✓	
EIF4B	0.53	0.64	☐	170.303	✓		✓	✓	Intergenic
ELAVL1		-0.31	☐	27.440	✓		☐	✓	
PABPC1	0.72	0.44	✓	448.025	✓	N	✓	✓	
PPIE		-0.27	☐	13.827	✓		☐	✓	
TIA1	0.34	0.41	✓	46.934	✓		☐	✓	
TIAL1	0.25		☐	40.593	☐		☐	✓	

*1 LogFC reported only if padj < 0.05
*2 scRNA expression in ACE+ and TMPRSS2+ lung cells: dataset GSE122960
*3 PPI Map: Experimental map of protein-protein interactions between human and viral proteins (Gordon *et al.*, 2020)
*4 Preprint: Experimental study revealing proteins interacting with SARS-CoV-2 RNA in a human liver cell line (Schmidt *et al.*, 2020)
*5 Conserved in SARS-CoV-2 genomes

A subset of viral genome variants correlate with increased COVID-19 severity

To test whether any viral sequence variants were associated with a change in disease severity in human hosts, we analyzed 1,511 complete SARS-CoV-2 genomes that had associated clinical metadata. The FDR-corrected statistical results from this analysis revealed four nucleotide

variations that were significantly associated with a change in viral pathogenesis. Three of these nucleotide changes resulted in non-synonymous variations at the amino acid level, while the last one was silent at the amino acid level. The first position was a $T \rightarrow G$ (L37F) substitution located in the Nsp6 coding region (p < 1.48E-5), the second position was a $C \rightarrow T$ (P323L) substitution located in the RNA-dependent RNA polymerase coding region (p < 2.01E-4), the third position was an $A \rightarrow G$ (D614G) substitution located in the spike coding region (p < 1.61E-4), and the fourth was a synonymous $C \rightarrow T$ substitution located in the Nsp3 coding region (p < 1.77E-4). As a further validation step, we performed the same analysis comparing viral sequence variants against potential confounders, such as the biological sex or age group of the patients. These comparisons validated that these four positions were only identified as significant in the results of the disease severity analysis.

Discussion

Airway epithelial cells are the primary entry points for respiratory viruses and therefore constitute the first producers of inflammatory signals that, in addition to their antiviral activity, promote the initiation of the innate and adaptive immune responses. Here, we report the results of a complementary panel of analyses that enable a better understanding of host-pathogen interactions which contribute to SARS-CoV-2 replication and pathogenesis in the human respiratory system. Moreover, we propose already established along with novel human factors exclusively detected in SARS-CoV-2 infected cells by our analyses that might be relevant in the context of COVID-19 and which are worth being further investigated at an experimental level (Figure 4).

The *CSF2* gene, which encodes the Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF), was among the most highly up-regulated genes in SARS-CoV-2 infected cells. GM-CSF induces survival and activation in mature myeloid cells such as macrophages and neutrophils. However, GM-CSF is considered more proinflammatory than other members of its family, such as G-CSF, and is associated with tissue hyper-inflammation [50]. In accordance with our results, high levels of GM-CSF were found in the blood of severe COVID-19 patients [81], and several clinical trials are planned using agents that either target GM-CSF or its receptor [51]. GM-CSF, together with other proinflammatory cytokines such as IL-6, TNF, IFNg, IL-7 and IL-18, is associated with the cytokine storm present in a hyperinflammatory disorder named hemophagocytic lymphohistiocytosis (HLH) which presents with organ failure [12]. Moreover, cytokines related to cytokine release syndrome (such as IL-1A/B, IL-6, IL-10, IL-18, and TNFA), showed increased positive association to the severity of the disease in the blood from COVID-19 patients [47]. Another proinflammatory cytokine specifically upregulated in SARS-CoV-2 infected cells was *IL-32*, which together with *CSF2*, promotes the release of TNF and IL-6 in a continuous positive loop and therefore contribute to this cytokine storm [86]. Interestingly, *IL-6, IL-7* and *IL-18* were found to be upregulated in two of the four data sets of SARS-CoV-2 infected cells. Moreover, not only upregulation, but also a shift in isoform usage of *IL-6* was detected in NHBE and A549 infected cells. A shift in 5' UTR usage in the presence of SARS-CoV-2 may be attributed to indirect host cell signaling cascades that trigger changes in transcription and splicing activity, which could also explain the overall increase in *IL-6* expression.

SERPINA3, a gene coding for an essential enzyme in the regulation of leukocyte proteases, is also induced by cytokines [28]. This was the only gene consistently upregulated in all cell line samples infected with SARS-CoV-2 and absent from the other datasets. Even though this was previously proposed as a promising candidate for the inhibition of viral replication, to date no experiments were carried out to validate this hypothesis [13]. Another interesting candidate gene, which has not been implicated experimentally in respiratory viral infections and was upregulated in our analysis, was *VNN2*. Vanins are involved in proinflammatory and oxidative processes, and VNN2 plays a role in neutrophil migration by regulating b2 integrin [54]. In contrast, the downregulated genes included *SNX8*, which has been previously reported in RNA virus-triggered induction of antiviral genes [13, 26]; and *FKBP5*, a known regulator of NF-kB activity [27]. These results suggest that the

SARS-CoV-2 virus tends to indirectly target specific genes involved in genome replication and host antiviral immune response without eliciting a global change in cellular transcript processing or protein production.

Figure 4. Overview of human factors specific to SARS-CoV-2 infection detected by our analyses. This includes human RBPs whose binding sites are enriched and conserved in the SARS-CoV-2 genome but not in the genomes of related viruses; and genes, isoforms and metabolites that are consistently altered in response to SARS-CoV-2 infection of lung epithelial cells but not in infection with the other tested viruses; ECM (extracellular matrix).

One of the first and most important antiviral responses is the production of type I Interferon (IFN). This protein induces the expression of hundreds of Interferon Stimulated Genes (ISGs), which in turn serve to limit virus spread and infection. Moreover, type I IFN can directly activate immune cells such as macrophages, dendritic cells and NK cells as well as induce the release of proinflammatory cytokines by other cell types [33]. Signaling pathway analysis showed that type I IFN response was greatly impacted in SARS-CoV-2 infected cells (A549 and Calu-3 cells at a MOI of 0.2 and 2 respectively). In the same direction, a higher expression of *PRDM1* (Blimp-1) that we observed in the SARS-CoV-2 infected cells, could also contribute to the critical regulation of IFN signaling cascades; interestingly, the TE family LTR13, which was also upregulated upon SARS-CoV-2 infection, is enriched in PRDM1 binding sites [76]. Therefore, it is possible that regulatory factors involved in IFN and immune response in the context of SARS-CoV-2 infection could also be attributed to TE transcriptional activation. In the same direction, we detected the

upregulation of several TE families in SARS-CoV-2 infected cells that have been previously ₃₅₇
implicated in immune regulation. Moreover, 16 upregulated families were specific to SARS-CoV-2 ₃₅₈
infection in Calu-3 and A549 cell lines. The MER41B family, for instance, is known to contribute to ₃₅₉
interferon gamma inducible binding sites (bound by STAT1 and/or IRF1) [14]. Functional ₃₆₀
enrichment analysis of nearby genes were in accordance with these findings, since several immunity ₃₆₁
related terms were enriched along with "progressive pulmonary impairment". In parallel, TEs seem ₃₆₂
to be co-regulated with phospholipid metabolism, which directly affects the Pi3K/AKT signaling ₃₆₃
pathway, central to the immune response and which were detected in our functional enrichment and ₃₆₄
metabolism flux analyses. ₃₆₅

RBPs are another example of host regulatory factors involved either in the response of human ₃₆₆
cells to SARS-CoV-2 or in the manipulation of human machinery by the virus. We aimed at finding ₃₆₇
RBPs which potentially interact with SARS-CoV-2 genomes in a conserved and specific way. Five of ₃₆₈
the proteins predicted to be interacting with the viral genome by our pipeline (EIF4B, hnRNPA1, ₃₆₉
PABPC1, PABPC4, and YBX1) were experimentally shown to bind to SARS-CoV-2 RNA in an ₃₇₀
infected human liver cell line, based on a recent preprint [65]. ₃₇₁

Among the RBPs whose potential binding sites were enriched and conserved within the ₃₇₂
SARS-CoV-2 virus genomes is the EIF4B, suggesting that the SARS-CoV-2 virus protein translation ₃₇₃
could be EIF4B-dependent. We also detected the upregulation of *EIF4B* in A549 and Calu-3 cells, ₃₇₄
which might indicate that this protein is sequestered by the virus and therefore the cells need to ₃₇₅
increase its production. Moreover, this protein was predicted to interact specifically with the ₃₇₆
intergenic region upstream of the gene encoding the SARS-CoV-2 membrane (M) protein, one of ₃₇₇
four structural proteins from this virus. ₃₇₈

Another conserved RBP, which was also upregulated in infected cells, is the Poly(A) Binding ₃₇₉
Protein Cytoplasmic 1 (PABPC1), which has well described cellular roles in mRNA stability and ₃₈₀
translation. PABPC1 has been previously implicated in multiple viral infections. The activity of ₃₈₁
PABPC1 is modulated to inhibit host protein transcript translation, promoting viral RNA access to ₃₈₂
the host cell translational machinery [70]. Importantly, the 3' UTR region of SARS-CoV2 is also ₃₈₃
enriched in binding sites of the PABPC1 and the PPIE RBPs, the latter of which is known to be ₃₈₄
involved in multiple processes, including mRNA splicing [9,32]. Interestingly, PABPC1 and PABPC4 ₃₈₅
interact with the SARS-CoV-2 N protein, which stabilizes the viral genome [24]. This raises the ₃₈₆
possibility that the viral genome, N protein, and human PABP proteins may participate in a joint ₃₈₇
protein-RNA complex that assists in viral genome stability, replication, and/or ₃₈₈
translation [1,57,60,69,70]. ₃₈₉

An interesting result was that the binding motifs for hnRNPA1, which has been shown to ₃₉₀
interact with other coronavirus genomes, were enriched specifically in the 3'UTR of SARS-CoV-2 ₃₉₁
even though they were depleted in the genome overall. The hnRNPA1 protein was described to ₃₉₂
interact more in particular with multiple sequence elements including the 3'UTR of the Murine ₃₉₃
Hepatitis Virus (MHV), and to participate in both transcription and replication of this ₃₉₄
virus [30,42,66]. This particular gene, along with *hnRNPA2B1*, were downregulated in Calu-3 cells ₃₉₅
and in contrast to the previous examples of upregulated genes, could denote a specific response of ₃₉₆
the human cells to control viral replication. ₃₉₇

Cross referencing the results from our statistical analysis of $\sim 5\%$ of the available genomes (₃₉₈
$\sim 1,500$ out of $> 27,000$ in GISAID) with clinical metadata revealed interesting new insights. ₃₉₉
Indeed, the $D \rightarrow G$ mutation at amino acid position 614 in the Spike protein found in our analysis ₄₀₀
has recently been proposed to have increased viral infectivity [37]. In addition, this same mutation ₄₀₁
has also been associated with an increase in the case fatality rate [6], however, these hypotheses need ₄₀₂
further verification. The P323L mutation in the RNA-dependent RNA polymerase (RdRP) has been ₄₀₃
identified previously, although in that study it was associated with changes in geographical location ₄₀₄
of the viral strain [56]. Finally, the L37F mutation in the Nsp6 protein has been reported to be ₄₀₅
located outside of the transmembrane domain [11], being present at a high frequency [83], and ₄₀₆
proposed to negatively affect protein structure stability [8]. Our statistics may contain bias based on ₄₀₇
the number of genome sequences being collected earlier *versus* later in the pandemic, genomes ₄₀₈

lacking clinical outcome metadata, and in the case of the Spike D614G a potential increase of fitness ₄₀₉
associated with this mutation. However, the fact that more than one of our predictions has also been ₄₁₀
detected by different studies justifies future wet lab experiments to compare the effect of the other ₄₁₁
identified mutations. ₄₁₂

Conclusion ₄₁₃

Overall, our analyses identified sets of statistically significant host genes, isoforms, regulatory ₄₁₄
elements, and other interactions that likely contribute to the cellular response during infection with ₄₁₅
SARS-CoV-2. Furthermore, we detected potential binding sites for human RBPs that are conserved ₄₁₆
across SARS-CoV-2 genomes, along with a subset of variants in the viral genome that correlate well ₄₁₇
with disease severity in SARS-CoV-2 infection. To our knowledge, this is the first work where a ₄₁₈
computational meta-analysis was performed to predict host factors that play a role in the specific ₄₁₉
pathogenesis of SARS-CoV-2, distinct from other respiratory viruses. ₄₂₀

We envision that applying this publicly available workflow will yield important mechanistic ₄₂₁
insights in future analyses on emerging pathogens. Similarly, we expect that the results for ₄₂₂
SARS-CoV-2 will contribute to ongoing efforts in the selection of new drug targets and the ₄₂₃
development of more effective prophylactics and therapeutics to reduce virus infection and ₄₂₄
replication with minimal adverse effects on the human host. ₄₂₅

Materials and Methods ₄₂₆

Datasets ₄₂₇

Two datasets were downloaded from the Gene Expression Omnibus (GEO) database, hosted at the ₄₂₈
National Center for Biotechnology Information (NCBI). The first dataset, GSE147507 [10], includes ₄₂₉
gene expression measurements from three cell lines derived from the human respiratory system ₄₃₀
(NHBE, A549, Calu-3) infected either with SARS-CoV-2, influenza A virus (IAV), respiratory ₄₃₁
syncytial virus (RSV), or human parainfluenza virus 3 (HPIV3). The second dataset, GSE150316, ₄₃₂
includes RNA-seq extracted from formalin fixed, paraffin embedded (FFPE) histological sections of ₄₃₃
lung biopsies from COVID-19 deceased patients and healthy individuals. This dataset encompasses a ₄₃₄
variable number of biopsies per subject, ranging from one to five. Given its limitations, we only ₄₃₅
utilized the second dataset for differential expression analysis. ₄₃₆

The reference genome sequences of SARS-CoV-2 (NC_045512), RaTG13 (MN996532.1), and ₄₃₇
SARS-CoV (NC_004718.3) were downloaded from NCBI. Additionally, a list of known RBPs and ₄₃₈
their PWMs were downloaded from ATtRACT (https://attract.cnic.es/download). Finally, all ₄₃₉
SARS-CoV-2 complete genomes collected from humans and that had disease severity information ₄₄₀
were downloaded from GISAID on 19 May, 2020 [67]. ₄₄₁

RNAseq data processing and differential expression analysis ₄₄₂

Data was downloaded from SRA using sra-tools (v2.10.8; https://github.com/ncbi/sra-tools) and ₄₄₃
transformed to fastq with fastq-dump. FastQC (v0.11.9; https://github.com/s-andrews/FastQC) ₄₄₄
and MultiQC (v1.9) [20] were employed to assess the quality of the data used and the need to trim ₄₄₅
reads and/or remove adapters. Selected datasets were mapped to the human reference genome ₄₄₆
(GENCODE Release 19, GRCh37.p13) utilizing STAR (v2.7.3a) [17]. Alignment statistics were used ₄₄₇
to determine which datasets should be included in subsequent steps. Resulting SAM files were ₄₄₈
converted to BAM files employing samtools (v1.9) [41]. Next, read quantification was performed ₄₄₉
using StringTie (v2.1.1) [58] and the output data was postprocessed with an auxiliary Python script ₄₅₀
provided by the same developers to produce files ready for subsequent downstream analyses. For the ₄₅₁
second gene expression dataset, raw counts were downloaded from GEO. DESeq2 (v1.26.0) [45] was ₄₅₂
used in both cases to identify differentially expressed genes (DEGs). Finally, an exploratory data ₄₅₃

analysis was carried out based on the transformed values obtained after applying the variance   454
stabilizing transformation [3] implemented in the vst() function of DESeq2 [46]. Hence, principal   455
component analysis (PCA) was performed to evaluate the main sources of variation in the data and   456
remove outliers.   457

Gene ontology enrichment analysis   458

The DEGs produced by DESeq2 with an absolute Log2FC > 1 and FDR-adjusted p-value < 0.05   459
were used as input to a general GO enrichment analysis [5, 74]. Each term was verified with a   460
hypergeometric test from the GOstats package (v2.54.0) [21] and the p-values were corrected for   461
multiple-hypothesis testing employing the Bonferroni method [40]. GO terms with a significant   462
adjusted p-value of less than 0.05 were reduced to representative non-redundant terms with the use   463
of REVIGO [71].   464

Host signaling pathway enrichment   465

The DEG lists produced by DESeq2 with an absolute Log2FC > 1 and FDR-adjusted p-value < 0.05   466
were used as input to the SPIA algorithm to identify significantly affected pathways from the R   467
graphite library [63, 73]. Pathways with Bonferroni-adjusted p-values less than 0.05 were included in   468
downstream analyses. The significant results for all comparisons from publicly available data from   469
KEGG, Reactome, Panther, BioCarta, and NCI were then compiled to facilitate downstream   470
comparison. Hypergeometric pathway enrichments were performed using the Database for   471
Annotation, Visualization and Integrated Discovery (DAVID, v6.8) [29].   472

Integration of transcriptomic analysis with the human metabolic network   473

To detect increased and decreased fluxes of metabolites we projected the transcriptomic data onto   474
the human reconstructed metabolic network Recon (v2.04) [75]. First, we ran EBSeq [38] on the   475
gene count matrix generated in the previous steps. Then, we used the output of EBSeq containing   476
posterior probabilities of a gene being DE (PPDE) and the Log2FC as input to the Moomin   477
method [61] using default parameters. Finally, we enumerated 250 topological solutions in order to   478
construct a consensus solution for each of the datasets tested.   479

Isoform Analysis   480

Using transcript quantification data from StringTie as input, we identified isoform switching events   481
and their predicted functional consequences with the IsoformSwitchAnalyzeR R package   482
(v1.11.3) [78]. In summary, we filtered for isoforms that experienced $|\geq 30\%|$ switch in usage for   483
each gene and were corrected for false discovery rate (FDR) with a q-value < 0.05. Following   484
filtering for significant isoforms, we externally predicted their coding capabilities, protein structure   485
stability, peptide signaling, and shifts in protein domain usage using The Coding-Potential   486
Assessment Tool (CPAT) [79], IUPred2 [18], SignalP [2] and Pfam tools respectively [19]. These   487
external results were imported back into IsoformSwitchAnalyzeR and used to further identify   488
alternative splicing events and functional consequences as well as visualize the overall and individual   489
effects of isoform switching data. Specifically, to calculate differential analysis between samples,   490
isoform expression and usage were measured by the isoform fraction (IF) value, which quantifies the   491
individual isoform expression level relative to the parent gene's expression level: 492

$$IF = \frac{isoform\ expression}{gene\ expression}$$

By proxy, the difference in isoform usage between samples (dIF) measures the effect size between   493
conditions and is calculated as follows: 494

$$dIF = IF2\text{--}IF1$$

dIF was measured on a scale of 0 to 1, with 0 = no (0%) change in usage between conditions and　495
1 = complete (100%) change in usage. The sum of dIF values for all isoforms associated with one　496
gene is equal to 1. Gene expression data was imported from the aforementioned DESeq2 results.　497
The top 30 isoforms per dataset comparison were identified by ranking isoforms by gene switch　498
q-value, i.e. the significance of the summation of all isoform switching events per gene between mock　499
and infected conditions.　500

Transposable Element Analysis　501

TE expression was quantified using the TEcount function from the TEtools software [39]. TEcount　502
detects reads aligned against copies of each TE family annotated from the reference genome. DETEs　503
in infected vs mock conditions were detected using DEseq2 (v1.26.0) [45] with a matrix of counts for　504
genes and TE families as input. Functional enrichment of nearby genes (upstream 5kb and　505
downstream 1kb of each TE copy within the human genome) was calculated with GREAT [49] using　506
options "genome background" and "basal + extension". We only selected occurrences statistically　507
significant by region binomial test.　508

Identification of putative binding sites for human RBPs on the　509
SARS-CoV-2 genome　510

The list of RBPs downloaded from ATtRACT was filtered to human RBPs. The list was further　511
filtered to retain PWMs obtained through competitive experiments and drop PWMs with very high　512
entropy. This left 205 PWMs for 102 human RBPs. The SARS-CoV-2 reference genome sequence　513
was scanned with the remaining PWMs using the TFBSTools R package (v1.20.0) [72]. A minimum　514
score threshold of 90% was used to identify putative RBP binding sites.　515

Enrichment analysis for putative RBP binding sites　516

The sequences of the SARS-CoV-2 genome, 5'UTR, 3'UTR, intergenic regions and negative sense　517
molecule were each scrambled 1,000 times. Each of the 1,000 scrambled sequences was scanned for　518
RBP binding sites as described above. The number of binding sites for each RBP was counted, and　519
the mean and standard deviation of the number of sites was calculated for each RBP, per region,　520
across all 1,000 simulations. A minimum FDR-adjusted p-value of 0.01 was taken as the cutoff for　521
enrichment. This analysis was repeated with the reference genomes of SARS-CoV and RaTG13 .　522

Conservation analysis for putative RBP binding sites　523

The multiple sequence alignment of 27,592 SARS-CoV-2 genome sequences was downloaded from　524
GISAID [67] on May 19th 2020. For each putative RBP binding site, we selected the corresponding　525
columns of the multiple sequence alignment. We then counted the number of genomes in which the　526
sequence was identical to that of the reference genome.　527

Viral genotype-phenotype correlation　528

All complete SARS-CoV-2 genomes from GISAID, together with the GenBank reference sequence,　529
were aligned with MAFFT (v7.464) within a high-performance computing environment using 1　530
thread and the –nomemsave parameter [53]. Sequences responsible for introducing excessive gaps in　531
this initial alignment were then identified and removed, leaving 1,511 sequences that were then used　532
to generate a new multiple sequence alignment. The disease severity metadata for these sequences　533

was then normalized into four categories: severe, moderate, mild, and unknown. Next, the sequence ₅₃₄
data and associated metadata were used as input to the meta-CATS algorithm to identify aligned ₅₃₅
positions that contained significant differences in their base distribution between 2 or more disease ₅₃₆
severities [59]. The Benjamini-Hochberg multiple hypothesis correction was then applied to all ₅₃₇
positions [7]. The top 50 most significant positions were then evaluated against the annotated ₅₃₈
protein regions of the reference genome to determine their effect on amino acid sequence. ₅₃₉

Code availability ₅₄₀

Code for these analyses is available at https://github.com/vaguiarpulido/covid19-research. ₅₄₁

Supporting Information ₅₄₂

Supplementary Figure 1. Isoform Analysis. ₅₄₃
Supplementary File 1. Zipped file containing Complete DEG tables. ₅₄₄
Supplementary File 2. Zipped file containing GO for each dataset. ₅₄₅
Supplementary File 3. TE family count/differential expression. ₅₄₆
Supplementary File 4. GREAT Analysis (complete and per family). ₅₄₇
Supplementary Table 1. Merged tables (Specific genes in SARS-CoV-2) ₅₄₈
Supplementary Table 2. Supporting information for Figure 2, consisting of functional ₅₄₉
enrichment specific to SARS-CoV-2. ₅₅₀
Supplementary Table 3. Pathway enrichment for each dataset (SPIA and DAVID merged into ₅₅₁
one file). ₅₅₂
Supplementary Table 4. Metabolic fluxes predicted for each dataset using Moomin. ₅₅₃
Supplementary Table 5. Isoform analysis. ₅₅₄
Supplementary Table 6. Putative binding sites for human RBPs on the SARS-CoV-2 genome. ₅₅₅
Supplementary Table 7. Enrichment of binding motifs for human RBPs on the SARS-CoV-2 ₅₅₆
genome. ₅₅₇
Supplementary Table 8. Conservation of binding motifs for human RBPs across genome ₅₅₈
sequences of SARS-CoV-2 isolates. ₅₅₉
Supplementary Table 9. Biological evidence associated with putative SARS-CoV-2 interacting ₅₆₀
human RBPs. ₅₆₁
Supplementary Table 10. Enrichment of binding motifs for human RBPs on the SARS-CoV ₅₆₂
genome ₅₆₃
Supplementary Table 11. Enrichment of binding motifs for human RBPs on the RaTG13 genome ₅₆₄

Funding ₅₆₅

The authors received no specific funding to support this work. ₅₆₆

Acknowledgments ₅₆₇

We would like to thank the Virtual BioHackathon on COVID-19 that took place during April 2020 ₅₆₈
(https://github.com/virtual-biohackathons/covid-19-bh20) for fostering an environment that ₅₆₉
triggered this collaboration and in particular the Gene Expression group for the fruitful discussions. ₅₇₀
This work was performed using the computing facilities of the CC/PRABI/LBBE in France and the ₅₇₁
France Génomique e-infrastructure (ANR-10-INBS-09-08), the HPC facilities of the University of ₅₇₂
Luxembourg [77]. We would also like to thank Slack for providing us with free access to the ₅₇₃
professional version of the platform. ₅₇₄
We would also like to thank Slack for providing us with free access to the professional version of the ₅₇₅
platform. ₅₇₆

Conflicts of Interest

A.L. is an employee of NVIDIA Corporation.

References

1. P. Ahlquist, A. O. Noueiry, W.-M. Lee, D. B. Kushner, and B. T. Dye. Host factors in positive-strand RNA virus genome replication. *J. Virol.*, 77(15):8181–8186, Aug. 2003.

2. J. J. Almagro Armenteros, K. D. Tsirigos, C. K. Sønderby, T. N. Petersen, O. Winther, S. Brunak, G. von Heijne, and H. Nielsen. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat. Biotechnol.*, 37(4):420–423, Apr. 2019.

3. S. Anders and W. Huber. Differential expression analysis for sequence count data. *Genome Biol.*, 11(10):R106, Oct. 2010.

4. K. G. Andersen, A. Rambaut, W. I. Lipkin, E. C. Holmes, and R. F. Garry. The proximal origin of SARS-CoV-2. *Nat. Med.*, 26(4):450–452, Apr. 2020.

5. M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, and G. Sherlock. Gene ontology: tool for the unification of biology. the gene ontology consortium. *Nat. Genet.*, 25(1):25–29, May 2000.

6. M. Becerra-Flores and T. Cardozo. SARS-CoV-2 viral spike G614 mutation exhibits higher case fatality rate. *Int. J. Clin. Pract.*, page e13525, May 2020.

7. Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: A practical and powerful approach to multiple testing, 1995.

8. D. Benvenuto, S. Angeletti, M. Giovanetti, M. Bianchi, S. Pascarella, R. Cauda, M. Ciccozzi, and A. Cassone. Evolutionary analysis of SARS-CoV-2: how mutation of Non-Structural protein 6 (NSP6) could affect viral autophagy. *J. Infect.*, 81(1):e24–e27, July 2020.

9. K. Bertram, D. E. Agafonov, W.-T. Liu, O. Dybkov, C. L. Will, K. Hartmuth, H. Urlaub, B. Kastner, H. Stark, and R. Lührmann. Cryo-EM structure of a human spliceosome activated for step 2 of splicing. *Nature*, 542(7641):318–323, Feb. 2017.

10. D. Blanco-Melo, B. E. Nilsson-Payant, W.-C. Liu, S. Uhl, D. Hoagland, R. Møller, T. X. Jordan, K. Oishi, M. Panis, D. Sachs, T. T. Wang, R. E. Schwartz, J. K. Lim, R. A. Albrecht, and B. R. tenOever. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell*, 181(5):1036–1045.e9, May 2020.

11. Y. Cárdenas-Conejo, A. Liñan-Rico, D. A. García-Rodríguez, S. Centeno-Leija, and H. Serrano-Posada. An exclusive 42 amino acid signature in pp1ab protein provides insights into the evolutive history of the 2019 novel human-pathogenic coronavirus (SARS-CoV-2). *J. Med. Virol.*, 92(6):688–692, June 2020.

12. S. J. Carter, R. S. Tattersall, and A. V. Ramanan. Macrophage activation syndrome in adults: recent advances in pathophysiology, diagnosis and treatment. *Rheumatology*, 58(1):5–17, Jan. 2019.

13. D. Chasman, K. B. Walters, T. J. S. Lopes, A. J. Eisfeld, Y. Kawaoka, and S. Roy. Integrating transcriptomic and proteomic data using predictive regulatory network models of host response to pathogens. *PLoS Comput. Biol.*, 12(7):e1005013, July 2016.

14. E. B. Chuong, N. C. Elde, and C. Feschotte. Regulatory evolution of innate immunity through co-option of endogenous retroviruses. *Science*, 351(6277):1083–1087, Mar. 2016.

15. E. B. Chuong, N. C. Elde, and C. Feschotte. Regulatory activities of transposable elements: from conflicts to benefits. *Nat. Rev. Genet.*, 18(2):71–86, Feb. 2017.

16. Ö. Deniz, M. Ahmed, C. D. Todd, A. Rio-Machin, M. A. Dawson, and M. R. Branco. Endogenous retroviruses are a source of enhancers with oncogenic potential in acute myeloid leukaemia. *Nat. Commun.*, 11(1):3506, July 2020.

17. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T. R. Gingeras. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1):15–21, Jan. 2013.

18. Z. Dosztányi, V. Csizmok, P. Tompa, and I. Simon. IUPred: web server for the prediction of intrinsically unstructured regions of proteins based on estimated energy content. *Bioinformatics*, 21(16):3433–3434, Aug. 2005.

19. S. El-Gebali, J. Mistry, A. Bateman, S. R. Eddy, A. Luciani, S. C. Potter, M. Qureshi, L. J. Richardson, G. A. Salazar, A. Smart, E. L. L. Sonnhammer, L. Hirsh, L. Paladin, D. Piovesan, S. C. E. Tosatto, and R. D. Finn. The pfam protein families database in 2019. *Nucleic Acids Res.*, 47(D1):D427–D432, Jan. 2019.

20. P. Ewels, M. Magnusson, S. Lundin, and M. Käller. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 32(19):3047–3048, Oct. 2016.

21. S. Falcon and R. Gentleman. Using GOstats to test gene lists for GO term association. *Bioinformatics*, 23(2):257–258, Jan. 2007.

22. T. S. Fung and D. X. Liu. Human coronavirus: Host-Pathogen interaction. *Annu. Rev. Microbiol.*, 73:529–557, Sept. 2019.

23. G. Giudice, F. Sánchez-Cabo, C. Torroja, and E. Lara-Pezzi. ATtRACT-a database of RNA-binding proteins and associated motifs. *Database*, 2016, Apr. 2016.

24. D. E. Gordon, G. M. Jang, M. Bouhaddou, J. Xu, K. Obernier, M. J. O'Meara, J. Z. Guo, D. L. Swaney, T. A. Tummino, R. Hüttenhain, R. M. Kaake, A. L. Richards, B. Tutuncuoglu, H. Foussard, J. Batra, K. Haas, M. Modak, M. Kim, P. Haas, B. J. Polacco, H. Braberg, J. M. Fabius, M. Eckhardt, M. Soucheray, M. J. Bennett, M. Cakir, M. J. McGregor, Q. Li, Z. Z. C. Naing, Y. Zhou, S. Peng, I. T. Kirby, J. E. Melnyk, J. S. Chorba, K. Lou, S. A. Dai, W. Shen, Y. Shi, Z. Zhang, I. Barrio-Hernandez, D. Memon, C. Hernandez-Armenta, C. J. P. Mathy, T. Perica, K. B. Pilla, S. J. Ganesan, D. J. Saltzberg, R. Ramachandran, X. Liu, S. B. Rosenthal, L. Calviello, S. Venkataramanan, Y. Lin, S. A. Wankowicz, M. Bohn, R. Trenker, J. M. Young, D. Cavero, J. Hiatt, T. Roth, U. Rathore, A. Subramanian, J. Noack, M. Hubert, F. Roesch, T. Vallet, B. Meyer, K. M. White, L. Miorin, D. Agard, M. Emerman, D. Ruggero, A. García-Sastre, N. Jura, M. von Zastrow, J. Taunton, O. Schwartz, M. Vignuzzi, C. d'Enfert, S. Mukherjee, M. Jacobson, H. S. Malik, D. G. Fujimori, T. Ideker, C. S. Craik, S. Floor, J. S. Fraser, J. Gross, A. Sali, T. Kortemme, P. Beltrao, K. Shokat, B. K. Shoichet, and N. J. Krogan. A SARS-CoV-2-Human Protein-Protein interaction map reveals drug targets and potential Drug-Repurposing. *bioRxiv*, Mar. 2020.

25. GTEx Consortium. Human genomics. the Genotype-Tissue expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science*, 348(6235):648–660, May 2015.

26. W. Guo, J. Wei, X. Zhong, R. Zang, H. Lian, M.-M. Hu, S. Li, H.-B. Shu, and Q. Yang. SNX8 modulates the innate immune response to RNA viruses by regulating the aggregation of VISA. *Cell. Mol. Immunol.*, Sept. 2019.

27. M. Hinz, M. Broemer, S. Ç. Arslan, A. Otto, E.-C. Mueller, R. Dettmer, and C. Scheidereit. Signal responsiveness of IκB kinases is determined by cdc37-assisted transient interaction with hsp90. *J. Biol. Chem.*, 282(44):32311–32319, Nov. 2007.

28. S. Horváth and K. Mirnics. Immune system disturbances in schizophrenia. *Biol. Psychiatry*, 75(4):316–323, Feb. 2014.

29. D. W. Huang, B. T. Sherman, and R. A. Lempicki. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.*, 4(1):44–57, 2009.

30. P. Huang and M. M. Lai. Heterogeneous nuclear ribonucleoprotein a1 binds to the 3'-untranslated region and mediates potential 5'-3'-end cross talks of mouse hepatitis virus RNA. *J. Virol.*, 75(11):5009–5017, June 2001.

31. J. Ito, R. Sugimoto, H. Nakaoka, S. Yamada, T. Kimura, T. Hayano, and I. Inoue. Systematic identification and characterization of regulatory elements derived from human endogenous retroviruses. *PLoS Genet.*, 13(7):e1006883, July 2017.

32. M. S. Jurica, L. J. Licklider, S. R. Gygi, N. Grigorieff, and M. J. Moore. Purification and characterization of native spliceosomes suitable for three-dimensional structural analysis. *RNA*, 8(4):426–439, Apr. 2002.

33. N. Kadowaki and Y.-J. Liu. Natural type I interferon-producing cells as a link between innate and adaptive immunity. *Hum. Immunol.*, 63(12):1126–1132, Dec. 2002.

34. R. J. Khan, R. K. Jha, G. M. Amera, M. Jain, E. Singh, A. Pathak, R. P. Singh, J. Muthukumaran, and A. K. Singh. Targeting SARS-CoV-2: a systematic drug repurposing approach to identify promising inhibitors against 3c-like proteinase and 2'-o-ribose methyltransferase. *J. Biomol. Struct. Dyn.*, pages 1–14, Apr. 2020.

35. D. Kim, J.-Y. Lee, J.-S. Yang, J. W. Kim, V. N. Kim, and H. Chang. The architecture of SARS-CoV-2 transcriptome. *Cell*, 181(4):914–921.e10, May 2020.

36. K. K. Kojima. Human transposable elements in repbase: genomic footprints from fish to humans. *Mob. DNA*, 9:2, Jan. 2018.

37. B. Korber, W. M. Fischer, S. Gnanakaran, H. Yoon, J. Theiler, W. Abfalterer, N. Hengartner, E. E. Giorgi, T. Bhattacharya, B. Foley, K. M. Hastie, M. D. Parker, D. G. Partridge, C. M. Evans, T. M. Freeman, T. I. de Silva, C. McDanal, L. G. Perez, H. Tang, A. Moon-Walker, S. P. Whelan, C. C. LaBranche, E. O. Saphire, D. C. Montefiori, A. Angyal, R. L. Brown, L. Carrilero, L. R. Green, D. C. Groves, K. J. Johnson, A. J. Keeley, B. B. Lindsey, P. J. Parsons, M. Raza, S. Rowland-Jones, N. Smith, R. M. Tucker, D. Wang, and M. D. Wyles. Tracking changes in SARS-CoV-2 spike: Evidence that D614G increases infectivity of the COVID-19 virus. *Cell*, July 2020.

38. N. Leng, J. A. Dawson, J. A. Thomson, V. Ruotti, A. I. Rissman, B. M. G. Smits, J. D. Haag, M. N. Gould, R. M. Stewart, and C. Kendziorski. EBSeq: an empirical bayes hierarchical model for inference in RNA-seq experiments. *Bioinformatics*, 29(8):1035–1043, Apr. 2013.

39. E. Lerat, M. Fablet, L. Modolo, H. Lopez-Maestre, and C. Vieira. TEtools facilitates big data expression analysis of transposable elements and reveals an antagonism between their activity and that of piRNA genes. *Nucleic Acids Res.*, 45(4):e17, Feb. 2017.

40. K. Lesack and C. Naugler. An open-source software program for performing bonferroni and related corrections for multiple comparisons. *J. Pathol. Inform.*, 2:52, Dec. 2011.

41. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, and R. Durbin. The sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16):2078–2079, Aug. 2009.

42. H. P. Li, X. Zhang, R. Duncan, L. Comai, and M. M. Lai. Heterogeneous nuclear ribonucleoprotein A1 binds to the transcription-regulatory region of mouse hepatitis virus RNA. *Proc. Natl. Acad. Sci. U. S. A.*, 94(18):9544–9549, Sept. 1997.

43. Z. Li and P. D. Nagy. Diverse roles of host RNA binding proteins in RNA virus replication. *RNA Biol.*, 8(2):305–315, Mar. 2011.

44. M. Liao, Y. Liu, J. Yuan, Y. Wen, G. Xu, J. Zhao, L. Cheng, J. Li, X. Wang, F. Wang, L. Liu, I. Amit, S. Zhang, and Z. Zhang. Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19, 2020.

45. M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.*, 15(12):550, 2014.

46. M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.*, 15(12):550, 2014.

47. C. Lucas, P. Wong, J. Klein, T. B. R. Castro, J. Silva, M. Sundaram, M. K. Ellingson, T. Mao, J. E. Oh, B. Israelow, T. Takahashi, M. Tokuyama, P. Lu, A. Venkataraman, A. Park, S. Mohanty, H. Wang, A. L. Wyllie, C. B. F. Vogels, R. Earnest, S. Lapidus, I. M. Ott, A. J. Moore, M. C. Muenker, J. B. Fournier, M. Campbell, C. D. Odio, A. Casanovas-Massana, A. Obaid, A. Lu-Culligan, A. Nelson, A. Brito, A. Nunez, A. Martin, A. Watkins, B. Geng, C. Kalinich, C. Harden, C. Todeasa, C. Jensen, D. Kim, D. McDonald, D. Shepard, E. Courchaine, E. B. White, E. Song, E. Silva, E. Kudo, G. DeIuliis, H. Rahming, H.-J. Park, I. Matos, J. Nouws, J. Valdez, J. Fauver, J. Lim, K.-A. Rose, K. Anastasio, K. Brower, L. Glick, L. Sharma, L. Sewanan, L. Knaggs, M. Minasyan, M. Batsu, M. Petrone, M. Kuang, M. Nakahata, M. Linehan, M. H. Askenase, M. Simonov, M. Smolgovsky, N. Sonnert, N. Naushad, P. Vijayakumar, R. Martinello, R. Datta, R. Handoko, S. Bermejo, S. Prophet, S. Bickerton, S. Velazquez, T. Alpert, T. Rice, W. Khoury-Hanold, X. Peng, Y. Yang, Y. Cao, Y. Strong, R. Herbst, A. C. Shaw, R. Medzhitov, W. L. Schulz, N. D. Grubaugh, C. Dela Cruz, S. Farhadian, A. I. Ko, S. B. Omer, A. Iwasaki, and Y. I. Team. Longitudinal analyses reveal immunological misfiring in severe covid-19. *Nature*, 2020.

48. M. G. Macchietto, R. A. Langlois, and S. S. Shen. Virus-induced transposable element expression up-regulation in human and mouse host cells. *Life Science Alliance*, 3(2), Feb. 2020.

49. C. Y. McLean, D. Bristor, M. Hiller, S. L. Clarke, B. T. Schaar, C. B. Lowe, A. M. Wenger, and G. Bejerano. GREAT improves functional interpretation of cis-regulatory regions. *Nat. Biotechnol.*, 28(5):495–501, May 2010.

50. H. M. Mehta, M. Malandra, and S. J. Corey. G-CSF and GM-CSF in neutropenia. *J. Immunol.*, 195(4):1341–1349, Aug. 2015.

51. P. Mehta, D. F. McAuley, M. Brown, E. Sanchez, R. S. Tattersall, J. J. Manson, and HLH Across Speciality Collaboration, UK. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*, 395(10229):1033–1034, Mar. 2020.

52. N. Muralidharan, R. Sakthivel, D. Velmurugan, and M. M. Gromiha. Computational studies of drug repurposing and synergism of lopinavir, oseltamivir and ritonavir binding with SARS-CoV-2 protease against COVID-19. *J. Biomol. Struct. Dyn.*, pages 1–6, Apr. 2020.

53. T. Nakamura, K. D. Yamada, K. Tomii, and K. Katoh. Parallelization of MAFFT for large-scale multiple sequence alignments. *Bioinformatics*, 34(14):2490–2492, July 2018.

54. T. Nitto and K. Onodera. Linkage between coenzyme a metabolism and inflammation: roles of pantetheinase. *J. Pharmacol. Sci.*, 123(1):1–8, Sept. 2013.

55. J. K. Pace, 2nd and C. Feschotte. The evolutionary history of human DNA transposons: evidence for intense activity in the primate lineage. *Genome Res.*, 17(4):422–432, Apr. 2007.

56. M. Pachetti, B. Marini, F. Benedetti, F. Giudici, E. Mauro, P. Storici, C. Masciovecchio, S. Angeletti, M. Ciccozzi, R. C. Gallo, D. Zella, and R. Ippodrino. Emerging SARS-CoV-2 mutation hot spots include a novel RNA-dependent-RNA polymerase variant. *J. Transl. Med.*, 18(1):179, Apr. 2020.

57. C. Perez, C. McKinney, U. Chulunbaatar, and I. Mohr. Translational control of the abundance of cytoplasmic poly(a) binding protein in human cytomegalovirus-infected cells. *J. Virol.*, 85(1):156–164, Jan. 2011.

58. M. Pertea, G. M. Pertea, C. M. Antonescu, T.-C. Chang, J. T. Mendell, and S. L. Salzberg. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat. Biotechnol.*, 33(3):290–295, Mar. 2015.

59. B. E. Pickett, M. Liu, E. L. Sadat, R. B. Squires, J. M. Noronha, S. He, W. Jen, S. Zaremba, Z. Gu, L. Zhou, C. N. Larsen, I. Bosch, L. Gehrke, M. McGee, E. B. Klem, and R. H. Scheuermann. Metadata-driven comparative analysis tool for sequences (meta-CATS): an automated process for identifying significant sequence variations that correlate with virus attributes. *Virology*, 447(1-2):45–51, Dec. 2013.

60. C. Polacek, P. Friebe, and E. Harris. Poly(A)-binding protein binds to the non-polyadenylated 3' untranslated region of dengue virus and modulates translation efficiency. *J. Gen. Virol.*, 90(Pt 3):687–692, Mar. 2009.

61. T. Pusa, M. G. Ferrarini, R. Andrade, A. Mary, A. Marchetti-Spaccamela, L. Stougie, and M.-F. Sagot. MOOMIN - mathematical exploration of 'omics data on a MetabolIc network. *Bioinformatics*, 36(2):514–523, Jan. 2020.

62. P. A. Reyfman, J. M. Walter, N. Joshi, K. R. Anekalla, A. C. McQuattie-Pimentel, S. Chiu, R. Fernandez, M. Akbarpour, C.-I. Chen, Z. Ren, R. Verma, H. Abdala-Valencia, K. Nam, M. Chi, S. Han, F. J. Gonzalez-Gonzalez, S. Soberanes, S. Watanabe, K. J. N. Williams, A. S. Flozak, T. T. Nicholson, V. K. Morgan, D. R. Winter, M. Hinchcliff, C. L. Hrusch, R. D. Guzy, C. A. Bonham, A. I. Sperling, R. Bag, R. B. Hamanaka, G. M. Mutlu, A. V. Yeldandi, S. A. Marshall, A. Shilatifard, L. A. N. Amaral, H. Perlman, J. I. Sznajder, A. C. Argento, C. T. Gillespie, J. Dematte, M. Jain, B. D. Singer, K. M. Ridge, A. P. Lam, A. Bharat, S. M. Bhorade, C. J. Gottardi, G. R. S. Budinger, and A. V. Misharin. Single-Cell transcriptomic analysis of human lung provides insights into the pathobiology of pulmonary fibrosis. *Am. J. Respir. Crit. Care Med.*, 199(12):1517–1536, June 2019.

63. G. Sales, E. Calura, D. Cavalieri, and C. Romualdi. graphite - a bioconductor package to convert pathway topology to gene network. *BMC Bioinformatics*, 13:20, Jan. 2012.

64. C. D. Schmid and P. Bucher. MER41 repeat sequences contain inducible STAT1 binding sites. *PLoS One*, 5(7):e11425, July 2010.

65. N. Schmidt, C. A. Lareau, H. Keshishian, R. Melanson, M. Zimmer, L. Kirschner, J. Ade, S. Werner, N. Caliskan, E. S. Lander, J. Vogel, S. A. Carr, J. Bodem, and M. Munschauer. A direct RNA-protein interaction atlas of the SARS-CoV-2 RNA in infected human cells.

66. S. T. Shi, P. Huang, H. P. Li, and M. M. Lai. Heterogeneous nuclear ribonucleoprotein A1 regulates RNA synthesis of a cytoplasmic virus. *EMBO J.*, 19(17):4701–4711, Sept. 2000.

67. Y. Shu and J. McCauley. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill.*, 22(13), Mar. 2017.

68. A. F. Smit. Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr. Opin. Genet. Dev.*, 9(6):657–663, Dec. 1999.

69. R. W. P. Smith, R. C. Anderson, O. Larralde, J. W. S. Smith, B. Gorgoni, W. A. Richardson, P. Malik, S. V. Graham, and N. K. Gray. Viral and cellular mRNA-specific activators harness PABP and eIF4G to promote translation initiation downstream of cap binding. *Proc. Natl. Acad. Sci. U. S. A.*, 114(24):6310–6315, June 2017.

70. R. W. P. Smith and N. K. Gray. Poly(A)-binding protein (PABP): a common viral target. *Biochem. J*, 426(1):1–12, Feb. 2010.

71. F. Supek, M. Bošnjak, N. Škunca, and T. Šmuc. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One*, 6(7):e21800, July 2011.

72. G. Tan and B. Lenhard. TFBSTools: an R/bioconductor package for transcription factor binding site analysis. *Bioinformatics*, 32(10):1555–1556, 01 2016.

73. A. L. Tarca, S. Draghici, P. Khatri, S. S. Hassan, P. Mittal, J.-S. Kim, C. J. Kim, J. P. Kusanovic, and R. Romero. A novel signaling pathway impact analysis. *Bioinformatics*, 25(1):75–82, Jan. 2009.

74. The Gene Ontology Consortium and The Gene Ontology Consortium. The gene ontology resource: 20 years and still GOing strong, 2019.

75. I. Thiele, N. Swainston, R. M. T. Fleming, A. Hoppe, S. Sahoo, M. K. Aurich, H. Haraldsdottir, M. L. Mo, O. Rolfsson, M. D. Stobbe, S. G. Thorleifsson, R. Agren, C. Bölling, S. Bordel, A. K. Chavali, P. Dobson, W. B. Dunn, L. Endler, D. Hala, M. Hucka, D. Hull, D. Jameson, N. Jamshidi, J. J. Jonsson, N. Juty, S. Keating, I. Nookaew, N. Le Novère, N. Malys, A. Mazein, J. A. Papin, N. D. Price, E. Selkov, Sr, M. I. Sigurdsson, E. Simeonidis, N. Sonnenschein, K. Smallbone, A. Sorokin, J. H. G. M. van Beek, D. Weichart, I. Goryanin, J. Nielsen, H. V. Westerhoff, D. B. Kell, P. Mendes, and B. Ø. Palsson. A community-driven global reconstruction of human metabolism. *Nat. Biotechnol.*, 31(5):419–425, May 2013.

76. M. Trizzino, A. Kapusta, and C. D. Brown. Transposable elements generate regulatory novelty in a tissue-specific fashion. *BMC Genomics*, 19(1):468, June 2018.

77. S. Varrette, P. Bouvry, H. Cartiaux, and F. Georgatos. Management of an academic hpc cluster: The ul experience. In *Proc. of the 2014 Intl. Conf. on High Performance Computing & Simulation (HPCS 2014)*, pages 959–967, Bologna, Italy, July 2014. IEEE.

78. K. Vitting-Seerup and A. Sandelin. IsoformSwitchAnalyzeR: analysis of changes in genome-wide patterns of alternative splicing and its functional consequences. *Bioinformatics*, 35(21):4469–4471, Nov. 2019.

79. L. Wang, H. J. Park, S. Dasari, S. Wang, J.-P. Kocher, and W. Li. CPAT: Coding-Potential assessment tool using an alignment-free logistic regression model. *Nucleic Acids Res.*, 41(6):e74, Apr. 2013.

80. W. Wen, W. Su, H. Tang, W. Le, X. Zhang, Y. Zheng, X. Liu, L. Xie, J. Li, J. Ye, L. Dong, X. Cui, Y. Miao, D. Wang, J. Dong, C. Xiao, W. Chen, and H. Wang. Immune cell profiling of COVID-19 patients in the recovery stage by single-cell sequencing. *Cell Discov*, 6:31, May 2020.

81. D. Wu and X. O. Yang. TH17 responses in cytokine storm of COVID-19: An emerging target of JAK2 inhibitor fedratinib. *J. Microbiol. Immunol. Infect.*, 53(3):368–370, June 2020.

82. R. Yan, Y. Zhang, Y. Li, L. Xia, Y. Guo, and Q. Zhou. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science*, 367(6485):1444–1448, Mar. 2020.

83. C. Yin. Genotyping coronavirus SARS-CoV-2: methods and implications. *Genomics*, Apr. 2020.

84. P. Zhou, X.-L. Yang, X.-G. Wang, B. Hu, L. Zhang, W. Zhang, H.-R. Si, Y. Zhu, B. Li, C.-L. Huang, H.-D. Chen, J. Chen, Y. Luo, H. Guo, R.-D. Jiang, M.-Q. Liu, Y. Chen, X.-R. Shen, X. Wang, X.-S. Zheng, K. Zhao, Q.-J. Chen, F. Deng, L.-L. Liu, B. Yan, F.-X. Zhan, Y.-Y. Wang, G.-F. Xiao, and Z.-L. Shi. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, 579(7798):270–273, Mar. 2020.

85. Y. Zhou, Y. Hou, J. Shen, Y. Huang, W. Martin, and F. Cheng. Network-based drug repurposing for novel coronavirus 2019-nCoV/SARS-CoV-2. *Cell Discov*, 6:14, Mar. 2020.

86. Y. Zhou and Y. Zhu. Important role of the IL-32 inflammatory network in the host response against viral infection. *Viruses*, 7(6):3116–3129, June 2015.