Genome-wide bioinformatic analyses predict key host and viral factors in SARS-CoV-2 pathogenesis

Mariana G. Ferrarini[1,§], Avantika Lal[2,§], Rita Rebollo[1], Andreas Gruber[3], Andrea Guarracino[4], Itziar Martinez Gonzalez[5], Taylor Floyd[6], Daniel Siqueira de Oliveira[7], Justin Shanklin[8], Ethan Beausoleil[8], Taneli Pusa[9], Brett E. Pickett[8,#] Vanessa Aguiar-Pulido[6,#]

1 University of Lyon, INSA-Lyon, INRA, BF2I, Villeurbanne, France
2 NVIDIA Corporation, Santa Clara, CA, USA
3 Oxford Big Data Institute, Nuffield Department of Medicine, University of Oxford, Oxford, UK
4 Centre for Molecular Bioinformatics, Department of Biology, University Of Rome Tor Vergata, Rome, Italy
5 Amsterdam UMC, Amsterdam, The Netherlands
6 Center for Neurogenetics, Weill Cornell Medicine, Cornell University, New York, NY, USA
7 Laboratoire de Biométrie et Biologie Evolutive, Université de Lyon; Université Lyon 1; CNRS; UMR 5558, Villeurbanne, France
8 Brigham Young University, Provo, UT, USA
9 Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Belvaux, Luxembourg

§: These authors contributed equally
#: Corresponding authors
Keywords: SARS-CoV-2, COVID-19, gene expression, RNA-seq, RNA-binding proteins, host-pathogen interaction, transcriptomics

Abstract

The novel betacoronavirus named Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) caused a worldwide pandemic (COVID-19) after initially emerging in Wuhan, China. Here we applied a novel, comprehensive bioinformatic strategy to public RNA sequencing and viral genome sequencing data, to better understand how SARS-CoV-2 interacts with human cells. To our knowledge, this is the first meta-analysis to predict host factors that play a specific role in SARS-CoV-2 pathogenesis, distinct from other respiratory viruses. We identified differentially expressed genes, isoforms and transposable element families specifically altered in SARS-CoV-2 infected cells. Well-known immunoregulators including *CSF2*, *IL-32*, *IL-6* and *SERPINA3* were differentially expressed, while immunoregulatory transposable element families were overexpressed. We predicted conserved interactions between the SARS-CoV-2 genome and human RNA-binding proteins such as hnRNPA1, PABPC1 and eIF4b, which may play important roles in the viral life cycle. We also detected four viral sequence variants in the spike, polymerase, and nonstructural proteins that correlate with severity of COVID-19. The host factors we identified likely represent important mechanisms in the disease profile of this pathogen, and could be targeted by prophylactics and/or therapeutics against SARS-CoV-2.

Graphical Abstract

Introduction  ₁

In December of 2019 a novel betacoronavirus that was named Severe Acute Respiratory Syndrome ₂
Coronavirus 2 (SARS-CoV-2) emerged in Wuhan, China [4, 29]. This virus is responsible for causing ₃
the coronavirus disease of 2019 (COVID-19) and, by August 18 of 2020, it had already infected more ₄
than 21 million people worldwide, accounting for at least 750 thousand deaths ₅
(https://covid19.who.int). The SARS-CoV-2 genome is phylogenetically distinct from the ₆
SARS-CoV and Middle East Respiratory Syndrome (MERS) betacoronaviruses that caused human ₇
outbreaks in 2002 and 2012 respectively [79, 87]. Based on its high sequence similarity to a ₈
coronavirus isolated from bats [88], SARS-CoV-2 is hypothesized to have originated from bat ₉
coronaviruses, potentially using pangolins as an intermediate host before infecting humans [39]. ₁₀

SARS-CoV-2 infects human cells by binding to the angiotensin-converting enzyme 2 (ACE2) ₁₁
receptor [85]. Recent studies have sought to understand the molecular interactions between ₁₂
SARS-CoV-2 and infected cells [24], unfortunately, only a few have quantified gene expression ₁₃
changes in patient samples or cultured lung-derived cells infected by this virus [10, 46, 83]. These ₁₄
studies are essential to understanding the mechanisms of pathogenesis and immune response which ₁₅
can facilitate the development of treatments for COVID-19 [35, 54, 89]. ₁₆

Viruses generally trigger a drastic host response during infection. A subset of these specific ₁₇
changes in gene regulation is associated with viral replication, and therefore can pinpoint potential ₁₈
drug targets. In addition, transposable element (TE) overexpression has been observed upon viral ₁₉
infection [50], and TEs have been actively implicated in gene regulatory networks related to ₂₀
immunity [15]. Moreover, SARS-CoV-2 is a virus with a positive-sense, single-stranded, monopartite ₂₁
RNA genome. Such viruses are known to co-opt host RNA-binding proteins (RBPs) for diverse ₂₂
processes including viral replication, translation, viral RNA stability, assembly of viral protein ₂₃
complexes, and regulation of viral protein activity [22, 45]. ₂₄

In this work we identified a signature of altered gene expression that is consistent across ₂₅
published datasets of SARS-CoV-2 infected human lung cells. We present extensive results from ₂₆
functional analyses (signaling pathway enrichment, biological functions, transcript isoform usage, ₂₇
metabolic flux prediction, and TE overexpression) performed upon the genes that are differentially ₂₈
expressed during SARS-CoV-2 infection [10]. We also predict specific interactions between the ₂₉
SARS-CoV-2 RNA genome and human RBPs that may be involved in viral replication, transcription ₃₀
or translation, and identify viral sequence variations that are significantly associated with increased ₃₁
pathogenesis in humans. Knowledge of these molecular and genetic mechanisms is important to ₃₂
understand the SARS-CoV-2 pathogenesis and to improve the future development of effective ₃₃
prophylactic and therapeutic treatments. ₃₄

Results

We designed a comprehensive bioinformatics workflow to identify relevant host-pathogen interactions using a complementary set of computational analyses (Figure 1). First, we carried out an exhaustive analysis of differential gene expression in the only publicly available dataset to date from human lung cells infected with SARS-CoV-2 along with other respiratory viruses. We identified gene, isoform- and pathway-level responses that specifically characterize SARS-CoV-2 infection. Second, we predicted putative interactions between the SARS-CoV-2 RNA genome and human RBPs. Third, we identified a subset of these human RBPs which were also differentially expressed in response to SARS-CoV-2. Finally, we predicted four viral sequence variants that could play a role in disease severity.

Figure 1. Overview of the bioinformatic workflow applied in this study.

SARS-CoV-2 infection elicits a specific gene expression and pathway signature in human cells

We wanted to identify genes that were differentially expressed across multiple SARS-CoV-2 infected samples and not in samples infected with other respiratory viruses. As a primary dataset, we selected GSE147507 [10], which includes gene expression measurements from three cell lines derived from the human respiratory system (NHBE, A549, Calu-3) infected either with SARS-CoV-2, influenza A virus (IAV), respiratory syncytial virus (RSV), or human parainfluenza virus 3 (HPIV3), with different multiplicity of infection (MOI). We also analyzed an additional dataset GSE150316, which includes RNA-seq extracted from formalin fixed, paraffin embedded (FFPE) histological sections of lung biopsies from COVID-19 deceased patients and healthy individuals (see Figure 2A and Materials and Methods for further details).

Hence, we retrieved 41 differentially expressed genes (DEGs) that showed significant and consistent expression changes in at least three datasets from cell lines infected with SARS-CoV-2, and that were not significantly affected in cell lines infected with other viruses within the same dataset (Supplementary Table 1A). To these, we added 23 genes that showed significant and consistent expression changes in two of four cell line datasets infected with SARS-CoV-2 and at least

one lung biopsy sample from a SARS-CoV-2 patient. Results coming from FFPE sections were less ₆₃
consistent presumably due to the collection of biospecimens from different sites within the lung. ₆₄
Thus, the final set consisted of 64 DEGs: 48 up-regulated and 16 downregulated of which 38 had an ₆₅
absolute Log2FC > 1 in at least one dataset (relevant genes from this list are shown in Table 1). ₆₆

SERPINA3, an antichymotrypsin which was proposed as an interesting candidate for the ₆₇
inhibition of viral replication [13], was the only gene specifically upregulated in the four cell line ₆₈
datasets tested (Table 1). Other interesting up-regulated genes were the amidohydrolase *VNN2*, the ₆₉
pro-fibrotic gene *PDGFB*, the beta-interferon regulator *PRDM1* and the proinflammatory cytokines ₇₀
CSF2 and *IL-32*. *FKBP5*, a known regulator of NF-kB activity, was among the consistently ₇₁
downregulated genes. We also generated additional lists of DEGs that met different filtering criteria ₇₂
(Supplementary Table 1B, see Supplementary File 1 for the complete DEG results for each dataset). ₇₃

In order to better understand the underlying biological functions and molecular mechanisms ₇₄
associated with the observed DEGs, we performed a hypergeometric test to detect statistically ₇₅
significant overrepresented gene ontology (GO) terms among the DEGs having an absolute Log2FC ₇₆
> 1 in each dataset separately. ₇₇

Table 1. Log2FC for selected genes that showed significant up-or down-regulation in SARS-CoV-2 ₇₈
infected samples (FDR-adjusted p-value < 0.05), and not in samples infected with the other viruses ₇₉
tested. Log2FC values are only provided for statistically significant samples. ₈₀

Gene	Cell Type and MOI				Biopsies	
	A549 MOI 0.2	A549	Calu-3 MOI 2	NHBE	Case 1	Case 3
VNN2		6.18	0.42	6.13		
CSF2		3.56	7.30	2.70		
WNT7A		4.99	0.79	0.45		
PDZK1IP1		1.72	0.70	2.28		
SERPINA3	0.49	1.39	0.77	1.44		
RHCG		1.51	2.02	1.33	2.53	
IL32		1.64	1.23	1.21		
PDGFB		1.91	1.75	1.00		
ALDH1A3		1.09	1.32	0.39		
TLR2		1.63	0.89	0.84		
G0S2		0.66	3.79	0.83		
NRCAM		0.73	1.82	0.78		
SERPINB1		0.61	1.17	0.72		
PRDM1		0.82	3.49	0.59		
MT-TN		0.55	1.70	0.33		
ATF4		0.79	1.07	0.26		
BHLHE40		0.75	1.56	0.18		
PTPN12	0.48	0.97	1.23			
GPCPD1	0.36	0.94	1.69			
DUSP16	0.33	0.41	1.43			
FKBP5		-0.39		-0.36	-1.47	-2.14
DAP		-0.18	-0.61		-1.16	
FECH		-0.27	-0.36		-1.54	
MT-CYB		-0.30	-0.26		-3.68	
EIF4A1		-0.33	-0.63		-1.85	
POLE4	-0.23	-0.82	-1.24			
DDX39A	-0.23	-1.27	-0.54			
CENPP		-0.36	-0.40	-0.38		
TMEM50B		-0.48	-0.59	-0.53		
HPS1	-0.28	-0.31	-0.62			
SNX8	-0.30	-0.43	-0.56			

₈₁

₈₂

Figure 2. Overview of the RNA-seq based results specific to SARS-CoV-2 which were not detected in the other viral infections (IAV, HPIV3 and RSV). (A) Representation of the RNA-seq studies used in our analyses. (B) Non-redundant functional enrichment of DEGs. Here we report a subset of non-redundant reduced terms consistently enriched in more than one SARS-COV-2 cell line which were not detected in the other viruses' datasets. We added generic categories of immunity, metabolism, cellular processes and signaling/epigeneitcs to the GO terms as colored dots. (C) Top 20 differentially expressed isoforms (DEIs) in SARS-CoV-2 infected samples. Y-axis denotes the differential usage of isoforms (dIF) whereas x-axis represents the overall log2FC of the corresponding gene. Thus, DEIs also detected as DEGs by this analysis are depicted in blue. (D) The upper right diagram depicts different manners by which TE family overexpression might be detected. While TEs may indeed be autonomously expressed, the old age of most TEs detected points toward either being part of a gene (exonization or alternative promoter), or a result of pervasive transcription. We report the functional enrichment for neighboring genes of differentially expressed TEs (DETEs) specifically upregulated in SARS-CoV-2 Calu-3 and A549 cells (MOI 2). The same categories used in subfigure (B) were attributed to the GO terms reported here.

Consistent with the findings of Blanco-Melo et al. [10], GO enrichment analysis returned terms 98
associated with immune system processes, Pi3K/AKT signaling pathway, response to cytokine, 99
stress and virus, among others (see Supplementary File 2 for complete results). In addition, we 100
report 285 GO terms common to at least two cell line datasets infected with SARS-CoV-2, and 101
absent in the response to other viruses (Figure 2B, Supplementary Table 2A), including neutrophil 102
and granulocyte activation, interleukin-1-mediated signaling pathway, proteolysis, and stress 103
activated signaling cascades. 104

Next, we wanted to pinpoint intracellular signaling pathways that may be modulated specifically 105
during SARS-CoV-2 infection. A robust signaling pathway impact analysis (SPIA) enabled us to 106
identify 30 pathways, including many involved in the host immune response, that were significantly 107
enriched among differentially expressed genes in at least one virus-infected cell line dataset 108
(Supplementary Table 3). More importantly, we predicted four pathways to be specific to 109
SARS-CoV-2 infection and observed that the significant pathways differed by cell type and 110
multiplicity of infection. The significant results included only one term common to A549 (MOI 0.2) 111
and Calu-3 cells (MOI 2), namely the interferon alpha/beta signaling. Additionally, we found the 112
amoebiasis (A549 cells, MOI 0.2), the p75(NTR)-mediated and the trka receptor signaling pathways 113
(A549 cells, MOI 2) as significantly impacted. 114

We also used a classic hypergeometric method as a complementary approach to our SPIA 115
pathway enrichment analysis. While there were generally higher numbers of significant results using 116
this method, we observed that the vast majority of enriched terms (FDR < 0.05) described 117
infections with various pathogens, innate immunity, metabolism, and cell cycle regulation 118
(Supplementary Table 3). Interestingly, we were able to detect enriched KEGG pathways common to 119
at least two SARS-CoV-2 infected cell types and absent from the other virus-infected datasets 120
(Figure 2B, Supplementary Table 2B). These included pathways related to infection, cell cycle, 121
endocytosis, signalling pathways, cancer and other diseases. 122

SARS-CoV-2 infection results in altered lipid-related metabolic fluxes 123

To integrate the gene expression changes with metabolic activity in response to virus infection, we 124
projected the transcriptomic data onto the human metabolic network [77]. This analysis detected 125
common decreased fluxes in inositol phosphate metabolism in both A549 and Calu-3 cells infected 126
with SARS-CoV-2 at a MOI of 2 (Supplementary Table 4). The consensus solution (obtained taking 127
into account the enumeration of all solutions) in A549 cells (MOI 2) also recovered decreased fluxes 128
in several lipid pathways: fatty acid, cholesterol, sphingolipid, and glycerophospholipid. In addition, 129
we detected an increased flux common to A549 and Calu-3 cell lines in reactive oxygen species (ROS) 130
detoxification, in accordance with previous terms recovered from functional enrichment analyses. 131

SARS-CoV-2 infection induced an isoform switch of genes associated 132
with immunity and mRNA processing 133

We wanted to analyze changes in transcript isoform expression and usage associated with 134
SARS-CoV-2 infection, as well as to predict whether these changes might result in altered protein 135
function. We identified isoforms experiencing a switch in usage greater than or equal to 30% in 136
absolute value, and retrieved those with a Bonferroni-adjusted p-value less than 0.05. After 137
calculating the difference in isoform usage (dIF) per gene (in each condition), we performed 138
predictive functional consequence and alternative splicing analyses for all isoforms globally as well as 139
at the individual gene level. 140

We observed 3,569 differentially expressed isoforms (DEIs) across all samples (Supplementary 141
Figure 1A, Supplementary Table 5A). Results indicate that isoforms from A549 cells infected with 142
RSV, IAV and HPIV3 exhibited significant differences in biological events such as complete open 143
reading frame (ORF) loss, shorter ORF length, intron retention gain and decreased sensitivity to 144
nonsense mediated decay (Supplementary Figure 1B). These conditions also displayed various 145
changes in splicing patterns, ranging from loss of exon skipping events, changes in usage of 146

alternative transcription start and termination sites, and decreased alternative 5' and 3' splice sites   147
(Supplementary Figure 1C).   148

In contrast, isoforms from SARS-CoV-2 infected samples displayed no significant global changes   149
in biological consequences or alternative splicing events between conditions (Supplementary Figures   150
1A and 1B respectively). Trends indicated transcripts in SARS-CoV-2 samples experienced decreases   151
in ORF length, numbers of domains, coding capability, intron retention and nonsense mediated   152
decay (Supplementary Figure 1A). These biological consequences may result from increased multiple   153
exon skipping events and alternative transcription start sites via alternative 5' acceptor sites   154
(Supplementary Figure 1B). While not significant, these trends implicate that the SARS-CoV-2 virus   155
may globally trigger host cell machinery to generate shorter isoforms that, while not shuttled for   156
degradation, either do not produce functional proteins or produce alternative aberrant proteins not   157
utilized in non-SARS-CoV-2 tissue conditions.   158

Despite the lack of global biological consequence and splicing changes, individual isoforms from   159
SARS-CoV-2 infected samples experienced significant changes in gene expression and isoform usage   160
(Figure 2C). Top-expressing genes were associated with cellular processes such as immune response   161
and antiviral activity (*IFI44L*, *IL6*, *MX1*, *TRIM5*), transcription and mRNA processing (*DDX10*,   162
HNRNPA3F6, *JMJD7*, *ZNF487*, *ZNF599*) and cell cycle and survival (*BCL2L2-PABPN1*, *CDCA3*)   163
(Supplementary Table 5B). Similarly, significant genes from non-SARS-CoV-2 samples were   164
associated with processes such as immune cell development and response (*ADCY7*, *BATF2*, *C9orf72*,   165
ETS1, *GBP2*, *IFIT3*), transcription regulation and DNA repair (*ABHD14B*, *ATF3*, *IFI16*,   166
POLR2J2, *SMUG1*, *ZNF19*, *ZNF639*), mitochondrial function (*ATP5E*, *BCKDH8*, *TST*, *TXNRD2*),   167
and GTPase activity (*GBP2*, *RAP1GAP*, *RGS20*, *RHOBTB2*) (Supplementary Figure 1D,   168
Supplementary Table 5B).   169

Upon further inspection, we noticed that *IL-6*, a gene encoding a cytokine involved in acute and   170
chronic inflammatory responses, displayed 3 and 4-fold increases in expression in NHBE and A549   171
cells, respectively (infected with a MOI of 2) (Supplementary Figure 1B). To date, the Ensembl   172
Genome Reference Consortium has identified 9 *IL-6* isoforms in humans, with the traditional   173
transcript having 6 exons (*IL6-204*), 5 of which contain coding elements. NHBE cells expressed 4   174
known *IL-6* isoforms, while A549 cells expressed 1 unknown and 6 known isoforms. When evaluating   175
the actual isoforms used across conditions, NHBE cells used 3 out of 4 isoforms observed, while A549   176
cells used all 7 observed isoforms. Isoform usage is evaluated based on isoform fraction (IF), or the   177
percentage of an isoform found relative to all other identified isoforms associated with a specific gene.   178
For example, in the case of NHBE SARS-CoV-2 samples, the IF for the *IL6-201* isoform = 0.75,   179
IL6-204 = 0.05, *IL6-206* = 0.09, *IL6-209* = 0.06, and the sum of these IF values = 0.95, or 95%   180
usage of the *IL-6* gene. Both SARS-CoV-2 samples exhibited exclusive usage of non-canonical   181
isoform *IL6-201*, and inversely, mock samples almost exclusively utilized the *IL6-204* transcript. In   182
NHBE infected cells, isoform *IL6-201* experienced a significant increase in usage (dIF = 0.75) and   183
IL6-204 a significant decrease in usage (dIF = -0.95) when compared to mock conditions. Similarly,   184
isoform *IL6-201* in A549 infected cells experienced an increase in usage (dIF = 0.58), while uses of   185
all other isoforms remained non-significant in comparison to mock conditions.   186

Overexpression of TE families close to immune-related genes upon   187
SARS-CoV-2 infection   188

In order to estimate the expression of TE families and their possible roles in SARS-CoV-2 infection,   189
we mapped the RNA-seq reads against all annotated human TE families and detected DETEs   190
(Supplementary File 3). We found 68 common TE families upregulated in SARS-CoV-2 infected   191
A549 and Calu-3 cells (MOI 2). From this list, we excluded all TE families detected in A549 cells   192
infected with the other viruses. This allowed us to identify 16 families that were specifically   193
upregulated in Calu-3 and A549 cells infected with SARS-CoV-2 and not in the other viral infections.   194

The 16 families identified were MER77B, MamRep4096, MLT2C2, PABL_A, Charlie9, MER34A,   195
L1MEg1, LTR13A, L1MB5, MER11C, MER41B, LTR79, THE1D-int, MLT1I, MLT1F1,   196

MamRep137. Most of the TE families uncovered are ancient elements, incapable of transposing, or ₁₉₇
harboring intrinsic regulatory sequences [37, 57, 70]. Eleven of the 16 TE families specifically ₁₉₈
upregulated in SARS-COV-2 infected cells are long terminal repeat (LTR) elements, and include well ₁₉₉
known TE immune regulators. For instance, the MER41B (primate specific TE family) is known to ₂₀₀
contribute to Interferon gamma inducible binding sites (bound by STAT1 and/or IRF1) [14, 66]. ₂₀₁
Other LTR elements are also enriched in STAT1 binding sites (MLT1L) [14], or have been shown to ₂₀₂
act as cellular gene enhancers (LTR13A [16, 32]). ₂₀₃

Given the propensity for the TE families detected to impact nearby gene expression, we further ₂₀₄
investigated the functional enrichment of genes near upregulated TE families (+- 5kb upstream, 1kb ₂₀₅
downstream). We detected GO functional enrichment of several immunity-related terms (e.g. MHC ₂₀₆
protein complex, antigen processing, regulation of dendritic cell differentiation, T-cell tolerance ₂₀₇
induction), metabolism related terms (such as regulation of phospholipid catabolic process), and ₂₀₈
more interestingly a specific human phenotype term called "Progressive pulmonary function ₂₀₉
impairment" (Figure 2D). Even though we did not limit our search only to neighboring genes which ₂₁₀
were also DE, we found several similar (and very specific) enriched terms in both analyses, for ₂₁₁
instance related to immune response, endosomes, endoplasmic reticulum, vitamin (cofactor) ₂₁₂
metabolism, among others. This result supports the idea that some responses during infection could ₂₁₃
be related to TE-mediated transcriptional regulation. Finally, when we searched for enriched terms ₂₁₄
related to each one of the 16 families separately, we also detected immunity related enriched terms ₂₁₅
such as regulation of interleukins, antigen processing, TGFB receptor binding and temperature ₂₁₆
homeostasis (Supplementary File 4). It is important to note that given the old age of some of the ₂₁₇
TEs detected, overexpression might be associated with pervasive transcription, or inclusion of TE ₂₁₈
copies within unspliced introns (see upper box in Figure 2D). ₂₁₉

The SARS-CoV-2 genome is enriched in binding motifs for 40 human ₂₂₀ RBPs, most of them conserved across SARS-CoV-2 genome isolates ₂₂₁

Our next aim was to predict whether any host RNA binding proteins interact with the viral genome. ₂₂₂
To do so, we first filtered the AtTRACT database [23] to obtain a list of 102 human RBPs and 205 ₂₂₃
associated Position Weight Matrices (PWMs) describing the sequence binding preferences of these ₂₂₄
proteins. We then scanned the SARS-CoV-2 reference genome sequence to identify potential binding ₂₂₅
sites for these proteins. Figure 3 illustrates our analysis pipeline. ₂₂₆

We identified 99 human RBPs with 11,897 potential binding sites in the SARS-CoV-2 ₂₂₇
positive-sense genome. Since the SARS-CoV-2 genome produces negative-sense intermediates as part ₂₂₈
of the replication process [36], we also scanned the negative-sense molecule, where we found 11,333 ₂₂₉
potential binding sites for 96 RBPs (Supplementary Table 6). ₂₃₀

To find RBPs whose binding sites occur in the SARS-CoV-2 genome more often than expected by ₂₃₁
chance, we repeatedly scrambled the genome sequence to create 1,000 simulated genome sequences ₂₃₂
with an identical nucleotide composition to the SARS-CoV-2 genome sequence (30% A, 18% C, 20% ₂₃₃
G, 32% T). We used these 1,000 simulated genomes to determine a background distribution of the ₂₃₄
number of binding sites found for a specific RBP. This allowed us to pinpoint RBPs with ₂₃₅
significantly more or significantly fewer binding sites in the actual SARS-CoV-2 genome than ₂₃₆
expected based on the background distribution (two-tailed z-test, FDR-corrected P < 0.01). To ₂₃₇
retrieve RBPs whose motifs were enriched in specific genomic regions, we also repeated this analysis ₂₃₈
independently for the SARS-CoV-2 5'UTR, 3'UTR, intergenic regions, and for the sequence from the ₂₃₉
negative sense molecule. Motifs for 40 human RBPs were found to be enriched in at least one of the ₂₄₀
tested genomic regions, while motifs for 23 human RBPs were found to be depleted in at least one of ₂₄₁
the tested regions (Supplementary Table 7). ₂₄₂

We next examined whether any of the 6,936 putative binding sites for these 40 enriched RBPs ₂₄₃
were conserved across SARS-CoV-2 isolates. We found that 6,581 putative binding sites, ₂₄₄
representing 34 RBPs, were conserved across more than 95% of SARS-CoV-2 genome sequences in ₂₄₅
the GISAID database (≥ 26,213 out of 27,592 genomes). However, this is of limited significance as ₂₄₆

RBP binding sites in coding regions are likely to be conserved due to evolutionary pressure on protein sequences rather than RBP binding ability. We therefore repeated this analysis focusing only on putative RBP binding sites in the SARS-CoV-2 UTRs and intergenic regions. We found 124 putative RBP binding sites for 21 enriched RBPs in the UTRs and intergenic regions. Of these, 50 putative RBP binding sites for 17 RBPs were conserved in >95% of the available genome sequences; 6 in the 5'UTR, 5 in the 3'UTR, and 39 in intergenic regions (Supplementary Table 8).

Figure 3. Workflow and selected results for analysis of potential binding sites for human RNA-binding proteins in the SARS-CoV-2 genome.

Subsequently, we interrogated publicly available data to validate the putative SARS-CoV-2 / RBP interactions (Supplementary Table 9). According to GTEx data [25], 39 of the 40 enriched RBPs and all 23 of the depleted RBPs were expressed in human lung tissue. Further, 31 of 40 enriched RBPs and 22 of 23 depleted RBPs were co-expressed with the ACE2 and TMPRSS2 receptors in single-cell RNA-seq data from human lung cells (GSE122960; [25, 64]), indicating that

they are present in cells that are susceptible to SARS-CoV-2 infection. We next checked whether any of these RBPs are known to interact with SARS-CoV-2 proteins and found that human poly-A binding proteins C1 and C4 (PABPC1 and PABPC4) bind to the viral N protein [24]. Thus, it is conceivable that these RBPs interact with both the SARS-CoV-2 RNA and proteins. Finally, we combined these results with our analysis of differential gene expression to identify SARS-CoV-2 interacting RBPs that also show expression changes upon infection. The results of this analysis are summarized for selected RBPs in Table 2.

Table 2. Selected conserved human RBPs predicted to interact with the SARS-CoV-2 genome along with experimental information.

RPB	DE Analysis*1			Experimental evidence in human datasets				RBP binding site prediction	
	A549 LogFC	Calu-3 LogFC	SARS-CoV-2 Specific DEG	GTEx Lung Tissue (TPM)	scRNA*2	PPI Map*3	Interaction with viral RNA*4	Conserved*5	Region
HNRNPA1		-0.32	☐	331.336	✔		✔	✔	
HNRNPA2B1	-1.08	-0.29	☐	539.829	✔		✔	✔	
PABPC1	0.72	0.44	✔	448.025	✔	N	✔	✔	**3'UTR**
PABPC4	0.30	-0.28	☐	103.082	✔	N	✔	✔	
PPIE		-0.27	☐	13.827	✔		☐	✔	
CELF5	0.56		☐	0.079	✔		☐	✔	
FMR1		0.75	☐	21.435	✔		☐	✔	**5'UTR**
RBM24	0.34		☐	1.412	☐		☐	✔	
EIF4B	0.53	0.64	☐	170.303	✔		✔	✔	
ELAVL1		-0.31	☐	27.440	✔		☐	✔	
PABPC1	0.72	0.44	✔	448.025	✔	N	✔	✔	**Intergenic**
PPIE		-0.27	☐	13.827	✔		☐	✔	
TIA1	0.34	0.41	✔	46.934	✔		☐	✔	
TIAL1	0.25		☐	40.593	☐		☐	✔	

*1 LogFC reported only if padj < 0.05
*2 scRNA expression in ACE+ and TMPRSS2+ lung cells: dataset GSE122960
*3 PPI Map: Experimental map of protein-protein interactions between human and viral proteins (Gordon *et al*., 2020)
*4 Preprint: Experimental study revealing proteins interacting with SARS-CoV-2 RNA in a human liver cell line (Schmidt *et al*., 2020)
*5 Conserved in SARS-CoV-2 genomes

Motif enrichment in SARS-CoV-2 differs from related coronaviruses

We repeated the above analysis to calculate the enrichment and depletion of RBP-binding motifs in the genomes of two related coronaviruses: the SARS-CoV virus (Supplementary Table 10) that caused the SARS outbreak in 2002-2003, and RaTG13 (Supplementary Table 11), a bat coronavirus with a genome that is 96% identical with that of SARS-CoV-2 [4, 88].

We found that the pattern of enrichment and depletion of RBP binding motifs in SARS-CoV-2 is different from that of the other two viruses. Specifically, the SARS-CoV-2 genome is uniquely enriched for binding sites of CELF5 in its 5'UTR, PPIE on its 3'UTR, and ELAVL1 in the viral negative-sense RNA molecule. These three proteins are involved in RNA metabolism and are important for RNA stability (ELAVL1, CELF5) and processing (PPIE). Despite the high sequence identity between the two genomes, the single binding site for CELF5 on the SARS-CoV-2 5'UTR is conserved in 97% of available SARS-CoV-2 genome sequences but absent in the 5'UTR of RaTG13.

A subset of viral genome variants correlate with increased COVID-19 severity

To test whether any viral sequence variants were associated with a change in disease severity in human hosts, we analyzed 1,511 complete SARS-CoV-2 genomes that had associated clinical metadata. The FDR-corrected statistical results from this analysis revealed four nucleotide variations that were significantly associated with a change in viral pathogenesis. Three of these nucleotide changes resulted in non-synonymous variations at the amino acid level, while the last one was silent at the amino acid level. The first position was a $T \to G$ (L37F) substitution located in the Nsp6 coding region (p < 1.48E-5), the second position was a $C \to T$ (P323L) substitution located in the RNA-dependent RNA polymerase coding region (p < 2.01E-4), the third position was an $A \to G$ (D614G) substitution located in the spike coding region (p < 1.61E-4), and the fourth was a synonymous $C \to T$ substitution located in the Nsp3 coding region (p < 1.77E-4). As a further validation step, we performed the same analysis comparing viral sequence variants against potential confounders, such as the biological sex or age group of the patients. These comparisons validated that these four positions were only identified as significant in the results of the disease severity analysis.

Discussion

Airway epithelial cells are the primary entry points for respiratory viruses and therefore constitute the first producers of inflammatory signals that, in addition to their antiviral activity, promote the initiation of the innate and adaptive immune responses. Here, we report the results of a complementary panel of analyses that enable a better understanding of host-pathogen interactions which contribute to SARS-CoV-2 replication and pathogenesis in the human respiratory system. Moreover, we propose already established along with novel human factors exclusively detected in SARS-CoV-2 infected cells by our analyses that might be relevant in the context of COVID-19 and which are worth being further investigated at an experimental level (Figure 4).

The *CSF2* gene, which encodes the Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF), was among the most highly up-regulated genes in SARS-CoV-2 infected cells. GM-CSF induces survival and activation in mature myeloid cells such as macrophages and neutrophils. However, GM-CSF is considered more proinflammatory than other members of its family, such as G-CSF, and is associated with tissue hyper-inflammation [52]. In accordance with our results, high levels of GM-CSF were found in the blood of severe COVID-19 patients [84], and several clinical trials are planned using agents that either target GM-CSF or its receptor [53]. GM-CSF, together with other proinflammatory cytokines such as IL-6, TNF, IFNg, IL-7 and IL-18, is associated with the cytokine storm present in a hyperinflammatory disorder named hemophagocytic lymphohistiocytosis (HLH) which presents with organ failure [12]. Moreover, cytokines related to cytokine release syndrome (such as IL-1A/B, IL-6, IL-10, IL-18, and TNFA), showed increased positive association to the severity of the disease in the blood from COVID-19 patients [49]. Another proinflammatory cytokine specifically upregulated in SARS-CoV-2 infected cells was *IL-32*, which together with *CSF2*, promotes the release of TNF and IL-6 in a continuous positive loop and therefore contribute to this cytokine storm [90]. Interestingly, *IL-6*, *IL-7* and *IL-18* were found to be upregulated in two of the four data sets of SARS-CoV-2 infected cells. Moreover, not only upregulation, but also a shift in isoform usage of *IL-6* was detected in NHBE and A549 infected cells. A shift in 5' UTR usage in the presence of SARS-CoV-2 may be attributed to indirect host cell signaling cascades that trigger changes in transcription and splicing activity, which could also explain the overall increase in *IL-6* expression.

SERPINA3, a gene coding for an essential enzyme in the regulation of leukocyte proteases, is also induced by cytokines [28]. This was the only gene consistently upregulated in all cell line samples infected with SARS-CoV-2 and absent from the other datasets. Even though this was previously proposed as a promising candidate for the inhibition of viral replication, to date no experiments were

carried out to validate this hypothesis [13]. Another interesting candidate gene, which has not been   337
implicated experimentally in respiratory viral infections and was upregulated in our analysis, was   338
VNN2. Vanins are involved in proinflammatory and oxidative processes, and VNN2 plays a role in   339
neutrophil migration by regulating b2 integrin [56]. In contrast, the downregulated genes included   340
SNX8, which has been previously reported in RNA virus-triggered induction of antiviral   341
genes [13, 26]; and *FKBP5*, a known regulator of NF-kB activity [27]. These results suggest that the   342
SARS-CoV-2 virus tends to indirectly target specific genes involved in genome replication and host   343
antiviral immune response without eliciting a global change in cellular transcript processing or   344
protein production.   345

Figure 4. Overview of human factors specific to SARS-CoV-2 infection detected by our analyses. This   348
includes human RBPs whose binding sites are enriched and conserved in the SARS-CoV-2 genome but not in   349
the genomes of related viruses; and genes, isoforms and metabolites that are consistently altered in response   350
to SARS-CoV-2 infection of lung epithelial cells but not in infection with the other tested viruses; ECM   351
(extracellular matrix).   352

One of the first and most important antiviral responses is the production of type I Interferon   355
(IFN). This protein induces the expression of hundreds of Interferon Stimulated Genes (ISGs), which   356
in turn serve to limit virus spread and infection. Moreover, type I IFN can directly activate immune   357
cells such as macrophages, dendritic cells and NK cells as well as induce the release of   358
proinflammatory cytokines by other cell types [34]. Signaling pathway analysis showed that type I   359
IFN response was greatly impacted in SARS-CoV-2 infected cells (A549 and Calu-3 cells at a MOI   360

of 0.2 and 2 respectively). In the same direction, a higher expression of *PRDM1* (Blimp-1) that we observed in the SARS-CoV-2 infected cells, could also contribute to the critical regulation of IFN signaling cascades; interestingly, the TE family LTR13, which was also upregulated upon SARS-CoV-2 infection, is enriched in PRDM1 binding sites [78]. Therefore, it is possible that regulatory factors involved in IFN and immune response in the context of SARS-CoV-2 infection could also be attributed to TE transcriptional activation. In the same direction, we detected the upregulation of several TE families in SARS-CoV-2 infected cells that have been previously implicated in immune regulation. Moreover, 16 upregulated families were specific to SARS-CoV-2 infection in Calu-3 and A549 cell lines. The MER41B family, for instance, is known to contribute to interferon gamma inducible binding sites (bound by STAT1 and/or IRF1) [14]. Functional enrichment analysis of nearby genes were in accordance with these findings, since several immunity related terms were enriched along with "progressive pulmonary impairment". In parallel, TEs seem to be co-regulated with phospholipid metabolism, which directly affects the Pi3K/AKT signaling pathway, central to the immune response and which were detected in our functional enrichment and metabolism flux analyses.

RBPs are another example of host regulatory factors involved either in the response of human cells to SARS-CoV-2 or in the manipulation of human machinery by the virus. We aimed at finding RBPs which potentially interact with SARS-CoV-2 genomes in a conserved and specific way. Five of the proteins predicted to be interacting with the viral genome by our pipeline (EIF4B, hnRNPA1, PABPC1, PABPC4, and YBX1) were experimentally shown to bind to SARS-CoV-2 RNA in an infected human liver cell line, based on a recent preprint [67].

Among the RBPs whose potential binding sites were enriched and conserved within the SARS-CoV-2 virus genomes is the EIF4B, suggesting that the SARS-CoV-2 virus protein translation could be EIF4B-dependent. We also detected the upregulation of *EIF4B* in A549 and Calu-3 cells, which might indicate that this protein is sequestered by the virus and therefore the cells need to increase its production. Moreover, this protein was predicted to interact specifically with the intergenic region upstream of the gene encoding the SARS-CoV-2 membrane (M) protein, one of four structural proteins from this virus.

Another conserved RBP, which was also upregulated in infected cells, is the Poly(A) Binding Protein Cytoplasmic 1 (PABPC1), which has well described cellular roles in mRNA stability and translation. PABPC1 has been previously implicated in multiple viral infections. The activity of PABPC1 is modulated to inhibit host protein transcript translation, promoting viral RNA access to the host cell translational machinery [72]. Importantly, the 3' UTR region of SARS-CoV2 is also enriched in binding sites of the PABPC1 and the PPIE RBPs, the latter of which is known to be involved in multiple processes, including mRNA splicing [9, 33]. Interestingly, PABPC1 and PABPC4 interact with the SARS-CoV-2 N protein, which stabilizes the viral genome [24]. This raises the possibility that the viral genome, N protein, and human PABP proteins may participate in a joint protein-RNA complex that assists in viral genome stability, replication, and/or translation [1, 59, 62, 71, 72].

An interesting result was that the binding motifs for hnRNPA1, which has been shown to interact with other coronavirus genomes, were enriched specifically in the 3'UTR of SARS-CoV-2 even though they were depleted in the genome overall. The hnRNPA1 protein was described to interact more in particular with multiple sequence elements including the 3'UTR of the Murine Hepatitis Virus (MHV), and to participate in both transcription and replication of this virus [31, 44, 68]. This particular gene, along with *hnRNPA2B1*, were downregulated in Calu-3 cells and in contrast to the previous examples of upregulated genes, could denote a specific response of the human cells to control viral replication.

Cross referencing the results from our statistical analysis of $\sim 5\%$ of the available genomes ($\sim 1,500$ out of $> 27,000$ in GISAID) with clinical metadata revealed interesting new insights. Indeed, the $D \rightarrow G$ mutation at amino acid position 614 in the Spike protein found in our analysis has recently been proposed to have increased viral infectivity [38]. In addition, this same mutation has also been associated with an increase in the case fatality rate [6], however, these hypotheses need

further verification. The P323L mutation in the RNA-dependent RNA polymerase (RdRP) has been 413
identified previously, although in that study it was associated with changes in geographical location 414
of the viral strain [58]. Finally, the L37F mutation in the Nsp6 protein has been reported to be 415
located outside of the transmembrane domain [11], being present at a high frequency [86], and 416
proposed to negatively affect protein structure stability [8]. Our statistics may contain bias based on 417
the number of genome sequences being collected earlier *versus* later in the pandemic, genomes 418
lacking clinical outcome metadata, and in the case of the Spike D614G a potential increase of fitness 419
associated with this mutation. However, the fact that more than one of our predictions has also been 420
detected by different studies justifies future wet lab experiments to compare the effect of the other 421
identified mutations. 422

Conclusion 423

Overall, our analyses identified sets of statistically significant host genes, isoforms, regulatory 424
elements, and other interactions that likely contribute to the cellular response during infection with 425
SARS-CoV-2. Furthermore, we detected potential binding sites for human RBPs that are conserved 426
across SARS-CoV-2 genomes, along with a subset of variants in the viral genome that correlate well 427
with disease severity in SARS-CoV-2 infection. To our knowledge, this is the first work where a 428
computational meta-analysis was performed to predict host factors that play a role in the specific 429
pathogenesis of SARS-CoV-2, distinct from other respiratory viruses. Noteworthy, we used the only 430
study available to date in which cells were infected with SARS-CoV-2 along with other viruses. 431
Other studies have focused on the sequencing of fluids (blood or bronchoalveolar lavage fluid), and 432
may not reflect directly the cellular response to this virus. The availability of new high quality, well 433
annotated datasets of patients and cells infected with SARS-CoV-2 will be valuable for the 434
verification of our predictions. We envision that applying our publicly available workflow will yield 435
important mechanistic insights in future analyses on emerging pathogens. Similarly, we expect that 436
the results for SARS-CoV-2 will contribute to ongoing efforts in the selection of new drug targets 437
and the development of more effective prophylactics and therapeutics to reduce virus infection and 438
replication with minimal adverse effects on the human host. 439

Materials and Methods 440

Datasets 441

Two datasets were downloaded from the Gene Expression Omnibus (GEO) database, hosted at the 442
National Center for Biotechnology Information (NCBI). The first dataset, GSE147507 [10], includes 443
gene expression measurements from three cell lines derived from the human respiratory system 444
(NHBE, A549, Calu-3) infected either with SARS-CoV-2, influenza A virus (IAV), respiratory 445
syncytial virus (RSV), or human parainfluenza virus 3 (HPIV3). The second dataset, GSE150316, 446
includes RNA-seq extracted from formalin fixed, paraffin embedded (FFPE) histological sections of 447
lung biopsies from COVID-19 deceased patients and healthy individuals. This dataset encompasses a 448
variable number of biopsies per subject, ranging from one to five. Given its limitations, we only 449
utilized the second dataset for differential expression analysis. 450

The reference genome sequences of SARS-CoV-2 (NC_045512), RaTG13 (MN996532.1), and 451
SARS-CoV (NC_004718.3) were downloaded from NCBI. Additionally, a list of known RBPs and 452
their PWMs were downloaded from ATtRACT (https://attract.cnic.es/download). Finally, all 453
SARS-CoV-2 complete genomes collected from humans and that had disease severity information 454
were downloaded from GISAID on 19 May, 2020 [69]. 455

RNAseq data processing and differential expression analysis   456

Data was downloaded from SRA using sra-tools (v2.10.8; https://github.com/ncbi/sra-tools) and   457
transformed to fastq with fastq-dump. FastQC (v0.11.9; https://github.com/s-andrews/FastQC)   458
and MultiQC (v1.9) [20] were employed to assess the quality of the data used and the need to trim   459
reads and/or remove adapters. Selected datasets were mapped to the human reference genome   460
(GENCODE Release 19, GRCh37.p13) utilizing STAR (v2.7.3a) [17]. Alignment statistics were used   461
to determine which datasets should be included in subsequent steps. Resulting SAM files were   462
converted to BAM files employing samtools (v1.9) [43]. Next, read quantification was performed   463
using StringTie (v2.1.1) [60] and the output data was postprocessed with an auxiliary Python script   464
provided by the same developers to produce files ready for subsequent downstream analyses. For the   465
second gene expression dataset, raw counts were downloaded from GEO. DESeq2 (v1.26.0) [47] was   466
used in both cases to identify differentially expressed genes (DEGs). Finally, an exploratory data   467
analysis was carried out based on the transformed values obtained after applying the variance   468
stabilizing transformation [3] implemented in the vst() function of DESeq2 [48]. Hence, principal   469
component analysis (PCA) was performed to evaluate the main sources of variation in the data and   470
remove outliers.   471

Gene ontology enrichment analysis   472

The DEGs produced by DESeq2 with an absolute Log2FC > 1 and FDR-adjusted p-value < 0.05   473
were used as input to a general GO enrichment analysis [5, 76]. Each term was verified with a   474
hypergeometric test from the GOstats package (v2.54.0) [21] and the p-values were corrected for   475
multiple-hypothesis testing employing the Bonferroni method [42]. GO terms with a significant   476
adjusted p-value of less than 0.05 were reduced to representative non-redundant terms with the use   477
of REVIGO [73].   478

Host signaling pathway enrichment   479

The DEG lists produced by DESeq2 with an absolute Log2FC > 1 and FDR-adjusted p-value < 0.05   480
were used as input to the SPIA algorithm to identify significantly affected pathways from the R   481
graphite library [65, 75]. Pathways with Bonferroni-adjusted p-values less than 0.05 were included in   482
downstream analyses. The significant results for all comparisons from publicly available data from   483
KEGG, Reactome, Panther, BioCarta, and NCI were then compiled to facilitate downstream   484
comparison. Hypergeometric pathway enrichments were performed using the Database for   485
Annotation, Visualization and Integrated Discovery (DAVID, v6.8) [30].   486

Integration of transcriptomic analysis with the human metabolic network   487

To detect increased and decreased fluxes of metabolites we projected the transcriptomic data onto   488
the human reconstructed metabolic network Recon (v2.04) [77]. First, we ran EBSeq [40] on the   489
gene count matrix generated in the previous steps. Then, we used the output of EBSeq containing   490
posterior probabilities of a gene being DE (PPDE) and the Log2FC as input to the Moomin   491
method [63] using default parameters. Finally, we enumerated 250 topological solutions in order to   492
construct a consensus solution for each of the datasets tested.   493

Isoform Analysis   494

Using transcript quantification data from StringTie as input, we identified isoform switching events   495
and their predicted functional consequences with the IsoformSwitchAnalyzeR R package   496
(v1.11.3) [81]. In summary, we filtered for isoforms that experienced $|\geq 30\%|$ switch in usage for   497
each gene and were corrected for false discovery rate (FDR) with a q-value < 0.05. Following   498
filtering for significant isoforms, we externally predicted their coding capabilities, protein structure   499

stability, peptide signaling, and shifts in protein domain usage using The Coding-Potential ₅₀₀
Assessment Tool (CPAT) [82], IUPred2 [18], SignalP [2] and Pfam tools respectively [19]. These ₅₀₁
external results were imported back into IsoformSwitchAnalyzeR and used to further identify ₅₀₂
alternative splicing events and functional consequences as well as visualize the overall and individual ₅₀₃
effects of isoform switching data. Specifically, to calculate differential analysis between samples, ₅₀₄
isoform expression and usage were measured by the isoform fraction (IF) value, which quantifies the ₅₀₅
individual isoform expression level relative to the parent gene's expression level: ₅₀₆

$$IF = \frac{isoform\ expression}{gene\ expression}$$

By proxy, the difference in isoform usage between samples (dIF) measures the effect size between ₅₀₇
conditions and is calculated as follows: ₅₀₈

$$dIF = IF2\text{--}IF1$$

dIF was measured on a scale of 0 to 1, with 0 = no (0%) change in usage between conditions and ₅₀₉
1 = complete (100%) change in usage. The sum of dIF values for all isoforms associated with one ₅₁₀
gene is equal to 1. Gene expression data was imported from the aforementioned DESeq2 results. ₅₁₁
The top 30 isoforms per dataset comparison were identified by ranking isoforms by gene switch ₅₁₂
q-value, i.e. the significance of the summation of all isoform switching events per gene between mock ₅₁₃
and infected conditions. ₅₁₄

Transposable Element Analysis ₅₁₅

TE expression was quantified using the TEcount function from the TEtools software [41]. TEcount ₅₁₆
detects reads aligned against copies of each TE family annotated from the reference genome. DETEs ₅₁₇
in infected vs mock conditions were detected using DEseq2 (v1.26.0) [47] with a matrix of counts for ₅₁₈
genes and TE families as input. Functional enrichment of nearby genes (upstream 5kb and ₅₁₉
downstream 1kb of each TE copy within the human genome) was calculated with GREAT [51] using ₅₂₀
options "genome background" and "basal + extension". We only selected occurrences statistically ₅₂₁
significant by region binomial test. ₅₂₂

Identification of putative binding sites for human RBPs on the ₅₂₃
SARS-CoV-2 genome ₅₂₄

The list of RBPs downloaded from ATtRACT was filtered to human RBPs. The list was further ₅₂₅
filtered to retain PWMs obtained through competitive experiments and drop PWMs with very high ₅₂₆
entropy. This left 205 PWMs for 102 human RBPs. The SARS-CoV-2 reference genome sequence ₅₂₇
was scanned with the remaining PWMs using the TFBSTools R package (v1.20.0) [74]. A minimum ₅₂₈
score threshold of 90% was used to identify putative RBP binding sites. ₅₂₉

Enrichment analysis for putative RBP binding sites ₅₃₀

The sequences of the SARS-CoV-2 genome, 5'UTR, 3'UTR, intergenic regions and negative sense ₅₃₁
molecule were each scrambled 1,000 times. Each of the 1,000 scrambled sequences was scanned for ₅₃₂
RBP binding sites as described above. The number of binding sites for each RBP was counted, and ₅₃₃
the mean and standard deviation of the number of sites was calculated for each RBP, per region, ₅₃₄
across all 1,000 simulations. A minimum FDR-adjusted p-value of 0.01 was taken as the cutoff for ₅₃₅
enrichment. This analysis was repeated with the reference genomes of SARS-CoV and RaTG13 . ₅₃₆

Conservation analysis for putative RBP binding sites 537

The multiple sequence alignment of 27,592 SARS-CoV-2 genome sequences was downloaded from 538
GISAID [69] on May 19th 2020. For each putative RBP binding site, we selected the corresponding 539
columns of the multiple sequence alignment. We then counted the number of genomes in which the 540
sequence was identical to that of the reference genome. 541

Viral genotype-phenotype correlation 542

All complete SARS-CoV-2 genomes from GISAID, together with the GenBank reference sequence, 543
were aligned with MAFFT (v7.464) within a high-performance computing environment using 1 544
thread and the –nomemsave parameter [55]. Sequences responsible for introducing excessive gaps in 545
this initial alignment were then identified and removed, leaving 1,511 sequences that were then used 546
to generate a new multiple sequence alignment. The disease severity metadata for these sequences 547
was then normalized into four categories: severe, moderate, mild, and unknown. Next, the sequence 548
data and associated metadata were used as input to the meta-CATS algorithm to identify aligned 549
positions that contained significant differences in their base distribution between 2 or more disease 550
severities [61]. The Benjamini-Hochberg multiple hypothesis correction was then applied to all 551
positions [7]. The top 50 most significant positions were then evaluated against the annotated 552
protein regions of the reference genome to determine their effect on amino acid sequence. 553

Code availability 554

Code for these analyses is available at https://github.com/vaguiarpulido/covid19-research. 555

Supporting Information 556

Supplementary Figure 1. Isoform Analysis. 557
Supplementary File 1. Zipped file containing Complete DEG tables. 558
Supplementary File 2. Zipped file containing GO for each dataset. 559
Supplementary File 3. TE family count/differential expression. 560
Supplementary File 4. GREAT Analysis (complete and per family). 561
Supplementary Table 1. Merged tables (Specific genes in SARS-CoV-2) 562
Supplementary Table 2. Supporting information for Figure 2, consisting of functional 563
enrichment specific to SARS-CoV-2. 564
Supplementary Table 3. Pathway enrichment for each dataset (SPIA and DAVID merged into 565
one file). 566
Supplementary Table 4. Metabolic fluxes predicted for each dataset using Moomin. 567
Supplementary Table 5. Isoform analysis. 568
Supplementary Table 6. Putative binding sites for human RBPs on the SARS-CoV-2 genome. 569
Supplementary Table 7. Enrichment of binding motifs for human RBPs on the SARS-CoV-2 570
genome. 571
Supplementary Table 8. Conservation of binding motifs for human RBPs across genome 572
sequences of SARS-CoV-2 isolates. 573
Supplementary Table 9. Biological evidence associated with putative SARS-CoV-2 interacting 574
human RBPs. 575
Supplementary Table 10. Enrichment of binding motifs for human RBPs on the SARS-CoV 576
genome 577
Supplementary Table 11. Enrichment of binding motifs for human RBPs on the RaTG13 genome 578

Funding   579

The authors received no specific funding to support this work.   580

Acknowledgments   581

We would like to thank the Virtual BioHackathon on COVID-19 that took place during April 2020   582
(https://github.com/virtual-biohackathons/covid-19-bh20) for fostering an environment that   583
triggered this collaboration and in particular the Gene Expression group for the fruitful discussions.   584
This work was performed using the computing facilities of the CC/PRABI/LBBE in France and the   585
France Génomique e-infrastructure (ANR-10-INBS-09-08), the HPC facilities of the University of   586
Luxembourg [80]. We would also like to thank Slack for providing us with free access to the   587
professional version of the platform.   588
We would also like to thank Slack for providing us with free access to the professional version of the   589
platform.   590

Conflicts of Interest   591

A.L. is an employee of NVIDIA Corporation.   592

References

1. P. Ahlquist, A. O. Noueiry, W.-M. Lee, D. B. Kushner, and B. T. Dye. Host factors in positive-strand RNA virus genome replication. *J. Virol.*, 77(15):8181–8186, Aug. 2003.

2. J. J. Almagro Armenteros, K. D. Tsirigos, C. K. Sønderby, T. N. Petersen, O. Winther, S. Brunak, G. von Heijne, and H. Nielsen. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat. Biotechnol.*, 37(4):420–423, Apr. 2019.

3. S. Anders and W. Huber. Differential expression analysis for sequence count data. *Genome Biol.*, 11(10):R106, Oct. 2010.

4. K. G. Andersen, A. Rambaut, W. I. Lipkin, E. C. Holmes, and R. F. Garry. The proximal origin of SARS-CoV-2. *Nat. Med.*, 26(4):450–452, Apr. 2020.

5. M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, and G. Sherlock. Gene ontology: tool for the unification of biology. the gene ontology consortium. *Nat. Genet.*, 25(1):25–29, May 2000.

6. M. Becerra-Flores and T. Cardozo. SARS-CoV-2 viral spike G614 mutation exhibits higher case fatality rate. *Int. J. Clin. Pract.*, page e13525, May 2020.

7. Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: A practical and powerful approach to multiple testing, 1995.

8. D. Benvenuto, S. Angeletti, M. Giovanetti, M. Bianchi, S. Pascarella, R. Cauda, M. Ciccozzi, and A. Cassone. Evolutionary analysis of SARS-CoV-2: how mutation of Non-Structural protein 6 (NSP6) could affect viral autophagy. *J. Infect.*, 81(1):e24–e27, July 2020.

9. K. Bertram, D. E. Agafonov, W.-T. Liu, O. Dybkov, C. L. Will, K. Hartmuth, H. Urlaub, B. Kastner, H. Stark, and R. Lührmann. Cryo-EM structure of a human spliceosome activated for step 2 of splicing. *Nature*, 542(7641):318–323, Feb. 2017.

10. D. Blanco-Melo, B. E. Nilsson-Payant, W.-C. Liu, S. Uhl, D. Hoagland, R. Møller, T. X. Jordan, K. Oishi, M. Panis, D. Sachs, T. T. Wang, R. E. Schwartz, J. K. Lim, R. A. Albrecht, and B. R. tenOever. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell*, 181(5):1036–1045.e9, May 2020.

11. Y. Cárdenas-Conejo, A. Liñan-Rico, D. A. García-Rodríguez, S. Centeno-Leija, and H. Serrano-Posada. An exclusive 42 amino acid signature in pp1ab protein provides insights into the evolutive history of the 2019 novel human-pathogenic coronavirus (SARS-CoV-2). *J. Med. Virol.*, 92(6):688–692, June 2020.

12. S. J. Carter, R. S. Tattersall, and A. V. Ramanan. Macrophage activation syndrome in adults: recent advances in pathophysiology, diagnosis and treatment. *Rheumatology*, 58(1):5–17, Jan. 2019.

13. D. Chasman, K. B. Walters, T. J. S. Lopes, A. J. Eisfeld, Y. Kawaoka, and S. Roy. Integrating transcriptomic and proteomic data using predictive regulatory network models of host response to pathogens. *PLoS Comput. Biol.*, 12(7):e1005013, July 2016.

14. E. B. Chuong, N. C. Elde, and C. Feschotte. Regulatory evolution of innate immunity through co-option of endogenous retroviruses. *Science*, 351(6277):1083–1087, Mar. 2016.

15. E. B. Chuong, N. C. Elde, and C. Feschotte. Regulatory activities of transposable elements: from conflicts to benefits. *Nat. Rev. Genet.*, 18(2):71–86, Feb. 2017.

16. Ö. Deniz, M. Ahmed, C. D. Todd, A. Rio-Machin, M. A. Dawson, and M. R. Branco. Endogenous retroviruses are a source of enhancers with oncogenic potential in acute myeloid leukaemia. *Nat. Commun.*, 11(1):3506, July 2020.

17. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T. R. Gingeras. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1):15–21, Jan. 2013.

18. Z. Dosztányi, V. Csizmok, P. Tompa, and I. Simon. IUPred: web server for the prediction of intrinsically unstructured regions of proteins based on estimated energy content. *Bioinformatics*, 21(16):3433–3434, Aug. 2005.

19. S. El-Gebali, J. Mistry, A. Bateman, S. R. Eddy, A. Luciani, S. C. Potter, M. Qureshi, L. J. Richardson, G. A. Salazar, A. Smart, E. L. L. Sonnhammer, L. Hirsh, L. Paladin, D. Piovesan, S. C. E. Tosatto, and R. D. Finn. The pfam protein families database in 2019. *Nucleic Acids Res.*, 47(D1):D427–D432, Jan. 2019.

20. P. Ewels, M. Magnusson, S. Lundin, and M. Käller. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 32(19):3047–3048, Oct. 2016.

21. S. Falcon and R. Gentleman. Using GOstats to test gene lists for GO term association. *Bioinformatics*, 23(2):257–258, Jan. 2007.

22. T. S. Fung and D. X. Liu. Human coronavirus: Host-Pathogen interaction. *Annu. Rev. Microbiol.*, 73:529–557, Sept. 2019.

23. G. Giudice, F. Sánchez-Cabo, C. Torroja, and E. Lara-Pezzi. ATtRACT-a database of RNA-binding proteins and associated motifs. *Database*, 2016, Apr. 2016.

24. D. E. Gordon, G. M. Jang, M. Bouhaddou, J. Xu, K. Obernier, M. J. O'Meara, J. Z. Guo, D. L. Swaney, T. A. Tummino, R. Hüttenhain, R. M. Kaake, A. L. Richards, B. Tutuncuoglu, H. Foussard, J. Batra, K. Haas, M. Modak, M. Kim, P. Haas, B. J. Polacco, H. Braberg, J. M. Fabius, M. Eckhardt, M. Soucheray, M. J. Bennett, M. Cakir, M. J. McGregor, Q. Li, Z. Z. C.

Naing, Y. Zhou, S. Peng, I. T. Kirby, J. E. Melnyk, J. S. Chorba, K. Lou, S. A. Dai, W. Shen, Y. Shi, Z. Zhang, I. Barrio-Hernandez, D. Memon, C. Hernandez-Armenta, C. J. P. Mathy, T. Perica, K. B. Pilla, S. J. Ganesan, D. J. Saltzberg, R. Ramachandran, X. Liu, S. B. Rosenthal, L. Calviello, S. Venkataramanan, Y. Lin, S. A. Wankowicz, M. Bohn, R. Trenker, J. M. Young, D. Cavero, J. Hiatt, T. Roth, U. Rathore, A. Subramanian, J. Noack, M. Hubert, F. Roesch, T. Vallet, B. Meyer, K. M. White, L. Miorin, D. Agard, M. Emerman, D. Ruggero, A. García-Sastre, N. Jura, M. von Zastrow, J. Taunton, O. Schwartz, M. Vignuzzi, C. d'Enfert, S. Mukherjee, M. Jacobson, H. S. Malik, D. G. Fujimori, T. Ideker, C. S. Craik, S. Floor, J. S. Fraser, J. Gross, A. Sali, T. Kortemme, P. Beltrao, K. Shokat, B. K. Shoichet, and N. J. Krogan. A SARS-CoV-2-Human Protein-Protein interaction map reveals drug targets and potential Drug-Repurposing. *bioRxiv*, Mar. 2020.

25. GTEx Consortium. Human genomics. the Genotype-Tissue expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science*, 348(6235):648–660, May 2015.

26. W. Guo, J. Wei, X. Zhong, R. Zang, H. Lian, M.-M. Hu, S. Li, H.-B. Shu, and Q. Yang. SNX8 modulates the innate immune response to RNA viruses by regulating the aggregation of VISA. *Cell. Mol. Immunol.*, Sept. 2019.

27. M. Hinz, M. Broemer, S. Ç. Arslan, A. Otto, E.-C. Mueller, R. Dettmer, and C. Scheidereit. Signal responsiveness of IκB kinases is determined by cdc37-assisted transient interaction with hsp90. *J. Biol. Chem.*, 282(44):32311–32319, Nov. 2007.

28. S. Horváth and K. Mirnics. Immune system disturbances in schizophrenia. *Biol. Psychiatry*, 75(4):316–323, Feb. 2014.

29. C. Huang, Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu, X. Gu, Z. Cheng, T. Yu, J. Xia, Y. Wei, W. Wu, X. Xie, W. Yin, H. Li, M. Liu, Y. Xiao, H. Gao, L. Guo, J. Xie, G. Wang, R. Jiang, Z. Gao, Q. Jin, J. Wang, and B. Cao. Clinical features of patients infected with 2019 novel coronavirus in wuhan, china. *Lancet*, 395(10223):497–506, Feb. 2020.

30. D. W. Huang, B. T. Sherman, and R. A. Lempicki. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.*, 4(1):44–57, 2009.

31. P. Huang and M. M. Lai. Heterogeneous nuclear ribonucleoprotein a1 binds to the 3'-untranslated region and mediates potential 5'-3'-end cross talks of mouse hepatitis virus RNA. *J. Virol.*, 75(11):5009–5017, June 2001.

32. J. Ito, R. Sugimoto, H. Nakaoka, S. Yamada, T. Kimura, T. Hayano, and I. Inoue. Systematic identification and characterization of regulatory elements derived from human endogenous retroviruses. *PLoS Genet.*, 13(7):e1006883, July 2017.

33. M. S. Jurica, L. J. Licklider, S. R. Gygi, N. Grigorieff, and M. J. Moore. Purification and characterization of native spliceosomes suitable for three-dimensional structural analysis. *RNA*, 8(4):426–439, Apr. 2002.

34. N. Kadowaki and Y.-J. Liu. Natural type I interferon-producing cells as a link between innate and adaptive immunity. *Hum. Immunol.*, 63(12):1126–1132, Dec. 2002.

35. R. J. Khan, R. K. Jha, G. M. Amera, M. Jain, E. Singh, A. Pathak, R. P. Singh, J. Muthukumaran, and A. K. Singh. Targeting SARS-CoV-2: a systematic drug repurposing approach to identify promising inhibitors against 3c-like proteinase and 2'-o-ribose methyltransferase. *J. Biomol. Struct. Dyn.*, pages 1–14, Apr. 2020.

36. D. Kim, J.-Y. Lee, J.-S. Yang, J. W. Kim, V. N. Kim, and H. Chang. The architecture of SARS-CoV-2 transcriptome. *Cell*, 181(4):914–921.e10, May 2020.

37. K. K. Kojima. Human transposable elements in repbase: genomic footprints from fish to humans. *Mob. DNA*, 9:2, Jan. 2018.

38. B. Korber, W. M. Fischer, S. Gnanakaran, H. Yoon, J. Theiler, W. Abfalterer, N. Hengartner, E. E. Giorgi, T. Bhattacharya, B. Foley, K. M. Hastie, M. D. Parker, D. G. Partridge, C. M. Evans, T. M. Freeman, T. I. de Silva, C. McDanal, L. G. Perez, H. Tang, A. Moon-Walker, S. P. Whelan, C. C. LaBranche, E. O. Saphire, D. C. Montefiori, A. Angyal, R. L. Brown, L. Carrilero, L. R. Green, D. C. Groves, K. J. Johnson, A. J. Keeley, B. B. Lindsey, P. J. Parsons, M. Raza, S. Rowland-Jones, N. Smith, R. M. Tucker, D. Wang, and M. D. Wyles. Tracking changes in SARS-CoV-2 spike: Evidence that D614G increases infectivity of the COVID-19 virus. *Cell*, July 2020.

39. T. T.-Y. Lam, N. Jia, Y.-W. Zhang, M. H.-H. Shum, J.-F. Jiang, H.-C. Zhu, Y.-G. Tong, Y.-X. Shi, X.-B. Ni, Y.-S. Liao, W.-J. Li, B.-G. Jiang, W. Wei, T.-T. Yuan, K. Zheng, X.-M. Cui, J. Li, G.-Q. Pei, X. Qiang, W. Y.-M. Cheung, L.-F. Li, F.-F. Sun, S. Qin, J.-C. Huang, G. M. Leung, E. C. Holmes, Y.-L. Hu, Y. Guan, and W.-C. Cao. Identifying SARS-CoV-2-related coronaviruses in malayan pangolins. *Nature*, 583(7815):282–285, July 2020.

40. N. Leng, J. A. Dawson, J. A. Thomson, V. Ruotti, A. I. Rissman, B. M. G. Smits, J. D. Haag, M. N. Gould, R. M. Stewart, and C. Kendziorski. EBSeq: an empirical bayes hierarchical model for inference in RNA-seq experiments. *Bioinformatics*, 29(8):1035–1043, Apr. 2013.

41. E. Lerat, M. Fablet, L. Modolo, H. Lopez-Maestre, and C. Vieira. TEtools facilitates big data expression analysis of transposable elements and reveals an antagonism between their activity and that of piRNA genes. *Nucleic Acids Res.*, 45(4):e17, Feb. 2017.

42. K. Lesack and C. Naugler. An open-source software program for performing bonferroni and related corrections for multiple comparisons. *J. Pathol. Inform.*, 2:52, Dec. 2011.

43. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, and R. Durbin. The sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16):2078–2079, Aug. 2009.

44. H. P. Li, X. Zhang, R. Duncan, L. Comai, and M. M. Lai. Heterogeneous nuclear ribonucleoprotein A1 binds to the transcription-regulatory region of mouse hepatitis virus RNA. *Proc. Natl. Acad. Sci. U. S. A.*, 94(18):9544–9549, Sept. 1997.

45. Z. Li and P. D. Nagy. Diverse roles of host RNA binding proteins in RNA virus replication. *RNA Biol.*, 8(2):305–315, Mar. 2011.

46. M. Liao, Y. Liu, J. Yuan, Y. Wen, G. Xu, J. Zhao, L. Cheng, J. Li, X. Wang, F. Wang, L. Liu, I. Amit, S. Zhang, and Z. Zhang. Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19, 2020.

47. M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.*, 15(12):550, 2014.

48. M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.*, 15(12):550, 2014.

49. C. Lucas, P. Wong, J. Klein, T. B. R. Castro, J. Silva, M. Sundaram, M. K. Ellingson, T. Mao, J. E. Oh, B. Israelow, T. Takahashi, M. Tokuyama, P. Lu, A. Venkataraman, A. Park, S. Mohanty, H. Wang, A. L. Wyllie, C. B. F. Vogels, R. Earnest, S. Lapidus, I. M. Ott, A. J. Moore, M. C. Muenker, J. B. Fournier, M. Campbell, C. D. Odio, A. Casanovas-Massana, A. Obaid, A. Lu-Culligan, A. Nelson, A. Brito, A. Nunez, A. Martin, A. Watkins, B. Geng, C. Kalinich, C. Harden, C. Todeasa, C. Jensen, D. Kim, D. McDonald, D. Shepard, E. Courchaine, E. B. White, E. Song, E. Silva, E. Kudo, G. DeIuliis, H. Rahming, H.-J. Park,

I. Matos, J. Nouws, J. Valdez, J. Fauver, J. Lim, K.-A. Rose, K. Anastasio, K. Brower, L. Glick, L. Sharma, L. Sewanan, L. Knaggs, M. Minasyan, M. Batsu, M. Petrone, M. Kuang, M. Nakahata, M. Linehan, M. H. Askenase, M. Simonov, M. Smolgovsky, N. Sonnert, N. Naushad, P. Vijayakumar, R. Martinello, R. Datta, R. Handoko, S. Bermejo, S. Prophet, S. Bickerton, S. Velazquez, T. Alpert, T. Rice, W. Khoury-Hanold, X. Peng, Y. Yang, Y. Cao, Y. Strong, R. Herbst, A. C. Shaw, R. Medzhitov, W. L. Schulz, N. D. Grubaugh, C. Dela Cruz, S. Farhadian, A. I. Ko, S. B. Omer, A. Iwasaki, and Y. I. Team. Longitudinal analyses reveal immunological misfiring in severe covid-19. *Nature*, 2020.

50. M. G. Macchietto, R. A. Langlois, and S. S. Shen. Virus-induced transposable element expression up-regulation in human and mouse host cells. *Life Science Alliance*, 3(2), Feb. 2020.

51. C. Y. McLean, D. Bristor, M. Hiller, S. L. Clarke, B. T. Schaar, C. B. Lowe, A. M. Wenger, and G. Bejerano. GREAT improves functional interpretation of cis-regulatory regions. *Nat. Biotechnol.*, 28(5):495–501, May 2010.

52. H. M. Mehta, M. Malandra, and S. J. Corey. G-CSF and GM-CSF in neutropenia. *J. Immunol.*, 195(4):1341–1349, Aug. 2015.

53. P. Mehta, D. F. McAuley, M. Brown, E. Sanchez, R. S. Tattersall, J. J. Manson, and HLH Across Speciality Collaboration, UK. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*, 395(10229):1033–1034, Mar. 2020.

54. N. Muralidharan, R. Sakthivel, D. Velmurugan, and M. M. Gromiha. Computational studies of drug repurposing and synergism of lopinavir, oseltamivir and ritonavir binding with SARS-CoV-2 protease against COVID-19. *J. Biomol. Struct. Dyn.*, pages 1–6, Apr. 2020.

55. T. Nakamura, K. D. Yamada, K. Tomii, and K. Katoh. Parallelization of MAFFT for large-scale multiple sequence alignments. *Bioinformatics*, 34(14):2490–2492, July 2018.

56. T. Nitto and K. Onodera. Linkage between coenzyme a metabolism and inflammation: roles of pantetheinase. *J. Pharmacol. Sci.*, 123(1):1–8, Sept. 2013.

57. J. K. Pace, 2nd and C. Feschotte. The evolutionary history of human DNA transposons: evidence for intense activity in the primate lineage. *Genome Res.*, 17(4):422–432, Apr. 2007.

58. M. Pachetti, B. Marini, F. Benedetti, F. Giudici, E. Mauro, P. Storici, C. Masciovecchio, S. Angeletti, M. Ciccozzi, R. C. Gallo, D. Zella, and R. Ippodrino. Emerging SARS-CoV-2 mutation hot spots include a novel RNA-dependent-RNA polymerase variant. *J. Transl. Med.*, 18(1):179, Apr. 2020.

59. C. Perez, C. McKinney, U. Chulunbaatar, and I. Mohr. Translational control of the abundance of cytoplasmic poly(a) binding protein in human cytomegalovirus-infected cells. *J. Virol.*, 85(1):156–164, Jan. 2011.

60. M. Pertea, G. M. Pertea, C. M. Antonescu, T.-C. Chang, J. T. Mendell, and S. L. Salzberg. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat. Biotechnol.*, 33(3):290–295, Mar. 2015.

61. B. E. Pickett, M. Liu, E. L. Sadat, R. B. Squires, J. M. Noronha, S. He, W. Jen, S. Zaremba, Z. Gu, L. Zhou, C. N. Larsen, I. Bosch, L. Gehrke, M. McGee, E. B. Klem, and R. H. Scheuermann. Metadata-driven comparative analysis tool for sequences (meta-CATS): an automated process for identifying significant sequence variations that correlate with virus attributes. *Virology*, 447(1-2):45–51, Dec. 2013.

62. C. Polacek, P. Friebe, and E. Harris. Poly(A)-binding protein binds to the non-polyadenylated 3' untranslated region of dengue virus and modulates translation efficiency. *J. Gen. Virol.*, 90(Pt 3):687–692, Mar. 2009.

63. T. Pusa, M. G. Ferrarini, R. Andrade, A. Mary, A. Marchetti-Spaccamela, L. Stougie, and M.-F. Sagot. MOOMIN - mathematical exploration of 'omics data on a MetabolIc network. *Bioinformatics*, 36(2):514–523, Jan. 2020.

64. P. A. Reyfman, J. M. Walter, N. Joshi, K. R. Anekalla, A. C. McQuattie-Pimentel, S. Chiu, R. Fernandez, M. Akbarpour, C.-I. Chen, Z. Ren, R. Verma, H. Abdala-Valencia, K. Nam, M. Chi, S. Han, F. J. Gonzalez-Gonzalez, S. Soberanes, S. Watanabe, K. J. N. Williams, A. S. Flozak, T. T. Nicholson, V. K. Morgan, D. R. Winter, M. Hinchcliff, C. L. Hrusch, R. D. Guzy, C. A. Bonham, A. I. Sperling, R. Bag, R. B. Hamanaka, G. M. Mutlu, A. V. Yeldandi, S. A. Marshall, A. Shilatifard, L. A. N. Amaral, H. Perlman, J. I. Sznajder, A. C. Argento, C. T. Gillespie, J. Dematte, M. Jain, B. D. Singer, K. M. Ridge, A. P. Lam, A. Bharat, S. M. Bhorade, C. J. Gottardi, G. R. S. Budinger, and A. V. Misharin. Single-Cell transcriptomic analysis of human lung provides insights into the pathobiology of pulmonary fibrosis. *Am. J. Respir. Crit. Care Med.*, 199(12):1517–1536, June 2019.

65. G. Sales, E. Calura, D. Cavalieri, and C. Romualdi. graphite - a bioconductor package to convert pathway topology to gene network. *BMC Bioinformatics*, 13:20, Jan. 2012.

66. C. D. Schmid and P. Bucher. MER41 repeat sequences contain inducible STAT1 binding sites. *PLoS One*, 5(7):e11425, July 2010.

67. N. Schmidt, C. A. Lareau, H. Keshishian, R. Melanson, M. Zimmer, L. Kirschner, J. Ade, S. Werner, N. Caliskan, E. S. Lander, J. Vogel, S. A. Carr, J. Bodem, and M. Munschauer. A direct RNA-protein interaction atlas of the SARS-CoV-2 RNA in infected human cells.

68. S. T. Shi, P. Huang, H. P. Li, and M. M. Lai. Heterogeneous nuclear ribonucleoprotein A1 regulates RNA synthesis of a cytoplasmic virus. *EMBO J.*, 19(17):4701–4711, Sept. 2000.

69. Y. Shu and J. McCauley. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill.*, 22(13), Mar. 2017.

70. A. F. Smit. Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr. Opin. Genet. Dev.*, 9(6):657–663, Dec. 1999.

71. R. W. P. Smith, R. C. Anderson, O. Larralde, J. W. S. Smith, B. Gorgoni, W. A. Richardson, P. Malik, S. V. Graham, and N. K. Gray. Viral and cellular mRNA-specific activators harness PABP and eIF4G to promote translation initiation downstream of cap binding. *Proc. Natl. Acad. Sci. U. S. A.*, 114(24):6310–6315, June 2017.

72. R. W. P. Smith and N. K. Gray. Poly(A)-binding protein (PABP): a common viral target. *Biochem. J*, 426(1):1–12, Feb. 2010.

73. F. Supek, M. Bošnjak, N. Škunca, and T. Šmuc. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One*, 6(7):e21800, July 2011.

74. G. Tan and B. Lenhard. TFBSTools: an R/bioconductor package for transcription factor binding site analysis. *Bioinformatics*, 32(10):1555–1556, 01 2016.

75. A. L. Tarca, S. Draghici, P. Khatri, S. S. Hassan, P. Mittal, J.-S. Kim, C. J. Kim, J. P. Kusanovic, and R. Romero. A novel signaling pathway impact analysis. *Bioinformatics*, 25(1):75–82, Jan. 2009.

76. The Gene Ontology Consortium and The Gene Ontology Consortium. The gene ontology resource: 20 years and still GOing strong, 2019.

77. I. Thiele, N. Swainston, R. M. T. Fleming, A. Hoppe, S. Sahoo, M. K. Aurich, H. Haraldsdottir, M. L. Mo, O. Rolfsson, M. D. Stobbe, S. G. Thorleifsson, R. Agren, C. Bölling, S. Bordel, A. K. Chavali, P. Dobson, W. B. Dunn, L. Endler, D. Hala, M. Hucka, D. Hull, D. Jameson, N. Jamshidi, J. J. Jonsson, N. Juty, S. Keating, I. Nookaew, N. Le Novère, N. Malys, A. Mazein, J. A. Papin, N. D. Price, E. Selkov, Sr, M. I. Sigurdsson, E. Simeonidis, N. Sonnenschein, K. Smallbone, A. Sorokin, J. H. G. M. van Beek, D. Weichart, I. Goryanin, J. Nielsen, H. V. Westerhoff, D. B. Kell, P. Mendes, and B. Ø. Palsson. A community-driven global reconstruction of human metabolism. *Nat. Biotechnol.*, 31(5):419–425, May 2013.

78. M. Trizzino, A. Kapusta, and C. D. Brown. Transposable elements generate regulatory novelty in a tissue-specific fashion. *BMC Genomics*, 19(1):468, June 2018.

79. K. W. Tsang, P. L. Ho, G. C. Ooi, W. K. Yee, T. Wang, M. Chan-Yeung, W. K. Lam, W. H. Seto, L. Y. Yam, T. M. Cheung, P. C. Wong, B. Lam, M. S. Ip, J. Chan, K. Y. Yuen, and K. N. Lai. A cluster of cases of severe acute respiratory syndrome in hong kong. *N. Engl. J. Med.*, 348(20):1977–1985, May 2003.

80. S. Varrette, P. Bouvry, H. Cartiaux, and F. Georgatos. Management of an academic hpc cluster: The ul experience. In *Proc. of the 2014 Intl. Conf. on High Performance Computing & Simulation (HPCS 2014)*, pages 959–967, Bologna, Italy, July 2014. IEEE.

81. K. Vitting-Seerup and A. Sandelin. IsoformSwitchAnalyzeR: analysis of changes in genome-wide patterns of alternative splicing and its functional consequences. *Bioinformatics*, 35(21):4469–4471, Nov. 2019.

82. L. Wang, H. J. Park, S. Dasari, S. Wang, J.-P. Kocher, and W. Li. CPAT: Coding-Potential assessment tool using an alignment-free logistic regression model. *Nucleic Acids Res.*, 41(6):e74, Apr. 2013.

83. W. Wen, W. Su, H. Tang, W. Le, X. Zhang, Y. Zheng, X. Liu, L. Xie, J. Li, J. Ye, L. Dong, X. Cui, Y. Miao, D. Wang, J. Dong, C. Xiao, W. Chen, and H. Wang. Immune cell profiling of COVID-19 patients in the recovery stage by single-cell sequencing. *Cell Discov*, 6:31, May 2020.

84. D. Wu and X. O. Yang. TH17 responses in cytokine storm of COVID-19: An emerging target of JAK2 inhibitor fedratinib. *J. Microbiol. Immunol. Infect.*, 53(3):368–370, June 2020.

85. R. Yan, Y. Zhang, Y. Li, L. Xia, Y. Guo, and Q. Zhou. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science*, 367(6485):1444–1448, Mar. 2020.

86. C. Yin. Genotyping coronavirus SARS-CoV-2: methods and implications. *Genomics*, Apr. 2020.

87. A. M. Zaki, S. van Boheemen, T. M. Bestebroer, A. D. M. E. Osterhaus, and R. A. M. Fouchier. Isolation of a novel coronavirus from a man with pneumonia in saudi arabia. *N. Engl. J. Med.*, 367(19):1814–1820, Nov. 2012.

88. P. Zhou, X.-L. Yang, X.-G. Wang, B. Hu, L. Zhang, W. Zhang, H.-R. Si, Y. Zhu, B. Li, C.-L. Huang, H.-D. Chen, J. Chen, Y. Luo, H. Guo, R.-D. Jiang, M.-Q. Liu, Y. Chen, X.-R. Shen, X. Wang, X.-S. Zheng, K. Zhao, Q.-J. Chen, F. Deng, L.-L. Liu, B. Yan, F.-X. Zhan, Y.-Y. Wang, G.-F. Xiao, and Z.-L. Shi. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, 579(7798):270–273, Mar. 2020.

89. Y. Zhou, Y. Hou, J. Shen, Y. Huang, W. Martin, and F. Cheng. Network-based drug repurposing for novel coronavirus 2019-nCoV/SARS-CoV-2. *Cell Discov*, 6:14, Mar. 2020.

90. Y. Zhou and Y. Zhu. Important role of the IL-32 inflammatory network in the host response against viral infection. *Viruses*, 7(6):3116–3129, June 2015.