# Nonlinear effects of intrinsic dynamics on temporal encoding in a model of avian auditory cortex

**Authors:**  Christof Fehrman[1], Tyler D Robbins[2], and C Daniel Meliza[1,3,*]

**Affiliations:**  [1] Psychology Department, [2], Cognitive Science Program, [3] Neuroscience Graduate Program, University of Virginia, Charlottesville VA 22904, USA

[*] Corresponding author. Email: cdm8j@virginia.edu

## Abstract

Neurons exhibit diverse intrinsic dynamics, which govern how they integrate synaptic inputs to produce spikes. Intrinsic dynamics are often plastic during development and learning, but the effects of these changes on stimulus encoding properties are not well known. To examine this relationship, we simulated auditory responses to zebra finch song using a linear-dynamical cascade model, which combines a linear spectrotemporal receptive field with a dynamical, conductance-based neuron model, then used generalized linear models to estimate encoding properties from the resulting spike trains. We focused on the effects of a low-threshold potassium current ($K_{LT}$) that is present in a subset of cells in the zebra finch caudal mesopallium and is affected by early auditory experience. We found that $K_{LT}$ affects both spike adaptation and the temporal filtering properties of the receptive field. Interestingly, the direction of the effects depended on the temporal modulation tuning of the linear (input) stage of the cascade model, indicating a strongly nonlinear relationship. These results suggest that small changes in intrinsic dynamics in tandem with differences in synaptic connectivity can have dramatic effects on the tuning of auditory neurons.

## Introduction                                                                                    15

Neurons have diverse, nonlinear dynamics. Many brain regions contain multiple kinds of neurons     16
with different spike waveforms and spiking patterns (Bal and Oertel, 2001; Sivaramakrishnan and     17
Oliver, 2001; Ascoli et al., 2008), and there is substantial variation even within well-defined cell     18
types (Toledo-Rodriguez et al., 2004; Schulz et al., 2006; Bomkamp et al., 2019). Intrinsic dynamics     19
can be modified by activity and experience (Ross et al., 2017; Daou and Margoliash, 2020; Chen     20
and Meliza, 2020), which may be an important mechanism for learning (Titley et al., 2017). This     21
physiological diversity has been known for many decades (Llinás, 1988) and can be modeled on a     22
detailed, biophysically realistic level (Padmanabhan and Urban, 2010; Tripathy et al., 2015), but     23
our understanding of how intrinsic dynamics affect neural computations in many systems has     24
remained surprisingly qualitative.                                                                 25

One reason why biophysical models are difficult to apply to processes at the algorithmic and     26
computational levels is their complexity. A simple, single-compartmental model that can produce     27
common physiological behaviors like bursting, adaptation, or rebound spiking, is a system of     28
around ten or more nonlinear differential equations, with fifty or more parameters (e.g., Rothman     29
and Manis, 2003; Meliza et al., 2014). These parameters correspond to specific aspects of the cell     30
biology (such as membrane capacitance or sodium channel density), which makes them easy to     31
interpret and, in some cases, possible to measure directly. However, the relationships between the     32
parameters and the observable behaviors of the neuron are highly nonlinear, making it difficult to     33
constrain them statistically. It is difficult and time-consuming to fit dynamical models to biological     34
data (Druckmann et al., 2008; Van Geit et al., 2008; Toth et al., 2011; Vavoulis et al., 2012), and     35
there is little consensus on the appropriate methods or even whether there are globally optimal     36
solutions (Prinz et al., 2004). Moreover, access to the intracellular voltage is needed, through a     37
low-resistance electrode or an optical sensor (Huys and Paninski, 2009), which greatly limits the     38
number of neurons that can be modeled within the context of a circuit, and almost always requires     39
the use of *ex vivo* preparations that cannot be presented with realistic stimuli.                   40

As a consequence, many studies of function in neural systems have emphasized phenomenological     41
models that omit most of the biophysical and dynamical features of spike generation in exchange     42

for computational tractability (Keat et al., 2001; Jolivet et al., 2004; Kobayashi et al., 2009; Izhikevich, 2003). One of the simplest examples is the generalized linear model (GLM), which represents spiking as an inhomogeneous Poisson process with a conditional intensity that depends only on a linear function of the stimulus and spiking response in the recent past (Pillow et al., 2005). In contrast to more realistic models, the GLM is a staple of statistics, with a well-defined likelihood function that is concave everywhere, guaranteeing that a global optimum can be found (Paninski et al., 2004). The GLM also has established techniques for regularization, which is necessary when stimuli have naturalistic (i.e., highly correlated) distributions (Theunissen et al., 2001; Schwartz et al., 2006).

Because of its simplicity and probabilistic formulation, a GLM can be thought of as a representation of a neuron's encoding properties; that is, an abstract view of how the cell transforms sensory stimuli into spike trains. Surprisingly, although GLMs have been successfully used to model encoding in a number of different sensory systems (Pillow et al., 2005; Calabrese et al., 2011), and there have been several studies using GLMs to predict and characterize more complex spiking models (Ostojic and Brunel, 2011; Pozzorini et al., 2015; Weber and Pillow, 2017), to our knowledge there has not been any attempt to relate the GLM to more detailed, dynamical models with realistic sensory inputs. As a result, it is difficult to predict how natural, pathological, or experience-dependent variations in voltage-gated channels are likely to affect sensory processing.

In this study, we examined the relationship between intrinsic dynamics and encoding properties in the context of auditory processing in songbirds. Encoding models, including GLMs, have been employed extensively to study this system (Theunissen et al., 2000; Sen et al., 2001; Nagel and Doupe, 2008; Woolley et al., 2009; Calabrese et al., 2011), but until recently, there have been no data on the intracellular physiology of the constituent neurons. Using whole-cell patch recordings from slices, we have found that the caudal mesopallium (CM), a cortical-level auditory area (Wang et al., 2010; Jarvis et al., 2013), has diverse, experience-dependent intrinsic dynamics (Chen and Meliza, 2018, 2020). Most of the putatively excitatory neurons fire repetitively when depolarized, but a substantial fraction only fire at stimulus onsets. This phasic firing behavior is correlated with strong outward rectification that activates at low voltages, and it can be pharmacologically converted to tonic firing by blocking low-threshold potassium currents ($K_{LT}$). The proportion of phasic neurons

changes over development, along with expression of Kv1.1, a low-threshold potassium channel, and both depend on exposure to a complex acoustic environment early in life.

The dependence of phasic firing on auditory experience suggests that intrinsic plasticity (i.e., a change in the expression or properties of voltage-gated currents, rather than synaptic currents) plays a critical role in development for songbirds, but for all the reasons noted above, the functional significance remains unclear. Here, we took a simulation-based approach to ask how changing the magnitude of low-threshold potassium currents in a dynamical model would affect encoding properties, as estimated with a GLM.

We simulated auditory responses using a linear-dynamical cascade model (Bjoring and Meliza, 2019), which combines a linear spectrotemporal receptive field (RF) with a single-compartment biophysical model (Fig. 1A). The dynamical model we used can produce phasic or tonic responses depending on a single parameter that governs the maximal conductance of a low-threshold potassium current ($g_{KLT}$) (Fig. 1B–C). As shown previously, increasing this parameter causes neurons to attenuate their responses to low-frequency components of a broadband driving current (Chen and Meliza, 2018) and to be less responsive to random synaptic-like noise (Bjoring and Meliza, 2019). Here, to quantify the effects of these changes on encoding, we simulated responses of the linear-dynamical model to zebra finch songs, convolving the stimuli with representative RFs (Woolley et al., 2009) and using the output as an external driving current to the dynamical system. We used the spike trains produced by these simulations to fit GLMs (Fig. 2) and then compared estimates for the RF and spike-history parameters to determine how $K_{LT}$ influenced how the model was representing the acoustic structure of the stimulus.

## Results

### Univariate white-noise stimulus

As a proof of principle, we began with an example using a white-noise stimulus drawn from a univariate Gaussian distribution. The absence of temporal correlations in this stimulus is ideal for obtaining unbiased estimates of the GLM parameters, allowing us to determine how intrinsic

**Fig. 1. Linear-dynamical cascade model.** (A) The linear stage of the model consists of the convolution of a stimulus with a receptive field. The output of the convolution is input as a driving current ($I_{stim}(t)$) into a biophysical model (B) that represents membrane voltage dynamics as a system of ordinary differential equations. A stimulus-independent current with a pink-noise spectral distribution ($I_{noise}(t)$) is added to produce intertrial variability. (C) The model is numerically integrated to produce a simulated voltage trace. Multiple trials are simulated by keeping $I_{stim}(t)$ the same from trial to trial, while drawing new values for $I_{noise}(t)$.

**Fig. 2. Schematic of parameter estimation for generalized linear model.** The data to be fit comprise a stimulus, which can be a univariate time series or a multivariate spectrogram (as shown here), and a spiking response. The model represents the response as an inhomogeneous Poisson process with a conditional intensity that depends on the convolution of the stimulus with a receptive field (**K**) and the convolution of the response with a spike-history filter, which was parameterized as the sum of two exponential decays representing short-term ($\alpha_1$) and long-term ($\alpha_2$) adaptation or facilitation. Not shown is a constant offset $\omega$, which governs the baseline probability of firing. These model parameters are estimated by regularized maximum likelihood.

dynamics affect encoding in a best-case scenario.                                                    98

We generated data for fitting the GLM by providing 100 s of white noise as input to two linear-      99
dynamical cascade (LDC) models that had the same RF but different dynamics. The dynamical            100
stage of the model was based on our previous work in the zebra finch caudal mesopallium (Chen        101
and Meliza, 2018; Bjoring and Meliza, 2019). The tonic model lacks $K_{LT}$ and has a higher capacitance,   102
whereas the phasic model includes $K_{LT}$ and has a lower capacitance (see Methods for parameter    103
values). These models reproduce the responses to step currents (Fig. 3A) and broadband currents      104
seen in slices. Both LDC models produced similar responses to the white noise stimulus, but the      105
phasic model tended to have narrower peaks of activity (Fig. 3B–C).                                   106

In general, parameter estimates are only interpretable to the extent that the model is a good fit to  107
the data. We checked the goodness of fit by comparing the responses of the LDC model and the         108
fitted GLM to a new white-noise stimulus. The output of the GLM was an excellent prediction of        109
the dynamical model's response (Fig. 3B–C). Indeed, the correlations between the average firing       110
rates for LDC data and GLM prediction (tonic: $r = 0.96$; phasic: $r = 0.84$) were comparable to the  111
correlations between average rates of even and odd trials in the data (tonic: 0.94; phasic: 0.90)—as  112
good as could be expected given the intrinsic variability of the data. Thus, at least for white-noise 113
stimuli, the linear spike-history filter and static nonlinearity of the GLM can closely approximate   114
the dynamical nonlinearity of a single-compartment biophysical model. This allows us to interpret     115
the GLM parameters as meaningful descriptions of the encoding properties of the more complex          116
model.                                                                                                117

The LDC and GLM both have receptive fields that are convolved with the stimulus to produce            118
a signal that modulates the probability of spiking. When a GLM is fit using data from an LDC          119
model, we expect the estimated RF to resemble the RF used to generate the data, but not exactly.      120
Indeed, differences between the input and estimated RFs will reflect the effects of the intrinsic     121
dynamics. One expected effect is from the filtering properties of the membrane. In the GLM, firing    122
probability depends on a static, exponential function of the convolved stimulus (Fig. 2). In the LDC  123
model, the output of the convolution enters as a current that contributes linearly to the derivative  124
of the membrane voltage. The capacitance and conductance of the membrane act as an additional,       125
lowpass filter, so we would expect the estimated RF to be a lowpass-filtered version of the input RF. 126

**Fig. 3. GLM estimates for exemplar tonic and phasic models with univariate white-noise stimulus.**
(A) Voltage responses of tonic and phasic models to high- and low-amplitude injected current steps (shown in bottom row). The tonic model exhibits depolarization block to strong currents but fires repetitively to weak currents, whereas the phasic model only fires a single spike to all suprathreshold current levels. (B) Top, response of the tonic dynamical model to a white-noise stimulus. The input RF is shown in D. Middle, raster plots of spike times from 10 trials with the same stimulus but varying $I_{\text{noise}}(t)$. Black ticks correspond to the output of the dynamical model and colored ticks are the predictions of a GLM fit to a different set of data from this model. Bottom, spike rate histograms (bin size = 10 ms) for 50 trials from the dynamical model (black) and the GLM (yellow). Only a subset of the full test data is shown. (C) Like B, but for the model with phasic dynamics. The stimulus, RF, and noise level were the same. (D) Estimated RFs from the GLMs compared to the input RF of the dynamical model. To indicate posterior uncertainty in the estimates, individual samples from the MCMC sampler are shown in light gray, and the median is overlaid in color. (E) Posterior distributions of baseline firing rate ($\omega$) and spike-history filter parameters ($\alpha_1$ and $\alpha_2$). The top panels in each column show marginal distributions for individual parameters, and the panels in the lower left corner show joint distributions for each pair of parameters. Note that more positive values of $\alpha_1$ and $\alpha_2$ correspond to stronger adaptation (i.e., a negative correlation with past spiking).

In the time domain, the effect of the membrane would be to stretch the RF out in time. In fact, what 127
we observed was that the estimated RFs were either very close to the input RF (Fig. 3D, top) or 128
compressed in time (Fig. 3D, bottom), corresponding to a relative boosting of higher frequencies. 129
This would not be possible for a model with a purely passive membrane; therefore, it must be the 130
active, voltage-gated currents that are shifting the model's temporal encoding properties. This 131
temporal distortion, which is consistent with the bandpass characteristics of $K_{LT}$ (Meng et al., 2012; 132
Chen and Meliza, 2018), will be explored further in subsequent analyses. 133

Intrinsic dynamics also affected the spike-history filter. Unlike the RF, the parameters for the 134
spike-history filter do not correspond to specific parameters in the LDC model; however, we 135
expect them to reflect the effects of currents that are activated by spiking. As seen in Fig. 3E, the 136
spike-history filter was stronger on both short ($\alpha_1$) and long ($\alpha_2$) timescales for data from the tonic 137
model compared to the phasic one. The posterior uncertainty in these parameter estimates was 138
low compared to the difference between dynamical models. This means that the spiking patterns 139
produced by phasic and tonic cells are sufficiently different, at least for this kind of stimulus and 140
amount of data, to observe changes in a single biophysical parameter. 141

**Multivariate birdsong stimulus** 142

Having demonstrated that the GLM can be used to analyze the encoding properties of a dynamical 143
model, we turned to a more realistic scenario using natural birdsong as the stimulus. The dynamics 144
remained the same as in the univariate case, but the linear stage was replaced with a spectrotempo- 145
ral RF. The stimulus, which consisted of 40 s of song from multiple zebra finches, was converted 146
to a spectrogram and convolved with the RF, summing across spectral channels. This produced a 147
univariate time series that entered into the dynamics as an external current. 148

We used RFs that were representative of the diversity found in cortical-level auditory neurons. 149
RF structure can be analyzed in terms of the modulation transfer function (MTF), a 2-D Fourier 150
transform of the RF that shows its joint spectral and temporal tuning (Woolley et al., 2005). Most 151
of the neurons in the zebra finch primary auditory pallium have MTFs with power along either 152
the spectral or temporal axis, indicating that they can be tuned to narrow spectral bands or to 153

rapid modulations of the temporal envelope, but only rarely to both (Woolley et al., 2009). This ₁₅₄ distribution is similar to the modulation spectrum of zebra finch song (Woolley et al., 2005) and ₁₅₅ at least partly reflects the statistics of early auditory experience (Moore and Woolley, 2019). Here, ₁₅₆ we simulated responses using 58 synthetic RFs drawn from this distribution (Woolley et al., 2009). ₁₅₇ Each RF was combined with the tonic and phasic dynamical models, so that we could quantify ₁₅₈ the effects of $K_{LT}$ across RF types and determine if there was any interaction with RF structure. ₁₅₉ As before, the simulated responses were used to estimate GLM parameters, but with the addition ₁₆₀ of elastic-net regularization to account for the high sparsity of the RFs and the autocorrelations ₁₆₁ present in the birdsong stimulus (see Methods). ₁₆₂

We begin by examining three examples representative of the distribution. As will be seen, the ₁₆₃ temporal characteristics of the input RF have a consistent effect on encoding properties, so we have ₁₆₄ denoted these three examples in terms of their temporal modulation transfer functions (tMTFs): ₁₆₅ wideband (WB), bandpass-low (BP-L), and bandpass-high (BP-H). As seen in Fig. 4A–F, the fitted ₁₆₆ GLMs had good predictive performance for both the phasic and tonic models and across all ₁₆₇ three input RFs, with high correlations between the spike rate histograms produced by the LDC ₁₆₈ and GL models to a novel birdsong stimulus. Thus, even with many more parameters and an ₁₆₉ autocorrelated stimulus, the GLM is still a good tool for analyzing the encoding properties of the ₁₇₀ dynamical models. ₁₇₁

As with the univariate case, the estimated RFs were qualitatively similar to the input RFs, but with ₁₇₂ distortions in the temporal profile. Most of the estimated RFs appeared to be smeared in time ₁₇₃ and with stronger and longer suppressive periods. Some of the distortions were consistent across ₁₇₄ tonic and phasic models, but there were also differences between the two dynamical models that ₁₇₅ reflect the effects of $K_{LT}$. We analyzed these effects by looking at the tMTFs, which are calculated ₁₇₆ by summing the 2D Fourier transform of the RFs across the spectral dimension (Fig. 4G). These ₁₇₇ plots show how well the model neuron is able to encode temporal modulations in the stimulus as a ₁₇₈ function of frequency. All of the estimated RFs were tuned to frequencies below 100 Hz, which is ₁₇₉ about the fastest temporal modulation rate found in zebra finch song (Singh and Theunissen, 2003). ₁₈₀ Although some of the input RFs had the potential to represent faster modulations, these frequencies ₁₈₁ were attenuated in the estimated RFs, probably because of the passive filtering properties of the ₁₈₂

**Fig. 4. GLM estimates for exemplar tonic and phasic models with zebra finch song stimuli.** (A) Receptive field parameters and responses for a model with tonic dynamics and a spectrally narrowband, temporally wideband RF. Top left, input RF in the LDC model. Top right, estimated RF from GLM. Note the temporal smearing and the broad suppression at longer lags in the estimated RF. Middle, examples of spiking responses to zebra finch song from the LDC model (top, black ticks) and the fitted GLM (bottom, red). Bottom, corresponding spike rate histograms (50 trials) for the LDC and GLM (product-moment correlation: $r_{WB} = 0.86$). (B-C) RFs and responses for models with tonic dynamics and BP-L (B) or BP-H RFs (C), same format as in (A). The GLM accurately predicted the firing rate of the LDC for these parameter values ($r_{BP-L} = 0.93$, $r_{BP-H} = 0.86$). (D-F) RFs and responses for models with the same RFs as in (A-C), but with phasic dynamics ($r_{WB} = 0.92$, $r_{BP-L} = 0.95$, $r_{BP-H} = 0.83$). All prediction correlations were high considering the underlying spiking variability in the even and odd trials of the LDC (product-moment correlations: $tonic_{WB} = 0.92$, $tonic_{BP-L} = 0.89$, $tonic_{BP-H} = 0.83$; $phasic_{WB} = 0.93$, $phasic_{BP-H} = 0.90$, $phasic_{BP-H} = 0.91$). (G) Posterior distributions of $\alpha_1$ and $\alpha_2$ comparing dynamical models for each RF. (H) Temporal modulation transfer functions of input RFs, tonic model estimates, and phasic model estimates for each of the three input RFs. Power is normalized relative to the peak for each spectrum.

membrane and the statistics of the stimulus. The main differences between the dynamical models      183
were in the attenuation of low frequencies.  Strikingly, the effects of the dynamics on lowpass      184
attention varied across RFs. For the WB input, the estimated tMTF was more bandpass in the phasic      185
model compared to the tonic model, while the opposite was true for the BP-L and BP-H inputs.      186
Thus, not only does $K_{LT}$ change the temporal encoding properties of the neuron, but this effect is      187
different depending on the filtering properties of the inputs (i.e., the input tMTF).      188

The posterior distributions for the spike-history parameters were broader than for the univariate      189
examples (Fig. 4H), indicating that the estimates are more poorly constrained by the data. This      190
was expected, given that the stimulus was shorter and more correlated. Although this reduced our      191
ability to resolve effects of dynamics on these parameters for single neurons, as the next section      192
will show, the trends in these examples were consistent across the larger sample of RFs.      193

As with the RF temporal structure, the spike-history filter parameters were affected by the inter-      194
action of RF type and dynamics.  In general, phasic models had stronger adaptation than tonic      195
models, as indicated by larger values of $\alpha_1$ or $\alpha_2$ (Fig. 4F). This effect was in the opposite direction      196
from what we saw in the univariate case (Fig. 3E), where tonic neurons had larger values of $\alpha_1$      197
and $\alpha_2$. This discrepancy presumably reflects differences in the stimulus statistics, because the      198
univariate example RF was qualitatively similar to the temporal profile of the example RFs. As has      199
been reported previously, neuron models fit to white-noise stimuli produce poor predictions to      200
natural stimuli (Theunissen et al., 2001). The univariate GLMs produced good predictions because      201
they were fit and tested with white-noise stimuli, but the parameter estimates do not generalize      202
to other kinds of stimuli.  Like many other natural stimuli, a key feature of birdsong is that the      203
temporal envelope is dominated by low frequencies. These slow oscillations produce sustained      204
periods of excitation or inhibition that drive the dynamical model into regimes where adaptive      205
processes come more strongly into play. This nonlinear interaction between stimulus statistics and      206
dynamics likely also explains why the effect of $K_{LT}$ varied across the example RFs. For the WB      207
and BP-L models, phasic dynamics (i.e., increased $K_{LT}$) caused $\alpha_1$ to increase while $\alpha_2$ remained      208
constant, but for the BP-H models, phasic dynamics did not have a large effect on either parameter.      209

## Interaction of intrinsic dynamics and RF temporal filtering

Based on these examples, we hypothesized that the key contributor to these interactions was the temporal profile of the input RF, in particular whether there was a negative lobe at longer lags. In the modulation frequency domain, this lobe corresponds to bandpass filtering. The parametric, Gabor-based model we used to generate the RFs (Woolley et al., 2009) represents this feature by a single parameter, the temporal phase ($P_t$), which is 0 for the WB example and $\frac{\pi}{2}$ for the BP-L and BP-H examples. Approximately half (26/58) of the RFs in our larger sample, those with modulation power primarily along the spectral axis, had $P_t$ of 0, whereas the RFs with power along the temporal modulation axis (32/58) had $P_t$ of $\frac{\pi}{2}$.

The performance of GLMs fit to data from the larger set of RFs was consistently good, with high correlations between the spike-rate histograms of the LDC and GL models for the tonic$_{\text{WB}}$ ($r = 0.74 \pm 0.11$), tonic$_{\text{BP}}$ ($r = 0.92 \pm .02$), phasic$_{\text{WB}}$ ($r = 0.84 \pm 0.04$), and phasic$_{\text{BP}}$ ($r = 0.93 \pm 0.05$) groups, that were comparable to the correlations between the even and odd trials of the LDC data for the tonic$_{\text{WB}}$ ($r = 0.90 \pm 0.05$), tonic$_{\text{BP}}$ ($r = 0.87 \pm 0.03$), phasic$_{\text{WB}}$ ($r = 0.92 \pm 0.01$), and phasic$_{\text{BP}}$ ($r = 0.90 \pm 0.02$) models. Performance was slightly lower for the tonic$_{\text{WB}}$ data, but the reason for this was not clear.

The results from the larger sample of RFs were consistent with our hypothesis. We looked first at the effects of dynamics on RF temporal structure, specifically the extent to which the estimated tMTF (which represents how the full LDC model encodes stimuli) was attenuated at low frequencies compared to the input tMTF ($\Delta l$). In Fig. 4G, $\Delta l$ corresponds to the difference between the black line and blue or yellow line at $f = 0$ with maximum power set to 1. Positive values of $\Delta l$ indicate that the estimated RF is more bandpass (i.e., responds less to low-frequency modulations) compared to the input RF. Negative values indicate that encoding of lower frequencies is boosted. As shown in Fig. 5, for models with WB temporal tuning, phasic dynamics attenuated low frequencies, in comparison to the matching tonic models (LMM: $b_1 = -0.38$, $n = 52$). For neurons with BP temporal tuning, the effect was the opposite: phasic dynamics caused low frequencies to be less attenuated compared to the matching tonic models ($b_1 = 0.22$, $n = 64$). In other words, across a broad range of RFs, $K_{\text{LT}}$ consistently causes neurons with broadly tuned inputs to become more

selective for higher-frequency features, but causes neurons that already have narrowly tuned inputs   238

to become more responsive to lower frequencies.   239

**Fig. 5. Phasic dynamics attenuate low-frequency temporal modulations for wideband RFs but enhance them for bandpass RFs.** Lowpass attenuation was defined as the difference in the ratios between the power at $f = 0$ and the peak power of the temporal modulation spectrum (as in Fig. 4H) of the input RF and GLM estimated RF. The y-axis shows the difference between this value for the input RF and the estimated RF. Positive values indicate that the estimated RF is more bandpass in its temporal filtering properties compared to the input RF, while negative values indicate the estimated RFs were more lowpass. For each RF, lowpass attenuation estimates for the phasic and tonic models are connected by a black dotted line. The bold dotted line shows the differences in the mean lowpass attenuation estimates (enlarged black dot) between RF types for a given model. The linear mixed effects model (LMM) with the interaction between RF type and dynamics fits significantly better than the LMM with main effects only (LMM: $\chi^2(1) = 81.8$, $p < 0.001$)

Similarly, just as we saw with the example models, the adaptation parameters also depended   240

on RF temporal structure and dynamics. As shown in Fig 6A, the general trend was for phasic   241

models to have lower spontaneous firing rates and stronger adaptation, but the effects differed   242

by RF type. Spontaneous firing rates ($\omega$; Fig 6B) were strongly suppressed by $K_{LT}$ in WB models   243

(LMM: $b_1 = -1.10$, $n = 52$) but not in BP models ($b_1 = -0.17$, $n = 64$, Fig. 6B). Short-timescale   244

adaptation ($\alpha_1$; Fig 6C) was stronger for models with phasic dynamics ($b_1 = -0.59$, $n = 116$) and   245

BP RFs ($b_2 = -0.29$, $n = 116$), but the effect of $K_{LT}$ was consistent across both kinds of RFs (i.e.,   246

no interaction). There was an interaction for longer-timescale adaptation ($\alpha_2$; Fig 6D), which was greater for BP models with phasic dynamics ($b_1 = 0.03$, $n = 64$) compared to the corresponding tonic models, but showed the opposite effect for WB models ($b_1 = -0.11$, $n = 52$). Note that in contrast to the univariate example, $\alpha_2$ estimates were consistently negative, which corresponds to a baseline facilitation (i.e., past spikes are associated with an increased probability of firing).

## Nonlinear, nonmonotonic effects of $K_{LT}$ on encoding properties

Up to this point, intrinsic dynamics have been dichotomized into tonic and phasic firing. For step currents, this dichotomy reflects a bifurcation in the dynamics: below a critical value of $g_{KLT}$, spiking is repetitive, but above this value, it occurs only at the stimulus onset (Rothman and Manis, 2003; Meng et al., 2012). For broadband current stimuli, however, the effects of $g_{KLT}$ are more graded (Chen and Meliza, 2018). To test whether $K_{LT}$ affects encoding properties in a continuous or binary manner, we simulated responses using LDC models with values of $g_{KLT}$ that varied in steps of 1 nS over a range of 0 to 50 nS (with capacitance kept constant at 60 pF), which encompasses the bifurcation in this model from tonic to phasic firing. For simplicity, we used only the three example receptive fields shown in Fig. 4 (WB, BP-L, and BP-H). Using the same birdsong stimulus, we fit GLMs to data from these simulations and examined how lowpass attenuation and adaptation were affected.

The performance of the GLM was good across all levels of $g_{KLT}$ (Fig. 7A), though for the WB RF, there was a drop in performance for larger $g_{KLT}$ values. Because this was not associated with a decrease in intertrial variability (as indicated by correlations between even and odd trials in the data; Fig. 7B), it suggests that the LDC model is more difficult to approximate with a GLM for some combinations of parameters. Overall, however, the predicted spike trains remained highly accurate, allowing resulting parameter estimates to be meaningfully interpreted.

Consistent with what we observed with dichotomized dynamics, the effects of $K_{LT}$ on RF temporal structure, spontaneous firing rate, and adaptation depended on RF type (Fig. 7C–F). With the exception of spontaneous firing rate (Fig. 7D), the trajectories of the parameters as $g_{KLT}$ increased were nonlinear and, in the case of $\alpha_1$, sometimes not even monotonic. However, there was little

**Fig. 6. Firing rate and spike-history parameter estimates depend on RF structure and dynamics.** (A) Point estimates of $\omega$, $\alpha_1$, and $\alpha_2$ GLM parameters for phasic (blues) and tonic (yellows) models by RF type. Across the diagonal are the marginal distributions for each of the parameters, with the joint distributions on the off-diagonal. (B) Scatter plot of parameter estimates showing paired phasic and tonic models (as in Fig. 5). For each RF, the phasic and tonic model parameter estimates are connected by a black dotted line. The bold dotted lines show the differences in the mean parameter estimates between RF types for a given model. The LMM with the interaction is a significantly better fit than the LMM with only main effects for $\omega$ (LMM: $\chi^2(1) = 45.9$, $p < 0.001$) and $\alpha_2$ ($\chi^2(1) = 42.4$, $p < 0.001$) but not for $\alpha_1$ ($\chi^2(1) = 0.16$, $p = 0.69$).

**Fig. 7.  Effects of low-threshold potassium conductance ($g_{KLT}$) on GLM parameters are nonlinear and depend on RF structure.** (A) Correlation coefficients between the spike-rate histograms of the LDC and GL models as a function of $g_{KLT}$. The RF$_{\text{BP-L}}$ (purple) and RF$_{\text{BP-H}}$ (orange) models were largely unaffected in their fit by $g_{KLT}$, while the RF$_{\text{WB}}$ (red) models showed a dependence on $g_{KLT}$. (B) Correlation between the even and odd trials of the LDC model as a function of $g_{KLT}$. (C–F) Lowpass attenuation, $\omega$, $\alpha_1$, and $\alpha_2$ estimates as a function of $g_{KLT}$ for the three exemplar RFs.

evidence of bifurcation, which would have appeared as a sharp discontinuity between two stable   274
regimes. These results confirm that the effects of intrinsic dynamics on encoding properties are   275
highly nonlinear, with a strong dependence on the statistics of the stimulus and the tuning of the   276
inputs.   277

## Discussion   278

These data demonstrate how intrinsic dynamics can affect the temporal encoding properties of   279
cortical-level auditory neurons. Although this effect is not unexpected, to our knowledge it has   280
not yet been quantitatively characterized. Our approach was to simulate zebra finch auditory   281
responses with a biophysically realistic linear-dynamical cascade model and then estimate encoding   282
properties using GLMs, which are statistically robust and easy to interpret. This allowed us to   283
modulate intrinsic dynamics by changing the parameter values that correspond to specific cellular   284
mechanisms and explore the effects on receptive fields and spike-history adaptation.   285

We focused on a low-threshold potassium current ($K_{LT}$), which is expressed in a subset of neurons   286
in zebra finch CM. In a previous study, we used broadband current injections to show that $K_{LT}$   287
affects temporal integration, causing neurons to reject low-frequency fluctuations and respond   288
selectively to frequencies around the maximum temporal modulation rate of zebra finch song   289
(Chen and Meliza, 2018). This filtering effect is reproduced by the dynamical model used here.   290
However, the current stimuli used to build the model were artificial and unrepresentative of   291
the stimulus-driven synaptic activity CM neurons would receive in vivo. Thus, to predict how   292
variation in $K_{LT}$ might affect auditory responses to vocal communications in this species, we drove   293
the dynamical model with an injected current that was the result of convolving natural zebra finch   294
song with a spectrotemporal RF, which we term the "input RF". Input RFs, which represent a   295
linear approximation of the preprocessing performed by the neuron's presynaptic partners, were   296
randomly drawn from a published distribution of RFs found in zebra finch Field L (Woolley et al.,   297
2009), the major source of ascending auditory input to CM (Vates et al., 1996; Wang et al., 2010).   298
This allowed us to predict which effects of the dynamics would be consistent across the population   299
and which would depend on tuning of the inputs.   300

The estimated RFs, which we interpret as the features of the stimulus that neurons encode in their    301
spiking outputs, reflected the statistics of the stimulus, the filtering properties of the input RFs,    302
and the dynamics of spiking. Estimated RFs qualitatively resembled input RFs but were distorted    303
in time. Analyzing these distortions using temporal modulation transfer functions (Fig. 4), we    304
found that most (90/116) of the model neurons were less responsive to high frequencies than their    305
inputs; we expected this effect from the lowpass filtering associated with passive leak currents. $K_{LT}$,    306
in contrast, primarily affected low frequencies in the tMTF. To our surprise, the sign of the effect    307
depended on the input tMTF, specifically how broadly tuned it was. Wideband tMTFs became more    308
bandpass, with less power at low frequencies, just as in our previous study (Chen and Meliza, 2018).    309
Bandpass tMTFs, however, became more lowpass, indicating that $K_{LT}$ was effectively boosting    310
responses to low frequencies in the stimulus. It is unclear how this happens.    311

$K_{LT}$ also affected the spike-history filter component of the GLM. Here the effects were more    312
consistent across RF types, though there were some weak but significant interactions (Fig. 6B).    313
Perhaps more surprisingly, the shape of the input RF itself influenced estimates of these parameters:    314
$\alpha_1$ estimates were higher for BP neurons whereas $\alpha_2$ was higher for WB neurons. As a result,    315
there was significant overlap in the population distributions of tonic and phasic neurons, such    316
that it would be unlikely one could infer whether a cell was tonic or phasic from the spike-history    317
parameters alone. Unfortunately, this means that intracellular recordings are still needed to    318
characterize intrinsic dynamics.    319

When dynamical neuron models are stimulated with step currents, $g_{KLT}$ is a bifurcation parameter    320
with a critical value that determines whether the cell can spike repetitively (tonic firing) or not    321
(phasic firing). We found that using more realistic currents, there is little evidence of bifurcation    322
in encoding properties, which changed smoothly as we varied $g_{KLT}$ (Fig. 7). These relationships    323
nonetheless tended to be quite nonlinear, indicating that neurons can in principle achieve dramatic    324
changes in functional response properties with only small changes in the expression or localization    325
of a single type of channel. This sensitivity is what makes intrinsic dynamics a useful target for    326
plasticity. We recently showed that CM neurons express more Kv1.1 and become more phasic    327
during the peak of the critical period for song memorization, but only in finches raised in the    328
complex acoustic environment of a colony (Chen and Meliza, 2020). In rats, exposure to dynamically    329

modulated noise causes neurons in primary auditory cortex to shift their tuning away from the       330
spectrotemporal modulation frequencies of the noise (Homma et al., 2020). Taken together with       331
our study, it may be that $K_{LT}$ is responsible for changes in sensory filtering during auditory       332
development. This mechanism could, in concert with synaptic plasticity, allow neurons to modulate       333
their responses to temporally complex acoustic environments during critical periods of auditory       334
development.       335

This study was limited to the effects of manipulating a single biophysical parameter ($g_{KLT}$) on       336
encoding of a single kind of auditory stimulus (zebra finch song), but the same approach could       337
easily be adapted to other auditory areas and sensory systems that exhibit diverse or plastic       338
intrinsic dynamics. We have shown that GLMs can accurately predict the spiking responses of       339
more complex, more biophysically realistic models across different kinds of stimuli, receptive       340
fields, and dynamical regimes. However, GLMs remain a linear approximation, and care is needed       341
in interpreting the parameter estimates, which are not linear or independent functions of the       342
underlying dynamics. Given the nonlinear kinetics of most voltage-gated currents, we expect that       343
the relationships between intrinsic dynamics and encoding properties will be complex and often       344
counterintuitive in most systems, but that there will be much to learn in each system about how       345
intrinsic dynamics reflect the computational tasks and constraints that need to be solved.       346

# Methods       347

## Stimuli Design       348

For univariate models, the stimulus consisted of 100 s of Gaussian white noise sampled at 1 kHz.       349
For multivariate models, the stimulus consisted of zebra finch song motifs recorded from 30       350
adult males in our colony. Each motif was normalized to the same RMS amplitude and repeated       351
twice, padding with at least 50 ms microphone noise at the beginning to avoid transients in the       352
convolution. The total duration of the stimulus was 63.7 s, of which 12.7 s was reserved for testing       353
performance. Spectrograms of the stimuli were calculated using a gammatone filter bank (Slaney,       354
1998) with a window size of 2.5 ms and 20 spectral channels between 1.0 and and 8.0 kHz, and a       355

step size of 1.0 ms.     356

## Receptive Field Construction     357

The univariate receptive field was generated from the difference of two gamma functions ($y(t) =$     358
$\frac{t^5}{\Gamma(6)\tau^6} \exp \frac{-t}{\tau}$) with time constants of 16 and 32 ms and an amplitude ratio of 1.5. Spectro-temporal     359
receptive fields (RFs) were parameterized as the outer product of two Gabor functions multiplied     360
by a scalar amplitude:     361

$$RF(t, f) = AH(t) \otimes G(f), \tag{1}$$
$$H(t) = \exp(-0.5[(t - t_0)/\sigma_t]^2) \cdot \cos(2\pi \cdot \Omega_t(t - t_0) + P_t),$$
$$G(f) = \exp(-0.5[(f - f_0)/\sigma_f]^2) \cdot \cos(2\pi \cdot \Omega_f(f - f_0) + P_f).$$

where $H$ is the temporal dimension of the RF, $G$ is the spectral dimension, $t_0$ is the latency, $f_0$ is the     362
peak frequency, $\sigma_t$ and $\sigma_f$ are the temporal and spectral bandwidths, $\Omega_t$ and $\Omega_f$ are the temporal     363
and spectral modulation frequencies, $P_t$ is the temporal phase (either 0 or $2\pi$), $P_f$ is the frequency     364
phase (set to 0 for all RFs), and $A$ is the amplitude. The temporal dimension $H$ had a duration of 50     365
ms with a 1 ms resolution, while the frequency dimension $G$ had 20 channels between 1 and 8 kHz.     366
We generated 59 RFs by sampling randomly from the distributions given in (Woolley et al., 2009) as     367
representative of empirically recorded RFs in primary areas of the zebra finch auditory pallium.     368
The amplitude parameter $A$ was initially set to 1 for all of the RFs, but was adjusted to between     369
1.5–2 for 7/59 models so that they would fire at least at 1 Hz on average. It was not possible to     370
meet this criterion for one RF out of the 59, so it was discarded, leaving a total of 58 RFs used for     371
analyses.     372

## Linear-Dynamical Cascade Model

Auditory responses were simulated with a model consisting of a linear, time-invariant stage that drives a conductance-based, single-compartment dynamical stage (Bjoring and Meliza, 2019). The voltage dynamics were governed by the sum of the external driving currents $I_{stim}(t)$ and $I_{noise}(t)$ and the intrinsic currents, which included a leak current and various voltage-gated currents:

$$C_m \frac{dV}{dt} = g_l(E_l - V) + g_{Na}m^3h(E_{Na} - V) + g_K n^4(E_K - V) + \cdots + I_{stim}(t) + I_{noise}(t) \qquad (2)$$

Each voltage-gated current depended on a maximal conductance $g_X$, the reversal potential for the ion species conducted by the channel $E_X$, and one or more gating variables (e.g., $m$, $h$, $n$). For all, the dynamics of the voltage-gated conductances were defined by first-order kinetics; for example,

$$\frac{dm}{dt} = \frac{m_\infty(V) - m}{\tau_m(V)}. \qquad (3)$$

The driving current, $I_{\text{stim}}(t)$ was produced by convolving the stimulus with the RF. For the univariate white noise simulation, the white noise stimulus was convolved with the 1-dimensional filter with respect to time. For the multivariate simulations, each spectral channel was convolved with the corresponding channel of the RF and the results were summed to produce a univariate time series. Multiple trials ($n = 50$) were simulated by adding pink noise to $I_{\text{stim}}(t)$, with a signal-to-noise ratio of 4.

The dynamical stage of the model was initially adapted for the zebra finch caudal mesopallium (Chen and Meliza, 2018) from the ventral cochlear nucleus model of Rothman and Manis (Rothman and Manis, 2003). This model can produce phasic or tonic responses to step currents depending on the maximal conductance of a low-threshold potassium current ($g_{KLT}$). When $g_{KLT}$ is low, the model neuron produces sustained responses to weak and moderate depolarizations; when $g_{KLT}$ is high, the model only fires at the onset of the current step. The model parameter values used

here are shown in Table 1. Each RF was paired with a tonic and a phasic model. To examine how

encoding properties change over the full range of $g_{KLT}$ values, we started with the tonic model

parameters and increased $g_{KLT}$ from 0 nS to 50 nS in steps of 1 nS.

**Table 1. Parameter values for biophysical models.**

| Parameter | Tonic | Phasic |
|---|---|---|
| $C_m$ (pF) | 60 | 40 |
| $E_l$ (mV) | –75 | –75 |
| $g_l$ (nS) | 1.3 | 1.3 |
| $E_{Na}$ (mV) | 55 | 55 |
| $g_{Na}$ (nS) | 750 | 750 |
| $E_K$ (mV) | –82 | –82 |
| $g_{KDR}$ (nS) | 0 | 0 |
| $g_{KHT}$ (nS) | 95 | 95 |
| **$g_{KLT}$** (nS) | 0 | 50 |
| $g_{KM}$ (nS) | 0 | 0 |
| $g_{KA}$ (nS) | 30 | 30 |
| $E_h$ (mV) | –43 | –43 |
| $g_h$ (nS) | 0.5 | 0.5 |

Symbols: $C_m$, capacitance; $E_l$, leak current reversal potential; $g_l$, leak conductance; $E_{Na}$, sodium reversal potential; $g_{Na}$ (maximum) sodium conductance; $E_K$, potassium reversal potential; $g_{KDR}$, delayed-rectifier potassium conductance; $g_{KHT}$, high-threshold potassium conductance; $g_{KLT}$, low-threshold potassium conductance; $g_{KM}$, M-type (slowly activating) potassium conductance; $g_{KA}$, A-type (slowly inactivating) potassium conductance; $E_h$, reversal potential for h-type (hyperpolarization-activated, cation-nonselective) current; $g_h$, h-type conductance. Bold highlights parameter differences in tonic and phasic models.

The dynamical model simulation code was generated using spyks (https://github.com/melizalab/ spyks; version 0.6.10), and the dynamics were integrated using a 5th-order Runge-Kutta algorithm with an adaptive error tolerance of $1 \times 10^{-5}$ and an interpolated step size of 0.025 ms. The output of the integration was converted to spike times by thresholding the voltage at -20 mV.

## Generalized Linear Models

A generalized linear model (GLM) (Pillow et al., 2005; Calabrese et al., 2011) was fit to the spike trains produced by the linear dynamical cascade models (Fig. 2). The conditional intensity of the model was given by:

$$\lambda(t) = exp(-\omega + \mathbf{K} * x(t) + h * y_{\text{hist}}(t)) \tag{4}$$

where $\lambda(t)$ is the conditional intensity at time $t$, $\exp(-\omega)$ corresponds to the baseline firing rate of the GLM, $\mathbf{K}$ is the RF, which is convolved with the song spectrogram $x$, and $h$ is the spike adaptation filter, which is convolved with the spike train history $y_{\text{hist}}(t)$. Note that we use $f_1 * f_2(t)$ to denote the convolution of two functions with respect to time. The full RF was a $20 \times 50$ matrix (20 spectral channels by 50 time bins of 1 ms). To reduce the number of parameters and avoid overfitting, $\mathbf{K}$ was parameterized with a rank-2 approximation; that is, the product of a $20 \times 2$ spectral filter and a $2 \times 50$ temporal filter (Thorson et al., 2015). The parameter count in the temporal dimension was further reduced by projecting into a basis set consisting of 12 raised cosine functions (Pillow et al., 2005). This basis set achieves good temporal resolution in the time immediately following a spike, with the resolution smoothly decreasing at long time intervals. The spike-history filter $h$ was parameterized in a basis set of two exponential functions:

$$h(t) = \alpha_1 \cdot \exp(\frac{t}{\tau_1}) + \alpha_2 \cdot \exp(\frac{t}{\tau_2}) \tag{5}$$

where $\tau_1$ and $\tau_2$ are time constants corresponding to short (10 ms) and long (200 ms) timescales, and $\alpha_1$ and $\alpha_2$ are the coefficients. This parameterization was chosen based on the multiadaptive timescale model, which is closely related to the GLM and has been shown to be capable of reproducing a broad range of intrinsic dynamics (Kobayashi et al., 2009; Yamauchi et al., 2011)

The GLMs were fit to data from the first 80% of the stimuli. The log-likelihood function of the GLM is given by

$$\log \mathcal{L}(\theta | t_0, \ldots, t_n) = \sum_{i=0}^{n} \log \lambda(t_i | \theta) + \int_0^T \lambda(s | \theta) ds, \tag{6}$$

where $t_i$ is the time of the $i$th spike, $n$ is the number of spikes in the experiment, $T$ is the final 421
time point of the experiment, and $\theta$ represents the free parameters (Rasmussen, 2018). Because 422
the stimulus is highly correlated and the RF is expected to be sparse, we used elastic-net regu- 423
larization to constrain the RF parameter estimates. Elastic-net regularization is combination of 424
ridge regression and the least absolute shrinkage and selection operator (LASSO). Ridge regression 425
introduces an $L_2$ penalization parameter ($\nu_2$) to account for multicollinearity, which is inherently 426
present in the highly correlated structure of the song spectrogram. The LASSO introduces an 427
$L_1$ penalization parameter ($\nu_1$) to shrink small correlations to zero and acts as a feature selection 428
algorithm, enforcing RF sparseness. A cost function was given by: 429

$$C = \nu_1 \|\mathbf{k}\|_1 + \nu_2 \|\mathbf{k}\|_2 - \log \mathcal{L}(\theta | t_0, \dots, t_n) \tag{7}$$

where $\|\mathbf{k}\|_1$ and $\|\mathbf{k}\|_2$ are the L1-norm and L2-norm of $\mathbf{K}$ (reshaped into a 1-D vector), respectively. 430
Since the log-likelihood function is concave and is guaranteed to be free of local maxima (Paninski 431
et al., 2004), we simultaneously estimated the parameters ($\omega, \mathbf{K}, \alpha_1, \alpha_2$) by minimizing the cost 432
function, which was done by using the nonlinear conjugate gradient method scipy function 433
'fmin_ncg' (version 1.3.0) (Virtanen et al., 2020). Theano (version 1.0.4) (Al-Rfou et al., 2016) was 434
used to symbolically derive the gradient and Hessian of the cost function and dynamically generate 435
C code to evaluate them. The regularization coefficients ($\nu_1, \nu_2$) and the factorization rank $D$ were 436
chosen using 4-fold cross-validation on the estimation data. 437

We quantified the uncertainty in the maximum-likelihood estimates ($\omega, \mathbf{K}, \alpha_1, \alpha_2$) by sampling 438
from the joint posterior distribution using emcee (version 2.2.1), a Python implementation of an 439
affine-invariant ensemble Markov chain Monte Carlo sampler (Foreman-Mackey et al., 2013). to 440
estimate the posterior uncertainty in the parameter estimates, $p(\theta | t_0, \dots, t_n) \propto p(\theta) L(\theta | t_0, \dots, t_n)$. 441
The log of the prior probability $p(\theta)$ was set to the elastic-net penalty (Eq. 7) using the values of $\nu_1$ 442
and $\nu_2$ obtained through cross-validation, and the log-likelihood was as in Eq. (6). An ensemble of 443
1000 chains was initialized with random values centered around the maximum-likelihood estimate 444
and given a burn-in of 2500–6000 steps. After this period, each chain was sampled one more time 445

to give a set of 1000 independent samples from $p(\theta|t_0, \ldots, t_n)$. For population-level analyses, the 446
final value of the GLM ($\omega, \mathbf{K}, \alpha_1, \alpha_2$) parameters were the median value of their respective posterior 447
distributions due to the symmetric bell-shaped curve of the posteriors. These values were very 448
close to the initial ML point estimates, so we did not sample from the posterior for the analyses 449
shown in Fig. 7. 450

To quantify performance, we generated posterior predictive distributions of spike trains from the 451
fitted GLMs, with time discretized to $\Delta = 0.5$ ms. At such short time scales, the conditional rate 452
$\lambda(t) \cdot \Delta$ could be approximated as a Bernoulli trial at each time bin which was used to produce spike 453
train responses from the GLMs. In each trial, we drew a sample from the posterior distribution, so 454
the intertrial variability reflects not only the intrinsic variance of the Bernoulli distribution but the 455
uncertainty in the parameter estimates as well. Performance was quantified as the product-moment 456
correlation between the spike-rate histograms (50 trials, 10 ms bins) for the data and the prediction 457
on the 20% of the stimulus reserved for testing. As a baseline measure of intrinsic variability, we 458
calculated the product-moment correlation between even and odd trials in the data (i.e., from the 459
linear-dynamical cascade model); however, we did not explicitly correct performance scores. 460

**Lowpass Attenuation** 461

The estimated RF parameters were projected back into a linear time basis and reshaped into a 462
$20 \times 50$ matrix. To obtain the temporal modulation transfer function (tMTF), a 2-dimensional 463
Fourier transform was performed on the RF, summing across the spectral dimension (including 464
positive and negative frequencies). The Fourier transform was calculated using the numpy package 465
in Python, with zero-padding and the application of a Hanning window in the temporal profile to 466
avoid edge effects. RF lowpass attenuation was quantified as: 467

$$\Delta l = \frac{P_0}{P_{max}} - \frac{\tilde{P}_0}{\tilde{P}_{max}} \tag{8}$$

where $P_0$ is the power for the zero frequency of the input tMTF, $P_{max}$ is the maximum power of the 468

input tMTF, $\tilde{P}_0$ is the power at the zero frequency of the estimated tMTF, and $\tilde{P}_{max}$ is the maximum ⁠469

power of the estimated tMTF. Positive values of $\Delta l$ indicate that the estimated RF responds more ⁠470

weakly to low modulation frequencies compared to the input RF, whereas negative values indicate ⁠471

that the estimated RF is more responsive to low frequencies. ⁠472

## Linear Mixed Effects Models ⁠473

Given the nested, repeated-measures nature of the experimental design (each input RF was used ⁠474

with tonic and phasic dynamical models), we used a random-intercepts LMM with input RF as a ⁠475

random effect. All LMMs were estimated using the lme4 (version 1.1.21) R package, which does not ⁠476

return p-values for parameter estimates due to unreliability issues (Bates et al., 2015). To determine ⁠477

statistical significance, we therefore took a model-comparison approach where nested LMMs of ⁠478

increasing complexity were compared against each other. Three candidate models were fit: random ⁠479

effects (variance components) only, random effects and main fixed effects, and random effects with ⁠480

main effects and interactions. Restricted maximum likelihood (REML) parameter estimation gives ⁠481

unbiased LMM estimates, however the LMMs cannot be compared as nested models (Pinheiro ⁠482

and Bates, 2006) and we therefore used maximum likelihood estimation (MLE) to generate LMMs. ⁠483

Candidate LMMs were compared across three fit statistics: AIC, BIC, and chi-squared. Lower ⁠484

values of AIC and BIC indicate better relative fit. The null hypothesis of the chi-squared test is that ⁠485

the more complicated model is not a better fit to the data than the less complicated model. ⁠486

The variance-components model was given by the equation: ⁠487

$$y_{ij} = b_0 + u_j + e_{ij} \tag{9}$$

$$u_j \sim \mathcal{N}(0, \sigma_u)$$

$$e_{ij} \sim \mathcal{N}(0, \sigma_e)$$

where $y_{ij}$ is the observed value of the dependent variable for the $i$th type of neuron model (tonic ⁠488

or phasic) and $j$th input RF type (WB or BP), $b_0$ is a fixed intercept, $u_j$ is the value of the random ⁠489

intercept of the $j$th RF type, and $e_{ij}$ is the error term for the for the LMM. Both $u_j$ and $e_{ij}$ are assumed to be normally distributed with a mean of zero and a constant variance of $\sigma_u$ and $\sigma_e$ respectively. This LMM essentially tests if the differences we see in the dependent variable are solely due to the random effects of each input RF rather than neuron model or RF type. For all LMM analyses, tonic neuron models and BP RFs were coded as 1, and phasic neuron models and WB RFs were coded as 0.

The main-effects model was given by the equation:

$$y_{ij} = b_0 + b_1 M_i + b_2 R_j + u_j + e_{ij} \tag{10}$$

where $y_{ij}$, $b_0$, $u_j$, and $e_{ij}$ are defined identically as above, $b_1$ is the fixed effect of $M_i$, the $i$th neuron model type, and $b_2$ is the fixed effect for $R_j$, the $j$th input RF type.

The interactions model is identical to the main-effects model, with the addition of a fixed effect $b_3$ of the multiplicative interaction between neuron model and RF type, with the equation given by:

$$y_{ij} = b_0 + b_1 M_i + b_2 R_j + b_3 M_i R_j + u_j + e_{ij} \tag{11}$$

If the interactions model was found to be the best fit to the data, simple-effects models were estimated using REML since these LMMs were not compared to any other candidate models. Simple effects models were calculated by subsetting the data by RF type and estimating a LMM with RF as a random intercept and neuron model type as a fixed effect. For each RF type, the LMM equation is given by:

$$y_{ij} = b_0 + b_1 M_i + u_j + e_{ij} \tag{12}$$

**Table 2. Results of $\Delta l$ LMM comparison**

| Model | AIC | BIC | $\chi^2$(df) | p-value |
|---|---|---|---|---|
| Variance Components | 28.45 | 36.71 | | |
| + Main Effects | 5.79 | 19.56 | 26.66 (2) | <.001 |
| + Interaction | -74.03 | -57.50 | 81.82 (1) | <.001 |

Reductions in AIC and BIC, as well as statistically significant chi-squared tests indicated that the LMM with main effects and interaction was the best candidate model for describing the data.

**Table 3. Results of $\omega$ LMM comparison**

| Model | AIC | BIC | $\chi^2$(df) | p-value |
|---|---|---|---|---|
| Variance Components | 260.77 | 269.03 | | |
| + Main Effects | 225.04 | 238.81 | 39.73 (2) | <.001 |
| + Interaction | 181.19 | 197.72 | 45.85 (1) | <.001 |

Reductions in AIC and BIC, as well as statistically significant chi-squared tests indicated that the LMM with main effects and interaction was the best candidate model for describing the data.

**Table 4. Results of $\alpha_1$ LMM comparison**

| Model | AIC | BIC | $\chi^2$(df) | p-value |
|---|---|---|---|---|
| Variance Components | 759.45 | 767.71 | | |
| + Main Effects | 665.64 | 679.41 | 97.8097 (2) | <.001 |
| + Interaction | 667.48 | 684.00 | 0.1617 (1) | .69 |

There were reductions in AIC and BIC, as well as statistically significant chi-squared tests for the main effects model compared to the variance components model. However, there were increases in AIC and BIC, as well as a non-statistically significant chi-squared test, for the main effects and interaction model. Therefore, the LMM with main effects only was the best candidate model for describing the data.

**Table 5. Results of $\alpha_2$ LMM comparison**

| Model | AIC | BIC | $\chi^2$(df) | p-value |
|---|---|---|---|---|
| Variance Components | -215.60 | -207.34 | | |
| + Main Effects | -238.64 | -224.87 | 27.03 (2) | <.001 |
| + Interaction | -279.03 | -262.51 | 42.39 (1) | <.001 |

Reductions in AIC and BIC, as well as statistically significant chi-squared tests indicated that the LMM with main effects and interaction was the best candidate model for describing the differences the data.

## Acknowledgments

We thank Margot Bjoring for assistance in model development and critical feedback, Laura Jamison  507
for suggestions in statistical analysis design, and Jacy Zanussi for thoughtful discussion and critical  508
feedback. This work was supported in part by funding from the National Science Foundation  509
(IOS-1942480), the Thomas F. and Kate Miller Jeffress Memorial Trust, and The Hartwell Foundation.  510

# References

Al-Rfou R, Alain G, Almahairi A, Angermueller C, Bahdanau D, Ballas N, et al. (2016) Theano: A Python framework for fast computation of mathematical expressions. *arXiv e-prints* abs/1605.02688.

Ascoli GA, Alonso-Nanclares L, Anderson SA, Barrionuevo G, Benavides-Piccione R, Burkhalter A, et al. (2008) Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex. *Nat Rev Neurosci* 9:557–568.

Bal R, Oertel D (2001) Potassium currents in octopus cells of the mammalian cochlear nucleus. *J Neurophysiol* 86:2299–2311.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4. *J Stat Soft* 67:1–48.

Bjoring MC, Meliza CD (2019) A low-threshold potassium current enhances sparseness and reliability in a model of avian auditory cortex. *PLoS Comput Biol* 15:e1006723.

Bomkamp C, Tripathy SJ, Bengtsson Gonzales C, Hjerling-Leffler J, Craig AM, Pavlidis P (2019) Transcriptomic correlates of electrophysiological and morphological diversity within and across excitatory and inhibitory neuron classes. *PLoS Comput Biol* 15:e1007113.

Calabrese A, Schumacher JW, Schneider DM, Paninski L, Woolley SMN (2011) A generalized linear model for estimating spectrotemporal receptive fields from responses to natural sounds. *PLoS ONE* 6:e16104.

Chen AN, Meliza CD (2018) Phasic and tonic cell types in the zebra finch auditory caudal mesopallium. *J Neurophysiol* 119:1127–1139.

Chen AN, Meliza CD (2020) Experience- and Sex-Dependent Intrinsic Plasticity in the Zebra Finch Auditory Cortex during Song Memorization. *J Neurosci* 40:2047–2055.

Daou A, Margoliash D (2020) Intrinsic neuronal properties represent song and error in zebra finch vocal learning. *Nat Comms* 11:1–17.

Druckmann S, Berger TK, Hill S, Schürmann F, Markram H, Segev I (2008) Evaluating automated

parameter constraining procedures of neuron models by experimental and surrogate data. *Biol Cybern* 99:371–379.

Foreman-Mackey D, Hogg DW, Lang D, Goodman J (2013) emcee: the mcmc hammer. *Publications of the Astronomical Society of the Pacific* 125:306.

Homma NY, Hullett PW, Atencio CA, Schreiner CE (2020) Auditory cortical plasticity dependent on environmental noise statistics. *Cell Rep* 30:4445–4458.

Huys QJM, Paninski L (2009) Smoothing of, and parameter estimation from, noisy biophysical recordings. *PLoS Comput Biol* 5:e1000379.

Izhikevich EM (2003) Simple model of spiking neurons. *IEEE Trans Neural Netw* 14:1569–1572.

Jarvis ED, Yu J, Rivas MV, Horita H, Feenders G, Whitney O, et al. (2013) Global view of the functional molecular organization of the avian cerebrum: mirror images and functional columns. *J Comp Neurol* 521:3614–3665.

Jolivet R, Lewis TJ, Gerstner W (2004) Generalized integrate-and-fire models of neuronal activity approximate spike trains of a detailed model to a high degree of accuracy. *J Neurophysiol* 92:959–976.

Keat J, Reinagel P, Reid RC, Meister M (2001) Predicting every spike: a model for the responses of visual neurons. *Neuron* 30:803–817.

Kobayashi R, Tsubo Y, Shinomoto S (2009) Made-to-order spiking neuron model equipped with a multi-timescale adaptive threshold. *Front Comput Neurosci* 3:9.

Llinás RR (1988) The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function. *Science* 242:1654–1664.

Meliza CD, Kostuk M, Huang H, Nogaret A, Margoliash D, Abarbanel HDI (2014) Estimating parameters and predicting membrane voltages with conductance-based neuron models. *Biol Cybern* 108:495–516.

Meng X, Huguet G, Rinzel J (2012) Type III Excitability, Slope Sensitivity and Coincidence Detection. *Discrete Contin Dyn Syst Ser A* 32:2729–2757.

Moore JM, Woolley SM (2019) Emergent tuning for learned vocalizations in auditory cortex. *Nat Neurosci* 22:1469–1476.

Nagel KI, Doupe AJ (2008) Organizing principles of spectro-temporal encoding in the avian primary auditory area field L. *Neuron* 58:938–955.

Ostojic S, Brunel N (2011) From spiking neuron models to linear-nonlinear models. *PLoS Comput Biol* 7:e1001056.

Padmanabhan K, Urban NN (2010) Intrinsic biophysical diversity decorrelates neuronal firing while increasing information content. *Nat Neurosci* 13:1276–1282.

Paninski L, Pillow JW, Simoncelli EP (2004) Maximum likelihood estimation of a stochastic integrate-and-fire neural encoding model. *Neural Comput* 16:2533–2561.

Pillow JW, Paninski L, Uzzell VJ, Simoncelli EP, Chichilnisky EJ (2005) Prediction and decoding of retinal ganglion cell responses with a probabilistic spiking model. *J Neurosci* 25:11003–11013.

Pinheiro J, Bates D (2006) *Mixed-effects models in S and S-PLUS* Springer Science & Business Media.

Pozzorini C, Mensi S, Hagens O, Naud R, Koch C, Gerstner W (2015) Automated High-Throughput Characterization of Single Neurons by Means of Simplified Spiking Models. *PLoS Comput Biol* 11:e1004275–29.

Prinz AA, Bucher D, Marder E (2004) Similar network activity from disparate circuit parameters. *Nat Neurosci* 7:1345–1352.

Rasmussen JG (2018) Lecture notes: Temporal point processes and the conditional intensity function. *arXiv preprint arXiv:1806.00221* .

Ross B, Barat M, Fujioka T (2017) Sound-making actions lead to immediate plastic changes of neuromagnetic evoked responses and induced $\beta$-band oscillations during perception. *J Neurosci* 37:5948–5959.

Rothman JS, Manis PB (2003) The roles potassium currents play in regulating the electrical activity of ventral cochlear nucleus neurons. *J Neurophysiol* 89:3097–3113.

Schulz DJ, Goaillard JM, Marder E (2006) Variable channel expression in identified single and electrically coupled neurons in different animals. *Nat Neurosci* 9:356–362.

Schwartz O, Pillow JW, Rust NC, Simoncelli EP (2006) Spike-triggered neural characterization. *J. Vis.* 6:484–507.

Sen K, Theunissen FE, Doupe AJ (2001) Feature analysis of natural sounds in the songbird auditory forebrain. *J Neurophysiol* 86:1445–1458.

Singh NC, Theunissen FE (2003) Modulation spectra of natural sounds and ethological theories of auditory processing. *J Acoust Soc Am* 114:3394–3411.

Sivaramakrishnan S, Oliver DL (2001) Distinct K currents result in physiologically distinct cell types in the inferior colliculus of the rat. *J Neurosci* 21:2861–2877.

Slaney M (1998) Auditory toolbox. *Interval Research Corporation, Tech. Rep* 10.

Theunissen FE, David SV, Singh NC, Hsu A, Vinje WE, Gallant JL (2001) Estimating spatio-temporal receptive fields of auditory and visual neurons from their responses to natural stimuli. *Network* 12:289–316.

Theunissen FE, Sen K, Doupe AJ (2000) Spectral-temporal receptive fields of nonlinear auditory neurons obtained using natural sounds. *J Neurosci* 20:2315–2331.

Thorson IL, Liénard J, David SV (2015) The Essential Complexity of Auditory Receptive Fields. *PLoS Comput Biol* 11:e1004628.

Titley HK, Brunel N, Hansel C (2017) Toward a Neurocentric View of Learning. *Neuron* 95:19–32.

Toledo-Rodriguez M, Blumenfeld B, Wu C, Luo J, Attali B, Goodman PH, Markram H (2004) Correlation maps allow neuronal electrical properties to be predicted from single-cell gene expression profiles in rat neocortex. *Cereb Cortex* 14:1310–1327.

Toth BA, Kostuk M, Meliza CD, Margoliash D, Abarbanel HDI (2011) Dynamical estimation of neuron and network properties I: variational methods. *Biol Cybern* 105:217–237.

Tripathy SJ, Burton SD, Geramita M, Gerkin RC, Urban NN (2015) Brain-wide analysis of electro-

physiological diversity yields novel categorization of mammalian neuron types. *J Neurophysiol* 113:3474–3489.

Van Geit W, De Schutter E, Achard P (2008) Automated neuron model optimization techniques: a review. *Biol Cybern* 99:241–251.

Vates GE, Broome BM, Mello CV, Nottebohm F (1996) Auditory pathways of caudal telencephalon and their relation to the song system of adult male zebra finches. *J Comp Neurol* 366:613–642.

Vavoulis DV, Straub VA, Aston JAD, Feng J (2012) A self-organizing state-space-model approach for parameter estimation in hodgkin-huxley-type models of single neurons. *PLoS Comput Biol* 8:e1002401.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. (2020) SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nat Meth* pp. 261–272.

Wang Y, Brzozowska-Prechtl A, Karten HJ (2010) Laminar and columnar auditory cortex in avian brain. *PNAS* 107:12676–12681.

Weber AI, Pillow JW (2017) Capturing the dynamical repertoire of single neurons with generalized linear models. *Neural Comput* 29:3260–3289.

Woolley SMN, Fremouw TE, Hsu A, Theunissen FE (2005) Tuning for spectro-temporal modulations as a mechanism for auditory discrimination of natural sounds. *Nat Neurosci* 8:1371–1379.

Woolley SMN, Gill PR, Fremouw T, Theunissen FE (2009) Functional groups in the avian auditory system. *J Neurosci* 29:2780–2793.

Yamauchi S, Kim H, Shinomoto S (2011) Elemental spiking neuron model for reproducing diverse firing patterns and predicting precise firing times. *Front Comput Neurosci* 5:42.