Infection of human lymphomononuclear cells by SARS-CoV-2

Marjorie C Pontelli*+[1], Italo A Castro*+[1], Ronaldo B Martins[1], Flávio P Veras[2], Leonardo La Serra[1], Daniele C Nascimento[2], Ricardo S Cardoso[1], Roberta Rosales[6], Thais M Lima[1], Juliano P Souza[1], Diego B Caetité[2], Mikhael H F de Lima[2], Juliana T Kawahisa[2], Marcela C Giannini[2,3], Letícia P Bonjorno[2,3], Maria I F Lopes[2,3], Sabrina S Batah[8], Li Siyuan[8], Rodrigo L Assad[2,3], Sergio C L Almeida[2,3], Fabiola R Oliveira[2,3], Maíra N Benatti[2,3], Lorena L F Pontes[7], Rodrigo C Santana[2,4], Fernando C Vilar[2,4], Maria A Martins[2,5], Thiago M Cunha[2], Rodrigo T Calado[7], José C Alves-Filho[2], Dario S Zamboni[2,6], Alexandre Fabro[8], Paulo Louzada-Junior[2,3], Rene D R Oliveira[2,3], Fernando Q Cunha[2], Eurico Arruda*[1].

1 Virology Research Center, Ribeirao Preto Medical School; 2 Center of Research in Inflammatory Diseases (CRID), 3 Divisions of Clinical Immunology; 4 Infectious Diseases, 5 Intensive Care Unit; 6 Department of Cell and Molecular Biology and Pathogenic Bioagents, Ribeirao Preto Medical School; 7 Blood Center of Ribeirao Preto, Ribeirao Preto, Brazil; 8 Department of Pathology, Ribeirao Preto Medical School, University of Sao Paulo, Ribeirao Preto, Sao Paulo, Brazil

+Contributed equally to the study
*Corresponding authors:

Marjorie Cornejo Pontelli; Virology Research Center, Ribeirao Preto Medical School, 3900 Bandeirantes Av, 14049-900, Ribeirao Preto –SP – Brazil. Tel.: +55 16 33154508, Email: marpontelli@gmail.com

Italo Araujo Castro; Virology Research Center, Ribeirao Preto Medical School, 3900 Bandeirantes Av, 14049-900, Ribeirao Preto –SP – Brazil. Tel.: +55 16 33154508, Email: italo.a.castro@gmail.com

Eurico Arruda; Virology Research Center, Ribeirao Preto Medical School, 3900 Bandeirantes Av, 14049-900, Ribeirao Preto –SP – Brazil. Tel.: +55 16 33153337, Email: eaneto@fmrp.usp.br

Abstract. Although SARS-CoV-2 severe infection is associated with a
hyperinflammatory state, lymphopenia is an immunological hallmark, and
correlates with poor prognosis in COVID-19. However, it remains unknown if
circulating human lymphocytes and monocytes are susceptible to SARS-CoV-2
infection. In this study, SARS-CoV-2 infection of human peripheral blood
mononuclear cells (PBMCs) was investigated both *in vitro* and *in vivo*. We found
that *in vitro* infection of whole PBMCs from healthy donors was productive of
virus progeny. Results revealed that monocytes, as well as B and T
lymphocytes, are susceptible to SARS-CoV-2 active infection and viral
replication was indicated by detection of double-stranded RNA. Moreover, flow
cytometry and immunofluorescence analysis revealed that SARS-CoV-2 was
frequently detected in monocytes and B lymphocytes from COVID-19 patients,
and less frequently in CD4$^+$T lymphocytes. The rates of SARS-CoV-2-infected
monocytes in PBMCs from COVID-19 patients increased over time from
symptom onset. Additionally, SARS-CoV-2-positive monocytes and B and
CD4+T lymphocytes were detected by immunohistochemistry in post mortem
lung tissue. SARS-CoV-2 infection of blood circulating leukocytes in COVID-19
patients may have important implications for disease pathogenesis, immune
dysfunction, and virus spread within the host.
56
57

Introduction

In December 2019, a new coronavirus emerged as the cause of a severe acute respiratory disease named Coronavirus-related disease 2019 (COVID-19). The virus that spilled over to humans in China was classified in the family *Coronaviridae*, genus *Betacoronavirus*, and was named Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), for its similarity to SARS-CoV [1].

Since its emergence, SARS-CoV-2 has spread to 185 countries/political regions and infected more than 11 million people worldwide, with a death toll of approximately 500,000 cases. The main clinical features of COVID-19 are fever, dry cough, dyspnea and myalgia, but some patients rapidly evolve to severe respiratory distress syndrome [2].

Previous studies have shown that inflammatory cytokine storm and lymphocytopenia are important markers of severe COVID-19 cases, with severe functional exhaustion of TCD4+ and TCD8+ lymphocytes [2–4]. Interestingly, peripheral blood mononuclear cells (PBMCs) from COVID-19 patients showed upregulation of autophagy and apoptosis pathways [5], suggesting that dampening of the immune system by SARS-CoV-2 infection may have a strong impact on the clinical outcome of severe COVID-19.

A decrease in circulating lymphocytes has been associated with poor COVID-19 outcome, but it is still unclear whether that lymphopenia is directly due to SARS-CoV-2 infection of lymphocytes with consequent cell death. SARS-CoV-2 interacts with target cells via binding of its major surface glycoprotein spike (S) with the angiotensin-converting enzyme 2 (ACE2) present in the cell membrane [6]. ACE2-independent cell-entry has also been reported and could be an alternative mechanism of SARS-CoV-2 entry in cells with low ACE2 expression [7]. Cleavage of the S protein is required for efficient entry of SARS-CoV-2, which is accomplished by transmembrane serine protease TMPRSS2 [6].

In addition to respiratory disease, COVID-19 patients frequently develop gastrointestinal symptoms, which is in keeping with the high expression of TMPRSS2 and ACE2 documented in enterocytes [8]. Furthermore, the SARS-

CoV-2 antigen was found *post mortem* in the spleen and lymph nodes with
pathological signs of damage. In these organs, monocytes do contain viral
antigens, but it was not clear whether this was due to active viral replication or
phagocytosis, nor if monocytes become infected before reaching secondary
lymphoid tissues [9]. While SARS-CoV-2 causes viremia, until now, infectious
SARS-CoV-2 was not successfully isolated from peripheral blood in COVID-19
patients, and it is suggested that the virus in blood may be cell-associated [5, 9].
In this study, we investigated the susceptibility and permissiveness of human
peripheral blood mononuclear cells (PBMC) to SARS-CoV-2. We found that
PBMCs are susceptible and permissive to SARS-CoV-2 infection, both *in vivo*
and *ex vivo*, which seems to play a direct role in the reduction of circulating
lymphocytes.
103
104 **Patients and Methods**
105
Ethical statement and COVID-19 patients. The study was approved by the
National Ethics Committee (CONEP, CAAE: 30248420.9.0000.5440 and
31797820.8.0000.5440). A total of 29 hospitalized patients were enrolled, all
with clinical and radiological features of COVID-19 and confirmed SARS-CoV-2
infection by RT-PCR in respiratory secretions, with detection of specific IgM or
IgG antibodies to SARS-CoV-2. Clinical features, laboratory results and drug
therapies are summarized in **Supplementary Table 1**. For all comparisons, 12
age and gender-matching healty controls were also enrolled. Written informed
consent was obtained for both patients and healthy controls.
115
Production of mouse anti-SARS-CoV-2 hyperimmune serum. Male C57Bl/6
mice were bred and maintained under specific pathogen-free conditions at the
animal facility of the Ribeirão Preto Medical School (FMRP) at University of São
Paulo. The protocol for production of mouse hyperimmune serum were carried
out with 8-week-old male mice following the institutional guidelines on ethics in
animal experiments and was approved by the University of São Paulo Ethics
Committee for Animal Experimental Research - CETEA (Protocol no. 001/2020-
1). To immunize animals, virus stock was inactivated by adding formaldehyde to

a final concentration of 0.2%, and incubated overnight at 37°C. Then, virus was

purified by ultracentrifugation (10% sucrose cushion, 159.000 × g for 1h). The

pellet was resuspended with Phosphate Buffer Saline (PBS) 1x and stored at -

20°C. In order to confirm inactivation, titration of the inactivated product was

done both by $TCID_{50}$ and by plaque assay in Vero-E6 cells with 5-day

incubation, without any cytopathic effects. Three C57Bl/6 mice were inoculated

intramuscularly with an emulsion containing the equivalent of 10^6 $TCID_{50}$ of

inactivated SARS-CoV-2 in complete Freund's adjuvant (CFA, BD, cat. 263810)

diluted 1:1 in PBS. Boosts were given with inactivated SARS-CoV-2 in

incomplete Freund's adjuvant (without *M. tuberculosis,* IFA, BD, cat. 263910) on

days 7 and 14 after the first immunization. One week after the last dose,

animals were euthanized with an excess of anesthetics xylazine (60 mg/kg) and

ketamine (300 mg/kg), following exsanguination by cardiac puncture. Animal

serum conversion was evaluated by indirect immunofluorescence using slide

preparations of SARS-CoV-2 infected Caco-2 cells, fixed with 4%

paraformaldehyde and AlexaFluor 488-labelled rabbit anti-mouse secondary

antibody. Coverslips were analyzed using an optic microscope (Olympus

BX40).

Isolation of peripheral blood mononuclear cells (PBMCs). Human PBMCs

were isolated from COVID-19 patients or healthy donors by density gradient

using Percoll (GE Healthcare, cat. 17-5445-01), as previously described [10,

11]. PBMCs were washed, resuspended in RPMI 1640 supplemented with 10%

fetal bovine serum (FBS) and kept on ice until further use.

Virus and cell lines. The passage 1 (P1) of SARS-CoV-2 Brazil/SPBR-02/2020

isolate obtained in Vero-E6 from a COVID-19 patient in Sao Paulo was kindly

provided by Prof. Edison Durigon (ICB-USP). P1 was diluted 1:1000 in

Dulbecco's modified Eagle's medium (DMEM) and inoculated in Vero-E6 cells

monolayers to produce the P2 stock. For stock titration, serial 10-fold dilutions

were inoculated in quadruplicate monolayers of Vero-E6 cells and incubated at

37°C in 5% CO_2. On the fourth day of incubation, the presence of cytopathic

effect (CPE) was recorded **(Supplementary Fig 1A)** and titers were expressed

as the 50% tissue culture infectious dose ($TCID_{50}$), using the Reed-Muench

method. All experiments involving SARS-CoV-2 propagation were done in

biosafety level 3 laboratory.

***In vitro* infection of PBMCs.** For these experiments, 10^6 PBMCs from 5

healthy donors were infected with SARS-CoV-2 (MOI=1) in RPMI with 0% FBS

at RT for 1 h under orbital agitation. Next, cells were pelleted at 300 × g, the

inoculum was washed and replaced by RPMI with 2% FBS and cells were

incubated at 37°C in 5% CO_2. As controls, equivalent quantities of cells were

exposed to UV-inactivated SARS-CoV-2 and treated in the same way. We also

used a control consisting of cells treated with 20 mM of NH_4Cl starting 20 min

before infection and maintained throughout the entire incubation period.

Supernatants from PBMCs were collected at 0, 6, 12, 24 and 48 h post-infection

and subjected to serial ten-fold dilutions to determine virus titers by $TCID_{50}$, as

previously described. In parallel, PBMCs were treated with 0.5ug/ml Camostat

(Sigma Aldrich, cat. SML005) or with 10 uM of anti-ACE2 antibody (Rhea

Biotech, cat. IM-0060) starting 1 hour before infection. Then, innoculum was

washed, the media containing the different treatments were replaced, and the

PBMCs were kept for 24h at 37°C in 5% CO_2.

RNA extraction and real-time RT-PCR. SARS-CoV-2 RNA detection was

done with primer-probe sets for SARS-CoV-2 according to the USA-CDC

protocol, targeting the virus N1 gene, and using the RNAse-P housekeeping

gene as control, by one-step real-time RT-PCR. Total RNA was extracted with

Trizol® (Invitrogen, CA, EUA) from 250µL of homogenized cell pellets and

supernatants from in vitro assays. All real-time PCR assays were done on a

Step-One Plus thermocycler (Applied Biosystems, Foster City, CA, USA).

Briefly, after Trizol® extraction, 100 ng of RNA was used for genome

amplification with N1 primers (20 µM) and probe (5 µM), and TaqPath 1-Step

qRT-PCR Master Mix (Applied Biosystems, Foster City, CA, USA), with the

following parameters: 25°C for 2 min, 50°C for 15 min, 95°C for 2 min, followed

by 45 cycles of 94 °C for 5 s and 60°C for 30s. Viral loads of SARS-COV-2 were

determined using a standard curve prepared with a plasmid containing a 944bp

amplicon, which includes all three targets for the sets of primers/probes

designed by CDC protocol (N1, N2 and N3), inserted into a TA cloning vector

(PTZ57R/T CloneJetTM Cloning Kit Thermo Fisher®). Results of viral RNA

quantifications by one-step qRT-PCR were plotted with GraphPad® Prism 8.4.2

software.

Indirect immunofluorescence staining of SARS-CoV-2 infected cells.

Coverslips pre-treated with poly-lysine 0.1% (Sigma-Aldrich, cat. P8920) were

incubated with isolated PBMCs from patients or healthy donors at 37ºC, 20

minutes for cell adherence. After that, coverslip-containing cells were fixed with

4% paraformaldehyde (PFA) in PBS for 15 minutes, and then washed 3 times

with PBS. To detect viral antigens in cells, we used serum from a recovered

COVID-19 patient, which was first tested for specificity by immunofluorescence

in SARS-CoV-2 infected Vero CCL81 cells (**Supplementary Fig 1B).** In

addition, for each experiment using the referred serum we included cells from

healthy donors or non-infected cells. As an isotype control of this serum, we

used a human serum collected in 2016. Biotin-conjugated anti-human IgG

(Sigma-Aldrich, cat. B-1140) was used as the secondary antibody, followed by

amplification with the TSA Cyanine 3 System (Perkin Elmer, NEL704A001KT),

following the manufacturer's protocol. To determine the phenotype of SARS-

CoV-2-infected cells, we used primary antibodies for CD4 (Abcam cat.

ab133616), CD8 (Abcam cat. ab4055), CD14 (Abcam cat. ab133335), CD19

(Abcam cat. ab134114), CD20 (Abcam cat. ab103573). For detection of virus

replication, we used a mouse anti-dsRNA J2 (dsRNA; English & Scientific

Consulting Kft, Hungary), which binds to dsRNA of 40 bp or longer. Secondary

antibodies used were polyclonal anti-rabbit conjugated with 488 (Thermo Fisher

cat. A21202), 594 (Abcam cat. ab150116) or 647 (Abcam cat. ab150079). The

Golgi complex and nuclei staining were carried out using a mouse anti-GM130

(BD cat. 610822) and 4′,6-diamidino-2-phenylindole dihydrochloride dye (DAPI,

Thermo Fisher cat. 62248), respectively.

Confocal microscopy. PBMCs from confirmed COVID-19 patients and from

healthy control donors were stained with human serum containing antibodies to

SARS-CoV-2, and with commercial antibodies to the different cell phenotypes,

followed by the appropriate secondary antibodies. Preparations were analyzed

in a Zeiss Confocal 780 microscope in a Tile 3x3 in a single focal plane. The

quantity of SARS-CoV-2-positive cells of different phenotypes was quantified by

using the analyze particles tool from Fiji by ImageJ.

Flow cytometry. Unseparated whole blood leukocyte samples from COVID-19

patients or healthy donors infected in vitro with SARS-CoV-2 were surface

stained with Fixable Viability Dye eFluor™ 780 (eBioscience) and monoclonal

antibodies specific for CD3 (APC eBioscience cat. 17-0036-42), CD4 (PerCP-

Cy5.5 BD cat. 560650), CD8 (PE-Cy7 BD cat. 557746), CD19 (APC BioLegend

cat. 302212), CD14 (PerCP Abcam cat. ab91146), CD16 (PE eBioscience cat.

12-0168-42), CCR2 (BV BioLegend cat. 357210) for 30 min at 4ºC, according to

manufacturer's instructions. Detection of SARS-CoV-2 by flow cytometry was

performed with BD Cytofix/Cytoperm™ kit to enable access to intracellular

antigens using mouse polyclonal antibody raised against formalin-inactivated

SARS-CoV-2, as described early in this manuscript, for 15 min at 4°C. To

ensure that viral detection was specific for replicating intracellular viruses,

additional preparations of infected cells were stained without permeabilization.

Treatment with trypsin for 60 min on ice after infection to remove surface-bound

viral particles was also included as a second control (**Supplementary Fig 2)**.

SARS-CoV-2 antibodies were detected with secondary anti-Mouse Alexa488.

Surface phosphatidylserine (PS) staining was carried out in whole blood using

ApoScreen AnnexinV-FITC apoptosis kit (SouthernBiotech cat 10010-02),

following manufacturer's guidelines. All data were acquired using a Verse or

Canto flow cytometers (BD Biosciences) and subsequent analysis was done

using FlowJo (TreeStar) software. Gating strategies are illustrated in

Supplementary Fig 3.

Serial immunohistochemistry. Tissue sections from paraffin-embedded lung

fragments obtained from two COVID-19 fatal cases were tested by

immunohistochemistry (IHC) using anti-SARS-CoV-2 polyclonal antibody for in

situ detection of SARS-CoV-2. Sequential immunoperoxidase labeling and

erasing (SIMPLE) [12] was then performed to determine the

immunophenotypes of SARS-CoV-2 infected cells, using antibodies to CD4

(Abcam cat. ab133616), CD20 (Abcam cat. ab103573), CD14 (Abcam cat.

ab133335) and IL-6 (BD cat. 554400). After each round of staining, slides were

scanned using a VS120 ScanScope (Olympus) under 400x magnification.

Images were pseudocolored and overlaid in the first image of the preparation

counterstained with hematoxylin using ImageJ v1.50b (NIH, USA) and Adobe

Photoshop CS5 software (Adobe Systems, San Jose, CA, USA). Lung paraffin-

embedded tissue obtained from a fatal case of hantavirus infection in 2016 was

used as a negative control for SARS-CoV-2 staining.

Statistical analysis. All descriptive statistics, patient stratification, and positive

cell frequencies were done using GraphPad Prism Software, version 6.0.

Correlation analysis, one-way ANOVA, two-way ANOVA, linear regressions,

Holm-Sidak, and Bonferroni post-tests were also performed using GraphPad

Prism. Values of P <0.05 were considered significant, as described in all

figures.

Results

SARS-CoV-2 infection of human PBMCs is productive. Considering that human lymphocyte and monocyte lineages are susceptible to SARS-CoV-2 infection *in vitro*, we sought to determine whether primary cultures of human PBMCs could also be infected. Therefore, PBMCs from five healthy donors were infected *in vitro* at a MOI=1. After 0, 6, 12, 24 and 48 hpi, supernatants were harvested, and virus progeny was titrated. SARS-CoV-2 titers peaked between 6 and 12 hpi, resulting in a 100-fold increase from the initial input, and decreased steadily thereof (**Fig 1A**). As expected, induction of general intracellular alkalization by treatment with NH_4Cl reduced progeny production by approximately 10x (p=0.017). Interestingly, virus progeny production was not entirely abolished by NH_4Cl treatment, suggesting an entry pathway alternative to endosomal acidification in PBMCs (**Fig 1B**).

Even though expression of ACE2 is minimal in human PBMCs in general [13, 14], we evaluated the viral production after blocking ACE2 and TMPRSS2. Virus titers obtained after Camostat blockage of TMPRSS2 were not significantly different from those obtained without the treatment (**Fig 1C**), suggesting that PBMC infection is not dependent on TMPRSS2. Conversely, the blockage of ACE2 with anti-ACE2 antibody resulted in reduction, but not abrogation of SARS-CoV-2 progeny production after 24 hpi (p=0.0216) (**Fig 1C**), indicating that SARS-CoV-2 can infect human PBMCs independently of ACE2.

Coronavirus replication entails the formation of abundant double-stranded RNAs (dsRNA) in the cytoplasm of infected cells, and thus its intracellular detection is a reliable marker of viral replication. Therefore, infected PBMCs were stained for SARS-CoV-2 and dsRNA and analyzed by confocal microscopy. Most SARS-CoV-2-positive cells were also positive for dsRNA, and rates of double-positive cells counted at 6 hours post-infection followed a pattern that roughly matched the accumulation of progeny (**Fig. 1D**). The dsRNA staining was seen as clear puncta in SARS-CoV-2-infected cells, in a pattern suggestive of virus factories.

Monocytes and T lymphocytes are the main targets of SARS-CoV-2 *in vitro* infection. To determine the susceptibility of circulating leukocytes to SARS-CoV-2, PBMCs from five healthy donors were infected (MOI=1), and analyzed the intracellular expression of SARS-CoV-2 antigens by flow cytometry. After 24 hpi, SARS-CoV-2 was detected in all immunophenotyped cells (**Fig 2A**). Monocytes were the most susceptible cell type, showing significant SARS-CoV-2 antigen staining (44.3%, p=0.039) (**Fig 2B**). In addition to monocytes, T CD4$^+$ (14.2%, p=0.028), CD8$^+$ (13.5%, p=0.019) and B lymphocytes (7.58%) were also susceptible to SARS-2 infection (**Fig 2C**). Staining for SARS-CoV-2 was significantly reduced in cells treated with NH$_4$Cl, suggesting that acidification is important for in vitro infection of PBMCs.

Infection of T lymphocytes leads to cell death by apoptosis. The COVID-19-related lymphocytopenia has been well described as a strong indicator of severe clinical outcomes in patients. Since we found both T CD4$^+$ and CD8$^+$ cells susceptible to SARS-2 infection *in vitro*, we investigated the presence of cell death in SARS-CoV-2-infected PBMCs from 5 healthy donors by the expression of translocated phosphatidylserine (PS) on the cell surface 24 hours post-infection by analizing its binding to annexin V (**Fig 3**). Despite the basal annexin V staining (CD4$^+$ mean 6.24%, CD8$^+$ mean 12.36%) seen in non-infected cells (**Fig 3A**), strong staining was observed both in live T CD4$^+$ (70.88%, p=0.0001) and CD8^{++} lymphocytes (39.72%, p=0.0009) (**Fig 3B**). When cells were analyzed independently of Live/Dead staining, differences were still significant and even increased for CD8$^+$ (59.64%, p=0.0001) (**Supplementary Fig 4**), indicating that a considerable percentage of Annexin V-positive CD8$^+$ cells were already dead. No significant differences were observed in cell death between cells infected in the presence or absence of NH$_4$Cl during infection. These results indicated that infection of human PBMCs by SARS-CoV-2 sharply increased the expression of apoptosis markers in T lymphocytes.

Circulating immune cells from COVID-19 patients are infected by SARS-CoV-2. During April 7th to June 18th, we enrolled 22 COVID-19 patients that were admitted to the intensive care unit (ICU), presenting a moderate to severe

disease. Clinical and demographic characteristics of enrolled patients are listed

in Supplementary Table 1. Blood samples were collected at admission in the

ICU. To check for SARS-CoV-2 infection in PBMCs from COVID patients, we

analyzed PBMCs prepared from the whole blood of 22 patients and 11 healthy

donors by flow cytometry (**Fig 4A**) with staining for SARS-CoV-2 antigens. Cells

from COVID-19 patients showed significant expression of SARS-CoV-2

antigens (7.68%±1.56 p=0.008) in comparison with cells from healthy donors

(**Fig 4B**). Interestingly, not all COVID-19 patients showed expressive staining

for SARS-CoV-2, and rates of SARS-CoV-2-positive cells ranged from 0.16 to

33.9% (**Fig 4B**). Additionally, PMBCs from 15 COVID-19 patients were tested

for the SARS-CoV-2 genome by real-time RT-PCR. Viral genome was detected

in 8 out of 15 PBMC samples (53.3%), with mean viral load of $3.8x10^4$ copies

per μg of RNA (**Supplementary Table 2**). Immunophenotyping of cells from

COVID-19 patients indicated that the highest proportion of SARS-CoV-2-

positive cells was found in B lymphocytes (42.73%±4.3). Although susceptible

to *in vitro* infection, we were not able to find significant numbers of SARS-CoV-2

positive T cells in PBMCs from COVID-19 patients by flow cytometry. Similarly

to what was observed by the *in vitro* experiments, monocytes (CD14$^+$) from

patients were found to be positive for SARS-CoV-2 in a high percentage

(14.19%±15.26). Inflammatory monocytes (CD14$^+$CCR2$^+$ and

CD14$^+$CD16$^+$CCR2$^+$) were positive for SARS-CoV-2 antigen in rates

significantly higher in comparison with healthy controls (18.73%±18.46 and

14.78%±15.5, respectively) (**Fig 4C**). To confirm the results obtained by flow

cytometry, immunofluorescence was done for SARS-CoV-2 antigens in PBMCs

isolated from COVID-19 patients. Some staining of SARS-CoV-2 with variable

intensity was observed in CD19 and CD14 cells in PBMCs from COVID-19

patients, with no discernible fluorescent signal seen in PBMCs from healthy

donors (**Fig 4D**). Despite what was observed by FC experiments, some IF

staining was found in CD4 T lymphocytes, and after extensively screening, very

few T CD8 cells were found to be positive for IF (**Supplementary Fig 5**).

Since the detection of SARS-CoV-2 in patients was found to be variable

(Figure 4C), we selected 15 COVID cases to analyze individual differences in

rates of SARS-CoV-2-positive cells. Patients were stratified based on the time

of sample collection after symptoms onset, and SARS-CoV-2-positive cell

frequencies were plotted on a heatmap for all cell immunophenotypes analyzed

(**Fig 4E**). It became clear that rates of SARS-CoV-2-positive B lymphocytes

were high throughout the entire dataset. In contrast, rates of SARS-CoV-2-

positive monocytes were higher after following time progression after symptoms

onset (**Fig 4E**). Frequencies of SARS-CoV-2-positive cells correlated positively

with the length of time of COVID-19 progression after symptoms onset,

especially for inflammatory $CD14^+CCR2^+$ monocytes (r=0.442 p=0.044) (**Fig**

4F).

To confirm whether SARS-CoV-2 was actively replicating in PBMCs from

COVID-19 patients, we analyzed the presence of dsRNA in SARS-CoV-2-

positive cells of different immunophenotypes by immunofluorescence and

confocal microscopy. Remarkably, dsRNA staining was found in most SARS-

CoV-2-positive cell subsets, $CD4^+$ T lymphocytes, B lymphocytes, and

monocytes (**Fig 5**). Altogether, these data confirm that SARS-CoV-2 infects

circulating white blood cells from COVID-19 patients, and the frequencies of

SARS-CoV-2-positive monocytes in the peripheral blood increase with time of

onset of symptoms.

Infected inflammatory monocytes are detected post mortem in lung

tissues from COVID-19 patients. The respiratory tract is the classical entry

route of coronaviruses in mammalian hosts. Therefore, we checked if the same

infected cell immunophenotypes found in PBMCs could also be found by

immunohistochemistry in the lungs of COVID-19 patients obtained *post mortem*.

Post mortem lung specimens from COVID-19 patients revealed abundant

staining for SARS-CoV-2, especially throughout the entire bronchovascular

axes and alveolar-capillary barriers. Control lung specimens showed no staining

(**Supplementary Fig 6**). Upon staining for SARS-CoV-2, slides were scanned,

the staining was erased, and re-stained sequentially for the surface antigens

CD4, CD20, and CD14. The serial immunolabelling indicated that $CD4^+$ T

lymphocytes, B lymphocytes, and monocytes express SARS-CoV-2 antigens

(**Fig 6**) in the lungs of COVID-19 cases. Additionally, due to its well-known role

in lung tissue damage in COVID-19, IL-6-positive cells were also searched for

and, interestingly, several $CD14^+$ monocytes expressing IL-6 were also positive

for SARS-CoV-2 (**Fig 6C-E**), indicating that inflammatory monocytes in lungs of
COVID-19 patients can also be infected with SARS-CoV-2.
411
412   **Discussion**
413
It has been well accepted that several SARS-CoV-2 strategies to escape
innate immune sensing, coupled with dysregulation of immune responses in
early phases of infection, drive a cytokine storm that is a hallmark of severe
COVID-19 [15–17]. Importantly, lymphopenia has also been recognized as a
feature of severe infection by SARS-CoV-2. Postmortem examination of
spleens and lymph nodes showed the presence of SARS-CoV-2 in those
organs, infecting ACE2-expressing macrophages and causing important tissue
damage [10].
SARS-CoV-2 detection in tissues far from the entry sites in the
respiratory tract, without exuberant viremia, suggests that SARS-CoV-2 may
reach target organs by alternative ways. One possibility could be the infection of
leukocytes that could serve as "Trojan horses" transporting the virus to
secondary infection sites. In that regard, we have recently reported that SARS-
CoV-2 infects neutrophils, which could also act as Trojan horses carrying
SARS-CoV-2 to neutrophil infiltrated tissues [18]. However, until now, it has
been unclear whether SARS-CoV-2 infects PBMCs in vivo, thus creating a
possibility of them being Trojan horses of viral dissemination. To address this
question, we first infected PBMCs from healthy donors *in vitro* with SARS-CoV-
2, as a preliminary way to check for their susceptibility and permissiveness to
the virus. Virus production in PBMCs peaked at 12 hpi, reaching titers 100-fold
the initial input, with steady decay thereafter until 48 hpi. The presence of
dsRNA in SARS-CoV-2 infected PBMCs in the first few hours after infection
provides further evidence that the virus replicates, yet modestly, in PBMCs in
vitro. These results are in keeping with reports of SARS-CoV infection of human
PBMCs [19]. Moreover, the treatment of PBMCs with ammonium chloride,
which elevates the pH and prevents organelle acidification, significantly reduced
but did not abrogate SARS-CoV-2 replication, consistent with an alternative
acidification-independent pathway.

The immunophenotyping of PBMCs infected in vitro with SARS-CoV-2
revealed that CD14[+], CD4[+], CD8[+] and CD19[+] cells were susceptible. Primary
human monocytes have been reported as susceptible to MERS-CoV and, more
recently, to SARS-CoV-2 [20, 21]. In contrast, another recent study did not
report PBMC infection by SARS-CoV-2 in vitro, possibly due to the low MOI
used [22], coupled with the reported reduced expression of ACE2 by
lymphocytes [14]. Despite that, we found that blockade of ACE2 partially
reduced SARS-CoV-2 titers in supernatants of infected PBMCs, suggesting that
among PBMCs there are ACE2-expressing cell types that contribute to the total
virus progeny production. However, ACE2 blockade does not eliminate virus
production, what strongly suggests the existence of ACE2-independent
mechanisms of infection in lymphohematopoietic cells.
A recent report indicated that SARS-CoV-2 spike protein can interact
with surface CD147, which could be an alternative virus receptor, in a way
similar to what was observed for SARS-CoV [7, 23]. The transmembrane
glycoprotein CD147, also known as Basignin, is expressed in some subsets of
T lymphocytes [24], and thus could play a role in SARS-CoV-2 entry in these
cells as well.
An intense T cell depletion in peripheral blood is seen in up to 85% of
severe COVID-19 patients [2, 25]. Furthermore, T cells from COVID-19 patients
show considerable levels of exhaustion markers [3, 4], and transcriptome
analysis of their BBMCs indicated upregulation of genes involved in apoptosis
and p53-signalling pathways [5]. These data suggests that SARS-CoV-2
infection could induce cell death by apoptosis in PBMCs, what could also
happen in inflamed secondarily infected organs of COVID-19 patients. Of note,
lymphopenia was also described in Middle East Respiratory Syndrome (MERS)
patients, in whom MERS-CoV can directly infect human primary T lymphocytes
and induce T-cell apoptosis through extrinsic and intrinsic pathways [20].
Annexin V staining showed that SARS-CoV-2 infection of PBMCs caused
increased translocation of phosphatidylserine (PS) to the cell surface of both
CD4[+] and CD8[+] T lymphocytes. The translocation of PS and subsequent
scrambling of lipid membrane asymmetry is indicative of late-stage apoptosis
[26]. Importantly, in the presence of NH_4Cl, SARS-CoV-2 infection significantly
increased annexin V labelling, suggesting that even at reduced levels of

replication, SARS-CoV-2 can trigger apoptosis in lymphocytes. Taken together,

the data indicate that SARS-CoV-2 infection of lymphocytes causes cell death,

which may concur to the observed lymphopenia. The association of

lymphopenia with poor prognosis may be related to the death of specific T-cell

subsets, which may result in loss of immune response regulatory components,

and drive a cytokine storm that can crosstalk with neutrophil NETosis [27]. Also,

it can be related to increased IL-6 and Fas-FasL interactions [10], resulting in

severe lymphoid tissue alterations [28].

In addition to in vitro infection, SARS-CoV-2 was also detected in PBMCs

from COVID-19 patients, more prominently in B lymphocytes and

subpopulations of monocytes. The predominance of B lymphocytes as target

cells of SARS-CoV-2 infection in vivo, in contrast to what was seen in PBMCs

infected in vitro, suggests that the susceptibility of different lymphocyte subsets

in natural SARS-CoV-2 infection may depend on ACE2-independent alternative

virus entry mechanisms. These findings corroborate previous observations that

SARS-CoV enters B lymphocytes and monocyte-derived cells via a FcγRII-

dependent pathway, which is facilitated by the presence of antibodies [29, 30].

The present results were obtained based on one-time sampling of patients who

were enrolled at different times of COVID-19 evolution, what may explain the

heterogeneity in rates of SARS-CoV-2-positive cells of different

immunophenotypes observed among them. Accordingly, SARS-CoV-2 RNA

was not detected in PBMCs from all, but in 53% of patients, indicating that

SARS-CoV-2 infection in PBMCs may be variable, depending on host factors

still unidentified, or present only in later phases of COVID-19, as suggested by

the positive correlation between time from symptoms onset and frequency of

SARS-CoV-2 positive cells in PBMCs. A possible explanation for an increase in

SARS-CoV-2 susceptible cells over time could be an increase in ACE2

expression, triggered by type I IFN [31]. In this context, it is noteworthy that the

replication of SARS-CoV in PBMCs was not sustained for long periods [19, 32].

To the best of our knowledge, this is the first report of circulating

lymphoid cells positive for SARS-CoV-2, and the presence of dsRNA indicates

that these cells are targets of virus replication. This may considerably impact

the cells immune competence during COVID-19 and may help cell-associated

SARS-CoV-2 spread to secondary infection sites.

SARS-CoV-2 recruits important inflammatory infiltrate in the lungs,
containing diverse immune cell types that bear close contact with SARS-CoV-2-
infected lung cells, such as pneumocytes and alveolar macrophages [33]. In the
present study, we found $CD4^+$ T and B lymphocytes and, importantly, also IL-6-
expressing inflammatory monocytes positive for SARS-CoV-2 infiltrating the
lung tissue from fatal cases of COVID-19. Further studies will be required to
clarify whether SARS-CoV-2-positive lympho-mononuclear cells become
infected in the lung or enter the affected tissue from the bloodstream already
containing the virus. Regardless of where the immune cells become infected by
SARS-CoV-2, their presence in the peripheral blood can impact directly on virus
dissemination, delivering the infectious virus to secondary sites of infection.

Inflammatory monocytes play a significant role in the immunopathology
of COVID-19 [15, 16] and ICU patients have high levels of circulating
CD14+CD16+ inflammatory monocytes, which correlates with unfavorable
outcomes [34, 35]. Increased expression of CCR2 and other inflammatory
markers by monocytes leads to the infiltration of tissues with high expression of
the correspondent CCL2 chemokine [15]. Interestingly, inflammatory monocytes
with the same profile were found abundantly in bronchoalveolar lavage fluids
from patients with severe COVID-19 [36]. Based on that, our data suggest that
CD14+CCR2+ and CD14+CD16+CCR2+ infected monocytes could act as
Trojan horses and traffic viruses to secondary sites of infection, where SARS-
CoV-2 causes severe tissue damage. Additional lymphoid cell recruitment to
damaged tissues may further contribute to lymphopenia [9].

Overall, the infection of lymphomononuclear cells by SARS-CoV-2 in
peripheral blood from patients with COVID-19 has important consequences for
pathogenesis of this multifaceted disease, including possible compromises of
immune cell functions, and helping the virus to reach immune-privileged
secondary sites of infection.

Conflict of interests
The authors declare none.

Acknowledgements

543   This study was supported by the Brazilian National Research Council
544 (CNPq grant numbers 310100/2017-8, 403201/2020-9 and INCTC
545 465539/2014-9) and the Sao Paulo State Research Foundation (FAPESP grant
546 numbers 2013/16349-2 and 2014/02438-6). Dr. Pontelli was funded by CNPq
547 grant number 380849/2020-8. The authors also thank Dimensions Sciences, a
548 Non-Profit Organization that granted Dr. Castro a research scholarship while
549 this study was conducted.

551
552
553

References

[1]   Zhou, P.; Yang, X. Lou; Wang, X. G.; Hu, B.; Zhang, L.; Zhang, W.; Si, H. R.; Zhu, Y.; Li, B.; Huang, C. L.; et al. A Pneumonia Outbreak Associated with a New Coronavirus of Probable Bat Origin. *Nature*, **2020**, *579* (7798), 270–273. https://doi.org/10.1038/s41586-020-2012-7.

[2]   Huang, C.; Wang, Y.; Li, X.; Ren, L.; Zhao, J.; Hu, Y.; Zhang, L.; Fan, G.; Xu, J.; Gu, X.; et al. Clinical Features of Patients Infected with 2019 Novel Coronavirus in Wuhan, China. *Lancet*, **2020**, *395* (10223), 497–506. https://doi.org/10.1016/S0140-6736(20)30183-5.

[3]   Diao, B.; Wang, C.; Tan, Y.; Chen, X.; Liu, Y.; Ning, L.; Chen, L.; Li, M.; Liu, Y.; Wang, G.; et al. Reduction and Functional Exhaustion of T Cells in Patients With Coronavirus Disease 2019 (COVID-19). *Front. Immunol.*, **2020**, *11* (May), 1–7. https://doi.org/10.3389/fimmu.2020.00827.

[4]   Zheng, H. Y.; Zhang, M.; Yang, C. X.; Zhang, N.; Wang, X. C.; Yang, X. P.; Dong, X. Q.; Zheng, Y. T. Elevated Exhaustion Levels and Reduced Functional Diversity of T Cells in Peripheral Blood May Predict Severe Progression in COVID-19 Patients. *Cell. Mol. Immunol.*, **2020**, *17* (5), 541–543. https://doi.org/10.1038/s41423-020-0401-3.

[5]   Xiong, Y.; Liu, Y.; Cao, L.; Wang, D.; Guo, M.; Jiang, A.; Guo, D.; Hu, W.; Yang, J.; Tang, Z.; et al. Transcriptomic Characteristics of Bronchoalveolar Lavage Fluid and Peripheral Blood Mononuclear Cells in COVID-19 Patients. *Emerg. Microbes Infect.*, **2020**, *9* (1), 761–770. https://doi.org/10.1080/22221751.2020.1747363.

[6]   Hoffmann, M.; Kleine-Weber, H.; Schroeder, S.; Kruger, N.; Herrler, T.; Erichesen, S.; Schiergens, T. S.; Wu, N.-H.; Nitsche, A.; Muller, M. A.; et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*, **2020**, *181* (16), 1–9.

[7]   Wang, K.; Chen, W.; Chen, Z.-N. SARS-CoV-2 Invades Host Cells via a Novel Route: CD147-Spike ProteiN. *bioRxiv*, **2012**, *91* (5), 1–10. https://doi.org/10.1101/2020.03.14.988345.

[8]   Zang, R.; Gomez Castro, M. F.; McCune, B. T.; Zeng, Q.; Rothlauf, P. W.; Sonnek, N. M.; Liu, Z.; Brulois, K. F.; Wang, X.; Greenberg, H. B.; et al. TMPRSS2 and TMPRSS4 Promote SARS-CoV-2 Infection of Human Small Intestinal Enterocytes. *Sci. Immunol.*, **2020**, *5* (47). https://doi.org/10.1126/sciimmunol.abc3582.

[9]   chen, yongwen; Feng, Z.; Diao, B.; Wang, R.; Wang, G.; Wang, C.; Tan, Y.; Liu, L.; Wang, C.; Liu, Y.; et al. The Novel Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Directly Decimates Human Spleens and Lymph Nodes. *medRxiv*, **2020**, *2*, 2020.03.27.20045427. https://doi.org/10.1101/2020.03.27.20045427.

[10]   Rios-Santos, F.; Alves-Filho, J. C.; Souto, F. O.; Spiller, F.; Freitas, A.; Lotufo, C. M. C.; Soares, M. B. P.; Dos Santos, R. R.; Teixeira, M. M.; Cunha, F. D. Q. Down-Regulation of CXCR2 on Neutrophils in Severe Sepsis Is Mediated by Inducible Nitric Oxide

Synthase-Derived Nitric Oxide. *Am. J. Respir. Crit. Care Med.*, **2007**, *175* (5), 490–497.

https://doi.org/10.1164/rccm.200601-103OC.

[11] Alves-Filho, J. C.; Freitas, A.; Souto, F. O.; Spiller, F.; Paula-Neto, H.; Silva, J. S.;

Gazzinelli, R. T.; Teixeira, M. M.; Ferreira, S. H.; Cunha, F. Q. Regulation of Chemokine

Receptor by Toll-like Receptor 2 Is Critical to Neutrophil Migration and Resistance to

Polymicrobial Sepsis. *Proc. Natl. Acad. Sci. U. S. A.*, **2009**, *106* (10), 4018–4023.

https://doi.org/10.1073/pnas.0900196106.

[12] Glass, G.; Papin, J. A.; Mandell, J. W. SIMPLE: A Sequential Immunoperoxidase

Labeling and Erasing Method. *J. Histochem. Cytochem.*, **2009**, *57* (10), 899–905.

https://doi.org/10.1369/jhc.2009.953612.

[13] Qi, F.; Qian, S.; Zhang, S.; Zhang, Z. Single Cell RNA Sequencing of 13 Human Tissues

Identify Cell Types and Receptors of Human Coronaviruses. *Biochem. Biophys. Res.*

Commun., **2020**, *526* (1), 135–140. https://doi.org/10.1016/j.bbrc.2020.03.044.

[14] Uhlén, M.; Fagerberg, L.; Hallström, B. M.; Lindskog, C.; Oksvold, P.; Mardinoglu, A.;

Sivertsson, Å.; Kampf, C.; Sjöstedt, E.; Asplund, A.; et al. Tissue-Based Map of the

Human Proteome. *Science (80-.).*, **2015**, *347* (6220).

https://doi.org/10.1126/science.1260419.

[15] Zhou, Y.; Fu, B.; Zheng, X.; Wang, D.; Zhao, C.; Qi, Y.; Sun, R.; Tian, Z.; Xu, X.; Wei, H.

Pathogenic T-Cells and Inflammatory Monocytes Incite Inflammatory Storms in Severe

COVID-19 Patients. *Natl. Sci. Rev.*, **2020**, No. March, 1–5.

https://doi.org/10.1093/nsr/nwaa041.

[16] Mehta, P.; McAuley, D. F.; Brown, M.; Sanchez, E.; Tattersall, R. S.; Manson, J. J.

COVID-19: Consider Cytokine Storm Syndromes and Immunosuppression. *Lancet*,

2020, *395* (10229), 1033–1034. https://doi.org/10.1016/S0140-6736(20)30628-0.

[17] Vabret, N.; Britton, G. J.; Gruber, C.; Hegde, S.; Kim, J.; Kuksin, M.; Levantovsky, R.;

Malle, L.; Moreira, A.; Park, M. D.; et al. Immunology of COVID-19: Current State of the

Science. *Immunity*, **2020**, *1*. https://doi.org/10.1016/j.immuni.2020.05.002.

[18] Veras, F.; Pontelli, M. C.; Cunha, F. SARS-CoV-2 Triggered Neutrophil Extracellular

Traps (NETs) Mediate COVID-19 Pathology. *medRxiv*, **2020**.

[19] Li, L.; Wo, J.; Shao, J.; Zhu, H.; Wu, N.; Li, M.; Yao, H.; Hu, M.; Dennin, R. H. SARS-

Coronavirus Replicates in Mononuclear Cells of Peripheral Blood (PBMCs) from SARS

Patients. *J. Clin. Virol.*, **2003**, *28* (3), 239–244. https://doi.org/10.1016/S1386-

6532(03)00195-1.

[20] Chu, H.; Zhou, J.; Wong, B. H. Y.; Li, C.; Chan, J. F. W.; Cheng, Z. S.; Yang, D.; Wang,

D.; Lee, A. C. Y.; Li, C.; et al. Middle East Respiratory Syndrome Coronavirus Efficiently

Infects Human Primary T Lymphocytes and Activates the Extrinsic and Intrinsic

Apoptosis Pathways. *J. Infect. Dis.*, **2016**, *213* (6), 904–914.

https://doi.org/10.1093/infdis/jiv380.

[21] Codo, A. C.; Davanzo, G. G.; Monteiro, L. D. B.; Souza, G. F. De; Muraro, S. P.;

Carregari, V. C.; Alberto, C.; Biagi, O. De; Crunfli, F.; Jimenez, J. L.; et al. Elevated

Glucose Levels Favor SARS-CoV-2 Infection and Monocyte Response through a HIF-

1α/Glycolysis Dependent Axis.

[22] Banerjee, A.; Nasir, J. A.; Budylowski, P.; Yip, L.; Aftanas, P.; Christie, N.; Ghalami, A.;

Baid, K.; Raphenya, A. R.; Hirota, J. A.; et al. Isolation, Sequence, Infectivity, and

Replication Kinetics of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerg.*

Infect. Dis., **2020**, *26* (9). https://doi.org/10.3201/eid2609.201495.

[23] Chen, Z.; Mi, L.; Xu, J.; Yu, J.; Wang, X.; Jiang, J.; Xing, J.; Shang, P.; Qian, A.; Li, Y.; et

al. Function of HAb18G/CD147 in Invasion of Host Cells by Severe Acute Respiratory

Syndrome Coronavirus. *J. Infect. Dis.*, **2005**, *191* (5), 755–760.

https://doi.org/10.1086/427811.

[24] Solstad, T.; Bains, S. J.; Landskron, J.; Aandahl, E. M.; Thiede, B.; Taskén, K.;

Torgersen, K. M. CD147 (Basigin/Emmprin) Identifies FoxP3 +CD45RO +CTLA4 +-

Activated Human Regulatory T Cells. *Blood*, **2011**, *118* (19), 5141–5151.

https://doi.org/10.1182/blood-2011-02-339242.

[25] Chen, N.; Zhou, M.; Dong, X.; Qu, J.; Gong, F.; Han, Y.; Qiu, Y.; Wang, J.; Liu, Y.; Wei,

Y.; et al. Epidemiological and Clinical Characteristics of 99 Cases of 2019 Novel

Coronavirus Pneumonia in Wuhan, China: A Descriptive Study. *Lancet*, **2020**, *395*

(10223), 507–513. https://doi.org/10.1016/S0140-6736(20)30211-7.

[26] Suzuki, J.; Denning, D. P.; Imanishi, E.; Horvitz, H. R.; Nagata, S. Xk-Related Protein 8

and CED-8 Promote Phosphatidylserine Exposure in Apoptotic Cells. *Science (80-.).*,

2013, *341* (6144), 403–406. https://doi.org/10.1126/science.1236758.

[27] Barnes, B. J.; Adrover, J. M.; Baxter-Stoltzfus, A.; Borczuk, A.; Cools-Lartigue, J.;

Crawford, J. M.; Daßler-Plenker, J.; Guerci, P.; Huynh, C.; Knight, J. S.; et al. Targeting

Potential Drivers of COVID-19: Neutrophil Extracellular Traps. *J. Exp. Med.*, **2020**, *217*

(6), 1–7. https://doi.org/10.1084/jem.20200652.

[28] Blanco-Melo, D.; Nilsson-Payant, B. E.; Liu, W. C.; Uhl, S.; Hoagland, D.; Møller, R.;

Jordan, T. X.; Oishi, K.; Panis, M.; Sachs, D.; et al. Imbalanced Host Response to

SARS-CoV-2 Drives Development of COVID-19. *Cell*, **2020**, *181* (5), 1036-1045.e9.

https://doi.org/10.1016/j.cell.2020.04.026.

[29] Kam, Y. W.; Kien, F.; Roberts, A.; Cheung, Y. C.; Lamirande, E. W.; Vogel, L.; Chu, S.

L.; Tse, J.; Guarner, J.; Zaki, S. R.; et al. Antibodies against Trimeric S Glycoprotein

Protect Hamsters against SARS-CoV Challenge despite Their Capacity to Mediate

FcγRII-Dependent Entry into B Cells in Vitro. *Vaccine*, **2007**, *25* (4), 729–740.

https://doi.org/10.1016/j.vaccine.2006.08.011.

[30] Jaume, M.; Yip, M. S.; Cheung, C. Y.; Leung, H. L.; Li, P. H.; Kien, F.; Dutry, I.;

Callendret, B.; Escriou, N.; Altmeyer, R.; et al. Anti-Severe Acute Respiratory Syndrome

Coronavirus Spike Antibodies Trigger Infection of Human Immune Cells via a PH- and

Cysteine Protease-Independent Fc R Pathway. *J. Virol.*, **2011**, *85* (20), 10582–10597.

https://doi.org/10.1128/jvi.00671-11.

[31] Ziegler, C. G. K.; Allon, S. J.; Nyquist, S. K.; Mbano, I. M.; Miao, V. N.; Tzouanas, C. N.;

Cao, Y.; Yousif, A. S.; Bals, J.; Hauser, B. M.; et al. SARS-CoV-2 Receptor ACE2 Is an
Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific
Cell Subsets across Tissues. *Cell*, **2020**, 1016–1035.
https://doi.org/10.1016/j.cell.2020.04.035.
[32]   Yilla, M.; Harcourt, B. H.; Hickman, C. J.; McGrew, M.; Tamin, A.; Goldsmith, C. S.;
Bellini, W. J.; Anderson, L. J. SARS-Coronavirus Replication in Human Peripheral
Monocytes/Macrophages. *Virus Res.*, **2005**, *107* (1), 93–101.
https://doi.org/10.1016/j.virusres.2004.09.004.
[33]   Roosecelis B. Martines; Ritter, J. M.; Matkovic, E.; Gary, J.; Bollweg, B. C.; Bullock, H.;
Goldsmith, C. S.; Silva-Flannery, L.; Seixas, J. N.; Reagan-Steiner, S.; et al. Pathology
and Pathogenesis of SARS-CoV-2 Associated with Fatal Coronavirus Disease, United
States. *Emerg. Infect. Dis.*, **2020**, *26*. https://doi.org/10.3201/eid2609.202095.
[34]   Zhang, D.; Guo, R.; Lei, L.; Liu, H.; Wang, Y.; Wang, Y.; Dai, T.; Zhang, T.; Lai, Y.;
Wang, J.; et al. COVID-19 Infection Induces Readily Detectable Morphological and
Inflammation-Related Phenotypic Changes in Peripheral Blood Monocytes, the Severity
of Which Correlate with Patient Outcome. *medRxiv*, **2020**, 2020.03.24.20042655.
https://doi.org/10.1101/2020.03.24.20042655.
[35]   Zhou, Y.; Fu, B.; Zheng, X.; Wang, D.; Zhao, C. In Severe Pulmonary Syndrome
Patients of a New Coronavirus. **2020**.
[36]   Liao, M.; Liu, Y.; Yuan, J.; Wen, Y.; Xu, G.; Zhao, J.; Cheng, L.; Li, J.; Wang, X.; Wang,
F.; et al. Single-Cell Landscape of Bronchoalveolar Immune Cells in Patients with
COVID-19. *Nat. Med.*, **2020**, *26* (6), 842–844. https://doi.org/10.1038/s41591-020-0901-
9.
697
698

Figure legends

Figure 1. Human primary blood cells are susceptible and permissive to

SARS-CoV-2. Blood from five healthy donors was collected and PBMCs were

separated by Ficoll density gradient. Cells were infected with SARS-CoV-2

Brazil/SPBR-02/2020 (MOI-1) and cultured for 48 h. (A) Overtime virus progeny

production from PBMCs infected with SARS-CoV-2. Supernatants from cultured

PBMCs were collected at each time point and titrated by $TCID_{50}$. The small

symbols represent individual values (5 healthy donors) and error bars depict

standard deviation. (B) SARS-CoV-2 progeny titers in supernatants of infected

PBMCs at 24 and 48 hpi, with and without treatment with 20mM NH_4Cl. (C)

Effects of blocking SARS-CoV-2 cell receptor ACE2 and TMPRSS2 on virus

progeny production. Infected PBMCs were exposed to antibody anti-ACE2 or

Camostat and virus progeny was titrated in supernatants at 24 hpi. (D)

Immunostaining for dsRNA in PBMCs cultured on poly-lysine –coated

coverslips 6h after SARS-CoV-2 infection. Cells were fixed, immunostained for

SARS-CoV-2 (red), dsRNA (cyan) and analyzed by confocal microscopy.

Statistical analysis was performed using one-way or two-way ANOVA. Tukey's

or Holm-Sidak post-tests were applied when suitable. P values < 0.05 were

considered significant. Magnification: 63x. Scale bars 10 μM.

Figure 2. SARS-CoV-2 differentially infects subsets of human PBMCs in

vitro. (A) Representative flow cytometry plots of PBMCs infected with SARS-

CoV-2 (24 hpi) in the presence or absence of 20mM NH_4Cl with gating in live

$CD14^+$, $CD4^+$, $CD8^+$ or $CD19^+$ cells. Representative histograms of the

fluorescence for each condition in comparison with the proper controls. The

gray-shaded curve indicates secondary antibody Alexa488 signal background,

while the dashed curve indicates the background signal in mock-infected cells.

The light- and dark-colored curves indicate respectively cells infected in the

presence and absence of NH_4Cl. Percentages of SARS-CoV-2-infected

monocytes (B) and lymphocytes (C), showing the average mean fluorescent

intensity (MFI) in the panels on the right and the frequency (%) of SARS-CoV-2-

infected cells in the panels on the left. Mean ± s.d. is indicated on the bar

graphs. Significance was determined by one-way ANOVA with Bonferroni's

post-test.

Figure 3. SARS-CoV-2 infection of PBMCs increases expression of

phosphatidylserine (PS) in T lymphocytes. (A) Representative flow

cytometry plots of live $CD4^+$ and $CD8^+$ T cells positive for Annexin V staining in

PBMCs from five heathy donors 24h after infection with SARS-CoV-2 (MOI= 1).

(B) Percentages of live lymphocytes positive for Annexin V and expressing PS

in the cell surface after SARS-CoV-2 *in vitro* infection. Mean ± s.d. is indicated

for all bar graphs. Significance was determined by one-way ANOVA and

Bonferroni's post-test.

Figure 4: Detection of SARS-CoV-2 in PBMCs from hospitalized COVID-19

patients. (A) Representative flow cytometry plots indicating SARS-CoV-2

positivity of PBMCs from COVID-19 patients in comparison with isotype control

and healthy donors. (B) Percentages of SARS-CoV-2-infected cells from

COVID-19 patients (n=22) compared with background signal from healthy

donors (n=12) cells. Results were compared by unpaired *t*-test (p=0.008). (C)

Percentages of SARS-CoV-2-infected cells considering the different

immunophenotypes, in COVID-19 patients (n=22). (D) Immunofluorescence of

PBMCs from COVID-19 patients labeling for SARS-CoV-2 (red), nuclei (blue)

and immunophenotypes CD4, CD19 or CD14 (green). Scale bars: 50 μM (E)

Heat-map indicating SARS-CoV-2-positive cell frequencies for each

immunophenotype, stratified by time from symptoms onset (Patient

number/symptoms onset (days)). Data was plotted individually for each COVID-

19 patient analyzed. (F) Correlation and linear regression analysis between time

after symptoms onset and frequencies of SARS-CoV-2-positive cells. Both 'p'

and 'r' values are indicated in the graphs. The best-fit line is displayed in all the

graphs, while the light-color area represents the confidence interval. P values

<0.05 were considered significant.

Figure 5. Peripheral blood cells naturally infected by SARS-CoV-2 from

COVID-19 patients presents double-stranded RNA, a replication

intermediate. PBMC from (A) healthy donors or (B) COVID-19 patients were

isolated and put on coverslips pre-treated with poly-lysyne. Cells were fixed and

stained for SARS-CoV-2 (red), immune phenotypes as CD4, CD19 or CD14

(green), dsRNA (cyan) and nuclei (blue). Immunofluorescence was examined

using confocal microscopy. In the bottom left corner of each channel, an inset of

the labelling phenotype is shown. Representative images for each

immunophenotype, where at least two patients were analyzed. Magnification

63x. Scale bar 10 um.

Figure 6. SARS-CoV-2 is detected in diverse immune cell types in COVID-

19 lungs. (A, F and I) SARS-CoV-2 staining pseudocolored in green with

hematoxylin counterstaining. (B, G, J) Staining for the immunophenotypes

CD14, CD20 and CD4, respectively, pseudocolored in red. (D): Staining for IL-

6, pseudocolored in magenta. (C, E, H and K) Overlaid layers from the previous

sequential rounds of staining, with superimposed staining indicated in yellow.

(c', e', h' and k') Insets from the respective previous panels. Scale bars: 50 µM.

Figure S1. Validation of SARS-CoV-2 detection with human convalescent
serum. Vero cells were infected with SARS-CoV-2 (MOI=1) or mock infected
and incubated for 48h. (A) Phase-contrast microscopy of uninfected (left panel)
and SARS-CoV-2-infected Vero cell monolayer showing cytopathic effect.
Magnification 400×. (B) Immunofluorescence of Vero cells infected with SARS-
CoV-2 or mock-infected at 48hpi, when cells were fixed and stained for GM130
(red), virus (green) and nuclei (DAPI). Scale bar 10 µm.

Figure S2. Flow cytometry (FC) of SARS-CoV-2-infected PBMCs from
healthy with labeling for SARS-CoV-2. PMBCs from healthy donors infected
in vitro (MOI=1) were analyzed by FC using mouse polyclonal anti-SARS-CoV-2
with and without cell permeabilization. Treatment with trypsin to remove
surface-bound viral particles was used as an additional control. (A)
Representative histograms of surface and intracellular staining for SARS-CoV-
2, with SARS-CoV-2-infected cells in red and trypsin-treated infected cells in
black. (B) Comparison of intracellular and surface staining of infected cells
treated or not with trypsin, and non-infected cells in percentages on the left and
MFI on the right.

Figure S3. Gating strategies used for immunophenotyping of SARS-CoV-
2-infected cells. (A) Cells were initially gated to exclude doublets and to
exclude dead cells, using Live/Dead APC/H7 and CD3 staining. Next, detection
of SARS-CoV-2 antigens in live T lymphocytes was defined based on the
background secondary antibody signal (Alexa488) and signal obtained in
healthy donors (flow plots and representative histograms). The same strategy
was used for CD19+ B lymphocytes. (B) Live monocytes were initially gated as

described for lymphocytes. Next, expression of CD14 and CD16 was used to
define circulating monocyte subpopulations. Expression of CCR2 by CD14$^+$ and
CD14$^+$CD16$^+$ cells was used to define inflammatory monocytes. Among every
defined subpopulation, expression of SARS-CoV-2 antigens was defined in
comparison with secondary antibody background and healthy donors staining
(flow plots and representative histograms).
814
815   **Figure S4. Percentage of lymphocytes expressing phosphatydilserine (PS)**
816   **on the surface after in vitro infection with SARS-CoV-2**. Cells were analyzed
817   independently of Live/Dead APC/H7 staining. Mean ± s.d. is shown for all bar
818   graphs. Significance was determined by one-way ANOVA and Bonferroni's
819   post-test was applied.
820
Figure S5. **T CD8 lymphocytes are rarely detected with SARS-CoV-2.** (A)
PBMC from COVID-19 patients were put on coverslips pre-treated with poly-
lysyne, fixed and stained for SARS-CoV-2 (red), CD8 (green) and nuclei (blue).
Coverslips were analyzed in epifluorescence microscopy. Magnification 400x.
Scale bar 20 um. (B) dsRNA detection in CD8 cells. PBMC was labeled as
described in (a) and dsRNA (cyan) using an anti-J2 antibody. At the bottom left
corner an inset is shown. Coverslips were analyzed in confocal microscopy.
Magnification 63x. Scale bar 10 um.
829
Figure S6. Immunohistochemistry for SARS-CoV-2 antigens in *post*
***mortem* lungs from COVID-19.** (A) *Post mortem* lung fragment from a
hantavirus fatal case obtained in 2016, as a negative control for SARS-CoV-2
staining. (B) Staining for SARS-CoV-2 in lung from COVID-19 fatal case. (b')
Individual cells showing strong cytoplasmic staining for SARS-CoV-2 antigens
in detail. Scale bars: 50 µM.
836
837
838

839 **Figure 1**

845 **Figure 2**

849 **Figure 3**

855 **Figure 4**

859 **Figure 5**

A

863 **Figure 6**

868 **Figure S1**

870
871
872

Figure S2

Figure S3

881 **Figure S4**

Figure S5

889 **Figure S6**

891
892
893

894 **Supplementary Tables**

896 **Supplementary Table 1. COVID-19 patient characteristics**

Demographics		%
Number	29	
Age (years)	61.89$\pm$17.10	
Hospital day	9.35$\pm$4.138	
Female	13	44%
Comorbidities		
Hypertension	16	55%
Diabetes	13	44%
Obesity	13	44%
Lung disease	4	14%
History of smoking	9	31%
Heart disease	7	24%
Kidney disease	2	7%
History of stroke	2	7%
Cancer	4	14%
Autoimmune diseases	2	7%
Immune deficiency	2	7%
Laboratorial findings		
CRP (mg/dL)[*]	14.41$\pm$8.11	
D-Dimers (µg/mL)[**]	2,244$\pm$1,698	
LDH (U/L)[#]	749,4$\pm$490,5	
Ferritin (ng/mL)[&]	1,985$\pm$2836	
Haemoglobin (g/dL)	12.16$\pm$2.62	
Neutrophils (cell/mm^3)	7,521$\pm$4,952	
Lymphocytes (cell/mm^3)	1,666$\pm$1,286	
Platelets (count/mm^3)	231,009$\pm$136,433	
Image findings (n)	29	100%
Medications		
Antibiotics	29	100%
Heparin	29	100%
Antimalarial	3	10%
Oseltamivir	11	37%
Glucocorticoids	16	55%
Respiratory status		
Mechanical ventilation	24	83%
Nasal-cannula oxygen	27	93%
Room air	0	
pO$_2$	70.36$\pm$45.64	
SatO$_2$	82.71$\pm$18.58	
Outcome		
Deaths	9	31%

897 [*]*CRP: C-reactive protein (Normal value <0.5 mg/dL);* [**]*D-dimers (NV <0.5 µg/mL);* [#]*LDH: lactic*
898 *dehydrogenase (Normal range: 120-246 U/L);* [&]*Ferritin (NR: 10-291 ng/mL)*

Supplementary Table 2. Viral loads of SARS-CoV-2 in PBMCs from COVID-19 patients tested by real-time RT-PCR

Patient ID	Mean viral load Genome copies/total RNA (µg)
P39	68560
P42	22770
P43	20782
P45	36507
P46	22140
P47	94736
P48	26455
P50	16396
Mean±s.d.	38543±28114

906