Intrinsic excitability mechanisms of neuronal ensemble formation

Tzitzitlini Alejandre-García, Samuel Kim, Jesús Pérez-Ortega and Rafael Yuste

Neurotechnology Center, Dept. Biological Sciences, Columbia University, New York, NY 10027, USA

Running Head: Mechanisms of neuronal ensembles

Correspondence to:
Rafael Yuste
902 NWC Building 550 West 120 Street, Box 4822, New York, NY, 10027.
Email: rmy5@columbia.edu

Number of pages (Title page, Abstract, Introduction, Methods, Results, Discussion): 30

Number of figures: 9

Number of words: Abstract (), Introduction (), Discussion ()

Conflict of Interest: The authors declare no competing financial interests

Acknowledgments: We thank James Holland, for his assistance and members of the Yuste Lab for useful comments. Supported by R01EY011787 and R01MH115900. T.A and J.P. have postdoctoral fellowships from the National Council of Science and Technology from Mexico (CONACYT). The authors have no competing financial interests to declare. S.K. and R.Y. conceived the project. S.K. and T.A. performed experiments and T.A. and R.Y. wrote the paper. T.A., S.K. and J.P. analyzed the data. All authors planned experiments, discussed results and edited the paper. R.Y. assembled and directed the team and secured funding and resources.

Significance Statement:

We investigated the synaptic and cellular mechanisms underlying the formation of neuronal ensembles, i.e., spontaneously coactive groups of neurons. Using *in vitro* electrophysiology and optogenetic in slices of mouse neocortex we replicated a protocol that generates ensembles *in vivo*. After optogenetic and electrical stimulation, we observed biphasic synaptic plasticity and, unexpectedly, major increases in excitability, input resistance, and reductions in firing threshold. The increased excitability can explain the observed synaptic plasticity. Our results reveal a major role for intrinsic excitability in the establishment of neuronal ensembles.

Abstract

Neuronal ensembles are coactive groups of cortical neurons, found in spontaneous and evoked

activity, that can mediate perception and behavior. To understand the mechanisms that lead to

the formation of ensembles, we co-activated optogenetically and electrically layer 2/3 pyramidal

neurons in brain slices from mouse visual cortex, in animals from both sexes, replicating *in vitro*

an optogenetic protocol to generate ensembles *in vivo*. Using whole-cell and perforated patch-

clamp pair recordings we find that, after optogenetic or electrical stimulation, coactivated

neurons increase their correlation in spontaneous activity, a hallmark of ensemble formation.

Coactivated neurons showed small biphasic changes in presynaptic plasticity, with an initial

depression followed by a potentiation after a recovery period. Unexpectedly, optogenetic and

electrical stimulation-induced significant increases in frequency and amplitude of spontaneous

EPSPs, even after single-cell stimulation. In addition, we observed strong and persistent

increases in neuronal excitability after stimulation, with increases in membrane resistance and

reduction in spike threshold. A pharmacological agent that blocks changes in membrane

resistance can revert this effect. These significant increases in excitability may partly explain the

observed biphasic synaptic plasticity. We propose that cell-intrinsic changes in excitability are

involved in the formation of neuronal ensembles. We propose an "iceberg" model, by which

increased neuronal excitability makes subthreshold connections suprathreshold, enhancing the

effect of already existing synapses, and generating a new neuronal ensemble.

Introduction

The function of the brain is anchored on the activity of groups of connected neurons, forming

microcircuits (Schüz and Braitenberg, 2001; Shepherd, 2004). Specifically, groups of coactive

neurons, or ensembles (also known as assemblies, synfire chains, or attractors), found in both

spontaneous and evoked activity, which exhibit synchronous or correlated activity (Lorente De

Nó, 1938; Hebb, 1949; Hopfield, 1982; Abeles, 1991; Cossart et al., 2003; Buzsáki, 2010; Miller

et al., 2014a; Yuste, 2015; Carrillo-Reid et al., 2016, 2019; Marshel et al., 2019), could mediate

memory formation (Hebb, 1949) and functional brain states (Hoshiba et al., 2017). In the

cerebral cortex, these coactive states arise from reverberating patterns of multineural activity,

generated by recurrent connectivity, and can occur spontaneously (Churchland and Sejnowski,

1992; Lorente de Nó, 1933). In mouse visual cortex, cortical ensembles are causally related to

memory storage and percepts (Carrillo-Reid et al., 2019; Marshel et al., 2019; Miller et al.,

2014). Moreover, disorganized neuronal ensembles are observed in mouse models of

schizophrenia (Hamm et al., 2017).

A hypothesis of how a neuronal ensemble (i.e., a neuronal assembly) could be formed was

initially proposed by Hebb in 1949. Assemblies could be formed by synaptic plasticity among

coactive neurons, as a consequence of the Hebbian rule, whereby the "persistent and repeated

activation of connected neurons induce metabolic changes or growth of processes in one or both

cells that strengthen those connections". One synaptic mechanism that fulfills Hebb's rule is

long-term potentiation (LTP), a long-lasting increase in synaptic efficacy after a high frequency

burst electrical stimulation, and postulated to be the cellular correlate of memory and learning

(Bliss and Gardner-Medwin, 1973). With LTP, increased synaptic weights enhances preferential

connectivity that leads to an increased flow of recurrent activity within ensembles (Hoshiba et

al., 2017). Consistent with this, synchronous stimulation of groups of neurons 50 to 100 times,

using *in vivo* two-photon optogenetics, bound them together into an "imprinted" ensemble that

became spontaneously coactive after the optogenetic imprinting (Carrillo-Reid et al., 2016).

However, no direct evidence has yet demonstrated that the formation of ensembles depends on

changes in synaptic connections.

As an alternative hypothesis, recent experiments have revealed widespread activity-dependent

cell-intrinsic mechanisms in cortical neurons, suggesting that the formation and stability of an

ensemble could be due to cell-autonomous intrinsic mechanisms, such as increases in neuronal

excitability (Ryan et al., 2015; Titley et al., 2017; Gill et al, 2020, Grassellil et al, 2020). This

hypothesis is consistent with experiments where specific patterns of activity in the hippocampus

recalls stored memories, under protein synthesis inhibitors, which should prevent LTP (Ryan et

al., 2015; Tonegawa et al., 2015). Moreover, individual Purkinje cells can acquire and represent

temporal pattern of inputs in a cell intrinsic manner, without synaptic plasticity (Johansson et al.,

2014). Thus, ensemble consolidation could be due to changes excitability state of the neurons,

instead of, or in addition to, reinforcement of its synapses (Disterhoft and Oh, 2006; Pignatelli et

al., 2019).

To explore mechanisms of ensemble formation, we optogenetic and electrically stimulated

neurons in mouse neocortical slices, replicating *in vitro* the optogenetic protocol to imprint

ensembles *in vivo* (Carrillo-Reid et al., 2016). We then used whole-cell and perforated patch

recordings, from individual neurons and pairs, to characterize electrophysiologically the

stimulated neurons. We find moderate biphasic changes in synaptic properties, in the midst of

major and generalized increases in cell-intrinsic excitability, that can, at least partly, explain the

synaptic changes observed. We conclude that changes in excitability play a major role in

ensemble formation.

Materials and Methods

All procedures were performed by following the U.S. National Institutes of Health and Columbia

University Institutional Animal Care and Use Committee guidelines (IACUC, Protocol #AC-

AAAV3464). Experiments were carried out on C57BL/6 transgenic mice (Vglut1-Cre, Jackson

Laboratories; RRID:IMSR_JAX:00064) and electroporated mice CD-1 (Charles River) of both

sexes at postnatal 4-8 weeks-old. Animals were housed on a 12h light-dark cycle with food and

water ad libitum.

Viral injection: After 1 month postnatal, animals were anesthetized with 2% isoflurane on a

head-fixed stereotactic apparatus. After sterilizing the incision site, the skin was opened and,

using an FG 1/4 dental drill, a small hole goes thin in the skull over the visual primary cortex

(2.5 mm lateral and 0.3 mm anterior from the lambda, 200 μm from pia). We injected 300 nL of

the virus at a rate of 30-40 nL/s, using a microsyringe pump (Micro 4), and a Hamilton (7653-

01) glass pipette. Once the pipette was placed, we waited 5 min before and after the virus

injection and closed the scalp with sutures. Viruses injected were: AAVDJ-CaMKIIa-

C1V1(E162T)-TS-p2A-EYFP-WPRE (Stanford University Gene Vector and Virus Core). Mice

were used for electrophysiology experiments 2–3 weeks postinjection. These mice were used

only to measure evoked EPSC of optogenetically stimulated neurons.

***In utero* cortical electroporation:** Pregnant mice (CD-1, Charles River) were placed in a sterile

environment following (dal Maschio et al., 2012). We performed a laparotomy, under 2%

isoflurane anesthesia, and injected (0.5 mg/µL) of pCAG ChroMe-mRuby-ST (Mardinly et al.,

2018), ~1µL per mouse with a glass pipette, in the left lateral ventricle of embryonic day 17

pups. After that, we placed the electrodes on the head (5 mm diameter platinum disk electrodes,

(Nepa Gene #CUY650P5) to electroporate with a set of three electrical pulses for pore formation

(50V; duration, 10 ms; intervals, 50ms) and then 3 electrical pulses for transferring the plasmid

(8V; duration, 10 ms; interval, 50 ms). Finally, the embryos were returned to the abdominal

cavity and left for their normal development until postnatal day 30.

Brain slices: Mice were anesthetized with ketamine/xylazine, after transcardial perfusion

procedure and posterior cervical dislocation, to obtain brain sagittal slices as described (Ting et

al., 2018). Brains were quickly dissected and cooled in a continuously gassed (95% O_2 and 5%

CO_2) icy 194 Sucrose, 30 NaCl, 2.6 KCl, 26 $NaHCO_3$, 1.2 NaH_2PO4, 10 glucose, 2 $MgSO_4$, and

0.2 $CaCl_2$ Titrate pH to 7.3–7.4 with HCl. 300 µm thick sagittal slices were cut on a Leica

VT1200 S vibratome (Leica Biosystems) and allowed to recover for 1 hr at 34°C aCSF solution

(in mM): 195 NaCl, 2.5 KCl, 1.2 NaH_2PO4, 26 $NaHCO_3$, 12 glucose, 1 thiourea, 1 Na-ascorbate,

1 Na-pyruvate, 2 $CaCl_2$, and 1 $MgSO_4$. Titrate pH to 7.3–7.4. Finally, slices were transferred to

room temperature.

Patch-Clamp Recordings: Brain slices were carefully placed in the recording chamber on an

upright microscope (Olympus, BX50WI), and continuously perfused with gassed (95% O2 and

5% CO2) aCSF at 5 ml/min. Electrophysiological recordings were performed at 30 degrees,

using patch pipettes of borosilicate glass (World Precision Instruments). Pipettes were pulled

with a micropipette puller (DMZ-Universal puller) to a 5 – 6 MOhm as final resistances.

For whole-cell and perforated patch-clamp, patch pipettes were filled with intracellular solution

containing the following (in mM): 130 K-gluconate, 20 KCl, 10 HEPES, 10 Na_2-

phosphocreatine, 4 Mg-ATP, 0.03 Na_2-GTP, and titrated to pH 7.3 with KOH. Gramicidin

(sigma Aldrich) was dissolved in dimethyl sulfoxide (1mg /10 µl DMSO) and then dissolved

with intracellular solution ~20 ng/ml. To allow membrane sealing, the tip of the pipette was first

filled with clean solution and then the pipette was back-filled with a gramicidin-containing

solution.

In each case, the patch pipettes were placed in contact with the cell with a MPC-200

micromanipulator (Sutter Instrument). Stimulation and data acquisition were sampled at 10 kHz

and low pass filtered at 4kHz using a Multiclamp 700B amplifier (Molecular Devices) and Im-

Patch© open-access software http://impatch.ifc.unam.mx. Recordings were discontinued when

leaking current was > 25 pA or seal resistance < 1 GΩ. Recordings were analyzed with custom

routines in MATLAB.

Imaging. To identify target brain region, we used an upright microscope with a 4X/0.10 NA air

objective (Olympus) before switching to 60X/0.90 W NA water immersion objective to confirm

ChroME or C1V1 opsin expression on target cells. For ChroME, the mRuby3 fluorescent

reporter was excited with monochromatic light transmitted through a fiber optic into the

microscope (Olympus 100 w high-pressure mercury burner model BH2RFLT3). Emitted

fluorescence was band-passed with an Olympus U-49006 ET CY5 filter set: 620/60 excitation

filter, 520 dichroic mirrors, and 700/75 emission filter. For C1V1, the EYFP fluorescent reporter

was visualized via a 500/24 ex, 520 dichroic mirrors, and a 542/27 em (Semrock). Fluorescence

images were acquired (50 ms exposure; 10 Hz) using a camera (Orca-ER C4742-95,

Hamamatsu) and shutter (UNIBLITZ model VCM-D1) controlled by HCImage software

(Hamamatsu). We visualized individual neurons' fluorescence deep (100 μm) into the brain slice

in layer 2/3, evidence of C1V1 or mRuby3 ChroMe expression.

Optogenetic stimulation. Optogenetic stimulation of ChroMe opsin was performed with a 470

nm fiber-coupled LED (M470F1, Thorlabs), fiber optic cannula (M79L01, Thorlabs), and LED

driver (M00329012, Thorlabs). Optogenetic stimulation of C1V1 was performed using a 617 nm

fiber-coupled LED (M617F2, Thorlabs), fiber optic cannula (CFM14L10, Thorlabs), and LED

driver (DC2200, Thorlabs). The optogenetic stimulation protocol consisted of trains of 10 Hz, 5

ms light pulses for 4 s followed by a 10 s rest, we repeated the same stimulus during 15 to 80

min to mimic stimulation conditions previously used *in vivo* experiments (Carrillo-Reid et al.,

2016) to co-stimulate neurons.

Only neurons that exhibited action potentials in response to each LED pulse were included in the

data analysis of optogenetically stimulated neurons. LED intensity applied throughout the

protocol was the lowest necessary for inducing action potentials. For optogenetically stimulated

neurons, parameters were monitored before and 15 to 80 min after the stimulation protocol.

Optogenetic stimulation protocol was stopped every 15 or 20 min to monitor parameters, and

then restarted. For unstimulated neurons, the same parameters were monitored at the beginning

and then every 15 – 20 min for 80 min approximately.

We observed that neurons expressing opsin often exhibited small alterations in intrinsic

properties. Specifically, the firing rate was higher in neurons with opsins than in control cells,

and the membrane resistance and membrane potential become higher and more variable than

neurons without opsin expression.

Experimental design. To co-activate neurons (Carrillo-Reid et al., 2016), we delivered C1V1 or

ChroME opsin in pyramidal neurons (Mardinly et al., 2018). No significant differences were

found in experiments with both opsins and data were pooled together. We localized layer 2/3

pyramidal neurons in primary visual cortex of brain slices expressing opsin by virus injection

and *in utero* electroporation (Figure 1A). To test whether the co-stimulation of connected

neurons reinforced or created new connections (Carrillo-Reid et al., 2016), we examined pairs of

connected and unconnected neurons (Figure 1B-C). We recorded and characterized evoked

excitatory postsynaptic currents and potentials (EPSCs and EPSPs). We also recorded passive

and active electrical properties of neurons using perforated patch-clamp recording. After

measuring these parameters in control condition, we began the optogenetic or electrical

stimulation protocol (Figure 1D and E). For optogenetic stimulation, action potentials were

generated by trains of 10 Hz, 5 ms light pulses for 4 s followed by a 10 s rest (Figure 1D). For

electrical stimulation protocol, action potentials were generated with 10 Hz trains, 5-10 ms

depolarizing currents pulses 400 to 600 pA for 4 s followed by a 10 s rest (Figure 1E). During

these experiments, we noticed that the time after optogenetic or electrical stimulation was

important, immediately after stimulation, the membrane activity of stimulated neurons became

unstable, particularly during the first 3 min after stimulation. After that, membrane potential

recovered. Thus, we evaluated membrane parameters 3-5 min post-stimulation and synaptic

parameters 10 min post-stimulation.

Statistical Analysis. Statistical details are showing in each figure legends. Group data are

expressed as median ± sem. Wilcoxon and Mann-Whitney test were used for nonparametric

analysis. Differences between two groups were considered significant when $*p < 0.05$, $**p <$

0.01 and $***p < 0.001$. All statistics were performed using statistical functions in MATLAB.

Results

Spontaneous activity becomes correlated in simultaneously stimulated neurons.

A neuronal ensemble can be defined as a group of neurons with correlated spontaneous and evoked activity (Buzsáki, 2010; Cossart et al., 2003; Ikegaya, 2004; Sasaki et al., 2007; Shepherd, 2010; Stringer et al., 2019; Yuste, 2015). Previous results have demonstrated that optogenetic coactivation of neurons can generate new ensembles *in vivo* (Carrillo-Reid et al., 2016). To explore the cellular and circuit mechanisms underlying this phenomenon, we studied the electrophysiological mechanisms that led to building of imprinted ensembles, replicating *in vitro* with the activation protocols that generates ensembles *in vivo,* optogenetic or electrical stimulation. Experiments were performed in layer 2/3 pyramidal neurons expressing the opsin ChroME in slices from primary visual cortex of adult mice of both sexes (Figure 1A).

The result of optogenetic imprinting *in vivo* was an increase in the correlation of spontaneous activity of stimulated neurons, indicative of the formation of a new ensemble (Carrillo-Reid et al., 2016). To explore if our *in vitro* stimulation protocols were effective in generating ensembles, we used paired recordings to explore whether simultaneous and repeated optogenetic or electrical activation neurons induced changes in the correlation of spontaneous activity. We recorded pairs of closely located neurons and found that 25% of recorded pairs at distances of 20 - 30 µm, were monosynaptically connected (7/28 pairs), consistent with previous reports (Markram et al., 1997; Holmgren et al., 2003; Song et al., 2005; Lefort et al., 2009; Ko et al., 2011; Levy and Reyes, 2012; Cossell et al., 2015) (Figure 1A-C). In agreement with previous studies, we found higher values of correlation of spontaneous activity between connected neuronal pairs (0.31 ± 0.16; n = 7) than between unconnected neuronal pairs (-0.04 ± 0.37; n =

21; p = 0.049 by Mann-Whitney test, Figure 1F) (Cossell et al., 2015; Ko et al., 2011; Yoshimura

et al., 2005). However, ~50% (10/21) of the unconnected neuronal pairs still had relatively high

correlations, comparable to connected neuronal pairs (0.32 ± 0.18; n = 9; all unconnected pairs

with correlation > zero; Figure 1B). Notably, after optogenetic or electrical stimulation (Figure

1D and E), the correlation of spontaneous activity between neurons increased significantly, in

both connected and unconnected pairs (0.02 ± 0.06 to 0.2 ± 0.05; n = 13 pairs, 2 connected, 11

unconnected; p = 0.049 by Wilcoxon; Figure 1G). No significant differences in correlations were

found in unstimulated pairs (0.05 ± 0.1 to 0.03 ± 0.1, n = 11 pairs, p = 0.7 Wilcoxon, Figure 1H).

We conclude that optogenetic and electrical stimulation increase the correlation of spontaneous

activity among neurons, a hallmark of neuronal ensemble formation.

Figure 1. Optogenetic and electrical stimulation increases correlations of spontaneous activity

(**A**) Example image of Ruby3 reporter fluorescent of ST-ChroMe opsin expression in L2/3 pyramidal neurons in primary visual cortex. Brain slices were obtained from *in-utero* electroporated mice. Scale bar: 40 μM. (**B**) I*n-vitro* paired recording for evaluating monosynaptic connectivity between neurons. Perforated patch-clamp recording of presynaptic action potentials elicited by 100 pA of current injections (500 ms), followed by identification of

a monosynaptic connection, generating postsynaptic potentials time-locked to presynaptic spikes.

(**C**) Spontaneous activity recording of two pairs. Top: current-clamp spontaneous activity of

unconnected neurons (correlation coefficient = 0.33). Bottom: Representative paired recording

from neurons that were connected bi-directionally (correlation coefficient = 0.49). (**D**)

Experimental design of optogenetic stimulation protocol. Perforated current-clamp recording of

two neurons with opsin expression in blue: 1 to 30 min of 10 Hz train, 5 ms light pulses for 4 s

followed by 10 s of rest. (**E**) Experimental design of electrical stimulation protocol. Perforated

current-clamp recording of two neurons without opsin expression in gray: 1 to 30 min of 10 Hz

train, 5 ms current pulses for 4 s followed by 10 s of rest. (**F**) Correlation coefficients of

connected and no connected neurons. Correlation coefficients were calculated for the first 3

minutes simultaneous recording in each pair; triangles represent correlation values of pairs

showed in C. Correlation coefficients of connected (n = 7 pairs) vs. no connected (n = 21 pairs)

showed significant differences p = 0.049 by Mann Whitney; 15 mice. (**G**) Correlation between

pairs increased after continued co-stimulation. The graph shows increase in the correlation

coefficient before and after 20 to 30 min of continuous optogenetic (blue) or electrical

stimulation (red) (p = 0.004 by Wilcoxon test; n = 13 pairs, 12 mice). (**H**) Correlation before and

after 20 to 30 min without stimulation (gray) (p = 0.7 by Wilcoxon; n = 11 pairs, 8 mice).

Triangles in the graph represent synaptically connected pairs.

Lack of significant plasticity of synaptic currents after optogenetic stimulation

The creation of ensembles *in vivo* after optogenetic stimulation (Carrillo-Reid et al., 2016),

suggested that ensembles could be built by Hebbian synaptic plasticity, by potentiation of

existing synapses of coactivated neurons. To explore this, we performed dual whole-cell and

perforated patch-clamp recordings of connected neurons, and examined whether persistent and

repeated optogenetic or electrical activation induced synaptic plasticity (Cossell et al., 2015;

Hebb, 1949). Action potentials were induced in the presynaptic neuron with depolarizing

currents (10 pulses; 400 to 600 pA, 5 ms, 20 Hz trains). For measuring synaptic currents,

postsynaptic neurons were recorded in voltage-clamp, in cero current injection. We classified

connections as monosynaptic if EPSC latency was 2 ± 1 ms, relative to the presynaptic spike

(Figure 2A; n = 7, 5 mice). To normalize EPSC amplitudes, postsynaptic current values in the

action potential start time were considered as baseline. Current values for the first EPSC ($EPSC_1$)

during the stimulation trains, including failures, were averaged. A connection was defined if the

average evoked $EPSC_1$ had an amplitude larger than 1.5 S.D. of the noise. Control recordings

were performed with unconnected neurons (n = 21 pairs), and the optogenetic protocol did no

generate any new detectable synaptic connections.

In pairs of connected neurons, we examined if optogenetic stimulation induced changes in

monosynaptic currents, pooling data from whole-cell (n = 5) and perforated patch-clamp (n = 2)

recordings. The average amplitude of monosynaptic ESPCs in control condition agreed with

previous reports (Yoshimura et al., 2005; Feldmeyer et al., 2006; Morishima et al., 2011;

Sempere-Ferràndez et al., 2019). Average $EPSC_1$ peak currents of optogenetically stimulated

neurons showed no significant differences, albeit with a trend towards amplitude depression right

after the stimulation, followed with a tendency towards potentiation, after a 20 minutes rest

period (before: 13 ± 7 pA; after: 9 ± 7 pA; p = 0.2 Wilcoxon; 6 mice; Figure 2B, C). Control

$EPSC_1$ amplitudes of unstimulated neurons also did not show significant difference before ($13 \pm$

2 pA) and after 30 min (12 ± 2 pA; p = 0.1 Wilcoxon; n = 11, whole-cell recordings, n = 7;

perforated patch-clamp, n = 5; 6 mice). To evaluate changes in short-term plasticity, we

calculated the paired-pulse ratio (PPR), i.e. the ratio of amplitudes of the second to the first

EPSC in the train ($EPSC_2/EPSC_1$). We integrated data from whole-cell (n = 5) and perforated

patch-clamp recordings (n = 2), and found that PPR did not significantly change with the

stimulation (before: 0.95 ± 0.09 PPR; after: 0.7 ± 0.1 PPR; p = 0.4 Wilcoxon; n = 7, 5 mice;

Figure 2D). Similarly, control unstimulated neurons in whole-cell (n = 5) and perforated patch-

clamp recordings (n = 7) did not show significant changes in PPR before (1.08 ± 0.12) and after

30 min (1. 06 ± 0.12; p = 0.7 Wilcoxon; n = 12; 7 mice).

Biphasic plasticity of synaptic currents after electrical stimulation

We carried out similar experiments with electrical stimulation of connected pairs of neurons,

thus avoiding possible side effects caused by opsin expression (see Methods) or the optogenetic

stimulation of other neurons expressing opsin in the field of view. Using perforated patch-clamp,

we recorded and electrically stimulated pairs of connected pyramidal neurons. After electrical

stimulation, $EPSC_1$ amplitudes decreased significantly (from 14 ± 2 pA to 8 ± 1 pA; p = 0.016 by

Wilcoxon; n = 7; 6 mice; Figure 2E). After pausing the stimulation for 20 min, the amplitude of

evoked ESPCs recovered, with a moderate potentiation. Specifically, for electrically stimulated

neurons, $EPSC_1$ amplitude recovered significantly from 8 ± 1 to 20 ± 1 pA after 20 min of post-

stimulation rest (p = 0.015 by Wilcoxon; n = 7). The amplitude after the recovery period was

moderately larger than the EPSCs amplitude before stimulation (14 ± 2 to 20 ± 1 pA; p = 0.015

by Wilcoxon; Figure 2E). These results are consistent with the trends found in pairs of

optogenetic stimulated neurons, where $EPSC_1$ amplitude after a post-stimulation recovery period

increased from 9 ± 7 pA to 19 ± 3 pA, although no statistical comparison was possible (n = 2;

Figure 2C). In contrast, control unstimulated neurons did not show a significant change (before:

13 ± 2 pA; after 30 min: 12 ± 2 pA; p = 0.6 by Wilcoxon; n = 8; 6 mice). We also measured the

PPR of electrical stimulated neurons before and after electrical stimulation, finding no significant

differences (before: 1.3 ± 0.15 PPR; after: 1.03 ± 0.18 PPR; p = 0.7 Wilcoxon; n = 7, 5 mice).

No difference in PPR was observed before and after 20 min post-stimulation rest (1.3 ± 0.13; p =

0.1 by Wilcoxon; Figure 2E). Similarly, the PPR of unstimulated neurons did not change before

(1.4 ± 0.1) and after 30 min without stimulation (1.3 ± 0.15; p = 0.5 by Wilcoxon; n = 7; 5 mice)

(Figure 2F).

Biphasic plasticity of synaptic potentials after stimulation

To further explore potential postsynaptic mechanisms of ensemble formation, in a parallel set of

experiments we used current-clamp to examine postsynaptic potentials (EPSPs) in connected

pairs, comparing evoked EPSP amplitudes before and after optogenetic and electrical stimulation

(Figure 2G). We combined data from optogenetically (n = 2, blue) and electrically-stimulated

pairs (n = 7, gray). In good agreement with the voltage-clamp data, the average peak amplitude

of the first EPSP ($EPSP_1$) significantly decreased after stimulation (0.43 ± 0.2 mV to 0.28 ± 0.2

mV; p = 0.004 by Wilcoxon; n = 9; 7 mice; Figure 2H). As with EPSC measurements, EPSPs

recovered and became potentiated after 20 min of post-stimulation rest to 0.8 ± 0.2 mV (recovery

compared with before: p = 0.008; and after stimulation: p = 0.004 by Wilcoxon). Control

unstimulated neurons did not show significant differences in EPSP amplitude before (0.44 ± 0.2

mV) and after 30 min (0.42 ± 0.2 mV; p = 0.7 by Wilcoxon; n = 10; 6 mice).

In conclusion, paired recordings revealed that the coactivation of neurons generates a moderate

biphasic synaptic plasticity, with an initial small depression, followed by a rebound potentiation,

after several minutes of inactivity.

Figure 2. Effect of optogenetic and electrical stimulation in evoked EPSCs and EPSPs.

(**A**) Representative paired whole-cell recording of synaptically connected neurons. Top: current-

clamp recording of presynaptic action potentials in response to 10 current injections (5 ms each;

20 Hz). Bottom: voltage-clamp recording of evoked EPSC before (black) and after (blue) 30 min

of optogenetic stimulation. Each trace is average of 30 successive responses evoked by

presynaptic current injection. (**B**) Representative paired recording of evoked EPSCs (perforated

patch-clamp). Top: current-clamp recording of presynaptic action potentials induced by the

positive current steps of the I-V curve (20 - 120 pA). Bottom: voltage-clamp recording of evoked

EPSCs before and after optogenetic stimulation and after 20 min of rest post-stimulation.

Thickest line is the average of successive responses to the first presynaptic action potentials for

every current step of the I-V curve. (**C**) $EPSC_1$ amplitudes without significant changes before

and after optogenetic stimulation. Left: mean $EPSC_1$ amplitude before and after optogenetic

stimulation. Open dots correspond to whole-cell recordings, n = 5; fill dots correspond to

perforated patch-clamp recordings; n = 2. Medians and blue box plot: p = 0.2 by Wilcoxon; n = 7

neurons, 5 mice). Right: Unstimulated neurons, n = 7 neurons with whole-cell and n = 5 neurons

with perforated patch-clamp. Medians and white box plots; p = 0.1 by Wilcoxon; n = 12

postsynaptic neurons, 8 mice. (**D**) Comparison of mean Paired Pulse Ratio (PPR). Left: Mean

PPRs before and after optogenetic stimulation and after 20 min of rest post-stimulation. PPR

show no significant differences between before and after, p = 0.4; Wilcoxon; n = 7 cells, 5 mice.

Right: PPR of unstimulated neurons did not show significant differences, p = 0.6 by Wilcoxon, n

= 12 cells, 8 mice). (**E**) The amplitude of evoked $EPSC_1$ after electrical stimulation. Comparable

to optogenetic stimulation, electrical stimulation protocol consisted of 30 min of 10 Hz train, 5

ms current pulses for 4 s followed by 10 s of rest in two neurons simultaneously, this protocol

was done in brain slices without opsin expression and using only perforated patch-clamp. Left:

the average $EPSC_1$ amplitude decreased after electrical stimulation (p = 0.015 by Wilcoxon; n =

7 postsynaptic neurons, 5 mice). $EPSC_1$ amplitudes were recovered with a moderate increase,

after 20 min of rest post-stimulation regarding after condition (p = 0.015), and also compared to

before stimulation (p = 0.015). Right: $EPSC_1$ amplitude of unstimulated neurons (p = 0.7 by

Wilcoxon; n = 8 neurons, 5 mice). (**F**) PPR remains unchanged: before and after electrical

stimulation (p = 0.7 Wilcoxon; n = 8 cells, 5 mice); also, after electrical stimulation and rest

post-stimulation (p = 0.2); comparison before and rest post-stimulation did not show differences

either (p = 0.1), as well as PPR of unstimulated neurons; p = 0.6 by Wilcoxon, n = 12 cells, 8

mice. (**G**) Representative paired recording of evoked EPSPs (perforated patch-clamp). Top:

current-clamp recording of presynaptic action potentials. Bottom: current-clamp recording of

evoked EPSPs before, after optogenetic stimulation and, after 20 min of rest post-stimulation.

Thickest lines are the average of the successive responses to presynaptic action potentials. (**H**)

Comparison between mean $EPSP_1$ amplitude after optogenetic stimulation (blue dots; n = 2) or

after electrical stimulation (gray dots; n = 7). Left: average $EPSP_1$ amplitude decreased after

stimulation protocol (p = 0.004 by Wilcoxon; n = 9 neurons, 5 mice). $EPSP_1$ amplitudes were

recovered with a moderate increase after 20 min of rest post-stimulation compared to after (p =

0.004 by Wilcoxon) and also, compared to before stimulation protocol (p = 0.008 by Wilcoxon).

Right: $EPSP_1$ amplitude of unstimulated neurons (p = 0.7 by Wilcoxon; n = 8 postsynaptic

neurons, 5 mice).

Biphasic synaptic plasticity in local circuit after optogenetic stimulation

To explore synaptic plasticity in the local circuit of optogenetically stimulated neurons, we also

performed voltage-clamp and current-clamp recordings of evoked EPSCs and EPSPs in neurons

without opsin expression but located in the same field of view (Figure 3A). Assuming a

photostimulated area by the LED of 150 µM in radius, we estimated that the typical number of

superficial neurons with opsin expression, and potentially activated by the light, was ~20 cells.

Thus, optical stimulation could lead to a significant increase in the activity of the local circuit. To

monitor the effect of the stimulation on the local circuit, we used perforated patch-clamp from

non-expressing pyramidal neurons, to measure EPSCs or EPSPs evoked by the LED, in both

voltage and current-clamp. Light-evoked EPSCs or EPSPs were analyzed if the latency was >2

ms relative to LED pulse onset (Figure 3A; n = 7, 5 mice). In non-expressing cells, population

light-evoked $EPSC_1$ (22 ± 3 pA) and $EPSP_1$ amplitudes (2 ± 0.1 mV) were significantly higher

than unitary evoked $EPSC_1$ (14 ± 2 pA; p = 0.04 by Mann-Whitney) or $EPSP_1$ (0.43 ± 0.2 mV; p

= 0.0006 by Mann-Whitney) found in connected pairs. This is likely because LED stimulated

several neurons in the field simultaneously, generating a compound EPSC and EPSP.

After the optogenetic stimulation protocol (10 Hz of 5 ms pulse for 4s every 10s; 30 min), the

average light-evoked $EPSC_1$ showed a significant diminution in amplitude (22 ± 3 to 11 ± 1 pA;

p = 0.015 by Wilcoxon; n = 7; 6 mice; Figure 3B), and a recovery with potentiation after 20 min

of rest post-stimulation (40 ± 3 pA; p = 0.015, comparing with before; p = 0.015, comparing

after stimulation by Wilcoxon). Similarly, light-evoked EPSPs showed a significant decrease

after 30 min optogenetic stimulation (2 ± 0.1 mV to 1.3 ± 0.2 mV; p = 0.015 by Wilcoxon; n = 7;

6 mice; Figure 3C), and also recovered with a moderate increase in amplitude after 20 min of rest

post-stimulation (2.5 ± 0.3 mV; p = 0.015, comparing with before; p = 0.015, comparing with

after optogenetic stimulation by Wilcoxon). These results, in excellent agreement with the

connected pair data, reveals that the biphasic synaptic plasticity is also found in unstimulated

neurons, thus suggesting that it is a circuit-wide phenomenon.

Figure 3. Effect of optogenetic stimulation in the synaptic plasticity of the local circuit

(**A**) Activation of expressing opsin neuronal population in the slice by LED pulses allowed the

detection of evoked EPSCs and EPSPs in non-expressing pyramidal neurons. Voltage-clamp

recording (top) and current-clamp recording (bottom) of a representative non-expressing neuron.

Evoked EPSCs and EPSPs before (gray); after 30 min of optogenetic stimulation protocol

(green), and 20 min post stimulation (dark green). (**B**) The amplitude of population light-evoked

EPSC$_1$ (left) and (**C**) light-evoked EPSP$_1$ (right) decreased after 30 min of optogenetic

stimulation (p = 0.01 and p = 0.01 respectively, by Wilcoxon; n = 7 postsynaptic neurons; 7

mice), they were recovered with a moderate increase after 20 min of rest (p = 0.01 and p = 0.01

respectively).

Increased of spontaneous synaptic inputs after optogenetic and electrical stimulation

To further explore the effect of the optogenetic and electrical stimulation on population activity,

we measured spontaneous EPSP from recorded neurons, and compared them before and after

stimulation (Figure 4A). To measure spontaneous EPSPs, we low-pass filtered current-clamp

recordings, subtracting membrane potential oscillations, to obtain comparable baselines of EPSP

amplitudes (Juárez-Vidales et al., 2021). We detected EPSPs with amplitudes between 0.3 mV

and 10 mV and analyzed changes in the frequency of spontaneous EPSP before and after

optogenetic stimulation (Figure 4A-B). The frequency of spontaneous EPSP significant

increased from 90 ± 9 EPSPs/min before to 127 ± 19 after optogenetic stimulation (p = 0.004 by

Wilcoxon; n = 10; 5 mice) (Figure 4D left). We also observed increases in EPSP amplitude (0.38

± 0.01 to 0.41 ± 0.02 mV; p = 0.04) (Figure 4E left). In control recordings (Figure 4D-E right),

non-expressing pyramidal neurons showed no statistically differences in the frequency of

spontaneous EPSPs (93 ± 25 to 64 ± 18; p = 0.06), or in EPSP amplitude (0.37 ± 0.004 to 0.37 ±

0.004 mV; p = 0.5; n = 11; 8 mice).

Similarly, electrical stimulated neurons showed a significant increase in the frequency of

spontaneous EPSPs from 80 ± 12 to 118 ± 17 EPSPs/min (p = 0.001 by Wilcoxon; n = 13; 8

mice) and EPSP amplitude (0.37 ± 0.03 to 0.40 ± 0.03 mV; p = 0.03) (Figure 3F-G left). In

unstimulated neurons (Figure 3F-G right), we found no statistically difference in the frequency

of spontaneous EPSPs (98 ± 18 to 93 ± 15; p = 0.4), or any significant differences in EPSP

amplitude (0.37 ± 0.01 to 0.38 ± 0.01 mV; p = 0.4; n = 11; 8 mice).

These results reveal major changes in spontaneous synaptic inputs after optogenetic and

electrical stimulation. The increase in spontaneous EPSPs amplitude could partly explain their

increased frequencies, as smaller EPSPs will become detectable. Importantly, since increases in

spontaneous EPSP frequency and amplitude also occur even after electrical stimulation of

individual neurons, our results demonstrate that neurons become more sensitive to synaptic

inputs after stimulation, in a cell intrinsic manner.

Figure 4. Optogenetic and electrical stimulation amplifies spontaneous EPSPs.

(**A**) Representative perforated patch-clamp recording of a neuron in current-clamp. Top: spontaneous EPSPs of a neuron before and after optogenetic stimulation protocol. Bottom: Section in the top trace shows spontaneous EPSPs amplitude > 0.3 mV that was identified before (gray) and after (blue) optogenetic stimulation protocol in a one-minute recording. Arrow shows a tentative EPSP < 0.3 mV that was not detected by the threshold (0.3 mV). (**B**) Frequency histogram of spontaneous EPSPs amplitudes shows that after optogenetic stimulation, the number of events increased as well as the number of events with higher amplitudes. (**C**) Frequency histogram of spontaneous EPSPs amplitudes shows that the number of events of a non-expressing pyramidal neuron remained unchanged before and after optogenetic stimulation of expressing neuronal population. (**D**) The number of spontaneous EPSPs increased in optogenetically stimulated neurons but not in non-expressing neurons. Left: Optogenetic stimulation of neurons: blue median and box plot; p = 0.004 by Wilcoxon; 8 mice. Right: Number of spontaneous EPSPs of non-expressing pyramidal neurons before and after

optogenetic stimulation protocol: green median and box plot; p = 0.06 by Wilcoxon; 5 mice. (**E**)

The amplitude of spontaneous EPSPs increased in optogenetically stimulated neurons but not in

non-expressing neurons. Right: Optogenetic stimulated neurons: blue median and box plot; p =

0.04 by Wilcoxon. Left: Non-expressing neurons: green median and box plot; p = 0.5 by

Wilcoxon. (**F**) The number of spontaneous EPSPs increased after electrical stimulation.

Electrical stimulated neurons: red medians and box plot; p = 0.0012 by Wilcoxon; 9 mice.

Unstimulated neurons: white medians and box plot; p = 0.4 by Wilcoxon; 9 mice. (**G**) The

amplitude of spontaneous EPSPs increased after electrical stimulation. Electrical stimulated

neurons: red median and box plot; p = 0.003 by Wilcoxon. Unstimulated neurons: white medians

and box plots; p = 0.4 by Wilcoxon.

Optogenetic stimulation increases neuronal excitability

Our results demonstrated that optogenetic and electrical stimulation, even of individual neurons,

could induce changes in intrinsic membrane properties, or cellular excitability. This is in

agreement with an extensive literature that has described non-synaptic cellular plasticity in a

variety of experimental preparations (Abraham et al., 2019; Malik and Chattarji, 2012; Marder

and Goaillard, 2006; Paz et al., 2009; Ryan et al., 2015; Titley et al., 2017; Xu, 2005; Yang and

Santamaria, 2016). To explore these mechanisms, we evaluated active and passive intrinsic

electrophysiological parameters such as current injection-dependent firing, frequency, membrane

resistance, membrane potential, and firing threshold. We recorded individual neurons with

perforated patch-clamp before and after optogenetic and electrical stimulation (Figure 5).

We first measured membrane potential dependency to current injections with I-V plots in

pyramidal neurons with opsin expression. To do this, we kept the membrane potential at -70 mV

and applied series of 500 ms current injections ranging from -100 pA to 160 pA, with 20 pA

alternating positive and negative steps (black traces; Fig. 5A). We first observed a significant

increase in the evoked number of spikes after optogenetic stimulation (blue traces; 5 ± 1 and $6 \pm$

1 spikes; $p = 0.015$ by Wilcoxon; $n = 12$; 8 mice; Figure 5D-a). These increases in the firing rate

were not observed in unstimulated neurons (5 ± 1 and 5 ± 1; $p = 1$ by Wilcoxon; $n = 7$; 6 mice;

Figure 5D-b). We also noticed that neurons expressing opsin exhibited a small alteration in their

intrinsic properties. In particular, the firing rate became higher than the neurons without opsins in

control condition (5 ± 1 firing rate of neurons with opsin expression, 3.5 ± 1 neurons without

opsin, but this difference was not significant $p = 0.1$ by Mann-Whitney; $n = 7$; 6 mice).

Neurons needed at least 15 - 20 min of optogenetic or electrical stimulation to increase their

excitability, but, after their firing rate increased, it was stable for the 80 min duration of the

experiment (6 ± 1 to 6 ± 1 spikes; $p = 0.5$ by Wilcoxon; $n = 7$; 5 mice; Figure 5B). The long-

lasting increase in firing rate did not require continued stimulation, since after pausing

optogenetic stimulation for 30 min, it remained elevated (6 ± 1 to 6 ± 1 spikes; $p = 0.8$ by

Wilcoxon; $n = 7$; 5 mice; Figure 5C). Similar results were found for the average instantaneous

firing frequency, which increased after optogenetic stimulation (from 15 ± 3 to 17 ± 3 Hz; $p =$

0.015 by Wilcoxon; $n = 12$; 8 mice; Figure 5E-a. Unstimulated neurons did not show significant

differences (from 16 ± 3 to 16 ± 3 Hz; $p = 0.5$ by Wilcoxon; $n = 7$; 5 mice; Figure 5E-b). Finally,

to evaluate the modification of firing properties that could facilitate burst induction, we measured

the mean frequency of the two first spikes for each current step, finding a significant increase

(before: 24 ± 3 Hz; after optogenetic stimulation: 27 ± 3 Hz; $p = 0.015$ Wilcoxon; Figure 5F-a),

whereas unstimulated neurons did not show significant differences after 30 minutes of recordings

(25 ± 6 and 24 ± 6 Hz; $p = 0.2$ by Wilcoxon; $n = 7$; 5 mice; Figure 5F-b). To rule out that these

changes in firing properties were due to the illumination, we recorded neurons without opsin

expression in perforated patch-clamp, before and after optogenetic stimulation. We did not detect

any significant change in current injection firing-dependence (4 ± 1 and 4 ± 1 spikes; $p = 1$, by

Wilcoxon; $n = 7$; 5 mice), nor in the average instantaneous frequency (7 ± 2 Hz before; and 7 ± 2

Hz after; $p = 0.6$ by Wilcoxon). These data, similar to those from unstimulated neurons,

demonstrate that the LED illumination does not induce changes in cellular excitability.

Terfenadine reverts optogenetic stimulation increases neuronal excitability

These results revealed that optogenetic stimulation produced a robust increase in current-

dependent firing. To pharmacologically explore potential mechanisms underlying these changes,

we used terfenadine, a pharmacological agent that blocks currents mediated by Ether-a-go-go

Related Gene (ERG) channels, and, in neocortical pyramidal cells, prevents persistent firing and

increases in membrane resistance (Cui and Strowbridge, 2018). Indeed, the increases in firing

rate and frequencies induced by optogenetic stimulation reverted to control conditions after the

application of 10µM terfenadine (5 ± 1 and 4 ± 1 spikes; $p = 0.015$ by Wilcoxon; $n = 7$; 5 mice;

Figure 5D-c). No differences were found comparing with before stimulation and when

terfenadine was applied (4 ± 1 and 4 ± 1 spikes; $p = 0.06$ by Wilcoxon; $n = 7$; 5 mice). However,

in unstimulated neurons without opsin, terfenadine generated a reduction in firing rate (4 ± 1

before; and 3 ± 1 terfenadine $p = 0.03$ by Wilcoxon; $n = 7$; 5 mice). We concluded the increase in

excitability due to optogenetic stimulation can be reverted by terfenadine. This could be due to

blockage of baseline conductances, such as ERG channels or other targets like histamine1, kir 6,

and Kv11.1 (Carter et al., 1985).

Electrical stimulation increases neuronal excitability

After characterizing studying the changes in firing properties due to optogenetic activation, we

explored if single-cell electrical stimulation generated similar alterations. Electrical stimulation

via a patch pipette avoids possible side effects produced by the optogenetic stimulation, or by the

activation of an undetermined number of opsin-expressing neurons. Electrical stimulation of one

neuron at the time also rules out synaptic effects. In these single cell electrical stimulation

experiments, we observed similar effects as in optogenetic stimulation: increases in firing rate

(before 4 ± 1 spikes; after 5 ± 1 spikes; p = 0.015 by Wilcoxon; n = 22; 12 mice; Figure 5D-d);

increases in average frequency (8 ± 2 Hz before; 10 ± 2 Hz after; p = 0.015 by Wilcoxon; Figure

5E-d); and increases in frequency of the first two spikes (14 ± 3 Hz before; 17 ± 3 Hz after; p =

0.015 by Wilcoxon; Figure 5F-d). Unstimulated neurons did not show any changes in firing rate

(before 4 ± 1; after 4 ± 1 spikes; p = 1 by Wilcoxon; n = 10; 8 mice; Figure 5D-e); or average

frequency (9 ± 2 Hz before; and 9 ± 2 Hz after; p = 0.015 by Wilcoxon; Figure 5E-e); nor in the

first frequency (15 ± 3 Hz before; and 15 ± 3 Hz after; p = 1 by Wilcoxon; Figure 5F-e).

The consistency in results from optogenetic and electrical stimulation indicate that the increases

in cellular excitability are not due to opsin expression or optogenetic protocol, and is a property

of individual neurons that can be induced by stimulation of individual cells.

Figure 5. Neuronal excitability increases after optogenetic or electrical stimulation.

(**A**) Current dependence firing rate increases after optogenetic or electrical stimulation protocol.

This returns to basal conditions after the application of terfenadine. Representative membrane

voltage recordings with perforated patch-clamp (top) in response to 500 ms series of 20 pA

current steps (20 to 120 pA, bottom). Black traces show membrane potential in response to

current steps before stimulation. Blue traces represent membrane voltage recordings of neurons

with opsin expression and responses after 30 min of optogenetic stimulation. The green traces

show the effect of the terfenadine (10 µM) in the membrane potential post-stimulation. (**B**)

Decrease in the rheobase after stimulation is stable. Current-dependent firing rate increased in a

representative neuron after 20 min of continuous optogenetic stimulation. This firing rate did

show more changes even after 80 min of continued stimulation ($p = 0.4$ by Wilcoxon; $n = 7$

neurons, 5 mice). (**C**) Decrease in the rheobase after stimulation is long-lasting. Gradual increase

in the firing rate of a representative neuron after 15 min of optogenetic stimulation and then after

30 min. 30 min post-stimulation (rest), the neurons kept their membrane excitability increased (p

$= 0.4$ by Wilcoxon; $n = 7$ neurons, 5 mice). (**D**) Neuronal activity increase after optogenetic or

electrical stimulation. For every graph (a - e), the thicker black lines are average spikes of all

neurons before stimulation protocol. Thicker lines color for the average spikes correspond to a)

blue, optogenetic stimulation; b) blue, 30 minutes without stimulation in neurons with opsin

expression; c) green, terfenadine (10µM) effect after optogenetic stimulation; d) red, electrical

stimulation in neurons without opsin expression; e) red, 30 minutes without stimulation in

neurons without opsin expression. Data represent mean ± SEM. *$p < 0.05$ by Wilcoxon. (**E**)

Same as **D** but for average frequency for each current step. *$p < 0.05$ by Wilcoxon. (**F**) The

frequency of the first two action potentials increased for the lower current steps after stimulation.

Same as **D** but for the average of the first two action potentials frequency. *$p < 0.05$ by

Wilcoxon.

Optogenetic and electrical stimulation increase membrane resistance

An increase in excitability could be due to an increase in membrane resistance (Marder and

Goaillard, 2006). To calculate membrane resistance, we measured membrane potential changes

produced by hyperpolarizing current pulses of low intensity (from 40 to -100 pA; 500 ms

duration, applied every second), before and after optogenetic stimulation (Figure 6A). Membrane

input resistance was measured from the voltage-current (I-V) relationship (Figure 6B). We

observed a significant increase in input resistance from 278 ± 18 mΩ before to 305 ± 22 mΩ

after optogenetic stimulation (p = 0.0001 by Wilcoxon; n = 14; 8 mice; Figure 6C-a).

Meanwhile, resistances of unstimulated neurons, with opsin expression, did not show any change

before (318 ± 23 mΩ) and 30 minutes after (309 ± 24 mΩ; p = 0.9 by Wilcoxon; n = 8; 8 mice,

Figure 6C-b). These increases were reverted by the addition of terfenadine (from 172 ± 15 mΩ

after stimulation to 153 ± 16 mΩ after terfenadine; p = 0.02 by Wilcoxon; n = 9; 7 mice; Figure

6C-c). Also, the addition of terfenadine brought back the resistance to its pre-stimulation values,

158 ± 15 mΩ before stimulation to 153 ± 16 mΩ with terfenadine; p = 0.9 by Wilcoxon.

Furthermore, electrical stimulation in non-expressing neurons also induced a significant increase

in the input resistance (from 176 ± 9 to 185 ± 9 mΩ; p = 0.0003 by Wilcoxon; n = 10; 8 mice;

Figure 6C-d). Unstimulated neurons did not display changes in membrane resistance (before 171

± 11 to after 174 ± 8 mΩ; p = 0.3 by Wilcoxon; n = 10; 8 mice; Figure 6C-e). Control

experiment with non-expressing neurons did not show any alteration in membrane resistance

after optogenetic stimulation of the slices (before 176 ± 20 mΩ; after 179 ± 20 mΩ; p = 0.3 by

Wilcoxon).

It has been reported that membrane resting potential can also change through increases in

excitability, resulting in depolarization (Debanne et al., 2019; Mellor, 2002; Ross and Soltesz,

2001). To explore this, we calculated membrane resting potential as the median of spontaneous

membrane activity with zero current injection. We observed that membrane resting potential

remained stable under all condition: before and after optogenetic stimulation (-73 ± 2 to -72 ± 1

mV; p = 0.4 by Wilcoxon; n = 10; Figure 6D-a); unstimulated neurons (-73 ± 2 to -75 ± 2 mV; p

= 0.8 by Wilcoxon; n = 13; Figure 6D-b). The application of terfenadine after stimulation also

did not reveal significant changes in membrane potential (-78 ± 2 to -77 ± 2 mV; p = 0.8 by

Wilcoxon; n = 8; Figure 6D-c) and no changes before and after electrical stimulation (-80 ± 1 to -

79 ± 1 mV; p = 0.9 by Wilcoxon; n = 13; Figure 6D-d). This lack of resting membrane potential

changes during intrinsic neuronal excitability agrees with previous reports (Cudmore and

Turrigiano, 2004; Disterhoft et al., 1986; Pignatelli et al., 2019). We conclude that the increase in

excitability induced by optogenetic or electrical stimulation is associated with increases in

membrane resistance but not with membrane potential depolarization.

Figure 6. Membrane input resistance increases after optogenetic or electrical stimulation

(**A**) Membrane resistance increases after optogenetic or electrical stimulation; these returns to

control condition after terfenadine application. Representative membrane voltage recordings with

perforated patch-clamp in response 500 ms series of 20 pA current pulses from 40 to -100 pA

(bottom). Black traces show membrane potential dependence before optogenetic stimulation, and

blue traces correspond to 30 min after (**B**) I-V plots show the relation between current injection

and the membrane potential (Ohm's law: Resistance = V/I). For every graph (a - c), black dots

are the average voltage membrane response to current steps before the stimulation protocol. Blue

average voltage membrane corresponds to a) optogenetic stimulation; b) green, the effect of

terfenadine (10μM) after optogenetic stimulation; c) red, electrical stimulation in neurons

without opsin expression. Data represent mean ± SEM. (**C**) Membrane resistance increased after

optogenetic or electrical stimulation. a) Optogenetic stimulated neurons: median and blue box

plot; $p = 0.0001$ by Wilcoxon; n = 14 neurons, 9 mice. b) Unstimulated neurons with opsin

expression: median and gray box plots; $p = 0.7$ by Wilcoxon, n = 8 neurons, 5 mice. c)

Terfenadine (10μM) effect after optogenetic stimulation: median and green box plot; $p = 0.003$

by Wilcoxon; n = 8 neurons, 5 mice. d) Electrical stimulated neurons: median and red box plot; p

$= 0.00004$ by Wilcoxon; n = 22 neurons, 15 mice. e) Unstimulated neurons without opsin

expression: median and gray box plot; $p = 0.5$ by Wilcoxon; n = 11 neurons, 8 mice. (**D**) Resting

membrane potential remain without changes after stimulation. a) Optogenetic stimulated

neurons: median and blue box plot; $p = 0.4$ by Wilcoxon; n = 10 neurons. b) Unstimulated

neurons: median and gray box plots; $p = 0.8$ by Wilcoxon; n = 13 neurons. c) Terfenadine

(10μM) effect after stimulation: median and green box plot; $p = 0.8$ by Wilcoxon; n = 8 neurons.

d) Electrical stimulated neurons: median and red box plot; $p = 0.9$ by Wilcoxon; n = 13.

Lowering of action potential threshold after optogenetic and electrical stimulation

The changes in frequencies of the first two spikes that we observed could also be due to changes

spike threshold. Burst firing depends on the interplay between the afterhyperpolarization (AHPs)

and afterdepolarizations (ADPs) that follow the first action potential (Brumberg et al., 2000;

Mahon and Charpier, 2012; Paz et al., 2009). To explore this, we evaluated the dynamics of

action potentials using phase plot analysis. We used the first spikes from 60 pA current injections

in the I-V dataset; because during this current step, most neurons consistently displayed action

potentials before and after optogenetic or electrical stimulation protocol. Action potentials were

detected after setting a threshold of >0.00015 in the first derivative of membrane potential

waveform, a value that corresponds to the firing threshold, and voltage/time values (mV/ms)

were plotted as a function of membrane potential (mV). In this analysis, action potentials are

represented as a loop in which the starting point represents the firing threshold (Figure 7A). After

optogenetic stimulation, the majority of phase plots displayed a leftward offset, making evident a

lowering in firing threshold (-41± 1 before to -43.7 ± 1 mV after; p = 0.0012 by Wilcoxon; n =

13, 8 mice; Figure 7B-a). This change was partially reverted by terfenadine (10 µM; -45 ± 1 mV

after stimulation to -44 ± 1 mV after terfenadine; p = 0.05 by Wilcoxon; n = 8, 6 mice; Figure

7B-c). We found no statistically significant differences in firing threshold of unstimulated

neurons, when comparing the beginning and the end of the recording (-41 ± 2 and -41 ± 2 mV

respectively; p = 0.7 by Wilcoxon; n = 8 cells, 5 mice; Figure 7B-b). Also, non-expressing

neurons did not show any alteration of firing threshold after optogenetic stimulation of the slices

(before -40 ± 1.4 mV to -40 ± 1.5 mV after; p = 0.7 by Wilcoxon). Thus, the LED light did not

induce damage or alterations in this cell-intrinsic parameter.

We also documented a similar lowering of the threshold voltage in electrically stimulated

neurons (before -42 ± 0.8 mV; after 44 ± 0.8 mV; p = 0.00004 by Wilcoxon; n = 22 cells; Figure

7B-d). Unstimulated neurons displayed no changes in threshold voltage (-42 ± 0.7 mV at the first

min and -43 ± 0.8 mV 30 min later; p = 0.5 by Wilcoxon; n = 11, 8 mice; Figure 7B-e).

The combine increase in membrane resistance and reduction of firing threshold, can explain the

increase in neuronal excitability previously observed. Thus, optogenetic and electrical

stimulation significantly increases action potential firing in response to similar synaptic inputs,

facilitating the formation of a neuronal ensemble.

Figure 7. Spike threshold decreases after optogenetic and electrical stimulation

(A) Firing threshold shifts to more negative potentials after optogenetic and electrical

stimulation. Left: the firing threshold was taken to the first action potential generated with 60 pA

current step: before (black), after 30 min of optogenetic stimulation (blue) and, terfenadine post-

stimulation (green). Right: phase plot of action potentials: before (black), after (blue), and

terfenadine (green). The arrow shows the firing threshold position. (**B**) Firing threshold was

detected for every cell before and after: a) optogenetic stimulation, median and blue box plot (p

= 0.0009 by Wilcoxon, n = 14 cells, 9 mice); b) Unstimulated neurons with opsin expression,

median and gray box plots (p = 0.7 by Wilcoxon; n = 8 cells, 5 mice). c) Terfenadine (10µM)

post-stimulation, median and green box plot; p = 0.003 by Wilcoxon; n = 8 neurons, 5 mice;

firing threshold before stimulation and after terfenadine (p = 0.1 by Wilcoxon); d) electrical

stimulation, median and red box plot (p = 0.00004 by Wilcoxon; n = 22 cells, 15 mice). e)

Unstimulated neurons without opsin, median and gray box plot (p = 0.7 by Wilcoxon; n = 11

cells, 8 mice).

671   **Discussion**

The traditional view of how a neuronal ensemble (i.e., an assembly) is created is by

strengthening its synaptic connections after repeated coactivation (Bliss and Lømo, 1973;

Carrillo-Reid et al., 2016; Cossell et al., 2015; Hebb, 1949; Hoshiba et al., 2017). However,

several reports have questioned whether synaptic plasticity is the only mechanism of circuit

plasticity, proposing instead cell-intrinsic mechanisms (Angelo et al., 2012; Brown et al., 2019;

Disterhoft and Oh, 2006; Marder and Goaillard, 2006; Ryan et al., 2015; Zhang and Linden,

2003). Consistent with this, an in contradiction with our predictions that synaptic plasticity

mediates neuronal ensembles (Carrillo-Reid et al, 2016), we describe that simultaneous

optogenetic or electrical stimulation of neurons *in vitro* increases the excitability of the

stimulated cells. Our experimental procedures, similar to those that build ("imprint") neuronal

ensembles *in vivo* (Carrillo-Reid et al., 2016), lead to increases in correlations of spontaneous

activity, a hallmark of a neuronal ensemble, just as it does *in vivo*. The cellular and/or synaptic

mechanism induced by this optogenetic and electrical activation could contribute to neuronal

ensemble development (Penn and Shatz, 1999; Zhang and Poo, 2001), flexibility and plasticity

(Clopath et al., 2017; Kirkwood et al., 1996; Mrsic-Flogel et al., 2007; Pérez-Ortega et al., 2021;

Wandell and Smirnakis, 2009; Wiesel and Hubel, 1963).

Role of synaptic mechanisms in ensemble formation

Our stimulation protocols induced a moderate biphasic plasticity of connected neurons and also

in the local circuit, with an initial depression followed by a recovery potentiation. This agrees

with reports that EPSP amplitude can decrease after high stimulation frequencies, but recovers

from depression by using lower stimulation rates (Feldmeyer et al., 2006). This is consistent with

the hypothesis that persistent optogenetic or electrical stimulation reduces the readily releasable

pool of synaptic vesicles in the presynaptic neuron inducing, first, synaptic depression and, after

a period of rest, a rebound which is revealed as a potentiation (Feldmeyer et al., 2006;

Schneggenburger et al., 2002). But, in addition to these synaptic changes, our results demonstrate

that persistent and simultaneous stimulation robustly increases membrane resistance. Thus,

neuronal excitability could increase the efficiency of existing synaptic connections between

neurons in an ensemble, without significant short-term or long-term synaptic plasticity.

Our results appear in contradiction with the hypothesis that the formation of an ensemble is due

to increases in synaptic connectivity of synchronous neurons (Hebb, 1949; Cossell et al., 2015;

Hoshiba et al., 2017). Consistent with this hypothesis, there is preferential connectivity between

neurons with similar receptive fields (Cossell et al., 2015; Ko et al., 2011; Yoshimura et al.,

2005). Increased synaptic connectivity also probably underlies pattern completion in cortical

ensembles (Carrillo-Reid et al., 2019; Pignatelli et al., 2019), and a few strong connections could

drive local excitation of the majority of neurons with weak connectivity (Cossell et al., 2015;

Holmgren et al., 2003; Ko et al., 2011; Lefort et al., 2009; Levy and Reyes, 2012; Markram et

al., 1997; Song et al., 2005). Based on previous studies, we expected to observe synaptic

efficiency changes (Bliss and Gardner-Medwin, 1973) between connected neurons after

ensemble formation. Using paired recordings, we find similar rates of connectivity between

neuronal pairs as previous reports (Cossell et al., 2015; Holmgren et al., 2003; Ko et al., 2011;

Lefort et al., 2009; Levy and Reyes, 2012; Markram et al., 1997; Song et al., 2005). And, also

consistent with many reports, connected neurons display higher correlation in spontaneous

activity than unconnected ones (Cossell et al., 2015; Ko et al., 2011; Yoshimura et al., 2005).

Interestingly, our results also showed that 50% of unconnected neurons can be as correlated as

connected neurons. However, contrary to our own expectations, we did not detect any new

connections formed after optogenetic stimulation previously unconnected neurons. Moreover,

rather than synaptic potentiation, we found depression of evoked EPSCs and EPSPs. Although

also detect a moderate potentiation after 20 min of rest, and we cannot rule out that Hebbian

synaptic plasticity contributes to this potentiation, it is possible that this could instead be due to a

recovery of the ready releasable pool, together with increased synaptic efficacy due to changes in

postsynaptic and presynaptic excitability. Indeed, we find that spontaneous EPSPs and EPSCs

become amplified in electrically stimulated neurons. A larger synaptic input could also naturally

result from an overall increase in neuronal firing too, since increased neuronal excitability would

result in increase in neurotransmitter release. This agrees with reports that reveal that synaptic

inputs can be enhanced by intrinsic membrane mechanisms (Hu et al., 2002; Li et al., 2020;

Manabe et al., 1992; Marder and Goaillard, 2006; Narayanan and Johnston, 2008; Turrigiano et

al., 1998). For all these reasons, we suspect synaptic plasticity can be explained by the effects of

the stimulation protocols in the ready releasable pool and intrinsic changes in excitability in the

stimulated neurons and the network.

Role of intrinsic mechanisms in ensemble formation

Changes in intrinsic properties are known to modulate EPSP amplitude, for example, by the

modification of membrane conductance induced by postsynaptic stimulation (Hu et al., 2002; Li

et al., 2020; Manabe et al., 1992; Marder and Goaillard, 2006; Narayanan and Johnston, 2008;

Turrigiano et al., 1998). These changes might be due to modification in intrinsic membrane

conductances (Li et al., 2020; Manabe et al., 1992; Narayanan and Johnston, 2008), and they

could amplify synaptic inputs and improve the accuracy of synaptic integration of neurons

(Desai et al., 1999; Mahon and Charpier, 2012; Malik and Chattarji, 2012; Sjöström et al., 2008;

Turrigiano et al., 1994).

Confirming the importance of intrinsic mechanisms, in our experiments, robust increases in

firing rate were easily detected in optogenetic and electrically stimulated neurons. Even though

we have noticed that the expression of opsin by itself can have small effects in excitability,

opsin-expressing neurons still had major changes in intrinsic properties after optogenetic

stimulation, and very similar changes were observed in non-expressing neurons after electrical

stimulation. This rules out that the changes in excitability observed were due to the opsin

expression. An increase in cellular activity through cell-intrinsic mechanisms after prolonged

firing agrees with many functional studies in vertebrates, using different experimental

excitability protocols (Abraham et al., 2019; Ryan et al., 2015; Titley et al., 2017). In fact,

increases in excitability are also found following LTP induction in visual cortical neurons

(Cudmore and Turrigiano, 2004), in hippocampal CA1 pyramidal cells (Xu, 2005), and after

behavior training in CA1 (Disterhoft et al., 1986). Intrinsic excitability is also modified when an

animal is exposed to novel sensorial experience (Brown et al., 2019) or after environmental

enrichment (Valero-Aracama et al., 2015).

The intensity-dependent and persistent increase in firing rate after stimulation protocol could be

due to changes in membrane resistance and firing threshold. The synergy between these

mechanisms could make neurons reach firing threshold with smaller depolarizations. Thus, an

increase in excitability would be particularly relevant for lower current injections generated by

weaker inputs. We propose that, after optogenetic stimulation, neurons shift to a more excitable

state, revealing weaker synaptic inputs and connections, like an iceberg emerging out of the

water (Figure 8). Therefore, the same unchanged synaptic inputs could bring neurons to

threshold and induce increased output firing after repetitive stimulation, improving their synaptic

efficiency in postsynaptic pyramidal neurons (Debanne et al., 2019; Lisman et al., 2018; Nicoll

et al., 1993; Titley et al., 2017; Zhang and Linden, 2003). Thus, neuronal optogenetic and

electrical stimulation might alter the circuit without changing the synapses themselves, and this

phenomenon could occur quickly.

Mechanisms of increased excitability in ensemble formation

Modifications in sub- and suprathreshold membrane conductances that initiate an action potential

or a burst could underlie increased excitability (Brumberg et al., 2000). In terms of passive

properties, our results coincide with many studies that have shown alterations in input resistance

as the mechanism responsible for increases in excitability (Aizenman and Linden, 2000; Aou et

al., 1992; Armano et al., 2000; Brown et al., 2019; Disterhoft et al., 1986; Marder and Goaillard,

2006; Woody et al., 1991; Xu, 2005). Activity-dependent increases in input resistance have been

shown in hippocampal neurons. There, the increase in excitability could be potentiated by

suppression of G protein-coupled inwardly rectifying K^+ channels (GIRK channels) (Valero-

Aracama et al., 2015); or by cAMP-responsive element-binding protein (CREB)-dependent

control of excitability by reducing K^+ conductance (Lisman et al., 2018; Yu et al., 2017).

Additionally, the contribution of H-channels in intrinsic membrane properties (Campanac et al.,

2008; Fan et al., 2005). However, that voltage sag, which corresponds to current-H, is largely

absent in layer 2/3 pyramidal neurons (Kalmbach et al., 2018; van Aerde and Feldmeyer, 2015).

Our experiment showed only a few neurons that displayed a modest current-H, but it was not

affected by optogenetic or electrical stimulation (not shown).

Moreover, recent studies show that persisting firing could be mediated by an Ether-à-gogo

Related Gene (ERG) K^+ channel in neocortical pyramidal neurons (Cui and Strowbridge, 2018;

Debanne et al., 2019). In this view, calcium entry induced by repetitive activation could

modulate leak potassium currents in the neurons, which will generate persistent activity.

Consistent with this, in our experiments, the blockage of ERG channels by terfenadine returned

the firing rate and the increased input resistance induced by optogenetic and electrical

stimulation to basal conditions. Thus, ERG channels appear to have a tonic activity in pyramidal

layer 2/3 neurons.

The cell-intrinsic mechanisms that induce excitability are cell-type and also, stimulation

protocol-dependent (Angelo et al., 2012). Hence, different changes in action potential firing have

been identified. One of the most common changes is a reduction in after-hyperpolarization

(AHP) amplitude, described in hippocampal neurons (Disterhoft et al., 1986; Disterhoft and Oh,

2006; Malik and Chattarji, 2012; Pignatelli et al., 2019; Zhang and Linden, 2003). In our

experiments, we found a significant decrease in spike threshold as the most robust change in

optogenetic and electrical stimulated neurons. This parameter also determines the firing

frequency of neurons associated with voltage-dependent sodium or calcium currents. Changes in

threshold appear to be usual in cortical neurons after intrinsic plasticity induction (Brumberg et

al., 2000; Cudmore and Turrigiano, 2004; Mahon and Charpier, 2012; Paz et al., 2009),

hippocampal neurons (Malik and Chattarji, 2012; Valero-Aracama et al., 2015; Xu, 2005), and

cerebellar Purkinje cells (Aizenman and Linden, 2000; Armano et al., 2000). Consistent with

this, visual deprivation increases the spike threshold in the pyramidal neurons of the visual

cortex, as an example of a mechanism that reduces neuronal excitability (Brown et al., 2019). It

has been proposed that regulation of the persistent sodium current could underlie the lowering of

spike threshold, throughout the modulation of protein kinase C (Astman et al., 1998; Valero-

Aracama et al., 2015); or protein kinase A activation-dependent on calcium influx (Cudmore and

Turrigiano, 2004). Also, potassium channels Kv1 can regulate the action potential threshold

(Feria Pliego and Pedroarena, 2020). Therefore, the intrinsic plasticity by optogenetic stimulation

that we report in pyramidal neurons from the visual cortex could result from modifications of

voltage-gated ion channels and inward rectification by potassium channels. Future studies will

examine the exact molecular mechanisms that underlying these increases in excitability.

Contribution of intrinsic excitability and synaptic plasticity to circuit plasticity

The apparently contradictory evidence supporting Hebbian synaptic plasticity or intrinsic

mechanisms of circuit modification and ensemble formation could be reconciled if one considers

the experimental protocol used, as well as temporal and mechanistic factors. As a note of

caution, we should mention that the stimulation protocols we used are artificial, and the

imprinting of ensembles with optogenetics or electrical stimulation may not be representative of

the natural ensembles found and created in the cortex under physiological conditions. Thus, the

balance of mechanisms we find may differ with *in vivo* ensembles, under naturalistic learning

conditions. In particular, the optogenetic imprinting model for ensemble formation we used

occurs quickly *in vivo*, as soon as we can measure it, within a few minutes. We also see fast

changes in neuronal correlations *in vitro* after optogenetic and electrical stimulation. But this

type of imprinted ensembles could be different from the circuit modification that occur during

learning, which, for visual tasks, can take several days or weeks in the same circuit we study

(Carrillo-Reid et al., 2019), and which could engage long-term plasticity, Hebbian or not

Hebbian. In addition, intrinsic plasticity can underlie or contribute with Hebbian plasticity

(Debanne et al., 2019). Thus, behavioral training could induce neuronal excitability through

Hebbian mechanisms (Disterhoft et al., 1986; Malik and Chattarji, 2012; Valero-Aracama et al.,

2015). So, circuit plasticity could be due to the interplay of synaptic and intrinsic mechanisms

and this balance could be different for short and long-term plasticity. In this view that

synthesizes all the different types of evidence, ensembles could be quickly formed by fast

intrinsic mechanisms but then be modified by slower synaptic plasticity, perhaps in relation with

learning.

To conclude, our work demonstrates that significant increases in cellular excitability by intrinsic

mechanisms are found during ensemble formation induced by optogenetic and electrical

stimulation. The possibility to modify input and output patterns in a postsynaptic neuron through

changes in its firing properties could promote synaptic efficacy (Brumberg et al., 2000; Lisman,

1997) and increase the correlation between recruited neurons, as we have demonstrated. This

could provide an ideal setting for Hebbian synaptic plasticity than could occur more slowly.

Intrinsic excitability could also implement temporary stimulus-response mappings and might

play a role in rapid cognitive flexibility (Pang and Fairhall, 2019). Faster "online" plasticity via

intrinsic excitability could have a broad impact on network dynamics and could serve as a

critical information-storage mechanism that may contribute to memory formation (Lisman et al.,

2018; Marder and Goaillard, 2006; Pérez-Ortega et al., 2021; Pignatelli et al., 2019; Xu, 2005)

and to the generation of intrinsic circuit states, such as for ensembles and circuit attractors

(Buzsáki, 2010; Carrillo-Reid et al., 2019, 2016; Hopfield, 1982; Miller et al., 2014).

Figure 8. Iceberg model of ensemble formation.

(**A**) Emergence of ensembles after increases in neuronal excitability. Response of a neural circuit to a test input, before and after neurons are coactivated and increase their excitability. Color corresponds to membrane potential response to a synaptic input: white is resting membrane potentials, gray are subthreshold responses, and red are suprathreshold ones, with firing of action potentials. After stimulation, neurons become more excitable, so the same inputs induce some of them to fire (red cells), while other cells have increased subthreshold responses to the input (gray cells). The model can explain how an ensemble is formed but does not assume any changes in numbers of strength of synapses.

(**B**) Left: Iceberg emergence model: An iceberg keeps underwater. (Right) But if its weight decreases, the iceberg emerges above water. Weight (or density) is an intrinsic property of the iceberg and, by changing it, the iceberg changes its response to the same environment. Likewise, for a neuronal ensemble, membrane resistance and firing threshold are intrinsic neuronal

866   properties that can be modified, and they enhance its response to the same excitatory input

867   intensity, resulting in an increased depolarization and generation of action potentials.

869   **References**

Abeles M. 1991. Corticonics. Cambridge University Press.
doi:10.1017/CBO9780511574566
Abraham WC, Jones OD, Glanzman DL. 2019. Is plasticity of synapses the mechanism of
long-term memory storage? *npj Science of Learning* **4**:9. doi:10.1038/s41539-019-
0048-y
Aizenman CD, Linden DJ. 2000. Rapid, synaptically driven increases in the intrinsic
excitability of cerebellar deep nuclear neurons. *Nature Neuroscience* **3**.
doi:10.1038/72049
Angelo K, Rancz EA, Pimentel D, Hundahl C, Hannibal J, Fleischmann A, Pichler B,
Margrie TW. 2012. A biophysical signature of network affiliation and sensory
processing in mitral cells. *Nature* **488**:375–378. doi:10.1038/nature11291
Aou S, Woody C, Birt D. 1992. Increases in excitability of neurons of the motor cortex of
cats after rapid acquisition of eye blink conditioning. *The Journal of Neuroscience* **12**.
doi:10.1523/JNEUROSCI.12-02-00560.1992
Armano S, Rossi P, Taglietti V, D'Angelo E. 2000. Long-Term Potentiation of Intrinsic
Excitability at the Mossy Fiber–Granule Cell Synapse of Rat Cerebellum. *The Journal
of Neuroscience* **20**. doi:10.1523/JNEUROSCI.20-14-05208.2000
Astman N, Gutnick MJ, Fleidervish IA. 1998. Activation of Protein Kinase C Increases
Neuronal Excitability by Regulating Persistent Na + Current in Mouse Neocortical
Slices. *Journal of Neurophysiology* **80**:1547–1551. doi:10.1152/jn.1998.80.3.1547
Bliss TVP, Gardner-Medwin AR. 1973. Long-lasting potentiation of synaptic transmission
in the dentate area of the unanaesthetized rabbit following stimulation of the perforant
path. *The Journal of Physiology* **232**:357–374. doi:10.1113/jphysiol.1973.sp010274
Bliss TVP, Lømo T. 1973. Long-lasting potentiation of synaptic transmission in the dentate
area of the anaesthetized rabbit following stimulation of the perforant path. *The
Journal of Physiology* **232**. doi:10.1113/jphysiol.1973.sp010273
Brown APY, Cossell L, Margrie TW. 2019. Visual Experience Regulates the Intrinsic
Excitability of Visual Cortical Neurons to Maintain Sensory Function. *Cell Reports*
27:685-689.e4. doi:10.1016/j.celrep.2019.03.073
Brumberg JC, Nowak LG, McCormick DA. 2000. Ionic Mechanisms Underlying Repetitive
High-Frequency Burst Firing in Supragranular Cortical Neurons. *Journal of
Neuroscience* **20**:4829–4843. doi:10.1523/JNEUROSCI.20-13-04829.2000
Buzsáki G. 2010. Neural syntax: cell assemblies, synapsembles, and readers. *Neuron*
68:362–85. doi:10.1016/j.neuron.2010.09.023

Campanac E, Daoudal G, Ankri N, Debanne D. 2008. Downregulation of Dendritic Ih in CA1 Pyramidal Neurons after LTP. *Journal of Neuroscience* **28**. doi:10.1523/JNEUROSCI.1411-08.2008

Carrillo-Reid L, Han S, Yang W, Akrouh A, Yuste R. 2019. Controlling Visually Guided Behavior by Holographic Recalling of Cortical Ensembles. *Cell* **178**:447-457.e5. doi:10.1016/j.cell.2019.05.045

Carrillo-Reid L, Yang W, Bando Y, Peterka DS, Yuste R. 2016. Imprinting and recalling cortical ensembles. *Science* **353**:691–694. doi:10.1126/science.aaf7560

Carter CA, Wojciechowski NJ, Hayes JM, Skoutakis VA, Rickman LA. 1985. Terfenadine, a Nonsedating Antihistamine. *Drug Intelligence & Clinical Pharmacy* **19**. doi:10.1177/106002808501901103

Churchland PS, Sejnowski TJ. 1992. The computational brain., The computational brain., Computational neuroscience. Cambridge, MA, US: The MIT Press.

Clopath C, Bonhoeffer T, Hübener M, Rose T. 2017. Variance and invariance of neuronal long-term representations. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**. doi:10.1098/rstb.2016.0161

Cossart R, Aronov D, Yuste R. 2003. Attractor dynamics of network UP states in the neocortex. *Nature* **423**:283–8. doi:10.1038/nature01614

Cossell L, Iacaruso MF, Muir DR, Houlton R, Sader EN, Ko H, Hofer SB, Mrsic-Flogel TD. 2015. Functional organization of excitatory synaptic strength in primary visual cortex. *Nature* **518**:399–403. doi:10.1038/nature14182

Cudmore RH, Turrigiano GG. 2004. Long-Term Potentiation of Intrinsic Excitability in LV Visual Cortical Neurons. *Journal of Neurophysiology* **92**:341–348. doi:10.1152/jn.01059.2003

Cui ED, Strowbridge BW. 2018. Modulation of Ether-à-Go-Go Related Gene (ERG) Current Governs Intrinsic Persistent Activity in Rodent Neocortical Pyramidal Cells. *Journal of Neuroscience* **38**:423–440. doi:10.1523/JNEUROSCI.1774-17.2017

dal Maschio M, Ghezzi D, Bony G, Alabastri A, Deidda G, Brondi M, Sato SS, Zaccaria RP, di Fabrizio E, Ratto GM, Cancedda L. 2012. High-performance and site-directed in utero electroporation by a triple-electrode probe. *Nature Communications* **3**:960. doi:10.1038/ncomms1961

Debanne D, Inglebert Y, Russier M. 2019. Plasticity of intrinsic neuronal excitability. *Current Opinion in Neurobiology* **54**:73–82. doi:10.1016/j.conb.2018.09.001

Desai NS, Rutherford LC, Turrigiano GG. 1999. Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nature Neuroscience* **2**. doi:10.1038/9165

Disterhoft JF, Coulter DA, Alkon DL. 1986. Conditioning-specific membrane changes of rabbit hippocampal neurons measured in vitro. *Proceedings of the National Academy of Sciences* **83**:2733–2737. doi:10.1073/PNAS.83.8.2733

Disterhoft JF, Oh MM. 2006. Learning, aging and intrinsic neuronal plasticity. *Trends in Neurosciences* **29**:587–599. doi:10.1016/j.tins.2006.08.005

Fan Y, Fricker D, Brager DH, Chen X, Lu H-C, Chitwood RA, Johnston D. 2005. Activity-dependent decrease of excitability in rat hippocampal neurons through increases in Ih. *Nature Neuroscience* **8**. doi:10.1038/nn1568

Feldmeyer D, Lübke J, Sakmann B. 2006. Efficacy and connectivity of intracolumnar pairs of layer 2/3 pyramidal cells in the barrel cortex of juvenile rats. *The Journal of Physiology* **575**. doi:10.1113/jphysiol.2006.105106

Feria Pliego JA, Pedroarena CM. 2020. Kv1 potassium channels control action potential firing of putative GABAergic deep cerebellar nuclear neurons. *Scientific Reports* **10**:6954. doi:10.1038/s41598-020-63583-7

Hamm JP, Peterka DS, Gogos JA, Yuste R. 2017. Altered Cortical Ensembles in Mouse Models of Schizophrenia. *Neuron* **94**:153-167.e8. doi:10.1016/j.neuron.2017.03.019

Hebb DO. 1949. The Organization of Behavior, New York: Wiley. doi:10.1016/S0361-9230(99)00182-3

Holmgren C, Harkany T, Svennenfors B, Zilberter Y. 2003. Pyramidal cell communication within local networks in layer 2/3 of rat neocortex. *The Journal of Physiology* **551**. doi:10.1113/jphysiol.2003.044784

Hopfield JJ. 1982. Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences of the United States of America* **79**:2554–8. doi:10.1073/pnas.79.8.2554

Hoshiba Y, Wada T, Hayashi-Takagi A. 2017. Synaptic Ensemble Underlying the Selection and Consolidation of Neuronal Circuits during Learning. *Frontiers in Neural Circuits* **11**. doi:10.3389/fncir.2017.00012

Hu H, Vervaeke K, Storm JF. 2002. Two forms of electrical resonance at theta frequencies, generated by M-current, h-current and persistent Na$^+$ current in rat hippocampal pyramidal cells. *The Journal of Physiology* **545**. doi:10.1113/jphysiol.2002.029249

Ikegaya Y. 2004. Synfire Chains and Cortical Songs: Temporal Modules of Cortical Activity. *Science* **304**:559–564. doi:10.1126/science.1093173

Johansson F, Jirenhed D-A, Rasmussen A, Zucca R, Hesslow G. 2014. Memory trace and timing mechanism localized to cerebellar Purkinje cells. *Proceedings of the National Academy of Sciences* **111**:14930–14934. doi:10.1073/pnas.1415371111

Juárez-Vidales J de J, Pérez-Ortega J, Lorea-Hernández JJ, Méndez-Salcido F, Peña-Ortega F. 2021. Configuration and dynamics of dominant inspiratory multineuronal activity patterns during eupnea and gasping generation in vitro. *Journal of Neurophysiology* **125**. doi:10.1152/jn.00563.2020

Kalmbach BE, Buchin A, Long B, Close J, Nandi A, Miller JA, Bakken TE, Hodge RD, Chong P, de Frates R, Dai K, Maltzer Z, Nicovich PR, Keene CD, Silbergeld DL, Gwinn RP, Cobbs C, Ko AL, Ojemann JG, Koch C, Anastassiou CA, Lein ES, Ting JT. 2018. h-Channels Contribute to Divergent Intrinsic Membrane Properties of Supragranular Pyramidal Neurons in Human versus Mouse Cerebral Cortex. *Neuron* **100**. doi:10.1016/j.neuron.2018.10.012

Kirkwood A, Rioult MG, Bear MF. 1996. Experience-dependent modification of synaptic plasticity in visual cortex. *Nature* **381**. doi:10.1038/381526a0

Ko H, Hofer SB, Pichler B, Buchanan KA, Sjöström PJ, Mrsic-Flogel TD. 2011. Functional specificity of local synaptic connections in neocortical networks. *Nature* **473**:87–91. doi:10.1038/nature09880

Lefort S, Tomm C, Floyd Sarria J-C, Petersen CCH. 2009. The Excitatory Neuronal Network of the C2 Barrel Column in Mouse Primary Somatosensory Cortex. *Neuron* **61**. doi:10.1016/j.neuron.2008.12.020

Levy RB, Reyes AD. 2012. Spatial Profile of Excitatory and Inhibitory Synaptic Connectivity in Mouse Primary Auditory Cortex. *Journal of Neuroscience* **32**. doi:10.1523/JNEUROSCI.5158-11.2012

Li B, Routh BN, Johnston D, Seidemann E, Priebe NJ. 2020. Voltage-Gated Intrinsic Conductances Shape the Input-Output Relationship of Cortical Neurons in Behaving Primate V1. *Neuron* **107**:185-196.e4. doi:10.1016/j.neuron.2020.04.001

Lisman J, Cooper K, Sehgal M, Silva AJ. 2018. Memory formation depends on both synapse-specific modifications of synaptic strength and cell-specific increases in excitability. *Nature Neuroscience* **21**:309–314. doi:10.1038/s41593-018-0076-6

Lisman JE. 1997. Bursts as a unit of neural information: making unreliable synapses reliable. *Trends in Neurosciences* **20**:38–43. doi:10.1016/S0166-2236(96)10070-9

Lorente De Nó R. 1938. Analysis of the activity of the chains of internuncial neurons. *Journal of Neurophysiology* **1**:207–244. doi:10.1152/jn.1938.1.3.207

Lorente de Nó R. 1933. Studies on the structure of the cerebral cortex. I. The area entorhinalis. *Journal für Psychologie und Neurologie*.

Mahon S, Charpier S. 2012. Bidirectional Plasticity of Intrinsic Excitability Controls Sensory Inputs Efficiency in Layer 5 Barrel Cortex Neurons in Vivo. *Journal of Neuroscience* **32**. doi:10.1523/JNEUROSCI.0415-12.2012

Malik R, Chattarji S. 2012. Enhanced intrinsic excitability and EPSP-spike coupling accompany enriched environment-induced facilitation of LTP in hippocampal CA1 pyramidal neurons. *Journal of Neurophysiology* **107**:1366–1378. doi:10.1152/jn.01009.2011

Manabe T, Renner P, Nicoll RA. 1992. Postsynaptic contribution to long-term potentiation revealed by the analysis of miniature synaptic currents. *Nature* **355**. doi:10.1038/355050a0

Marder E, Goaillard J-M. 2006. Variability, compensation and homeostasis in neuron and network function. *Nature Reviews Neuroscience* **7**:563–574. doi:10.1038/nrn1949

Mardinly AR, Oldenburg IA, Pégard NC, Sridharan S, Lyall EH, Chesnov K, Brohawn SG, Waller L, Adesnik H. 2018. Precise multimodal optical control of neural ensemble activity. *Nature Neuroscience* **21**:881–893. doi:10.1038/s41593-018-0139-8

Markram H, Lübke J, Frotscher M, Roth A, Sakmann B. 1997. Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex. *The Journal of Physiology* **500**. doi:10.1113/jphysiol.1997.sp022031

Marshel JH, Kim YS, Machado TA, Quirin S, Benson B, Kadmon J, Raja C, Chibukhchyan A, Ramakrishnan C, Inoue M, Shane JC, McKnight DJ, Yoshizawa S, Kato HE, Ganguli S, Deisseroth K. 2019. Cortical layer-specific critical dynamics triggering perception. *Science (New York, NY)* **365**. doi:10.1126/science.aaw5202

Mellor J. 2002. Mediation of Hippocampal Mossy Fiber Long-Term Potentiation by Presynaptic Ih Channels. *Science* **295**. doi:10.1126/science.1064285

Miller J -e. K, Ayzenshtat I, Carrillo-Reid L, Yuste R. 2014. Visual stimuli recruit intrinsically generated cortical ensembles. *Proceedings of the National Academy of Sciences* **111**:E4053–E4061. doi:10.1073/pnas.1406077111

Morishima M, Morita K, Kubota Y, Kawaguchi Y. 2011. Highly Differentiated Projection-Specific Cortical Subnetworks. *Journal of Neuroscience* **31**. doi:10.1523/JNEUROSCI.0772-11.2011

Mrsic-Flogel TD, Hofer SB, Ohki K, Reid RC, Bonhoeffer T, Hübener M. 2007. Homeostatic Regulation of Eye-Specific Responses in Visual Cortex during Ocular Dominance Plasticity. *Neuron* **54**. doi:10.1016/j.neuron.2007.05.028

Narayanan R, Johnston D. 2008. The h Channel Mediates Location Dependence and
Plasticity of Intrinsic Phase Response in Rat Hippocampal Neurons. *Journal of*
Neuroscience **28**. doi:10.1523/JNEUROSCI.0835-08.2008

Nicoll A, Larkman A, Blakemore C. 1993. Modulation of EPSP shape and efficacy by
intrinsic membrane conductances in rat neocortical pyramidal neurons in vitro. *The*
Journal of Physiology **468**. doi:10.1113/jphysiol.1993.sp019795

Pang R, Fairhall AL. 2019. Fast and flexible sequence induction in spiking neural networks
via rapid excitability changes. *eLife* **8**. doi:10.7554/eLife.44324

Paz JT, Mahon S, Tiret P, Genet S, Delord B, Charpier S. 2009. Multiple forms of activity-
dependent intrinsic plasticity in layer V cortical neurones in vivo. *The Journal of*
Physiology **587**:3189–3205. doi:10.1113/jphysiol.2009.169334

Penn AA, Shatz CJ. 1999. Brain Waves and Brain Wiring: The Role of Endogenous and
Sensory-Driven Neural Activity in Development. *Pediatric Research* **45**:447–458.
doi:10.1203/00006450-199904010-00001

Pérez-Ortega J, Alejandre-García T, Yuste R. 2021. Long-term stability of cortical
ensembles. *eLife* **10**. doi:10.7554/eLife.64449

Pignatelli M, Ryan TJ, Roy DS, Lovett C, Smith LM, Muralidhar S, Tonegawa S. 2019.
Engram Cell Excitability State Determines the Efficacy of Memory Retrieval. *Neuron*
101:274-284.e5. doi:10.1016/j.neuron.2018.11.029

Ramón y Cajal S. 1909. Histologie du système nerveux de l'homme & des vertébrés.
Paris : Maloine,. doi:10.5962/bhl.title.48637

Ross ST, Soltesz I. 2001. Long-term plasticity in interneurons of the dentate gyrus.
Proceedings of the National Academy of Sciences **98**:8874.
doi:10.1073/pnas.141042398

Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S. 2015. Engram cells retain memory
under retrograde amnesia. *Science* **348**:1007–1013. doi:10.1126/science.aaa5542

Sasaki T, Matsuki N, Ikegaya Y. 2007. Metastability of Active CA3 Networks. *Journal of*
Neuroscience **27**:517–528. doi:10.1523/JNEUROSCI.4514-06.2007

Schneggenburger R, Sakaba T, Neher E. 2002. Vesicle pools and short-term synaptic
depression: lessons from a large synapse. *Trends in Neurosciences* **25**:206–212.
doi:10.1016/S0166-2236(02)02139-2

Schüz A, Braitenberg V. 2001. Cerebral Cortex: Organization and FunctionInternational
Encyclopedia of the Social & Behavioral Sciences. Elsevier. pp. 1634–1640.
doi:10.1016/B0-08-043076-7/03443-4

Sempere-Ferràndez A, Martínez S, Geijo-Barrientos E. 2019. Synaptic mechanisms
underlying the intense firing of neocortical layer 5B pyramidal neurons in response to
cortico-cortical inputs. *Brain Structure and Function* **224**. doi:10.1007/s00429-019-
01842-8

Shepherd G. 2010. Handbook of Brain Microcircuits. Oxford University Press.
doi:10.1093/med/9780195389883.001.0001

Shepherd GM. 2004. The Synaptic Organization of the Brain, The Synaptic Organization of
the Brain. Oxford University Press. doi:10.1093/acprof:oso/9780195159561.001.1

Sjöström PJ, Rancz EA, Roth A, Häusser M. 2008. Dendritic Excitability and Synaptic
Plasticity. *Physiological Reviews* **88**. doi:10.1152/physrev.00016.2007

Song S, Sjöström PJ, Reigl M, Nelson S, Chklovskii DB. 2005. Highly Nonrandom Features of Synaptic Connectivity in Local Cortical Circuits. *PLoS Biology* **3**. doi:10.1371/journal.pbio.0030068

Stringer C, Pachitariu M, Steinmetz N, Carandini M, Harris KD. 2019. High-dimensional geometry of population responses in visual cortex. *Nature* **571**:361–365. doi:10.1038/s41586-019-1346-5

Ting JT, Lee BR, Chong P, Soler-Llavina G, Cobbs C, Koch C, Zeng H, Lein E. 2018. Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-glucamine Protective Recovery Method. *Journal of Visualized Experiments* e53825. doi:10.3791/53825

Titley HK, Brunel N, Hansel C. 2017. Toward a Neurocentric View of Learning. *Neuron* **95**:19–32. doi:10.1016/j.neuron.2017.05.021

Tonegawa S, Pignatelli M, Roy DS, Ryan TJ. 2015. Memory engram storage and retrieval. *Current Opinion in Neurobiology* **35**:101–109. doi:10.1016/j.conb.2015.07.009

Turrigiano G, Abbott L, Marder E. 1994. Activity-dependent changes in the intrinsic properties of cultured neurons. *Science* **264**. doi:10.1126/science.8178157

Turrigiano GG, Leslie KR, Desai NS, Rutherford LC, Nelson SB. 1998. Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature* **391**. doi:10.1038/36103

Valero-Aracama MJ, Sauvage MM, Yoshida M. 2015. Environmental enrichment modulates intrinsic cellular excitability of hippocampal CA1 pyramidal cells in a housing duration and anatomical location-dependent manner. *Behavioural Brain Research* **292**:209–218. doi:10.1016/J.BBR.2015.05.032

van Aerde KI, Feldmeyer D. 2015. Morphological and Physiological Characterization of Pyramidal Neuron Subtypes in Rat Medial Prefrontal Cortex. *Cerebral Cortex* **25**. doi:10.1093/cercor/bht278

Wandell BA, Smirnakis SM. 2009. Plasticity and stability of visual field maps in adult primary visual cortex. *Nature Reviews Neuroscience* **10**. doi:10.1038/nrn2741

Wiesel TN, Hubel DH. 1963. EFFECTS OF VISUAL DEPRIVATION ON MORPHOLOGY AND PHYSIOLOGY OF CELLS IN THE CAT'S LATERAL GENICULATE BODY. *Journal of Neurophysiology* **26**:978–993. doi:10.1152/jn.1963.26.6.978

Woody CD, Gruen E, Birt D. 1991. Changes in membrane currents during Pavlovian conditioning of single cortical neurons. *Brain Research* **539**. doi:10.1016/0006-8993(91)90688-R

Xu J. 2005. Activity-Dependent Long-Term Potentiation of Intrinsic Excitability in Hippocampal CA1 Pyramidal Neurons. *Journal of Neuroscience* **25**:1750–1760. doi:10.1523/JNEUROSCI.4217-04.2005

Yang Z, Santamaria F. 2016. Purkinje cell intrinsic excitability increases after synaptic long term depression. *Journal of Neurophysiology* **116**. doi:10.1152/jn.00369.2016

Yoshimura Y, Dantzker JLM, Callaway EM. 2005. Excitatory cortical neurons form fine-scale functional networks. *Nature* **433**. doi:10.1038/nature03252

Yu X-W, Oh MM, Disterhoft JF. 2017. CREB, cellular excitability, and cognition: Implications for aging. *Behavioural Brain Research* **322**:206–211. doi:10.1016/J.BBR.2016.07.042

Yuste R. 2015. From the neuron doctrine to neural networks. *Nature Reviews Neuroscience* **16**:487–497. doi:10.1038/nrn3962

1130   Zhang LI, Poo M. 2001. Electrical activity and development of neural circuits. *Nature*
1131   *Neuroscience* **4**. doi:10.1038/nn753
1132   Zhang W, Linden DJ. 2003. The other side of the engram: experience-driven changes in
1133   neuronal intrinsic excitability. *Nature Reviews Neuroscience* **4**. doi:10.1038/nrn1248
1134
1135