Defending as a unit: sex- and context-specific territorial defence in a duetting bird

Lucia Mentesana[1]†, Maria Moiron[1,2]†, Ernesto Guedes[3], Enzo Cavalli[3], Bettina Tassino[3],

Nicolas M. Adreani[1,4]*†

[1] Max Planck Institute for Ornithology, Seewiesen, Germany.

[2] Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175 Campus CNRS, Montpellier,

France.

[3] Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay.

[4] Konrad Lorenz Research Center, University of Vienna, Grünau im Almtal, 4643, Austria

* Author for correspondence: nicolas.adreani@univie.ac.at

† Contributed equally to the manuscript

Abstract

Behaviours such as territorial defence represent functionally integrated traits that underlie

multiple behavioural variables such as physical and acoustic responses. Characterizing the

multivariate structure of such traits is fundamental to understand their evolution. In bird species

that form stable pair bonds and are territorial year-round, both sexes are expected to defend

their territory; however, the role that each sex plays in defending their shared territory remains

largely unknown. Evidence for the sex-roles during territorial defence is mixed and sex- and

context-specific characterizations of territorial defence embracing the multivariate nature of the

trait are currently lacking. Here we investigated sex- and context-specific variation in a

hypothesised latent variable called "territorial defence" and tested whether duets were part of

territorial defence in a wild population of rufous hornero (*Furnarius rufus*). To do so, we

combined a simulated territorial intrusion approach during nest building and provisioning

contexts with a structural equation modelling approach. Our results showed that, in males and

females, the six measured behavioural variables were linked by a single latent trait, territorial

defence, in both contexts. Flights over the decoy and duet songs were equally good proxies of

territorial defence. Although males were defending more the territory than females, pair

members showed a positive correlation in their behaviour. The structural equation modelling

framework enabled us to capture a complex correlation pattern among behavioural variables,

expanding upon a classic body of research on territorial defence. Thus, the combination of

classical behavioural approaches with sophisticated statistical analyses brings new exciting

possibilities to the field of behavioural ecology.

Keywords: territorial aggression, duets, behavioural character, phenotypic integration, female
aggression, structural equation analysis

Significance statement

Territorial defence is a key behaviour in territorial species as it plays a major role in an
individual's reproductive success and survival. Additionally, territorial defence has been
proposed as one possible evolutionary driver of duetting behaviour, one of the most fascinating
vocal behaviours in birds. As behaviours are evolutionary characters, they must be studied in a
multivariate framework. In this study we focused on characterizing territorial defence during a
simulated territorial intrusion in an integrative manner using a classical territorial intrusion
framework. We did so in male and female rufous horneros (Aves: *Furnaridae*) across two
breeding contexts, while simultaneously testing theoretical predictions about the role of
duetting behaviour as key part of territorial defence. Overall, our study provides for the first
time a sex- and context-comparison of the multivariate, latent variable "territorial defence" in
duetting birds, while highlighting the potential of combining field behavioural approaches with
structural equation modelling.
49

Introduction

Behaviours, such as territorial defence, are typically studied by simultaneously measuring different observable variables (e.g. "number of attacks", "latency of response", "proximity" measures, and/or "vocal responses"; Wingfield 1994; Bollen 2002). Extensive empirical research over the last decades has focused on analyzing male and, to a lesser extent, female territorial defence behaviour using different approaches. Evidence generally shows that multiple components of an animal' defence response in a territory intrusion are intercorrelated (e.g. Huntingford 1976; Sprenger et al. 2012). When these multiple behaviours are functionally related, they might be considered expressions of a single evolutionary character (Araya-Ajoy and Dingemanse 2013). Their study should therefore not be addressed by means of bivariate correlations but integrate the multivariate nature of the behaviour by quantifying unobserved, biologically-relevant latent variables (Houle et al. 2011; Carter and Feeney 2012; Araya-Ajoy and Dingemanse 2013). One way to characterize "territorial defence" as an evolutionary character while also quantifying which of the observed behavioural variables should be considered expressions of it, is to apply a Structural Equation Modelling approach (SEM; e.g. Card and Little 2007). This statistical framework allows to explore complex correlation patterns among multiple behavioural variables, and to test *a priori* defined hypotheses of how multiple observed behavioural variables are linked by the unmeasured latent variable (Araya-Ajoy and Dingemanse 2013).

Territorial defence behaviour has been widely studied in diverse organisms from insects to several groups of vertebrates (reviewed in Smith and Blumstein 2008), likely because of its impact on fitness (Stamps and Krishnan 1997; Smith and Blumstein 2008). During territory defence, aggressive interactions can be beneficial for both males and females because an intrusion of a conspecific into the breeding territory might, for example, lead to loss of limited resources (Stamps and Krishnan 1997; Garcia and Arroyo 2002). Nevertheless, the sex-specific contribution to territory defence differs among species according to variation in mating systems and parental care (Emlen and Oring 1977; Clutton-Brock and Vincent 1991; Owens and Thompson 1994). In birds, among those species that form stable pair bonds and are territorial year round, it is expected that males and females equally contribute to territory defence (Greenberg and Gradwohl 1983). In line with this prediction, in dot-winged antwrens (*Microrhopia quixensis*) and in the purple-crowned and red-backed fairy-wrens (*Malurus coronatus, Malurus melanocephalus*) both sexes contribute to the same extent to defend their territory (Greenberg and Gradwohl 1983; Hall and Peters 2008; Dowling and Webster 2016). However, these findings were not observed in other antbird species (*Phaenostictus mcleannani,*

Willis 1972; *Cercomacra tyrannina*; Morton and Derrickson 1996; *Hylophylax naevioides*; Bard et al. 2002; *Myrmeciza longipes*; Fedy and Stutchbury 2005), in the zenaida dove (*Zenaida aurit*a; Quinard and Cézilly 2012) and in the rufous hornero (*Furnarius rufus;* Diniz et al. 2018), where males engaged more in defensive interactions than females. In addition, whilst intensity of territorial defence can be positively correlated within pairs in some species (especially on those that duet e.g. Logue 2005; Hall and Peters 2008), in others the opposite relationship is true (e.g. zenaida dove; Quinard and Cézilly 2012). Therefore, the generality of sex-specific territorial defence as well as its intensity from both members of a pair remains poorly understood in species that are socially monogamous and territorial year-round. Furthermore, it also remains largely unknown whether the same observed behavioural variables characterize the latent variable territorial defence in males and females and across different breeding contexts. For instance, it is still an open question whether territorial defence is equally characterized by physical and vocal behaviours in both sexes and across contexts. Studies describing territorial defence as a latent variable using SEM and explaining broader patterns of territorial defence across sexes or contexts, will help to shed light on these questions and further our understanding of the evolution of male and female defence of territory.

While physical displays are considered the main defensive responses, the role of vocal displays as defensive signals remains under discussion (see Searcy and Beecher 2009; Naguib and Mennill 2010). Among the vocal displays that take place during agonistic interactions, perhaps the most fascinating one is duetting – occurring in around 18% of avian species worldwide (Tobias et al. 2016). Duets are defined as coordinated vocal interactions between two individuals - usually a male and female of a pair - that occur with a given temporal precision (Farabaugh 1982; Hall 2004). Duets are hypothesised to represent an important component of territorial defence (Langmore 1998; Hall 2004). In particular, the "joint territorial defence" hypothesis, proposed as one evolutionary driver of duetting behaviour (Wickler and Seibt 1980), postulates that duets allow pairs to cooperatively defend resources from conspecific intruders (Robinson 1949; reviewed by Hall 2004). A central prediction of the "joint territorial defence" hypothesis is that duets are threatening signals, stronger than solo songs (Hall 2004). To date, few studies investigated duetting in the context of territorial defence across different life-history stages (Topp and Mennill 2008; Odom et al. 2017; Quirós-Guerrero et al. 2017; Sosa-López et al. 2017; Diniz et al. 2018). The few that did so used mainly three methods: i) context criterion (i.e. which compares responses towards acoustic stimuli that represent different contexts, like only male/female solo songs, only duet songs or only heterospecific songs; e.g. Dowling and Webster 2016), ii) response criterion (i.e. which compares responses

with and without a playback stimulus e.g. Hall and Peters 2008), and iii) correlation methods
(i.e. which applies correlation techniques and principal component transformations on
behavioural data, e.g. Kolof and Mennil 2013). However, none of the abovementioned methods
allow to test the role that duets play in the context of territorial defence while embracing the
multivariate nature of this traits as a latent variable.

We conducted simulated territorial intrusions (STI) in the territory of focal pairs of
rufous horneros during two contexts in the breeding season: nest building and chick
provisioning. The rufous hornero, hereafter hornero, is a single brooded furnarid bird species
that is widely distributed throughout southern South America (Fraga 1980). Horneros are
territorial year round, socially monogamous (Fraga 1980; Diniz et al. 2018) and both members
of the pair are involved in defending their territory (Fraga 1980; Diniz et al. 2018). Indeed, all
breeding behaviours studied in horneros so far are performed in an equitable and coordinated
manner between sexes, such as incubation, parental care-related activities and even territorial
defence in non-breeding context (Fraga 1980; Massoni et al. 2012; Diniz et al. 2020). Also,
previous studies on this species suggest that duets have a territorial function (Diniz et al. 2018,
2019, 2020). However, these studies were either acoustic-centered (Diniz et al. 2018, 2019) or
carried out during a non-breeding season (Diniz et al. 2020), and none of them considered the
multivariate nature of territorial defence.

The main goals of our study were first to characterize the multivariate nature of
territorial defence in male and female horneros, and to test whether duets were indeed part of
the defence displays during a territorial intrusion. Our second goal was to quantify phenotypic
variation across sexes, breeding contexts and pair members. Our third goal was to evaluate the
level of coordination between sexes during territorial defence. To do so, we constructed a series
of structural equation models where we tested three hypotheses of potential associations among
the behavioural variables: model 1 hypothesised that each territorial defence behaviour is
independent and not part of a functional unit or evolutionary character; model 2 hypothesises
that one latent variable, "territorial defence", underlies the relationships between all behavioural
variables; and model 3 hypothesises that all behavioural variables except number of duet songs
are linked by the latent variable "territorial defence". These models were therefore specifically
constructed to test the "joint territorial defence" hypothesis (Hall 2009). According to this
hypothesis, for our first aim, we predicted duets to be part of the latent trait 'territorial defence'
and to be more relevant than solo songs. We also predicted that territorial defence will be
characterized by the same behavioural traits (i.e. number of duets, number of flights over the
decoy, times spent within 5m of the decoy, number of solo songs, time spent on nest) in males

and females. Second, we predicted that in our STIs males would defend more their territories
than females. This was based on the notion that, although in neotropical birds there is mixed
evidence for the sexual difference in territorial defence, a recent study reported that male
horneros engaged more in defending their territories than females (Diniz et al. 2018). Further,
because in horneros territory take-over is expected to be a stronger driver of aggression than
paternity loss (i.e. extra-pair paternity levels are ~ 3%; Diniz et al. 2019), we predicted higher
levels of territory defence earlier in the nest building than in the provisioning context (see also
Demko and Mennill 2018). Finally, we predicted both members of the pair to positively
correlate their territory defence behaviors (e.g. Diniz et al. 2020).
161

Materials and Methods
Field site and experimental procedures
We studied pairs of horneros in two periods during 2016 on the campus of INIA "Las Brujas"
(National Institute of Agricultural Research), department of Canelones, Uruguay (34°40' S,
56°20' W; 0-35 m a.s.l.). Behavioural assays were carried out during "nest-building" (i.e. when
pairs were observed finishing their nests and females were in their fertile period; August 23rd –
September 27th), and "provisioning" periods (i.e. when pairs were observed feeding their young;
November 7th – December 6th). Overall, we observed 39 males and 38 females during nest
building and 25 males and 24 females during provisioning. Each pair was tested only one time
(i.e. either during the nest-building or during the provisioning period). It was not possible to
record data blind because our study involved focal animals in the field. All the behavioural
assays were performed between 07h00 and 13h00. We quantified territorial defence behaviours
by performing simulated territorial intrusions (STI) in the territory of a focal pair. Once the
focal pair was identified, a stuffed decoy of a hornero together with playback sounds was
presented 10 metres away from the pair's nest. Using two decoys could have been more
realistic; however, we could not do this for ethical reasons. Every STI playback lasted 20
minutes and consisted of randomly selected stimuli from a pool of ten male solo songs, ten
duets and ten audio files containing 7-15 seconds of silence. The auditory stimuli for each STI
were randomly selected to avoid pseudo-replication of the acoustic component across territories
(e.g. Apfelbeck et al. 2011), to avoid a behavioural bias towards specific songs or duets, and to
elicit comparable behavioural responses across territories. Our approach hinged on the notion
that horneros are suboscines and do not learn their vocalizations (Freeman et al. 2017). This is,
compared to oscines, the acoustic variability of songs and duets across individuals is low
(Freeman et al. 2017). We played silence tracks of different length to avoid habituation in the

focal birds. All playback sounds were "wav" files recorded in Uruguay and were obtained from
the database of Xeno-canto (www.xeno-canto.org) and normalized in amplitude. All the sounds
were broadcasted from a single speaker (Douglas and Mennil 2010). Although a multiple-
speaker approach would have been more realistic, because horneros frequently perform their
duets while close to each other, it is unlikely that our set-up introduced a bias in the STIs. We
simultaneously recorded the behaviour of each individual of the pair during the 20 minutes of
STI (mean ± SE; 19.96 ± 0.23 min). Two observers performed the observations from a distance
of 15 metres using digital voice recorders (Philips VoiceTraicer DVT1200 and Olympus Digital
Recorder VN-733 PC). The focal bird was randomly assigned to each observer. The following
measures were recorded: 1) response latency (time between start of playback and first approach
within 10 metres from the dummy), 2) time spent within five metres of the decoy, 3) average
time spent on the nest during a visit, 4) number of solo songs, 5) number of duet songs, and 6)
number of flights over the decoy (i.e. flights directed to and over the decoy). Regarding the
variable 'number of duets', we initially considered the fact that both males and females can
initiate the duet (Diniz et al, 2018). However, during our STIs there were seldom cases in which
the female initiated the duet, and none were during nest building. For this reason, we only
considered the number of duets as a joint variable across sexes in our models. Additionally, the
solo songs in males represent instances in which females decided not to join in the duet. The
sex of each bird could be determined from the acoustic signature of each individual because the
vocal contribution of each sex in the duet is dimorphic (Roper 2005). As part of a different
project, birds were captured after the STI and the sex was verified by PCR (sex was correctly
assigned by the observers in 96.3% of the cases for those individuals to whom the sex could be
assigned acoustically and were trapped in the nets; for details see Adreani et al. 2018).
209
Statistical analyses
(i) Structural equation modelling
First, we applied a structural equation modelling (SEM) approach to study three *a priori*
hypotheses of relationships among the six behavioural variables quantified during the simulated
territorial intrusion (i.e. response latency, time spent within five metres of the decoy, time spent
on the nest, number of solo songs, number of duet songs, and number of flights over the decoy;
Fig. 1). Of the three models for each sex and context, model 1 represents a (biologically
unrealistic) "null" expectation (i.e. each defensive behaviour is independent and not part of a
functional unit or evolutionary character); model 2 hypothesises that one latent variable,
"territorial defence", underlies the relationships between the six behavioural variables; and

model 3 hypothesises that all behavioural variables except number of duet songs are linked by

the latent variable "territorial defence". While more complex structural models could be

constructed (i.e. including trade-offs between behavioural variables), the present framework is

the one that allows for a straightforward testing the "joint territorial defence" hypothesis (Hall

2009). We estimated each structural equation model separately for males and females because

we only have one measure of the number of duet songs from a single pair and, therefore, it is

not possible to disentangle the sex-differences in number of duets on the latent variable

territorial defence. We also estimated each model separately for the two breeding contexts (i.e.

nest building *vs.* provision context). The formulation of these four different sets of models

allowed us to qualitatively assess whether there were differences between sexes and breeding

contexts (nest building and provisioning) in the structure and strength of the hypothesized latent

variable. Therefore, besides characterizing the latent variable structure, we were also interested

in qualitatively investigating differences between sexes and contexts in path loadings across

models (i.e. whether behavioural traits maintain their rank differences among path loadings).

We also constructed a single model for each sex in both breeding contexts, where the 12

different behavioural variables were modelled simultaneously. However, we decided to present

here the separated models, one for each breeding context, because the full model (i.e., with the

12 variables) is likely over-parametrized (i.e., there was a compromise between the complexity

of the SEM models fitted and the number of observations given the number of variables tested

in each SEM). See Supplementary Material for further details on the full model (Table S1-2

and Fig S1).

To test the relative fit of each alternative biological hypotheses, we first estimated the

matrix of phenotypic correlations of all the behavioural variables for each combination of sex

and breeding context. The correlation matrix was constructed using Spearman coefficients

obtained with the R package "stats" in R v. 3.3.3 (R Core Team 2013). Data on "response

latency" were missing for one out of 38 individual females during nest building and data for

"time spent on the nest" were missing for two out of 24 females during provisioning. We

assigned the average population phenotypic value of each trait to those individuals with missing

values (note that a "complete-case analysis" did not change our findings, results not shown).

We then implemented each among-trait correlation matrix in the R-package "sem" and tested

the different SEM hypotheses. We statistically compared each model's fit using the Akaike

Information Criterion for small sample sizes (AICc; Burnham and Anderson 2004) and

evaluated their relative support based on AICc differences relative to the best-fitting model

(ΔAICc). We also present values for the Goodness of Fit Index (GFI), which represents the

proportion in the improvement of the overall fit of a given hypothesised model compared to the
independence model. GFI values range from 0 (poor fit) to 1 (perfect fit), being considered as
satisfactory when it is higher than 0.9.
257
(ii) Univariate mixed-effect models
As a second step, we investigated sources of variation in territorial defence behaviour in our
population of horneros using a series of univariate mixed models. This step was necessary
because we were also interested in investigating sex-differences in territorial defence. However,
as we only had a single measure of number of duets per pair, we could not test for sources of
variation in territorial defence using our multivariate SEM approach. Thus, we ran two
univariate mixed-effect models fitting number of flights over the decoy and number of duets as
the response variables, respectively. Though we had measured various potential proxies of
territorial defence (detailed above), we used number of flights over decoy and number of duet
songs because they consistently had the highest value in path loading across all models (for a
further discussion on the rationale of this approach, see Araya-Ajoy and Dingemanse 2013).
Breeding context (nest building vs. provisioning), sex (male vs. female) and their interaction,
time of the day (i.e. moment of the day when the territorial intrusion was simulated, expressed
in decimal fractions of hours after sunrise and mean centred), and observer identity (observer 1
vs. 2) were included as fixed effects in the univariate mixed-effect models. Time of day was
mean centred, such that the fixed-effect intercept of the model was estimated for the behavioural
trait on the average time (following Dingemanse and Dochtermann 2013). In the model with
"number of flights over the decoy" as a response variable, we fitted random intercepts for pair
identity ("Pair identity"; $n = 63$ levels). In the model with "number of duet songs", we did not
include this random effect because we did not have repeated measures of duet song frequency
for the same pair identity. Both response variables, number of flights over the decoy and number
of duet songs, were modelled with Poisson errors with a log-link function. In both models, we
included an observation level random effect to account for over-dispersion (Harrison 2014).
The analyses were performed using the R packages "lme4" (Bates et al. 2014) and "arm"
(Gelman and Yu-Sung 2015). We used the "sim" R function to simulate posterior distributions
of the model parameters. Based on 5000 simulations, we extracted the mean value and 95%
Credible Intervals (CrI) of the posterior distributions. Model fit was assessed by visual
inspection of the residuals. Assessment of statistical support was obtained from the posterior
distribution of each parameter (Zuur 2016). We considered an effect "strongly supported" if
zero was not included within the 95% CI, and "moderately supported" if the point estimate was

skewed away from zero while its 95% CI simultaneously overlapped zero. Estimates centred

on zero were viewed as strong support for the absence of an effect.

(iii) Correlation between male and female contribution to territorial defence

We studied the correlation pattern between male and female territorial defence to investigate

whether defence response within a pair was positively correlated. To do so, we estimated the

Spearman correlation coefficient of "number of flights over the decoy" between males and

females from a single pair. This correlation was calculated by including in our analysis all pairs

observed during both contexts ($n = 91$ pairs), given that we found no differences in the number

of flights over the decoy between the nest building and provisioning contexts (see below). The

correlation test was performed using the R-package "stats" in R v. 3.3.3 (R Core Team 2013).

Results

(i) Territorial defence as a latent trait and the role of duetting

The behavioural variables assayed during the simulated territorial intrusion (i.e. response

latency, time spent within five metres of the decoy, time spent on the nest, number of solo

songs, number of duet songs, and number of flights over the decoy) were, to a varying extent,

correlated with each other; both across sexes and contexts (Table S3, S4). Overall, horneros

with shorter latency of response to the territorial intrusion spent more time within five metres

of the decoy and on the nest, sang more solo and duet songs, and flew more often over the

decoy, suggesting the existence of the hypothesised latent variable "territorial defence" linking

the six behavioural variables.

AICc model comparison identified the SEM model 2 as the best one (among the models

we fitted) explaining the structure of the phenotypic variables across the four different set of

models (Table 1). Model 2 represented an overarching latent variable ("territorial defence")

linking the expression of all behavioural variables, including the number of duet songs (Fig.

1A). Furthermore, the number of duet songs together with the number of flights over the decoy

had consistently the highest values of path loading in males and females for both breeding

context (Fig. 1B, Table S5). We thus considered them both equally good predictors for territory

defence in horneros. Furthermore, a standard index of model fit ("Goodness of Fit Index")

considered satisfactory our best fitting model across all models (i.e., GFI values for Model 2

were around or above 0.90 across all models, Table 1).

(ii) Effect of sex, breeding context and time of the day on territorial defence

We did not find strong evidence that horneros differed in the number of duet songs between the

nest building and provisioning context. While the effect size is moderately supported, the

evidence is weak due to large uncertainty (Table 2; Fig. 2A). We did not find differences in the

number of duets explained by time of the day (Table 2). Regarding the number of flights over

the decoy, males were on average defending more (i.e. flew more times over the decoy) than

females during both breeding contexts (Table 2; Fig. 2B). However, we again found weak

evidence that horneros differed in the number of flights over the decoy between the nest

building and provisioning context, and there were also no sex-specific differences between the

two breeding contexts (i.e. the effect sizes are relatively large but estimated with large

uncertainty, therefore the support is moderate; Table 2, Fig. 2B). Furthermore, we observed

moderate effects of time of day and observer identity in our model (the estimates include zero

in their 95% CrI, but the effect sizes are considerable).

(iii) Defending as a unit: correlation between male and female territorial defence

We investigated whether the defensive response to a territory intrusion of an individual was

correlated with the response expressed by its partner. We found that territorial defence of males

and females within pairs was strongly positively correlated ($\rho = 0.67$, $p < 0.0001$). Thereby,

within a single pair, male and female had matching levels of defensive response (Fig. 3).

Discussion

We studied territorial defence in both male and female horneros and the role of duets during

nest building and provisioning contexts. By using structural equation modelling, we were able

to demonstrate that six observed behavioural variables (i.e. response latency, time spent within

five metres of the decoy, average time spent on the nest during a visit, number of solo songs,

number of duet songs, and number of flights over the decoy) were linked by an unmeasured

latent trait "territorial defence", both across sexes and contexts (Fig. 1). We also found that the

number of flights over the decoy and duet songs were the variables with highest path loading

to territorial defence. We then proceed to study independently those two traits with highest path

loadings and showed that males were defending territories more strongly than were females

during both breeding contexts, even though we only found weak evidence for differences

between contexts within each sex (owing to large uncertainty in our estimate, Fig. 2). Lastly

and as expected, we observed a strong positive correlation in territorial defence between males

and females within the pair (Fig. 3).

The correlation structure of the latent variable territorial defence was similar in males

and females, providing for the first time a sex-comparison of the latent variable "territorial

defence" in birds. We also observed that the sign and magnitude of the path loadings between

breeding contexts (i.e. nest building and provisioning) were very similar (Fig. 1B). Our results

thus suggest the existence of a common structure in defensive behaviour during territorial

intrusion in horneros, potentially underlined by a sex- and context-independent mechanism that

is affecting all behavioural variables in a similar manner. However, to draw general

evolutionary patterns of territorial defence it would be necessary to investigate whether the

described latent variable is under selection as an integrated trait (i.e. functional module) and

whether the same structure among traits is observed in other life-history stages (e.g. outside the

breeding season) or in different ecological contexts (e.g. as territorial defence against predators

or nest-parasitic species). Importantly, by using a structural equation modelling approach we

were able to reveal complex relationships for multiple traits that would have otherwise not been

possible to capture. Traditional statistical approaches such as multiple regression analysis or

principal component analysis (PCA) are not suitable to evaluate different *a priori* defined

hypotheses while accounting for trait correlation. PCAs are defined purely on the basis of

mathematical associations between the traits and so their biological meaning can be challenging

to interpret or even nonexistent. However, SEM provides a more interpretable method of

viewing variation among correlated variables. Although SEM will ultimately be defined by

mathematical associations like PCAs, with SEM, one makes use of biological information to fit

the correlation structure to be tested among variables. Therefore, SEM has a clear advantage

over PCA in terms of making biological inferences from the data. Lastly, another advantage of

using a structural equation modelling approach is ralated to data collection methods. The fact

that the number of flights over the decoy and duets songs were the variables with the highest

path loadings to territorial defence indicates that by measuring only these two observable

behaviours, and not all six, researchers should have a good estimation of territorial defence – at

least in those studies that aim to quantify territorial defence towards conspecifics in horneros.

Nonetheless, a good biological use of the approach would require a validation of the method

for each new species where territorial defence is to be characterized.

In the context of territorial defence, duets have been mostly studied as a single trait

independently of other complementary or related behaviours in multiple bird species (e.g. Hall

and Peters 2008; Dowling and Webster 2016; Odom et al. 2017; Quirós-Guerrero et al. 2017).

Here we explicitly tested for the first time whether duets are a behavioural variable linked by a

latent trait, "territorial defence". We did so by combining a classical STI approach with

structural equation modelling. One of the predictions of the "joint territorial defence"
hypothesis is that duets should play a more important role than solo songs (Hall 2009). As
expected, in our study duets represented an important response during territorial defence for
both sexes and breeding contexts. They were stronger than solo songs and were as relevant as
other physical traits like the number of flights over the decoy. Our results are in line with
previous findings in the species suggesting the territorial function of duets in hornero, overall
providing evidence for the "joint territorial defence" hypothesis (Diniz et al. 2018, 2019, 2020).
While previous studies tested this hypothesis by means of different approaches, the strength of
our study resides in the application of a comprehensive method that accounts for the
multivariate nature of territorial defence behaviours.

Males defended more their territories than females during both breeding contexts. At
first glance, this is not surprising given that an unequal sex contribution of territorial defence
has been previously reported in bird species that are socially monogamous and maintain
territories year round (e.g. Willis 1972; Morton and Derrickson 1996; Bard et al. 2002; Fedy
and Stutchbury 2005; Quinard and Cézilly 2012). In the specific case of horneros, however,
male and female have been reported to contribute equally in most of the behaviours studied to
date (Fraga 1980; Massoni et al. 2012; Diniz et al. 2020). However, there is strong evidence for
sexual differences in singing-related traits independent of season (Diniz et al. 2018).
Additionally, the observation that males engaged more in aggressive interactions than females
has only been recently described (Diniz et al. 2018). Thus, our findings confirm and expand
this observation with a standardized field test applied to a multivariate framework beyond song
production. One explanation for the observed sex-differences might be a division of labour
between members of a pair (e.g. Morton et al. 2000). For instance, males might invest more
resources (i.e., time and energy) in actively defending their territory or nest (e.g. physical
attacking the intruder), whereas females might focus on different activities (e.g. predator
vigilance, guarding the nest against parasitic species). Another factor potentially explaining our
results is that males and females might face different physiological (breeding) costs (e.g.
Nilsson and Råberg 2001). In fact, during nest building (when females are close to egg laying)
females have a poorer oxidative condition than males and are more sensitive to STIs, suggesting
a sex-specific physiological cost of territorial defence (Mentesana and Adreani 2020). Lastly,
our findings could also be influenced by the way the territorial intrusions were performed (i.e.
with one single dummy). While plausible, this explanation seems unlikely given that horneros
are monomorphic in body size and plumage colouration (Diniz et al. 2016) and the playbacks
consisted of vocalizations from both sexes.

We did not find strong support for our prediction that the levels of territorial defence

were higher during the nest building than in the provisioning context (Table 2). Given that

extra-pair levels are very low in this species (~3%, Diniz et al. 2019), one of the main

assumptions of our prediction was that territory take-over was higher during the fertile period

of the females, i.e., during nest building than during provisioning (Gill et al. 2007; Demko and

Mennill 2018). It is possible that for horneros it is more beneficial to maintain constant levels

of territorial defence in order to hold the territory year-round than the potential benefits of extra-

pair paternity (Warner and Hoffman 1980). This might be especially the case when population

densities are high, where comparable territorial defence can be expected across different life-

stages as we observed in the horneros. Further research will help to shed light on these context-

specific patterns.

Male and female aggression were strongly and positively correlated within the pair

despite sex-specific differences in territorial defence. Our results are in line with previous

findings of coordinated territorial defense on rufous horneros outside the breeding season

(Diniz et al. 2020) and more generally with other studies showing that duetting birds were more

collaborative within the pair than non-duetting species (see Logue 2005).Although our study

cannot directly address the evolutionary relevance of pairs being positively correlated in their

behaviours (e.g., fitness consequences), our findings suggest that exhibiting a joint territorial

defence might be an important mechanism of pair bonding or pair stability (Wickler and Seibt

1980). In this direction, our study raises the question of whether pairs of horneros that show

similar territorial defence levels would experience increased reproductive benefits (Schuett et

al. 2010). Indeed, it is known from other bird species that pairs exhibiting comparably high

levels of territory defence towards conspecifics attain higher reproductive success (e.g. in

eastern blue birds, *Sialia sialis;* Harris and Siefferman 2014).Therefore, investigating patterns

of selection on assortative mating in pairs of horneros poses an exciting avenue for future

research.

Conclusion

This work expands upon a classical body of research on territorial defence. We demonstrated

that six observed behavioural variables quantified during a simulated territorial intrusion were

linked by an unmeasured latent trait "territorial defence". In particular, the number of flights

over the decoy and the number of duet songs were the variables with highest path loadings to

the latent variable "territorial defence". Furthermore, this study fills an important gap in our

knowledge about the role of duets. We provided support for the hypothesis that avian duets are

a key component in the joint territory defence. Indeed, we showed that duets represented a

stronger response of territory defence than solo songs, and that their importance was

comparable to physical traits. Our study also highlights the importance of using more

integrative, multivariate approaches to study behavioural traits. By applying a structural

equation modelling framework, we were able to evaluate *a priori* hypotheses of how different

behavioural variables were linked by an unmeasured latent trait. Such complex patterns would

have not been possible to capture using traditional statistical approaches such as principal

component analyses Hence, the combination of a classical behavioural approach like simulated

territorial intrusions with structural equation modelling brings new exciting possibilities into

the field of behavioural ecology.

Authors' contributions

NMA, LM and MM share first authorship and names are ordered at random. NMA and LM conceived the study and designed the study. BT provided logistic support. NMA, LM, EG and EC collected the data. NMA, LM, and MM analyzed the data. NMA, LM, and MM wrote the manuscript with input from all authors.

Acknowledgments

We thank Klaus Pichler for his help to prepare the STI and all the "INIA Las Brujas" staff for supporting us with accommodation and equipment during fieldwork. We also thank Facultad de Ciencias and the Ethology lab from Universidad de la República and Juan Carlos Reboreda and the "Laboratorio de Ecología y Comportamiento Animal" at the Universidad de Buenos Aires for the logistical support, and Pablo Tubaro from the Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia' (MACN) for providing us with the mounted hornero. We are grateful to Manfred Gahr and Michaela Hau for their valuable support. We also thank Yimen Araya-Ajoy, Glenn Cockburn, Luke Eberhart-Phillips and Wolfgang Wickler for constructive criticism on previous versions of the manuscript. Finally, we want to thank the anonymous reviewers for constructive feedback on the manuscript.

Availability of data

The datasets generated and/or analysed during the current study are available in the open repository: Mendeley Data (https://data.mendeley.com/datasets/7ztwn539jd/1).

Compliance with ethical standards

Ethical approval

The experimental procedures of this study have approval by the Ethics Committee of Animal Experimentation (CEUA) of the Facultad de Ciencias of the Universidad de la República, Uruguay (Protocol number 186, file 2400-11000090-16).

Conflict of interest

The authors declare they have no conflicts of interest.

Funding

This work was funded by the International Max Planck Research School (IMPRS) for Organismal Biology, and by Idea Wild that provided field equipment.

References

Adreani NM, Goymann W, Mentesana L (2018) Not one hormone or another: Aggression
differentially affects progesterone and testosterone in a South American ovenbird. Horm Behav
105:104–109

Apfelbeck B, Goymann W (2011) Ignoring the challenge? Male black redstarts (*Phoenicurus*
ochruros) do not increase testosterone levels during territorial conflicts but they do so in
response to gonadotropin-releasing hormone. Proc R Soc Lond B 278:3233–3242

Araya-Ajoy YG, Dingemanse NJ (2013) Characterizing behavioural "characters": an
evolutionary framework. Proc R Soc B 281:20132645

Bard SC, Hau M, Wikelski M, Wingfield JC (2002) Vocal distinctiveness and response to
conspecific playback in the spotted antbird, a neotropical suboscine. Condor 104:387-394

Bates D, Mächler M, Bolker B, Walker S (2014) Fitting linear Mixed-Effects Models using
lme4, https://cran.r-project.org/web/packages/lme4/index.html

Bollen KA (2002) Latent variables in psychology and the social sciences. Annu Rev Psychol
53:605–634

Burnham KP, Anderson DR (2004) Multimodel inference. Sociol Methods Res 33:261–304

Carter AJ, Feeney WE (2012) Taking a comparative approach: Analysing personality as a
multivariate behavioural response across species. PLoS ONE 7:e42440

Card NA, Little TD (2007) Studying agression with structural equation modeling. In: Flannery
DJ, Vazsonyi AT, Waldman ID (eds) The Cambridge handbook of violent behavior and
aggression. Cambridge University Press New York, NY, pp 727-739

Clutton-Brock TH, Vincent ACJ (1991) Sexual selection and the potential reproductive rates
of males and females. Nature 351:58–60

Demko AD, Mennill DJ (2018) Male and female signaling behavior varies seasonally during

territorial interactions in a tropical songbird. Behav Ecol Sociobiol 72:84

Dingemanse NJ, Dochtermann NA (2013) Quantifying individual variation in behaviour:
mixed-effect modelling approaches. J Anim Ecol 82:39–54

Diniz P, Júnior EF da S, Webster MS, Macedo RH (2018) Duetting behavior in a Neotropical
ovenbird: sexual and seasonal variation and adaptive signaling functions. J Avian Biol 49:jav-
01637

Diniz P, Macedo RH, Webster MS (2019) Duetting correlates with territory quality and
reproductive success in a suboscine bird with low extra-pair paternity. Auk 136:1-13

Diniz P, Rech G, Ribeiro PH, Webster MS, Macedo RH (2020) Partners coordinate territorial
defence against simulated intruders in a duetting ovenbird. Ecol Evol 10:81-89

Diniz P, Ribeiro PHL, Rech GS, Macedo RH (2016) Monochromatism, cryptic sexual
dimorphism and lack of assortative mating in the Rufous Hornero, *Furnarius rufus albogularis*.
Emu 116:294–300

Dowling J, Webster MS (2016) An experimental test of duet function in a fairy-wren (*Malurus*)
with moderate cuckoldry rates. Behav Ecol 27:228–236

Emlen ST, Oring LW (1977) Ecology, sexual selection, and the evolution of mating systems.
Science 197:215–223

Farabaugh SM (1982) The ecological and social significance of duetting.
In: Kroodsma DE, Miller EH (eds) Acoustic Communication in Birds, vol. 2. Academic Press,
New York, pp 85–124

Fedy BC, Stutchbury BJM (2005) Territory defence in tropical birds: are females as aggressive
as males? Behav Ecol Sociobiol 58:414–422

Fraga RM (1980) The breeding of Rufous Horneros (*Furnarius rufus*). Condor 82:58–68

Freeman BG, Montgomery GA, Schluter D (2017) Evolution and plasticity: Divergence of song
discrimination is faster in birds with innate song than in song learners in Neotropical passerine
birds. Evolution 71:2230-2242

Garcia JT, Arroyo BE (2002) Intra- and interspecific agonistic behaviour in sympatric harriers
during the breeding season. Anim Behav 64:77–84

Gelman A, Yu-Sung S (2015) arm: Data analysis using regression and multilevel/hierarchical
models. R package version 1.8-5, https://cran.r-project.org/web/packages/arm/arm.pdf

Gill SA, Alfson ED, Hau M (2007) Context matters: female aggression and testosterone in a
year-round territorial neotropical songbird (*Thyrothorus leucotis*). Proc R Soc Lond B
274:2187–2194

Greenberg R, Gradwohl J (1983) Sexual roles in the dot-winged antwren (*Microrhopias*
quixensis), a tropical forest passerine. Auk 100:920–925

Hall ML (2004) A review of hypotheses for the functions of avian duetting. Behav Ecol
Sociobiol 55:415–430

Hall ML (2009) A review of vocal duetting in birds. Adv Stud Behav 40:67–121

Hall ML, Peters A (2008) Coordination between the sexes for territorial defence in a duetting
faire-wren. Anim Behav 76:65–73

Harris MR, Siefferman L (2014) Interspecific competition influences fitness benefits of
assortative mating for territorial aggression in eastern bluebirds (*Sialia sialis*). PLoS ONE
9:e88668

Harrison XA (2009) Using observation-level random effects to model overdispersion in count
data in ecology and evolution. PeerJ 2:e616

Houle D, Pélabon C, Wagner GP, Hansen TF (2011) Measurement and meaning in biology. Q
Rev Biol 86:3-34

Huntingford FA (1976) An investigation of the territorial behaviour of the three-spined

stickleback (*Gasterosteus aculeatus*) using principal components analysis. Anim

Behav 24:822-834

Koloff J, Mennill DJ (2013) The responses of duetting antbirds to stereo duet playback provide

support for the joint territory defence hypothesis. Ethology 119:462–471

Korner-Nievergelt F, Roth T, von Felten S, Guélat J, Almasi B, Korner-Nievergelt P (2015)

Bayesian data analysis in ecology using linear models with R, BUGS, and Stan. Academic

Press, New York

Langmore NE (1998) Functions of duet and solo songs of female birds. Trends Ecol Evol

13:136–140

Massoni V, Reboreda JC, López GC, Aldatz MF (2012) High coordination and equitable

parental effort in the Rufous Hornero. Condor 114:564–570

Mentesana L, Adreani NM (2020) Is aggression costly? Acute aggressive behavior increases

oxidative stress independently of testosterone. bioRxiv,

https://doi.org/10.1101/2020.04.06.027029

Morton ES, Derrickson KC (1996) Song ranging by the dusky antbird, *Cercomacra tyrannina*:

ranging without song learning. Behav Ecol Sociobiol 39:195–201

Morton ES, Derrickson KC, Stutchbury BJM (2000) Territory switching behavior in a

sedentary tropical passerine, the dusky antbird (*Cercomacra tyrannina*). Behav Ecol 11:648–

653

Naguib M, Mennill DJ (2010) The signal value of birdsong: empirical evidence suggests song

overlapping is a signal. Anim Behav 80:e11-e15

Nilsson JÅ, Råberg L (2001) The resting metabolic cost of egg laying and nestling feeding in

great tits. Oecologia 128:187–192

Odom KJ, Logue DM, Studds CE, Monroe MK, Campbell SK, Omland KE (2017) Duetting

behavior varies with sex, season, and singing role in a tropical oriole (*Icterus icterus*). Behav

Ecol 28:1256–1265

Owens IPF, Thompson DBA (1994) Sex differences, sex ratios and sex roles. Proc R Soc Lond

B 258:93–99

Quinard A, Cézilly F (2012) Sex roles during conspecific territorial defence in the Zenaida

dove, *Zenaida aurita*. Anim Behav 83:47–54

Quirós-Guerrero E, Janeiro MJ, Lopez-Morales M, Cresswell W, Templeton CN (2017)

Riverside wren pairs jointly defend their territories against simulated intruders. Ethology

123:949–956

R Core Team (2013) R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria, http://www.R-project.org

Robinson A (1949) The biological significance of bird song in Australia. Emu 48:291–315

Roper JJ (2005) Sexually distinct songs in the duet of the sexually monomorphic Rufous

Hornero. J Ornithol 76:234–236

Schuett W, Tregenza T, Dall SRX (2010) Sexual selection and animal personality. Biol Rev

85:217–246

Searcy WA, Beecher MD (2009) Song as an aggressive signal in songbirds. Anim

Behav 78:1281-1292

Smith BR, Blumstein DT (2008) Fitness consequences of personality: a meta-analysis. Behav

Ecol 19:448–455

Sosa-López JR, Mennill DJ, Renton K (2017) Sexual differentiation and seasonal variation in

response to conspecific and heterospecific acoustic signals. Ethology 123:460–466

Sprenger D, Dingemanse NJ, Dochtermann NA, Theobald J, Walker SP (2012) Aggressive females become aggressive males in a sex-changing reef fish. Ecol Lett 15:986-992

Stamps JA, Krishnan VV (1997) Functions of fights in territory establishment. Am Nat 150:393–405

Tobias JA, Sheard C, Seddon N, Meade A, Cotton AJ, Nakagawa S (2016) Territoriality, social bonds, and the evolution of communal signaling in birds. Front Ecol Evol 4:74

Topp SM, Mennill DJ (2008) Seasonal variation in the duetting behaviour of rufous-and-white wrens (*Thryothorus rufalbus*). Behav Ecol. Sociobiol 62:1107–1117

Warner RR, Hoffman SG (1980) Population density and the economics of territorial defence in a coral reef fish. Ecology 61:772–780

Wickler W, Seibt U (1980) Vocal dueting and the pair bond. Z Tierpsychol 52:217–226

Willis EO (1972) The Behavior of ocellated antbirds. Ornithol Monogr 10:1–162

Wingfield JC (1994) Regulation of territorial behavior in the sedentary song sparrow, *Melospiza melodia morphna*. Horm Behav 28:1–15

Table and Figure captions

Table 1 Results of model comparison using Akaike Information Criterion for small sample

sizes (AICc) values to compare our three candidate models. Smaller AICc values are given to

models that better fit the data. Models whose AICc values differ from that of the top model

(ΔAICc) by more than 2 are considered to lack explanatory power relative to the top model.

We also present values for the Goodness of Fit Index (GFI). The best-supported hypothesis is

printed in boldface

Table 2 Sources of variation in "number of flights over the decoy" and "number of duets" in

horneros. Breeding context (nest building vs provisioning), sex (female vs male) and their

interaction; time of the day (hours after sunrise, mean centred); and observer identity (Observer

1 vs 2) were fitted as fixed effects. Pair identity and an observation-level parameter were fitted

as random effects. Both response variables were modelled with Poisson error. We present

estimates of fixed (β) and random (σ^2) parameters with their 95% Credible Intervals (CrI) in

brackets. The reference category for the categorical variable sex is "female"; for breeding

context, "nest building"; and for observer identity is "observer 1"

Fig. 1 (A) Three models (hypotheses) explaining the correlation structure among behavioural

variables assayed during a simulated territory intrusion in the nest building and provisioning

context in wild horneros. Model 1 hypothesises trait independence (null model); model 2

hypothesises a latent variable ("territorial defence") linking all behavioural variables, whereas

model 3 hypothesises a latent variable ("territorial defence") linking all behavioural variables

except "number of duets". Unidirectional arrows represent directional relationships between

traits. Solid lines represent relationships present across all models; and the dashed line

represents a relationship expressed in a specific model structure. Path "a" is only active in model

2. (B) Path loadings of model 2 for males and females in both breeding contexts. Squares and

circles represent the estimated mean, and error bars represent the standard error of the mean

Fig. 2 Number of (A) duets and (B) flights over decoy of hornero pairs during nest building

and provisioning context. Grey symbols represent raw data. The mean estimates of the posterior

distributions (black symbols) as well as the 95% credible intervals (error bars) are also shown

Fig. 3 Female-male correlation of territorial defence, using number of flights over the decoy as

proxy. The black line represents the regression line and the dashed line is the reference line

with a slope of 1

Table 1

Sex – breeding context	SEM models								
	Model 1			**Model 2**			**Model 3**		
	AICc	ΔAICc	GFI	AICc	ΔAICc	GFI	AICc	ΔAICc	GFI
female - nest building	72.64	44.53	0.60	**28.11**	**0.00**	**0.90**	51.11	23.00	0.79
female - provisioning	55.74	19.43	0.58	**36.31**	**0.00**	**0.89**	54.06	17.75	0.78
male - nest building	30.53	11.60	0.81	**18.93**	**0.00**	**0.95**	30.99	12.06	0.87
male - provisioning	62.17	16.11	0.54	**46.06**	**0.00**	**0.84**	49.40	3.34	0.80

735 **Figure 1**

 Table 2

| | **Flights over decoy** | **Duets** |
Fixed effects	**β (95% CrI)**	**β (95% CrI)**
Intercept	2.33 (1.87, 2.78)	2.44 (2.21, 2.69)
Sex	0.58 (0.32, 0.83)	--
Breeding context	-0.28 (-0.88, 0.31)	-0.11 (-0.52, 0.28)
Sex × Breeding context	0.22 (-0.19, 0.64)	--
Time of day	-0.19 (-0.45, 0.05)	-0.07 (-0.26, 0.13)
Observer Identity	-0.16 (-0.36, 0.04)	--
Random effects	**σ2 (95%CrI)**	**σ2 (95%CrI)**
Pair Identity	1.02 (0.75, 1.36)	--
Observation–level parameter	0.18 (0.14, 0.24)	0.44 (0.33, 0.60)

745 **Figure 2**

748   **Figure 3**