Single molecule dynamics at a bacterial replication fork after nutritional downshift

Rogelio Hernández-Tamayo[1,2#], Hannah Schmitz[1,2] and Peter L. Graumann[1,2 #].

[1]SYNMIKRO, LOEWE Center for Synthetic Microbiology, Marburg, Germany,

[2]Department of Chemistry, Philipps Universität Marburg, Marburg, Germany,

KEY WORDS: replication, single molecule microscopy, DNA primase, helicase, stringent response, *Bacillus subtilis*.

RUNNING TITLE: Dynamics of DNA primase and helicase at the single molecule level

ADDRESS FOR CORRESPONDENCE:
[#]Peter L. Graumann, peter.graumann@synmikro.uni-marburg.de
[#]Rogelio Hernández-Tamayo, rogelio.hernandez@synmikro.uni-marburg.de

SYNMIKRO, LOEWE Center for Synthetic Microbiology and Department of Chemistry, Philipps Universität Marburg, Marburg, Germany, Hans-Meerwein-Straße 6, 35032 Marburg, Germany. Phone number: +49-6421 28 22210

ABSTRACT

Replication forks must respond to changes in nutrient conditions, especially in bacterial cells. By investigating the single molecule dynamics of replicative helicase DnaC, DNA primase DnaG, and of lagging strand polymerase DnaE in the model bacterium *Bacillus subtilis* in response to transient replication blocks due to DNA damage, to inhibition of the replicative polymerase, or to downshift of serine availability, we show that proteins react differentially to the stress conditions. DnaG appears to be recruited to the forks by a diffusion and capture mechanism, becomes more statically associated after arrest of polymerase PolC, but binds much less often after fork blocks due to DNA damage or to nutritional downshift. These results indicate that binding of the alarmone ppGpp due to the stringent response prevents DnaG from binding to forks rather than blocking bound primase. Dissimilar behaviour of DnaG and of DnaE suggest that both proteins are recruited independently to the forks, rather than jointly. Turnover of all three proteins was increased during replication block after nutritional downshift, different from the situation due to DNA damage or polymerase inhibition, showing high plasticity of forks in response to different stress conditions. Forks persisted during all stress conditions, apparently ensuring rapid return to replication extension.

INTRODUCTION

DNA replication is a highly choreographed and tightly regulated event in the lifecycle of all cells. It is carried out by a dynamic, multi-protein complex known as the replisome (1), which precisely coordinates the action of several distinct factors to efficiently and rapidly couple DNA unwinding with high-fidelity nucleic acid synthesis (2-4). Importantly, DNA replication must respond to situations of changing environmental or developmental conditions, including response to damage in the template, and reduced nutritional capacity of cells (5). This is especially relevant for bacterial cells that are directly affected by changes in their surroundings (6-8).

DNA replication is inherently asymmetric; one daughter strand, termed the leading strand, is continually synthesized in the same direction as the unwinding of the DNA duplex. The other (lagging) strand is synthesized in the opposite direction in short intervals, giving rise to Okazaki fragments (9). The Gram-positive model organism *Bacillus subtilis* and the rest of the Firmicutes (low-G+C-content Gram-positive bacteria) use two essential DNA polymerases, DnaE and PolC, during replication (10,11). *In vivo*, PolC is the main replicative polymerase, while DnaE acts in lagging strand synthesis. *In vitro* reconstitution of the *B. subtilis* replisome has demonstrated that PolC is responsible for all leading strand synthesis as well as most lagging strand synthesis, whereas the more error prone and much slower DNA replicase DnaE (25-60 nt/s for DnaE compared to ~500 nt/s for PolC) plays a crucial role in initiating lagging strand synthesis. DnaE is important for extending the lagging strand RNA primer before handing off to PolC, which then completes replication of the Okazaki fragment (11,12). The synergistic relationship between two polymerases in the *B. subtilis* replisome resembles the synergy found in eukaryotic replication (11,13).

Replication of the lagging strand requires a specialized RNA polymerase, termed primase, to initiate each Okazaki fragment with a short oligoribonucleotide (14). In *B. subtilis* the primase, DnaG, is recruited to the replication fork by an interaction with the replicative helicase, DnaC, where it synthesizes an RNA primer every 1.5–2 kb (1). Working together, the helicase and primase unwind the DNA template and initiate thousands of regularly spaced Okazaki fragments to promote fork progression at a rate of 1000 bases per second in rapidly dividing bacterial cells.

The direct association of primase and helicase co-regulates their functions. For
example, primase increases both the ATPase and helicase activities of DnaC. Similarly,
DnaC can modulate the overall activity of DnaG, as well as the length of primers
synthesized by primase, and its initiation specificity (15). Each Okazaki fragments is
initiated by a short RNA primer by DnaG for extension by DnaE. Primase requires
interaction with the helicase to stimulate its RNA polymerase activity (16). In *E. coli* the
helicase-primase contact is established through the interaction between the C-terminal
domain of primase (17,18) and the N-terminal domains of the helicase (19). In *E. coli*, this
interaction is weak and transient, with a K_D in the low micromolar range, giving rise to fast
on/off kinetics (18). Yet, in *Geobacillus stearothermophilus*, helicase and primase form a
stable complex that can be isolated and crystallized (18,20).

Bacteria respond to nutrient downshift by the so-called stringent response. Non-
charged tRNAs arise when a shortage of any amino acid occurs, and bind to the A-site
on the ribosome, where they are sensed by RelA (21,22). This multifunctional enzyme in
turn is activated and converts GTP into ppGpp (guanosine 3',5'-bispyrophosphate) or
pppGpp (guanosine 3'-diphosphate 5'-triphosphate), a second messenger that triggers
many events including downregulation of translation, and increasing the synthesis of
enzymes for e.g. amino acid synthesis pathways (23). Interestingly, (p)ppGpp also binds
to DnaG, whereby DNA replication is greatly slowed down, or even stopped, until nutrient
shortage is overcome (24). The stringent response induces arrest of DNA replication in
B. subtilis and, to a lesser extent, in *E. coli* (25,26). Interference of (p)ppGpp with the
activity of DNA primase inhibits replication elongation in a dose-dependent manner and
adjusts elongation rate according to the nutritional status of the cell (27,28).

We were interested in finding out more exactly how replication forks are adapted
during nutritional downshift inducing the stringent response, and in comparing changes
in molecule dynamics occurring in response to DNA damage-induced blocks of replication
forks, or by specifically blocking PolC. By generating functional fluorescent protein fusions
to DnaC and to DnaG, and employing a partially characterized fluorescent protein fusion
to DnaE, we find distinct changes at the single molecule level to the three scenarios
described above, showing that replication forks have a high plasticity to deal with different
stress situations in a bacterial cell.

MATERIALS AND METHODS

MATERIALS AND METHODS

Bacterial Strains and Growth Conditions.

The bacterial strains and plasmids used in this study are listed in Table S1, and the

nucleotides are listed in Table S2. *Escherichia coli* strain DH5α (Stratagene) was used

for the construction and propagation of plasmids. All *Bacillus subtilis* strains were derived

from the wild-type strain BG214. Cells were grown in Luria-Bertani (LB) rich medium at

30°C. When needed, antibiotics were added at the following concentrations (in µg/ml):

ampicillin, 100; chloramphenicol, 5; spectinomycin, 100; kanamycin, 30. When required,

media containing mitomycin C (MMC), 50 µg/ml; 6(p-Hydroxyphenylazo)-uracil (HPUra),

50 µg/ml; or D,L-Serine Hydroxamate (SHX) 5 mg/ml were prepared by adding

appropriate volumes of a filter-sterilized solution.

Construction of strains

DnaE, DnaG and DnaC were visualized as a DnaE-mVenus, DnaG-mVenus and DnaC-

mVenus ("mV"), fusion proteins expressed at the original locus. The last 500 bp coding

for each gene were integrated into vector pSG1164-mVenus (29), using *Apa*I and *EcoR*I

restriction sites, and BG214 cells were transformed with this construct, selecting for Cm

resistance (leading to strains Table S1). For colocalization studies, DnaX-CFP was

integrated at *amyE* locus by the use of the plasmid pSG1192 and expression was

controlled by xylose addition (30). To investigate colocalization of DnaE, DnaG and DnaC,

the resulting strains PG3307, PG3322 and PG3323 (see Table S1) were transformed with

chromosomal DNA of strains leading to the expression of DnaX-CFP in parallel to DnaE-

mV, or to DnaG-mV, or to DnaC-mV.

Fluorescence Microscopy.

For fluorescence microscopy, *B. subtilis* cells were grown in LB at 30ºC under shaking

conditions until exponential growth. Conventional light microscopy was performed using

a Zeiss Observer Z1 (Carl Zeiss) with an oil immersion objective (100 x magnification,

1.45 numerical aperture, alpha Plan-FLUAR, Carl Zeiss) and a CCD camera (Cool SNAP

EZ, Photometrics). Data were processed using Metamorph 7.5.5.0 software (Molecular

Devices, Sunnyvale, CA, USA).

Single molecule microscopy and tracking.

Cells were spotted on coverslips (25 mm, Menzel) and covered using 1% agarose pads prepared before with fresh $S7_{50}$ minimal medium by sandwiching the agarose between two smaller coverslips (12 mm Marienfeld). All coverslips were cleaned before use by sonication in Hellmanex II solution (1% v/v) for 15 min followed by rinsing in distilled water and a second round of sonication in double distilled water. In contrast to the wide-field illumination used in conventional epifluorescence microscopy, the excitation laser beam used in our setup is directed to underfill the back aperture of the objective lens, generating a concentrated parallel illumination profile at the level of the sample, leading to a strong excitation followed by rapid bleaching of the fluorophores. When only a few unbleached molecules are present, their movement can be tracked. In addition, freshly synthesized and folded fluorophores become visible when the sample is excited again. When an observed molecule is bleached in a single step during the imaging, it is assumed to be a single molecule (4,31). Image acquisition was done continuously during laser excitation with the electron-multiplying CCD (EMCCD) camera iXon Ultra (Andor Technology, Belfast, UK). A total of 2,500 frames were taken per movie, with an exposure time of 20 ms (23 frames per second [fps]). The microscope used in the process was an Olympus IX71, with a ×100 objective (UAPON 100×OTIRF; numerical aperture [NA], 1.49; oil immersion). A 514-nm laser diode was used as excitation source, and the band corresponding to the fluorophore was filtered out. Of note, cells continued to grow after imaging, showing that there is little to no photodamage during imaging, while cells stop growing when exposed to blue light (below 480 nm). Acquired streams were loaded into Fiji ImageJ (32). Automated tracking of single molecules was done using the ImageJ plugin MtrackJ, or u-track 2.1.3 (33).

Diffusion analysis of single-molecule tracks.

Tracking analysis was done with u-track-2.1.3, which was specifically written for Matlab (MathWorks, Natick, MA, USA). Only trajectories consisting of a minimum of 5 frames were considered tracks and included for further analysis. A widely accepted method to analyse the diffusive behaviour of molecules is by using the mean squared displacement

(MSD)-versus-time-lag curve (34,35). This provides an estimate of the diffusion

coefficient as well as of the kind of motion, e.g., diffusive, sub-diffusive or directed.

However, the method requires that within a complete trajectory there be only one type of

homogeneous motion and that the trajectory is preferably of infinite length. To distinguish

immobile and mobile molecules from each other, we compare the frame-to-frame

displacement of all molecules in x and the y directions. Using a Gaussian mixture model

(GMM) to fit the probability density distribution function of all frame-to-frame

displacements, determine the standard deviations σ_1 and σ_2, as well as the percentages

F_1 and F_2 of the slow and the fast subfractions of molecules, respectively. Finally, the

diffusion constants were calculated according $D_i = \frac{\sigma^2}{2\Delta t}$, $(i = 1,2)$ where Δt is the time

interval between subsequent imaging frames.

Generation of heat maps, analyses of molecule dwell times, and visualization of slow

and fast tracks in a standardized cell are based on a custom-written Matlab script

(SMTracker) that is available on request (36). SMTracker can use particle tracking tools

u-track (33) and TrackMate (37) and computes the x- and y-coordinates of molecular

trajectories relative to the geometry of each cell, as obtained by the cell segmentation

tools MicrobeTracker (34) or Oufti (38).

Dwell times.

Dwell time is defined as the average duration that a particle stays inside a certain region.

Observing the trajectories in this manner could give insights, for example, on how long

the replication proteins are bound at the replication fork. For that matter, dwell time

calculations need as parameters the (circular) region, and the minimum number of steps

that a molecule should remain inside the region (1 step = 1 interval time). The procedure

operates in such a way that searches for the longest dwell events of the protein in each

trajectory. For T = $(C_1,...C_n)$ a trajectory is defined as a set of nodes $C_i = (x_i,y_i)$, and x_i,y_i the

nodes' coordinates in a cartesian axis, the circle $C(C_k ,R)$ is chosen, with R being the

radius that contains the maximum number of consecutive points of the trajectory. Then,

the amount of time the molecule stays is counted and the same track T excluding that

segment of trajectory, $T/(C_k,...C_{k+p})$, is again searched for more dwell events. The

procedure finishes when no more dwell events can be found. In our procedure, one gap

(point absent for one frame) or one point outside the circle that goes and comes back are
also considered to have remained inside the circle. The number of dwell events and their
frequency is plotted in a pdf-histogram, and the data are fitted to a single – or if
appropriate, multi-exponential decay, in order to distinguish up to two different
populations of dwell times events.
214
Statistical data analysis
The goodness of fits of the Gaussian mixture models was assessed using probability-
probability plots (pp-plots). Errors on the fitted parameters are given as 95% confidence
intervals, which were derived from the Jacobian matrix of the nonlinear optimization
process using the MATLAB™ function *nlparci.* To compare fraction sizes and diffusion
constants under different conditions and between different proteins, statistical hypothesis-
testing was performed using *Z*-tests. Differences in dwell time and step size distributions
were tested using a Kolmogorov-Smirnov 2-sample test. To assess the most likely
number of populations for each fit, we applied the Bayesian Information Criterion (BIC),
as detailed in (36). As usual, ** and *** stands for p-values lower than 0.01 and 0.05
respectively, while n.s. means statistically "not significant". Statistical hypothesis testing
and plotting was performed using SMTracker (36) and MATLAB™ custom written scripts.
227
228

RESULTS

We wished to gain insight into the changes of dynamics occurring at *Bacillus subtilis* replication forks in response to conditions inducing a transient block in DNA replication, including the response to nutritional downshift, termed stringent response. Our strategy was to employ serine hydroxamate (SHX) to induce the stringent response (SHX blocks serine tRNA synthetase, leading to an accumulation of uncharged serine tRNAs) (39) and to compare dynamics of replication proteins to those seen after addition of Mitomycin C (MMC), which induces DNA damage supposed to transiently block the progression of forks (11,40,41), or of 6(p-Hydroxyphenylazo)-uracil (HPUra), which reversibly binds to and inhibits DNA polymerase PolC, thereby blocking progression of replication (42).

We additionally generated functional mVenus fluorescent protein fusions to DnaC and to DnaG, and employed fusions to DnaE and to DnaX, previously shown to be able to functionally replace wild type proteins (43). We found that a fusion of mVenus to the C-terminus of DnaC or to that of DnaG, each expressed from the original gene locus under native transcriptional control, as sole source of the protein in the cell, did not negatively affect exponential growth of *B. subtilis* cells (Fig. S1A and C). Additionally, treatment of cells expressing each fusion with MMC, HPUra or SHX did not compromise viability (Fig. S1E). Note that DnaE-mVenus expressing cells showed a weak sensitivity towards SHX treatment (Fig. S1B and S1E). Using high resolution epifluorescence microscopy, we found that very similar to DnaX-CFP, DnaC-mVenus formed fluorescent foci within the cells (Fig. 1A). In 80% of cases, DnaX-CFP and DnaC-mVenus foci colocalized (in the rest of events, only one of the two signals was visible), showing that the helicase fusion is recruited to replication forks, as expected.

For DnaG, two principle scenarios could be envisioned: a) DnaG may come and go to forks by a diffusion/capture mechanism, which would lead to an exchange event every few thousand base pairs, b) DnaG might have binding sites at the forks (analogous to DnaA, the initiator of replication, (44)), which would lead to an enrichment at the forks and a concomitant enhanced replacement efficiency through a local pool of molecules. Fig. 1B shows that in most cells, DnaG-mVenus was dispersed throughout the cells, with some cells showing weak accumulations. Thus, at first sight, there does not seem to be an accumulation of DnaG at the forks.

Next, we treated cells with concentrations of MMC, HPUra or SHX that led to slowed

down growth rate but did not stop growth (Fig. S1A-C). We reasoned that these

concentrations were able to strongly act at the respective targets including replication

forks but did not lead to a large degree of cell death. Fig. 2 shows that DnaC-mVenus foci

appeared to be visually weaker after treatment with HPUra and SHX than after MMC

treatment. As we will move on to single molecule tracking (SMT) below, we will refrain

from quantifying at this place. For DnaG-mVenus, we noted the appearance of visible foci

after addition of MMC and of HPUra, but not in response to SHX (Fig. 2). While there

were qualitative changes in localization patterns, we clearly observed the continued

presence of foci representing replication forks for the three analysed conditions, indicating

that in general, forks persist through the three types of treatments.

Single molecule tracking reveals poor enrichment of DnaG at replication forks,

arguing for a diffusion capture mechanism for recruitment

We next turned towards SMT to quantify changes in protein dynamics at a single molecule

level, using an experimental setup that has been described before (43,45,46). We used

SMTracker software 1.5 (36) to analyse tracks that were generated using u-track

software (33). We employed 20 ms stream acquisition to ensure that even freely diffusing

molecules of DnaC (50.4 kDa plus mVenus) and of DnaG (68.6 kDa plus mVenus) would

be trackable. Fig. 3A and 3B show overlays of all frames of a typical movie (see movies

S1 and S2 for examples) of DnaC, revealing clear foci within cells that contain little

background, indicative of a high number of DnaC molecules being present at the forks.

Conversely, we observed barely detectable discrete accumulations of DnaG, but many

molecules localized throughout the cells. Thus, these two proteins show clearly distinct

localization patterns, indicating that DnaG only transiently and shortly associates with the

replication machinery, contrarily to DnaC. For quantification of molecule trajectories, we

employed Gaussian Mixture Modelling (GMM), which allows to directly compare molecule

dynamics between different proteins, or for a given protein between different growth

conditions: a common value for the diffusion constant "D" is found, which leaves

alterations only to occur in changes of the population size of molecules with a given

diffusion coefficient. GMM also allows to distinguish if the probability density function of

observed step sizes can be explained by a single Gaussian function, and thus by the presence of a single population of molecules having the same value for D, or by two or three different populations, which is tested by an r_2 analysis (47). Fig. S2 shows that for all three proteins, observed distributions could be well explained with the existence of two populations, but not of one. One fraction had a low diffusion constant, corresponding to molecules bound to replication forks, and one with a high value for D, characteristic of freely diffusing molecules, as was described before (43). Fig. 4, upper panel illustrates that DnaC featured the highest proportion of statically bound molecules, with 81% (Fig. S3), and only 19% diffusive molecules, as was expected for the helicase. DnaE showed a relatively balanced proportion of 58% for bound and 42% diffusive molecules (Fig. S3), while DnaG had the lowest population of static molecules (46%), and a larger amount (54%) of unbound ones. These data strongly suggest that DnaG comes and goes to forks, and is only present at very low, likely single molecule level at the replication machinery, suggesting a diffusion capture model for recruitment to the lagging strands, rather than an exchange with molecules pre-bound to other replication-associated proteins.

DnaC, DnaG and DnaE respond differentially to stress conditions at a single molecule level

Addition of MMC to growing cells did not strongly affect the size of DnaE populations, as was described before (43). If anything, DnaE turned to be slightly more stably associated with the forks, becoming more static from about 58 to 63% (Fig. S3). Conversely, both DnaC and DnaG revealed a decrease in the static population with a concomitant increase in unbound molecules (Fig. 4, second row of panels). For DnaC, the size of the static population observed during exponential growth (about 81%) was almost halved to 42%, while for DnaG, the static population decreased from ~ 46 to 29% (Fig. S3). Because chromosome segregation and thus replication continue after addition of MMC, albeit more slowly (48), likely based on frequent restart processes, it follows that DnaC molecules are more frequently exchanged during replication restart than polymerases. Unexpectedly, blocking of PolC by HPUra lead to an increase of fork-bound primase molecules (Fig. 4, third row of panels), while populations of DnaC remained unchanged, and DnaE became slightly more dynamic. The static fraction of DnaG almost doubled from ~ 46 to 82%, while

that of DnaC almost halved from ~ 81 to 42% (Fig. S3). Thus, DnaC and DnaG respond
in an opposite manner to blocking of polymerase activity. Interestingly, under this
condition, DnaE was partially lost from forks, with the static fraction dropping from ~ 58 to
49% (Fig. S3).

Induction of the stringent response induced a third pattern of changes, namely a
strong decrease in fork-bound, static molecules and thus a large increase in freely
diffusive molecules for all three replication proteins. We interpret these findings to mean
that a) stringently blocked forks become highly prone to protein exchange (in agreement
with weaker fluorescence observed using epifluorescence microscopy, Fig. 2), b) in spite
of a block in DnaG recruitment, forks do not completely disintegrate, but are likely ready
to rapidly return to activity when ppGpp levels are lowered.

The different behaviour of DnaG and DnaE at a single molecule level after the
different kinds of replication stress induced suggests that they are independently recruited
to replication forks. Were they generally recruited in an ensemble manner, we would have
expected similar shifts in bound and free populations of molecules during the same stress
conditions.

Nutritional downshift increases turnover of helicase, primase and polymerase at the forks

We wished to obtain more information on the exchange rates of the three replication
proteins under investigation at the replication machinery. We therefore introduced into
SMTracker an analytical tool to quantify the extent of molecules that show transitions
between mobile mode and static behaviour. Confined motion was defined as molecules
staying within a radius of 120 nm (about three times our localization error) for at least 9
steps (Fig. 5A). These states of static motion are indicated in red colour in Fig. 5. Note
that a confined track can be part of a longer track that changes between static and mobile
mode, or vice versa (Fig. 5A). Such "transition" tracks are shown in green in Fig. 5. Not
shown are tracks that are entirely mobile, and do not rest for 9 steps in a row (see Fig.
S4); note that even freely diffusive molecules can stochastically stop for shorter periods
of time. Conversely, confined motion occurs when a protein has restricted movement for
an extended amount of time, which is due to an interaction/binding to a much larger

subcellular structure. To locate these events, a confinement map tool was developed,
using the information given by the dwell time calculation and projecting events into a
standardized cell. A trajectory is considered to present confinement when it has at least
one dwell event (confined track), transitions (mixed behaviour) are changes between
confinement and mobility, freely diffusive track lack any kind of confinement. Fig. 5C
shows that confined motion of DnaC, DnaG and DnaE clusters in the cell centre where
replication forks are present, as expected. In agreement with the finding that DnaG shows
the smallest static fraction, it also showed the smallest extent of confined motion, while
DnaE showed the largest (Fig. 5C). Treatment with MMC visually increased the
percentage of molecules showing transitions, while HPUra did not appear to strongly
affect the ratio between confined tracks and those undergoing transitions (Fig. 5C). To
better quantify these findings, we scored tracks undergoing purely diffusive motion,
transitions or purely confined motion. The latter two fractions were set to 100%, and we
scored the change between confined and transitory molecules. Fig. 5B shows that MMC
treatment slightly increased transitions of DnaC at the forks, this was more pronounced
during blocking of polymerases by HPUra, in agreement with DnaC becoming more
mobile under this condition (Fig. 4 and S3). Most notably, transitions greatly increased
after addition of SHX, not only for DnaC, but also for DnaG and for DnaE (Fig. 5B and C),
revealing that during stringent response, all three proteins revealed highly increased
exchange rates at the forks, even though these were blocked for elongation. Intriguingly,
the pattern of localization of confined and transitory tracks changed markedly under
stringent conditions. All three proteins showed localization away from the cell centre
towards the periphery of cells (Fig. 5). Curious about this finding, we stained cells with
DAPI to study the subcellular localization of the chromosome(s). Fig. S5 shows regular
nucleoids during exponential growth, and condensed nucleoids following addition of
MMC, as has been reported before (49). Nucleoids were also more condensed after
addition of HPUra but were almost completely decondensed during the stringent
response (Fig. S5). Previously, it has been shown that 70S ribosomes occupy nucleoid-
free zones at the cell poles and at the cell periphery (50), which is lost upon inhibition of
transcription, where the nucleoids decondense (51). Our finding that the stringent
response also leads to a strong nucleoid de-condensation suggests that also under this

condition, the separation between RNA polymerase and ribosomes is lost. Our data also suggest that the central positioning of replication forks is lost, which agrees with the appearance of more randomly located forks seen in epifluorescence (Fig. 2). Thus, nutritional downshift not only strongly increases exchange of proteins at the replication forks, but also affects their subcellular localization.

Stress conditions differentially affect dwell times of DnaC, DnaG and DnaE.

Based on the above findings, we would have predicted lower dwell times of replication at the forks during nutritional downshift. We analysed times molecules dwell within a radius of 120 nm and used a two-population fit to analyse decay curves (Fig. S6). Fraction τ_1 shows shorter average dwell times and will mostly consist of mobile molecules that stochastically stay put for a short time. Fraction τ_2 will likely by composed of molecules residing at the replication machinery. Of note, although average track lengths (around 8 steps) were shorter than average dwell times determined, especially for the second fraction (τ_2) that shows long dwell times, there were enough tracks longer than average track length to allow for a correct extrapolation of average residence time. Because we used YFP-bleaching type SMT, our determined numbers are underestimates of actual dwell times *in vivo*. However, for the sake of comparison between different growth conditions, our estimates are useful to observe relative changes in dwell times. Fig. 6 shows that in agreement with our expectations, the dwell time of DnaC decreased during (transient) replication arrest due to MMC or HPUra treatment, and further decreased during the stringent response. For DnaG, we observed an increased residence time after addition of HPUra (Fig. 6) (Table S3), corresponding to the increase in the fraction of static molecules observed (Fig. 4). However, during stringent conditions, average residence time remained similar to values during unperturbed growth, which is unexpected. DnaE showed clearest changes in dwell times, which decreased from MMC to SHX treatment, where residence times were shortest.

Keeping in mind the caveat that our determined dwell times are underestimates, we can still observe that DnaC and DnaE show similar residence times during non-perturbed growth. Thus, as DnaE exchanges with every 1000 to 2000 base pairs synthesised at the

lagging strand, so do DnaC subunits appear to exchange at this DNA strand. We propose that this exchange is based on exchange of subunits from the hexameric helicase, which thus appear to exchange within a time frame of few seconds.

DISCUSSION

All cells need to adjust their decision when to replicate to the nutritional state of the cell, as well as to many other conditions and physiological requirements. Especially for bacteria, it is not only important to regulate initiation of replication, but also extension, because a run-out of e.g. nucleotides might be detrimental if replication fork speed was not down regulated. We have sought to shed light onto the question how *Bacillus subtilis*, a model organism especially for the large group of Gram-positive bacteria, adjusts replication at a single molecule level, in response to amino acid starvation. Interestingly, the architecture of *B. subtilis* replication forks is rather similar to that of eukaryotic cells, and dissimilar to other model bacteria such as *E. coli*, in that two replicative polymerases act at the lagging strand (43,52-54), besides other differences.

Recent studies of bacterial DNA replication have led to a picture of the replisome as an entity that freely exchanges DNA polymerases and displays intermittent coupling between the helicase and polymerase(s) (9). Challenging the textbook model of the polymerase holoenzyme acting as a stable complex coordinating the replisome, these observations suggest a role of the helicase as the central organizing hub. We show here that there is a high degree of plasticity in the interaction between the lagging strand polymerase and the replicative helicase upon association of the primase with the replisome. By combining epifluorescence and *in vivo* single-molecule assays, we demonstrate that replicative helicase DnaC, DNA primase DnaG, and lagging strand polymerase DnaE act differentially in response to transient replication blocks due to DNA damage, to inhibition of the replicative polymerase, or to downshift of serine availability (stringent response).

Addition of HPUra has been shown to block the activity of the replicative DNA polymerase in several bacterial species. Interestingly, we find more static binding of primase DnaG at the forks, and an increase in its dwell times, while DnaE becomes slightly more dynamic, and helicase DnaC even more so. These findings suggest that

blocking of PolC allows for completion of an Okazaki fragment and permits DnaG to
reinitiate binding, but slows down or blocks its turnover. Exchange of DnaC molecules
was found during exponential growth and in an increased manner during all three stress
conditions, when the progression of the forks was blocked or strongly reduced. We
interpret these findings to indicate that the hexameric helicase exchanges its subunits
within few second intervals, replacing them continuously, as is known from exchange of
e.g. rotor parts of the bacterial flagellum (55).

While DnaE became slightly more statically associated with replication forks during
MMC-induced DNA repair and less so during blocking of PolC via addition of HPUra,
DnaG became highly stabilized during block of PolC activity, and less statically positioned
during DNA repair. Thus, responses in single molecule dynamics were quantitatively
different between DnaG and DnaE during conditions of replication stress. Likewise, dwell
times changed in very different manners during stress conditions, indicating that both
proteins are recruited separately and independently from each other to initiate new
Okazaki fragments. Moreover, although DnaC, DnaG and DnaE have been shown to form
a stable complex *in vitro* (53), changes in single molecule dynamics of DnaC in response
to MMC or HPUra treatment were distinct from those of DnaG and DnaE, showing
intriguing plasticity in protein dynamics within this complex, and thus within replication
forks.

Most pronounced changes in single molecule dynamics were found after nutritional
downshift. All three proteins became much less statically associated with forks during the
stringent response, concomitant with a decrease in dwell times for DnaC and for DnaE,
and strongly increased turnover of binding/unbinding events. These findings show that
interaction of DnaG with the stringent response second messenger (p)ppGpp, which is
synthesized via RelA in response to binding of uncharged tRNAs at the ribosome A site,
strongly reduces binding of DnaG to the lagging strand, rather than blocking its activity
and stalling DnaG at the forks. Additionally, our findings show that while chromosomes
decondense during the stringent response, in contrast to condensation during fork block
via MMC or HPUra, replication forks persist, but feature increased protein turnover. This
will lead to slowing down of replication elongation, or even halt replication completely, but
allow for rapid regaining of extension, possibly without the need for restart.

We have recently shown that chromosome segregation and thus also DNA
replication, which occur concomitantly, are relatively robust against DNA damage-
induction via MMC or inhibition of DNA gyrase, continuing to follow a general pattern that
resembles that of directed diffusion (48). Here, we show that replication forks can be seen
to persist during MMC treatment, as well as during inhibition of PolC or of DnaG via
(p)ppGpp binding. Apparently, while featuring rapid exchange of subunits, the replication
machinery appears to be very robust against different kinds of stresses and can be tuned
down in speed after nutritional downshift. Clearly, nature has evolved a highly adaptable
and overall processive/stable machinery for one of the most central aspects of life,
duplication of the genetic information.

ACKNOWLEDGMENTS
This work was supported by the Center for Synthetic Microbiology, SYNMIKRO, at the
Philipps-Universität Marburg, by the research consortium MOSLA, funded by the LOEWE
Program of the state of Hessen, and by Deutsche Forschungsgemeinschaft (DFG)-
funded research consortium TRR 174.

AUTHOR CONTRIBUTIONS
R. H.-T. conceived of the study, performed all experiments, evaluated data and co-wrote
the manuscript. H.S. contributed to experiments shown in Fig. 4 and in Fig. S3 and helped
writing the manuscript. P.L.G. conceived of the study, helped to evaluate data and co-
wrote the manuscript.

COMPETING INTERESTS
The authors declare no competing financial or scientific interests.

REFERENCES

1. Sanders, G.M., Dallmann, H.G. and McHenry, C.S. (2010) Reconstitution of the B. subtilis replisome with 13 proteins including two distinct replicases. *Mol Cell*, **37**, 273-281.
2. McInerney, P., Johnson, A., Katz, F. and O'Donnell, M. (2007) Characterization of a triple DNA polymerase replisome. *Mol Cell*, **27**, 527-538.
3. Berkmen, M.B. and Grossman, A.D. (2006) Spatial and temporal organization of the Bacillus subtilis replication cycle. *Mol Microbiol*, **62**, 57-71.
4. Reyes-Lamothe, R., Sherratt, D.J. and Leake, M.C. (2010) Stoichiometry and architecture of active DNA replication machinery in Escherichia coli. *Science*, **328**, 498-501.
5. Li, Y., Schroeder, J.W., Simmons, L.A. and Biteen, J.S. (2017) Visualizing bacterial DNA replication and repair with molecular resolution. *Curr Opin Microbiol*, **43**, 38-45.
6. Million-Weaver, S., Samadpour, A.N. and Merrikh, H. (2015) Replication Restart after Replication-Transcription Conflicts Requires RecA in Bacillus subtilis. *J Bacteriol*, **197**, 2374-2382.
7. Paschalis, V., Le Chatelier, E., Green, M., Nouri, H., Kepes, F., Soultanas, P. and Janniere, L. (2017) Interactions of the Bacillus subtilis DnaE polymerase with replisomal proteins modulate its activity and fidelity (vol 7, 170146, 2017). *Open Biol*, **7**.
8. Seco, E.M. and Ayora, S. (2017) Bacillus subtilis DNA polymerases, PolC and DnaE, are required for both leading and lagging strand synthesis in SPP1 origin-dependent DNA replication. *Nucleic Acids Res*, **45**, 8302-8313.
9. Monachino, E., Jergic, S., Lewis, J.S., Xu, Z.Q., Lo, A.T.Y., O'Shea, V.L., Berger, J.M., Dixon, N.E. and van Oijen, A.M. (2020) A Primase-Induced Conformational Switch Controls the Stability of the Bacterial Replisome. *Mol Cell*, **79**, 140-154 e147.
10. Beattie, T.R., Kapadia, N., Nicolas, E., Uphoff, S., Wollman, A.J., Leake, M.C. and Reyes-Lamothe, R. (2017) Frequent exchange of the DNA polymerase during bacterial chromosome replication. *Elife*, **6**.
11. Hernandez-Tamayo, R., Oviedo-Bocanegra, L.M., Fritz, G. and Graumann, P.L. (2019) Symmetric activity of DNA polymerases at and recruitment of exonuclease ExoR and of PolA to the Bacillus subtilis replication forks. *Nucleic Acids Res*, **47**, 8521-8536.
12. Bruck, I., Goodman, M.F. and O'Donnell, M. (2003) The essential C family DnaE polymerase is error-prone and efficient at lesion bypass. *Journal of Biological Chemistry*, **278**, 44361-44368.
13. Siddiqui, K., On, K.F. and Diffley, J.F. (2013) Regulating DNA replication in eukarya. *Cold Spring Harb. Perspect. Biol.*, **5**.
14. Corn, J.E. and Berger, J.M. (2006) Regulation of bacterial priming and daughter strand synthesis through helicase-primase interactions. *Nucleic Acids Research*, **34**, 4082-4088.
15. Rannou, O., Le Chatelier, E., Larson, M.A., Nouri, H., Dalmais, B., Laughton, C., Janniere, L. and Soultanas, P. (2013) Functional interplay of DnaE polymerase, DnaG primase and DnaC helicase within a ternary complex, and primase to polymerase hand-off during lagging strand DNA replication in Bacillus subtilis. *Nucleic Acids Research*, **41**, 5303-5320.
16. Johnson, S.K., Bhattacharyya, S. and Griep, M.A. (2000) DnaB helicase stimulates primer synthesis activity on short oligonucleotide templates. *Biochemistry*, **39**, 736-744.

17. Tougu, K. and Marians, K.J. (1996) The extreme C terminus of primase is required for
interaction with DnaB at the replication fork. *Journal of Biological Chemistry*, **271**, 21391-
21397.
18. Oakley, A.J., Loscha, K.V., Schaeffer, P.M., Liepinsh, E., Pintacuda, G., Wilce, M.C., Otting,
G. and Dixon, N.E. (2005) Crystal and solution structures of the helicase-binding domain
of Escherichia coli primase. *J Biol Chem*, **280**, 11495-11504.
19. Bailey, S., Eliason, W.K. and Steitz, T.A. (2007) Structure of hexameric DnaB helicase and
its complex with a domain of DnaG primase. *Science*, **318**, 459-463.
20. Bird, L.E., Pan, H., Soultanas, P. and Wigley, D.B. (2000) Mapping protein-protein
interactions within a stable complex of DNA primase and DnaB helicase from Bacillus
stearothermophilus. *Biochemistry*, **39**, 171-182.
21. Steinchen, W. and Bange, G. (2016) The magic dance of the alarmones (p)ppGpp. *Mol.*
Microbiol., **101**, 531-544.
22. Loveland, A.B., Bah, E., Madireddy, R., Zhang, Y., Brilot, A.F., Grigorieff, N. and Korostelev,
A.A. (2016) Ribosome*RelA structures reveal the mechanism of stringent response
activation. *Elife*, **5**.
23. Schafer, H., Beckert, B., Frese, C.K., Steinchen, W., Nuss, A.M., Beckstette, M., Hantke, I.,
Driller, K., Sudzinova, P., Krasny, L. *et al.* (2020) The alarmones (p)ppGpp are part of the
heat shock response of Bacillus subtilis. *PLoS Genet*, **16**, e1008275.
24. Hauryliuk, V., Atkinson, G.C., Murakami, K.S., Tenson, T. and Gerdes, K. (2015) Recent
functional insights into the role of (p)ppGpp in bacterial physiology. *Nat Rev Microbiol*,
13, 298-309.
25. Autret, S., Levine, A., Vannier, F., Fujita, Y. and Seror, S.J. (1999) The replication
checkpoint control in Bacillus subtilis: identification of a novel RTP-binding sequence
essential for the replication fork arrest after induction of the stringent response. *Mol*
Microbiol, **31**, 1665-1679.
26. Levine, A., Autret, S. and Seror, S.J. (1995) A checkpoint involving RTP, the replication
terminator protein, arrests replication downstream of the origin during the Stringent
Response in Bacillus subtilis. *Mol Microbiol*, **15**, 287-295.
27. Wang, J.D., Sanders, G.M. and Grossman, A.D. (2007) Nutritional control of elongation of
DNA replication by (p)ppGpp. *Cell*, **128**, 865-875.
28. Denapoli, J., Tehranchi, A.K. and Wang, J.D. (2013) Dose-dependent reduction of
replication elongation rate by (p)ppGpp in Escherichia coli and Bacillus subtilis. *Mol*
Microbiol, **88**, 93-104.
29. Lucena, D., Mauri, M., Schmidt, F., Eckhardt, B. and Graumann, P.L. (2018) Microdomain
formation is a general property of bacterial membrane proteins and induces
heterogeneity of diffusion patterns. *BMC Biol.*, **16**, 97.
30. Feucht, A. and Lewis, P.J. (2001) Improved plasmid vectors for the production of multiple
fluorescent protein fusions in Bacillus subtilis. *Gene*, **264**, 289-297.
31. Plank, M., Wadhams, G.H. and Leake, M.C. (2009) Millisecond timescale slimfield imaging
and automated quantification of single fluorescent protein molecules for use in probing
complex biological processes. *Integr Biol (Camb)*, **1**, 602-612.

32. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B. *et al.* (2012) Fiji: an open-source platform for
biological-image analysis. *Nat Methods*, **9**, 676-682.
33. Jaqaman, K., Loerke, D., Mettlen, M., Kuwata, H., Grinstein, S., Schmid, S.L. and Danuser,
G. (2008) Robust single-particle tracking in live-cell time-lapse sequences. *Nature
Methods*, **5**, 695-702.
34. Michalet, X. (2011) Mean square displacement analysis of single-particle trajectories with
localization error: Brownian motion in an isotropic medium (vol 82, 041914, 2010). *Phys
Rev E*, **83**.
35. Vestergaard, C.L., Blainey, P.C. and Flyvbjerg, H. (2014) Optimal estimation of diffusion
coefficients from single-particle trajectories. *Phys Rev E*, **89**.
36. Rosch, T.C., Oviedo-Bocanegra, L.M., Fritz, G. and Graumann, P.L. (2018) SMTracker: a
tool for quantitative analysis, exploration and visualization of single-molecule tracking
data reveals highly dynamic binding of B. subtilis global repressor AbrB throughout the
genome. *Sci Rep*, **8**, 15747.
37. Tinevez, J.Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E.,
Bednarek, S.Y., Shorte, S.L. and Eliceiri, K.W. (2017) TrackMate: An open and extensible
platform for single-particle tracking. *Methods*, **115**, 80-90.
38. Paintdakhi, A., Parry, B., Campos, M., Irnov, I., Elf, J., Surovtsev, I. and Jacobs-Wagner, C.
(2016) Oufti: an integrated software package for high-accuracy, high-throughput
quantitative microscopy analysis. *Molecular Microbiology*, **99**, 767-777.
39. Samarrai, W., Liu, D.X., White, A.M., Studamire, B., Edelstein, J., Srivastava, A., Widom,
R.L. and Rudner, R. (2011) Differential responses of Bacillus subtilis rRNA promoters to
nutritional stress. *J Bacteriol*, **193**, 723-733.
40. Hernandez-Tamayo, R. and Graumann, P.L. (2019) Bacillus subtilis RarA forms damage-
inducible foci that scan the entire cell. *BMC Res Notes*, **12**, 219.
41. Romero, H., Torres, R., Hernandez-Tamayo, R., Carrasco, B., Ayora, S., Graumann, P.L. and
Alonso, J.C. (2019) Bacillus subtilis RarA acts at the interplay between replication and
repair-by-recombination. *DNA Repair (Amst)*, **78**, 27-36.
42. Brown, N.C. (1970) 6-(p-hydroxyphenylazo)-uracil: a selective inhibitor of host DNA
replication in phage-infected *Bacillus subtilis*. *Proc. Natl. Acad. Sci. U S A*, **67**, 1454-1461.
43. Hernandez-Tamayo, R., Oviedo-Bocanegra, L.M., Fritz, G. and Graumann, P.L. (2019)
Symmetric activity of DNA polymerases at and recruitment of exonuclease ExoR and of
PolA to the *Bacillus subtilis* replication forks. *Nucleic Acids Res.*, **47**, 8521-8536.
44. Soufo, C.D., Soufo, H.J., Noirot-Gros, M.F., Steindorf, A., Noirot, P. and Graumann, P.L.
(2008) Cell-cycle-dependent spatial sequestration of the DnaA replication initiator protein
in *Bacillus subtilis*. *Dev. Cell*, **15**, 935-941.
45. Schibany, S., Kleine Borgmann, L.A.K., Rosch, T.C., Knust, T., Ulbrich, M.H. and Graumann,
P.L. (2018) Single molecule tracking reveals that the bacterial SMC complex moves slowly
relative to the diffusion of the chromosome. *Nucleic Acids Res.*, **46**, 7805-7819.
46. Romero, H., Rosch, T.C., Hernandez-Tamayo, R., Lucena, D., Ayora, S., Alonso, J.C. and
Graumann, P.L. (2019) Single molecule tracking reveals functions for RarA at replication
forks but also independently from replication during DNA repair in *Bacillus subtilis*. *Sci.
Rep.*, **9**, 1997.

47. Rosch, T.C., Oviedo-Bocanegra, L.M., Fritz, G. and Graumann, P.L. (2018) SMTracker: a
tool for quantitative analysis, exploration and visualization of single-molecule tracking
data reveals highly dynamic binding of *B. subtilis* global repressor AbrB throughout the
genome. *Sci. Rep.*, **8**, 15747.
48. El Najjar, N., Geisel, D., Schmidt, F., Dersch, S., Mayer, B., Hartmann, R., Eckhardt, B., Lenz,
P. and Graumann, P.L. (2020) Chromosome Segregation in *Bacillus subtilis* Follows an
Overall Pattern of Linear Movement and Is Highly Robust against Cell Cycle Perturbations.
mSphere, **5**.
49. Sanchez, H., Carrasco, B., Cozar, M.C. and Alonso, J.C. (2007) *Bacillus subtilis* RecG branch
migration translocase is required for DNA repair and chromosomal segregation. *Mol.*
Microbiol., **65**, 920-935.
50. Lewis, P.J., Thaker, S.D. and Errington, J. (2000) Compartmentalization of transcription
and translation in *Bacillus subtilis*. *EMBO J.*, **19**, 710-718.
51. Mascarenhas, J., Weber, M.H. and Graumann, P.L. (2001) Specific polar localization of
ribosomes in Bacillus subtilis depends on active transcription. *EMBO Rep*, **2**, 685-689.
52. Seco, E.M. and Ayora, S. (2017) *Bacillus subtilis* DNA polymerases, PolC and DnaE, are
required for both leading and lagging strand synthesis in SPP1 origin-dependent DNA
replication. *Nucleic Acids Res.*, **45**, 8302-8313.
53. Rannou, O., Le Chatelier, E., Larson, M.A., Nouri, H., Dalmais, B., Laughton, C., Janniere, L.
and Soultanas, P. (2013) Functional interplay of DnaE polymerase, DnaG primase and
DnaC helicase within a ternary complex, and primase to polymerase hand-off during
lagging strand DNA replication in *Bacillus subtilis*. *Nucleic Acids Res.*, **41**, 5303-5320.
54. Sanders, G.M., Dallmann, H.G. and McHenry, C.S. (2010) Reconstitution of the *B. subtilis*
replisome with 13 proteins including two distinct replicases. *Mol. Cell*, **37**, 273-281.
55. Leake, M.C., Chandler, J.H., Wadhams, G.H., Bai, F., Berry, R.M. and Armitage, J.P. (2006)
Stoichiometry and turnover in single, functioning membrane protein complexes. *Nature*,
443, 355-358.
660
661
662

663 **TABLE 1.** Diffusion constants and percentages of static and mobile molecule fractions.

Strain	# cells	# tracks	D_a	D_{1b}	F_{1c}	D_{2d}	F_{2e}
Non-treated							
DnaE-mV	136	6058	0.032 ± 0.004	0.053 ± 0.005	58±2.2	0.75 ± 0.014	42±1.6
DnaC-mV	122	5303	0.039 ± 0.005	0.093 ± 0.004	81±1.7	0.82 ± 0.025	19±1.3
DnaG-mV	150	5893	0.153 ± 0.007	0.100 ± 0.005	46±1.9	0.88 ± 0.020	54±1.6
MMC-treated							
DnaE-mV	136	5275	0.027 ± 0.005	0.053 ± 0.005	63±1.7	0.75 ± 0.014	37±2.2
DnaC-mV	134	5139	0.107 ± 0.005	0.093 ± 0.004	42±1.6	0.82 ± 0.025	58±1.8
DnaG-mV	135	7097	0.174 ± 0.007	0.100 ± 0.005	29±1.4	0.88 ± 0.020	71±1.5
HPUra-treated							
DnaE-mV	121	4775	0.093 ± 0.007	0.053 ± 0.005	49±2.3	0.75 ± 0.014	51±1.2
DnaC-mV	134	5135	0.078 ± 0.005	0.093 ± 0.004	42±1.5	0.82 ± 0.025	58±1.5
DnaG-mV	110	5526	0.109 ± 0.007	0.100 ± 0.005	81±1.6	0.88 ± 0.020	19±1.1
SHX-treated							
DnaE-mV	158	5067	0.220 ± 0.007	0.053 ± 0.005	12±2.4	0.75 ± 0.014	88±1.9
DnaC-mV	171	6690	0.187 ± 0.003	0.093 ± 0.004	13±1.5	0.82 ± 0.025	87±1.4
DnaG-mV	157	6937	0.230 ± 0.005	0.100 ± 0.005	12±1.6	0.88 ± 0.020	88±1.2

665
666 D_a, average diffusion constant of all molecules ($\mu m^2 \cdot s^{-1}$).
667 D_{1b}, diffusion constant of static fraction ($\mu m^2 \cdot s^{-1}$).
668 F_{1c}, percentage of static molecules (%).
669 D_{2d}, diffusion constant of mobile fraction ($\mu m^2 \cdot s^{-1}$).
670 F_{2e}, percentage of mobile molecules (%).
671
672

LEGENDS TO THE FIGURES

Fig. 1 Colocalization of DnaC-mVenus, or of DnaG-mVenus with DnaX-CFP in *B.*

***subtilis* cells**. Epifluorescence microscopy of cells expressing A) both DnaC-mV and

DnaX-CFP, or B) DnaG-mV and DnaX-CFP, during exponential growth as sole source of

the proteins. Scale bars 5 µm and zoom panel 2 µm.

Fig. 2 Localization of DnaC-mVenus, DnaG-mVenus and DnaE-mVenus during

stress conditions. Epifluorescence microscopy of *B. subtilis* cells expressing DnaC-mV,

DnaG-mV or DnaE-mV as sole source of the proteins treated with different drugs as

indicated above the panel rows. Scale bars 2 µm.

Fig. 3 Sum of single molecule tracking movies of exponentially growing cells

expressing DnaC-mVenus or DnaG-mVenus. Insets in panels A and C show bright field

images, outlines of cells are indicated by white ovals. Panels A and C show heat maps of

localization, panels B and D fluorescence images. White bars 2 µm.

Fig. 4 Diffusion patterns of DnaE-mVenus, DnaC-mVenus and DnaG-mVenus.

Gaussian mixture model (GMM) analyses of frame-to-frame displacements in x- and y-

directions. Black lines represent the sum of the two Gaussian distributions. Dotted red

and blue lines represent the single Gaussian distributions corresponding to the static and

mobile fractions.

Fig. 5 Analyses of tracks for the percentage of transitions under stress conditions.

A) cartoon showing the mode of analyses of confined movement (red) versus transition

events (green). B) Bar plot showing changes between confined and transitory motion for

the three proteins under different stress conditions. C) Maps showing intracellular location

of confined motion (as defined by not leaving a radius of 120 nm for at least 9 steps) and

transition events for DnaC-mVenus, DnaG-mVenus and DnaE-mVenus, projected into a

standardized cell of 3 x 1 µm size.

Fig. 6 Dwell times. Cumulative distribution of residence times of DnaC-mVenus, DnaG-
mVenus and DnaE-mVenus strains, before and after treatment with MMC, HPUra or SHX.
Dwell times are estimated using an exponential decay model. Histograms show events
of resting fitted by a two-component exponential function. Bars represent long dwell times
of DNA-bound molecules. Dark blue bars untreated cells, brown MMC-treated cells, grey
HPUra-treated cells and blue light SHX-treated cells. As usual, ** and *** stands for p-
values lower than 0.01 and 0.05 respectively, while *n.s.* stands for statistically not
significant.
712

714
715   Fig 1.
716
717

718
719   Fig 2.
720
721

722
723   Fig 3.
724

726   Fig 4.

728
729   Fig 5.
730

731
732   Fig 6.

733
734
735
736