Dynamic Configuration of Coactive Micropatterns in the Default Mode Network during Wakefulness and Sleep

Yan Cui[1, #], Min Li[1, #], Bharat Biswal[1, 2, *], Wei Jing[1, 3], Changsong Zhou[4], Huixiao Liu[1], Daqing Guo[1, 6, *], Yang Xia[1], Dezhong Yao[1, 5, 6, *]

[1]The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for NeuroInformation, University of Electronic Science and Technology of China, Chengdu 611731, China

[2]Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ 07102, USA

[3]Department of Physiology, School of Basic Medicine and Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

[4]Department of Physics, Centre for Nonlinear Studies and Beijing-Hong Kong-Singapore Joint Centre for Nonlinear and Complex Systems (Hong Kong), Institute of Computational and Theoretical Studies, Hong Kong Baptist University, Kowloon Tong, Hong Kong

[5]School of Electrical Engineering, Zhengzhou University, Zhengzhou 450001, China

[6]Sichuan Institute for Brain Science and Brain-Inspired Intelligence, Chengdu 611731, China

[#]These authors contributed equally to the work.

[*]Corresponding authors: dqguo@uestc.edu.cn, dyao@uestc.edu.cn and bbiswal@gmail.com

Keywords: default mode network; coactive micropattern; wakefulness and sleep; up-down states; dynamic configuration

Abstract

The default mode network (DMN) is a prominent intrinsic network that is observable in many mammalian brains. However, few studies have investigated the temporal dynamics of this network based on direct physiological recordings. Herein, we addressed this issue by characterizing the dynamics of local field potentials (LFPs) from the rat DMN during wakefulness and sleep with an exploratory analysis. We constructed a novel coactive micropattern (CAMP) algorithm to evaluate the configurations of rat DMN dynamics and further revealed the relationship between DMN dynamics with different wakefulness and alertness levels. From the gamma activity (40-80 Hz) in the DMN across wakefulness and sleep, three spatially stable CAMPs were detected: a common low-activity level micropattern (cDMN), an anterior high-activity level micropattern (aDMN) and a posterior high-activity level micropattern (pDMN). A dynamic balance across CAMPs emerged during wakefulness and was disrupted in sleep stages. In the slow-wave sleep (SWS) stage, cDMN became the primary activity pattern, whereas aDMN and pDMN were the major activity patterns in the rapid eye movement sleep (REM) stage. Additionally, further investigation revealed phasic relationships between CAMPs and the up-down states of the slow DMN activity in the SWS stage. Our study revealed that the dynamic configurations of CAMPs were highly associated with different stages of wakefulness and provided a potential three-state model to describe the DMN dynamics for wakefulness and alertness.

Impact Statement

In the current study, a novel coactive micropattern method (CAMP) was developed to elucidate fast DMN dynamics during wakefulness and sleep. Our findings demonstrated that the dynamic configurations of DMN activity are specific to different wakefulness stages and provided a three-state DMN CAMP model to depict wakefulness levels, thus revealing a potentially new neurophysiological representation of alertness levels. This work could elucidate the DMN dynamics underlying different stages of wakefulness and have important implications for the theoretical understanding of the neural mechanism of wakefulness and alertness.

Introduction

Multimodal imaging studies of the human brain have discovered several intrinsic

connectivity networks (ICNs) coexist during the resting state (Beckmann, DeLuca,

Devlin, & Smith, 2005; Q. Liu, Farahibozorg, Porcaro, Wenderoth, & Mantini, 2017).

Dynamic switching within these ICNs have demonstrated a hierarchical structure over

time for brain activity at rest and have been significantly associated with cognitive

traits (M. D. Fox et al., 2016; Vidaurre, Smith, & Woolrich, 2017). This suggests that

brain activity is appropriately understood in terms of the dynamic configuration

among ICNs. These studies have mainly considered each ICN as a whole during brain

dynamics while ignoring the intrinsic dynamics of individual ICNs. Indeed, individual

ICN also exhibits strong fluctuations in brain activity, and different ICNs are believed

to dominate distinct cognitive functions (Rosazza & Minati, 2011). For a specific

brain function, further tracking the dynamic configuration of fluctuations in brain

activity at the single-ICN level might be critical for revealing the underlying

physiological mechanism.

The default mode network (DMN) is one of the important ICNs, and is typically

believed to be related to off-task internal mentations with high activity in the resting

state (Gusnard, Akbudak, Shulman, & Raichle, 2001; M E Raichle et al., 2001).

Recent studies have reported that the DMN is also engaged and displays positive

contributions during several higher cognition task performances, such as the Tower of

London task (D. Vatansever, Menon, Manktelow, Sahakian, & Stamatakis, 2015;

Deniz Vatansever, Manktelow, Sahakian, Menon, & Stamatakis, 2018). Changes in

wakefulness could also lead to alterations of DMN activity and connectivity in

humans and rodents (K. C. R. Fox, Foster, Kucyi, Daitch, & Parvizi, 2018; Lu et al.,

2012; Marcus E Raichle, 2015). DMN connectivity between the frontal and posterior

areas in the human brain was reduced during the slow wave sleep (SWS) stage with a

low level of wakefulness (Sämann et al., 2011). However, at sleep onset and

throughout the rapid eye movement sleep (REM) stage, regions in the human DMN

were shown to be persistently coupled (Horovitz et al., 2008; Larson-Prior et al.,

2009). These findings illustrated that DMN activity was functionally reorganized

during sleep and might further reflect levels of wakefulness and alertness.

Additionally, fast and ever-changing dynamics of DMN activity have also been

observed in various wakefulness levels in humans, implying that the temporal aspects

of spontaneous DMN activity might be associated with alertness levels (Kapogiannis,

Reiter, Willette, & Mattson, 2014; Panda et al., 2016). Thus, research investigating the

dynamic configuration of DMN in different stages of wakefulness using direct

physiological recordings is important.

The DMN could also be observable in rat brains, indicating its conservation in the

mammalian brain during evolution (Huang et al., 2016; Lu et al., 2012). Though there

exists several difference between the brain regions in rat DMN and human DMN, the

anatomical topologies of them are visually similar (Hsu et al., 2016; Marcus E

Raichle, 2015). In addition, neural activity of DMN regions also activated in awake

rats, and suppressing the activity in the core DMN regions could modulate rats'

normal behavior (Tu, Ma, Ma, Dopfel, & Zhang, 2020; Upadhyay et al., 2011).

Furthermore, the information flow within anterior and posterior DMN subsystems

exhibited various alterations in different sleep stages and mental disorders in rats (Cui

et al., 2018; Jing et al., 2017). The above findings imply that the DMN might carry a

core function that transcends across species. Therefore, employing animal models to

direct record physiological DMN signals is an effective way to investigate the

temporal dynamics of DMN activity.

Several neurophysiological studies have reported that during the deep sleep stage,

the neurons in various brain regions exhibited highly synchronized firing rates,

occurring at approximately 0.5-2 Hz (Crunelli & Hughes, 2010; Gretenkord et al.,

2020; Lőrincz et al., 2015). This specific activity pattern was identified as the

up-down state, and has been considered a biomarker of low-level wakefulness in deep

sleep (Jercog et al., 2017; Perez-Zabalza et al., 2020). Moreover, this up-down state

has been demonstrated for both neuron membrane potentials and local field potentials

(LFPs) (Holcman & Tsodyks, 2006) and characterizes the dynamics of slow

oscillations during deep sleep (Ji & Wilson, 2007; Lőrincz et al., 2015). However, the

existence of a physiological relationship between the up-down state and DMN

dynamics is a topic of active interest, that deserves further exploration.

In the present study, we developed and applied a new dynamic activity pattern

method to address these challenges. The new method, named the coactive

micropattern analysis (CAMP), decomposed the fast dynamic activity into several

intrinsic CAMPs and defined the brain dynamics through the constitutions and

transitions among these CAMPs. We employed the CAMP analysis to elucidate the

dynamic configurations of LFPs from rat DMN in different stages during wakefulness

and sleep. Our results illustrate a reorganized dynamic configurations of CAMPs for

fast DMN activity in different stages of wakefulness, implying that the dynamic

configurations of DMN micropatterns might provide underlying neural correlates for

the wakefulness levels observed during wakefulness and sleep.

Methods and Materials

Detailed descriptions of the experimental procedures and data acquisition are

described in the ***Supplementary Materials.*** Twenty-nine male Sprague-Dawley rats

were used in our study. The DMN signals were acquired by chronically implanting

fifteen electrodes into the brains of rats under deep anesthesia (Fig. 1 and Table 1).

The rat DMN contained the following bilateral structures: the orbital frontal cortex

(OFC), the rostral dorsal prelimbic cortex (PrL), the cingulate cortex (CG), the

retrosplenial cortex (RSC), the dorsal hippocampus (HIP), the temporal lobe cortex

(TE), the medial secondary visual cortex (V2) and the posterior parietal cortex (PPC).

According to their anatomical coordinates (Lu et al., 2012), the PrL, OFC and CG

regions were considered to be in the anterior subsystem of the DMN, whereas the

RSC, HIP, PPC, TE and V2 regions were in the posterior subsystem of the DMN. In

addition, we also implanted two electromyographic (EMG) electrodes bilaterally in

the dorsal neck muscles. After DMN electrode implantation surgery, all rats recovered

for approximately 2 weeks. During the recording session, the rats were placed in a

noise-attenuated chamber and were allowed to move freely without anesthesia. All the

signals for the LFP, the EMG and the videos signals were simultaneously recorded

and continuously monitored for 72 h.

The dataset used in the current study was selected from the last 24 h of the total

recording and was separated into three stages, including resting (AWAKE), SWS and

REM sleep stages. The AWAKE stage of rats was defined when the rats were standing

or sitting quietly with low-amplitude and mixed-frequency LFP activity and relatively

low and stable EMG activity. The SWS stage was the sleep duration when the rats

were sleeping with high-amplitude and low-frequency LFP activity and low-level

EMG activity. The REM sleep stage was the duration when the rats were sleeping

with sawtooth-pattern LFP activity and flat EMG activity. For each rat, 30 segments

in different stages were chosen, and each segment lasted 10 s (a total of 300 s of

LFPs). All experimental animal procedures were approved by the Institutional Animal

Care and Use Committee of the University of Electronic Science and Technology of

China.

Moreover, we proposed a novel CAMP method to track fast DMN dynamics

during wakefulness and sleep. Briefly, this method utilizes a point process approach

that combines the advantages of both microstate analysis and coactive pattern analysis

(X. Liu & Duyn, 2013; Michel & Koenig, 2018) and extracts CAMPs based on the

extreme values of envelope signals at a high temporal resolution. Five steps were

included in the CAMP algorithm. First, the original data were bandpass filtered into

the gamma frequency band (40-80 Hz) and then Hilbert transformation was used to

obtain the envelope signals (Fig. 2b). Second, the envelope signals were normalized

and downsampled to improve the signal-to-noise ratio (SNR) for further analysis (Fig.

2c). Third, the active points for each envelope signal channel were then defined as the

extreme points of the envelope signals, including local maximum and minimum

values. Afterwards, the coactive patterns (CAPs) of the brain, which were defined as

brain maps in which more than one brain region displayed active points at the same

time point, were introduced for all stages (Fig. 2d). Fourth, a k-means clustering

algorithm was applied to all the CAPs to decompose the CAMPs and the CAMP

index (Fig. 2e). Finally, the criterion based on squared Euclidean distance was applied

to update the CAMP and CAMP index (Fig. 2f). The last step was employed to

precisely determine the final spatial structures of all CAMPs and the CAMP index.

Using the CAMP method, we decomposed three stable CAMPs from gamma activity

during DMN dynamics to reveal the fast changes in DMN activity in different stages

of wakefulness. A detailed description of the CAMP analysis is provided in the

Supplementary Methods.

The CAMP method was separately applied to the DMN activity of each segment

in different stages for each rat and to the concatenated DMN activity of all rats and all

stages during wakefulness and sleep. The CAMPs extracted from each rat in different

stages were further employed to test their spatial stability across rats and wakefulness

levels using Pearson correlation method. The results were derived from the CAMPs

extracted from the concatenated DMN activities from all rats during wakefulness and

sleep unless otherwise described.

Results

Three CAMPs of gamma activity in the DMN during wakefulness and sleep

The CAMP analysis procedure developed in the present study is schematically

illustrated in Fig. 2 and described in more detail in the ***Supplementary Methods***. The

concatenated gamma activities in the DMNs of all rats and all stages during

wakefulness and sleep were decomposed into three distinct CAMPs, including a

common low-activity level micropattern (cDMN), an anterior high-activity level

micropattern (aDMN) and a posterior high-activity level micropattern (pDMN). In the

cDMN, all DMN regions exhibited similar and low levels of activity (mean

normalized activity: 0.2577 ± 0.0041, Fig. 2g), indicating a potential cooperation of

them in this type of CAMP. However, two different levels of activity were observed in

both the aDMN and pDMN with the aDMN exhibiting relatively higher levels of

activity in the anterior DMN regions (mean normalized activity: 0.3868 ± 0.0018) and

lower activity in the posterior DMN structures (mean normalized activity: $0.3050 \pm$

0.0060, Fig. 2h). In the pDMN, the posterior DMN structures displayed higher levels

of activity (mean normalized activity: 0.3793 ± 0.0145), whereas the anterior DMN

regions exhibited relatively lower levels of activity (mean normalized activity: 0.3073

± 0.0021, Fig. 2i). Accordingly, both the aDMN and pDMN were considered

high-activity micropatterns in DMN dynamics.

We separately decomposed the CAMPs for each rat in every wakefulness stage

individually and tested their reliability across rats and stages. All three CAMPs

exhibited high stability with large correlation coefficients among different rats during

wakefulness and sleep (mean correlation coefficients: r = 0.7451, r = 0.7535, r =

0.6684 for the AWAKE, SWS and REM sleep stages, respectively; Table 2). In

addition, the spatial structures of these CAMPs were also similar among the AWAKE,

SWS and REM sleep stages (mean correlation coefficients: r = 0.6229, r = 0.7882, r =

0.8600 for the cDMN, aDMN and pDMN, respectively; Table 3). These findings

demonstrated the high reliability and robustness of these CAMPs.

Temporal features and activity levels of each CAMP during wakefulness and

sleep

We computed several temporal measurements, including the total occurrence

(occurrence probability), total duration (duration probability) and mean duration, to

characterize the features and dynamics of these CAMPs during wakefulness and sleep.

All these features represented the temporal properties of these CAMPs in different

stages. Based on the comparisons, all features of cDMN displayed the largest values

in the SWS stage and the smallest values in the REM sleep stage, and the two

high-activity micropatterns (aDMN and pDMN) exhibited the largest values for all

features in the REM sleep stage and the smallest values in the SWS stage (Fig. 3a-3c).

These opposite alterations in features between low- and high-activity micropatterns

suggests that these two types of CAMPs might play different physiological roles for

wakefulness. Besides, all the features in three stages were remarkably different among

CAMPs, improving our knowledge of the changes in wakefulness levels during

wakefulness and sleep.

However, all of these CAMPs displayed different activities in DMN regions

during wakefulness and sleep. In particular, all DMN regions exhibited reduced

activity during SWS stages in all CAMPs (Fig. 3e-3g). Moreover, the posterior DMN

structures exhibited significantly reduced activity in the aDMN, while the anterior

DMN regions exhibited significantly reduced activity in the pDMN. All of these

regions showed relatively lower activity in the AWAKE stage, indicating a

preservation of the major activity in these two micropatterns during deep sleep (Fig.

3f-3g, red stars). However, all CAMPs displayed increased activity in most DMN

regions during the REM sleep stage. The activities in the HIP, OFC and RSC regions

were significantly increased during the REM sleep stage in all CAMPs, implying the

importance of these DMN regions for REM sleep (Fig. 3h-3j, red stars). In addition,

the mean activity level of each CAMP exhibited similar variation trends across

different stages of wakefulness. The lowest mean activity of CAMPs was observed in

the SWS stage, whereas the highest mean activity was observed in the REM sleep

stage (Fig. 3d).

The features and transitions of CAMPs during wakefulness and sleep

The configurations of these CAMPs involved in DMN dynamics in different

stages were also distinct (Fig. 4a-4c). All CAMPs presented similar features in the

AWAKE stage (occurrence probabilities: 32.12%, 34.12% and 33.75%; duration

probabilities: 31.68%, 34.15% and 34.17%; and mean duration: 29.79 ms, 24.41 ms

and 24.37 ms for the cDMN, aDMN and pDMN, respectively). No significant

differences of the features among three CAMPs were observed, indicating that their

roles were equivalent and that a dynamic balance in DMN activity might exist among

CAMPs at wakeful rest. However, the cDMN became the dominant activity pattern of

DMN dynamics in the SWS stage as it had the largest occurrence probabilities

(62.66%, 17.56% and 19.78% for the cDMN, aDMN and pDMN, respectively),

duration probabilities (61.47%, 18.02% and 20.51% for the cDMN, aDMN and

pDMN, respectively) and mean duration (55.58 ms, 21.89 ms and 22.78 ms for the

cDMN, aDMN and pDMN, respectively) among all CAMPs. The predominant

constituent of the low-activity micropattern suggests that all the DMN regions might

have been in a stage of low activity and that DMN activity preferred a silent pattern

during deep sleep. However, the two high-activity micropatterns were the main

CAMPs during the REM sleep stage. All the features of aDMN and pDMN were

significantly larger than those of cDMN (occurrence probabilities: 19.42%, 42.05%

and 38.53%; duration probabilities: 19.33%, 41.84% and 38.83%; and mean duration:

21.88 ms, 26.92 ms and 26.01 ms for the cDMN, aDMN and pDMN, respectively).

The greater percentage of high-activity micropatterns during REM sleep suggests a

reactivation of DMN activity in this stage. In addition, comparisons of features within

two high-activity micropatterns demonstrated that the aDMN displayed significantly

larger values for the three features, implying that it played a more important role in

REM sleep.

Furthermore, the temporal concatenations of these CAMPs (i.e., the CAMP

indices) in different stages also exhibited specific changes. We first performed a

randomization test to examine the transition structures of these CAMP indices in

different stages. The transitions among CAMPs occurred randomly in the AWAKE

stage (p = 0.8157), indicating that the transition probabilities (TPs) of pairs of CAMPs

in the resting stage were proportional to their occurrences. However, these transitions

did not occur randomly in the SWS (p<0.0001) or REM sleep stages (p<0.0001),

suggesting the stabilization of the CAMP index structures during the sleep cycle and

further implying the existence of several preferred transitions among CAMPs in both

SWS and REM sleep stages.

Next, we compared the TPs for pairs of CAMPs between the two sleep stages and

the AWAKE stage. We revealed similar TPs in the AWAKE stage (no significant

differences among all TPs, Fig. 4d), suggesting the presence of balanced transitions

among all CAMPs at rest. However, the TPs within the two high-activity

micropatterns showed significant reductions in the SWS stage, whereas those between

the high-activity micropatterns and the low-activity micropattern increased

significantly (Fig. 4e). These changes in TPs emphasized the functional role of

inhibitory activity in DMN regions in deep sleep. On the other hand, TPs in the REM

sleep stage displayed different alterations, including significantly increased TPs

within the high-activity micropatterns and remarkable decrease in TPs between the

high-activity micropatterns and the low-activity micropattern (Fig. 4f). The increased

transitions within two high-activity micropatterns revealed activation of DMN regions

during REM sleep. Based on these findings, the CAMP indices and the functional

roles of these CAMPs were specific for different stages. The alterations in DMN

activity during wakefulness and sleep might be attributed to the specific temporal

combinations of the CAMPs constituting the activity in different stages rather than the

spatial structures of CAMPs themselves, which were rather stable across different

stages.

Strong phasic relationships between CAMPs and up-down states in the SWS

stage

Up-down states are considered the predominant pattern of slow oscillations (0.5-2

Hz) during the SWS stage. Estimations of the phase distributions of each CAMP in

the anterior and posterior DMN slow activity regions with the Hilbert transformation

demonstrated that these CAMPs displayed strong phasic relationships with the

up-down states in the SWS stage. The cDMN preferred the down state of anterior

DMN activity (Fig. 5a, significant directionality: 1.97 π, red line) and the up state of

posterior DMN activity (Fig. 5d, significant directionality: 1.16 π, red line).

Additionally, both the aDMN and pDMN were phase locked to the up state of anterior

DMN activity (Fig. 5b, significant directionality: 1.21 π for aDMN; Fig. 5c,

significant directionality: 1.18 π for pDMN) and the down state of posterior DMN

activity (Fig. 5e, significant directionality: 0.23 π for aDMN; Fig. 5f, significant

directionality: 0.18 π for pDMN). These similar phasic relationships implied that two

high-activity micropatterns might belong to the same activity pattern of slow

oscillations during deep sleep. Thus, our proposed CAMPs may reflect the up-down

states of DMN slow activity in the SWS stage, and a close physiological association

existed between the up-down states with DMN dynamics.

Discussion

In the present study, we proposed a coactive micropattern (CAMP) algorithm to

reveal the dynamics of DMN based on the direct physiological recordings in rat

during wakefulness and sleep. Our results indicated that the fast dynamics of DMN

gamma activity could be decomposed into three different CAMPs. These CAMPs

exhibited stable spatial structures across wakefulness and sleep, while their dynamic

configurations were specific to different stages. In addition, all these CAMPs were

strongly phase locked to the up-down states of slow DMN activity in the SWS stage,

suggesting the temporal sequence of the neural relationship between up-down states

and DMN dynamics. Our findings described the distinct dynamic configurations of

DMN activity during wakefulness and sleep. The proposed a three-state model may

reveal a neural mechanism by which DMN dynamics mediated wakefulness and

alertness.

Physiological significance of the three CAMPs

Previous studies have reported a strong correlation between electrophysiological

gamma activity and blood oxygen level-dependent (BOLD) signals (N. K. Logothetis,

Pauls, Augath, Trinath, & Oeltermann, 2001; Nikos K. Logothetis, 2002; Magri,

Schridde, Murayama, Panzeri, & Logothetis, 2012; Scheering, Koopmans, Van

Mourik, Jensen, & Norris, 2016). In addition, DMN regions have also shown

deactivation at gamma frequency during the performance of external tasks in several

human electroencephalography (EEG) studies (Karim Jerbi et al., 2010; Ossandon et

al., 2011), indicating the importance of gamma oscillation in DMN activity. Hence,

we specifically focused on the fast dynamics of DMN gamma activity in the current

study. The gamma activity in the rat DMN was decomposed into three stable CAMPs

during wakefulness and sleep. We also showed the CAMPs decomposed from DMN

alpha (8-13Hz) and beta (13-30Hz) activity, which exhibited similar structures with

those found in gamma band (Supplementary Fig. 5). The differences across these

CAMPs further provided direct electrophysiological evidence that the DMN regions

might not be simultaneously activated. Besides, different CAMPs had distinct mean

durations. These phenomena revealed the differences in the activation times of

anterior and posterior DMN structures in the fast dynamics and further illustrated the

diversity in the latencies for both the excitation and inhibition of DMN regions (Brett

L. Foster, Mohammad Dastjerdi, 2012; Foster, Rangarajan, Shirer, & Parvizi, 2015).

Indeed, both human and animal studies found that the DMN structure could be

separated into two subnetworks, i.e., a parietal subnetwork and a prefrontal

subnetwork (Cui et al., 2018; Hagmann et al., 2008; Lu et al., 2012; Wu et al., 2017).

In the present study, we not only reinforced this finding from the aspect of fast DMN

dynamics but also provided a possible dynamic substrate for this separation of the

DMN structure. As a key component of the DMN, the orbital frontal cortex (OFC) has

historically been posited to integrate interoceptive and exteroceptive information from

multisensory stimuli to process information about the internal and external bodily

milieu (Ongur & Price, 2000). Accordingly, we hypothesized that the high-activity

micropattern aDMN might play an important role in making inferences and guiding

actions in a timely and environmentally relevant manner.

Furthermore, the retrosplenial cortex (RSC), another key area in the DMN, has

extensive connections with the hippocampal formation. The projections between the

RSC and hippocampal formation provide an important pathway that regulates learning,

memory and emotional behavior (Wyss & Vangroen, 1992). Furthermore, the

hippocampal formation is a limbic structure that forms direct or indirect connections

to other DMN regions. Therefore, the high-activity micropattern pDMN detected in

the present study might be associated with memory and emotional behavior.

Additionally, both the aDMN and pDMN were strongly phase locked to the up state

of anterior DMN activity and the down state of posterior DMN activity during the

SWS stage, indicating that they may reflect similar performances for the up-down

states of slow oscillations during DMN dynamics. Moreover, these two high-activity

micropatterns together accounted for more than 70% of the time in the resting state,

which helps explain why the brain requires high basal cerebral blood flow and

metabolism for spontaneous activity (Marcus E Raichle & Mintun, 2006). It should be

noted that the DMN structure could also be split into dorsal and ventral branches

according to the dorsal and medial temporal regions of RSC and hippocampus in

human brain (Chen, Glover, Greicius, & Chang, 2017; Shirer, Ryali, Rykhlevskaia,

Menon, & Greicius, 2012). However, the rat DMN is commonly divided into the

anterior and posterior subsystems for the anatomical difference with human DMN

(Marcus E Raichle, 2015).

We also observed a low-activity micropattern (i.e., cDMN) in DMN dynamics

that was widely distributed in all wakefulness stages. In the cDMN, all DMN regions

displayed lower activity, indicating that the cDMN could represent the silent state for

DMN activity in which all the DMN regions prefer relaxations and are prepared for

the next excitation. Moreover, the cDMN was the only coactive micropattern in which

all DMN regions operated in the same manner in DMN dynamics. Thus, the

appearance of cDMN suggested a working mode for DMN with low energy, but this

concept requires further study.

The balance of dynamic DMN configurations supports wakefulness and alertness

during wakefulness

Based on accumulating evidence, DMN activity is tightly correlated with

wakefulness levels in health and disease (Buckner, Andrews-Hanna, & Schacter, 2008;

Kapogiannis et al., 2014; Panda et al., 2016). In the AWAKE stage, all the CAMPs

exhibited similar features, and the dynamic transitions among them were not

significantly different. These similarities illustrated a balanced dynamic configuration

among these CAMPs during fast gamma activity in the DMN at rest. The DMN is a

key network involved in integrating high-order information from multiple sensory

modalities based on numerous projections from variable somatic cortex and core

limbic structures (HIP and amygdala) to the DMN regions (Heidbreder &

Groenewegen, 2003; Reep, Chandler, King, & Corwin, 1994). These projections

might provide the anatomical substrate for the correlation of DMN activity with

alertness levels, which are largely believed to be determined by global levels of

arousal regulated by the brainstem via the reticular activating system (RAS)

(Delano-Wood et al., 2015). Accordingly, the identified balance of DMN dynamics

might be a competitive product between the integration and differentiation of DMN

activity in maintaining wakefulness and alertness during resting state (Cavanna, Vilas,

Palmucci, & Tagliazucchi, 2018; Tononi, 2004). Furthermore, this balance of dynamic

configurations also indicated that the DMN might function in multistable regimes and

revealed the potential neural mechanism by which DMN activity supported

wakefulness and alertness in the resting state (Andrews-Hanna, 2012; Buckner et al.,

2008).

Functional reorganization of dynamic DMN configurations during sleep

Compared to the resting state, the SWS stage was consistently accompanied with

reduced brain activity, whereas commensurate brain activity has been reported in the

REM sleep stage (Horovitz et al., 2008). Consistent alterations in the average brain

activity associated with CAMPs during DMN dynamics were also observed in our

study, suggesting that the activities of CAMPs might also reveal the changes in

wakefulness during wakefulness and sleep. However, the reduced activities of all

CAMPs might not sufficiently explain the decrease in DMN activity observed during

deep sleep due to the stable spatial structures of these CAMPs during wakefulness and

sleep. The decrease in activity might result from the increased occurrence probability

of the cDMN and the decreased probabilities of other two high-activity CAMPs.

These inversely changed occurrence probabilities in different CAMPs revealed the

neural mechanism of reduced brain activity given that the DMN regions were shown

to prefer the low-activity state during deep sleep (Bazhenov, Timofeev, Steriade, &

Sejnowski, 2002; Diekelmann & Born, 2010).

The balance of dynamic DMN configurations was also disrupted during sleep,

indicating the functional reorganization of DMN dynamics. The reorganization of

DMN activity might be associated with the alteration of different wakefulness levels

in different sleep stages (Tononi, 2004). In the REM sleep stage, the dynamic

transitions between the aDMN and pDMN increased, indicating more communication

between anterior and posterior DMN regions. The communications displayed the

top-down and bottom-up mechanisms in DMN structure, both of which are important

for the information processing in the brain (Buschman & Miller, 2007; Theeuwes,

2010). Thus, we speculate that the communications between anterior and posterior

DMN regions might help elucidate the neurophysiological basis underlying the

preservation of the wakefulness level in the REM sleep stage.

In the SWS stage, the dynamic transitions between the low-activity micropattern

and two high-activity micropatterns increased significantly. Moreover, different types

of micropatterns corresponded to distinct up-down states in slow oscillations among

DMN regions. Accordingly, the transitions between the low-activity micropattern and

two high-activity micropatterns in DMN dynamics could be deemed as the transitions

within up-down states. The dominant transitions of up-down states in deep sleep

further suggested the physiological importance of these increased dynamic transitions.

However, the dynamic transitions within the two high-activity micropatterns

decreased in the SWS stage. These reductions supported our hypothesis that

communications between anterior and posterior DMN regions are important for levels

of wakefulness and alertness given that wakefulness and alertness are almost lost

during deep sleep. The loss of wakefulness and alertness might not be caused by the

change in a single type of dynamic transition within pairs of CAMPs. We

hypothesized that the balance of dynamic DMN configurations was the underlying

key neural mechanism supporting wakefulness and alertness, which emerged during

wakefulness and disappeared during sleep. The coordination and cooperation of all

CAMPs played a core role for the ability of the DMN in supporting wakefulness and

alertness.

Based on these findings, we propose a three-state model to describe the

relationship between DMN micropatterns and wakefulness levels observed during

wakefulness and sleep. As shown in Fig. 6, the three CAMPs involved in DMN

dynamics are the basis of this model, and their interactions refer to the underlying

mechanism regulating the wakefulness level observed in distinct stages. Equal

communications among the three CAMPs support conscious awareness in the

AWAKE stage. The communications between the low-activity micropattern (i.e.,

cDMN) and each high-activity micropattern (i.e., aDMN and pDMN) are important

for the SWS stage characterized by a low level of wakefulness. During the REM sleep

stage, communications within high-activity micropatterns are predominant.

According to the proposed three-state model, we hypothesize that preservation of

wakefulness and alertness not only requires information transition between anterior

and posterior DMN regions, but also need a state that all DMN regions remain silent

and relaxed. Information transition within anterior and posterior DMN regions is

mediated by up-down and bottom-up mechanisms and vital for supporting

wakefulness and alertness. The absence of this process could lead to the loss of

alertness in the SWS stage, and this process alone would result in the wakefulness

level of the REM sleep stage, which is more wakeful than the SWS stage and less

wakeful than the AWAKE stage. This phenomenon highlights the importance of the

silent pattern for all DMN regions during the resting state with wakefulness and

alertness.

Wakefulness and alertness in humans not only depend on the anterior-posterior

integration with DMN regions but also involves fronto-parietal task-positive

executive and attention networks. Moreover, the connections between DMN and task

positive networks also play critical roles in supporting wakefulness and alertness.

Given the limitation of neuroimaging measure, the current work only describes the

associations between DMN dynamics and wakefulness levels. The roles of

integrations between DMN and other networks could not be explored at present.

Further work could validate our model with DMN activity during the

wakefulness-sleep cycle and further track the roles of integrations among different

ICNs across different levels of wakefulness and alertness with human EEG signals.

Methodological perspectives

Consistent with the promising microstate analysis of EEG/LFP signals (Michel &

Koenig, 2018), the CAMP analysis reported herein also assumes that brain activity

consists of several distinct instantaneous patterns. The difference is that the CAMP

method focuses on the nature of brain activity in different regions and extracts

micropatterns from the envelope signals. Envelope signals imply temporal alterations

of brain power, and their decomposition directly reveals brain rhythm dynamics. In

addition, the coactive patterns analyzed in the CAMP method were selected based on

the distributions of extreme values in the envelope signals, which differs from the

method used in microstate analysis. Local extreme values in envelope signals

represent the instantaneous higher/lower activities of brain regions followed by

contrasting changes in activity. The derived coactive patterns were thus considered to

represent the activity patterns leading to inversion of activity among regions in

specific brain networks. Therefore, we postulate that the proposed CAMP method will

help researchers elucidate coactive micropatterns in specific brain networks and

reveal additional underlying information about fast brain dynamics.

Limitations

Although we reveal several interesting findings in the current work, some

limitations exist that should be taken into consideration. First, the rat DMN is slightly

anatomically different from the human DMN, and our findings in rat DMN dynamics

need to be validated in the human DMN during wakefulness and sleep. In addition,

the present work reveals the dynamic configurations of DMN activity exclusively in

the gamma band, and different frequency bands have distinct physiological roles. The

relationships between the DMN dynamics in other frequency bands with wakefulness

levels should be investigated in future studies.

Conclusion

The DMN is believed to be associated with neural mechanisms of wakefulness

and alertness levels, whereas fast dynamics of DMN activity based on direct

physiological recordings in different stages of wakefulness remain unclear. We

highlighted that the fast dynamics of DMN activity during wakefulness and sleep

shared structurally stable CAMPs, whereas their dynamic configurations were specific

to different levels of wakefulness. Our results indicated the reorganization of DMN

dynamics during wakefulness and sleep, and provided a three-state model to reveal

the fundamental neural associations between DMN activity and wakefulness levels.

Acknowledgments

We thank Professor Pedro A. Valdes-Sosa for the valuable discussions and

suggestions regarding this work.

Authorship Confirmation Statement

Y.X., D.Y., C.Z., and D.G. designed the research; Y.C., M.L., and W.J. performed

the research; Y.C., M.L., B.B., and H.L. analyzed the data; and C.Y., B.B., D.G., and

D.Y. wrote the paper.

Authors' Disclosure Statement

The authors declare that no competing financial interests exist.

Funding Statement

This study was supported by the National Natural Science Foundation of China

(grant nos. 81861128001, 61527815, 31771149, 61761166001, 61871420, 11975194

and 81901366), the Sichuan Science and Technology Program (grant no.

2018HH0003), and the 111 project (grant no. B12027).

References

Andrews-Hanna, J. R. (2012). The Brain's Default Network and its Adaptive Role in

Internal Mentation. *Neuroscientist*, *18*(3), 251–270.

https://doi.org/10.1016/j.neuroimage.2013.08.045.The

Bazhenov, M., Timofeev, I., Steriade, M., & Sejnowski, T. J. (2002). Model of

thalamocortical slow-wave sleep oscillations and transitions to activated states.

The Journal of Neuroscience, *22*(19), 1.

Beckmann, C. F., DeLuca, M., Devlin, J. T., & Smith, S. M. (2005). Investigations

into resting-state connectivity using independent component analysis.

Philosophical Transactions of the Royal Society B: Biological Sciences,

360(1457), 1001–1013. https://doi.org/10.1098/rstb.2005.1634

Brett L. Foster, Mohammad Dastjerdi,   and J. P. (2012). Neural populations in

human posteromedial cortex display opposing responses during memory and

numerical processing. *Proceedings of the National Academy of Sciences*, *109*,

15514–15519.

https://doi.org/10.1073/pnas.1206580109/-/DCSupplemental.www.pnas.org/cgi/

doi/10.1073/pnas.1206580109

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default

network: Anatomy, function, and relevance to disease. *Annals of the New York*

Academy of Sciences, *1124*, 1–38. https://doi.org/10.1196/annals.1440.011

Buschman, T. J., & Miller, E. K. (2007). Top-Down Versus Bottom-Up Control of

Attention in the Prefrontal and. *Science*, *315*(March), 1860–1863.

https://doi.org/10.1126/science.1138071

Cavanna, F., Vilas, M. G., Palmucci, M., & Tagliazucchi, E. (2018). Dynamic

functional connectivity and brain metastability during altered states of

consciousness. *NeuroImage*, *180*, 383–395.

505 https://doi.org/10.1016/j.neuroimage.2017.09.065

506 Chen, J. E., Glover, G. H., Greicius, M. D., & Chang, C. (2017). Dissociated patterns

507  of anti-correlations with dorsal and ventral default-mode networks at rest.

508  *Human Brain Mapping*, *38*(5), 2454–2465. https://doi.org/10.1002/hbm.23532

509 Crunelli, V., & Hughes, S. W. (2010). The slow (1 Hz) rhythm of non-REM sleep: A

510  dialogue between three cardinal oscillators. *Nature Neuroscience*, *13*(1), 9–17.

511  https://doi.org/10.1038/nn.2445

512 Cui, Y., Yu, S., Zhang, T., Zhang, Y., Xia, Y., Yao, D., & Guo, D. (2018). Altered

513  activity and information flow in the default mode network of pilocarpine-induced

514  epilepsy rats. *Brain Research*, *1696*, 71–80.

515  https://doi.org/10.1016/j.brainres.2018.05.012

516 Delano-Wood, L., Bangen, K. J., Sorg, S. F., Clark, A. L., Schiehser, D. M., Luc,

517  N., … Bigler, E. D. (2015). Brainstem white matter integrity is related to loss of

518  consciousness and postconcussive symptomatology in veterans with chronic mild

519  to moderate traumatic brain injury. *Brain Imaging and Behavior*, *9*(3), 500–512.

520  https://doi.org/10.1007/s11682-015-9432-2

521 Diekelmann, S., & Born, J. (2010). The memory function of sleep. *Nature Reviews.*

522  *Neuroscience*, *11*(2), 114–126. https://doi.org/10.1038/nrn2762

523 Foster, B. L., Rangarajan, V., Shirer, W. R., & Parvizi, J. (2015). Intrinsic and

524  task-dependent coupling of neuronal population activity in human parietal cortex.

525  *Neuron*, *86*(2), 578–590. https://doi.org/10.1016/j.neuron.2015.03.018

526 Fox, K. C. R., Foster, B. L., Kucyi, A., Daitch, A. L., & Parvizi, J. (2018). Intracranial

Electrophysiology of the Human Default Network. *Trends in Cognitive Sciences*.

https://doi.org/10.1016/j.tics.2018.02.002

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Essen, C. Van, Raichle, M.

E., … Raichle, M. E. (2016). Networks Linked references are available on

JSTOR for this article⬜: The human brain is intrinsically organized into

dynamic , anticorrelated functional networks. *Proceedings of the National*

Academy of Sciences of the United States of America, (May).

Gretenkord, S., Olthof, B. M. J., Stylianou, M., Rees, A., Gartside, S. E., & LeBeau, F.

E. N. (2020). Electrical stimulation of the ventral tegmental area evokes

sleep-like state transitions under urethane anaesthesia in the rat medial prefrontal

cortex via dopamine D1-like receptors. *European Journal of Neuroscience*, *52*(2),

2915–2930. https://doi.org/10.1111/ejn.14665

Gusnard, D. A., Akbudak, E., Shulman, G. L., & Raichle, M. E. (2001). Medial

prefrontal cortex and self-referential mental activity: Relation to a default mode

of brain function. *Proceedings of the National Academy of Sciences of the United*

States of America, *98*(7), 4259–4264. https://doi.org/10.1073/pnas.071043098

Hagmann, P., Cammoun, L., Gigandet, X., Meuli, R., Honey, C. J., Van Wedeen, J.,

& Sporns, O. (2008). Mapping the structural core of human cerebral cortex.

PLoS Biology, *6*(7), 1479–1493. https://doi.org/10.1371/journal.pbio.0060159

Heidbreder, C. A., & Groenewegen, H. J. (2003). The medial prefrontal cortex in the

rat: Evidence for a dorso-ventral distinction based upon functional and

anatomical characteristics. *Neuroscience and Biobehavioral Reviews*, *27*(6),

555–579. https://doi.org/10.1016/j.neubiorev.2003.09.003

Holcman, D., & Tsodyks, M. (2006). The emergence of up and down states in cortical

networks. *PLoS Computational Biology*, *2*(3), 174–181.

https://doi.org/10.1371/journal.pcbi.0020023

Horovitz, S. G., Fukunaga, M., De Zwart, J. A., Van Gelderen, P., Fulton, S. C.,

Balkin, T. J., & Duyn, J. H. (2008). Low frequency BOLD fluctuations during

resting wakefulness and light sleep: A simultaneous EEG-fMRI study. *Human*

Brain Mapping, *29*(6), 671–682. https://doi.org/10.1002/hbm.20428

Hsu, L.-M., Liang, X., Gu, H., Brynildsen, J. K., Stark, J. A., Ash, J. A., … Yang, Y.

(2016). Constituents and functional implications of the rat default mode network.

Proceedings of the National Academy of Sciences, *113*(31), E4541–E4547.

https://doi.org/10.1073/pnas.1601485113

Huang, S. M., Wu, Y. L., Peng, S. L., Peng, H. H., Huang, T. Y., Ho, K. C., & Wang,

F. N. (2016). Inter-Strain Differences in Default Mode Network: A Resting State

fMRI Study on Spontaneously Hypertensive Rat and Wistar Kyoto Rat.

Scientific Reports, *6*(February), 1–9. https://doi.org/10.1038/srep21697

Jercog, D., Roxin, A., Barthó, P., Luczak, A., Compte, A., & De La Rocha, J. (2017).

UP-DOWN cortical dynamics reflect state transitions in a bistable network.

ELife, *6*, 1–33. https://doi.org/10.7554/eLife.22425

Ji, D., & Wilson, M. A. (2007). Coordinated memory replay in the visual cortex and

hippocampus during sleep. *Nature Neuroscience*, *10*(1), 100–107.

https://doi.org/10.1038/nn1825

Jing, W., Guo, D., Zhang, Y., Guo, F., Valdés-Sosa, P. A., Xia, Y., & Yao, D. (2017).

Reentrant information flow in electrophysiological rat default mode network.

Frontiers in Neuroscience, *11*(FEB), 1–12.

https://doi.org/10.3389/fnins.2017.00093

Kapogiannis, D., Reiter, D. A., Willette, A. A., & Mattson, M. P. (2014). Dynamic

functional connectivity of the default mode network tracks daydreaming.

NeuroImage, *100*, 112–119.

https://doi.org/10.1016/j.neuroimage.2012.09.029.Posteromedial

Karim Jerbi, R., V. J., Tomas , , O., Dalal, S. S., Julien Jung, D. H., Minotti, L., …

Lachaux,   and J.-P. (2010). Exploring the electrophysiological correlates of the

default-mode network with intracerebral EEG. *Frontiers in Systems*

Neuroscience, *4*(June), 1–9. https://doi.org/10.3389/fnsys.2010.00027

Larson-Prior, L. J., Zempel, J. M., Nolan, T. S., Prior, F. W., Snyder, A., & Raichle,

M. E. (2009). Cortical network functional connectivity in the descent to sleep.

Proceedings of the National Academy of Sciences of the United States of

America, *106*(11), 4489–4494. https://doi.org/10.1073/pnas.0900924106

Liu, Q., Farahibozorg, S., Porcaro, C., Wenderoth, N., & Mantini, D. (2017).

Detecting large-scale networks in the human brain using high-density

electroencephalography. *Human Brain Mapping*, *38*(9), 4631–4643.

https://doi.org/10.1002/hbm.23688

Liu, X., & Duyn, J. H. (2013). Time-varying functional network information extracted

from brief instances of spontaneous brain activity. *Proceedings of the National*

Academy of Sciences of the United States of America, *110*(11), 4392–4397.

https://doi.org/10.1073/pnas.1216856110

Logothetis, N. K., Pauls, J., Augath, M., Trinath, T., & Oeltermann, A. (2001).

Neurophysiological investigation of the basis of the fMRI signal. *Nature*,

412(6843), 150–157. https://doi.org/10.1038/35084005

Logothetis, Nikos K. (2002). The neural basis of the blood-oxygen-level-dependent

functional magnetic resonance imaging signal. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *357*(1424), 1003–1037.

https://doi.org/10.1098/rstb.2002.1114

Lőrincz, M. L., Gunner, D., Bao, Y., Connelly, W. M., Isaac, J. T. R., Hughes, S. W.,

& Crunelli, V. (2015). A distinct class of slow (~0.2–2 Hz) intrinsically bursting

layer 5 pyramidal neurons determines UP/DOWN state dynamics in the

neocortex. *Journal of Neuroscience*, *35*(14), 5442–5458.

https://doi.org/10.1523/JNEUROSCI.3603-14.2015

Lu, H., Zou, Q., Gu, H., Raichle, M. E., Stein, E. A., & Yang, Y. (2012). Rat brains

also have a default mode network. *Proceedings of the National Academy of*

Sciences of the United States of America, *109*(10), 3979–3984.

https://doi.org/10.1073/pnas.1200506109

Magri, C., Schridde, U., Murayama, Y., Panzeri, S., & Logothetis, N. K. (2012). The

Amplitude and Timing of the BOLD Signal Reflects the Relationship between

Local Field Potential Power at Different Frequencies. *The Journal of*

Neuroscience, *32*(4), 1395–1407.

615 https://doi.org/10.1523/JNEUROSCI.3985-11.2012

616 Michel, C. M., & Koenig, T. (2018). EEG microstates as a tool for studying the

617 temporal dynamics of whole-brain neuronal networks: A review. *NeuroImage*,

618 *180*(November 2017), 577–593.

619 https://doi.org/10.1016/j.neuroimage.2017.11.062

620 Ongur, D., & Price, J. (2000). The Organization of Networks within the Orbital and

621 Medial Prefrontal Cortex of Rats, Monkeys and Humans. *Cerebral Cortex*, *10*(3),

622 206–219. https://doi.org/10.1093/cercor/10.3.206

623 Ossandon, T., Jerbi, K., Vidal, J. R., Bayle, D. J., Henaff, M.-A., Jung, J., … Lachaux,

624 J.-P. (2011). Transient Suppression of Broadband Gamma Power in the

625 Default-Mode Network Is Correlated with Task Complexity and Subject

626 Performance. *Journal of Neuroscience*, *31*(41), 14521–14530.

627 https://doi.org/10.1523/JNEUROSCI.2483-11.2011

628 Panda, R., Bharath, R. D., Upadhyay, N., Mangalore, S., Chennu, S., & Rao, S. L.

629 (2016). Temporal Dynamics of the Default Mode Network Characterize

630 Meditation-Induced Alterations in Consciousness. *Frontiers in Human*

631 *Neuroscience*, *10*(July), 1–12. https://doi.org/10.3389/fnhum.2016.00372

632 Perez-Zabalza, M., Reig, R., Manrique, J., Jercog, D., Winograd, M., Parga, N., &

633 Sanchez-Vives, M. V. (2020). Modulation of cortical slow oscillatory rhythm by

634 GABAB receptors: an in vitro experimental and computational study. *Journal of*

635 *Physiology*, *598*(16), 3439–3457. https://doi.org/10.1113/JP279476

636 Raichle, M E, MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., &

Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the*

National Academy of Sciences of the United States of America, *98*(2), 676–682.

https://doi.org/10.1073/pnas.98.2.676

Raichle, Marcus E. (2015). The Brain ' s Default Mode Network. *Annual Review*

Neuroscience, (April), 413–427.

https://doi.org/10.1146/annurev-neuro-071013-014030

Raichle, Marcus E, & Mintun, M. A. (2006). Brain work and brain imaging. *Annual*

Review of Neuroscience, *29*, 449–476.

https://doi.org/10.1146/annurev.neuro.29.051605.112819

Reep, R. L., Chandler, H. C., King, V., & Corwin, J. V. (1994). Rat posterior parietal

cortex: topography of corticocortical and thalamic connections. *Experimental*

Brain Research, *100*(1), 67–84. https://doi.org/10.1007/BF00227280

Rosazza, C., & Minati, L. (2011). Resting-state brain networks: Literature review and

clinical applications. *Neurological Sciences*, *32*(5), 773–785.

https://doi.org/10.1007/s10072-011-0636-y

Sämann, P. G., Wehrle, R., Hoehn, D., Spoormaker, V. I., Peters, H., Tully, C., …

Czisch, M. (2011). Development of the brain's default mode network from

wakefulness to slow wave sleep. *Cerebral Cortex*, *21*(9), 2082–2093.

https://doi.org/10.1093/cercor/bhq295

Scheering, R., Koopmans, P. J., Van Mourik, T., Jensen, O., & Norris, D. G. (2016).

The relationship between oscillatory EEG activity and the laminar-specific

BOLD signal. *Proceedings of the National Academy of Sciences of the United*

States of America, *113*(24), 6761–6766.

https://doi.org/10.1073/pnas.1522577113

Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V., & Greicius, M. D. (2012).

Decoding subject-driven cognitive states with whole-brain connectivity patterns.

Cerebral Cortex, *22*(1), 158–165. https://doi.org/10.1093/cercor/bhr099

Theeuwes, J. (2010). Top-down and bottom-up control of visual selection. *Acta*

Psychologica, *135*(2), 77–99. https://doi.org/10.1016/j.actpsy.2010.02.006

Tononi, G. (2004). *An information integration theory of consciousness. BMC*

neuroscience (Vol. 5). BioMed Central. https://doi.org/10.1186/1471-2202-5-42

Tu, W., Ma, Z., Ma, Y., Dopfel, D., & Zhang, N. (2020). Suppressing Anterior

Cingulate Cortex Modulates Default Mode Network and Behavior in Awake

Rats. *Cerebral Cortex*, 1–12. https://doi.org/10.1093/cercor/bhaa227

Upadhyay, J., Baker, S. J., Chandran, P., Miller, L., Lee, Y., Marek, G. J., … Day, M.

(2011). Default-Mode-Like Network Activation in Awake Rodents. *PLoS ONE*,

6(11). https://doi.org/10.1371/journal.pone.0027839

Vatansever, D., Menon, D. K., Manktelow, A. E., Sahakian, B. J., & Stamatakis, E. A.

(2015). Default mode network connectivity during task execution. *NeuroImage*,

122, 96–104. https://doi.org/10.1016/j.neuroimage.2015.07.053

Vatansever, Deniz, Manktelow, A., Sahakian, B. J., Menon, D. K., & Stamatakis, E.

A. (2018). Default Mode Network Engagement beyond Self-Referential Internal

Mentation. *Brain Connectivity*, *8*(4), 245–253.

https://doi.org/10.1089/brain.2017.0489

681 Vidaurre, D., Smith, S. M., & Woolrich, M. W. (2017). Brain network dynamics are

682   hierarchically organized in time. *Proceedings of the National Academy of*

683   *Sciences*, 201705120. https://doi.org/10.1073/pnas.1705120114

684 Wu, X. jie, Zeng, L. L., Shen, H., Yuan, L., Qin, J., Zhang, P., & Hu, D. (2017).

685   Functional network connectivity alterations in schizophrenia and depression.

686   *Psychiatry Research - Neuroimaging*, *263*(March), 113–120.

687   https://doi.org/10.1016/j.pscychresns.2017.03.012

688 Wyss, J. M., & Vangroen, T. (1992). Connections between the retrosplenial cortex

689   and the hippocampal-formation in the rat - a review. *Hippocampus*, *2*(1), 1–12.

690   https://doi.org/10.1002/hipo.450020102

692 **Correspondence Addresses**

693 Daqing Guo

694 The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for

695 NeuroInformation, University of Electronic Science and Technology of China,

696 Chengdu 611731, China

697 dqguo@uestc.edu.cn

698 Dezhong Yao

699 The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for

700 NeuroInformation, University of Electronic Science and Technology of China,

701 Chengdu 611731, China

702 dyao@uestc.edu.cn

Bharat Biswal

The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for

NeuroInformation, University of Electronic Science and Technology of China,

Chengdu 611731, China

bbiswal@gmail.com

Figure 1. The placement of 15 intracranial electrodes.

Figure 2. Schematic of the CAMP procedure and three CAMPs of gamma activity in the

DMN during wakefulness and sleep. (a) The original LFPs. (b) The envelope signals (blue

lines) were extracted by applying the Hilbert transform to the bandpass-filtered signals (gray

lines). (c) All the envelope signals were downsampled (blue lines), and the extreme values

were detected as the active points for each channel (red dots). The dotted lines suggest the

coactive points in which more than N (N=7 in the present study) active points were observed

across DMN regions. (d) The coactive patterns were the maps of activity of all DMN regions

at coactive points. (e) The k-means clustering algorithm was applied to all coactive patterns to

detect the CAMPs. (f) A criterion was employed to remove several coactive points and

increase the aggregation of the CAMPs. The final CAMPs and CAMP index detected in this

step were subjected to further analyses. (g) Spatial structure of the common low-activity level

micropattern (cDMN). (h) Spatial structure of the anterior high-activity level micropattern

724   (aDMN). (i) Spatial structure of the posterior high-activity level micropattern (pDMN).

Figure 3. Comparisons of the temporal features and activity levels of each CAMP during wakefulness and sleep. (a) Comparisons of the total occurrence of each CAMP in different stages of wakefulness. The dots represent the values obtained from 29 rats, and the black stars indicate significant differences with a corrected $p<0.001$. (b) Comparisons of the total duration. (c) Comparisons of the mean duration. (d) Comparisons of the mean DMN activity during wakefulness and sleep for different CAMPs. (e-j) Comparisons of activity in DMN nodes for different CAMPs across different stages of wakefulness: (e and h) cDMN, (f and i) aDMN, and (g and j) pDMN. Gray dots indicate decreased normalized activity and black dots indicate increased normalized activity. The size of the dot reflects the value of the difference, and the red stars indicate significance differences with a corrected $p<0.001$.

Figure 4. Characteristics of CAMPs and the transitions among them in different stages of

wakefulness during wakefulness and sleep. (a) Comparisons of the occurrence probability

for all CAMPs in the three stages. The black dots indicate the values of the occurrence

probability obtained from 29 rats in different CAMPs and stages. The black stars indicate

significant differences with a corrected p<0.001. (b) Comparisons of the duration probability.

(c) Comparisons of the mean duration. (d-f) The transition structures among CAMPs for the

AWAKE (d), SWS (e) and REM sleep stages (f). All the numbers indicate the mean TPs

calculated for the 29 rats and the standard deviation. The numbers in blue indicate a

significantly lower transition probability than observed in the AWAKE stage, and the numbers

in red indicate a significantly higher transition probability. The significance level is a

corrected p<0.001.

Figure 5. Phase locking relationship between each CAMP with slow oscillations in the SWS

stage. (a-c) The phase locking relationships between the cDMN (a), aDMN (b) and pDMN (c)

with the slow oscillations in anterior DMN regions. (d-f) The phase locking relationships

between the cDMN (d), aDMN (e) and pDMN (f) with the slow oscillations in posterior

DMN regions. The red lines showed the significant directionality with Rayleigh test p<0.001.

Figure 6. The three-state model of the alertness levels during wakefulness and sleep. The

AWAKE stage requires the cooperation of all three CAMPs, while the SWS stage requires

communications between the low-activity micropattern (cDMN) and the high-activity

micropatterns (aDMN or pDMN). The REM sleep stage requires interactions within the two

high-activity micropatterns.