Exosomes are predominantly loaded with mRNA transcript encoding cytoplasmic

proteins and exclude mRNA transcript encoding nuclear proteins

Shabirul Haque [1] and Sarah R. Vaiselbuh [1, 2]

1. The Feinstein Institute for Medical Research, Northwell Health, 350 Community Drive,

Manhasset, New York 11030. 2. Division of Hematology/Oncology-Department of

Pediatrics, Staten Island University Hospital at Northwell Health, New York.

Correspondence:

Sarah R. Vaiselbuh, MD, MBA

Associate Professor in Pediatrics & Molecular Medicine, Zucker School of Medicine

Assistant Professor, Center for Oncology & Cell Biology,

The Feinstein Institute of Medical Research

Director, Pediatric Hematology/Oncology, Children's Cancer Center,

Staten Island University Hospital at Northwell Health

E-mail: svaiselbuh@gmail.com

Communication and portal handling by:

Shabirul Haque, PhD

Feinstein Institute for Medical Research, Northwell Health, Manhasset, New York

Email: shaque@northwell.edu and shabirul7@gmail.com

Phone: 508-667-0785

Running title: Exosomes are rich with mRNA transcripts encoding cytoplasmic proteins.

Keywords: Exosomal mRNA; cytoplasmic proteins; nuclear proteins; exosomes

Abstract text count: 199; Main text count: 3,137; Table: 2;   Figures: 4

Abbreviations: Exo-CM: Exosomes derived from conditioned medium, Exo-cDNA: direct

conversion of exosomal RNA into cDNA.

Graphical Abstract

Abstract

Exosomes are nanovesicles (~30-150 nm diameters) released via an endocytic pathway

in almost all mammalian cell types. Exosomes are composed of a lipid bilayer membrane

that encloses RNA, miRNA, proteins and DNA. This manuscript unravels how exosome

cargo is collected by a highly precise process delineating two separate mRNA transcript

entities encoding cytoplasmic and nuclear proteins separately.

Ultracentrifuge isolated exosomes were directly converted into cDNA (Exo-cDNA),

by a method developed in our laboratory. Cellular RNA was extracted from each cell line

and cDNA was prepared (Cell-cDNA). We amplified mRNA transcripts translating

cytoplasmic proteins CD10 and CXCR4 and mRNA transcripts translating *nuclear*

proteins such as proliferating cell nuclear antigen (PCNA), CREB-BP, activation induced

cytidine deaminase (AID), and terminal deoxynucleotidyl transferase (TdT). We amplified

all four different mRNA transcripts (PCNA, CREB-BP, AID, and TdT) from *cellular* cDNA

but none from *exosomal* cDNA (Exo-cDNA). These findings suggest that exosomes carry

mRNA transcripts encoding *cytoplasmic* proteins only but mRNA transcripts encoding

nuclear proteins could not be detected. This important observation could prove to be

crucial for the exosome research community since it sheds light on one of the limitations

relating to the use of exosomes as biomarkers in cancer biology and other diseases.

1. Introduction

Extracellular vesicles such as exosomes are spherical nanoparticles derived from almost all kind of cells in normal and pathological conditions. The exosome diameter size varies from 30-150 nm, and they originate from the cytoplasm of the cell by inward budding of multivesicular bodies [1]. Exosomal content is rich in proteins, DNA, mRNA and non-coding RNA [2, 3]. Exosomes have been isolated *in vivo* from most bio-fluids such as saliva, serum, urine, breast milk, cerebrospinal fluid, and amniotic fluid [4-7]. In addition, exosomes have been isolated *in vitro* from conditioned medium (CM) of cultured cell lines by ultracentrifugation method [8]. The application of exosomes as biomarkers for disease in cancer research has gained momentum to facilitate early diagnosis, early relapse, risk stratification, and real-time monitoring of the disease in a real time strategy. Exosomes carry the molecular signature of the parent cells. Their major functions are being cell-to-cell communication and exchange of bio-active molecules (such as mRNA and miRNA, DNA, proteins etc.) into target or recipient cells. These exosomal cargo molecules are delivered by a process called 'exosomal shuttle RNA' (esRNA) which remains biologically functional in recipient cells [9].

Gene transcription is a multistep and multistage complex process. After mRNA transcription in the nucleus, mRNA transcripts pass through several nuclear modification/process (e.g. 5 prime capping, splicing, and 3 prime ends processing etc.) before being exported to the cytosol. These modifications direct the fate of mRNA transcripts (such as stability, life span, folding etc.) into the cytosol [10]. When processed mRNA transcripts get out of the nucleus they are exposed to the cytosolic milieu in the cytosol. The critical point is, if the mRNA transcripts are poorly processed, they fail to

survive and sustain in the cytosolic medium. Several proteins are responsible for nuclear processing of mRNA transcripts, as a result of which many proteins are adhered or anchored onto the mRNA transcripts. These adhered proteins on an mRNA transcript also regulate its export, localization, translation, and stability [10]. It has been established that exosomes collect its entire cargo such as proteins, lipids and coding and non-coding RNAs & DNA directly from the cytoplasmic and/or cytoplasmic membrane derived particles [9]. There are also a few reports which illuminates how different varieties of exosomal RNA species get packaged and loaded in a stoichiometric fashion depending on their source. Typically, serum-derived exosomes contain 30-75% miRNA, 10-20% mRNA, 5-30% rRNA while urine-derived exosomes showed 2-10% miRNA, 5-12% mRNA, 30-60% rRNA [11]. Another study showed that within exosomal RNA, rRNA (28S and 18S fragment) are the most abundant (accounting for ~97%) and 2% are known RNA (miRNA, mRNA etc.), while 1% are unknown genes [12]. It appears from literature that the knowledge on exosomal RNA content is quite conflicting and limited, and as exosome research is still evolving, investigators are trying to learn about packaging of mRNA transcripts into exosomes.

We hypothesized that selection of mRNA transcripts by the exosomes during exosomal biogenesis is a very precise process. Exosomes selectively entrap mRNA transcripts encoding cytoplasmic proteins, per exclusion of mRNA transcripts that encode nuclear proteins.

2. Material and Methods

2. 1. Cell lines and reagents. Cell lines of acute lymphatic leukemia (ALL), SUP-B15 (cat# ATCC® CRL-1929 ™), and JM1 cells (cat # ATCC® CRL-10423™) were obtained from ATCC. CL-01 cells obtained (Chiorazzi lab, The Feinstein Institute for Medical research, Northwell Health, Manhasset, NY). Agarose (cat # CA3510-8, Denville Scientific, Inc.). Tris-borate-EDTA buffer (cat # 28355, Thermo Scientific), SYBR Safe (cat # S33102, Invitrogen), 6x loading dye (cat # R0611, Crystalgen). RNase Inhibitor (4 unit/µl) (cat # 1055213, Qiagen), rRNasin Rnase inhibitor (40 U/µl) (Cat # N251B, Promega), RT-Superscript III (200 U/µl) (Cat # 18080-044, Invitrogen), random hexamer primers (cat # SO142, Thermo fisher scientific).

2. 2. Cell culture. JM1 cells were expanded and cultured in IMDM supplemented with 0.05 mM β-mercaptoethanol, 10 % FBS, and 1x Penicillin-Streptomycin. SUP-B15 cells were cultured in Iscove's Modified Dulbecco's Medium (IMDM), supplemented with 1.5 g/L sodium bicarbonate, 0.05 mM β-mercaptoethanol, 20% fetal bovine serum (not heat inactivated), and 1x Penicillin-Streptomycin. CL-01 cells were expanded and cultured in RPMI-1640 supplemented with 10 % FBS, and 1x Penicillin-Streptomycin. Each cell lines were incubated in a humidified incubator at 37°C with 5% CO_2.

2. 3. Depletion of exosomes from FBS (Exosome-free FBS). Fetal bovine serum (FBS) are abundantly loaded with exosomes. FBS was converted into exosome-free FBS by centrifugation and ultracentrifugation method [8]. In short, FBS was subjected to centrifugation for 10 min at 300 xg. Supernatant was collected and centrifuged for another

10 min at 2000 xg. Supernatant was collected again and centrifuged for another 30 min at 10,000 xg. Supernatant was then placed into ultra-centrifugation for overnight at 100,000 xg. FBS supernatant was collected next day and labeled as exo-free FBS. Each centrifugation step was carried out at 4°C [8].

2. 4. Exosome isolation from conditioned medium of cell lines. Cell line was cultured in cell culture medium supplemented with Exo-free FBS (FBS depleted with exosomes). Cells (1×10^6/ml) were plated in 10 ml total volume in a 100 mm tissue culture dish. After 48 hours, CM of each cell line was harvested by centrifugation and filtered through a 0.22 micron filter system to remove cell debris. Filtered CM subjected for exosome isolation by the method of ultracentrifugation.

2. 5. Exosome isolation. Exosomes were isolated by the method of centrifugation and followed by ultracentrifugation [8]. Human serum and conditioned medium of cell lines were subjected to centrifugation for 10 min at 300 xg. Collected supernatant was again centrifuged for 10 min at 2000 X g. Supernatant was collected and centrifuged for 30 min at 10,000 xg. Then supernatant was ultra-centrifuged for 120 min at 100,000 xg. Pellets after ultracentrifugation were re-suspended in PBS and ultra-centrifuged for another 120 min at 100,000 xg. Pellets containing exosomes were harvested and reconstituted in around 100-200 μl ice-cold PBS. Each centrifugation and ultra-centrifugation steps were carried out at 4°C. Exosomal protein content was estimated by BCA protein assay kit (Bio-Rad) and stored at -80°C for further usage.

2. 6. Direct conversion of exosomes into cDNA (Exo-cDNA). Exosomes were directly converted into cDNA by following a method published by Haque and Vaiselbuh [13].

2. 7. Cellular RNA isolation and cDNA synthesis. Total RNA was extracted from SUP-B15, JM1, and CL-01 cells by the method of Trizol (Invitrogen). RNA concentration and quality was analyzed by NanoDrop ND1000 spectrophotometer. We took cellular RNA (2-5 μg) for cDNA synthesis. We used Oligo-dT primers (Invitrogen), M-MLV reverse transcriptase (Invitrogen) and followed manufacturer's protocol for cDNA synthesis.

2. 8. Gene amplification by two-step PCR. Gene amplification such as β-actin, CD10, CXCR4, PCNA, CREB-BP, AID, and TdT was carried out by two-step PCR [13]. In short, 1st round of PCR was carried out using AccuPrime™ Pfx DNA Polymerase. 2nd round of PCR was carried out using Ampli Taq Gold. Both rounds of PCR are ran under the following conditions: denaturation at 95°C for 30 seconds, annealing at 56°- 60°C for 60 seconds, and extension step at 72°C for 60 seconds for 35 cycles. Cell-cDNA or Exo-cDNA (2 μl) were used as template for the first round of PCR. Around 1 – 2 μl of the PCR product of the first round was used as template for the second round of PCR, AmpliTaq Gold was used to a final volume of 30 μl. Brief details of the primers such as gene name, gene bank, sequences, and annealing temperature are mentioned (**Table 1)**.

2. 9. Agarose gel electrophoresis. Briefly, 1.5% agarose was prepared in 1x Tris-borate-EDTA buffer. SYBR Safe was added to the melted agarose gel before polymerization. PCR product (5.0 μl) were mixed with 6x loading dye, mixed product were

loaded on to gel and ran for 20-25 minutes at 200 V. The gel was exposed and image

were captured by Bio-Rad gel doc system (Bio-Rad, Hercules, California, USA).

3. Results

3.1. Characterization of exosomes by Nanoparticle Tracking Analysis (NTA).

Exosomes were purified from CM by ultra-centrifugation method. Optimum diameter size

of isolated exosomal vesicles was confirmed by Nanoparticle Tracking Analysis (NTA–

Malvern) (**Fig. 1 A**). Representative NTA-deduced image of exosomes shown (**Fig. 1 B**).

Enrichment of CD63 on exosomes and no expression of calnexin (negative exosomes

marker) shown in our published article (13).

3. 2. Amplication of CD10 and CXCR4 from Cell-cDNA/Exo-cDNA. Both the CD10 and

CXCR4 proteins are localized within the cytoplasm of the cell. Cell-cDNA and Exo-cDNA

from the leukemia cell lines JM1 and SUP-B15 and control normal B-cells CL-01 were

used as templates to amplify CD10 and CXCR4 in two-step PCR (**Fig. 2**). We chose

CD10 as a marker of the above mentioned leukemia cell lines with CL-01 as negative

control. Cellular expression of CD10 was amplified in JM1 and SUP-B15 Cell-cDNA but

at exosomal level (Exo-cDNA), CD10 expression could not be detected after a first round

of PCR (**Fig. 2 A, upper panel**). After second round of PCR, CD10 expression was

detected in both Cell-cDNA and Exo-cDNA (**Fig. 2 A, lower panel**). CXCR4 was amplified

by PCR using Cell-cDNA obtained from JM1, SUP-B15, and CL-01 cells but remained

undetectable in Exo-cDNA in a first round of PCR (**Fig. 2 B, lower panel**). A second

round of PCR showed amplification of CXCR4 expression in both Cell-cDNA and Exo-

cDNA (**Fig. 2 B, lower panel**). Endogenous control, β-actin was considered internal

control for quality confirmation of Cell-cDNA and Exo-cDNA.

3. 3. Amplication of PCNA and CREB-BP from Cell-cDNA/Exo-cDNA. Both PCNA and

CREB-BP are nuclear proteins commonly used as cell markers. PCNA and CREB-BP

mRNA were amplified by 1^{st} and 2^{nd} rounds of PCR utilizing Cell-cDNA and Exo-cDNA

from each cell lines (JM1, SUP-B15, and CL-01) (**Fig. 3**). In a first round of PCR, PCNA

mRNA expression at cellular level (Cell-cDNA) was amplified in all three cell lines but

could not amplify at exosomal level (Exo-cDNA) (**Fig. 3 A, upper panel**). Similarly after a

second round of PCR, PCNA mRNA expression was detected at cellular level (Cell-

cDNA) but exosomal level (Exo-cDNA) did not show PCNA mRNA amplification (**Fig. 3**

A, lower panel). When we tested CREB-BP amplification, a comparable pattern was

observed: a first round of PCR showed CREB-BP mRNA amplified product from all three

cell lines derived Cell-cDNA, but Exo-cDNA did not produce a detectable level of PCR

product (**Fig. 3 B, upper panel**). A second round of PCR demonstrated consistent CREB-

BP mRNA amplification of Cell-cDNA, in contrast to Exo-cDNA that again could not

amplify a detectable amount of CREB-BP (**Fig. 3 B, lower panel**). Our findings are

suggestive of the fact that nuclear mRNA transcripts are not present in the cDNA derived

from exosomes. We confirmed this observation by using additional mRNA transcripts

that encode other *nuclear* proteins such as PCNA, p53, VBRAF600 and TdT and each

time, Exo-cDNA failed to amplify. Endogenous control (β-actin) was used as quality

confirmation of Cell-cDNA and Exo-cDNA.

3. 4. Amplication of AID and TdT from Cell-cDNA/Exo-cDNA. We tested two more proteins (AID and TdT) which are enzymatic in nature and expressed predominantly in the nucleus. PCR amplification of AID and TdT mRNA expression was carried out and electrophoresed on an agarose gel (**Fig. 4**). A first round of PCR showed AID mRNA amplification of Cell-cDNA derived from JM1, SUP-B15 and CL-01 cell lines but Exo-cDNA could not produce any AID PCR product and therefore could not be detected on agarose (**Fig. 4 A, upper panel**). The normal B cell line, CL-01, is known to overexpress AID. A second round of PCR demonstrated nuclear protein AID mRNA amplification with Cell-cDNA, but Exo-cDNA could not amplify any detectable amount of AID product (**Fig. 4 A, lower panel**). Similarly, PCR amplification of TdT mRNA encoding this nuclear protein failed to amplify cDNA derived from exosomes (**Fig. 4 B**). CL-01 cells were used as a negative control for TdT expression and housekeeping gene (β-actin) was used as confirmation of Cell-cDNA and Exo-cDNA.

Altogether, we have looked at five genes encoding cytoplasmic proteins (CD10, CD34, actin, CXCR4, HLA-DR) and five genes encoding nuclear proteins (p53, PCNA, TdT, CREB-BP, and AID). PCR amplification of all genes are summarized individually as positive and negative in cellular and exosomal cDNA (**Table 2**).

4. Discussion

Cell biology is defined as the study of the structure and compartments within the cell, while molecular biology can be defined as the study of cellular content within those compartments by analyzing the composition, structure and interaction of cellular molecules such as RNA, protein, DNA etc. The study of biogenesis of exosomes helps

us to understand the processing of exosomes by focusing closely on how both fields of

biology overlap. This manuscript discusses how the cellular compartmental origin of

exosomes delineates and discerns their very specific nature of mRNA transcripts. Our

data demonstrate that exosomes encapsulate mRNA transcripts that encode *cytoplasmic*

proteins and avoids mRNA transcript encoding *nuclear* proteins.

Exosomes were extracted from the CM of B-cell lines (JM1, SUP-B15, and CL-01)

and characterized by NTA system, showing that most of the exosome particle diameter

size was 65 nm. Others have reported that a particle size of 30-100 nm in diameter

identifies exosomes [14, 15]. We chose to amplify CD10 and CXCR4 mRNA by PCR

because both are cytoplasmic proteins. CD10 is expressed by early B, pro-B and pre-B

lymphocytes in acute lymphocytic leukemia (ALL) and lymph node germinal centers [16].

CD10 is demonstrated to be cytoplasmic in nature and oriented as a transmembrane

region [17]. CD10 is a common ALL antigen and an important cell surface marker in the

diagnosis of pediatric ALL. CD10 and CD34 combined expression has prognostic

importance and is associated with a better outcome in childhood ALL [18].

CXCR4 is predominantly a cytoplasmic protein but gets shuttle into the nucleus in

cancer settings, considered as poor prognostic event [19]. CXCR4 is a well-established

biomarker in both solid and liquid tumors. Augmented CXCR4 expression in tumors is

associated with low overall survival rate and in general is considered as a poor prognostic

feature in cancer [20]. In addition, high expression of CXCR4 as a biomarker in breast

cancer, predicts axillary lymph node metastasis [21, 22], and higher risk for bone

metastatic disease [23]. CXCR4 along with CXCL12 have been studied as predictive

biomarkers of glioma recurrence [24]. Based on studies mentioned, CXCR4 represents a

useful prognostic biomarker in cancer [25], CD10 also studied and reported as diagnostic

marker in follicular carcinoma (80% patients are positive) and thyroid carcinoma (77%),

is not detected in normal thyroid tissue [26]. Based on above reports and evidences we

can propose that amplification of exosomal CXCR4 and CD10 has great prospective in

early diagnosis or development of diagnostic marker, and prognostic markers in both

cancer settings (solid and liquid tumors).

Both acute lymphocytic leukemia cell lines (JM1, and SUP-B15), and the normal

B cell line (CL-01) showed amplification of the cytosolic mRNA transcript of CD10 and

CXCR4 expression after a second rounds of PCR amplified from both Cell-cDNA and

Exo-cDNA. CL-01 has a low expression of CD10, and therefore could not be detected

after the first round of PCR.

We chose another set of predominant *nuclear* proteins (PCNA and CREB-BP) for

mRNA amplification by two-step PCR. PCNA is a homo-trimer and acts as a scaffold to

recruit proteins involved in DNA repair, DNA replication, and chromatin remodeling

and epigenetics system [27]. It is a nuclear protein and also an indicator for cell

proliferation [28]. CREB-BP is exclusively expressed and localized in the perinuclear and

nuclear region [29]. CREB-BP is an ubiquitously expressed gene and mainly participates

in the activation of several different transcription factors which helps in embryonic

development, growth control, and homeostasis of cells. We found that first and second

rounds of PCR showed amplification PCNA and CREB-BP in Cell-cDNA but these two

nuclear gene markers could not be detected in Exo-cDNA.

Furthermore, two other nuclear enzymes (AID and TdT) were selected for PCR

amplification. Major functions of AID include somatic hypermutation, gene conversion,

and class-switch recombination of immunoglobulin genes in B cells of the immune system

[30, 31]. AID is highly expressed in germinal center B cells and contributes to the

generation of both protective antibodies against microbes and high affinity, class-switched

pathogenic auto-antibodies [32]. AID is also a nuclear protein and localized to the sub-

nuclear region [33]. There is also evidence that AID shuttles between the nucleus and

the cytoplasm like apolipoprotein B [34].

TdT is a unique DNA polymerase enzyme, expressed in a variety of cells such as

immature pre-B, pre-T lymphoid cells, ALL and lymphoma cells [16]. TdT catalyzes

addition of N-nucleotides to the V, D, and J exons of the TCR and BCR genes

during antibody gene recombination which lead to junctional diversity [35]. TdT is a

nuclear protein as reported [36]. In our study, we observed that Cell-cDNA of JM1 and

SUP-B15 cells amplified AID and TdT in first and second rounds of PCR but again no

amplification of AID and TdT was detected when Exo-cDNA was utilized as a template

for PCR. Cell-cDNA derived CL-01 cells could not amplify TdT which was expected, since

TdT is a marker for malignant B cells and is absent in normal B cells. We did not find

mRNA transcripts encoding any of the tested nuclear proteins (TdT, CREB-BP, AID, and

PCNA) in the exosomes. These findings intrigued us as to why exosomes are choosing

one set of mRNA transcripts and avoiding entrapment of another set of mRNA transcripts

during its process of packaging and biogenesis? Structurally, the size of the nucleus is

tremendously smaller than the whole cell with the nuclear volume accounting for only 8%

of the cellular volume [37]. As a result, cells may have tendencies to conserve mRNA

transcripts responsible for encoding nuclear proteins. Moreover, Hocine S and

colleagues [10] reported that when mRNA transcripts migrate out of the nucleus after

nuclear processing, many proteins which are adhered and tagged on to the mRNA transcripts regulate pathway and fate of the mRNA transcripts. These finding provide a platform to interpret and explain that there may be potentially two separate sets of mRNA transcripts, which are tagged by different sets of proteins directing mRNA transcripts, such that one set of proteins direct mRNA transcripts to enter into the exosomes (mRNA transcripts encoding cytoplasmic proteins) while another set of proteins direct mRNA transcripts to not enter into the exosomes (mRNA transcripts encoding nuclear proteins).

5. Conclusion

In conclusion, we show that exosomes carry mRNA transcripts encoding cytoplasmic proteins but not mRNA transcripts encoding nuclear proteins. Since exosomes are widely used as biomarkers for cancer and other diseases, we urge researchers in biomedical medicine to take our observation into account to possibly avoid premature negative conclusions. A selection of exosomal mRNA transcripts encoding *nuclear* proteins for biomarkers studies in exosomes, will not be expressed and therefore might not be the right choice for a biomarker study. In contrast, exosomal RNA transcripts of *cytoplasmic* proteins such as CXCR4/CD10 and others would be our recommended choice as potential exosomal biomarkers in various diseases conditions.

Acknowledgments: This work was supported by the Pediatric Oncology Fund at Staten Island University Hospital, The DiMartino Family Foundation and Island Auto group. We also thanks Malvern Instruments for the characterization of exosomes by NTA.

Author's contribution: S.H. conceived the concept, conducted the experiments,

analyzed data and wrote the manuscript. S.V. discussed that data for analysis and

reviewed the manuscript.

Conflict of interests: The authors have no competing interests to declare.

References:

[1] M. Yanez-Mo, P.R. Siljander, Z. Andreu, A.B. Zavec, F.E. Borras, E.I. Buzas, K. Buzas, E. Casal, F. Cappello,
J. Carvalho, E. Colas, A. Cordeiro-da Silva, S. Fais, J.M. Falcon-Perez, I.M. Ghobrial, B. Giebel, M. Gimona,
M. Graner, I. Gursel, M. Gursel, N.H. Heegaard, A. Hendrix, P. Kierulf, K. Kokubun, M. Kosanovic, V. Kralj-
Iglic, E.M. Kramer-Albers, S. Laitinen, C. Lasser, T. Lener, E. Ligeti, A. Line, G. Lipps, A. Llorente, J. Lotvall,
M. Mancek-Keber, A. Marcilla, M. Mittelbrunn, I. Nazarenko, E.N. Nolte-'t Hoen, T.A. Nyman, L. O'Driscoll,
M. Olivan, C. Oliveira, E. Pallinger, H.A. Del Portillo, J. Reventos, M. Rigau, E. Rohde, M. Sammar, F.
Sanchez-Madrid, N. Santarem, K. Schallmoser, M.S. Ostenfeld, W. Stoorvogel, R. Stukelj, S.G. Van der
Grein, M.H. Vasconcelos, M.H. Wauben, O. De Wever, Biological properties of extracellular vesicles and
their physiological functions, J Extracell Vesicles 4 (2015) 27066.
[2] B.K. Thakur, H. Zhang, A. Becker, I. Matei, Y. Huang, B. Costa-Silva, Y. Zheng, A. Hoshino, H. Brazier, J.
Xiang, C. Williams, R. Rodriguez-Barrueco, J.M. Silva, W. Zhang, S. Hearn, O. Elemento, N. Paknejad, K.
Manova-Todorova, K. Welte, J. Bromberg, H. Peinado, D. Lyden, Double-stranded DNA in exosomes: a
novel biomarker in cancer detection, Cell Res 24(6) (2014) 766-9.
[3] C. Thery, L. Zitvogel, S. Amigorena, Exosomes: composition, biogenesis and function, Nat Rev Immunol
2(8) (2002) 569-79.
[4] C. Admyre, S.M. Johansson, K.R. Qazi, J.J. Filen, R. Lahesmaa, M. Norman, E.P. Neve, A. Scheynius, S.
Gabrielsson, Exosomes with immune modulatory features are present in human breast milk, J Immunol
179(3) (2007) 1969-78.
[5] J. Nilsson, J. Skog, A. Nordstrand, V. Baranov, L. Mincheva-Nilsson, X.O. Breakefield, A. Widmark,
Prostate cancer-derived urine exosomes: a novel approach to biomarkers for prostate cancer, Br J Cancer
100(10) (2009) 1603-7.
[6] G. Rabinowits, C. Gercel-Taylor, J.M. Day, D.D. Taylor, G.H. Kloecker, Exosomal microRNA: a diagnostic
marker for lung cancer, Clin Lung Cancer 10(1) (2009) 42-6.
[7] J.P. Hegmans, M.P. Bard, A. Hemmes, T.M. Luider, M.J. Kleijmeer, J.B. Prins, L. Zitvogel, S.A. Burgers,
H.C. Hoogsteden, B.N. Lambrecht, Proteomic analysis of exosomes secreted by human mesothelioma
cells, Am J Pathol 164(5) (2004) 1807-15.
[8] C. Thery, S. Amigorena, G. Raposo, A. Clayton, Isolation and characterization of exosomes from cell
culture supernatants and biological fluids, Curr Protoc Cell Biol Chapter 3 (2006) Unit 3.22.
[9] H. Valadi, K. Ekstrom, A. Bossios, M. Sjostrand, J.J. Lee, J.O. Lotvall, Exosome-mediated transfer of
mRNAs and microRNAs is a novel mechanism of genetic exchange between cells, Nat Cell Biol 9(6) (2007)
654-9.
[10] S. Hocine, R.H. Singer, D. Grunwald, RNA processing and export, Cold Spring Harb Perspect Biol 2(12)
(2010) a000752.
[11] M. Li, E. Zeringer, T. Barta, J. Schageman, A. Cheng, A.V. Vlassov, Analysis of the RNA content of the
exosomes derived from blood serum and urine and its potential as biomarkers, Philos Trans R Soc Lond B
Biol Sci 369(1652) (2014).

[12] P. Jenjaroenpun, Y. Kremenska, V.M. Nair, M. Kremenskoy, B. Joseph, I.V. Kurochkin, Characterization
of RNA in exosomes secreted by human breast cancer cell lines using next-generation sequencing, PeerJ
1 (2013) e201.
[13] S. Haque, S.R. Vaiselbuh, Exosomes molecular diagnostics: Direct conversion of exosomes into the
cDNA for gene amplification by two-step polymerase chain reaction, J Biol Methods 5(3) (2018) e96.
[14] E. van der Pol, F.A. Coumans, A.E. Grootemaat, C. Gardiner, I.L. Sargent, P. Harrison, A. Sturk, T.G. van
Leeuwen, R. Nieuwland, Particle size distribution of exosomes and microvesicles determined by
transmission electron microscopy, flow cytometry, nanoparticle tracking analysis, and resistive pulse
sensing, J Thromb Haemost 12(7) (2014) 1182-92.
[15] V.S. Chernyshev, R. Rachamadugu, Y.H. Tseng, D.M. Belnap, Y. Jia, K.J. Branch, A.E. Butterfield, L.F.
Pease, 3rd, P.S. Bernard, M. Skliar, Size and shape characterization of hydrated and desiccated exosomes,
Anal Bioanal Chem 407(12) (2015) 3285-301.
[16] M. Perez-Andres, B. Paiva, W.G. Nieto, A. Caraux, A. Schmitz, J. Almeida, R.F. Vogt, Jr., G.E. Marti, A.C.
Rawstron, M.C. Van Zelm, J.J. Van Dongen, H.E. Johnsen, B. Klein, A. Orfao, Human peripheral blood B-cell
compartments: a crossroad in B-cell traffic, Cytometry B Clin Cytom 78 Suppl 1 (2010) S47-60.
[17] W. Cheuk, J.K. Chan, Subcellular localization of immunohistochemical signals: knowledge of the
ultrastructural or biologic features of the antigens helps predict the signal localization and proper
interpretation of immunostains, Int J Surg Pathol 12(3) (2004) 185-206.
[18] N. Dakka, H. Bellaoui, N. Bouzid, M. Khattab, Y. Bakri, A. Benjouad, CD10 AND CD34 expression in
childhood acute lymphoblastic leukemia in Morocco: clinical relevance and outcome, Pediatr Hematol
Oncol 26(4) (2009) 216-31.
[19] H. Yasuoka, M. Tsujimoto, K. Yoshidome, M. Nakahara, R. Kodama, T. Sanke, Y. Nakamura,
Cytoplasmic CXCR4 expression in breast cancer: induction by nitric oxide and correlation with lymph node
metastasis and poor prognosis, BMC Cancer 8 (2008) 340.
[20] J.J. Ma Y, Jhang J, CXCR4 expression as a predictive biomarker for esophageal cancer Int J Clin Exp
Med 11(9) (2016) 22055-22061
[21] L.C. Ding Y, Yang Q, Chen C, Cheng Z, The Significance of CXCR4 Expression for the Prediction of Lymph
Node Metastasis in Breast Cancer Patients, Chin J Clin Oncol 5 (2008) 83-86.
[22] N. Cabioglu, M.S. Yazici, B. Arun, K.R. Broglio, G.N. Hortobagyi, J.E. Price, A. Sahin, CCR7 and CXCR4
as novel biomarkers predicting axillary lymph node metastasis in T1 breast cancer, Clin Cancer Res 11(16)
(2005) 5686-93.
[23] F. Andre, W. Xia, R. Conforti, Y. Wei, T. Boulet, G. Tomasic, M. Spielmann, M. Zoubir, N. Berrada, R.
Arriagada, G.N. Hortobagyi, M.C. Hung, L. Pusztai, S. Delaloge, S. Michiels, M. Cristofanilli, CXCR4
expression in early breast cancer and risk of distant recurrence, Oncologist 14(12) (2009) 1182-8.
[24] W. Tang, X. Wang, Y. Chen, J. Zhang, Y. Chen, Z. Lin, CXCL12 and CXCR4 as predictive biomarkers of
glioma recurrence pattern after total resection, Pathol Biol (Paris) 63(4-5) (2015) 190-8.
[25] A. Krieg, J.C. Riemer, L.A. Telan, H.E. Gabbert, W.T. Knoefel, CXCR4--A Prognostic and
Clinicopathological Biomarker for Pancreatic Ductal Adenocarcinoma: A Meta-Analysis, PLoS One 10(6)
(2015) e0130192.
[26] C. Tomoda, R. Kushima, E. Takeuti, K. Mukaisho, T. Hattori, H. Kitano, CD10 expression is useful in the
diagnosis of follicular carcinoma and follicular variant of papillary thyroid carcinoma, Thyroid 13(3) (2003)
291-5.
[27] G.L. Moldovan, B. Pfander, S. Jentsch, PCNA, the maestro of the replication fork, Cell 129(4) (2007)
665-79.
[28] S.S. Citterio S, Levi M, Colombo BM, Sparvoli E, PCNA and total nuclear protein content as markers of
cell proliferation in pea tissue Journal of Cell Science 102 (1992) 71-78
[29] T.E. Shaiken, A.R. Opekun, Dissecting the cell to nucleus, perinucleus and cytosol, Sci Rep 4 (2014)
4923.

[30] R. Bransteitter, P. Pham, M.D. Scharff, M.F. Goodman, Activation-induced cytidine deaminase
deaminates deoxycytidine on single-stranded DNA but requires the action of RNase, Proc Natl Acad Sci U
S A 100(7) (2003) 4102-7.
[31] H. Arakawa, H. Saribasak, J.M. Buerstedde, Activation-induced cytidine deaminase initiates
immunoglobulin gene conversion and hypermutation by a common intermediate, PLoS Biol 2(7) (2004)
E179.
[32] J. Liu, E. Xiong, H. Zhu, H. Mori, S. Yasuda, K. Kinoshita, T. Tsubata, J.Y. Wang, Efficient Induction of Ig
Gene Hypermutation in Ex Vivo-Activated Primary B Cells, J Immunol 199(9) (2017) 3023-3030.
[33] Y. Hu, I. Ericsson, B. Doseth, N.B. Liabakk, H.E. Krokan, B. Kavli, Activation-induced cytidine deaminase
(AID) is localized to subnuclear domains enriched in splicing factors, Exp Cell Res 322(1) (2014) 178-92.
[34] S. Ito, H. Nagaoka, R. Shinkura, N. Begum, M. Muramatsu, M. Nakata, T. Honjo, Activation-induced
cytidine deaminase shuttles between nucleus and cytoplasm like apolipoprotein B mRNA editing catalytic
polypeptide 1, Proc Natl Acad Sci U S A 101(7) (2004) 1975-80.
[35] L.A. Bentolila, S. Olson, A. Marshall, F. Rougeon, C.J. Paige, N. Doyen, G.E. Wu, Extensive junctional
diversity in Ig light chain genes from early B cell progenitors of mu MT mice, J Immunol 162(4) (1999)
2123-8.
[36] C.L. Benedict, S. Gilfillan, J.F. Kearney, The long isoform of terminal deoxynucleotidyl transferase
enters the nucleus and, rather than catalyzing nontemplated nucleotide addition, modulates the catalytic
activity of the short isoform, J Exp Med 193(1) (2001) 89-99.
[37] M.D. Huber, L. Gerace, The size-wise nucleus: nuclear volume control in eukaryotes, J Cell Biol 179(4)
(2007) 583-4.

463 **Figures with legend**

A

B

467 **Figure 1**

468 **Characterization of exosomes by Nanoparticle Tracking Analysis (NTA).**

469 Purified exosomes were passed through NTA system. **A.** Graph shows the heterogeneity

470 of diameter size of the exosomes particle. **B.** Shows representative image of exosomes.

Figure 2

Amplification of cytoplasmic CD10 and CXCR4 from Cell-cDNA and Exo-cDNA.

Agarose gel elecprohporesis were carried out after 1st and 2nd round of PCR amplification as shown in upper and lower panel respectively. **A.** CD10 mRNA expression in Cell-cDNA and Exo-cDNA in indicated cell lines (JM1, SUP-B15, and CL-01), upper panel (first round of PCR), lower panel (seocnd round of PCR). **B.** CXCR4 mRNA expression in Cell-cDNA and Exo-cDNA in indicated cell lines (JM1, SUP-B15, and CL-01), upper panel (first round of PCR), lower panel (seocnd round of PCR). β-actin housekeeping gene as quality control.

Figure 3

Amplification of nuclear PCNA and CREB-BP from Cell-cDNA and Exo-cDNA.

Agarose gel elecprohporesis were carried out after 1st and 2nd round of PCR

amplification as shown in upper and lower panel respectively. **A.** PCNA mRNA expression

in Cell-cDNA and Exo-cDNA in indicated cell lines (JM1, SUP-B15, and CL-01), upper

panel (first round of PCR), lower panel (seocnd round of PCR). **B.** CREB-BP mRNA

expression in Cell-cDNA and Exo-cDNA in indicated cell lines (JM1, SUP-B15, and CL-

01), upper panel (first round of PCR), lower panel (seocnd round of PCR). β-actin

housekeeping gene as quality control.

Figure 4

Amplification of nuclear enzymes AID and TdT from Cell-cDNA and Exo-cDNA.

Execution of agarose gel elecprohporesis was carried out after 1st and 2nd rounds of

PCR amplification as shown in upper and lower panel respectively. **A.** AID mRNA

expression in Cell-cDNA and Exo-cDNA in indicated cell lines (JM1, SUP-B15, and CL-

01), upper panel (first round of PCR), lower panel (seocnd round of PCR). **B.** TdT mRNA

expression in Cell-cDNA and Exo-cDNA in indicated cell lines (JM1, SUP-B15, and CL-

01). Represntative data is seocnd round of PCR product. β-actin as housekeeping gene.

Table 1: Details of human primers

Gene	GenBank	Primer sequences	Annealing Temp	Product Size (bp)
CXCR4	AY242129.1	For: 5'-AATGAGGCCACAACAAACATCACA-3' Rev: 5'-ATCCAGACGCCAACATAGAC-3'	58 °C	464
CD10	NM_000902.3	For: 5'-CTGTGACAATGATCGCACTCTATG-3' Rev: 5'-GATTCCAGTGCATTCATAGTAATC-3'	60 °C	587
PCNA	NM_182649.1	For: 5'-TTTCTAGGTCTCAGCCGGTC-3' Rev: 5'-GCAAATTCACCAGAAGGCAT-3'	55 °C	557
CREB-BP	NM_004380.2	For: 5'- CCAGTGACAAGCGAAACCAA-3' Rev: 5'- GGCTGGTTCATGTAGGGGAG-3'	56 °C	181
AID	AB040431.1	For: 5'-AGGCAAGAAGACACTCTGGACACC-3' Rev: 5'-GTGACATTCCTGGAAGTTGC-3'	58 °C	645
TdT	AB046378.1	For: 5'-GCCGTCAGTGTGCTGGTTAAAGAGG- 3' Rev: 5'-TCTGCTTTGAGGAATATCCTCTTGG-3'	60 °C	535
β-actin	NM_001101.4	For: 5'-GTCCTCTCCCAAGTCCACACA-3' Rev: 5'-CTGGTCTCAAGTCAGTGTACAGGTAA-3'	60 °C	240

Table 2: Total number of cytoplasmic and nuclear coding genes

cDNA type	Cell types	Cytoplasmic protein coding genes					Nuclear protein coding genes				
		CD10	CD34	Actin	CXCR4	HLA-DR	p53	PCNA	TdT	CREB-BP	AID
Cell-cDNA	JM1	+	+	+	+	+	+	+	+	+	+
Cell-cDNA	SUP-B156	+	+	+	+	+	+	+	+	+	+
Cell-cDNA	CL-01	+	-	+	+	+	+	+	-	+	+
Exo-cDNA	JM1	+	+	+	+	+	-	-	-	-	-
Exo-cDNA	SUP-B156	+	+	+	+	+	-	-	-	-	-
Exo-cDNA	CL-01	+	-	+	+	+	-	-	-	-	-