Single-nuclei transcriptomics of schizophrenia prefrontal cortex primarily implicates neuronal subtypes

Benjamin C. Reiner, PhD[1,4,*], Richard C. Crist, PhD[1,4], Lauren M. Stein, PhD[1], Andrew E. Weller, MD[1], Glenn A. Doyle, PhD[1], Gabriella Arauco-Shapiro, MS[1], Gustavo Turecki, MD, PhD[2] ,Thomas N. Ferraro, PhD[3], Matthew R. Hayes, PhD[1], and Wade H. Berrettini, MD, PhD[1]

[1] Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania. [2] McGill Group for Suicide Studies, Douglas Mental Health University Institute, McGill University, Montreal, QC H4H 1R3, Canada. [3] Department of Biomedical Sciences, Cooper Medical School of Rowan University, Camden, NJ. [4] These authors contributed equally.

Running title: Single nuclei RNAseq in the Schizophrenia cortex

*Corresponding author

Benjamin C. Reiner, PhD

125 S. 31st St., room 2213

Philadelphia, PA 19104

Phone: (215)-573-0278

Email: bcreiner@pennmedicine.upenn.edu

Abstract (limit 150-250) - 150

Transcriptomic studies of bulk neural tissue homogenates from persons with schizophrenia and controls have identified differentially expressed genes in multiple brain regions. However, the brain's heterogeneous nature prevents identification of relevant cell types. This study analyzed single-nuclei transcriptomics of ~275,000 nuclei from frozen human postmortem dorsolateral prefrontal cortex samples from males with schizophrenia (n = 12) and controls (n = 14). 4,766 differential expression events were identified in 2,994 unique genes in 16 of 20 transcriptomically-distinct cell populations. ~96% of differentially expressed genes occurred in five neuronal cell types, and differentially expressed genes were enriched for genes associated with schizophrenia and bipolar GWAS loci. Downstream analyses identified cluster-specific enriched gene ontologies, KEGG pathways, and canonical pathways. Additionally, microRNAs and transcription factors with overrepresented neuronal cell type-specific targets were identified. These results expand our knowledge of disrupted gene expression in specific cell types and permit new insight into the pathophysiology of schizophrenia.

Introduction (limit 1500; article excluding abstract 3500) - 478

Schizophrenia is a chronic psychotic illness affecting ~1% of the population worldwide. Transcriptomic studies utilizing bulk homogenates of frozen human postmortem brain tissue from persons with schizophrenia and controls have identified differentially expressed genes (DEGs) in the amygdala[1], hippocampus[2, 3], superior temporal gyrus[4], anterior cingulate cortex[5, 6], and dorsolateral prefrontal cortex (dlPFC)[7, 8], with the largest study identifying ~4,800 DEGs associated with schizophrenia in the dlPFC[9]. However, the heterogenous cellular composition of the bulk homogenates prevents identification of the specific cell types in which relevant genes are differentially regulated and expressed. Two studies used laser capture microdissection followed with transcriptomic analysis by microarray to elucidate the effect of schizophrenia on the transcriptome of individual neural cell types. Examination of layer 3 and 5 pyramidal neurons

in the dlPFC identified ~1,400 DEGs[10] in the context of schizophrenia, whereas a study of

parvalbumin positive (PVALB+) interneurons in the dlPFC identified ~900 DEGs[11]. A substantial

portion of the DEGs identified in these studies were not detected in previous examinations of

bulk homogenates of the same brain regions, suggesting that examination of transcriptomic

changes associated with schizophrenia at the level of neural cellular subpopulations is

necessary to fully appreciate the neuropathophysiology of the disorder[10, 11].

Laser capture microdissection studies of human postmortem brain tissue are limited by

their ability to examine a small number of cell types in a targeted fashion, relatively low

throughput, and the pooling of cells, which loses the variability of the transcriptome between

cells and may collapse transcriptomically-distinct subpopulations. Recent advances in single

nuclei RNA sequencing (snRNAseq) allow for simultaneous transcriptomic profiling of

thousands of nuclei, across all neural cell types in frozen human postmortem brain homogenate,

with simultaneous indexing of transcripts at the sample, nucleus, and individual transcript level

(unique molecular identifier, UMI). The utility of this approach for human postmortem study is

supported by evidence suggesting that single cells and their nuclei have similar transcriptomes,

with ~98% of transcripts having the same relative levels[12]. snRNAseq has identified cell type-

specific transcriptomic changes in human postmortem brain samples from Alzheimer's

disease[13], autism[14], multiple sclerosis[15], and major depressive disorder[16].

In this study, we performed snRNAseq of ~275,000 nuclei from dlPFC of individuals with

schizophrenia (n = 12) and controls (n = 14). We chose to examine the dlPFC due to the

evidence of dlPFC dysfunction in schizophrenia[17]. We identified 4,766 DEGs in 16 of 20

transcriptomically-distinct cell populations. ~96% of the DEGs occurred in five neuronal cell

types. The DEGs were enriched for genes associated with schizophrenia GWAS loci and

overrepresented in gene ontologies and KEGG pathways previously associated with the

pathophysiology of schizophrenia. Canonical pathway analysis identified cluster-specific

alterations in metabolic and cell signaling pathways, and microRNA, transcription factor, and

upstream regulator analyses identified putative regulators of cluster-specific DEG. Taken

together, the results of this study help elucidate the cell type-specific transcriptomic and

neurobiological changes that underlie schizophrenia.

Materials and Methods - 1036

Brain Samples

This study was approved by the University of Pennsylvania Institutional Review Board.

Fresh frozen postmortem dlPFC tissue from male individuals with schizophrenia (n = 14) and

controls (n =14) were obtained from the Douglas-Bell Canada Brain Bank at McGill University,

the Human Brain and Spinal Fluid Resource Center at UCLA and the New South Wales Brain

Tissue Resource Center. Schizophrenia cases were individuals who were clinically diagnosed

with schizophrenia using DSM-IV criteria and controls were individuals without history of

psychiatric disease who died of non-central nervous system-related reasons. All reported age,

sex, ethnicity, postmortem interval, and prefrontal cortex pH data are based on associated

medical records (Supplementary Table 1). Gray matter samples from the dlPFC were dissected

by trained neuroanatomists at their respective brain banks.

10x Library Preparation, Sequencing, and Quality Control

Nuclei were isolated from frozen postmortem dlPFC (~30mg) using a modified version of

a previously described protocol[16] (see Supplemental Methods). Microfluidics capture and

sequencing library preparation was performed with the 10x Genomics Chromium Single Cell 3'

GEM, Library and Gel Bead Kit v3.0 at the Children's Hospital of Philadelphia Center for Applied

Genomics per manufacturer's instructions. To achieve a target capture of ~10,000 nuclei per

sample, ~20,000 nuclei per sample were loaded. Libraries were sequenced in pools of eight on

Illumina NovaSeq 6000 S2 flow cells. Pools contained schizophrenia and control samples, to

minimize any batch effects. CellRanger version 3.1 was used to align reads to the hg38 pre-

mRNA transcriptome. Filtered read count matrices for all subjects and nuclei were merged into

a single Seurat object for subsequent quality control and clustering using Seurat version 3.1.

For initial quality control assessment, the distributions of the numbers of genes and UMIs were

determined. Nuclei with the lowest 1% of genes (< 470 genes) were removed, as they were

unlikely to be informative in downstream analyses. Likewise, nuclei in the top 1% of UMI count

(UMI > 60,335) were removed to reduce the presence of multiplets in downstream analyses.

Finally, nuclei with >10% of reads from mitochondrial genes were excluded and mitochondrial

transcripts were removed from the dataset[18].

Calculation of PCs, Clustering, and Cell Type Annotation

Transcript counts were normalized to 10,000 counts per subject and scaled. Variably

expressed genes were identified with the FindVariableFeatures function in Seurat using the

mean.var.plot selection method and analyzing only genes with mean scaled expression

between 0.003 and 2. These parameters identified 2,486 highly variable genes, which were

used to generate principal components (PCs). Clustering was performed in Seurat using the first

50 PCs. Initial clustering was performed at a resolution of 0.25. Two schizophrenia samples did

not cluster with the other 26 samples and were removed as outliers. The dataset was

reclustered and two cell populations with low mean UMIs were removed. Six clusters with

>90% of nuclei coming from ≤2 subjects were also removed, with remaining clusters having

<30% of nuclei coming from ≤2 subjects. Major cell types and neuronal subtypes were identified

using known cell type markers and methods described in Nagy et al.[19] (see Supplemental

Methods). Two clusters with mixed cell type markers were removed for a final total of 20

clusters.

<u>Differential Gene Expression Analysis</u>

Count data were normalized to one million reads, extracted from Seurat, and converted

to log2 counts per million (cpm). Metadata and cpm were merged to form a SingleCellAssay

object for each cluster. Genes expressed in <20% of the nuclei in a cluster were excluded from

downstream analyses. Differential expression analysis between cases and controls was

performed using the *MAST* R package[20] by fitting the following linear mixed model:

m <- zlm(~casestatus + gdr + age + sex + batch + (1|subject), sca, parallel = TRUE, method =

"glmer", ebayes = FALSE, silent=TRUE)

Case status, age, the capture and sequencing batch, and the number of genes detected in each

nucleus (gdr) were included as fixed effects. Subject was included as a random effect to

account for correlations between the nuclei coming from a single person. To optimize the

random and fixed effects coefficients in the penalized iteratively reweighted least squares step,

the integer scalar in the *lme4* R package was set equal to zero, as previously described[14]

(https://github.com/DmitryVel/KriegsteinLab/blob/master/snRNAseq_DGE.R). Likelihood Ratio

Test was performed in *MAST* to test for differences between the model with and without

schizophrenia case status, identifying gene expression differences associated with

schizophrenia. DEGs were defined as those that were a) statistically significant after multiple

testing correction with false discovery rate (FDR) = 0.1 and b) had at least a 10% difference in

expression between case and controls ($\log_2$ fold change ≥0.14).

<u>Overrepresentation of GWAS loci</u>

To determine if schizophrenia, bipolar disorder, and Alzheimer's disease GWAS loci

were overrepresented in cluster-specific DEGs, MAGMA [21] was used to identify significant

genes using GWAS summary statistics. Cluster-specific overrepresentation of GWAS loci was

determined by performing a hypergeometric test using the overlapping list of genes that were

checked for differential expression in each cluster and were significant by MAGMA analysis.

<u>Enrichment Analysis with FUMA</u>

Overrepresentation of brain expressed genes was determined by comparing cluster-

specific DEGs to brain region-specific transcriptome data from the Genotype-Tissue Expression

(GTEx) project (v8)[22] using FUMA[22, 23]. Similarly, overrepresentation of brain expressed genes

throughout the human life span was determined by comparing cluster-specific DEGs to the

BrainSpan transcriptomics data [24]. Identification of gene ontologies, KEGG pathways,

transcription factor targets, and microRNA targets that were significantly overrepresented in

cluster-specific DEGs was analyzed using FUMA and data from the Molecular and Signatures

Database (MsigDB) v7.0 [25]. For all analyses, all genes, except the MHC region, from Ensembl

v92 were used for the gene background and all results were corrected for multiple testing using

a Bonferroni correction (α=0.05).

<u>Canonical pathways and upstream regulators</u>

Canonical pathway and upstream regulator analysis was conducted using Ingenuity

Pathway Analysis[26]. DEGs from each of the primary neuronal clusters were analyzed

independently using the Core Analysis function and a list of DEG Ensembl IDs, *MAST* raw p-

values, FDR corrected p-values (i.e. q-values), and $\log_2$ fold change. The analyses from the five

clusters were compared using the Compare Analyses function. P-values were calculated using

a right-tailed Fisher's exact test and corrected for multiple testing using a Benjamini-Hochberg

correction ($\alpha=0.1$).

Statistics

Statistical calculations were performed with SPSS (v24).

Results – 1100

Single nuclei RNA sequencing and identification of cell types

To better understand the effects of schizophrenia on the neural transcriptome, we

utilized snRNAseq to profile dlPFC nuclei from frozen human postmortem brain samples from

12 individuals with schizophrenia and 14 controls (Supplementary Table 1). Schizophrenia and

control groups did not differ in mean age, postmortem interval, or pH (Supplementary Table 1

and Supplementary Fig. 1). snRNAseq was performed using the 10x Genomics platform and

sequenced to an average depth of ~492 million reads per sample. We initially identified 361,681

nuclei, with average medians of ~2,987 genes and ~6,886 UMI per nucleus (Supplementary

Table 2). The number of sequencing reads per sample, number of nuclei per sample, average

sequencing reads per nuclei per sample, median genes per nuclei per sample, and median UMI

per nuclei per sample were consistent between schizophrenia and control groups

(Supplementary Fig. 2). RIN value was unavailable for all samples, so RNA quality was

evaluated by examining sequencing-derived surrogates of RNA quality from CellRanger. RNA

quality did not differ between schizophrenia and control groups, with no difference in the fraction

of reads mapped confidently to the genome, intergenic regions, exonic regions, intragenic

regions, or the transcriptome (all Mann-Whitney U p-values >0.28; Supplementary Table 3).

After quality control measures were applied, we identified 273,050 high confidence nuclei:

145,120 from control samples and 127,930 from schizophrenia samples. These nuclei were

used for all further analyses.

To identify nuclei from transcriptomically-distinct cellular populations, we used the

snRNAseq counts for nuclei from all individuals to perform unbiased clustering using Seurat[27],

and 20 cellular clusters were identified (Fig. 1a and Supplementary Fig. 3). Clusters were

annotated by expression of known cellular subtype markers and all major neural cell types were

identified (Fig. 1a-b). The schizophrenia and control groups were consistent for the number of

nuclei from major cell types (Fig. 1c) and the number of nuclei per cluster (Fig. 1d & 1e).

Excitatory neuron (ExNeuro) and inhibitory neuron (InNeuro) clusters were further annotated

using known markers for cortical layers and defined subpopulations (Fig. 1b and Supplemental

Fig. 4a-b).

Differential expression

Cell type-specific changes in nuclear transcript levels were compared between the

schizophrenia and control groups using a hurdle model, a linear mixed model, in MAST[20]. 4,766

differential expression events were identified in 2,994 unique DEGs (q-value <0.1; expression

change ≥10%; Supplementary Table 4). DEGs were detected in 16 of 20 cellular clusters, with

35.3% of differential expression events being upregulation and 64.7% downregulation (Fig. 2a).

The number of DEGs per cluster was unrelated to the number of nuclei per cluster (r(18)=0.16,

p=0.50, Pearson's Correlation; Supplementary Fig. 5). Of the unique DEGs, 96.2% occurred in

five neuronal cell types (primary neuronal cell types), including four clusters of ExNeuro (cluster

10 layer 5 HTR2C+ ExNeuro, cluster 12 layer 5 ExNeuro, cluster 13 layer 4/5 ExNeuro, and

cluster 14 layer 2/3 ExNeuro) and a PVALB+ InNeuro cluster (cluster 3; Fig. 2b).

<u>DEGs associated with GWAS loci</u>

MAGMA was used to determine if cell type-specific DEGs were enriched for genes

identified in GWAS of schizophrenia[28] or bipolar disorder[29], because previous studies identified

substantial heritability overlap[30]. Schizophrenia GWAS loci-associated genes were

overrepresented in four of five primary neuronal cell types, with cluster 12 also showing

overrepresentation of bipolar disorder GWAS genes (Fig. 2c and Supplementary Table 5).

Supporting the specificity of these observations, no enrichment was detected for Alzheimer's

disease GWAS genes (Fig. 2c and Supplementary Table 5). Similarly, cell type-specific DEGs

were enriched for GWAS Catalog genes for schizophrenia in cluster 3 PVALB+ InNeuro and

cluster 13 layer 4/5 ExNeuro and for bipolar disorder in cluster 10 layer 5 HTR2C+, cluster 12

layer 5, and cluster 13 layer 4/5, and cluster 14 layer 2/3 ExNeuro (Supplementary Table 6).

<u>Overrepresentation of brain expressed and developmental genes</u>

Comparison of cluster-specific DEGs to brain region-specific transcriptomic data from

GTEx shows substantial overrepresentation of up and down regulated DEGs throughout the

brain for the primary neuronal clusters (Fig. 2d and Supplementary Table 7). Intriguingly, when

compared to BrainSpan transcriptomic data, overrepresentation of cluster-specific DEGs was

concentrated at eight to thirteen post-conception weeks and two years of age (Fig. 2e and

Supplementary Table 8), suggesting a putative neurodevelopmental relevance.

<u>Gene Ontologies and KEGG pathways</u>

To understand the neurobiological consequences of the DEGs in the primary neuronal

clusters, FUMA was used to identify gene ontologies (GO; Supplementary Tables 9) and KEGG

pathways (Supplementary Tables 10) in which DEGs from each cell cluster were over- or under-

represented. GO and KEGG analysis identified cluster-specific representation changes of DEGs

in ontologies and pathways previously associated with the pathophysiology of schizophrenia,

including those related to mitochondrial function. These results suggest that cluster-specific

DEGs may underlie some cell type-specific neurobiological changes associated with

schizophrenia neuropathophysiology.

<u>Canonical Pathways</u>

To identify metabolic and cell signaling pathways that are likely to be altered in the

primary neuronal clusters, canonical pathway analysis was performed with Ingenuity Pathway

Analysis (IPA) using the cluster-specific DEGs. Shared and unique canonical pathways were

identified for the primary neuronal clusters (Supplementary table 11). The top five shared

canonical pathways for the primary neuronal clusters, as determined by p-value, are presented

in Fig. 3a. While the primary neuronal clusters shared predicted disruptions of pathways, the

DEGs underlying the effect for each cellular cluster were a combination of shared and unique

DEGs (Fig. 3b and Supplementary table 11). These results suggest that a combination of

shared and cluster-specific transcriptome alterations drive overall pathway dysfunction.

<u>Upstream transcription factors, microRNAs, and regulatory networks</u>

To identify putative mechanisms for the alterations in gene expression observed in the

primary neuronal clusters, we assessed overrepresentation of transcription factor

(Supplementary table 12) and microRNA (Supplementary table 13) targets among cluster-

specific DEGs. Unique transcription factor and microRNA targets were identified for each of the

primary neuronal clusters, suggesting potential cell type-specific mechanism for transcriptional

dysregulation. Additionally, a subset of transcription factors and microRNAs had significantly

overrepresented targets across the primary neuronal clusters, suggesting a potential shared

etiology. Unique and shared putative upstream regulatory networks were also identified for all

clusters (Supplementary table 14). The top five shared upstream network master regulators for

the primary neuronal clusters, as determined by p-value, are presented in Fig. 4a. These

predictions were driven by shared and cluster-specific DEGs (Fig. 4b). These results are

supported by the direct detection of predicted alterations for a portion of the upstream

regulators. For example, *DDX5* function was predicted to be inhibited in four of five primary

neuronal clusters (Fig. 4a) and *DDX5* was found to be significantly downregulated in both

cluster 3 PVALB+ InNeuro and cluster 13 layer 4/5 ExNeuro (Supplementary table 14).

Discussion – 885

This study examined the transcriptome of ~275,000 single nuclei from the dlPFC of

persons with schizophrenia and controls. The median UMI and gene counts per nucleus were

approximately double those of three recent human postmortem cortical snRNAseq studies,

although the ratio of median UMI count to median gene count was similar[14-16]. Differences

between versions 2.0 and 3.0 of the 10x Genomics gene expression assay explain at least a

portion of our increased gene and UMI yield. The identification of 20 transcriptomically distinct

cellular clusters is consistent with other snRNAseq studies of human postmortem cortex[13, 14, 16]

and the number of clusters per major cell type closely approximates those of a recent studies of

dlPFC, including identification of a single *HTR2C*+ cluster of pyramidal ExNeuro and two

PVALB+ clusters of InNeuro[16]. 4,766 differential expression events were detected in 16 of 20

cellular clusters, with ~96% of DEGs occurring in five neuronal cell types. Prior evidence

suggests that GWAS loci and gene sets associated with schizophrenia are primarily expressed

in a limited subset of neurons, including PVALB+ InNeuro and glutamatergic pyramidal

neurons[31]. Similarly, GWAS loci-related genes were overrepresented in the DEGs of cluster 3

PVALB+ InNeuro and the cluster 10 layer 5 *HTR2C*+, cluster 12 layer 5, and cluster 14 layer 2/3

ExNeuro. Cluster 12 layer 5 ExNeuro also had an overrepresentation of bipolar disorder GWAS

loci, fitting prior knowledge that schizophrenia and bipolar disorder share genetic risk loci. These

results support the hypothesis that common genetic variants associated with schizophrenia are

relevant in specific sets of neuronal cell types and schizophrenia-related transcriptomic

alterations are primarily limited to these cells. Several reports provide fairly consistent evidence

of presynaptic marker decreases for frontal cortical fast-spiking parvalbumin +- GABAergic

interneurons, coupled with increased postsynaptic GABAA receptors, both of which may be

consistent with partial loss of GABAergic inhibition of glutamatergic pyramidal neurons[32, 33]. The

large number of DEGs in both cell types provides support for dysfunction of a frontal cortex

GABAergic-glutamatergic circuit. No DEG occurred in more than five cellular clusters and that

no DEG was present in all the clusters of any multi-clustered major cell type underscores the

importance of utilizing single nuclei/single cell approaches to neural transcriptomics.

The substantial overrepresentation of cluster-specific DEGs during critical

neurodevelopmental timepoints, 8 to 13 post-conception weeks and 2 years of age (Fig. 2b)

supports hypotheses about schizophrenia as a neurodevelopmental disorder [34-36]. Our analyses

identified a substantial number of GO terms and KEGG and canonical pathways related to

energy metabolism and oxidative stress in the primary neuronal clusters and prior works have

hypothesized the prenatal and early developmental dysregulation of oxidative stress may play a

role in the development of schizophrenia, particularly in PVALB+ neurons (reviewed[37]). Taken

together, these data suggest potential windows for PVALB+ InNeuro oxidative stress targeted

interventions.

Human postmortem studies are identifying increasing numbers of shared and brain

region-specific differentially expressed microRNAs that are associated with schizophrenia[38],

including a global increase in microRNA levels[39]. While the experimental approach of this study

was unable to directly detect alterations in microRNA expression, complementary approaches to

identifying cell type-specific microRNA target overrepresentation yield overlapping predictions of

microRNAs known to regulate brain function. For example, miR-424 (aka miR-322) targets were

predicted to be overrepresented in the cluster 3 PVALB+ InNeuro and cluster 10 layer 5

HTR2C+ and cluster 13 layer 4/5 ExNeuro. miR-424 is known to regulate BDNF expression[40]

and literature evidence supports a role for alterations in BDNF expression in schizophrenia

pathogenesis[41]. Taken together, these data suggest that the neuronal cell type-specific

microRNA identified in this study may warrant further investigation.

Several limitations of this study must be noted. First, the relatively small number of

postmortem samples analyzed increases the possibility that the subjects are not representative

of the broader populations. Replication in a larger sample, including female samples, will be

essential for these results. Second, schizophrenia patients frequently have comorbidities (e.g.

smoking, obesity) that are less common in control individuals, presenting analytical confounds.

Similarly, schizophrenia patients usually have a history of chronic antipsychotic treatment,

whereas controls do not. Thus, it is impossible to know at present whether any of the identified

DEGs reflect causality or response to chronic pharmacotherapy. It may also be possible to

address this issue by studying postmortem brains of persons with schizophrenia who never had

antipsychotic treatment. However, at least in the United States, these patients are uncommon.

Third, transcriptome-based methods such as snRNAseq have the potential to miss relevant

genes that are regulated primarily at the level of translation or splicing, which may also help to

shape transcriptomic architecture and be relevant to schizophrenia pathology. Finally, this

project was limited to a single brain region from individuals over 18 years of age. Therefore,

spatial and temporal changes in gene expression occurring over the course of the disease

would not be identified in our analysis. The findings from this study of dlPFC cannot be

extrapolated to other areas of the brain, justifying the need for more comprehensive studies.

There is also substantial evidence for neurodevelopmental origins in schizophrenia[42, 43],

suggesting there are relevant transcriptomic differences between cases and controls before

adulthood. In summary, we have begun to characterize transcriptome alterations in

schizophrenia at the level of single neural cells and extension of this work may provide a new

basis for the development of effective treatment strategies.

<u>Availability of Data</u>

Raw sequencing data and sample annotations are available at NCBI GEO accession #

GSE158516.

Acknowledgements

BCR is supported, in part, by a 2017 NARSAD Young Investigator Grant (#26634) from

the Brain and Behavior Research Foundation as the Patrick A. Coffer Investigator, funding for

which was generously provided by Ronald and Kathy Chandonais. BCR and AEW were

supported during part of this work by T32MH014654 (PI is WHB). This work was funded by a

National Institutes of Health grant to WHB (R01MH109260). LMS was supported by NIDDK F32

DK118818. MRH was supported by NIDDK R01 DK115762. The authors thank Rachel Kember

for her assistance with R. The authors would also like to thank Jane Gaisinsky for her help with

tissue preparation. Additionally, the authors thank all the participants, and their families, who

donated their brain tissue to research, for making these studies possible.

Author Contributions

BCR conceptualized this study and performed experiments. BCR and RCC performed

bioinformatics and wrote the manuscript. GA contributed to the performance of experiments and

manuscript preparation. LMS, AEW, GAD, TNF and MRH contributed to data interpretation and

manuscript preparation. WHB provided oversight and contributed to data interpretation and

manuscript preparation.

Competing Interests Statement

MRH receives separate financial support from Boehringer Ingelheim and Eli Lily & Co

that was not used for or related to these studies.

References

1.   Chang X, Liu Y, Hahn CG, Gur RE, Sleiman PMA, Hakonarson H. RNA-seq analysis of amygdala
tissue reveals characteristic expression profiles in schizophrenia. *Transl Psychiatry* 2017; **7**(8)**:**
e1203.

2.   Hwang Y, Kim J, Shin JY, Kim JI, Seo JS, Webster MJ *et al.* Gene expression profiling by mRNA
sequencing reveals increased expression of immune/inflammation-related genes in the
hippocampus of individuals with schizophrenia. *Transl Psychiatry* 2013; **3:** e321.

3.   Kohen R, Dobra A, Tracy JH, Haugen E. Transcriptome profiling of human hippocampus dentate
gyrus granule cells in mental illness. *Transl Psychiatry* 2014; **4:** e366.

4.   Wu JQ, Wang X, Beveridge NJ, Tooney PA, Scott RJ, Carr VJ *et al.* Transcriptome sequencing
revealed significant alteration of cortical promoter usage and splicing in schizophrenia. *PloS one*
2012; **7**(4)**:** e36351.

5.   Zhao Z, Xu J, Chen J, Kim S, Reimers M, Bacanu SA *et al.* Transcriptome sequencing and genome-
wide association analyses reveal lysosomal function and actin cytoskeleton remodeling in
schizophrenia and bipolar disorder. *Molecular psychiatry* 2015; **20**(5)**:** 563-572.

6.   Hong S, Chen X, Jin L, Xiong M. Canonical correlation analysis for RNA-seq co-expression
networks. *Nucleic acids research* 2013; **41**(8)**:** e95.

7.   Fillman SG, Cloonan N, Catts VS, Miller LC, Wong J, McCrossin T *et al.* Increased inflammatory
markers identified in the dorsolateral prefrontal cortex of individuals with schizophrenia.
Molecular psychiatry 2013; **18**(2)**:** 206-214.

8.   Fromer M, Roussos P, Sieberts SK, Johnson JS, Kavanagh DH, Perumal TM *et al.* Gene expression
elucidates functional impact of polygenic risk for schizophrenia. *Nat Neurosci* 2016; **19**(11)**:**
1442-1453.

9.   Gandal MJ, Zhang P, Hadjimichael E, Walker RL, Chen C, Liu S *et al.* Transcriptome-wide isoform-
level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science* 2018; **362**(6420).

10.   Arion D, Huo Z, Enwright JF, Corradi JP, Tseng G, Lewis DA. Transcriptome Alterations in
Prefrontal Pyramidal Cells Distinguish Schizophrenia From Bipolar and Major Depressive
Disorders. *Biol Psychiatry* 2017; **82**(8)**:** 594-600.

11.   Enwright Iii JF, Huo Z, Arion D, Corradi JP, Tseng G, Lewis DA. Transcriptome alterations of
prefrontal cortical parvalbumin neurons in schizophrenia. *Molecular psychiatry* 2018; **23**(7)**:**
1606-1613.

12.   Grindberg RV, Yee-Greenbaum JL, McConnell MJ, Novotny M, O'Shaughnessy AL, Lambert GM *et*
al. RNA-sequencing from single nuclei. *Proceedings of the National Academy of Sciences of the*
United States of America 2013; **110**(49)**:** 19802-19807.

13.   Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ *et al.* Single-cell
444   transcriptomic analysis of Alzheimer's disease. *Nature* 2019; **570**(7761)**:** 332-337.

446   14.   Velmeshev D, Schirmer L, Jung D, Haeussler M, Perez Y, Mayer S *et al.* Single-cell genomics
447   identifies cell type-specific molecular changes in autism. *Science* 2019; **364**(6441)**:** 685-689.

449   15.   Schirmer L, Velmeshev D, Holmqvist S, Kaufmann M, Werneburg S, Jung D *et al.* Neuronal
450   vulnerability and multilineage diversity in multiple sclerosis. *Nature* 2019; **573**(7772)**:** 75-82.

452   16.   Nagy C, Maitra M, Tanti A, Suderman M, Theroux JF, Davoli MA *et al.* Single-nucleus
453   transcriptomics of the prefrontal cortex in major depressive disorder implicates oligodendrocyte
454   precursor cells and excitatory neurons. *Nat Neurosci* 2020; **23**(6)**:** 771-781.

456   17.   Hoftman GD, Datta D, Lewis DA. Layer 3 Excitatory and Inhibitory Circuitry in the Prefrontal
457   Cortex: Developmental Trajectories and Alterations in Schizophrenia. *Biol Psychiatry* 2017;
458   **81**(10)**:** 862-873.

460   18.   Osorio D, Cai JJ. Systematic determination of the mitochondrial proportion in human and mice
461   tissues for single-cell RNA sequencing data quality control. *Bioinformatics* 2020.

463   19.   Nagy C, Maitra M, Tanti A, Suderman M, Theroux JF, Davoli MA *et al.* Single-nucleus
464   transcriptomics of the prefrontal cortex in major depressive disorder implicates oligodendrocyte
465   precursor cells and excitatory neurons. *Nat Neurosci* 2020.

467   20.   Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK *et al.* MAST: a flexible statistical
468   framework for assessing transcriptional changes and characterizing heterogeneity in single-cell
469   RNA sequencing data. *Genome biology* 2015; **16:** 278.

471   21.   de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS
472   data. *PLoS Comput Biol* 2015; **11**(4)**:** e1004219.

474   22.   Consortium GT, Laboratory DA, Coordinating Center -Analysis Working G, Statistical Methods
475   groups-Analysis Working G, Enhancing Gg, Fund NIHC *et al.* Genetic effects on gene expression
476   across human tissues. *Nature* 2017; **550**(7675)**:** 204-213.

478   23.   Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of
479   genetic associations with FUMA. *Nat Commun* 2017; **8**(1)**:** 1826.

481   24.   Kang HJ, Kawasawa YI, Cheng F, Zhu Y, Xu X, Li M *et al.* Spatio-temporal transcriptome of the
482   human brain. *Nature* 2011; **478**(7370)**:** 483-489.

484   25.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP. Molecular
485   signatures database (MSigDB) 3.0. *Bioinformatics* 2011; **27**(12)**:** 1739-1740.

487   26.   Kramer A, Green J, Pollard J, Jr., Tugendreich S. Causal analysis approaches in Ingenuity Pathway
488   Analysis. *Bioinformatics* 2014; **30**(4)**:** 523-530.

490   27.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd *et al.* Comprehensive
491   Integration of Single-Cell Data. *Cell* 2019; **177**(7)**:** 1888-1902 e1821.

493   28.   Pardinas AF, Holmans P, Pocklington AJ, Escott-Price V, Ripke S, Carrera N *et al.* Common
494   schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong
495   background selection. *Nature genetics* 2018; **50**(3)**:** 381-389.

497   29.   Stahl EA, Breen G, Forstner AJ, McQuillin A, Ripke S, Trubetskoy V *et al.* Genome-wide
498   association study identifies 30 loci associated with bipolar disorder. *Nature genetics* 2019; **51**(5)**:**
499   793-803.

501   30.   Bipolar D, Schizophrenia Working Group of the Psychiatric Genomics Consortium. Electronic
502   address drve, Bipolar D, Schizophrenia Working Group of the Psychiatric Genomics C. Genomic
503   Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell* 2018; **173**(7)**:**
504   1705-1715 e1716.

506   31.   Skene NG, Bryois J, Bakken TE, Breen G, Crowley JJ, Gaspar HA *et al.* Genetic identification of
507   brain cell types underlying schizophrenia. *Nature genetics* 2018; **50**(6)**:** 825-833.

509   32.   Benes FM. The GABA system in schizophrenia: cells, molecules and microcircuitry. *Schizophr Res*
510   2015; **167**(1-3)**:** 1-3.

512   33.   Dienel SJ, Lewis DA. Alterations in cortical interneurons and cognitive function in schizophrenia.
513   *Neurobiol Dis* 2019; **131:** 104208.

515   34.   Murray RM, Lewis SW. Is schizophrenia a neurodevelopmental disorder? *Br Med J (Clin Res Ed)*
516   1987; **295**(6600)**:** 681-682.

518   35.   Weinberger DR. The neurodevelopmental origins of schizophrenia in the penumbra of genomic
519   medicine. *World Psychiatry* 2017; **16**(3)**:** 225-226.

521   36.   Thapar A, Riglin L. The importance of a developmental perspective in Psychiatry: what do recent
522   genetic-epidemiological findings show? *Molecular psychiatry* 2020; **25**(8)**:** 1631-1639.

524   37.   Behrens MM, Sejnowski TJ. Does schizophrenia arise from oxidative dysregulation of
525   parvalbumin-interneurons in the developing cortex? *Neuropharmacology* 2009; **57**(3)**:** 193-200.

526
527   38.   Beveridge NJ, Cairns MJ. MicroRNA dysregulation in schizophrenia. *Neurobiol Dis* 2012; **46**(2)**:**
528   263-271.

529
530   39.   Beveridge NJ, Gardiner E, Carroll AP, Tooney PA, Cairns MJ. Schizophrenia is associated with an
531   increase in cortical microRNA biogenesis. *Molecular psychiatry* 2010; **15**(12)**:** 1176-1189.

532
533   40.   Eyileten C, Sharif L, Wicik Z, Jakubik D, Jarosz-Popek J, Soplinska A *et al.* The Relation of the
534   Brain-Derived Neurotrophic Factor with MicroRNAs in Neurodegenerative Diseases and Ischemic
535   Stroke. *Mol Neurobiol* 2021; **58**(1)**:** 329-347.

536
537   41.   Goren JL. Brain-derived neurotrophic factor and schizophrenia. *Ment Health Clin* 2016; **6**(6)**:**
538   285-288.

539
540   42.   Birnbaum R, Weinberger DR. Genetic insights into the neurodevelopmental origins of
541   schizophrenia. *Nature reviews Neuroscience* 2017; **18**(12)**:** 727-740.

542
543   43.   Rapoport JL, Giedd JN, Gogtay N. Neurodevelopmental model of schizophrenia: update 2012.
544   *Molecular psychiatry* 2012; **17**(12)**:** 1228-1238.

 Figure Legends

Fig. 1: snRNAseq and clustering. a The transcriptome profile of ~275,000 nuclei were utilized

for unbiased clustering and is presented as a uniform manifold approximation and projection

(UMAP) dimension reduction plot of all nuclei color coded by cluster. **b** Clusters were annotated

with genes known to be markers for major neural cell types. The size and color of dots is

proportional to the percentage of cells expressing the gene (Pct. Exp.) and the average

expression level of the gene (Avg. Exp.), respectively. The cluster numbers and colors are

matched to that of the UMAP. **c** The proportion of major cell types between the schizophrenia

and control groups. **d** The proportion of schizophrenia and control nuclei in each cluster. The

labels and numbers correspond to those of the UMAP and dot plot. **e** Cluster contribution by

individual sample. Colors correspond to those of the UMAP and dot plot. Abbreviations:

somatostatin (SST), parvalbumin (PVALB), vasoactive intestinal peptide (VIP), synaptic vesicle

glycoprotein 2C (SV2C), reelin (RELN), nuclear receptor subfamily 4 group A member 2

(NR4A2), sulfatase 2 (SULF2), 5-hydroxytryptamine receptor 2C (HTR2C), cut like homeobox 2

(CUX2), oligodendrocyte precursor cells (OPC).

Fig. 2: Differential expression. a The number of up or down regulated genes by cell type. **b**

UpSet plot of the number of unique differentially expressed genes (DEGs) that are unique or

shared between the five neuronal cell clusters that encompass the majority (~96%) of all DEGs.

c Overrepresentation of schizophrenia (SZ), bipolar disorder (BP), and Alzheimer's disease

(AD) GWAS loci in primary neuronal clusters. **d** Heatmap of the overrepresentation of cluster-

specific DEGs throughout the brain. **e** Heatmap of the overrepresentation of cluster-specific

DEGs throughout the neural life span.

Fig. 3: Canonical pathways. a Overrepresentation of cluster-specific DEGs in the top five

shared canonical pathways for the primary neuronal clusters. **b** Shared and unique cluster-

specific DEGs that underlie the oxidative phosphorylation canonical signaling pathway.

Fig. 4: Upstream regulators. a Overrepresentation of cluster-specific DEGs in the top five

predicted upstream master regulators. **b** Shared and unique cluster-specific DEGs that underlie

the prediction of DDX5 as an upstream regulator of cell type-specific DEGs.