Genetically Determined Strength of Natural Killer Cells is Enhanced by Adaptive HLA class I Admixture in East Asians

Zhihui Deng[1,2,+], Jianxin Zhen[1,3,+], Genelle F. Harrison[4,5,+], Guobin Zhang[1], Rui Chen[1], Ge Sun[1], Qiong Yu[1], Neda Nemat-Gorgani[6], Lisbeth A. Guethlein[6], Liumei He[1], Mingzhong Tang[7], Xiaojiang Gao[8], Siqi Cai[1], Jonathan A. Shortt[4], Christopher R. Gignoux[4], Mary Carrington[8], Hongyan Zou[1], Peter Parham[6], Wenxu Hong[9,*] and Paul J. Norman[4,5,**]

1. Immunogenetics Laboratory, Shenzhen Blood Center, Shenzhen, Guangdong 518035, P. R. of China.
2. Department of Transfusion Medicine, School of Laboratory Medicine and Biotechnology, Southern Medical University, Guangzhou, Guangdong 510515, P. R. of China.
3. Central Laboratory, Shenzhen Baoan Women's and Children's Hospital, Shenzhen, Guangdong 518100, P. R. of China.
4. Division of Biomedical Informatics and Personalized Medicine, University of Colorado, Anschutz Medical Campus, Aurora, CO 80045, USA.
5. Department of Immunology and Microbiology, University of Colorado, Anschutz Medical Campus, Aurora, CO 80045, USA.
6. Department of Structural Biology, Stanford University School of Medicine, Stanford, CA94305, USA.
7. Clinical Laboratory, Wuzhou Red Cross Hospital, Wuzhou, Guangxi 543002, P. R. of China.
8. Frederick National Laboratory for Cancer Research (FNLCR), Frederick, MD21702, USA.
9. Shenzhen Institute of Transfusion Medicine, Shenzhen Blood Center, Shenzhen, Guangdong 518035, P. R. of China.

[+] These authors contributed equally to this work

Correspondence: *Wenxu Hong: szbloodcenter@hotmail.com and **Paul J. Norman: paul.norman@cuanschutz.edu

Keywords: HLA class I; KIR; admixture; adaptive introgression; natural killer cells; infectious disease; East Asia.

Abstract

Human natural killer (NK) cells are essential for controlling infection, cancer and fetal development. NK cell functions are modulated by interactions between polymorphic inhibitory killer cell immunoglobulin-like receptors (KIR) and polymorphic HLA-A, -B and -C ligands expressed on tissue cells. All *HLA-C* alleles encode a KIR ligand and contribute to reproduction and immunity. In contrast, only some *HLA-A* and *-B* alleles encode KIR ligands and they focus on immunity. By high-resolution analysis of *KIR* and *HLA-A*, *-B* and *-C* genes, we show that the Chinese Southern Han are significantly enriched for interactions between inhibitory KIR and HLA-A and -B. This enrichment has had substantial input through population admixture with neighboring populations, who contributed *HLA class I* haplotypes expressing the KIR ligands B*46:01 and B*58:01, which subsequently rose to high frequency by natural selection. Consequently, over 80% of Southern Han *HLA* haplotypes encode more than one KIR ligand. Complementing the high number of KIR ligands, the Chinese Southern Han *KIR* locus combines a high frequency of genes expressing potent inhibitory KIR, with a low frequency of those expressing activating KIR. The Southern Han centromeric *KIR* region encodes strong, conserved, inhibitory HLA-C specific receptors, and the telomeric region provides a high number and diversity of inhibitory HLA-A and -B specific receptors. In all these characteristics, the Southern Han represent other East Asians, whose NK cell repertoires are thus enhanced in quantity, diversity and effector strength, likely through natural selection for resistance to endemic viral infections.

1 Introduction

Human leukocyte antigen (HLA) class I molecules are critical components of immunity, whose

extreme variation associates with resistance and susceptibility to infection, multiple immune-

mediated diseases and some cancers (Dendrou et al. 2018). *HLA class I* genes are located in

the *major histocompatibility complex* (*MHC)* of chromosome 6 and encode proteins that bind

peptide fragments derived from intracellular protein breakdown and transport them to the cell

surface. In doing so they can communicate to the adaptive immune system's T cells whether a

tissue cell is healthy, or unhealthy due to infection or cancer. Subsets of HLA class I allotypes

additionally contain an externally facing amino acid motif that binds killer cell

immunoglobulin-like receptors (KIR), facilitating interaction with natural killer (NK) cells of

innate immunity.

KIR are expressed on the surface of NK cells and regulate their functions through binding to

HLA class I ligands on other cells (Cooper et al. 2009; Long et al. 2013). The functions of

these interactions are crucial in immunity to aid recognition and elimination of infected or

tumorous tissue, and in reproduction to regulate placentation and fetal development (Parham

and Moffett 2013). In accordance with these critical and independent roles in human health,

KIR and their HLA class I ligands are subject to natural selection, mediating their exceptional

diversity across individuals, populations and species (Parham and Moffett 2013; Prugnolle et

al. 2005). Indeed, *KIR* and *MHC* are some of the fastest evolving genomic loci in higher

primates (Guethlein et al. 2015). Correlating with direct impact on both NK cell development

and effector function (Freud et al. 2017; Vivier et al. 2011), numerous studies have implicated

combinatorial diversity of *KIR* and *HLA class I* alleles with the course of specific infectious

and autoimmune diseases, as well as the success of transplantation (Boudreau and Hsu 2018;

Holzemer et al. 2017). Importantly, the quantity as well as quality of these interactions can

influence individual responses to infection (Boelen et al. 2018; Pelak et al. 2011). Thus, the

polymorphism of *KIR* and *HLA class I* has profound impact on human health. Under-explored

are the scale and characteristics of *KIR* and *HLA class I* combinatorial diversity worldwide,

and the processes that shape this diversity.

NK cells express overlapping subsets of KIR that are acquired stochastically during their

development (Andersson et al. 2009). During this process, the interaction of inhibitory KIR

with HLA class I KIR ligands broadens and strengthens subsequent effector functions of the

NK cell repertoire (Bjorkstrom et al. 2016; Hoglund and Brodin 2010; Saunders et al. 2015).

This education process matures some NK cells, allowing them to respond effectively to specific

instances of infection or cancer, and enhances the NK cell repertoire compared to those that

develop using other more conserved pairs of ligands and receptors. In this role, and also in

pregnancy where HLA-A and -B have no function, HLA-C is dominant because all expressed

HLA-C are KIR ligands (Guethlein et al. 2015). Four mutually exclusive sequence motifs

define the four HLA class I epitopes that are KIR ligands: C1 is carried by subsets of HLA-C

and HLA-B allotypes. C2 is carried by the other allotypes of HLA-C. Bw4 is carried by subsets

of HLA-A and -B allotypes. The A3/11 motif is carried by a subset of HLA-A allotypes (HLA-

A*03 and A*11). Thus, only some HLA-A and -B allotypes are KIR ligands and their main

role is likely to diversify the NK cell response to pathogens.

The *KIR* locus on chromosome 19q13.4 varies in gene content, containing up to eight genes

encoding inhibitory KIR and five encoding activating KIR (Wilson et al. 2000). Four of the

inhibitory KIR and four activating KIR have well-characterized HLA-A, -B or -C ligands. Two

broad groups of *KIR* haplotypes are present in every human population. *KIR A* haplotypes carry

all four of the HLA-class I specific inhibitory receptors and are associated with resistance to

infectious diseases (Bashirova et al. 2006). *KIR B* haplotypes are more variable in their gene

number, carrying two or more genes for inhibitory receptors as well as various activating

receptor genes, and favor fetal development (Parham and Moffett 2013). A recombination hotspot separates the *KIR* locus into two segments (Wilson et al. 2000). Two inhibitory receptors specific for HLA-C are encoded in the centromeric region, and two HLA-A and -B specific receptors are encoded in the telomeric region. Additional to gene content variation, polymorphism of both receptors and ligands can directly affect NK cell activity (Guethlein et al. 2015). Thus, by varying the number, density, specificity, strength or signaling properties of the receptor-ligand interaction, genetic variation of *KIR* and *HLA class I* can pre-determine functional differences in NK cell repertoires between individuals. This genetic diversity is substantial among populations, as demonstrated with high-resolution studies (Guethlein et al. 2015; Nemat-Gorgani et al. 2018). In such detailed analysis, Asian populations are under-represented.

Comprising 20% of the human population, the Chinese Han are the largest ethnic group in the world (Abdulla et al. 2009). The Han have a complex population history and are presently structured with the Northern and Southern Han forming two main subgroups that are separated geographically by the Yangtze River (Wen et al. 2004). The Southern Han originated through large scale population movement from the north ~1500 years ago, in parallel with admixture with resident and neighboring populations (Hellenthal et al. 2014; Wen et al. 2004). Importantly for the current study, the major genetic distinction between the Northern and Southern Han occurs in the *MHC*, and localizes to the region that spans *HLA-A*, *-B* and *-C* (Chen et al. 2016; Xu et al. 2009). The most significant component of this difference is the *A*33:03-B*58:01-C*03:02 HLA class I* haplotype, which is common in the Southern Han and remains conserved across multiple unrelated individuals (Chen et al. 2016). Such strong linkage disequilibrium is consistent with recent acquisition of this haplotype by admixture (Chen et al. 2016). This haplotype encodes two KIR ligands, HLA-B*58:01 and C*03:02 (Guethlein et al. 2015). Although less is known of *KIR* allele diversity in the Han, several

studies established that the genes characteristic of *KIR A* haplotypes are common, and

demonstrated differences in their distribution among the different Han groups and among other

resident populations (Bao et al. 2013; Wang et al. 2012; Yao et al. 2011). These studies also

confirmed that *KIR* and *HLA class I* combinatorial diversity is an important factor in pregnancy

syndromes, infectious disease, blood cancers and transplantation outcome in the Han. They

also uncovered both similarities and differences from the specific disease associations observed

in Europeans (Bao et al. 2016; Jiang et al. 2013; Long et al. 2015; Shen et al. 2016; Su et al.

2018). To investigate these findings, we have examined how demographic and evolutionary

processes have shaped combinatorial diversity of HLA class I and KIR in the Chinese Southern

Han.

Materials and methods

Study samples

Peripheral blood samples were collected from 306 unrelated healthy volunteer blood donors

from Shenzhen, Guangdong, China. All donors self-identified to be of Han ethnicity from

southern China. All subjects provided written informed consent for participation in the present

research, which was approved by the ethics review board of Shenzhen Blood Center, Shenzhen,

Guangdong, China.

Genomic DNA extraction

Genomic DNA was extracted from 400 μl of peripheral blood using a MegCore Nucleic Acid

Extractor (MegCore, Taiwan, China). DNA purity and concentration were tested by UV-

spectrophotometry using a Biophotometer (Eppendorf, Hamburg, Germany) and adjusted to a

concentration of 50-100 ng/μl.

High-resolution HLA-A, -B and -C genotyping

HLA-A, *-B* and *-C* genotyping was performed using the AlleleSEQR HLA sequencing-based

genotyping commercial kit (Atria Genetics, San Francisco, USA). According to the

manufacturer's instructions, exons 2-4 for *HLA-A*, *-B* and *-C* were sequenced in both directions

using an ABI 3730XL DNA sequencer (Applied Biosystems, Foster City, CA, USA). HLA

genotypes were assigned using the Assign 4.7 software (Conexio Genomics, Fremantle,

Australia). Samples giving ambiguous allele combinations by sequencing were further resolved

using HLA PCR-SSP (Olerup, Stockholm, Sweden).

High-resolution KIR genotyping

The presence or absence of *KIR2DL1*, *2DL2/3*, *2DL4*, *2DL5*, *2DS1*, *2DS2*, *2DS3*, *2DS4*, *2DS5*,

3DL1/S1, *3DL2* and *3DL3* was first determined for each individual using the 'KIR Ready Gene'

PCR-SSP kit (Inno-Train Diagnostik GmbH, Frankfurt, Germany). The *KIR* genes identified

using PCR-SSP were then subject to nucleotide sequencing of all exons (Deng et al. 2018).

Sequencing reactions were performed using ABI PRISM BigDye Terminator Cycle

Sequencing Ready reagents and analyzed using an ABI 3730 DNA Sequencer (Applied

Biosystems, Foster City, USA). *KIR* alleles were assigned using Assign 4.7 allele identification

software (Conexio Genomics, Fremantle, Australia), and release 2.6.1 (February 2015) of the

Immuno-Polymorphism database (IPD) (Robinson et al. 2015). When the sequencing results

gave ambiguous allele combinations, we used group-specific PCR primer pairs to amplify and

sequence the target alleles separately (Zhang and Deng 2016)

Novel KIR alleles

To confirm and fully characterize any novel allele identified during amplicon sequencing we

cloned and sequenced *KIR* transcripts. Further samples of peripheral blood samples were

collected, and total RNA isolated using the Maxwell 16 low elution volume simplyRNA Blood

Kit (Promega, Madison, USA). Complementary DNA (cDNA) was synthesized using the

Transcriptor First Strand cDNA Synthesis Kit (Roche, Basel, Switzerland). *KIR* transcripts

were amplified specifically from cDNA using primer pairs described previously (Yawata et al.

2006), with addition of KIR3DL3-specific primers (forward 5'-

GGTTCTTCTTGCTGGAGGGGC-3' and reverse 5'-TTACACGCTGGTATCTGTTGGGG-

3'). The amplified transcripts were cloned using the TA cloning kit (Takara, Dalian, China)

and at least three clones of any novel allele were sequenced. The sequences of novel *KIR* alleles

were submitted to GenBank and the IPD KIR database (Robinson et al. 2015) to obtain official

names.

Admixture Estimates

Whole genome SNP genotypes for Japanese (N = 104), Vietnamese (N = 99), Han from Beijing

(N = 103), Southern Han (N = 105), and Dai (N = 93) were obtained from the 1000 Genomes

Project (Auton et al. 2015). We used any SNPs having minor allele frequency >1% and

independent of other SNPs (linkage disequilibrium, $r^2 < 0.3$). Admixture was calculated for

chromosome 6 using the ADMIXTURE program (Alexander et al. 2009), with the

unsupervised option and k=3. Two regions were analyzed, the *MHC* (chr6:28,477,797–

33,448,354: 3,541 SNPs) and chromosome 6 excluding the *MHC* (84,898 SNPs). We selected

a K of 3 to represent the three primary ancestry groups in the region that are represented in the

1000 Genomes data: Japanese, South East Asian, and East Asian (Chen et al. 2016). *HLA class*

I alleles were obtained from the 1000 Genomes Project data (Gourraud et al. 2014). We

analyzed the Hondo Japanese (JPT), Vietnamese (KHV), Chinese Dai (CDX), Chinese

Southern Han (CHS), and Beijing Han (CHB) Validating their use for this purpose, the

correlation of the *HLA class I* allele frequencies between our study population and the CHS is

0.95 (p= 6.65^{-11}, Figure S1). Individuals were considered carriers if they had at least one copy

of the respective allele. Distributions of ancestry proportions for carriers and non-carriers of

specific *HLA* alleles were compared using a Wilcoxon test, using the wilcox.test function in R

(R Development Core Team 2008).

Estimates of nucleotide diversity

We used π (Nei and Takahata 1993) to measure the nucleotide diversity of haplotypes carrying

specific *HLA-B* alleles. We used the phased genomes of the Chinese Southern Han (CHS)

population available from the 1000 Genomes Project (Auton et al. 2015), and extracted the

genomic region containing the *HLA-B* and *-C* genes, with 500kbp flanking on each side. For

each carrier of a given allele, we identified (by sequence) and retained the haplotype

representing the allele of interest. For each given allele, we pooled all of the respective

haplotypes present in the population and calculated π in 100bp windows using VCFtools

(Danecek et al. 2011). Distributions of π values were compared between respective alleles with

a Wilcoxon test using the wilcox.test function in R.

Tests for positive selection affecting specific HLA class I alleles

We filtered 1,000 Genomes genotyping data of chromosome 6 from the CHS population to

remove non-biallelic and duplicated SNPs (Purcell et al. 2007), then phased using the program

Eagle (Loh et al. 2016). We used the program Selscan (Voight et al. 2006) to calculate

integrated haplotype statistic (iHS). The statistic is a measure of haplotype diversity associated

with a given genetic variant, where lower diversity and longer haplotypes correlate with

selection of that variant.

To determine if specific *HLA class I* alleles have been targeted by directional selection in the

Chinese Southern Han we again used the 1,000 Genomes SNP data from the CHS population.

SNPs within the following hg19 coordinates were used: *HLA-A*, Chr6: 29,910,089 –

29,913,770; *HLA-B*, Chr6: 31,321,648 – 31,325,007; *HLA-C*, Chr6: 31,236,517 – 31,239,917.

We phased haplotypes from individuals positive for each given *HLA class I* allele and aligned

them to reference sequences to identify the haplotype containing that allele. The alignments

were then used to identify 'tagging' SNPs that could be used to identify each given *HLA class*

I allele. The criteria for choosing tagging SNP alleles were that they must be present in every

individual carrying the corresponding *HLA class I* allele and that they must be absent from the

other *HLA class I* alleles in the analysis. We analyzed the alleles present on the 10 most frequent

HLA class I haplotypes that we observed in the Chinese Southern Han; *HLA-B*15:02* was

excluded because we were not able to identify unique tagging alleles on haplotypes carrying

this allele. For each tagging SNP, we calculated the integrative haplotype score (iHS) using

SelScan. We used the absolute value of iHS since derived alleles under selection will have a

negative value and ancestral alleles under selection will have a positive value (Szpiech and

Hernandez 2014). Using a Wilcoxon two-sample test, we examined whether the distributions of absolute iHS values differed between tagging SNPs of *HLA* alleles and SNPs of the full chromosome 6.

Haplotype and ligand frequencies

KIR and *HLA-A*, *-B* and *-C* allele frequencies were calculated from the observed genotypes. For individuals genotyped as homozygous for an allele of a given *KIR* that has presence/absence polymorphism, the number of copies present was determined by analyzing LD with alleles of the flanking genes. The subsequent genotype distributions for all loci were consistent with Hardy-Weinberg equilibrium. *KIR* haplotype frequencies were determined using PHASE II (Stephens and Donnelly 2003). The following parameters were used; -f1, -x5, and -d1, and from the output, the two haplotypes with highest probability were taken for each individual. Watterson's homozygosity F test was performed using Pypop software (Lancaster et al. 2007), with 10,000 replicates to calculate the normalized deviate F_{nd} test (Salamon et al. 1999).

HLA class I (A-B-C) haplotype frequencies were determined using the EM algorithm of 'Arlequin software version 3.5 (Excoffier and Lischer 2010). For comparison with other populations, we only used populations for which *HLA class I* genotype data were available from every individual sampled, and for which the resolution of genotyping was the same as the Southern Han described here. We therefore used the subset of populations described in the 13[th] International Histocompatibility Workshop and Conference report that have 50 or more *HLA-A*, *-B* and *-C* genotyped individuals (Meyer et al. 2007). These were supplemented with our own data from the Ga-Adangbe from Ghana in West Africa (Norman et al. 2013), KhoeSan from Southern Africa (Nemat-Gorgani et al. 2018), Yucpa from South America (Gendzekhadze et al. 2009), Europeans from the USA (Norman et al. 2016) and Hondo Japanese (Yawata et

al. 2006). We compared the proportion of *HLA class I* haplotypes encoding one KIR ligand to those encoding more than one KIR ligand across populations using a two-proportions Z test, using the prop.test function in R (R Development Core Team 2008).

Comparison of HLA class I and KIR ligand distributions.

Clustering based on allele frequencies: Any *HLA class I* allele occurring in fewer than two of the populations studied was excluded from this analysis. The allele frequencies of all three *HLA class I* genes were used for each population. Cluster dendrograms were constructed using R 3.4.3, hclust with 1,000 bootstrap values. The package used was fpc (Hennig 2020). Cluster dendrograms were constructed in the same manner using the frequencies of the *HLA class I* haplotypes encoding one, two or three KIR ligands.

Assessment of receptor/ligand quality and quantity

As described previously (Nemat-Gorgani et al. 2018), experimental data were used to determine the interacting pairs of KIR and HLA class I, which are listed in **Figure S2**. To determine the quantity of receptor/ligand interactions, the number of KIR/HLA allotype pairs that are known to interact were summed for each individual, and homozygous KIR or HLA allotypes were counted twice. To determine the diversity of interactions, the number of different KIR/HLA allotype pairs that are known to interact were summed for each individual (in this case homozygous allotypes were counted once). Populations were compared using unpaired t tests, using GraphPad software.

1 Results

2 *High Frequency of KIR ligands in the Southern Han*

All HLA-C and subtypes of HLA-A and -B allotypes are ligands for KIR, which are expressed

on the surface of NK cells to modulate their functions in immunity and reproduction. Within

human populations, *HLA class I* haplotypes tend to form a balance between those that encode

HLA-A or -B KIR ligands and those that do not (Guethlein et al. 2015). To determine if this

pattern is also observed in the Chinese Southern Han, we analyzed the *HLA-class I* genes of

306 healthy individuals. We identified 27 *HLA-A*, 54 *HLA-B* and 29 *HLA-C* alleles (Figure S3).

Each of these 110 alleles encodes a different HLA class I allotype, and 58 of them are known

KIR ligands (Figure 1A). The majority of 233 *HLA class I* haplotypes, including the ten most

frequent (Figure 1B), encode more than one KIR ligand (70.3% of distinct haplotypes; 81.8%

by frequency, Figure S4A). This observation is unusual and indicates the balance between

having and not having KIR ligands at HLA-A and -B is perturbed in the Chinese Southern Han.

To investigate the unusually high frequency of KIR ligands, we compared Southern Han *HLA*

class I haplotypes with those of sub-Saharan African, Oceanian, European and South American

populations that represent major modern human groups (Rosenberg et al. 2002; Tishkoff et al.

2009). In this data set, rather than the larger and more widely studied Hondo Japanese

population, a Ryukyu Japanese population was included because they more closely represent

the Japanese population prior to admixture with Han (Takeuchi et al. 2017). Among the eight

populations, 1,034 different *HLA class I* haplotypes were observed (Figure S4B). Six

populations have a similar distribution of KIR ligands, with each population having an

approximately equal frequency of *HLA class I* haplotypes carrying one and two KIR ligands,

and a smaller frequency of haplotypes carrying three KIR ligands (Figure 1C). Only Southern

Han and South Americans differed from this pattern, with the Han encoding more and the

Amerindians encoding less KIR ligands per haplotype than other populations (Figure 1C). The

difference in the proportion of *HLA class I* haplotypes encoding one versus more than one KIR ligand between the Southern Han and each of the other seven representative populations is statistically significant, as is that between Amerindians and the other populations (Two-proportions Z-test, Benjamini-Hocherg corrected p <0.001, Figure 1C). The allele frequency distribution of South American Amerindians was likely influenced by severe population bottlenecks, leading to a reduced genome-wide diversity compared with other populations (Fagundes et al. 2008; Raghavan et al. 2015), whereas the Han were not subject to severe population-specific bottleneck (Henn et al. 2012; Lu et al. 2016; Schiffels and Durbin 2014). To examine if the Chinese Southern Han are representative of other related populations, we examined groups from East Asia (Hondo Japanese and Korean) and Southeast Asia (Thai, Malay and Filipino). This analysis showed these populations also have a high frequency of *HLA class I* haplotypes encoding multiple KIR ligands (Figure 1D). Our analysis thus shows that East Asian and South East Asian *HLA class I* haplotypes encode more ligands for inhibitory KIR than the haplotypes of any other populations.

Despite having distinct population histories, the sub-Saharan African, Oceanic, European and Ryukyu Japanese populations all have a similar mean number of KIR ligands per *HLA* haplotype (Figure 1C). However, very few *HLA class I* haplotypes are shared by any of these populations. For example, only 19 of 369 haplotypes detected in Africans are present in more than one of the three African populations studied, and comparing the disparate Southern African Nama, Indigenous Australian and Ryukyu Japanese populations revealed just five haplotypes in common (Figure 1E).

The Chinese Southern Han acquired HLA haplotypes encoding multiple KIR ligands by admixture

Previous analyses suggested that specific *MHC* region haplotypes (which include the *HLA* genes) present in the Chinese Southern Han were obtained from the Northern Han through admixture (Chen et al. 2016). The most frequent HLA class I allotypes contributing to the enrichment of KIR ligands in the Chinese Southern Han are HLA-A*11, -A*24, -B*46, and -B*58 (Figure 1B). We therefore examined the relative contributions of admixture to the high frequency of these alleles in the Chinese Southern Han. For this analysis we considered known admixture events (Hellenthal et al. 2014; Wen et al. 2004; Xu et al. 2009) and drew upon the 1000 Genomes SNP and *HLA* genotype data (Auton et al. 2015; Gourraud et al. 2014) from Hondo Japanese, Vietnamese, Dai, Beijing Han, and Chinese Southern Han. Consistent with previous work examining whole-genome data (Takeuchi et al. 2017), in analyzing chromosome 6 we identified three primary genetic ancestries, corresponding to the Japanese, East Asian (Southern and Beijing Han) and South East Asian (Vietnamese and Dai) population groups (Figure 2A). That we identify a higher 'Japanese' component in the Beijing than Southern Han (36% vs 22%: Figure 2A) likely reflects the higher proportion in Beijing of Northern Han (Auton et al. 2015), a population from which we have no data for the current study. Supporting this observation, the greatest contribution from China to Japanese ancestry is from the Northern Han (Chen et al. 2016; Takeuchi et al. 2017).

We compared the relative proportions of the three genetic ancestries in the Chinese Southern Han within the *MHC* region of chromosome 6 to their proportions throughout chromosome 6 excluding the *MHC*. This analysis revealed a predominance of East Asian ancestry throughout the length of chromosome 6, including the *MHC* (Figure 2B). In carriers of *HLA-B*40:01*, the most frequent *HLA-B* allele in Chinese Southern Han, there is also clear East Asian ancestry throughout the length of chromosome 6 (Figure 2B). The proportion of East Asian ancestry is similar in *B*40:01* carriers than non-carriers (Wilcoxon test, p=0.98). By contrast, among *HLA-B*46:01* carriers, the *MHC* is primarily of South East Asian ancestry (Figure 2B) with

carriers having a significantly higher proportion of South East Asian genetic ancestry in the

MHC than outside the *MHC* (p = 2.7^{-6}), or within the *MHC* of non-carriers (p = 1.9^{-5}). Similarly,

among *HLA-B*58:01* carriers, the *MHC* region is primarily of Japanese ancestry (Figure 2B),

with carriers having a significantly higher proportion of Japanese genetic ancestry within the

MHC than outside the *MHC* (p = 2.4^{-4}) and compared with non- *B*58:01* carriers (p = 6.1^{-7}).

Excluding the *MHC,* carriers of any of these three alleles show East Asian ancestry along

chromosome 6 (Figure 2B). Further supporting the observed population structure as specific to

the *MHC* region, among the three ancestral groups the F_{ST} values range from 0.098 – 0.161,

compared with 0.012 – 0.017 for the remainder of chromosome 6.

Based on the analysis of *B*46* and *B*58*, we examined the proportions of genetic ancestry of

alleles that comprise the 10 most frequent *HLA class I* haplotypes observed in the Chinese

Southern Han. The primary genetic ancestry outside of the *MHC* region was determined as East

Asian for every allele studied (Figure 2B). Thus, for our comparisons, we determined the

primary genetic ancestry in the flanking *MHC* region for each allele and then determined the

relative proportion of that ancestry in the remainder of the *MHC*. This analysis identified six

haplotypes maintaining strong evidence of East Asian genetic ancestry both within the *MHC*

and throughout chromosome 6. These haplotypes include those that carry *A*11:01* and

*A*24:02*, as well as *B*40:01* (Figure 2C). It was shown previously that HLA-A*11 and -A*24

derive from introgression with archaic humans (Abi-Rached et al. 2011) and our results and

others (Gonzalez-Galarza et al. 2015; Solberg et al. 2008) thus suggest these haplotypes are

now endemic to East Asia. The analysis also identified four haplotypes having genetic ancestry

within the *MHC* that is distinct from the ancestry of the remainder of the chromosome (Figure

2C). For three of the haplotypes, which include the two most frequent haplotypes in the

population, this distinction is statistically significant ($p_{corr} < 0.01$). Two of these haplotypes

contain *B*46:01* and one contains *B*58:01* (Figure 2C). In total, four of five of the *HLA-B*

alleles that encode a KIR ligand and are present on these 10 most frequent haplotypes show increased evidence for admixture in the *MHC* region. By contrast, neither of the two *HLA-A* alleles that encode a KIR ligand show a genetic ancestry within the *MHC* that differed from the East Asian ancestry throughout chromosome 6. This finding suggests that the number of *HLA-B* genes encoding KIR ligands was enhanced in Chinese Southern Han by admixture with neighboring or displaced populations. In summary, these findings clearly show that the *B*46:01* and *B*58:01* alleles are present in Chinese Southern Han through admixture.

Positive selection favors HLA haplotypes expressing more than one KIR ligand in Chinese Southern Han

To investigate whether or not the admixed haplotypes were also subject to natural selection we examined further characteristics of their diversity and distribution. We first measured nucleotide diversity (π) of the genomic sequence flanking +/- 500kb of specific *HLA-B* alleles (Figure 3A). We found significantly reduced nucleotide diversity of haplotypes containing *HLA-B*46* compared to haplotypes containing *HLA-B*40* (mean π of *B*40* = 2.2 x 10^{-3}, *B*46* = 0.6 x 10^{-3}, Wilcoxon test, p = 1.24 x 10^{-12}). We also observed that haplotypes containing *B*58* have lower diversity than *B*40*, but this reduction was not statistically significant (mean π of *B*58* = 1.6 x 10^{-3}, Wilcoxon test, p =0.12). This reduced diversity suggests that *B*46* haplotypes have arisen in frequency in the Chinese Southern Han without accumulating mutations. To further explore this finding, we used the iHS statistic, which identifies genomic variants that have increased in frequency recently and rapidly under natural selection, so that their haplotypic background has not yet been diversified by recombination (Voight et al. 2006). We identified a strong signal of recent selection (iHS >= 99[th] percentile) that falls precisely in the *MHC* of the Chinese Southern Han (Figure 3B).

To investigate the patterns of selection specific to *HLA-B*46:01* and *B*58:01* haplotypes we

first identified SNPs that characterize those haplotypes and then compared their distribution of

iHS values to the distribution of all the SNPs of chromosome 6 (Figure 3C). For *B*46:01* the

mean absolute iHS of 3.38, was significantly higher than the mean for chromosome 6 of 0.785

(Wilcoxon two-sample test, p= 5.77^{-6}), as was the mean iHS for *B*58:01* (3.43, p= 1.8^{-3}).

Although the signal for *B*58:01* is weaker, there is a more distinct subset of SNPs having

extremely high iHS values (>99th percentile, Figure 3C), which could indicate recent selection

of an older haplotype, although it was not possible from our analysis to determine if the SNP

allele was ancestral or derived in each case. Interestingly, the mean iHS for *B*40:01* associated

SNPs was also significantly higher than the chromosome average (1.88, p= 7.7^{-6}). However,

fewer *B*40:01* specific SNPs had an iHS value in the 95th percentile than *B*58:01* or *B*46:01*

specific SNPs (30%, 50%, and 100% of SNPs respectively). Because the most frequent

B*40:01 containing haplotypes in the Chinese Southern Han carry either *A*11:01 or A*24:02,*

which are KIR ligands (Figure 1B), we extended the analysis to these alleles (Figure 3C).

Again, this analysis showed both *A*11:01* (mean = 2.8, Wilcoxon two sample test p= 1.45^{-11})

and *A*24:02* (mean = 1.37, p= 1.27^{-9}) associated SNPs have significantly higher iHS values

than the chromosome average, with *A*11:01* having a mean iHS that is above the 95th

percentile. Haplotypes carrying *HLA-A*11:01, A*24:02, B*46:01* or *B*58:01* were previously

identified to have unusually high LD in this population (Chen et al. 2016). This analysis

identified two other *HLA class I* alleles as having highly distinct signatures of directional

selection, A*30 and C*07. Whereas HLA-C*07 is known to interact strongly with KIR to

educate NK cells (Hilton et al. 2015a; Yawata et al. 2006), HLA-A*30 does not possess a KIR

ligand. Together, these findings thus illustrate that HLA class I in the Chinese Southern Han

has been targeted by natural selection and suggest that one major benefit has been to increase

the number of KIR ligands present in the population.

We next examined whether the observed distributions of *HLA class I* encoded KIR ligands were consistent with modern human population dispersal. Cluster analysis shows there are five groups of *HLA class I* frequency spectra that correspond to the broad population groups of African, European, Asian, Oceanian and American origin (Figure 3D). By contrast, three distinct and strongly supported groups cluster according to their proportions of haplotypes encoding one, two or three KIR ligands (Figure 3E). Notable examples are the Ryukyu Japanese and Indigenous Australian populations, who group with Asian populations when analyzed by *HLA class I* haplotype distribution (Figure 3D). By contrast, Ryukyu Japanese and Australians appear more similar to Africans and other groups when analyzed according to the number of KIR ligands encoded by their *HLA class I* haplotypes (Figure 3E). In a counter example, Northern Indian and Tuvan populations (Figure S4C) cluster with Europeans when analyzed by their *HLA class I* alleles, but with East Asians when analyzed by the number of KIR ligands. Thus, *HLA class I* allele frequency distributions are consistent with the origins of the populations studied and with the pattern of human dispersal out of Africa (Henn et al. 2012), whereas the number of KIR ligands encoded by *HLA class I* haplotypes is not. This finding suggests that the similar number of KIR ligands observed across populations is likely due to convergent evolution, because distinct *HLA class I* haplotypes produce similar distributions of KIR ligands. The findings thus also support our assessment that the unusual distribution of KIR ligands in East Asian populations is due to natural selection.

In summary, these results show that similar quantities of KIR ligands can be obtained using different subsets of *HLA class I* haplotypes, indicating there is pressure to maintain a certain balance of KIR ligands across populations, regardless of the background HLA allotype, and that this balance is perturbed in East Asia. Our observations show that successive rounds of admixture followed by natural selection favouring specific *HLA class I* haplotypes have increased the quantity of KIR/HLA interactions of populations in East Asia. To investigate the

characteristics of this receptor and ligand diversity, we next studied the *KIR* locus in the

Chinese Southern Han.

High frequency of inhibitory KIR allotypes in Southern Han

The *KIR* locus comprises genes encoding the four inhibitory and six activating KIR known to

bind polymorphic HLA class I ligands, and three that do not bind polymorphic HLA class I

(Guethlein et al. 2015). In total, we identified 116 *KIR* alleles, representing 101 KIR allotypes

(Figure S5). A total of 46 novel *KIR* alleles (39.7% of total *KIR* alleles detected) were

characterized (Figure S6) and 24.8% of the individual Han carried at least one novel allele.

Correcting for the number of individuals tested showed that the Southern Han are more diverse

than Amerindians and Oceanians, but less diverse than Europeans and Africans (Figure 4A).

KIR diversity of the Chinese Southern Han is thus consistent with genome-wide diversity when

compared to other populations (Campbell and Tishkoff 2008). The Chinese Southern Han have

70 centromeric and 91 telomeric *KIR* haplotype motifs that combine to form a minimum of 199

KIR haplotypes (Figure S7A-C). The majority are *KIR A* haplotypes (74.7%, Figure 4B),

including 8 of the 10 most frequent haplotypes (Figure 4C). This skewing towards *KIR A*

haplotypes is more pronounced in the centromeric region (87.9%) than the telomeric (79.7%)

region (Figure 4C, Figure S7A-B).

KIR A haplotypes encode all four inhibitory receptors that bind HLA class I ligands and either

one or no activating receptors (Wilson et al. 2000). Accordingly, among the *KIR* alleles

identified in the Chinese Southern Han, we observed high frequencies of those encoding strong

inhibitory receptors. Both KIR2DL1*003, a strongly-inhibiting allotype of KIR2DL1 (Bari et

al. 2009; Hilton et al. 2015a), and KIR2DL3*001, a strongly-inhibiting allotype of KIR2DL2/3

(Yawata et al. 2006), are common in the Chinese Southern Han, having frequencies of 73.5%

and 71.1% respectively (Figure S5). Also frequent are KIR3DL1*015, which is a strong

inhibitor on binding to the Bw4 ligand (Yawata et al. 2006), and KIR3DL2*002 that has high

expression but unknown functional properties (Figure S5). Noticeably scarce are inhibitory

KIR allotypes having mutations that prevent cell surface expression, of which there are many

examples (Bari et al. 2009; Hilton et al. 2015b; Pando et al. 2003; VandenBussche et al. 2006).

For instance, weakly-expressed KIR3DL1*004 is frequent in many populations (Norman et al.

2007), but absent from the Chinese Southern Han (Figure S5, and ref (Tao et al. 2014)). Also

rare in the Chinese Southern Han are alleles encoding inhibitory allotypes of reduced function,

such as KIR2DL1*004 (3.6%, Figure S5), which is common in other population groups (Bari

et al. 2009; Meenagh et al. 2008; Nemat-Gorgani et al. 2014; Norman et al. 2013; Vierra-Green

et al. 2012). Moreover, the frequencies of alleles encoding activating receptors are much lower

(4.4%-18%) than those encoding inhibitory receptors (91.5%-100%), an effect compounded

by presence of multiple non-functional activating KIR allotypes (Figure S5). Exceptional is

KIR2DS4, for which the frequencies of functional and non-functional allotypes are balanced

(55%:45%, Figure S5). These observations point to a strong requirement in the Southern Han

population for retaining high numbers of functional inhibitory KIR, but not activating KIR.

Directional selection reduced centromeric KIR region diversity in the Southern Han

In Chinese Southern Han, the *KIR3DL1/S1* and *KIR3DL2* genes encoding inhibitory NK cell

receptors specific for polymorphic HLA class I ligands, have two or three high frequency

alleles and multiple less frequent alleles (Figure S5). In contrast, *KIR2DL1* and *2DL2/3* also

encode inhibitory receptors but are each dominated by one high frequency allele (Figure 4D).

To explore this observation, we compared the observed homozygosity to the expected across

populations representing major ancestry groups from Europe, Africa, Asia, South America, and

Oceania, using the Ewens-Watterson test (Fnd). *KIR2DL1* and *KIR2DL2/3* are in the

centromeric region of the *KIR* locus, whereas *KIR3DL1/S1* and *KIR3DL2* are telomeric *KIR*

genes (Wilson et al. 2000). Overall the Southern Han show greater homozygosity compared to

other populations, which is more pronounced among centromeric than telomeric *KIR* genes and

statistically significant for *KIR2DL2/3* (Fnd = 3.1, P > 0.985, Figure 4E). This high-resolution

analysis of KIR alleles, complements recent analysis of genome-wide SNP data that identified

directional selection specifically in East Asian centromeric *KIR* (Augusto et al. 2019). The only

other population exhibiting directional selection in the centromeric *KIR* region is the Hondo

Japanese (Yawata et al. 2006). Thus, the profile observed for East Asian populations is distinct

from other populations. Together, these analyses suggest that directional selection reduced

sequence diversity of the centromeric *KIR* in the Southern Han, whereas the telomeric *KIR*

region retains some diversity. In addition, we observed a minimum of eleven different *KIR*

haplotypes having a duplication in the telomeric region (Figure S7D). The telomeric *KIR* have

greater allelic diversity than centromeric *KIR* in Chinese Han (Figure S7D), and these

duplication haplotypes have potential to further diversify the NK cell repertoire because both

allotypes of each gene are expressed (Beziat et al. 2013; Norman et al. 2009). We conclude

that the centromeric *KIR* region provides consistency to Chinese Southern Han NK cell

receptors, whereas the telomeric *KIR* region provides NK cell receptor diversity.

Interactions of KIR with HLA class I

NK cell function is modulated by interactions between KIR and their cognate ligands, HLA

class I molecules. While all HLA-C molecules are always ligands for KIR, only a sub-set of

HLA-A and -B molecules function as KIR ligands. We examined the impact of genetic

variation on the diversity and quantity of KIR/HLA class I interactions in the Chinese Southern

Han. Individuals have a mean of 6.7 different pairs of interacting KIR and HLA class I ligands.

These form a normal distribution in which individuals have from one to twelve interactions

(Shapiro -Wilk test, p = 0.147, Figure 5A). Such normal distributions are seen in other

populations (Nemat-Gorgani et al. 2014; Norman et al. 2013). To investigate the distinct HLA-

A and -B ligand distribution of the Southern Han we divided this analysis into its major

components of KIR interactions with HLA-C, and of KIR interactions with HLA-A and -B (Figure S2). In analyzing only the interactions with HLA-C, we find that functional diversity, as measured by the mean number of different receptor/ligand combinations per individual, is consistent with the overall genetic diversity of the populations studied. At the low end of the range are the Yucpa Amerindians with two different receptor/ligand interactions per individual. At the high end are the Southern African Nama with 4.5 different interactions (Figure 5B).

When the total number of interacting pairs of inhibitory KIR and HLA-C ligands is analyzed, the ranking remains the same but the difference across populations is reduced, ranging from 3.6 to 4.9 viable inhibitory KIR/HLA-C interactions per individual (Figure 5B). On this scale, the Chinese Southern Han are seen to have relatively low diversity and a similar number of interactions between inhibitory KIR and HLA-C to other populations. In sharp contrast, the Chinese Southern Han, together with the Hondo Japanese, have significantly higher number (t-test, $p<0.0001$) and diversity ($p<0.001$), of inhibitory KIR interactions with HLA-A and -B than any other population (Figure 5B). Thus, both the quantity and quality of interactions between inhibitory KIR and HLA-A and -B are enhanced in Southern Han and Hondo Japanese. We predict this will also be true for other East Asian populations.

Discussion

Our analysis shows that the geographic distribution of *HLA class I* alleles and haplotypes is consistent with human dispersal out of Africa and the distance of their geographical location from Africa. Despite the significant differences across human populations in the distributions of *HLA class I* alleles (Meyer et al. 2007), the distribution of KIR ligands is very similar. For *HLA-C*, all haplotypes encode one of two alternative KIR ligands, whereas for *HLA-A* and *HLA-B* there is a balance between haplotypes that either encode one or two KIR ligands, and haplotypes that encode no KIR ligand (Guethlein et al. 2015). Thus, a similar balance of KIR ligands is independently maintained in different human populations using very different HLA-A and -B allotypes. In sharp contrast, the Southern Han Chinese and other East Asian populations do not follow these patterns. The high frequencies of the *HLA class I* haplotypes shared by East Asian populations are an indication of their recent shared ancestry (Abdulla et al. 2009). We find a greater abundance of HLA-A and -B KIR ligands in East Asians than other populations, as well as a greater diversity of interactions between inhibitory KIR and HLA-A and -B. That our comparison included Southern African KhoeSan, whose genetic diversity is the highest among modern humans (Henn et al. 2011; Tishkoff et al. 2009), strongly suggests the high frequency and diversity of KIR ligands in East Asia is the result of natural selection.

The frequency of East Asian *HLA class I* alleles that derive from ancient humans by introgression was previously estimated to be 70-80% (Abi-Rached et al. 2011). The most common of these alleles are *HLA-A*11* and *HLA-A*24*, which encode KIR ligands. We show that more recently, *HLA-B*46:01* and *HLA-B*58:01*, which also encode KIR ligands, were specifically enhanced in frequency in Southern Han following admixture with local populations. HLA-B*46 is a good educator of NK cells (Yawata et al. 2006) and rose in frequency in South East Asia under positive selection (Abi-Rached et al. 2010). The haplotype that encodes HLA-B*58 likely arose in Northern Asia and although the signal is weaker, this

may have been selected both in the Northern and Southern Han. Consequently, *HLA-B*46:01* and *HLA-B*58:01* are the most frequent *HLA-B* alleles encoding KIR ligands, and distinguish the most frequent *HLA class I* haplotypes, in the Southern Han. Together with clear demonstration of natural selection recently targeting the *MHC* region, these findings all support the proposition that natural selection in East Asia favors *HLA class I* haplotypes carrying more than one KIR ligand and suggests there were two major waves of adaptive introgression involving these haplotypes. There is evidence for adaptive introgression of *HLA* alleles in other modern human populations (Busby et al. 2017; Rishishwar et al. 2015). For example, Bantu speakers from western central Africa expanded through new habitats and acquired *HLA* haplotypes from rainforest hunter-gatherer pygmies (Patin et al. 2017). Our findings may fit with recent work identifying a second wave of Denisovan-like admixture that is specific to East Asian populations (Browning et al. 2018). Thus, although we show the *HLA-B*46:01* and -*B*58:01* haplotypes were obtained by the Han from neighboring modern human populations, they were likely to have been acquired by those populations as a consequence of admixture with archaic humans.

Complementing the high number of HLA class I ligands, we find that in the Chinese Southern Han the number of inhibitory KIR is increased relative to other groups. These KIR allotypes are distinguished by their high expression, inhibitory strength and fine specificity for ligand (Boudreau et al. 2017; Hilton et al. 2015a; Saunders et al. 2016). Possessing higher numbers of inhibitory KIR leads to better effector function, and a higher number of inhibitory KIR ligands leads to larger numbers of circulating NK cells, stronger killing and greater diversity of the NK cell repertoire (Beziat et al. 2013; Brodin et al. 2009; Yawata et al. 2006). That the number of receptors (Pelak et al. 2011) and ligands (Thons et al. 2017) correlates with infection control, suggests the diverse NK cell repertoires of the Southern Han have likely evolved to combat infectious diseases common or endemic to East Asia. Although it is difficult to identify

the specific pathogen exposure history of the Chinese Southern Han, the most plausible candidates for causing selection pressure are viral infections that have established roles for KIR/HLA interaction during host defense (Abi-Rached et al. 2010; Bashirova et al. 2006). Such pathogens have been shown to be effective drivers of adaptive introgression and natural selection in human populations (Enard and Petrov 2018; Harrison et al. 2019). One example is nasopharyngeal carcinoma (NPC) caused by Epstein-Barr virus. HLA-A*11 offers protection from NPC (Tang et al. 2012), and the interaction of KIR3DL2 with HLA-A*11 is dependent on presentation of peptides derived from EBV (Hansasuta et al. 2004). Influenza is another key candidate, with highly virulent epidemics linked to the combination of dense population, agriculture and industrialization (Cao et al. 2009; Chen et al. 2006). Human specific viral hepatitis infections and arboviruses are also endemic to China and South East Asia, including Japanese encephalitis, Dengue and chikungunya (Bashirova et al. 2006; Khakoo et al. 2004; Naiyer et al. 2017; Petitdemange et al. 2011; Thons et al. 2017; Townsley et al. 2016). Consistent with these observations, *KIR A* has established roles in controlling virus infections (Bashirova et al. 2006; Khakoo et al. 2004), and we recently showed *KIR A* homozygosity protects from leukemia (Deng et al. 2019). Reproduction is also a major driver of selection, where *KIR AA*/*C2*$^+$*HLA-C* genotype is associated with preeclampisia (Parham and Moffett 2013). Thus, the low frequency of C2$^+$HLA in East Asia (Figure S3) likely allows the *KIR A* haplotype to reach high frequency (Nemat-Gorgani et al. 2018). High resolution analysis of KIR and HLA diversity will be critical for understanding these and other complex diseases.

In conclusion, our high-resolution analysis of KIR and HLA class I combinatorial diversity has uncovered a distinctive enhancement of the interactions between inhibitory KIR and HLA-A and -B in East Asians. These genetically determined distinctions likely underlie differences across human populations in their susceptibility to infections and immune-mediated diseases.

Supplemental Data

Supplemental Data include four figures and three Excel spreadsheets.

Acknowledgements

This study was supported by the Science, Technology and Innovation Commission of Shenzhen

Municipality (grant number: JCYJ20190806152001762 to ZD), the National Natural Science

Foundation of China (grant number: 81373158 to ZD) and the National Institutes of Health of

the USA (grant numbers: NIH R01 AI017892 to PP, and R56 AI151549 to PJN). We thank the

Chinese blood donors for generously providing DNA samples for this study.

Web Resources and Accession Numbers

The URLs for data, material and programs used herein are as follows:

The scripts used in the study are located at https://github.com/n0rmski/Han_Study/

ImmunoPolymorphism database (IPD), http://www.ebi.ac.uk/ipd/

International Histocompatibility Working Group (IHWG), www.ihwg.org/

The official IPD names (Robinson et al. 2015) and GenBank accession numbers for the *KIR*

sequences reported in this paper are:

(*KIR* prefix excluded for brevity) *2DL1*00304* (KT438851), *2DL1*00305* (KT438852),

*2DL1*030* (KP025959), *2DL1*031* (KP025960), *2DL1*033* (KT438853), *2DL1*034*

(KT438854), *2DL2*013* (KM017076), *2DL3*00109* (KF766495), *2DL3*00110* (KF766497),

*2DL3*025* (KF766496), *2DL3*026* (KF766498), *2DL3*027* (KF766499), *2DL3*028*

(KF766500), *2DL3*029* (KF766501), *2DL3*031* (KF849247), *2DL4*00503* (KT438855),

*2DL4*00504* (KT438856), *2DL4*032* (KT438858), *2DL4*033* (KT438859), *2DL4*034*

(KT438857), *2DL5A*022* (KT438863), *2DS2*009* (KT438862), *2DS4*00105* (KP025962),

*2DS4*017* (KP025961), *2DS4*018* (KP025963), *3DL1*01505* (KF849249), *3DL1*079*

(KF849250), *3DL2*00706* (KT899864), *3DL2*00707* (KT899868), *3DL2*083* (KT899867),

3DL2*084 (KT438861), *3DL2*091* (KT438860), *3DL2*093* (KT899866), *3DL2*099*

(KT899865), *3DL3*01003* (KU529275), *3DL3*02602* (KU529271), *3DL3*04802*

(KU529269), *3DL3*062* (KU529272), *3DL3*063* (KU529270), *3DL3*064* (KU529273),

3DL3*065 (KU529274), *3DS1*078* (KJ001806), *3DS1*082* (KJ001804), *3DS1*083*

(KJ001805), *3DS1*084* (KJ001807), *3DS1*085* (KJ365317).

Figure Legends

Figure 1. Chinese Southern Han *HLA class I* haplotypes express multiple KIR ligands

A. Pie charts show the frequency spectra for HLA-A, -B and -C allotypes of the Southern Han

cohort of 306 unrelated individuals (2N=612). Each pie segment represents one allotype.

Alternative sequence motifs in the α1 domain of the HLA class I molecule determine the four

epitopes recognized by different KIR, and which are also called KIR ligands. The A3/11

epitope is carried by HLA-A3 and -A11 (yellow colored pie segments); the Bw4 epitope is

carried by subsets of HLA-A and -B allotypes (green colored pie segments). The C1 epitope is

carried by a majority of HLA-C allotypes, as well as by HLA-B*46 and HLA-B*73 (red

colored pie segments). The C2 epitope is carried by all HLA-C allotypes that do not carry C1

(blue-colored pie segments). Grey-colored pie segments correspond to allotypes that are not

KIR ligands. Figure S2 lists all the HLA-A, -B and -C allotypes present in the study population

and shows which KIR ligand motifs they carry.

B. Shows the ten most frequent *HLA class I* haplotypes in the Southern Han and their

frequencies (2N=612). Colored shading indicates *HLA class I* alleles that encode KIR ligands,

as described in panel A.

C. (left) Bars show the combined frequencies of *HLA class I* haplotypes encoding one (blue),

two (gold) or three (green) KIR ligands in eight representative populations worldwide

(Southern, Western and Eastern Africa, Europe, Oceania, South America, Japan and Chinese

Southern Han). (Right) Heat-plot shows pairwise comparisons between populations of the

proportion of *HLA class I* haplotypes encoding one KIR ligand to those carrying two or more

KIR ligands. Colors correspond to -log^{10} of a Benjamini-Hochberg corrected p, as shown in the

key.

D. Heat-plot shows pairwise comparisons of Chinese Southern Han with five East/ South East Asian populations of the proportion of *HLA* haplotypes encoding one KIR ligand to those carrying two or more KIR ligands. Colors correspond to -$\log_{10}$ of a Benjamini-Hochberg corrected p, as shown in the key.

E. Venn diagrams show the distribution of *HLA class I* haplotypes within representative subsets of populations. The number of haplotypes in each overlapping region is given. The % values indicate the combined frequency of haplotypes unique to a population when compared to the other populations in the diagram.

Figure 2. HLA-B*46:01 and -B*58:01 were acquired by admixture into the Chinese Southern Han

A. Shown are the relative proportions of genetic ancestry among Asian populations from the 1000 genomes project, plotted by considering three ancestral groups (K = 3: Japanese (green), East Asian (red) and South East Asian (blue)).

B. Shown are the relative proportions for each of the three genetic ancestries in chromosome 6 (left) and within the *MHC* (right) for selected Chinese Southern Han individuals, shown from left to right; all individuals, *B*40:01* carriers, *B*46:01* carriers, *B*58:01* carriers.

C. Shown for each of the ten most frequent *HLA class I* haplotypes in the Chinese Southern Han is a comparison of mean admixture proportion of the genetic ancestry group that is most abundant in *MHC* compared to the proportion of that ancestry in chromosome 6 with *MHC* excluded. The size of the circle represents the -$\log_{10}$ (Benjamini-Hochberg corrected p) from a Wilcoxon test. The difference in size and shade between the two circles corresponds to extent

and direction, respectively, of any shift in the genetic ancestry proportions between the *MHC*

and the remainder of the chromosome 6.

Figure 3. Positive selection has targeted *HLA class I* genes in the Chinese Southern Han

A. Shown is the nucleotide diversity (π) of genomic sequence +/- 500kbp of the *HLA-B* and -

C genes for haplotypes containing the specific *HLA-B* alleles, *B*40*, *B*46* and *B*58*. π was

measured in windows of 100bp. *** p <0.001 by Wilcoxon test.

B. Manhattan plot shows the absolute iHS values above the 95th percentile calculated for

independent SNPs throughout chromosome 6 in Chinese Southern Han. The *MHC* region is

boxed.

C. Density plots show the distribution of absolute iHS values for chromosome 6 (top left, grey

shading), and for SNPs unique to haplotypes carrying specific *HLA-B* (left, cyan), *HLA-A*

(center, purple) and *HLA-C* (right, orange) alleles, as indicated in each plot. The number of

SNPs unique to each of the *HLA* alleles is shown in brackets next to the allele name. For each

allele, the distribution of iHS values was compared to the distribution on chromosome 6 using

a Wilcoxon two-sample test (* $p < 0.05$, ** $< 0.05^{-5}$, *** $< 0.05^{-10}$). Grey dashed line marks the

95th percentile of iHS values for chromosome 6 SNPs (=1.93). The density shown is the kernel

density estimate of the SNP counts associated with the distribution of absolute iHS values.

D. Shows cluster analysis of *HLA class I* allele frequencies from 26 populations. Vertical lines

at the left show the clusters identified when five groups were specified in the input parameters

(k=5) and the support (%) from 1,000 bootstrap replicates. Population names in green text

indicate sub-Saharan African populations, red text – East Asian, brown text – Northeast Asian

(Tuva) and South Asian (Indian), blue text – Oceanic, purple text – Amerindian (Yupik are

North Amerindian who back-migrated to Siberia (Raghavan et al. 2015), NaDene are North American). J. Hondo are Japanese from the major islands of Japan, J. Ryukyu are Japanese from Okinawa. The *HLA class I* haplotypes detected in each population are described in Figure S4D.

E. Shows cluster analysis of the combined frequencies of *HLA class I* haplotypes carrying 1, 2 or 3 KIR ligands. Vertical lines at the left show the clusters identified when three groups were specified (k=3) and the support (%) from 1,000 bootstrap replicates.

Figure 4. Directional selection on *centromeric KIR* in Southern Han

A. Shows the number of *KIR* alleles present in the Southern Han compared with other populations analyzed at comparable resolution; 75 individuals were selected at random from each population.

B. Shown are the combined frequencies of *KIR A* (red) and *KIR B* (blue) haplotypes, for the complete haplotypes, and for the *centromeric* and *telomeric* regions.

C. Shown are the ten most frequent complete, high-resolution, *KIR* haplotypes identified in the Chinese Southern Han population. *KIR A* haplotypes are shaded in red, *KIR B* haplotypes are shaded in blue. At the right is shown for each haplotype the number of individuals carrying the haplotype, and its frequency. All the haplotypes are shown in Figure S7.

D. Bar graph shows a summary *KIR* allele frequencies. The colors from blue to red correspond to the rank in frequency from highest (blue) to lowest (red). Full frequency distributions are shown in Figure S5.

E. Shown are normalized deviate values of Ewens-Watterson's F test (F_{nd}) in representative global populations. Positive values of Fnd indicate homozygosity, negative values indicate heterozygosity. An asterisk denotes significance ($P < 0.05$ or > 0.95) using the exact test (Salamon et al. 1999).

Figure 5. East Asians have a greater diversity of KIR interactions with HLA-A and -B than other populations.

A. Plot of the number of different interacting ligand/receptor allotype pairs observed per individual in the Southern Han.

B. Shows the mean number of different ligand/receptor allotype pairs per individual (left) and the mean total number of ligand/receptor allotype pairs per individual (right) for HLA-C (upper) and HLA-A and -B combined (lower). In the populations shown, *KIR* and *HLA class I* have been analyzed to similar high-resolution as described here for Southern Han. These populations comprise the Yucpa (Gendzekhadze et al. 2009), Japanese (Yawata et al. 2006), Māori (Nemat-Gorgani et al. 2014), European (Vierra-Green et al. 2012), Ghanaian (Norman et al. 2013) Nama (Nemat-Gorgani et al. 2018). Error bars are s.e.m. and p values are from a t-test.

References

Abdulla MA, Ahmed I, Assawamakin A, Bhak J, Brahmachari SK, Calacal GC, Chaurasia A, Chen CH, Chen J, Chen YT, Chu J, Cutiongco-de la Paz EM, De Ungria MC, Delfin FC, Edo J, Fuchareon S, Ghang H, Gojobori T, Han J, Ho SF, Hoh BP, Huang W, Inoko H, Jha P, Jinam TA, Jin L, Jung J, Kangwanpong D, Kampuansai J, Kennedy GC, Khurana P, Kim HL, Kim K, Kim S, Kim WY, Kimm K, Kimura R, Koike T, Kulawonganunchai S, Kumar V, Lai PS, Lee JY, Lee S, Liu ET, Majumder PP, Mandapati KK, Marzuki S, Mitchell W, Mukerji M, Naritomi K, Ngamphiw C, Niikawa N, Nishida N, Oh B, Oh S, Ohashi J, Oka A, Ong R, Padilla CD, Palittapongarnpim P, Perdigon HB, Phipps ME, Png E, Sakaki Y, Salvador JM, Sandraling Y, Scaria V, Seielstad M, Sidek MR, Sinha A, Srikummool M, Sudoyo H, Sugano S, Suryadi H, Suzuki Y, Tabbada KA, Tan A, Tokunaga K, Tongsima S, Villamor LP, Wang E, Wang Y, Wang H, Wu JY, Xiao H, Xu S, Yang JO, Shugart YY, Yoo HS, Yuan W, Zhao G, Zilfalil BA, Indian Genome Variation C (2009) Mapping human genetic diversity in Asia. Science 326:1541-5

Abi-Rached L, Jobin MJ, Kulkarni S, McWhinnie A, Dalva K, Gragert L, Babrzadeh F, Gharizadeh B, Luo M, Plummer FA, Kimani J, Carrington M, Middleton D, Rajalingam R, Beksac M, Marsh SG, Maiers M, Guethlein LA, Tavoularis S, Little AM, Green RE, Norman PJ, Parham P (2011) The shaping of modern human immune systems by multiregional admixture with archaic humans. Science 334:89-94

Abi-Rached L, Moesta AK, Rajalingam R, Guethlein LA, Parham P (2010) Human-specific evolution and adaptation led to major qualitative differences in the variable receptors of human and chimpanzee natural killer cells. PLoS Genet 6:e1001192

Alexander DH, Novembre J, Lange K (2009) Fast model-based estimation of ancestry in unrelated individuals. Genome Res 19:1655-64

Andersson S, Fauriat C, Malmberg JA, Ljunggren HG, Malmberg KJ (2009) KIR acquisition probabilities are independent of self-HLA class I ligands and increase with cellular KIR expression. Blood 114:95-104

Augusto DG, Norman PJ, Dandekar R, Hollenbach JA (2019) Fluctuating and Geographically Specific Selection Characterize Rapid Evolution of the Human KIR Region. Front Immunol 10:989

Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR (2015) A global reference for human genetic variation. Nature 526:68-74

Bao X, Wang M, Zhou H, Wu X, Yang L, Xu C, Yuan X, Zhang J, Li L, Wu D, He J (2013) Characterization of Killer cell immunoglobulin-like receptor (KIR) genotypes and haplotypes in Chinese Han population. Tissue Antigens 82:327-37

Bao X, Wang M, Zhou H, Zhang H, Wu X, Yuan X, Li Y, Wu D, He J (2016) Donor Killer Immunoglobulin-Like Receptor Profile Bx1 Imparts a Negative Effect and Centromeric B-Specific Gene Motifs Render a Positive Effect on Standard-Risk Acute Myeloid Leukemia/Myelodysplastic Syndrome Patient Survival after Unrelated Donor Hematopoietic Stem Cell Transplantation. Biol Blood Marrow Transplant 22:232-239

Bari R, Bell T, Leung WH, Vong QP, Chan WK, Das Gupta N, Holladay M, Rooney B, Leung W (2009) Significant functional heterogeneity among KIR2DL1 alleles and a pivotal role of arginine 245. Blood 114:5182-90

Bashirova AA, Martin MP, McVicar DW, Carrington M (2006) The killer immunoglobulin-like receptor gene cluster: tuning the genome for defense. Annu Rev Genomics Hum Genet 7:277-300

Beziat V, Traherne JA, Liu LL, Jayaraman J, Enqvist M, Larsson S, Trowsdale J, Malmberg KJ (2013) Influence of KIR gene copy number on natural killer cell education. Blood 121:4703-7

Bjorkstrom NK, Ljunggren HG, Michaelsson J (2016) Emerging insights into natural killer cells in human peripheral tissues. Nat Rev Immunol 16:310-20

Blokhuis JH, Hilton HG, Guethlein LA, Norman PJ, Nemat-Gorgani N, Nakimuli A, Chazara O, Moffett A, Parham P (2017) KIR2DS5 allotypes that recognize the C2 epitope of HLA-C are common among Africans and absent from Europeans. Immun Inflamm Dis

Boelen L, Debebe B, Silveira M, Salam A, Makinde J, Roberts CH, Wang ECY, Frater J, Gilmour J, Twigger K, Ladell K, Miners KL, Jayaraman J, Traherne JA, Price DA, Qi Y, Martin MP, Macallan DC, Thio CL, Astemborski J, Kirk G, Donfield SM, Buchbinder S, Khakoo SI, Goedert JJ, Trowsdale J, Carrington M, Kollnberger S, Asquith B (2018) Inhibitory killer cell immunoglobulin-like receptors strengthen CD8(+) T cell-mediated control of HIV-1, HCV, and HTLV-1. Sci Immunol 3

Boudreau JE, Giglio F, Gooley TA, Stevenson PA, Le Luduec JB, Shaffer BC, Rajalingam R, Hou L, Hurley CK, Noreen H, Reed EF, Yu N, Vierra-Green C, Haagenson M, Malkki M, Petersdorf EW, Spellman S, Hsu KC (2017) KIR3DL1/HLA-B Subtypes Govern Acute Myelogenous Leukemia Relapse After Hematopoietic Cell Transplantation. J Clin Oncol 35:2268-2278

Boudreau JE, Hsu KC (2018) Natural Killer Cell Education and the Response to Infection and Cancer Therapy: Stay Tuned. Trends Immunol

Brodin P, Lakshmikanth T, Johansson S, Karre K, Hoglund P (2009) The strength of inhibitory input during education quantitatively tunes the functional responsiveness of individual natural killer cells. Blood 113:2434-41

Browning SR, Browning BL, Zhou Y, Tucci S, Akey JM (2018) Analysis of Human Sequence Data Reveals Two

Pulses of Archaic Denisovan Admixture. Cell

Burian A, Wang KL, Finton KA, Lee N, Ishitani A, Strong RK, Geraghty DE (2016) HLA-F and MHC-I Open Conformers Bind Natural Killer Cell Ig-Like Receptor KIR3DS1. PLoS One 11:e0163297

Busby G, Christ R, Band G, Leffler E, Le QS, Rockett K, Kwiatkowski D, Spencer C (2017) Inferring adaptive gene-flow in recent African history. bioRxiv:205252

Campbell MC, Tishkoff SA (2008) African genetic diversity: implications for human demographic history, modern human origins, and complex disease mapping. Annu Rev Genomics Hum Genet 9:403-33

Cao B, Li XW, Mao Y, Wang J, Lu HZ, Chen YS, Liang ZA, Liang L, Zhang SJ, Zhang B, Gu L, Lu LH, Wang DY, Wang C (2009) Clinical features of the initial cases of 2009 pandemic influenza A (H1N1) virus infection in China. N Engl J Med 361:2507-17

Chen CH, Yang JH, Chiang CWK, Hsiung CN, Wu PE, Chang LC, Chu HW, Chang J, Song IW, Yang SL, Chen YT, Liu FT, Shen CY (2016) Population structure of Han Chinese in the modern Taiwanese population based on 10,000 participants in the Taiwan Biobank project. Hum Mol Genet 25:5321-5331

Chen H, Smith GJ, Li KS, Wang J, Fan XH, Rayner JM, Vijaykrishna D, Zhang JX, Zhang LJ, Guo CT, Cheung CL, Xu KM, Duan L, Huang K, Qin K, Leung YH, Wu WL, Lu HR, Chen Y, Xia NS, Naipospos TS, Yuen KY, Hassan SS, Bahri S, Nguyen TD, Webster RG, Peiris JS, Guan Y (2006) Establishment of multiple sublineages of H5N1 influenza virus in Asia: implications for pandemic control. Proc Natl Acad Sci U S A 103:2845-50

Chen R, Zhang GB, Deng ZH (2017) Description of the novel KIR2DL1*034 allele Identified from a southern Chinese Han individual. HLA 89:121-122

Cooper MA, Colonna M, Yokoyama WM (2009) Hidden talents of natural killers: NK cells in innate and adaptive immunity. EMBO Rep 10:1103-10

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G, Durbin R (2011) The variant call format and VCFtools. Bioinformatics 27:2156-8

Dendrou CA, Petersen J, Rossjohn J, Fugger L (2018) HLA variation and disease. Nat Rev Immunol 18:325-339

Deng Z, Zhao J, Cai S, Qi Y, Yu Q, Martin MP, Gao X, Chen R, Zhuo J, Zhen J, Zhang M, Zhang G, He L, Zou H, Lu L, Zhu W, Hong W, Carrington M, Norman PJ (2019) Natural Killer Cells Offer Differential Protection From Leukemia in Chinese Southern Han. Front Immunol 10:1646

Deng Z, Zhen J, Zhang G (2018) METHOD FOR SIMULTANEOUS SEQUENCE-BASED TYPING OF 14 FUNCTIONAL KILLER CELL IMMUNOGLOBULIN-LIKE RECEPTOR (KIR) GENES In Patent US (ed.). USA

Deng Z, Zhen J, Zhu B, Zhang G, Yu Q, Wang D, Xu Y, He L, Lu L (2015) Allelic diversity of KIR3DL1/3DS1 in a southern Chinese population. Hum Immunol 76:663-6

Deng ZH, Zhang GB, Chen R (2017) Description of the novel KIR3DL3*04802 allele identified in a southern Chinese Han individual. Hla 89:66-67

Dohring C, Scheidegger D, Samaridis J, Cella M, Colonna M (1996) A human killer inhibitory receptor specific for HLA-A1,2. J Immunol 156:3098-101

Enard D, Petrov DA (2018) Evidence that RNA Viruses Drove Adaptive Introgression between Neanderthals and Modern Humans. Cell 175:360-371.e13

Excoffier L, Lischer HE (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. Mol Ecol Resour 10:564-7

Fagundes NJ, Kanitz R, Eckert R, Valls AC, Bogo MR, Salzano FM, Smith DG, Silva WA, Jr., Zago MA, Ribeiro-dos-Santos AK, Santos SE, Petzl-Erler ML, Bonatto SL (2008) Mitochondrial population genomics supports a single pre-Clovis origin with a coastal route for the peopling of the Americas. Am J Hum Genet 82:583-92

Freud AG, Mundy-Bosse BL, Yu J, Caligiuri MA (2017) The Broad Spectrum of Human Natural Killer Cell Diversity. Immunity 47:820-833

Garcia-Beltran WF, Holzemer A, Martrus G, Chung AW, Pacheco Y, Simoneau CR, Rucevic M, Lamothe-Molina PA, Pertel T, Kim TE, Dugan H, Alter G, Dechanet-Merville J, Jost S, Carrington M, Altfeld M (2016) Open conformers of HLA-F are high-affinity ligands of the activating NK-cell receptor KIR3DS1. Nat Immunol 17:1067-74

Gendzekhadze K, Norman PJ, Abi-Rached L, Graef T, Moesta AK, Layrisse Z, Parham P (2009) Co-evolution of KIR2DL3 with HLA-C in a human population retaining minimal essential diversity of KIR and HLA class I ligands. Proc Natl Acad Sci U S A 106:18692-7

Gonzalez-Galarza FF, Takeshita LY, Santos EJ, Kempson F, Maia MH, da Silva AL, Teles e Silva AL, Ghattaoraya GS, Alfirevic A, Jones AR, Middleton D (2015) Allele frequency net 2015 update: new features for HLA epitopes, KIR and disease and HLA adverse drug reaction associations. Nucleic Acids Res 43:D784-8

Gourraud PA, Khankhanian P, Cereb N, Yang SY, Feolo M, Maiers M, Rioux JD, Hauser S, Oksenberg J (2014) HLA diversity in the 1000 genomes dataset. PLoS One 9:e97282

Graef T, Moesta AK, Norman PJ, Abi-Rached L, Vago L, Older Aguilar AM, Gleimer M, Hammond JA, Guethlein LA, Bushnell DA, Robinson PJ, Parham P (2009) KIR2DS4 is a product of gene conversion with KIR3DL2 that introduced specificity for HLA-A*11 while diminishing avidity for HLA-C. J Exp Med 206:2557-72

Guethlein LA, Norman PJ, Hilton HH, Parham P (2015) Co-evolution of MHC class I and variable NK cell receptors in placental mammals. Immunol Rev 267:259-82

Hansasuta P, Dong T, Thananchai H, Weekes M, Willberg C, Aldemir H, Rowland-Jones S, Braud VM (2004) Recognition of HLA-A3 and HLA-A11 by KIR3DL2 is peptide-specific. Eur J Immunol 34:1673-9

Harrison GF, Sanz J, Boulais J, Mina MJ, Grenier JC, Leng Y, Dumaine A, Yotova V, Bergey CM, Nsobya SL, Elledge SJ, Schurr E, Quintana-Murci L, Perry GH, Barreiro LB (2019) Natural selection contributed to immunological differences between hunter-gatherers and agriculturalists. Nat Ecol Evol 3:1253-1264

He LM, Deng ZH, Hu HY, Zhen JX, Liu ZH (2017) Identification and characterization of the novel KIR2DL1*030 allele by sequence-based typing in a southern Chinese Han individual. HLA 89:263-264

Hellenthal G, Busby GBJ, Band G, Wilson JF, Capelli C, Falush D, Myers S (2014) A genetic atlas of human admixture history. Science 343:747-751

Henn BM, Cavalli-Sforza LL, Feldman MW (2012) The great human expansion. Proc Natl Acad Sci U S A 109:17758-64

Henn BM, Gignoux CR, Jobin M, Granka JM, Macpherson JM, Kidd JM, Rodriguez-Botigue L, Ramachandran S, Hon L, Brisbin A, Lin AA, Underhill PA, Comas D, Kidd KK, Norman PJ, Parham P, Bustamante CD, Mountain JL, Feldman MW (2011) Hunter-gatherer genomic diversity suggests a southern African origin for modern humans. Proc Natl Acad Sci U S A 108:5154-62

Hennig C (2020) fpc: Flexible Procedures for Clustering. R package version 2.2-7. https://CRAN.R-project.org/package=fpcc.

Hilton HG, Guethlein LA, Goyos A, Nemat-Gorgani N, Bushnell DA, Norman PJ, Parham P (2015a) Polymorphic HLA-C Receptors Balance the Functional Characteristics of KIR Haplotypes. J Immunol 195:3160-70

Hilton HG, Norman PJ, Nemat-Gorgani N, Goyos A, Hollenbach JA, Henn BM, Gignoux CR, Guethlein LA, Parham P (2015b) Loss and Gain of Natural Killer Cell Receptor Function in an African Hunter-Gatherer Population. PLoS Genet 11:e1005439

Hoglund P, Brodin P (2010) Current perspectives of natural killer cell education by MHC class I molecules. Nat Rev Immunol 10:724-34

Holzemer A, Garcia-Beltran WF, Altfeld M (2017) Natural Killer Cell Interactions with Classical and Non-Classical Human Leukocyte Antigen Class I in HIV-1 Infection. Front Immunol 8:1496

Horowitz A, Djaoud Z, Nemat-Gorgani N, Blokhuis J, Hilton HG, Beziat V, Malmberg KJ, Norman PJ, Guethlein LA, Parham P (2016) Class I HLA haplotypes form two schools that educate NK cells in different ways. Sci Immunol 1:eaag1672

Jiang Y, Chen O, Cui C, Zhao B, Han X, Zhang Z, Liu J, Xu J, Hu Q, Liao C, Shang H (2013) KIR3DS1/L1 and HLA-Bw4-80I are associated with HIV disease progression among HIV typical progressors and long-term nonprogressors. BMC Infect Dis 13:405

Khakoo SI, Thio CL, Martin MP, Brooks CR, Gao X, Astemborski J, Cheng J, Goedert JJ, Vlahov D, Hilgartner M, Cox S, Little AM, Alexander GJ, Cramp ME, O'Brien SJ, Rosenberg WM, Thomas DL, Carrington M (2004) HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. Science 305:872-4

Lancaster AK, Single RM, Solberg OD, Nelson MP, Thomson G (2007) PyPop update--a software pipeline for large-scale multilocus population genomics. Tissue Antigens 69 Suppl 1:192-7

Liu J, Xiao Z, Ko HL, Shen M, Ren EC (2014) Activating killer cell immunoglobulin-like receptor 2DS2 binds to HLA-A*11. Proc Natl Acad Sci U S A 111:2662-7

Loh P-R, Palamara PF, Price AL (2016) Fast and accurate long-range phasing in a UK Biobank cohort. Nature genetics 48:811

Long EO, Kim HS, Liu D, Peterson ME, Rajagopalan S (2013) Controlling natural killer cell responses: integration of signals for activation and inhibition. Annu Rev Immunol 31:227-58

Long W, Shi Z, Fan S, Liu L, Lu Y, Guo X, Rong C, Cui X, Ding H (2015) Association of maternal KIR and fetal HLA-C genes with the risk of preeclampsia in the Chinese Han population. Placenta 36:433-7

Lu D, Lou H, Yuan K, Wang X, Wang Y, Zhang C, Lu Y, Yang X, Deng L, Zhou Y, Feng Q, Hu Y, Ding Q, Yang Y, Li S, Jin L, Guan Y, Su B, Kang L, Xu S (2016) Ancestral Origins and Genetic History of Tibetan Highlanders. Am J Hum Genet 99:580-594

Meenagh A, Gonzalez A, Sleator C, McQuaid S, Middleton D (2008) Investigation of killer cell immunoglobulin-like receptor gene diversity, KIR2DL1 and KIR2DS1. Tissue Antigens 72:383-91

Meyer D, Single RM, Mack SJ, Lancaster A, Nelson MP, Erlich HA, Fernandez-Vina M, Thomson G (2007) Single Locus Polymorphism of Classical HLA Genes. *In* Hansen JA (ed.) Immunobiology of the Human MHC: Proceedings of the 13th International Histocompatibility Workshop and Conference. IHWG press, Seattle, WA

Moesta AK, Graef T, Abi-Rached L, Older Aguilar AM, Guethlein LA, Parham P (2010) Humans differ from other hominids in lacking an activating NK cell receptor that recognizes the C1 epitope of MHC class I. J Immunol 185:4233-7

Moesta AK, Norman PJ, Yawata M, Yawata N, Gleimer M, Parham P (2008) Synergistic polymorphism at two positions distal to the ligand-binding site makes KIR2DL2 a stronger receptor for HLA-C than KIR2DL3. J Immunol 180:3969-79

Naiyer MM, Cassidy SA, Magri A, Cowton V, Chen K, Mansour S, Kranidioti H, Mbirbindi B, Rettman P, Harris
S, Fanning LJ, Mulder A, Claas FHJ, Davidson AD, Patel AH, Purbhoo MA, Khakoo SI (2017) KIR2DS2
recognizes conserved peptides derived from viral helicases in the context of HLA-C. Sci Immunol 2
Nei M, Takahata N (1993) Effective population size, genetic diversity, and coalescence time in subdivided
populations. Journal of Molecular Evolution 37:240-244
Nemat-Gorgani N, Edinur HA, Hollenbach JA, Traherne JA, Dunn PP, Chambers GK, Parham P, Norman PJ
(2014) KIR diversity in Maori and Polynesians: populations in which HLA-B is not a significant KIR
ligand. Immunogenetics 66:597-611
Nemat-Gorgani N, Hilton HG, Henn BM, Lin M, Gignoux CR, Myrick JW, Werely CJ, Granka JM, Moller M,
Hoal EG, Yawata M, Yawata N, Boelen L, Asquith B, Parham P, Norman PJ (2018) Different Selected
Mechanisms Attenuated the Inhibitory Interaction of KIR2DL1 with C2(+) HLA-C in Two Indigenous
Human Populations in Southern Africa. J Immunol
Norman PJ, Abi-Rached L, Gendzekhadze K, Hammond JA, Moesta AK, Sharma D, Graef T, McQueen KL,
Guethlein LA, Carrington CV, Chandanayingyong D, Chang YH, Crespi C, Saruhan-Direskeneli G,
Hameed K, Kamkamidze G, Koram KA, Layrisse Z, Matamoros N, Mila J, Park MH, Pitchappan RM,
Ramdath DD, Shiau MY, Stephens HA, Struik S, Tyan D, Verity DH, Vaughan RW, Davis RW, Fraser
PA, Riley EM, Ronaghi M, Parham P (2009) Meiotic recombination generates rich diversity in NK cell
receptor genes, alleles, and haplotypes. Genome Res 19:757-69
Norman PJ, Abi-Rached L, Gendzekhadze K, Korbel D, Gleimer M, Rowley D, Bruno D, Carrington CV,
Chandanayingyong D, Chang YH, Crespi C, Saruhan-Direskeneli G, Fraser PA, Hameed K,
Kamkamidze G, Koram KA, Layrisse Z, Matamoros N, Mila J, Park MH, Pitchappan RM, Ramdath DD,
Shiau MY, Stephens HA, Struik S, Verity DH, Vaughan RW, Tyan D, Davis RW, Riley EM, Ronaghi M,
Parham P (2007) Unusual selection on the KIR3DL1/S1 natural killer cell receptor in Africans. Nat Genet
39:1092-9
Norman PJ, Hollenbach JA, Nemat-Gorgani N, Guethlein LA, Hilton HG, Pando MJ, Koram KA, Riley EM, Abi-
Rached L, Parham P (2013) Co-evolution of Human Leukocyte Antigen (HLA) Class I Ligands with
Killer-Cell Immunoglobulin-Like Receptors (KIR) in a Genetically Diverse Population of Sub-Saharan
Africans. PLoS Genet 9:e1003938
Norman PJ, Hollenbach JA, Nemat-Gorgani N, Marin WM, Norberg SJ, Ashouri E, Jayaraman J, Wroblewski EE,
Trowsdale J, Rajalingam R, Oksenberg JR, Chiaroni J, Guethlein LA, Traherne JA, Ronaghi M, Parham
P (2016) Defining KIR and HLA Class I Genotypes at Highest Resolution via High-Throughput
Sequencing. Am J Hum Genet 99:375-91
Pando MJ, Gardiner CM, Gleimer M, McQueen KL, Parham P (2003) The protein made from a common allele of
KIR3DL1 (3DL1*004) is poorly expressed at cell surfaces due to substitution at positions 86 in Ig
domain 0 and 182 in Ig domain 1. J Immunol 171:6640-9
Parham P, Moffett A (2013) Variable NK cell receptors and their MHC class I ligands in immunity, reproduction
and human evolution. Nat Rev Immunol 13:133-44
Patin E, Lopez M, Grollemund R, Verdu P, Harmant C, Quach H, Laval G, Perry GH, Barreiro LB, Froment A,
Heyer E, Massougbodji A, Fortes-Lima C, Migot-Nabias F, Bellis G, Dugoujon JM, Pereira JB,
Fernandes V, Pereira L, Van der Veen L, Mouguiama-Daouda P, Bustamante CD, Hombert JM, Quintana-
Murci L (2017) Dispersals and genetic adaptation of Bantu-speaking populations in Africa and North
America. Science 356:543-546
Pelak K, Need AC, Fellay J, Shianna KV, Feng S, Urban TJ, Ge D, De Luca A, Martinez-Picado J, Wolinsky SM,
Martinson JJ, Jamieson BD, Bream JH, Martin MP, Borrow P, Letvin NL, McMichael AJ, Haynes BF,
Telenti A, Carrington M, Goldstein DB, Alter G, Immunology NCfHAV (2011) Copy number variation
of KIR genes influences HIV-1 control. PLoS Biol 9:e1001208
Petitdemange C, Becquart P, Wauquier N, Beziat V, Debre P, Leroy EM, Vieillard V (2011) Unconventional
repertoire profile is imprinted during acute chikungunya infection for natural killer cells polarization
toward cytotoxicity. PLoS Pathog 7:e1002268
Prugnolle F, Manica A, Charpentier M, Guégan JF, Guernier V, Balloux F (2005) Pathogen-driven selection and
worldwide HLA class I diversity. Curr Biol 15:1022-7
Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ,
Sham PC (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses.
Am J Hum Genet 81:559-75
R Development Core Team (2008) A language and environment for statistical computing. R Foundation for
Statistical Computing. Vienna, Austria
Raghavan M, Steinrucken M, Harris K, Schiffels S, Rasmussen S, DeGiorgio M, Albrechtsen A, Valdiosera C,
Avila-Arcos MC, Malaspinas AS, Eriksson A, Moltke I, Metspalu M, Homburger JR, Wall J, Cornejo
OE, Moreno-Mayar JV, Korneliussen TS, Pierre T, Rasmussen M, Campos PF, Damgaard Pde B,
Allentoft ME, Lindo J, Metspalu E, Rodriguez-Varela R, Mansilla J, Henrickson C, Seguin-Orlando A,
Malmstrom H, Stafford T, Jr., Shringarpure SS, Moreno-Estrada A, Karmin M, Tambets K, Bergstrom A,
Xue Y, Warmuth V, Friend AD, Singarayer J, Valdes P, Balloux F, Leboreiro I, Vera JL, Rangel-Villalobos
H, Pettener D, Luiselli D, Davis LG, Heyer E, Zollikofer CP, Ponce de Leon MS, Smith CI, Grimes V,

Pike KA, Deal M, Fuller BT, Arriaza B, Standen V, Luz MF, Ricaut F, Guidon N, Osipova L, Voevoda MI, Posukh OL, Balanovsky O, Lavryashina M, Bogunov Y, Khusnutdinova E, Gubina M, Balanovska E, Fedorova S, Litvinov S, Malyarchuk B, Derenko M, Mosher MJ, Archer D, Cybulski J, Petzelt B, Mitchell J, Worl R, Norman PJ, Parham P, Kemp BM, Kivisild T, Tyler-Smith C, Sandhu MS, Crawford M, Villems R, Smith DG, Waters MR, Goebel T, Johnson JR, Malhi RS, Jakobsson M, Meltzer DJ, Manica A, Durbin R, Bustamante CD, Song YS, Nielsen R, et al. (2015) POPULATION GENETICS. Genomic evidence for the Pleistocene and recent population history of Native Americans. Science 349:aab3884

Rishishwar L, Conley AB, Wigington CH, Wang L, Valderrama-Aguirre A, Jordan IK (2015) Ancestry, admixture and fitness in Colombian genomes. Sci Rep 5:12376

Robinson J, Halliwell JA, Hayhurst JD, Flicek P, Parham P, Marsh SG (2015) The IPD and IMGT/HLA database: allele variant databases. Nucleic Acids Res 43:D423-31

Rosenberg NA, Pritchard JK, Weber JL, Cann HM, Kidd KK, Zhivotovsky LA, Feldman MW (2002) Genetic structure of human populations. Science 298:2381-5

Salamon H, Klitz W, Easteal S, Gao X, Erlich HA, Fernandez-Vina M, Trachtenberg EA, McWeeney SK, Nelson MP, Thomson G (1999) Evolution of HLA class II molecules: Allelic and amino acid site variability across populations. Genetics 152:393-400

Saunders PM, Pymm P, Pietra G, Hughes VA, Hitchen C, O'Connor GM, Loiacono F, Widjaja J, Price DA, Falco M, Mingari MC, Moretta L, McVicar DW, Rossjohn J, Brooks AG, Vivian JP (2016) Killer cell immunoglobulin-like receptor 3DL1 polymorphism defines distinct hierarchies of HLA class I recognition. J Exp Med 213:791-807

Saunders PM, Vivian JP, O'Connor GM, Sullivan LC, Pymm P, Rossjohn J, Brooks AG (2015) A bird's eye view of NK cell receptor interactions with their MHC class I ligands. Immunol Rev 267:148-66

Schiffels S, Durbin R (2014) Inferring human population size and separation history from multiple genome sequences. Nat Genet 46:919-25

Shen M, Linn YC, Ren EC (2016) KIR-HLA profiling shows presence of higher frequencies of strong inhibitory KIR-ligands among prognostically poor risk AML patients. Immunogenetics 68:133-44

Solberg OD, Mack SJ, Lancaster AK, Single RM, Tsai Y, Sanchez-Mazas A, Thomson G (2008) Balancing selection and heterogeneity across the classical human leukocyte antigen loci: a meta-analytic review of 497 population studies. Hum Immunol 69:443-64

Stephens M, Donnelly P (2003) A comparison of bayesian methods for haplotype reconstruction from population genotype data. Am J Hum Genet 73:1162-9

Su N, Wang H, Zhang B, Kang Y, Guo Q, Xiao H, Yang H, Liao S (2018) Maternal natural killer cell immunoglobulin receptor genes and human leukocyte antigen-C ligands influence recurrent spontaneous abortion in the Han Chinese population. Exp Ther Med 15:327-337

Sun G, Wang C, Zhen J, Zhang G, Xu Y, Deng Z (2016) [Cloning and sequencing of KIR2DL1 framework gene cDNA and identification of a novel allele]. Zhonghua Yi Xue Yi Chuan Xue Za Zhi 33:694-7

Szpiech ZA, Hernandez RD (2014) selscan: an efficient multithreaded program to perform EHH-based scans for positive selection. Mol Biol Evol 31:2824-7

Takeuchi F, Katsuya T, Kimura R, Nabika T, Isomura M, Ohkubo T, Tabara Y, Yamamoto K, Yokota M, Liu X, Saw WY, Mamatyusupu D, Yang W, Xu S, Japanese Genome Variation C, Teo YY, Kato N (2017) The fine-scale genetic structure and evolution of the Japanese population. PLoS One 12:e0185487

Tang M, Lautenberger JA, Gao X, Sezgin E, Hendrickson SL, Troyer JL, David VA, Guan L, McIntosh CE, Guo X, Zheng Y, Liao J, Deng H, Malasky M, Kessing B, Winkler CA, Carrington M, De The G, Zeng Y, O'Brien SJ (2012) The principal genetic determinants for nasopharyngeal carcinoma in China involve the HLA class I antigen recognition groove. PLoS Genet 8:e1003103

Tao SD, He YM, Ying YL, He J, Zhu FM, Lv HJ (2014) KIR3DL1 genetic diversity and phenotypic variation in the Chinese Han population. Genes Immun 15:8-15

Thons C, Senff T, Hydes TJ, Manser AR, Heinemann FM, Heinold A, Heilmann M, Kim AY, Uhrberg M, Scherbaum N, Lauer GM, Khakoo SI, Timm J (2017) HLA-Bw4 80(T) and multiple HLA-Bw4 copies combined with KIR3DL1 associate with spontaneous clearance of HCV infection in people who inject drugs. J Hepatol 67:462-470

Tishkoff SA, Reed FA, Friedlaender FR, Ehret C, Ranciaro A, Froment A, Hirbo JB, Awomoyi AA, Bodo JM, Doumbo O, Ibrahim M, Juma AT, Kotze MJ, Lema G, Moore JH, Mortensen H, Nyambo TB, Omar SA, Powell K, Pretorius GS, Smith MW, Thera MA, Wambebe C, Weber JL, Williams SM (2009) The genetic structure and history of Africans and African Americans. Science 324:1035-44

Townsley E, O'Connor G, Cosgrove C, Woda M, Co M, Thomas SJ, Kalayanarooj S, Yoon IK, Nisalak A, Srikiatkhachorn A, Green S, Stephens HA, Gostick E, Price DA, Carrington M, Alter G, McVicar DW, Rothman AL, Mathew A (2016) Interaction of a dengue virus NS1-derived peptide with the inhibitory receptor KIR3DL1 on natural killer cells. Clin Exp Immunol 183:419-30

VandenBussche CJ, Dakshanamurthy S, Posch PE, Hurley CK (2006) A single polymorphism disrupts the killer Ig-like receptor 2DL2/2DL3 D1 domain. J Immunol 177:5347-57

VandenBussche CJ, Mulrooney TJ, Frazier WR, Dakshanamurthy S, Hurley CK (2009) Dramatically reduced

surface expression of NK cell receptor KIR2DS3 is attributed to multiple residues throughout the molecule. Genes Immun 10:162-73

Vierra-Green C, Roe D, Hou L, Hurley CK, Rajalingam R, Reed E, Lebedeva T, Yu N, Stewart M, Noreen H, Hollenbach JA, Guethlein LA, Wang T, Spellman S, Maiers M (2012) Allele-level haplotype frequencies and pairwise linkage disequilibrium for 14 KIR loci in 506 European-American individuals. PLoS One 7:e47491

Vivier E, Raulet DH, Moretta A, Caligiuri MA, Zitvogel L, Lanier LL, Yokoyama WM, Ugolini S (2011) Innate or adaptive immunity? The example of natural killer cells. Science 331:44-9

Voight BF, Kudaravalli S, Wen X, Pritchard JK (2006) A map of recent positive selection in the human genome. PLoS Biol 4:e72

Wang HD, Zhang FX, Shen CM, Wu YM, Lv YG, Xie ST, Yang G, Qin HX, Fan SL, Zhu BF (2012) The distribution of genetic diversity of KIR genes in the Chinese Mongolian population. Hum Immunol 73:1031-8

Wen B, Li H, Lu D, Song X, Zhang F, He Y, Li F, Gao Y, Mao X, Zhang L, Qian J, Tan J, Jin J, Huang W, Deka R, Su B, Chakraborty R, Jin L (2004) Genetic evidence supports demic diffusion of Han culture. Nature 431:302-5

Wilson MJ, Torkar M, Haude A, Milne S, Jones T, Sheer D, Beck S, Trowsdale J (2000) Plasticity in the organization and sequences of human KIR/ILT gene families. Proc Natl Acad Sci U S A 97:4778-83.

Xu S, Yin X, Li S, Jin W, Lou H, Yang L, Gong X, Wang H, Shen Y, Pan X, He Y, Yang Y, Wang Y, Fu W, An Y, Wang J, Tan J, Qian J, Chen X, Zhang X, Sun Y, Zhang X, Wu B, Jin L (2009) Genomic dissection of population substructure of Han Chinese and its implication in association studies. Am J Hum Genet 85:762-74

Yao Y, Shi L, Tao Y, Lin K, Liu S, Yu L, Yang Z, Yi W, Huang X, Sun H, Chu J, Shi L (2011) Diversity of killer cell immunoglobulin-like receptor genes in four ethnic groups in China. Immunogenetics 63:475-83

Yawata M, Yawata N, Draghi M, Little AM, Partheniou F, Parham P (2006) Roles for HLA and KIR polymorphisms in natural killer cell repertoire selection and modulation of effector function. J Exp Med 203:633-45

Zhang G, Deng Z (2016) [A strategy to clarify ambiguities during genotyping of functional KIR framework genes by sequencing-based typing among ethnic Hans from southern China]. Zhonghua Yi Xue Yi Chuan Xue Za Zhi 33:773-777

Zhang G, Deng Z (2017) [Allelic diversity of KIR2DL4 gene and identification of five novel alleles among southern Han Chinese population]. Zhonghua Yi Xue Yi Chuan Xue Za Zhi 34:270-274

Zhang GB, Chen R, Deng ZH (2017) Identification of the novel KIR2DL1*033 allele from a southern Chinese Han individual by sequencing-based typing. Hla 89:175-176

Zhen J, He L, Xu Y, Zhao J, Yu Q, Zou H, Sun G, Deng Z (2015) Allelic polymorphism of KIR2DL2/2DL3 in a southern Chinese population. Tissue Antigens 86:362-7

Zhen J, Zhang G, Yu Q, He L, Xu Y, Zou H, Deng Z (2017) [Genetic polymorphisms of KIR2DS4 gene among ethnic Hans from southern China]. Zhonghua Yi Xue Yi Chuan Xue Za Zhi 34:21-25

A	HLA class I Allotypes in Chinese Southern Han (2N=612)								
KIR Ligand	HLA-A			HLA-B			HLA-C		
	Total	A3/11	Bw4	Total	Bw4	C1	Total	C1	C2
Number of allotypes	27	6	4	54	18	1	29	21	8
Frequency (%)	-	34.6	17.1	-	34.8	14.2	-	84.8	15.2

B

Ten most frequent *HLA* haplotypes			%
*A*02:07*	*B*46:01*	*C*01:02*	7.68
*A*33:03*	*B*58:01*	*C*03:02*	5.72
*A*11:01*	*B*40:01*	*C*07:02*	4.58
*A*11:01*	*B*13:01*	*C*03:04*	3.27
*A*11:01*	*B*15:02*	*C*08:01*	2.12
*A*30:01*	*B*13:02*	*C*06:02*	1.96
*A*11:01*	*B*46:01*	*C*01:02*	1.96
*A*24:02*	*B*40:01*	*C*03:04*	1.63
*A*24:02*	*B*40:01*	*C*07:02*	1.47
*A*11:01*	*B*40:06*	*C*08:01*	1.47

C

	KIR gene													Southern Han	
	Centromeric KIR						*Telomeric KIR*								
h	3DL3	2DS2	2DL2/3	2DL5B	2DS3/5	2DL1	2DL4	3DL1/S1	2DL5A	2DS3/5	2DS1	2DS4	3DL2	obs.	%
1	*010		3*00101			*00302	*00102	*01502				*00101	*002	71	11.60
2	*008		3*00101			*00302	*00102	*01502				*00101	*002	56	9.15
3	*009		3*00101			*00302	*00102	*01502				*00101	*002	52	8.50
4	*010		3*00101			*00302	*011	*00501				*010	*010	26	4.25
5	*010		3*00101			*00302	*00801	*00101				*00301	*001	12	1.96
6	*010		3*00101			*00302	*00501	S1*01301	A*005	3*00201	*00201		*00701	11	1.80
7	*010		3*023			*00302	*00102	*01502				*00101	*002	10	1.63
8	*010		3*00101			*00302	*00501	S1*01301	A*001	5*00201	*00201		*00701	9	1.47
9	*010		3*00101			*00302	*00103	*020				*00101	*009	9	1.47
10	*010		3*00101			*00302	*00102	*01502				*00101	*039	9	1.47

A

Number of Distinct KIR+HLA class I
Interactions per Individual

B