1   Title: Flexible decision under ambiguous cues reverses

2   neural signaling from occipitotemporal to prefrontal cortex

5   Authors: Kaho Tsumura,[1] Keita Kosugi,[1] Ryuta Aoki,[2] Masaki Takeda,[2]

6   Junichi Chikazoe,[3] Kiyoshi Nakahara,[2] Koji Jimura[1,2]

9   Affiliations: [1] Department of Biosciences and Informatics,

10   Keio University, Yokohama, Japan

11   [2] Research Center for Brain Communication,

12   Kochi University of Technology, Kami, Japan

13   [3] Supportive Center for Brain Research,

14   National Institute for Physiological Sciences, Okazaki, Japan

Correspondence should be addressed to:

Koji Jimura, Ph.D.

Department of Biosciences and Informatics,

Keio University,

3-14-1 Hiyoshi Kohoku-ku,

Yokohama, 223-0061, Japan

Email: jimura@bio.keio.ac.jp

Phone: +81-45-566-1625

Abstract

Adaptation to changing environments involves the appropriate extraction of environmental information to achieve a behavioral goal. It remains unclear how behavioral flexibility is guided under situations where the relevant behavior is ambiguous. Using functional brain mapping of machine-learning decoders and directional functional connectivity, we show that brain-wide reversible neural signaling underpins behavioral flexibility in ambiguously changing environments. When relevant behavior is cued ambiguously during behavioral shifting, neural coding is attenuated in distributed cortical regions, but top-down signals from the prefrontal cortex complement the coding. When behavioral shifting is cued more explicitly, modality-specialized occipitotemporal regions implement distinct neural coding about relevant behavior, and bottom-up signals from the occipitotemporal region to the prefrontal cortex supplement the behavioral shift. These results suggest that our adaptation to an ever-changing world is orchestrated by the alternation of top-down and bottom-up signaling in the fronto-occipitotemporal circuit depending on the availability of environmental information.

Introduction

Executive control guides flexible adaptation to changing environments, and is most developed in humans throughout evolution (1, 2). Shifting between different types of behavior is one of the core executive control functions (3, 4), and switching paradigms have been often used to investigate behavioral flexibility and its underlying neural mechanisms. Previous neuropsychological and neuroimaging studies of human and non-human animals suggest a critical role of the prefrontal cortex in switching tasks and rules (5-15) (Fig. 1A).

Importantly, executive control depends on perceived information of external environments, and relevant information is appropriately extracted from the external environment to achieve a behavioral goal. Perception of sensory information from the external environment guides the course of action, which is referred to as perceptual decision-making (16, 17). It involves the extraction of goal-relevant information, which is integrated to form a relevant decision. Studies of perceptual decision-making have used behavioral tasks that demand discrimination of sensory stimuli involving perceptual uncertainty (18-20).

By manipulating perceptual uncertainty, neurophysiological and neuroimaging studies have examined cortical mechanisms of the perceptual decision-making. For example, the middle temporal (MT) region is known to play an important role in the perception of moving stimuli (18, 21-24). It has been also suggested that the fusiform face area (FFA) and parahippocampal place area (PPA) are associated with the perception of face (25-29) and place (27, 28, 30) stimuli, respectively. These collective results suggest that temporal and occipital regions play important roles in perceptual decision-making, and are functionally segmented depending on the modality of the stimulus (Fig. 1A).

In our daily life, goal-relevant information in external environments is not always evident. Such situation can be well illustrated by an incorporation of behavioral shifting and perceptual decision-making, in which the behavioral goal was indicated by a cue involving perceptual uncertainty. Notably, the task switching paradigm is composed of hierarchically structured configuration of a set of task rules, called task sets (7, 31-33), allowing the manipulation of the uncertainty of task cue at the multi-task (most abstract) tier (Fig. 1A). As such, the uncertainty of multi-task level information

may provide novel opportunities to examine the relationships between executive control and perceptual decision-making.

One potential approach to elucidate underlying neural mechanisms is to identify the signal contents of responsible brain regions; this has recently been demonstrated for perception by neural decoding techniques (34-38). In particular, prior neuroimaging studies have shown that mental and behavioral states can be decoded from neural coding by machine learning techniques that classify distributed patterns of brain activity (34-39). One commonly used technique is the support vector machine (SVM) (34, 40-43), which enables categorical discrimination by dividing multidimensional space composed of brain activation pattern using a linear hyperplane. Convolutional neural network (CNN) classifier (44, 45) is one of the deep neural network (DNN) classifiers consisting of multiple feature-aggregating layers, which allows more robust classification, especially for 2D-images. Importantly, recent technical advancements of CNN allow mapping that highlights image locations characterizing a classified image (46). The mapping technique may provide novel information about neural coding and functional localization of the brain.

The current study aimed to elucidate relationships between behavioral flexibility and perceptual decision-making under cue uncertainty, and to explore the underlying neural mechanisms (Fig. 1A). Functional MRI was administered while human participants performed a task-switching paradigm with a cue involving perceptual uncertainty. Standard univariate analysis identified brain regions associated with task switching, motion strength, and task modality. In order to elucidate causal network dynamics during task switching under cue ambiguity, we examined effective connectivity among the task-related brain regions. Finally, whole brain exploratory analyses based on machine learning techniques, CNN and SVM, were performed to identify brain regions that coded relevant task information.

Fig 1. Experimental design and behavioral results.

(A) Schematic illustration of a brain-wide model of interaction of behavioral flexibility and perceptual decision-making. Behavioral flexibility associated with prefrontal areas is subject to appropriate perception of the external world implemented in stimulus-modality-dependent occipitotemporal areas (top), and is guided by cue indicating relevant behavior at the top layer in the hierarchical structure of the behavioral set (bottom). (B/C) Behavioral task. (B) Stimuli. Cue stimulus indicates the task to be

performed (face or place task) and is composed of a set of white moving dots presented within a
donut-shaped display circle (indicated by dotted lines). The arrow indicates the motion direction of each
dot, and overall motion was either upward or downward. Upward and downward motion indicate face and
place task, respectively. Motion strength of the cue stimulus was manipulated by coherence of dot motion.
The target image was superimposed picture of face and place, which was presented at the center of screen.
Participants judged whether the face picture was male or female, or the place is indoor or outdoor,
depending on the task to be performed. (C) Task procedure. Trials with simultaneous presentation of a dot
cue and target (switch and repeat; N and N+M th trials) were followed by target-only trials (N+1, N+2
and N+M+1 th trials). Switch and repeat trials were dependent on the performed task prior to the cue
presentation. (D/E) Behavioral results. Accuracy (D) and reaction times (E) as a function of motion
coherence. Solid and dotted lines indicate face and place task, respectively, and orange, blue, and green
lines indicate switch, repeat, and target-only trials, respectively. Error bars indicate standard error of the
mean across participants.

Results

<u>Behavioral results.</u>

Human participants performed a task-switching paradigm (7, 31-33), in which they

alternated discrimination tasks for face and place stimuli (Figs. 1B-C). The relevant task

was indicated by a cue stimulus involving perceptual uncertainty, which was

manipulated by the motion strength of randomly moving dots.

Accuracy was lower in low-coherence (i.e., more uncertain) trials compared

to high-coherence trials [$F(1, 28) = 12.7$; $P < .005$; Fig. 1D], and became lower in

switch trials than in repeat trials [$F(1, 28) = 11.8$; $P < .005$]. Likewise, reaction times

(RTs) were longer in low-coherence trials compared to high-coherence trials [$F(1, 28) =

69.9$; $P < .001$; Fig. 1E], and were longer in switch trials than in repeat trials [$F(1, 28) =

43.2$; $P < .001$]. These behavioral results suggest that the current behavioral task

successfully manipulated task switching (3-5, 7, 8, 33) and perceptual decision-making

(17, 18, 21). The interaction effect of trial type (switch/repeat) and coherence levels did

not reach statistical significance [$F(1,28) = 2.1$; $P = 0.155$].

In trials without cue presentation, occurring after the switch and repeat trials

until the next cue trials (target only trial; Fig. 1C), accuracy was lower after lower

coherence cue trials [$F(1,28) = 16.3$; $P <. 001$; Fig 1D], but RTs remained unchanged

after lower coherence cue trials [$F(1,28) = 1.0$; $P = 0.317$], suggesting that switching

might not complete immediately after low coherence cue trials. RTs were longer in

place than face tasks [$F(1, 28) = 5.0$; $P <. 05$; Fig. 1E].

Fig 2. Whole-brain exploration activation and functional connectivity analyses
(A-C) Statistical activation map of univariate analysis and functional connectivity analysis. Maps are
overlaid onto the 3D surface of the brain. Hot and cool colors indicate positive and negative effects,
respectively. (A) Switch effect (switch minus repeat trials) (B) Motion coherence effect (high- vs.
low-coherence trials). Green solid line on the surface indicates the axial section on the right. (C) Task
effect (face minus place tasks in target only trials). lPFC: lateral prefrontal cortex; ITS: inferior temporal
sulcus; MT: middle temporal; FFA: fusiform face area; PPA: parahippocampal place area. (D/E)
Effective connectivity analysis. (D) Effective connectivity and extrinsic inputs modulated by the contrast
switch vs. repeat trials during face (left) and place (right) tasks. Red and blue arrows indicate connectivity
enhancements in switch and repeat trials, respectively. The values next to the arrows indicate the
magnitude of connectivity enhancements (positive: switch > repeat; negative: repeat > positive). (E)
Effective connectivity and extrinsic inputs modulated by motion coherence during face (left) and place

(right) tasks. Red and blue arrows indicate enhancements in high- and low-coherence trials, respectively.
The values next to the arrows indicate the magnitude of connectivity (positive: high coherence > low
coherence; negative: low coherence > high coherence). Arrows with solid line indicate statistically
significant connectivity (P < .05, uncorrected).

<u>Exploration of switch-related and stimulus-modality-dependent brain regions.</u>

We first explored brain regions associated with task switching, motion coherence, and

the perception of face and place based on univariate general linear model (GLM)

analysis. Figure 2A shows brain regions showing significant increases and decreases in

univariate brain activity during switch relative to repeat trials (P < .05 corrected with

cluster-wise family-wise error rate based on non-parametric permutation tests; see

Materials and Methods). Robust activation increases were observed in the left frontal

regions including the inferior frontal cortex (IFC), dorsolateral prefrontal cortex

(DLPFC), inferior frontal junction (IFJ), and pre-supplementary motor area (pre-SMA),

and in left parietal regions including posterior parietal cortex (PPC), consistent with

prior studies (5-11, 33). A full list of brain regions is shown in Table S1.

We then explored brain regions associated with motion coherence. Figure 2B

shows brain regions showing significant modulation of brain activity in relation to

motion coherence during the cue (i.e. switch and repeat) trials. In low-coherence trials,

activation was increased in multiple fronto-parietal regions including IFC, DLPFC, IFJ,

pre-SMA, and PPC (Fig. 2B and Table S2), consistent with a prior study (24). In

contrast, activation was greater in high-coherence trials in the MT region (Fig 2B *right*;

Table S2), which is also consistent with prior studies of perceptual decision-making for

motion (18, 21-24).

We next explored brain regions associated with face and place tasks (Fig. 2C

and Table S3). Consistent with prior studies of perception of face (25-29) and place (27,

28, 30), in the face task, activity was greater in the FFA, whereas in the place task,

increased activity was observed in the PPA.

These collective univariate activation results suggest that the current

univariate activation analysis successfully identified brain regions associated with task

switching in addition to the perception of face, place and motion stimulus, and that

those regions were cooperatively engaged in task switching when the cue stimulus

involved perceptual uncertainty.

<u>Reversal of functional connectivity depending on cue uncertainty.</u>

The whole brain exploratory analyses of univariate activation identified three

types of brain regions: 1) left PFC associated with task switching and perceptual

uncertainty (Figs. 2A-B); 2) MT region associated with motion coherence of task cue

(Fig. 2B); and 3) FFA and PPA associated with discrimination of face and place stimuli,

respectively (Fig. 2C). One possible mechanism to explain these regions playing

differential roles in task switching with cue uncertainty is that lPFC, MT, and FFA/PPA

mutually received or sent task-related signals during switching, which was modulated

by cue coherence, such that the task-related signal complemented the engagement of

these regions depending on cue uncertainty and the task to be performed.

In order to test this hypothesis, we performed an interregional effective

connectivity analysis based on dynamic causal modeling (DCM) that allows the

examination of directionality of task-related functional connectivity based on the

state-space model (see Materials and Methods).

We first examined the task-related effective connectivity during task

switching, and found that the connectivity was enhanced from the lPFC toward the FFA

during switch relative to repeat trials of the face task (i.e. switch-to-face versus

repeat-face trials; Fig. 2D *left*), and also enhanced from the lPFC toward the PPA in

switch relative to repeat trials of the place task (Fig. 2D *right*). These results are in line

with the well-known role of the left lPFC in behavioral flexibility (4-7), and suggest

top-down signaling from the prefrontal cortex to stimulus-modality-dependent

occipitotemporal regions during task switching.

We then asked a critical question whether the top-down signaling from the

lPFC to the stimulus-modality-dependent regions is modulated depending on the

uncertainty of the relevant task (i.e., multi-task level) that was manipulated by the

motion coherence of task cue. During face tasks with high-coherence cue, the

task-related effective connectivity was enhanced from the MT and FFA regions to the

lPFC, and the directionality of connectivity between these regions was reversed in

low-coherence trials (Fig. 2E *left*). Likewise, the effective connectivity was enhanced

from the MT and PPA regions to the lPFC during the place task with the high coherent

cue, and the directionality of the connectivity was also reversed in low-coherence trials

(Fig. 2E *right*).

In order to test the reliability and robustness of the results above, we

estimated the effective connectivity by 1) using an alternative estimation method (Figs.

S1A-B), 2) changing the number of the regions of interest (ROIs) in the models (Figs.

S1C-D), and 3) changing the definition of the ROIs (Figs. S1E-F) (see Materials and

Methods). Overall results were maintained, confirming that the connectivity results

were robust against estimation procedures, model structures, and model parameters.

Task-unrelated intrinsic connectivity is shown in Figs S1G-H.

These collective results of effective connectivity suggest that, when relevant

task was indicated ambiguously, top-down signal from prefrontal regions become

stronger toward stimulus-modality-dependent occipitotemporal regions (i.e. MT and

FFA/PPA for face/place tasks). On the other hand, when task cue information is more

evident, bottom-up signals from the occipitotemporal regions to the prefrontal regions

become stronger.

<u>Whole-brain decoding by a CNN classifier.</u>

Given the brain regions associated with tasks, switching tasks, and perceptual

decision-making identified by univariate activation analysis and their effective

connectivity mechanisms, we explored brain regions that code relevant task information

using a convolutional neural network (CNN) classifier (45). More specifically, we

examined whether brain activity patterns involve discriminable information about face

and place tasks during task switching with cue uncertainty, and then identified brain

regions involving critical information about the relevant task.

The current analysis used VGG16 (47) that was trained to classify a concrete

object image dataset provided by ImageNet (44) (http://www.image-net.org/; Fig. S2A).

We re-trained the VGG16-ImageNet model using flat whole cortical activation maps

(Figs. 3A and S2B) such that it classified face and place tasks based on fine-tuning [48]

(Fig. S2C). The retraining was performed based on cortical maps that were independent

of the tested maps. We retrained the model using flat maps during a working memory

(WM) task for face and place stimuli obtained from the Human Connectome Project

(HCP) (Fig. S2C *top*; see also Materials and Methods), followed by additional

retraining based on flat activation maps during target-only trials of the current task in

which only the face/place target stimulus was presented without the dot cue stimulus

(Figs. 1C and S2C *bottom*).

Classification accuracy for target-only trials was $82.1 \pm 5.0\%$ (mean $\pm$ SD

with 10-fold cross validation), which was significantly greater than chance level (P

< .001) (Fig S3A; see Materials and Methods). Interestingly, direct retraining of

VGG16-ImageNet model to classify the current target-only trial maps showed little

increase in accuracy (Fig. S3A). Training of a randomly initialized VGG model to

classify HCP WM maps was also not successful (Fig. S3B). Thus, these results

demonstrate that the current two-step retraining of the VGG16-ImageNet model was

sufficient for the CNN model to learn from small sample data sets of brain images.

269
Fig 3.   Convolutional neural network (CNN) classifier mapping
(A) CNN model was based on VGG16 pre-trained by ImageNet data. The model was retrained to classify
performed task (face or place). Retraining of brain maps was based on fine tuning from the 5th
convolution layer to the full-connected layers. (B) Classification accuracy for each task condition. *: P
< .005; **: P < .001. High: high-coherence (80%) trials; Mid: middle-coherence (40%) trials; Low:

low-coherence (20%) trials. (C) Weight gradients between convolution layers was aggregated and then
visualized to identify cortical regions involving more information to classify tasks. (D) Visualization of
aggregated weight contrasts for switch vs. repeat trials in face task (left) and place task (right). Statistical
maps are overlaid onto flat cortical anatomical images with a statistical threshold of |z| > 2.0 (top). Hot
and cool colors indicate statistical z-values of positive (switch > repeat) and negative (repeat > switch)
effect, respectively. Gray and black closed lines overlaid on flat map indicate brain regions significantly
activated during face and place tasks in univariate analysis, respectively (Fig 2C and S6 Fig).
Occipitotemporal regions in rectangular boxes with green broken lines were expanded below. Maps were
overlaid onto flat maps masked by the univariate activation contrast face vs. place tasks for target only
trials (gray and black closed lines in the top panels). The fusiform face area (FFA) and parahippocampal
place area (PPA) are indicated by arrow heads. (E) Visualization of weight contrast for coherence effect
in face task (left) and place task (right). Hot and cool colors indicate greater statistical values of weights
in high- and low-coherence trials, respectively. The formats are similar to those in panel (D). (F) Regions
of interests (ROIs) analysis. PPA and FFA ROIs were defined based on target-only trials, and weight
magnitudes were collected for each cue trial conditions and ROIs. All error bars in the figure indicate
standard errors of the means across participants.

Given that the CNN model successfully classified face and place tasks during

the target only trials with high accuracy, we then examined the classification accuracy

for cue trials (Fig. S2D). Accuracy was higher than chance level in switch trials at all

coherence levels [80% switch: $t(28) = 4.7$, $P < .001$; 40% switch: $t(28) = 5.5$, $P < .001$;

20% switch: $t(28) = 5.4$, $P < .001$] and 80%-repeat trials [$t(28) = 3.6$, $P < .005$] (Fig.

3B), which ensured those maps contained information about performed tasks. More

importantly, classification accuracy was higher in switch than in repeat trials [$F(1, 28) =$

10.9; $P < .005$], suggesting that cortical activation patterns involves more information

about task dimension in switch than in repeat trials, although the coherence effect was

absent [$F(1, 28) = 0.3$; $P = 0.6$].

We explored brain regions involving critical information to classify face and

place tasks by visualizing weights of convolution layers of the CNN model. We used

Grad-CAM (46) that aggregates weights across convolution layers, and highlights

image locations with greater weights when important information to classify the image

are involved (Fig. 3C). The weight maps were created for each of the tested images (i.e.

flat activation maps for cue trials), and then collected for each of the tasks (face/place),

switching conditions (switch/repeat), and coherence levels (high/mid/low) (see

Methods).

We first contrasted weight maps between switch and repeat trials, and

calculated pixel-wise group-level z-statistics with participants treated as a random effect

(Fig. 3D). Prefrontal, parietal and occipitotemporal areas including FFA and PPA (see

Figs. 2C and S4 for references) showed greater weights in switch trials than repeat trials

(Fig. 3D). In particular, weights became greater in trials switching to face task but not to

place task in the FFA (Fig. 3D *left*). On the other hand, greater weights were observed

in trials switching to the place task but not to face task in the PPA (Fig. 3D *right*). These

results suggest that modality-dependent FFA and PPA encode task-relevant information

to a greater degree during the switch to the task that demands stimulus discrimination of

optimal modality. Next we examined the coherence effect during cue trials by

calculating the weighted sum of weights for each pixel (Fig. 3E). The FFA and PPA

also showed greater weights in high-coherence trials than low-coherence trials in both

of the face and place tasks.

323    In order to statistically test dissociated weight patterns in the FFA and PPA

during cue trials, ROI analysis was performed (Fig. 3F). ROIs were defined based on

target only trials, independently of cue trials (see Materials and Methods). In the FFA,

weights were greater during the face task than the place task [$F(1, 28) = 15.2$, $P < 01$],

and in the PPA, weights were greater during the place task than the face task [$F(1, 28) =$

66.5, $P < .001$]. For weight magnitudes with optimal task-region relation (i.e., face task

in FFA and place task in PPA), weights became greater in switch relative to repeat trials

[$F(1, 28) = 6.4$, $P < .05$], and in high relative to low-coherence trials [$F(1, 28) = 13.4$, P

$< .01$ with linear contrast]; however, there was no without switching-by-coherence

interaction [$F(1, 28) = 0.1$, $P = .75$]. In a separate analysis, we used a

leave-one-subject-out procedure when retraining the classifier to classify the activation

maps of the target only trials, and then tested the remaining subject (Fig. S5). Overall

results are consistent, suggesting that subject-specific noises are not dominant in our

results.

337    These results suggest that the FFA and PPA involve more modality

specialized task-related pattern information in high-coherence trials. Thus,

modality-dependent occipitotemporal regions may encode relevant task information (i.e.

FFA for face task and PPA for place task), which is enhanced in switch trials with a

high-coherence task cue.

<u>Whole-brain decoding by SVM.</u>

In order to complement decoding and mapping by the CNN classifier, we performed

another decoding analysis using an SVM classifier. Similar to the CNN classifier

analysis above, the classifier was trained based on HCP WM task such that the classifier

discriminates the dimension of the tasks (face or place) (see Materials and Methods).

We then tested the experimental data to examine classification performance for face and

place tasks of the cue trials.

Fig 4. Multi-variate pattern analysis (MVPA) with support vector machine (SVM).
(A) Classification accuracy for each task condition with SVM classification in the whole brain. The
model was trained for classification of face and place tasks with the working memory task in the Human
Connectome Project (HCP) and tested with each condition of cue trials in the current study. Error bars
indicate standard error of the mean across participants. *: P < .05; **: P < .005; ***: P < .001. (B)
Visualization of weight assigned to the pixels for classification of face and place tasks in the working
memory task in the HCP. Formats are similar to those in Figs. 3D/E. (C-E) Statistical significance maps
for searchlight MVPA. Classifiers were trained to classify the performed task. Maps are overlaid onto a
3D surface of the brain and displayed from a ventral view. White and black closed lines overlaid onto the
3D surface of the brain indicate significant clusters for contrast face vs. place tasks in univariate analysis,
respectively (Fig. 2c). The fusiform face area (FFA) and parahippocampal place area (PPA) are indicated
by blue arrow heads. (C) Target-only trial. Hot and cool colors indicate statistical level for classification
accuracy relative to chance level. (D) Accuracy difference between switch and repeat trials. Hot and cool
colors indicate higher accuracy in switch and repeat trials, respectively. (E) Differential classification
accuracy depending on the coherence effect. Hot and cool colors indicate higher accuracy in high- and
low-coherence trials, respectively.

We found that accuracy was higher than chance level in switch trials at all coherence levels [80% switch: $t(28) = 4.5$, $P < .001$; 40% switch: $t(28) = 2.5$, $P < .05$; 20% switch: $t(28) = 3.3$, $P < .005$; Fig. 4A], and higher in switch than repeat trials [$F(1, 28) = 5.0$; $P < .05$], which is consistent with the CNN classifier results.

Weights of the SVM classifier were mapped onto 2D cortical surface of the brain in order to identify brain regions with greater weights to classify the face and place tasks. Occipitotemporal regions including the FFA and PPA showed prominent reverse directed weights (Fig. 4B), indicating that these regions involve important information to classify the two tasks. Notably, these maps are consistent with the CNN-based mapping, especially in the FFA and PPA (Figs. 3D/E). We also trained another SVM based on target-only trials in the current task, and found that the classification accuracy of cue trials (Fig. S6A) and weight maps (Fig. S6B) were consistent to those with the CNN classifier (Figs. 3B/D-E) and whole-brain SVM (Figs. 4A-B). We note that SVM weight maps (Figs. 4B and S6B) reflect a hyperplane calculated by training data (i.e. maps for HCP WM or current target-only trials), whereas CNN weight maps reflect degree of contribution to classify tested image (i.e. current cue trial maps) (see Discussion).

Decoding mapping by searchlight SVM.

The above CNN and SVM classifiers were based on pattern information of whole brain cortical regions. Another SVM analysis was also performed using the searchlight procedure (see Materials and Methods). By exploring across the whole brain, searchlight was used to identify brain regions where local image voxels involved pattern information about the performed task to classify face and place tasks. Again, the classifier in each searchlight was trained using HCP datasets, and thus the training and testing datasets were independent.

For target-only trials, classification accuracy was significantly higher in the FFA and PPA regions (Fig. 4C and Table S4), suggesting that modality-dependent occipitotemporal regions involve relevant task information; this result is consistent with the univariate analysis (Fig. 2C).

Voxel-wise classification accuracy maps were contrasted between switch vs. repeat trials for each participant, and group-level statistical tests were performed in

order to identify brain regions where discriminable pattern information is greater in

switch trials than in repeat trials. Occipitotemporal regions showed a significant effect

of switching (Fig. 4D and Table S5), indicating that, in these regions, classification

accuracy is higher in switch relative to repeat trials. Interestingly, these regions were

spatially located in-between the FFA and PPA, where pattern information of searchlight

classifiers may modestly involve both of face-related and place-related information,

possibly in a balanced manner. These regions also showed a coherence effect with

higher accuracy in high-coherence trials (Fig. 4E and Table S6). Notably, these results

were consistent with weight mapping of the CNN classifier (Figs. 3D-E).

410   It is important that these regions showed significantly higher classification

411   accuracy than chance-level in switch and repeat trials (Figs. 7A-B and Tables S7-8), and

412   cue trials at each coherence level (Figs. S7C-E and Tables S9-11). This assures that the

413   differential accuracies between switch and repeat trials, and across coherence levels

414   were attributable to accuracy enhancement in switch trials with the high-coherence cue.

415   These collective results suggest that occipitotemporal regions adjacent to

stimulus-modality-dependent FFA/PPA areas involves information about ongoing task,

and that the information amount is increased during task switching with more coherent

cue presentation. These results are also consistent with those of the classification

performance and mapping based on whole-brain CNN and SVM classifiers. Such

differential classification accuracy was not observed in fronto-parietal regions well

known to be involved in executive control (Figs. S7F-G and Tables S5-6), even when

the classifier was trained by target-only trials in the current experiment (Figs. S7H-I).

Discussion

The current study examined neural mechanisms during task switching under situation where task cue involved uncertainty. Task-related neural coding in FFA/PPA became more evident during task switching and also when the relevant task was cued more explicitly. When task-cue was distinct, the lPFC received task-related signals from the MT region and PPA/FFA, and the direction of the signal was reversed when the task cue involved more ambiguity. These results suggest a distributed cortical network of fronto-occipitotemporal regions for behavioral flexibility where task-related signal among these regions help to implement task representation depending on the ambiguities of external cue (Fig. 5).

Fig 5. Putative model of behavioral flexibility under perceptual uncertainty
Schematic diagrams for functional mechanism among the middle temporal (MT) region, fusiform face area (FFA), parahippocampal place area (PPA), and lateral prefrontal cortex (lPFC) during task switching with cue uncertainty. The arrows indicate signal directions.

<u>Neural mechanism for task switching and perceptual decision making.</u>

Prior work of task switching has examined switch-related neural mechanisms under situations where perceptual uncertainties were applied to target stimulus (24, 49-51). In contrast, the current study manipulated uncertainty of the task cue that involves multi-task information (Figs. 1A-C and 5). Notably, the task cue indicates relevant task dimension at the top tier of the hierarchical task set structure, and thus the current study

allowed us to elucidate higher-level cognitive functions governing task switching and

perceptual decision-making.

For putative mechanisms to achieve task switching under cue uncertainty,

three hypotheses are possible: 1) a unitary mechanism implements task switching under

perceptual uncertainty in a task cue; 2) distinct mechanisms for perceptual

decision-making and task switching interactively guides successful task switching; and

3) a hub-like region links the two distinct mechanisms. Behaviorally, an interaction

between switching and coherence levels was absent (Figs. 1D/E; see also Results); this

is consistent with the univariate imaging analysis and CNN classifier mapping showing

no interaction effect. The absence of an interaction effect, together with distinct brain

regions associated with cue/target perception and task switching (Figs. 2A-C and 3D-F),

suggests distributed mechanisms for task switching under cue ambiguities, which

supports the second hypothesis. Thus, by modulating effective connectivity and

interregional signaling depending on cue ambiguity, these regions cooperatively guide

behavioral flexibility (Fig. 5). This interpretation is also compatible with the central role

of frontal regions for flexible task control (52).

<u>Comparisons of classification and mapping among machine learning techniques.</u>

In the current study, whole brain exploration of task-related neural representation was

performed by three approaches based on three machine learning techniques, 1) CNN

classifier, 2) whole-brain cortical SVM, and 3) searchlight SVM.

CNN classified activation maps along task dimensions based on all pixels

across whole cortical regions, and the classification accuracy was higher in switch trials

than repeat trials. One novel signature of the current CNN classifier approach is that

brain regions involving critical information to classify cue trials were mapped by

aggregated weight gradients across convolution layers based on the Grad-CAM

technique. It is notable that this CNN weight mapping is available on image-by-image

basis for testing data, which is not the case for the SVM mapping using whole cortical

images. Then, the CNN weight mapping revealed that task representation in the FFA

during the face task and in the PPA during the place task was enhanced in switch and

more coherent trials than in repeat and less coherent trials. Increased task-related

activation (Figs. 2C and S4) may be associated with higher classification accuracy and

enhanced task representation.

Standard voxel-wise univariate GLM analysis identifies brain regions where

the MRI signal is differentiated between task conditions, but does not necessarily

indicate that identified brain regions are critical for task performance; this makes it hard

to identify brain regions playing an important role in cognitive functions (i.e., reverse

inference). In contrast, the current CNN classifier demonstrated that the visualization of

convolution layers of the classifier for brain activation is useful in identifying brain

regions that characterize task performance.

The analysis based on the CNN classifier was complemented by a standard

SVM analysis for whole brain cortical maps that tested identical map images. The

classification accuracy of cue trials was highly consistent between the two classifiers.

Additionally, SVM weight maps also showed differential weights in the FFA and PPA,

which is also consistent with the CNN classifier. One notable technical limitation of the

SVM is that weight mapping for testing classification for cue trials was unavailable,

unlike Grad-CAM of CNN classifier. Thus the SVM weight map indicates that the FFA

and PPA is critical to classify tasks for the HCP N-back working memory task or target

only trials in the current task (i.e. training data), but not necessarily for cue trials of the

current task (i.e. testing data). Nonetheless, together with differential classification

accuracy among cue conditions, the SVM weight maps suggest that pattern information

in the FFA and PPA is distinct (i.e. distant from separating hyperplane) in switch trials

than in repeat trials.

Another approach to identify brain regions that characterize task performance

is the searchlight SVM, which also allows whole-brain exploration of activation

patterns, but individual classifications were restricted in local brain regions (Figs 4C-E

and S7A-E). This is in contrast to the CNN classifier and whole-brain cortical SVM that

are trained and classify based on a whole brain image. Nonetheless, results were

complementary to those whole-brain-based classifiers in that 1) occipitotemporal

regions adjacent to the FFA and PPA were capable of task classification, and 2) the

classification performance in these regions became higher in switch and more coherent

trials.

Classifier training using independent open resource data.

One notable analysis procedure in the current machine-learning-based functional brain

mapping is that classifier was trained using an open resource dataset that was

independently collected from the current experiment. This procedure ensured

independence between the training and testing data.

Task switch and working memory may involve distinct cognitive control

demands, with the former related to behavioral flexibility (3, 4) and the latter related to

active maintenance and updates of goal-relevant information (53). However, the two

tasks used common visual stimulus categories (face and place); thus, perceptual

demands may involve some degree of commonality. Recognition demands for the

presented stimuli were also distinct: the current task-switching paradigm used

male-female and indoor-outdoor discriminations during face and place tasks,

respectively, but HCP working memory tasks used discrimination of identicalness to

past stimuli. Additionally, the current task used face-place superimposed stimuli, and

thus the identical stimulus set was used during face and place tasks, whereas HCP

working memory task used distinct visual stimulus sets during face and place blocks.

Thus task representation examined in the current study may reflect visual perception or

attention rather than low level visual features.

For the CNN classifier, training involved two steps: retraining of HCP

working memory maps, and additional retraining of target-only trials. CNN classifies

maps based on whole cortical areas including fronto-parietal regions in which

differential sub-regions are recruited during task switching (5-15, 31, 33) and working

memory (1, 53, 54). Thus additional retraining of the CNN model based on identical

recognition demands (i.e. target-only trials) was effective in classifying tasks to

optimize the CNN model for classification using whole cortical images. Distinct weight

differences in the parietal cortex (Figs. 3D-E) may partially be attributable to higher

performance with the additional retraining.

Because incremental training of HCP working memory trials and the target

only trials in the current study is irrelevant to SVM, these two datasets were separately

trained for whole brain cortical SVM, and weight maps were consistent especially in

occipitotemporal regions (Figs. 4B and S6B). Importantly, classification accuracy for

the cue conditions were consistent in SVMs with the two training datasets, and also

with the CNN classifier. The sample size was much smaller for the current target-only

trials than HCP dataset, but classification performance was comparable between those

two classifiers (Figs. 4A and S6A). Thus, SVM may thus not require a larger sample

size like the HCP data for training, while CNN training needed incremental training
even with large sets of image data.
The searchlight SVM using HCP working memory maps as training data
identified occipitotemporal regions spatially closed to the FFA/PPA showing higher
classification accuracy for target-only trials. Moreover, classification accuracy was
higher in high-coherence switch trials. Interestingly, these classification results were
absent in fronto-parietal regions, well known to be involved in executive control (Figs.
S7F/G), even when the searchlight classifier was trained by the target-only trials of the
current task (Figs. S7H/I). One possibility for this discrepancy is that control and
recognition demands are incompatible while perceptual modality is compatible in HCP
working memory trials, target only trials, and switch/repeat trials. Then the distinct
control and recognition demands might be reflected in classification incompatibility in
the fronto-parietal regions.
558
<u>Fronto-occipitotemporal network mechanisms.</u>
DCM analyses revealed top-down signal from the lPFC to the FFA or PPA, depending
on the task to be switched, and this top-down signaling is enhanced when task cue
involved ambiguity during switching (Fig. 5). The top-down mechanisms may reflect
supplemental attention to visual stimulus required to collect task cue information about
the tasks to be performed. The supplemental attention involving frontal engagement
may complement stimulus-modality-dependent activation in occipitotemporal regions
(55-58).
In contrast, bottom-up signaling with increased cue information may reflect
the conversion of sensory information to behavioral information through an information
stream from the visual sensory area to executive control areas (55, 56). Thus, when the
task cue was apparent, the bottom-up signal was strengthened because cue-related
information is more available, which may help to enhance task switching performance.
572

Materials and Methods

<u>Participants.</u>

Written informed consent was obtained from 30 healthy right-handed subjects (age

range: 18-22; 11 females). Experimental procedures were approved by the institutional

review board of Keio University and Kochi University of Technology. Participants

received 2000 yen for each of the training and scanning sessions. One participant was

excluded from analyses due to low behavioral performance; accuracy was lower than

30% in one of the experimental conditions.

<u>Behavioral procedures.</u>

The experiment consisted of two sessions administered on separate days. The first day

was a training session, in which participants practiced discrimination tasks (random dot

motion) and switching between two tasks (face and place tasks; see below for

behavioral procedures). On the second day, while fMRI scanning was administered, the

participants performed the switching paradigm identical to those practiced in the

training sessions.

<u>Stimuli.</u>

All stimuli were generated in Matlab version 2012a, using the Psychophysics Toolbox

(59) extension version 3.0.10, and were visually presented on a computer screen. The

current task cue stimuli were randomly moving dot stimuli similar to those used in a

previous study of perceptual decision-making (55). Each motion stimulus involved 60

dots moving inside a donut-shaped display patch with a white cross in the center of the

patch on a black background (Fig. 1B). The display patch and cross were centered on

the screen and extended from 6 to 12° degrees of visual angle (dva). Within the display

patch, every dot moved at the speed of 10 dva per second. Some dots moved coherently

toward one direction (upward or downward) while the others moved randomly. The

percentage of coherently moving dots determined the "motion coherence", which was

set to three levels (20, 40, and 80%).

Dot presentation was controlled to remove local motion signals following a

standard method for generating motion stimuli (18, 60-62). Namely, upon stimulus

onset, the dots were presented at new random locations on each of the first three frames.

They were relocated after two subsequent frames, such that the dots in frame 1 were

repositioned in frame 4, and the dots in frame 2 were repositioned in frame 5, and so on.

When repositioned, each dot was either randomly presented at the new location or

aligned with the pre-determined motion direction, depending on the pre-determined

motion strength on that trial. Each stimulus was composed of 18 video frames with a 60

Hz refresh rates (i.e. 300-msec presentation).

Within the center circle mask of the donut-shape motion stimulus, a

face/place superimposed stimulus was presented simultaneously (Fig. 1B). The face

image set consisted of an image of picture of 4 male and 4 female unfamiliar Japanese

faces, and the place image set consisted of 4 indoor and 4 outdoor unfamiliar places;

this resulted in 64 overlaid images.

<u>Task procedure.</u>

At the beginning of the task, a dot patch and face/place stimulus were simultaneously

presented. The direction of the dot patch (up or down) indicated the task to be

performed (discrimination of face or place). Depending on the motion direction,

participants were required to judge whether the face was male or female, or whether the

place was indoor or outdoor (Fig. 1B), and pressed the corresponding button with their

right thumb. Both of accuracy and speed were stressed. Stimulus was presented for 1.8

sec, followed by a 0.7 sec intertrial interval. If participants made an incorrect response,

or did not respond within 1.8 sec from the stimulus onset, feedback stimulus indicating

an error (X) was presented for 1.0 sec, followed by high-coherence (80%) cue trials for

the same task dimension. The cue trials immediately after the error were discarded from

analyses. Stimulus-response and cue-task associations for the two tasks were identical

on days 1 and 2, and counterbalanced across participants.

The trial with simultaneous cue/target presentation was followed by trials

with presentation of the face/place target stimulus without the dots cue stimulus (target

only trials). The target-only trials were repeated for 3-5 times (Fig. 1C). In the

target-only trials, participants were required to discriminate the center image stimulus

along the same dimension until the next task cue (moving dots) was presented. One task

block lasted for approximately for 90 secs, and 20-sec fixation blocks were inserted

between task blocks. Each functional run involved 3 task blocks and lasted for 305 secs.

The first trial at the beginning of each task block presented the dot cue with highest

coherence (80%), and was discarded from analysis.

<u>Practice procedure.</u>

On the first day, participants practiced the tasks outside of the scanner. They first

practiced a discrimination task for the moving dot stimulus, and were required to judge

the direction of overall motion (up or down) and to press the correct corresponding

button as quickly as possible. The response window was 1050 msec. Each practice run

involved 70 trials, and 5 runs were administered for each participant. The first 5 trials

and last 5 trials in each run used the highest coherence level (80%). Thus the middle 60

trials were composed of 20 trials for each of coherence levels (20, 40, or 80%).

The participants then practiced discrimination tasks for the face and place

stimulus (see above). Across task switching practice runs, the switching frequency and

coherence levels of the moving dot cue were manipulated such that the cue trials

gradually became more difficult (i.e. more switch trials with low-coherence cue).

Participants performed 8 practice runs for task switching.

<u>Behavioral procedure in scanning session.</u>

On the second day, after practicing task switching for one run, the participants

performed 9 runs of task switching with identical procedure as for day 1 (see above)

while functional MRI was administered. The frequency of switch and repeat trials and

coherence level were approximately equivalent across runs.

<u>Imaging procedure.</u>

MRI scanning was administered by a 3T MRI scanner (Siemens Verio, Germany) with

a 32ch head coil. Functional images were acquired using a multi-band acceleration

echo-planar imaging sequence (63) [repetition time (TR): 0.8 sec; echo time (TE): 30

msec; Flip angle (FA): 45 deg; 80 slices; slice thickness: 2 mm; in-plane resolution: 3 x

3 mm; multiband factor: 8]. Each functional run involved 385 volume acquisitions. The

first 10 volumes were discarded from analysis to take into account the equilibrium of

the longitudinal magnetization. High-resolution anatomical images were acquired using

an MP-RAGE T1-weighted sequence [TR: 2500 msec; TE = 4.32 msec; FA: 8 deg; 192

slices; slice thickness: 1 mm; in-plane resolution: 0.9 x 0.9 mm^2].

<u>Behavioral analysis.</u>

Trials were classified into three types in terms of switching: 1) task switch trials

presenting the random dot cue and target stimuli simultaneously, where the task to be

performed alternated (i.e., face to place or place to face); 2) task repeat trials presenting

the random dot cue and target stimuli simultaneously, where the same task was

repeated; and 3) target-only trials presented after the switch and repeat trials, and their

subsequent trials without random dot cue stimuli presentation (Fig. 1C). These trial

types were analyzed separately. Trials were also classified by task dimension (face or

place), and switch and repeat trials were examined at each coherence level (20, 40, or

80%). Accuracy and reaction times (RTs) were calculated for each trial condition, and

then compared. Statistical testing was performed based on repeated measures ANOVAs

implemented in SPSS Statistics 24 (IBM Corporation, NY USA).

Image preprocessing.

MRI data were analyzed using SPM12 software (http://fil.ion.ac.uk/spm/). All

functional images were initially temporally realigned across volumes and runs, and the

anatomical image was coregistered to a mean image of the functional images. The

functional images were subsequently spatially normalized to a standard MNI template

with normalization parameters estimated based on the anatomical scans. The images

were resampled into 2-mm isotropic voxels, and spatially smoothed with a 6-mm

full-width at half-maximum (FWHM) Gaussian kernel.

Imaging analysis: general linear model (GLM).

Single level analysis.

A GLM approach (64) was used to estimate parameter values for task events. The

events of interest were correct switch, repeat, and target-only trials. For switch and

repeat trials, the normalized (z-scored) coherence level of the dot stimuli was also added

as a parametrical effect of interest. Error trials in all conditions were separately coded in

GLM as nuisance effects. Those task events were time-locked to the onset of target

images and then convolved with canonical hemodynamic response function (HRF)

implemented in SPM. Additionally, six-axis head movement parameters, white-matter

signals, lateral-ventricle signals, and parametrical effect of reaction times normalized

across trials were also included in GLM as nuisance effects. The parameters were then

estimated for each voxel across the whole brain.

<u>Group-level analysis.</u>

Maps of parameter estimates were first contrasted within individual participants.

Contrast maps were collected from all participants, and subjected to a group-level

paired t-test. For the coherence effect, the contrast maps were subjected to a one-sample

group-mean test, with maps weighted and summed based on normalized coherence

levels. Voxel clusters were identified using an uncorrected threshold of $P < .001$ based

on voxel-wise t-statistics. The voxel clusters were tested for a significance with a

threshold of $P < .05$ corrected by family-wise error (FWE) rate based on permutation

methods (65) (5000 permutations) implemented in *randomise* in FSL suite

(http://fmrib.ox.ac.uk/fsl/). This group analysis procedure was validated to appropriately

control false positive rates in a prior study (66). Peaks of significant clusters were then

identified and listed on tables. If multiple peaks were identified within 12 mm in one

cluster, the most significant peak was retained. When exploring brain regions associated

with motion coherence, exploration was restricted within a mask obtained from

Neurosynth (67) (http://neurosynth.org/) for the search word 'motion' ($z > 3.0$, for

uniformity test), in order to ensure the extraction of motion-related regions, because the

current cue trials simultaneously presented face/place stimuli in addition to cues

indicating task switching tasks between face and place discrimination.

<u>Effective connectivity analysis.</u>

The current analysis was designed to test the hypothesis that functional connectivity

among brain regions associated with task switching, motion perception, face perception,

and place perception identified in univariate analysis (Figs. 2A-C) is modulated by task

manipulations and brain signals. Dynamic causal modeling (68) (DCM) analysis was

performed in order to examine functional connectivity mechanisms associated with task

switching under cue uncertainty (Figs. 2D-E). DCM allows us to explore the effective

connectivity among brain regions under the premise that the brain is a deterministic

dynamic system that is subject to environmental inputs and produces outputs based on

the space-state model. The model constructs a nonlinear system involving intrinsic

connectivity, task-induced connectivity, and extrinsic inputs. Parameters of the

nonlinear system are estimated based on fMRI signals (system states) and task events.

Four regions of interest were first defined based on univariate analysis and

prior studies of task switching and perceptual decision-making: 1) task switching [left

lateral prefrontal cortex (lPFC) (5-11, 33, 69); Fig. 2A]; 2) motion perception [middle

temporal (MT) (16-18, 21-24); Fig. 2B]; 3) face perception [fusiform face area (FFA)

(23-27); Fig. 2C]; 4) place perception [parahippocampal place area (PPA) (27, 28, 30);

Fig. 2C]. More specifically, meta-analysis maps were obtained from Neurosynth[64]

(http://neurosynth.org/) using a keyword search for "switching", "mt", "ffa", "place" to

obtain the meta-analysis maps for lPFC, MT, FFA, and PPA regions of interest (ROIs),

respectively. ROI images were then created with 6-mm radius spheres centered in the

peak coordinates in the meta-analysis activation maps thresholded above $z > 10$

(uniformity test).

Given these ROIs, we first tested whether the switch-related prefrontal region

sends (or receives) a task-related signal toward (or from) the stimulus-modality

dependent occipitotemporal region of the target (i.e. FFA/PPA) during task switching

(Fig. 2D). If this were the case, then we tested whether signaling between prefrontal and

occipitotemporal regions changed depending on the uncertainty of cue stimuli (Fig. 2E).

Signal time courses of four ROIs and regressors in events of interest were

extracted from first-level GLMs. The events of interest were cue (switch and repeat)

trials and target-only trials of each task. For switch and repeat trials, the contrast of the

two trials and normalized coherence level of the dot stimuli were added as a

parametrical effects of interest. Nuisance effects of head-motion, white matter signal,

ventricle signal, functional run, and contrast were subtracted out from the ROI

timecourses. The input matrix was U mean-centered.

For each trial effect, causal models were defined as those that differed in

external inputs and modulatory effects among ROIs. Because the current analysis

involved 2 or 3 ROIs (Figs. 2D-E), the tested models included 16 or 512 types (i.e. 2^2

inputs and 2^2 connection effects or 2^3 inputs and 2^6 connection effects). Connectivity

matrices reflecting 1) first-order connectivity, 2) effective change in coupling induced

by the inputs, and 3) extrinsic of inputs on MRI signal in ROIs were estimated for each

of 1the 6 or 512 models based on DCM analysis implemented in SPM12. Parametric

regressor (switch vs repeat / coherence) was used as an extrinsic effect for effective

connectivity between ROIs and ROIs inputs.

In order to estimate strength of effective connectivity, a Bayesian model

reduction method (70) was used. The reduction method allows the calculation of

posterior densities for all possible reduced models, which were then inverted to a fully

connected model. The reduced models were then supplemented by second-level

parametric empirical Bayes (70) (PEB) to apply empirical priors that remove subjects'

variability from each model.

Next, the parameters of these models were estimated based on Bayesian

model averaging (68) (BMA) to estimate group-level statistics. Because the current

analysis aimed to identify effective connectivity observed as an average across

participants, we used a fixed effect (FFX) estimation assuming that every participant

uses the same model. This is in contrast to using a random effect (RFX) estimation

assuming different participants use different models, which is often used to test group

differences in effective connectivity (71). The significance of connectivity was then

tested by thresholding at a posterior probability at the 95% confidence interval. We used

the uncorrected threshold, because the current analysis aimed to test if connectivity

between two specific brain regions was enhanced depending on task manipulation and

brain activity, not to explore one model involving connectivity among multiple brain

regions that best fits to the imaging and behavioral data.

Additionally, in order to test the robustness of the functional connectivity, we

performed supplemental analyses. We estimated model parameters 1) without an

empirical prior (Figs. S1C-D); 2) changing the number of the regions of interest (ROIs)

in the models (Figs. S1E-F), and 3) changing the definition of the ROIs (Figs. S1G-H).

When changing the ROI definition, we used a leave-one-out procedure; the centers of

ROIs of one participant were determined based on group-level univariate activation

maps of corresponding contrasts (i.e. lPFC: switch vs. repeat trials; MT: high vs. low

coherence trials; FFA/PPA: face vs. place of target-only trial) without the participant in

order to circumvent circular analysis. The ROIs were created as spheres with 6-mm

radius for individual participants. Complete results can be provided upon request.

<u>Convolutional neural network (CNN) classifier.</u>

In order to explore brain regions involving task-related neural representation, a

convolutional neural network classifier (CNN) (44, 45) was used. The current CNN

model was based on VGG16 (47), with five convolution layers for extracting image

features and two fully connected layers for binary classification. Initial parameters of

convolution layers were set to parameters pre-trained with concrete object images

provided from ImageNet (72) (Fig. S2A; http://www.image-net.org/).

The VGG16/ImageNet model is capable of classifying concrete object images

into 1,000 item categories. Importantly, it has been demonstrated that the pre-trained

model can learn novel image sets more efficiently than the non-trained model by tuning

convolution and fully connected layers and fully connected layers only (48, 73) (Figs.

3A and S2C). Thus, the current analysis retrained the pre-trained VGG16-ImageNet

model to classify brain activation maps.

Training data were single-subject 2nd level z-maps during the N-back

working memory task from the S1200 release of the Human Connectome Project (74,

75) (N = 992; HCP; http://www.humanconnectomeproject.org/). From each participant,

statistical z-maps for activation contrasts for face vs. fixation and place vs. fixation

(2-back and 0-back corrupted) were collected. The pre-trained VGG16 model was

retrained by the activation maps during the N-back working memory task such that the

model classifies face and place trials. We used gray-scaled flat 2D cortical maps (76)

provided from HCP (992 images; face: 496, place: 496; Fig. S2B) for dimensional

compatibility of images between VGG16-ImageNet and activation maps. The training

data set was divided into 10 subsets, and 9 subsets were used for retraining and the

remaining 1 set was used for validation, enabling a 10-fold cross-validation test. Model

training and testing was implemented using Keras (https://keras.io/) under Tensorflow

backend (https://www.tensorflow.org/) [input image size: 480 x 1280 pixels; batch size:

10; epoch: 50; learning rate: 0.0001; optimizer: Stochastic Gradient Descent (SGD);

Figs. 3A and S2C *top*].

After retraining of the HCP working memory maps, the model with highest

classification accuracy was further retrained to classify activity maps for face and place

tasks during target-only trials of the current dataset (Fig. S2C *bottom*). For each

functional run of each participant, a single-level GLM estimation was performed with

regressors identical to those in the univariate analysis as described above. The GLM

estimations were performed within standard MNI space. Activation z-maps for the

contrasts for face vs. fixation and place vs. fixation during correct target-only trials were

collected from each functional run. Activity maps for the contrast for face vs. fixation

and place vs. fixation were then gray-scaled and flattened such that these maps were

anatomically and geometrically identical to those from the HCP working memory task

using Connectome Workbench

(https://www.humanconnectome.org/software/connectome-workbench/). The training

dataset consisting of 522 images (261 face and 261 place maps from each of 9 runs of

29 participants) was divided into 10 subsets, enabling a 10-fold cross-validation test.

Given the limited number of images available from the current experiment,

this two-step retraining of the model was found to be effective when classifying current

tasks, because 1) training randomly initialized models failed in classifying the current

target-only trials (Fig. S3A) and in classifying the HCP working memory conditions

(Fig. S3B); 2) retraining VGG16-ImageNet was successful in classifying the HCP

working memory conditions (Fig. S3B); 3) retraining VGG16-ImageNet model failed in

classifying the current target-only trials (Fig. S3A); and 4) retraining

VGG16-ImageNet/HCP was successful when classifying the current face/place tasks

(Fig. S3A).

After learning of the target-only trials from the current dataset, the retrained

10 models were tested to classify activation maps during task switching and repeat trials

where the dot cue stimulus was presented to indicate the task to be performed. Testing

data was created based on a GLM analysis where switch and repeat trials at differential

coherence levels were coded separately. For each functional run of each participant, a

single-level GLM estimation was performed, and activation contrast z-maps for face vs.

fixation and place vs. fixation were collected during those correct switch and repeat

trials. Gray-scale 2D activation maps were created for the contrasts for face vs. fixation

and place vs. fixation during 6 types of cue presentation trials (switch/repeat x

high/middle/low coherence). Importantly, the testing data were independent of the 2 sets

of retraining data (HCP working memory and current target-only trials). The maps were

tested, and accuracy was averaged across cross-validation models within each

participant. A statistical test of classification accuracy was performed based on repeated

measures ANOVAs implemented in SPSS Statistics 24 (IBM Corporation, NY USA).

In a separate supplemental analysis, in order to examine whether the current

results were biased by subject-specific characteristics of image data, we used a

leave-one-subject-out procedure to retrain the CNN classifier to classify activation maps

of the target only trials, and then tested the remaining subject (Fig. S5).

<u>Visualization of convolution layer weights.</u>

In order to identify brain regions involving discriminative information to classify

performed tasks, Grad-CAM (46) was used. Grad-CAM visualizes the aggregation of

weight gradients between convolution layers, and highlights image locations critical for

classification on a pixel-by-pixel basis (Fig. 3C).

Aggregated weights were visualized onto 2D brain surface maps for each

tested map. For each participant, the weight maps were averaged across models and

maps within each of the 6 cue trial conditions (switch/repeat x high/middle/low

coherence cue). The averaged maps were contrasted between switch and repeat trials to

explore brain regions showing differential weights gradients between switch and repeat

trials within participants. For coherence the effect, three maps for coherence level trials

were weighted and summed based on behavioral accuracy estimated by sigmoid fitting

within participants. For each contrast, maps were collected from all participants, and

pixel-wise z-values were calculated treating participants as a random effect, and the

z-values were mapped on the 2D surfaces of the cortical areas.

In order to statistically test dissociable weight patterns during cue trials in the

FFA and PPA, ROI analyses were performed (Fig. 3F). ROIs were defined as

occipitotemporal regions showing greater activation in the contrast of face minus place

tasks or place minus face tasks (Figs. 2C and S4; Table S3), independently of the cue

trials. From each ROI, weight magnitudes were collected for each of the trial

conditions: task (face/place), switching (switch/repeat), and cue coherence

[high/middle/low: 80/40/20%], and then averaged within ROIs for each participant.

Statistical tests were then performed based on repeated measures ANOVA.

<u>Support vector machine (SVM) analysis for whole brain cortical regions.</u>

In order to supplement the CNN classifier analysis, multi-variate pattern analyses

(MVPA) based on SVM were performed. The SVM classifier was trained to perform

bivariate classification for face and place tasks. Training and testing were implemented

using scikit-learn package (https://scikit-learn.org/stable/) with a Tensorflow backend

(https://www.tensorflow.org/). We used a linear kernel, and adjusted C parameters (C =

0.1, 1.0, 10.0). As the overall results were maintained with the C adjustment, then we

reported the results with the default parameter (C = 1.0).

For classifier training, we used the image set of single-subject 2nd level

z-maps during the N-back working memory task (face vs. fixation and place vs.

fixation) obtained from the S1200 HCP (N = 992), which was identical to those used in

the CNN classifier training. The classifier was trained by the activation images, such

that it classified face and place tasks during the HCP N-back working memory task.

Weights of the trained classifier were mapped on the 2D cortical surface.

The testing dataset was also identical to those used in the CNN classifier

analysis: 2-D z-maps for activation contrast of switch and repeat trials (switch vs.

fixation and repeat vs. fixation) at each coherence level (20, 40, 80%) of the current

experiment (Fig. S2B). Testing activation images were subject to the trained classifier

for each cue trial condition of each participant, and classification accuracy was averaged

for each trial condition across participants (Fig. 4A). In a separate analysis, the SVM

classifier was trained based on single level z-maps during target only trials from the

current dataset (N = 29), and the identical image set was tested (Fig. S6).

<u>Searchlight SVM.</u>

In order to explore brain regions in which local activity patterns involves information

about a performed task (face/place), searchlight MVPA (77) was conducted. Bivariate

classification based on SVM was used to decode the performed task (face or place). A

searchlight procedure with a 5-voxel radius was used to provide a measure of decoding

accuracy in the neighborhood of each voxel. Training and testing were performed based

on the Decoding Toolbox (TDT; version 3.95;

https://sites.google.com/site/tdtdecodingtoolbox/). Again, training and testing data were

independent, based on different behavioral tasks and data sets (training: HCP working

memory; testing: current task switching with male/female or indoor/outdoor

judgements).

Training data were 3D single subject 2nd level z-maps during N-back working

memory task (N = 1,000) from HCP S1200 release (74, 75). Similar to the CNN

analysis, we used activation contrasts for face task vs. fixation and for place task vs.

fixation (2-back and 0-back corrupted). These activation contrasts were collected from
each participant, and the whole dataset was divided into 10 subsets (N = 100 each). For
each subset of the training data, a classifier in each searchlight was trained based on
these z-maps such that it classified face and place conditions during the HCP N-back
working memory task.

Test data were single-subject z-maps during switch, repeat and target only
trials of the current experiment. For each functional run of each participant, single-level
GLM estimation was performed with regressors identical to those in the univariate
analysis as described above. Activation maps for face vs. fixation and place vs. fixation
during switch and repeat correct trials were collected from each functional run. These
GLM analyses were performed within standard MNI space.

Another set of testing data was created based on a separate GLM analysis
with correct cue (switch and repeat) trials separately coded at each coherence level
(20/40/80%). For each functional run of each participant, a single-level GLM estimation
was performed, and activation maps for face vs. fixation and place vs. fixation were
collected for each coherence level.

For each training subset, the classifier was tested on whether it correctly
classified the performed task (face or place task) for each trial condition (i.e.
switch/repeat/target only and coherence level). Classification performance was then
collected from all functional runs and averaged within participants for each searchlight.
The performance of classification was calculated as the accuracy minus chance level for
bivariate classification. Accuracy maps were then averaged across testing data sets
within participants.

Accuracy maps for switch, repeat and target only trials were first averaged
across training subset models within participants, and averaged accuracy maps were
collected from all participants. Voxel-wise one-sample group-mean test was performed
for each trial condition, with a procedure similar to that in the univariate analysis as
stated above. In order to explore brain regions showing differential classification
accuracy between switch and repeat trials, voxel-wise group-level paired test was
performed, and significance was tested similarly.

Accuracy maps for each coherence level trials were also averaged across
training subset models within participants, and the averaged accuracy maps were
collected from all participants. Voxel-wise one-sample group-mean test was performed

for each coherence level, and significance was tested similarly. In order to examine

coherence effect, the maps for three coherence levels were weighted and summed based

on behavioral accuracy estimated by sigmoid fitting. Voxel-wise one-sample

group-mean test was performed, and significance was tested similarly.

In separate analyses, all group-level tests above were also performed for each

of the 10 training subset models, and we confirmed that overall results were consistent.

Acknowledgements

We thank Drs. Akira Funahashi, Shori Nishimoto, and Teppei Matsui for scientific

comments on the study and manuscript. We thank Ms. Maoko Yamanaka for

administrative assistance.

Funding

This study was supported by Kakenhi (Japan Society for the Promotion of Science)

19H04914, 17K01989, 17H05957, 17H00891, 26350986, and 26120711 to KJ; Kakenhi

18H04953 and 18H05140 to MT; 18H05017 to JC; 17H00891 to KN. This study was

also supported by a grant from Uehara Memorial Foundation, a grant from Takeda

Science Foundation, Keio Gijuku Academic Development Funds, and grant from Keio

Leading-edge Laboratory of Science and Technology to KJ.

Author contributions

K.T. and K.J. designed the experiment and study. K.T., R.A., K.N. and K.J. collected

the data. K.T., K.K. and K.J. analyzed the data. M.T. and J.C. contributed to the

analysis design of imaging data and to the development of machine learning classifier.

K.T., M.T., J.C., K.N. and K.J. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Data and code availability

The datasets and codes supporting the current study are available from the

corresponding author (Koji Jimura, jimura@bio.keio.ac.jp) on request.

References

1. E.K. Miller, J.D. Cohen, An integrative theory of prefrontal cortex function. *Annu. Rev. Neurosci.* **24**, 167-202 (2001).

2. G. Stoet, L.H. Snyder, Neural correlates of executive control functions in the monkey. *Trends Cogn. Sci.* **13**, 228-234 (2009).

3. D. A. Allport, E.A. Styles, S. Hsieh, *Attention and performance XV*: Conscious and nonconscious information processing (The MIT Press, Cambridge, 1994).

4. R.D. Rogers, S. Monsell, Cost of a predictable switch between simple cognitive tasks. *J. Exp. Psychol. Gen.* **124**, 207–231 (1995).

5. A. Dove, S. Pollmann, T. Schubert, C.J. Wiggins, D.Y. von Cramon, Prefrontal cortex activation in task switching: an event-related fMRI study. *Cognitive Brain Res.* **9**, 103-109 (2000).

6. M.F. Rushworth, R.E. Passingham, A.C. Nobre, Components of switching intentional set. *J. Cogn. Neurosci.* **14**, 1139–1150 (2002).

7. S.A. Bunge, J.D. Wallis, A. Parker, M. Brass, E.A. Crone, E. Hoshi, K. Sakai, Neural circuitry underlying rule use in human and nonnuman primates. *J. Neurosci.* **25**, 10347-10350 (2005).

8. E.A. Crone, C. Wendelken, S.E. Donohue, S.A. Bunge, Neural evidence for dissociable components of task-switching. *Cereb. Cortex* **16**, 475-486 (2006).

9. J. Derrfuss, M. Brass, J. Neumann, D.Y. von Cramon, Involvement of the inferior frontal junction in cognitive control: Meta-analyses of switching and Stroop studies. *Hum. Brain Mapp.* **25**, 22-34 (2005).

10. N. Yeung, L.E. Nystrom, J.A. Aronson, J.D. Cohen, Between-task competition and cognitive control in task switching. *J. Neurosci.* **26**, 1429–1438 (2006).

11. C. Kim, S.E. Cilles, N.F. Johnson, B.T. Gold, Domain general and domain preferential brain regions associated with different types of task switching: a meta-analysis. *Hum. Brain Mapp.* **33**, 130–142 (2012).

12. H. Malagon-Vina, S. Ciocchi, J. Passecker, G. Dorffner, T. Klausberger, Fluid network dynamics in the prefrontal cortex during multiple strategy switching. *Nat. Commun.* **9**, 309 (2018).

13. G. B. Bissonette, M. R. Roesch, Neurophysiology of rule switching in the corticostriatal circuit. *Neuroscience* **345**, 64-76 (2017).

14. E.F. Fouragnan, B.K.H. Chau, D. Folloni, N. Kolling, L. Verhagen, M.

Klein-Flügge, L. Tankelevitch, G.K. Papageorgiou, J-.F. Aubry, J. Sallet, M.F.S.

Rushworth. The macaque anterior cingulate cortex translates counterfactual choice

value into actual behavioral change. *Nat. Neurosci.* **22**, 797-808 (2019).

15. D.E. Nee, M., D'Esposito. The hierarchical organization of the lateral prefrontal

cortex. *eLife* **5**, e12112 (2016).

16. J.I. Gold, M.N. Shadlen, The neural basis of decision making. *Annu. Rev. Neurosci.*

30, 535-574 (2007).

17. T.D. Hanks, C. Summerfield, Perceptual decision making in rodents, monkeys, and

humans. *Neuron* **93**, 15-31 (2017).

18. W.T. Newsome, E.B.A. Pare, A selective impairment of motion perception

following lesions of the middle temporal visual area (MT). *J. Neurosci.* **8**,

2201-2211 (1988).

19. M. Corbetta, F.M. Miezin, S. Dobmeyer, G.L. Shulman, S.E. Petersen, Selective

and divided attention during visual discriminations of shape, color, and speed:

functional-anatomy by positron emission tomography. *J. Neurosci.* **11**, 2383-2402

(1991).

20. T. Kurikawa, T. Haga, T. Handa, R. Harukuni, T. Fuai, Neuronal stability in medial

frontal cortex set individual variability in decision-making *Nat. Neurosci.* **21**,

1764-1773 (2018).

21. M.N. Shadlen, K.H. Britten, W.T. Newsome, J.A Movshon, A computational

analysis of the relationship between neuronal and behavioral responses to visual

motion. *J. Neurosci.* **16**, 1486-1510 (1996).

22. M.S. Beauchamp, R.W. Cox, E.A. Deyoe, Graded effects of spatial and featural

attention on human area MT and associated motion processing areas. *J.*

Neurophysiol. **78**, 516-520 (1997).

23. A.C. Huk, R.F. Dougherty, D.J. Heeger, Retinotopy and functional subdivision of

human areas MT and MST. *J. Neurosci.* **22**, 7195-7205 (2002).

24. A.S. Kayser, D.T. Erickson, B.R. Bushsdaum, M. D'Esposito, Neural

representations of relevant and irrelevant features in perceptual decision making. *J.*

Neurosci. **30**, 15778-15789 (2010).

25. N. Kanwisher, J. McDermott, M.N. Chun, The fusiform face area: a module in

human extrastriate cortex specialized for face perception. *J. Neurosci.* **17**, 4302–

4311 (1997).

26.   G. McCarthy, A. Puce, J.C. Gore, T. Allison, Face-specific processing in the human

1062   fusiform gyrus. *J. Cogn. Neurosci.* **9**, 605-610 (1997).

1063   27.   A. Ishai, L.G. Ungerleider, A. Martin, J.L. Schouten, J.V. Haxby, Distributed

1064   representation of objects in the human ventral visual pathway. *Proc. Natl Acad. Sci.*

1065   *USA* **96**, 9379–9384 (1999).

1066   28.   A. Gazzaley, J.W. Cooney, K. McEvoy, R.T. Knight, M. D'Esposito, Top-down

1067   enhancement and suppression of the magnitude and speed of neural activity. *J.*

1068   *Cogn. Neurosci.* **17**, 507-517 (2005).

1069   29.   W.A. Freiwald, D.Y. Tsao, Functional compartmentalization and viewpoint

1070   generalization within the macaque face-processing system. *Science* **330**, 845-851

1071   (2010).

1072   30.   R. Epstein, N. Kanwisher, A cortical representation of the local visual environment.

1073   *Nature* **392**, 598–601 (1998).

1074   31.   E. Koechlin, C. Ody, F. Kouneiher, The architecture of cognitive control in the

1075   human prefrontal cortex. *Science* **302**, 1181-1185 (2003).

1076   32.   D. Badre, Cognitive control, hierarchy, and the rostro-caudal organization of the

1077   frontal lobes. *Trends Cogn. Sci.* **12**, 193-200 (2008).

1078   33.   K. Jimura, T.S. Braver,　Age-related shifts in brain activity dynamics during task

1079   switching. *Cereb. Cortex* **20**, 1420–1431 (2010).

1080   34.   Y. Kamitani, F. Tong, Decoding the visual and subjective contents of the human

1081   brain. *Nat. Neurosci.* **8**, 679–685 (2005).

1082   35.   J.D. Haynes, G. Rees, Decoding mental states from brain activity in humans. *Nat.*

1083   *Rev. Neurosci.* **7**, 523-534 (2006).

1084   36.   K.A. Norman, S.M. Polyn, G.J. Detre, J.V. Haxby, Beyond mind-reading:

1085   multi-voxel pattern analysis of fMRI data. *Trends Cogn. Sci.* **10**, 424-430 (2006).

1086   37.   S. Nishimoto, A.T. Vu, T. Naselaris, Y. Benjamini, B. Yu , J.L. Gallant,

1087   Reconstructing visual experiences from brain activity evoked by natural movies.

1088   *Curr. Biol.* **21**, 1641-1646 (2011).

1089   38.   J. Chikazoe, D. Lee, N. Kriegeskorte, A.K. Anderson, Distinct representation of

1090   basic taste qualities in human gustatory cortex. *Nat. Comm.* **10**, 1048 (2019).

1091   39.   X. Wang, X. Liang, Z, Jiang, B. A. Nguchu, Y. Zhou, Y. Wang, et al. Decoding and

1092   mapping task states of the human brain via deep learning. *Hum. Brain. Mapp.* **41**,

1093   1505-1519 (2020).

40. V.N. Vapnik, Statistical Learning Theory (Wiley, New York, 1998).

41. M. Misaki, Y. Kim, P.A. Bandettini, N. Kriegeskorte, Comparison of multivariate

classifiers and response normalization for pattern-information fMRI. *NeuroImage*

53, 103-118 (2010).

42. K. Jimura, R.A. Poldrack, Analyses of regional-average activation and multivoxel

pattern information tell complementary stories. *Neuropsychologia* **50**, 544-552

(2012).

43. K. Nakahara et al., Associative-memory representations emerge as shared spatial

patterns of theta activity spanning the primate temporal cortex. *Nat. Comm.* **7**,

11827 (2016).

44. A. Krizhevsky, I. Sutskever, G.E. Hinton, Imagenet classification with deep

convolutional neural network. *Adv. Neural Inf. Proc. Sys.* **25**, 1097-1105 (2012).

45. Y. LeCun, Y. Bengio, G. Hinton, Deep learning. *Nature* **521**, 436-444 (2015).

46. R.R. Selvaraju, et al. Grad-CAM: Visual Explanations from Deep Networks via

Gradient-Based Localization. *2017 IEEE International Conference on Computer*

Vision (ICCV), Venice, 618-626 (2017).

47. K. Simonyan, A. Zisserman, Very deep convolutional networks for large-scale

image recognition. Preprint at https://arxiv.org/pdf/1409.1556.pdf (2015).

48. J. Donahue et al., Decaf: A deep convolutional activation feature for generic visual

recognition. *Proceedings of the 31st International Conference on Machine*

Learning, PMLR **32**, 647-655 (2014).

49. V. Mante, D. Sussillo, K.V. Shenoy, W.T. Newsome, Context-dependent

computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78-84 (2013).

50. J. Zhang, N. Kriegeskorte, J.D. Carlin, J.B. Rowe, Choosing the rules: Distinct and

overlapping frontoparietal representations of task rules for perceptual decisions. *J.*

Neurosci. **33**, 11852-11862 (2013).

51. H. Kumano, Y. Suda, T. Uka, Context-dependent accumulation of sensory evidence

in the parietal cortex underlies flexible task switching. *J. Neurosci.* **36**, 12192–

12202 (2016).

52. M.W. Cole et al., Multi-task connectivity reveals flexible hubs for adaptive task

control. *Nat. Neurosci.* **16**, 1348–1355 (2013).

53. M. D'Esposito, B.R. Postle, The cognitive neuroscience of working memory. *Annu.*

Rev. Psychol. **66**, 115-142 (2015).

54. S.M. Courtney, L.G. Ungerleider, K. Keil, J.V. Haxby, Transient and sustained
activity in a distributed neural system for human working memory. *Nature* **386**,
608-611 (1997).

55. R. Desimone, J. Duncan, Neural mechanisms of selective visual-attention. *Annu.*
Rev. Neurosci. **18**, 193-222 (1995).

56. S. Kastner, L.G. Ungerleider, Mechanisms of visual attention in the human cortex.
Annu. Rev. Neurosci. **23**, 315-341 (2000).

57. M. Corbetta, G.L. Shulman, Control of goal-directed and stimulus-driven attention
in the brain. *Nat. Rev. Neurosci.* **3**, 201-215 (2002).

58. J.A. Lewis-Peacock, B.R. Postle, Temporary activation of long-term memory
supports working memory. *J. Neurosci.* **28**, 8765-8771 (2008).

59. D.H. Brainard, The psychophysics toolbox. *Spatial Vision* **10**, 433–436 (1997).

60. M.Y. Chen, K. Jimura, C.N. White, W.T. Maddox, R.A. Poldrack, Multiple brain
networks contribute to the acquisition of bias in perceptual decision-making. *Front.*
Neurosci. **9**, 63 (2015).

61. K.H. Britten, M.N. Shadlen, W.T. Newsome, J.A. Movshon, Responses of neurons
in macaque MT to stochastic motion signals. *Visual Neurosci.* **10**, 1157-1169
(1993).

62. J. Palmer, A.C. Huk, M.N. Shadlen, The effect of stimulus strength on the speed
and accuracy of a perceptual decision. *J. Vision* **5**, 376-404 (2005).

63. S. Moeller, E. Yacoub, C.A. Olman, E. Auerbach, J. Strupp, N. Harel, K. Uğurbil,
Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial
parallel imaging with application to high spatial and temporal whole-brain fMRI.
Magn. Reson. Med. **63**, 1144-1153 (2010).

64. D. B. Worsley, K. J. Friston, Analysis of fMRI time-series revisited again.
NeuroImage **2**, 173-181 (1995).

65. T.E. Nichols, A.P. Holmes, Nonparametric permutation tests for functional
neuroimaging: a primer with examples. *Hum. Brain Mapp.* **15**, 1-25 (2001).

66. A. Eklund, T.E. Nichols, H. Knutsson, Cluster failure: Inflated false positives for
fMRI. *Proc. Natl Acad. Sci. USA* **113**, 7900-7905 (2016).

67. T. Yarkoni, R.A. Poldrack, T.E. Nichols, D.C. van Essen, T.D. Wager, Large-scale
automated synthesis of human functional neuroimaging data. *Nat. Methods* **8**,
665-670 (2011).

68. K.J. Friston, L. Harrison, W. Penny, Dynamic causal modelling. *NeuroImage* **19**,
1273-1302 (2003).

69. S. Konishi et al., Hemispheric asymmetry in human lateral prefrontal cortex during
cognitive set shifting. *Proc. Natl Acad. Sci. USA* **99**, 7803-7808 (2002).

70. K.J. Friston et al., Bayesian model reduction and empirical Bayes for group (DCM)
studies. *NeuroImage* **128**, 413-431 (2016).

71. W.D. Penny et al., Comparing families of dynamic causal models. *PLoS Comput.*
Biol. **6**, e1,000709 (2010).

72. J. Deng et al,. ImageNet: A large-scale hierarchical image database. *2009 IEEE*
Conference on Computer Vision and Pattern Recognition, Miami, FL, 248-255
(2009).

73. S.J. Pan, Q.A. Yang, A Survey on Transfer Learning. *IEEE T. Knowl. Data En.* **22**,
1345-1359 (2010).

74. M.F. Glasser et al., The human connectome project's neuroimaging approach. *Nat.*
Neurosci. **19**, 1175-1178 (2016).

75. D.M. Barch et al., Function in the human connectome: task-fMRI and individual
differences in behavior. *NeuroImage* **80**, 169-189 (2013).

76. M.F. Glasser et al., A multi-modal parcellation of human cerebral cortex. *Nature*
536, 171-178 (2016).

77. N. Kriegeskorte, R. Goebel, P. Bandettini, Information-based functional brain
mapping. *Proc. Natl Acad. Sci. USA* **103**, 3863–3868 (2006).

1181

1   **Supplementary materials**

3   Title: Flexible decision under ambiguous cues reverses

4   neural signaling from occipitotemporal to prefrontal cortex

7   Authors: Kaho Tsumura,[1] Keita Kosugi,[1] Ryuta Aoki,[2] Masaki Takeda, [2]

8   Junichi Chikazoe,[3] Kiyoshi Nakahara,[2] Koji Jimura[1,2]

11   Affiliations: [1] Department of Biosciences and Informatics,

12   Keio University, Yokohama, Japan

13   [2] Research Center for Brain Communication,

14   Kochi University of Technology, Kami, Japan

15   [3] Supportive Center for Brain Research,

16   National Institute for Physiological Sciences, Okazaki, Japan

Correspondence should be addressed to:

Koji Jimura, Ph.D.

Department of Biosciences and Informatics,

Keio University,

3-14-1 Hiyoshi Kohoku-ku,

Yokohama, 223-0061, Japan

Email: jimura@bio.keio.ac.jp

Phone: +81-45-566-1625

Supplementary figures

cont.

38
Fig. S1. Effective connectivity analysis. (A-B) Parameters estimated without parametric empirical
Bayes. (A)   Switching effect. (B)   Coherence effect. (C-D) The numbers of ROIs in the model was
changed.   (C) Switching effect was examined in a model involving the lateral prefrontal cortex (lPFC),
fusiform face area (FFA), and parahippocampal place area (PPA) for both of the face and place tasks. (D)
Coherence effect was examined in a model involving lPFC, FFA, PPA, and middle temporal (MT)
regions. (E-F) The definition of ROIs. (E) Switching effect. (F) Coherence effect. The other analysis
procedures were identical to those in Figs. 2D-E. (G-H) Task unrelated intrinsic connectivity, estimations
of "A" matrix in the dynamic causal modeling. (G) Switching effect (H) coherence effect. Gray solid
lines indicate statistical significance.

48
49

Fig. S2. Training and testing procedure for the convolutional neural network classifier (CNN).

(A) VGG16 classifier pre-trained for concrete item pictures available from ImageNet. The classifier

consists of 5-tier convolution layers and 3-tier full-connected layers, and is capable of classifying pictures
of concrete items into 1,000 categories. (B) 3D surface maps (*left*) are mapped to 2D flat maps (*middle*).
For illustration purpose, colored areas are cortical subregions functionally and anatomically segmented in
prior work (ref. 42). Sample flattened 2D z-maps for the contrast face vs. fixation and place vs. fixation
(*right*). (C) Training procedures for 2D-flat images. ImageNet-pretrained VGG16 model (A) was first
retrained such that it classified activation maps for the face and place task during working memory task
distributed by Human Connectome Project (HCP) (top). The retrained model was further retrained such
that it classifies activation maps for the face and place task during target only trials in the current
experiment (*bottom*). (D) Testing procedure. The classifier trained by the two stepwise procedure (C) was
tested for classification of face and place tasks during cued trials involving switch and repeat trials with
different coherence levels.

Fig. S3. Classification accuracy of convolutional neural network during training epoch. Vertical and
horizontal axes indicate validation accuracy, and training epoch, respectively. (A) Validation accuracy for
face/place classification of the current target-only trials. Purple line indicates the learning curve of a
model trained based on the activation maps of target-only trials from a randomly initialized parameter.
Green line indicates the learning curve of a model re-trained by the activation maps of target-only trials
from pre-trained parameter by ImageNet. Red line indicates the learning curve of a model re-trained by
the activation maps of target-only trials from the pre-trained parameter by Human Connectome Project
(HCP) working memory maps and ImageNet (i.e. green line). Error bars indicate standard error of the
mean across cross-validation testing. (B) Validation accuracy for face/place classification of working
memory trials provided by HCP. Orange line indicates the learning curve of a model trained based on the
activation maps of working memory trials from a randomly initialized parameter. Blue line indicates the
learning curve of a model trained by activation maps of HCP working memory trials from a pre-trained
parameter by ImageNet. Error bars indicate standard error of the mean across cross-validation testing.

Face vs. Place

Fig. S4. Statistical activation z-map for contrast face vs. place tasks in univariate analysis. Data are derived from those shown in Fig. 2C, but maps are overlaid onto 2D-flat map of the cortical surface of the brain. Hot and cool colors indicate greater activity in face and place tasks, respectively. Closed lines indicate prominent activity during face (gray) and place (black), respectively, as identified by a group-level analysis (Table S3). The fusiform face area (FFA) and parahippocampal place area (PPA) are indicated by red arrow heads.

Fig. S5. Classification accuracy and weight maps of CNN classifier. A leave one-subject-out

procedure was used in training and validation of the CNN classifier. (A) Classification accuracy for each

task condition. (B) Visualization of aggregated weight contrasts for switch vs. repeat trials in face task

(*left*) and place task (*right*). Occipitotemporal regions in rectangular boxes with green broken lines were

expanded below. (C) Visualization of weight contrast for coherence effect in face task (*left*) and place

task (*right*). (D) Regions of interests (ROIs) analysis. All formats are similar to those in Figs. 3B/D-F.

101
102
Fig. S6. The support vector machine classifier trained by target-only trials. (A) Classification
accuracy for cue trials. Error bars indicate standard error of the mean across participants. Formats are
similar to those in Fig. 2B. (B) Weight map of SVM. Maps are overlaid onto flat maps of the cortical
brain. Hot and cool colors indicate high and low weights, respectively. Gray and black lines overlaid on
flat map indicate clusters significantly activated during face and place tasks in univariate analysis,
respectively (Figs. 2C and S4). FFA and PPA was indicated by red arrow heads.
109
110

111
Fig. S7. Searchlight multi-variate pattern analysis. Hot and cool colors indicate statistical level for
classification accuracy relative to chance level. Maps are overlaid onto 3D surface of the brain and
displayed from a ventral view. White and black closed lines overlaid onto 3D surface of the brain indicate
significant cluster for contrast face vs. place tasks in univariate analysis, respectively (Figs. 2C and S4).

The fusiform face area (FFA) and parahippocampal place area (PPA) are indicated by blue arrow heads. (A) Switch trial. (B) Repeat trial. (C) 20% coherence trial. (D) 40% coherence trial. (E) 80% coherence trial. (F) Accuracy difference between switch and repeat trials with a model trained by Human Connectome Project (HCP) data. Hot and cool colors indicate higher accuracy in switch and repeat trials, respectively. (G) Differential classification accuracy depending on coherence effect with a model trained by HCP data. Hot and cool colors indicate higher accuracy in high and low coherent trials, respectively. (H) Accuracy difference between switch and repeat trials with model trained by target-only trials of the current dataset. (I) Differential classification accuracy depending on the coherence effect with model trained data of target-only trials. The formats are similar to those in panel (F).

127   **Supplementary tables**

129   Table S1.   Brain regions showing significant signal increase and decrease in the contrast of switch vs.

130   repeat trials. Coordinates are listed in MNI space. Positive and negative z-values indicate increase in

131   switch and repeat trials, respectively. BA indicates Brodmann areas and is approximate.

Area	x	y	z	z-value	BA
Frontal cortex	-30	0	54	5.32	6
	-6	8	52	4.98	6
	-18	12	62	4.80	6
	12	12	68	4.66	6
	-42	0	48	4.52	6
	0	28	32	4.42	32
	-6	4	70	4.12	6
	10	18	28	4.00	32
	10	14	56	3.59	6
	-40	12	26	4.62	44
	-38	-2	36	3.61	6
	22	0	48	4.16	6
	38	0	40	3.63	6
Parietal cortex	-28	-70	36	5.62	39
	0	-36	26	5.49	23
	20	-60	28	5.55	31
	-4	-16	32	5.35	23
	-10	-68	44	4.98	7
	-30	-56	44	4.67	39
	-50	-40	50	4.63	40
	-38	-48	50	4.58	40
	-14	-68	28	4.39	7
	6	-38	38	4.02	23
	16	-68	42	4.00	7
	16	-66	54	3.87	7
	-36	-82	26	3.64	19
	-4	-44	12	3.63	30
	12	-42	28	3.33	23

134 Table S2.   Brain regions showing significant parametrical effect with motion coherence. Coordinates are
135 listed in MNI space. Positive and negative z-values indicate increase in high and low coherent trials,
136 respectively. *: corrected within motion-related brain regions (see Methods).

Area	x	y	z	z-value	BA
Frontal cortex	0	18	50	-5.90	8
	10	24	46	-5.59	8
	54	20	38	-5.43	9/44
	46	14	4	-5.43	44
	52	14	16	-5.21	44
	-56	12	28	-5.16	44
	36	4	54	-4.62	6
	48	6	34	-4.62	6
	-32	0	50	-4.49	6
	-38	10	32	-4.43	8
	8	30	28	-4.41	8
	42	24	28	-4.39	9
	2	42	-16	4.37	11
	4	30	-18	4.33	11
	20	12	46	-4.31	6
	-50	8	10	-4.25	44
	-42	24	26	-4.08	9/44
	38	12	20	-3.82	44
	40	36	-4	-3.82	47
	-8	46	-6	3.80	10
	24	8	62	-3.79	6
	34	34	34	-3.78	9
	-52	18	-6	-3.67	47
	48	38	14	-3.35	46
	-44	38	28	-3.21	9/10
Parietal cortex	-18	-56	74	4.70	7
	-16	-48	36	4.37	31
	-12	-68	30	-4.28	7
	-24	-64	38	-4.25	39
	-30	-52	46	-4.19	7/39
	14	-48	18	-4.19	23
	58	-46	36	-4.08	39/40
	-48	-40	48	-4.07	40
	26	-40	64	4.03	1/5
	20	-58	26	-4.02	23
	42	-50	52	-3.86	40
	-30	-42	56	3.85	5/7
	-40	-46	40	-3.66	39
	6	-46	32	3.63	23
	42	-46	38	-3.50	39
	32	-42	46	-3.27	7
Temporal cortex	-42	-54	-8	-5.45	37
	-30	-58	-10	-4.10	37
	56	-64	2	3.75*	37/19
Others	32	24	4	-5.66	insula
	-30	26	0	-5.56	insula
	0	-32	-2	-4.66	brain stem
	12	-18	-8	-4.30	brain stem
	8	-10	4	-3.83	thalamus
	-16	-28	-4	-3.60	thalamus

138 Table S3. Brain regions showing significant signal increase and decrease in the contrast of face vs.
139 place in target-only trials. Coordinates are listed in MNI space. Positive and negative z-values indicate
140 increase in face and place trials, respectively.

Area	x	y	z	z-value	BA
Frontal cortex	4	32	46	5.00	8
	38	28	12	4.97	45
	8	40	38	4.23	9
	46	2	28	3.96	6
	48	42	12	3.49	46
	48	18	40	3.40	8
	-6	36	36	3.38	8
Parietal cortex	-22	-74	44	-5.90	7
	-20	-58	12	-5.88	23
	-18	-70	58	-5.79	7
	-10	-76	50	-5.38	7
	18	-74	44	-4.78	7
	2	-60	30	4.60	31
	12	-68	62	-4.47	7
	40	-74	40	-4.37	39
	2	-48	18	4.09	23
Temporal cortex	26	-38	-14	-8.03	37
	-28	-44	-12	-7.46	37
	-28	-30	-20	-6.81	37
	42	-58	-14	5.72	37
	-44	-54	-20	4.91	37
	-10	-46	4	-4.38	18/30
	52	-64	0	4.06	37
	60	-64	-10	3.53	37
Occipital cortex	-40	-82	22	-6.69	19
	44	-72	-12	6.36	19
	40	-74	20	-6.27	19
	-48	-78	-10	5.75	19
	-20	-66	22	-5.72	19
	12	-46	4	-5.65	18
	18	-56	16	-5.61	18/23
	-26	-66	-12	-5.14	19
	24	-64	26	-4.18	19
	-32	-86	38	-4.08	19
	-48	-82	2	3.73	19
	-40	-70	-4	3.41	19
Others	-18	-6	-16	5.09	amygdala
	-30	2	-22	4.54	amygdala
	32	18	2	3.80	insula

142 Table S4. Brain regions showing significant accuracy in classifying task dimension (face/place) for
143 target-only trials. Coordinates are listed in MNI space. Positive z-values indicate higher accuracy than
144 chance level.

Area	x	y	z	z-value	BA
Frontal cortex	-10	36	42	5.66	8
	-6	24	32	4.05	32
Parietal cortex	24	-68	48	6.67	7
	-28	-70	26	6.57	39
	26	-82	46	6.40	7
	14	-78	56	6.10	7
	24	-68	66	5.78	7
	8	-64	22	5.60	31
	-24	-56	32	5.55	7/39
	-12	-46	24	5.40	23
	10	-84	46	5.34	7/19
	26	-54	28	5.23	23
	-10	-74	40	5.06	7
	-32	-78	38	4.96	39
	14	-60	62	4.86	7
	-26	-60	48	4.79	7
	42	-66	32	4.76	39
	36	-74	50	4.69	39
	-14	-64	56	4.68	7
	-14	-66	32	4.59	7
	50	-76	30	4.47	39
	-8	-76	52	4.44	7
	36	-82	26	4.41	19/39
	30	-68	32	4.32	7
Temporal cortex	32	-50	-18	7.86	37
	-28	-42	-20	7.60	37
	-40	-50	-10	7.46	37
	38	-38	-16	7.34	37
	-38	-32	-16	7.29	37
	-34	-26	-28	6.94	36
	-36	-46	4	6.40	21
	-40	-18	-20	6.36	20
	-40	-36	-2	6.28	21
	-26	-20	-20	6.19	36
	16	-40	0	6.00	18/30/36
	18	-26	-20	5.97	36
	14	-48	14	5.75	23
	44	-60	-16	5.56	37
	-34	8	-26	5.47	38
	-18	-48	2	5.43	30
	-48	-44	-24	5.18	37
	-10	-38	-4	5.01	36
	26	-14	-22	4.88	36
	-28	4	-42	4.82	38
	54	-50	-20	4.78	37
	54	-62	-8	4.75	37
	16	-8	-28	4.59	36
	6	-34	-4	4.28	36
	-38	-6	-20	4.12	22
Occipital cortex	34	-72	-12	7.26	19/37
	22	-70	12	6.96	17

	-44	-80	26	6.91	19
	16	-70	32	6.60	19
	22	-52	0	6.51	18/19
	30	-60	18	6.43	18
	-40	-68	26	6.37	19/39
	32	-84	-18	6.04	19
	-44	-76	14	6.01	19
	48	-76	8	6.00	19
	20	-64	0	5.95	19
	-30	-88	-10	5.94	18
	-32	-68	-10	5.90	19
	36	-84	-4	5.90	18
	20	-84	34	5.76	19
	46	-84	-14	5.67	19
	-32	-64	16	5.54	19
	-12	-64	20	5.43	18
	18	-90	22	5.41	18
	-32	-78	8	5.22	18
	-26	-56	-2	5.11	19
	-42	-86	0	4.95	18
	30	-96	10	4.50	18
	4	-60	8	4.49	18
	22	-76	-2	4.37	18
	54	-76	-4	4.35	19
	32	-84	8	4.27	19
Others	-24	-28	-8	7.66	hippocampus
	28	-42	4	6.64	hippocampus
	16	-46	-24	6.51	cerebellum
	-16	-32	-22	6.37	cerebellum
	-34	-68	-22	6.36	cerebellum
	-30	-56	-26	6.25	cerebellum
	38	-26	-8	6.14	hippocampus
	-4	-34	6	6.08	callosum
	26	-26	-8	5.63	hippocampus
	-48	-56	-32	5.59	cerebellum
	14	-58	-14	5.58	cerebellum
	-14	-14	-10	5.52	brain stem
	-26	-10	-10	5.48	hippocampus
	12	-38	-14	5.30	cerebellum
	-50	-68	-26	4.89	cerebellum
	10	-42	-40	4.88	cerebellum
	32	-32	-34	4.75	cerebellum
	-4	-24	-2	4.70	thalamus
	-22	-4	-22	4.70	hippocampus
	20	-70	-20	4.64	cerebellum
	16	-12	-12	4.63	hippocampus
	-32	6	-14	4.62	insula
	32	-72	-26	4.59	cerebellum
	16	2	-12	4.39	amygdala
	48	-56	-30	4.20	cerebellum
	28	-2	-14	4.15	amygdala
	-4	-38	-38	4.10	cerebellum
	8	-44	-56	3.96	cerebellum
	10	-54	-38	3.83	cerebellum
	-16	0	-4	5.05	globus pallidus

146 Table S5.   Brain regions showing significant accuracy difference for face-place classification between
147 switch and repeat trials. Coordinates are listed in MNI space. Positive and negative z-values indicate
148 higher accuracy in switch and repeat trials, respectively.

Area	x	y	z	z-value	BA
Parietal cortex	34	-62	22	4.08	39
	14	-66	22	3.75	31
	22	-56	28	3.26	23/31
Temporal cortex	36	-50	-18	5.79	37
	-30	-48	-16	5.54	37
	-36	-22	-28	5.35	36
	22	-42	-16	5.24	37
	-34	-34	-16	4.93	36
	44	-54	-4	4.71	37
	36	-62	-8	4.33	37
	54	-62	-12	4.30	37
	20	-30	-12	3.88	36
	-40	-48	0	3.84	37
	26	-58	-14	3.80	37
	-26	-20	-20	3.38	36
	-42	-24	-16	3.26	20
	44	-50	8	3.25	37
Occipital cortex	22	-72	12	5.01	17
	-36	-72	4	4.91	18/19
	32	-74	-4	4.48	19
	-42	-84	10	4.05	19
	36	-88	-4	3.98	18
	44	-80	2	3.96	19
	22	-48	0	3.92	18
	-30	-82	-2	3.90	18
	24	-72	28	3.78	19
	18	-62	-4	3.28	19
Others	12	-52	-12	4.23	cerebellum
	-38	-34	-4	3.80	caudate
	28	-46	-30	3.69	cerebellum
	30	-38	-2	3.30	hippocampus

151   Table S6.   Brain regions showing significant accuracy modulation for face and place tasks depending on
152   coherence level. Coordinates are listed in MNI space. Positive and negative z-values indicate higher
153   accuracy in high and low coherent trials, respectively.

Area	x	y	z	z-value	BA
Temporal cortex	-30	-62	-8	4.85	37
	-24	-38	-8	4.29	36
	-34	-52	-20	4.25	37
	-42	-40	-18	4.20	37
	-14	-28	-12	3.78	36
Occipital cortex	-18	-54	-8	3.78	19
Others	-38	-32	-4	4.18	caudate
	-6	-30	0	3.37	thalamus
	-34	-46	-32	3.17	cerebellum

156   Table S7.   Brain regions showing significant accuracy to classify task dimension (face/place) in switch
157   trials. Coordinates are listed in MNI space. Positive z-values indicate higher accuracy than chance level.

Area	x	y	z	z-value	BA
Parietal cortex	14	-66	20	6.45	31
	24	-54	10	5.48	23
	2	-52	20	5.43	23
	20	-70	34	5.35	19
	26	-58	32	5.10	31
	22	-74	52	5.08	7
	14	-54	26	4.77	23
	-46	-66	14	4.77	39
	-18	-52	22	4.27	23
	-14	-78	46	4.19	7
	42	-76	36	4.14	39
	50	-54	18	4.04	39
	34	-68	30	4.02	39
	-6	-64	20	3.92	31
	12	-70	60	3.87	7
	28	-60	62	3.84	7
	24	-58	46	3.68	7
	8	-70	44	3.64	7
	-34	-82	42	3.63	39
	34	-82	44	3.39	7
	-44	-68	38	3.37	39
	-8	-56	30	3.28	31
	34	-68	44	3.26	39
Temporal cortex	-32	-48	-12	8.39	37
	34	-50	-16	7.26	37
	-22	-28	-20	6.88	36
	-32	-38	-22	6.82	37
	16	-32	-8	6.73	36
	-34	-18	-24	6.64	36
	-42	-38	-4	6.51	21
	34	-64	-8	6.07	37
	34	-30	-24	5.61	37
	22	-32	-20	4.99	37
	42	-46	0	4.78	21
	54	-52	2	4.73	37
	8	-38	8	4.03	30
	58	-52	-20	4.03	37
	54	-68	-8	3.86	37
	42	-20	-28	3.59	20
Occipital cortex	16	-54	-4	6.28	19
	-34	-60	-4	6.18	19/37
	44	-58	-2	5.56	19/37
	26	-66	14	5.48	17/23
	-48	-84	8	5.46	19
	-36	-72	4	5.29	18/19
	32	-76	-10	5.24	19
	-28	-72	-8	5.13	19
	-48	-78	22	5.00	19
	-38	-74	28	4.98	19
	-20	-54	-2	4.90	19
	44	-80	2	4.77	19

18	-78	24	4.74	18/19
-32	-62	22	4.67	19
-34	-88	6	4.59	18
6	-66	8	4.54	18
22	-82	38	4.51	19
54	-68	22	4.44	19
-28	-68	12	4.30	17
56	-64	4	4.11	19/37
36	-94	-2	4.09	18
22	-90	22	4.03	19
22	-80	12	3.92	17
-28	-86	-10	3.75	18
-46	-86	-4	3.66	18/19
28	-90	-14	3.61	18
-18	-74	16	3.41	17
-24	-88	14	3.41	19
32	-76	4	3.32	18
-8	-46	2	3.25	18
-36	-84	20	3.22	19

Others				
	-26	-32	-6	6.96	hippocampus
	28	-26	-8	6.34	hippocampus
	20	-44	-18	6.30	cerebellum
	-34	-28	-14	6.20	hippocampus
	30	-46	-28	6.06	cerebellum
	28	-64	-20	5.94	cerebellum
	-16	-60	-20	5.36	cerebellum
	-22	-46	-28	5.14	cerebellum
	34	-36	0	4.90	caudate
	-22	-42	2	4.86	hippocampus
	40	-28	-8	4.46	caudate
	2	-56	0	4.28	cerebellum
	-32	-80	-24	4.02	cerebellum
	-36	-54	-28	3.67	cerebellum
	38	-16	-16	3.30	hippocampus
	-40	-66	-22	3.20	cerebellum

159   Table S8.   Brain regions showing significant accuracy to classify task dimension (face/place) in repeat
160   trials. Positive z-values indicate higher accuracy than chance level.

Area	x	y	z	z-value	BA
Parietal cortex	0	-50	16	4.84	23
	12	-42	18	4.58	30
	10	-62	26	4.53	31
	36	-50	4	3.88	30
	-14	-56	32	3.66	31
	-6	-58	8	3.62	23
	-22	-52	22	3.53	23
	28	-56	12	3.46	23
	-32	-50	4	3.32	30
Temporal cortex	30	-42	-22	6.04	37
	-34	-50	-10	6.01	37
	30	-56	-18	5.96	37
	36	-26	-20	5.34	36
	38	-38	-14	5.28	37
	-38	-32	-18	4.20	20/36/37
	24	-20	-26	4.15	36
	-30	-16	-26	4.02	36
	-42	-54	-20	3.77	37
	44	-26	-2	3.42	22
	38	-76	-16	3.39	37
	22	-48	-12	3.16	37
Occipital cortex	34	-66	-6	5.06	19/37
	-36	-66	-16	4.92	19
	-46	-78	6	4.83	19
	16	-54	-2	4.79	19
	-32	-94	-14	4.67	18
	34	-88	-2	4.46	18
	-40	-66	20	4.21	19/39
	30	-90	-18	4.18	18/19
	-28	-56	30	4.16	7
	-26	-70	16	4.12	17
	-24	-74	-14	4.09	18
	-14	-64	22	4.08	18
	-34	-80	-10	4.06	19
	24	-70	20	3.84	19
	14	-66	6	3.67	17
	46	-86	6	3.52	19
	-46	-80	24	3.15	19
Others	-34	-36	-4	5.78	caudate
	-26	-40	-26	5.55	cerebellum
	28	-26	-8	5.46	hippocampus
	22	-32	-24	5.16	cerebellum
	30	-12	-8	4.61	putamen
	16	-52	-24	4.38	cerebellum
	36	-38	2	4.30	caudate
	-30	-20	-14	4.15	hippocampus
	-30	-82	-22	3.98	cerebellum
	-24	-60	-24	3.82	cerebellum
	12	-38	-16	3.53	cerebellum
	18	-16	-16	3.47	hippocampus
	30	-36	-34	3.46	cerebellum

161   Table S9.   Brain regions showing significant accuracy to classify task dimension (face/place) in 20%

162   coherence trials. Coordinates are listed in MNI space. Positive z-values indicate higher accuracy than

163   chance level.

Area	x	y	z	z-value	BA
Frontal cortex	26	14	-18	4.54	47
	6	60	36	4.49	9
	14	54	44	3.98	9
	10	42	42	3.67	8/9
Parietal cortex	14	-66	22	4.87	31
	0	-52	18	4.81	23
	20	-52	30	4.63	23
	-48	-66	14	4.40	39
	-48	-76	26	4.40	39
	26	-60	10	4.07	23
	-40	-56	18	3.64	39
	4	-64	32	3.38	31
Temporal cortex	30	-56	-16	6.33	37
	-36	-50	-8	6.32	37
	-26	-46	-20	5.53	37
	32	-40	-24	4.85	37
	32	0	-30	4.49	36
	40	-32	-14	4.42	36
	-32	-54	4	4.19	18/30
	44	-24	-22	3.75	20
	30	14	-32	3.74	38
	58	-66	-12	3.68	37
	58	-58	2	3.39	37
	48	-68	-4	3.36	37
Occipital cortex	36	-64	-4	4.92	19
	-48	-78	8	4.89	19
	16	-54	-4	4.85	19
	32	-84	-4	4.66	18
	-24	-64	-12	4.60	19
	-16	-56	-4	4.41	19
	28	-60	22	4.29	18
	16	-42	0	4.07	18
	54	-72	6	4.03	19/37
	8	-60	10	3.94	18
	-34	-70	-2	3.87	19
	50	-82	0	3.76	19
	-32	-72	-18	3.51	19
	-44	-88	2	3.43	18
	-40	-84	36	3.38	19
Others	-32	-34	-10	6.06	hippocampus
	30	-70	-20	5.01	cerebellum
	-20	-30	-26	4.84	cerebellum
	16	-60	-18	4.71	cerebellum
	-32	-20	-20	4.60	hippocampus
	20	-48	-24	4.47	cerebellum
	22	-30	-26	4.41	cerebellum
	28	-28	-10	4.07	hippocampus
	32	-62	-30	3.52	cerebellum
	30	-42	4	3.52	hippocampus

164 Table S10. Brain regions showing significant accuracy to classify task dimension (face/place) in 40%
165 coherence trials. Coordinates are listed in MNI space. Positive z-values indicate higher accuracy than
166 chance level.

Area	x	y	z	z-value	BA
Parietal cortex	34	-60	-8	5.89	23
	6	-48	18	5.35	23
	-22	-68	48	5.10	7
	20	-46	10	5.08	30
	34	-46	4	4.84	30
	28	-68	50	4.75	7
	8	-60	20	4.75	31
	22	-58	42	4.54	7
	-8	-46	16	4.54	23
	32	-58	6	4.41	23
	-28	-64	60	4.34	7
	24	-52	20	4.32	23
	-16	-60	58	4.18	7
	-16	-56	22	4.17	23
	-28	-62	24	4.13	39
	-40	-74	30	4.08	39
	-2	-58	50	4.04	7
	46	-78	34	3.97	39
	6	-54	56	3.93	7
	-22	-82	46	3.91	7
	4	-46	50	3.74	31
	-14	-68	38	3.69	7
	36	-60	54	3.39	7
Temporal cortex	-34	-38	-10	6.99	36
	30	-52	-16	6.94	37
	-38	-50	-10	6.60	37
	-28	-34	-20	6.29	37
	-30	-60	-14	5.27	37
	38	-30	-24	5.11	37
	46	-42	-6	4.49	21
	-36	-38	-28	4.06	37
	-14	-26	-20	4.02	36
	-22	-50	-10	4.01	19/37
	44	-58	-22	3.82	37
Occipital cortex	-46	-80	10	6.51	19
	20	-70	14	5.35	17
	-34	-72	6	5.24	18/19
	-40	-68	16	5.01	19/39
	10	-60	8	4.97	18
	16	-60	-4	4.79	19
	18	-70	28	4.78	19
	-16	-76	22	4.73	18
	-32	-48	2	4.64	18/30
	-30	-82	-16	4.53	19
	-38	-66	-4	4.50	19
	-36	-82	-4	4.46	18
	36	-82	-2	4.43	18
	-18	-84	34	4.22	19
	30	-64	18	4.21	18
	54	-72	22	3.97	19

	38	-70	0	3.91	19
	32	-78	-14	3.74	19
	-34	-90	10	3.70	18
	28	-76	34	3.62	7/19
	4	-72	8	3.56	17
	28	-90	-12	3.32	18
	26	-70	0	3.20	19
Others	38	-32	-2	6.00	caudate
	-36	-72	-22	5.26	cerebellum
	26	-32	-2	5.25	hippocampus
	10	-48	-4	5.20	cerebellum
	16	-52	-16	5.16	cerebellum
	38	-24	-12	4.78	hippocampus
	14	-34	0	4.44	thalamus
	-32	-52	-30	4.20	cerebellum
	-18	-58	-24	4.13	cerebellum
	-12	-66	-12	4.10	cerebellum
	18	-34	-28	3.88	cerebellum
	4	-34	10	3.75	callosum
	10	-36	-18	3.32	cerebellum
	-32	-20	-16	3.30	hippocampus
	0	-46	4	3.16	cerebellum

169   Table S11.   Brain regions showing significant accuracy to classify task dimension (face/place) in 80%
170   coherence trials. Coordinates are listed in MNI space. Positive z-values indicate higher accuracy than
171   chance level.

Area	x	y	z	z-value	BA
Parietal cortex	18	-62	30	5.46	31
	26	-52	16	5.40	23
	18	-60	48	5.23	7
	2	-50	18	5.04	23
	-16	-52	22	5.01	23
	-20	-56	34	4.93	7
	14	-50	14	4.92	23
	34	-56	28	4.87	39
	6	-66	36	4.28	7
	12	-76	52	4.22	7
	-28	-66	44	4.20	7/39
	46	-62	24	4.05	39
	14	-66	62	4.04	7
	26	-60	58	3.88	7
	-10	-74	50	3.86	7
	-28	-62	64	3.80	7
	28	-60	38	3.71	7
	-12	-46	34	3.58	23
	26	-74	46	3.53	7
	-8	-70	32	3.46	7
	-28	-78	44	3.41	7
	-50	-74	28	3.30	39
Temporal cortex	34	-50	-16	7.50	37
	-32	-48	-12	7.16	37
	-32	-62	-10	6.60	37
	34	-60	-8	6.51	37
	-44	-38	-10	6.06	37
	20	-36	-6	5.91	36
	22	-24	-12	5.74	36
	-40	-46	0	5.67	21/37
	-32	-26	-20	5.26	36
	26	-36	-22	5.10	37
	42	-48	0	4.92	37
	-44	-24	-16	4.78	20
	60	-54	10	4.33	37
	52	-62	-18	4.20	37
	56	-64	2	4.19	37
	26	-8	-32	3.92	36
Occipital cortex	14	-66	18	6.47	18
	-44	-74	14	5.88	19
	18	-50	0	5.79	18
	16	-56	-12	5.41	19
	-34	-72	2	5.34	18/19
	-22	-52	0	5.33	18
	-26	-80	-14	5.25	19
	38	-92	-2	5.14	18
	34	-76	-2	4.97	18/19
	10	-60	8	4.90	18
	-14	-64	22	4.77	18
	44	-80	4	4.66	19

-36	-70	28	4.65	19
20	-66	4	4.61	18
30	-50	-2	4.55	19
-30	-92	-14	4.42	18
24	-72	24	4.36	19
-30	-84	2	4.34	18
-42	-84	30	4.25	19
-26	-70	14	4.21	17
-16	-78	22	4.16	18
18	-86	20	3.96	18
36	-68	14	3.92	19
16	-68	-10	3.84	18/19
16	-80	34	3.70	19
58	-68	22	3.56	19
-44	-80	-6	3.48	19
28	-80	8	3.33	18
-20	-66	-6	3.24	19
-38	-92	-4	3.20	18
Others -28	-28	-8	6.70	hippocampus
36	-40	-28	5.94	cerebellum
-22	-34	-26	5.66	cerebellum
14	-28	-24	5.26	brain stem
-26	-50	-26	5.26	cerebellum
32	-32	-6	5.06	caudate
16	-44	-24	4.57	cerebellum
-26	-14	-22	4.49	hippocampus
-10	-38	-22	4.23	cerebellum
-14	-62	-16	4.05	cerebellum
2	-36	4	3.92	brain stem
36	-78	-22	3.86	cerebellum
26	-14	-20	3.82	hippocampus
-32	-74	-24	3.78	cerebellum
-14	-12	-12	3.74	brain stem
2	-50	0	3.69	cerebellum
-18	-38	4	3.53	hippocampus
22	-66	-24	3.46	cerebellum