A Murine Model for the Transition of *Streptococcus pneumoniae* from Asymptomatic

Colonizer to Systemic Pathogen

Basma H. Joma[*1, 2], Nalat Siwapornchai[*1,] Vijay K. Vanguri[3], Anishma Shrestha[1], Sara E.

Roggensack[1, 4], Bruce A. Davidson[5], Albert K. Tai[6], Anders P. Hakansson[7], Simin N.

Meydani[8], John M. Leong[#1] and Elsa N. Bou Ghanem[#9]

* Co-first authors; B.H.J. and N.S. contributed equally to this work; the order of names was

chosen alphabetically

Co-corresponding authors

[1] Department of Molecular Biology and Microbiology at Tufts University School of Medicine,

Boston, Massachusetts, USA

[2] Graduate Program in Immunology, Tufts Graduate School of Biomedical Sciences, Boston,

USA

[3] UMass Memorial Health Care, University of Massachusetts Medical School, Worcester,

Massachusetts, USA

[4]Graduate Program in Molecular Microbiology, Tufts Graduate School of Biomedical

Sciences, Boston, USA

[5] Department of Anesthesiology, University at Buffalo School of Medicine, Buffalo, New York,

USA

[6]Department of Immunology, Tufts University School of Medicine, Boston, Massachusetts,

USA

[7] Department of Translational Medicine, Lund University, Malmö, Sweden

[8] Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston,

Massachusetts, USA

[9]Department of Microbiology and Immunology, University at Buffalo School of Medicine, Buffalo, New York, USA

Running Title: Murine Model of Influenza-Pneumococcal Co-infection

Address correspondence and reprint requests to John M. Leong (J.M.L.), Department of Molecular Biology and Microbiology, Tufts University School of Medicine, 136 Harrison Avenue, Boston, MA 02114 and Elsa Bou Ghanem (E.N.B.G.), Department of Microbiology and Immunology, University at Buffalo School of Medicine, 955 Main Street, Buffalo, New York, USA. E-mail addresses: john.leong@tufts.edu (J.M.L.) and elsaboug@buffalo.edu (E.N.B.G.)

Keywords: Co-infection, secondary bacterial pneumonia, *Streptococcus pneumoniae*, Influenza A, colonization, aging, neutrophils, inflammation

ABSTRACT

Streptococcus pneumoniae (pneumococcus) resides asymptomatically in the nasopharynx but, as a pathobiont, can progress from benign colonizer to lethal pulmonary or systemic pathogen. Both viral infection and aging are risk factors for serious pneumococcal infections. Previous work established a murine model that featured the movement of pneumococcus from the nasopharynx to the lung upon nasopharyngeal inoculation with influenza A virus (IAV) but did not fully recapitulate the severe disease associated with human co-infection. We built upon this model by establishing pneumococcal nasopharyngeal colonization in mice, then inoculating both the nasopharynx and lungs with IAV. In young (2 months) mice, co-infection triggered bacterial dispersal from the nasopharynx into the lungs, pulmonary inflammation, concomitant disease and mortality in a fraction of mice. Although neutrophils are critical for pneumococcal clearance, IAV infection was associated with inefficient killing of pneumococci by neutrophils *ex vivo* and these cells were dispensable for host resistance during co-infection. In old mice (20-22 months), co-infection resulted in earlier and more severe disease compared to young mice. Aging was not associated with greater bacterial spread from the nasopharynx to the lungs or bloodstream. Rather, old mice displayed a more rapid proinflammatory cytokine response, earlier pulmonary neutrophil influx, lung damage and abrogated neutrophil function. Thus, in this model, IAV fosters the transition of pneumococci from commensals to pathogens through modifying bacterial behavior in the nasopharynx, enhancing pulmonary infection, and compromising neutrophil function. This model faithfully replicates age-associated susceptibility and can provide insight into factors that influence pathobiont behavior in different hosts.

INTRODUCTION

Streptococcus pneumoniae (pneumococcus) is a Gram-positive pathobiont that typically resides asymptomatically in the nasopharynx of healthy individuals (1). It is hypothesized *S. pneumoniae* establishes an asymptomatic biofilm on the nasopharyngeal epithelium by attenuating the production of virulence factors and concomitant inflammation (2-5). However, when immunity is compromised, a common occurrence in aging (6), pneumococci can cause serious disease such as otitis media, pneumonia, meningitis and bacteremia (7). In humans, pneumococcal carriage is believed to be a prerequisite to invasive disease (8, 9). Bacterial isolates from invasive infections are genetically identical to those found in the nasopharynx of patients (9); this and other longitudinal studies have led to the suggestion that invasive disease often involves pneumococcal carriage in the upper respiratory tract (8, 9).

The rate of reported colonization is quite variable among adults and is confounded by differences in detection methods, but colonization may be more prevalent in the elderly (10). A meta-analysis of twenty nine published studies found that in individuals above 60 years of age, conventional culture showed 0-39% carriage, while 3-23% carriage was detected by molecular methods (11). Importantly carriage was higher among nursing home residents (11). In older adults, conventional culture methods estimated carriage to be <5% (12-14) but more recent data from European studies using molecular detection methods indicated that carriage in the elderly ranges from 10-22% (10, 15-17). Thus, carriage rates may be much higher than what was previously estimated in elderly individuals (12-14).

Advanced age increases the risk of invasive pneumococcal disease and pneumococcal pneumonia (7). Aging is associated with immunosenescence, the overall decline in immunity that accompanies aging as well as inflammaging, a low-grade chronic

inflammation that render the elderly more susceptible to pulmonary infections (18).

Polymorphonuclear leukocytes (PMNs), also known as neutrophils, are a crucial

determinant of age-related susceptibility to primary pneumococcal infection (19). These

cells are required to control bacterial burden at the start of infection (20-22). However,

aging is accompanied by impaired PMN anti-bacterial function (23, 24). In addition, aged

hosts experience exacerbated PMN pulmonary influx during primary pneumococcal

pneumonia (19, 25, 26), and persistence of PMNs in the airways beyond the first few days

leads to tissue destruction and systemic spread of infection (27, 28).

In addition to advanced age, epidemiological and experimental data show that

invasive pneumococcal infections are strongly associated with viral infection (29-31). The

risk of pneumococcal pneumonia is enhanced 100-fold by influenza A virus (IAV) infection

(32), resulting in the seasonal peak of pneumococcal disease during influenza outbreaks

(33). Further, *S. pneumoniae* is historically among the most common etiologies of

secondary bacterial pneumonia following influenza and associated with the most severe

outcomes (30, 34-36). Symptoms of secondary bacterial pneumonia include cough,

dyspnea, fever, muscle aches and when severe result in hospitalizations, respiratory

failure, mechanical ventilation, and can lead to death (30, 34-36).

IAV commonly infects the upper respiratory tract, and at this location viral infection

can enhance the nutritional environment for pneumococcus in the nasopharynx, leading to

greater bacterial loads and/or higher rates of bacterial acquisition (4, 37). IAV can also

directly bind the pneumococcal surface and enhance bacterial binding to the pulmonary

epithelium leading to increased colonization (38). In addition, viral infection of the

pulmonary epithelium induces the release of host components, such as adenosine

triphosphate and norepinephrine, that are sensed by biofilm-associated pneumococci,

triggering both the production of pneumococcal virulence factors and the dispersal of

bacteria (4, 37, 39, 40). This in turn facilitates bacterial spread to and colonization of the

lower respiratory tract (4). Murine models featuring sequential lung challenge with IAV first

followed by pneumococcus show the virus can also compromise pulmonary immune

defense against pneumococcus (29, 34, 41). In these models, IAV alters both the lung

environment and the immune response to enhance subsequent bacterial colonization and

tissue damage. For example, IAV pre-infection increases mucus production and fibrosis

and dysregulates ciliary function (34, 42, 43), thus impairing mechanical clearance of

invading bacteria. Viral enzymes, along with virus-elicited inflammation, result in the

exposure of epithelial proteins that promote pneumococcal adherence to and invasion of

host cells (41, 44-46). Further, IAV triggers type I and II interferon (IFN) responses that

impair both the recruitment and antibacterial function of phagocytes key to defense against

pneumococcus (47-50). The combined tissue damage and the compromise in immune

function render the lung more permissive for invasive *S. pneumoniae* infection (34, 45, 47).

These studies indicate that bacterial-viral synergy is multi-factorial and can occur at

different sites with the host.

Notably, advanced age and IAV infection appear to synergistically enhance

susceptibility to pneumococcal lung infections (6, 18, 51, 52). Indeed, individuals ≥ 65

years old account for 70-85% of deaths due to pneumonia and influenza (52). Interestingly,

elderly individuals with influenza-like symptoms were reported to have an increased

pneumococcal carriage rate of 30% (53). In animal models in which bacteria are directly

instilled into the lungs following influenza infection, aging is associated with increase

susceptibility of secondary pneumococcal pneumonia (51). Age-dependent changes in the

expression of key components of innate immune signaling contribute to disease in this co-

infection model (51). However, the factors, including those that are age-dependent, that

trigger the transition of pneumococci from benign colonizer to pathogen are poorly defined,

in part because small animal models that recapitulate the transition from asymptomatic

colonization to overt clinical illness are lacking.

Insight into events that occur in both the nasopharynx and lung and contribute to the

heightened susceptibility of the aged to serious disease upon pneumococcal/IAV

coinfection is needed to develop better therapeutic and preventative approaches. A current

murine model for the spread of nasopharyngeal pneumococci to the lung after viral

infection of the upper respiratory tract relies on initial bacterial colonization of the

nasopharynx followed by viral infection, but does not recapitulate the severe signs of

human clinical disease (4). To better investigate the transition of *S. pneumoniae* from

asymptomatic colonizers to invasive pathogens following IAV infection, as well as the

effect of host age on the disease process, we built upon this mouse model of *S.*

*pneumoniae/*IAV co-infection. The enhanced model incorporates the sequential

introduction of bacteria and virus and recapitulates the severe and age-exacerbated

clinical disease observed in humans.

RESULTS

Intranasal IAV inoculation of mice pre-colonized with *S. pneumoniae* strain TIGR4

does not result in disease.

Biofilm-grown *S. pneumoniae* are relatively less virulent and thus adapted to host

colonization rather than disease (2-4). A previously established murine model in

BALB/cByJ mice utilizes biofilm-grown *S. pneumoniae* strains D39 and 19F EF3030 to

establish heavy carriage in the nasopharynx (4). Then, two days after bacterial inoculation,

IAV is introduced into the nasal cavity and results in the spread of pneumococci from the

nasopharynx (NP) to the lung (4). We first recapitulated this model with *S. pneumoniae*

strain TIGR4, an invasive serotype 4 strain (54, 55) that we previously found to be highly

virulent in aged C57BL/6 mice (19). We intranasally (i.n.) inoculated young (8-10 weeks)

C57BL/6 (B6) mice with 1×10^6 colony forming units (CFU) of biofilm-generated *S.*

pneumoniae TIGR4 by delivering the bacteria in 10 µl (a volume that is unlikely to

inoculate the lung (56)) to the nares of non-anesthetized mice. Forty-eight hours later,

mice were i.n. inoculated with 10 µl containing 20 PFU of Influenza A (IAV) virus PR8.

(See Fig. S1A for general scheme). Control groups of mice were either infected with *S.*

pneumoniae or IAV alone as controls. However, under these conditions, mice did not

display signs of sickness, nor did bacteria spread into the lungs after seven days (not

shown).

To increase the likelihood of clinical disease, we repeated the experiment with a

five-fold higher (5×10^6 CFU) dose of bacteria and a 25-fold higher dose (500 PFU) of IAV.

Similar to earlier reports for other *S. pneumoniae* strains (D39 and 19F) (4), IAV co-

infection resulted in a 10-fold increase in *S. pneumoniae* TIGR4 in the nasal lavage fluid at

2 days post-IAV infection (Fig. S1B). In addition, IAV co-infection was associated with the

detection of bacteria in the lungs of 40% of mice, compared to none in the control group

infected with *S. pneumoniae* TIGR4 alone. This trend is consistent with the previous

BALB/cByJ mice model of coinfection (4), but did not reach statistical significance.

Furthermore, co-infection was not associated with weight loss when assessed over the

course of 4 days post-infection (Fig. S1C). We also scored mice for clinical signs of the

disease based on weight loss, activity, posture and breathing and ranging from healthy

[score = 0] to moribund [score = 25] and requiring euthanasia if the score was >9 as

previously described (57). As secondary pneumonia can occur several days following IAV

(41), we monitored the disease course up to 7 days, but did not detect disease symptoms

or death in any of the co-infected mice (100% survival and 0 daily clinical score for all

mice). Therefore, despite promoting bacterial dispersal from the nasopharynx into the

lungs, this model of *S. pneumoniae* TIGR4/IAV coinfection did not result in overt clinical

signs of disease.

Combined intranasal/intratracheal IAV inoculation of *S. pneumoniae*-colonized mice

results in bacterial dissemination and disease

IAV infection is not restricted to the upper respiratory tract, and can cause viral

pneumonia in a significant fraction of infected individuals (58, 59) that is likely to be crucial

for creating an environment in the lungs that is more permissive for bacterial infection (29,

34, 41). Indeed, viral lung infection diminishes pulmonary defenses against *S. pneumoniae*

and promotes secondary bacterial pneumonia (34, 41, 44-50). Delivery of IAV i.n. to

BALB/cByJ mice results in signs of viral pneumonia (4), but we found that pulmonary

access of inocula delivered via the nasopharynx is more restricted in B6 mice as compared

to BALB/c mice (unpublished observation), raising the possibility that the lack of disease

observed in co-infected B6 mice was due to the exclusive localization of virus in the nasal

cavity, with limited opportunity to alter systemic or pulmonary immunity.

To ensure delivery of IAV to both the nasopharynx and the lungs, we co-infected *S.*

pneumoniae-colonized B6 mice by delivering the virus by two routes. Mice were inoculated

i.n. with 5×10^6 CFU of biofilm grown S. *pneumoniae* TIGR4 and 48 hours later infected not

only with 500 PFU IAV i.n., but also 20 PFU i.t. to ensure pulmonary infection (Fig. 1A). No

mice in a control group inoculated i.n. with *S. pneumoniae* alone lost weight (Fig 1B),

displayed clinical signs of sickness (Fig 1C), or died (Fig 1D). A second control group,

inoculated i.n. and i.t. with IAV alone displayed no disease until after day 4, when they

exhibited weight loss (Fig 1B) and began succumbing to viral infection (Fig 1D). In

contrast, inoculation of IAV to animals pre-colonized with *S. pneumoniae* caused a

bacterial/viral co-infection that resulted in weight loss (Fig. 1B), clinical symptoms (Fig. 1C)

and death (Fig. 1D) that were detected starting day 2 post IAV introduction. At this time

point, a higher fraction of co-infected mice displayed signs of disease as compared to

controls infected with IAV only (40% vs 64%), and disease was more severe in sick mice,

although this did not reach statistical significance. In addition, the overall survival rate

among co-infected mice was significantly lower than mice singly infected with *S.*

pneumoniae alone (Fig. 1D). These signs of exacerbated disease were associated with

significantly higher bacterial burdens in the nasopharynx as well as translocation of *S.*

pneumoniae into the lung in comparison to mice colonized with the bacteria or mice

infected with IAV alone (Fig. 1E). These findings demonstrated that the new co-infection

model leads to disease in a fraction of young healthy mice, which may increase the

likelihood of detecting enhanced susceptibility in vulnerable hosts.

PMN depletion has no significant effect on the course of disease during IAV/*S.*

***pneumoniae* co-infection**

We previously found that in primary pneumococcal pneumonia, PMNs are required

to control bacterial numbers early in the infection process; however their persistence in the

lungs is detrimental to the host and can promote the infection at later time points (20). To

address the role of PMNs during co-infection, we treated young mice with PMN-depleting

anti-Ly6G antibody (1A8) one day prior to pneumococcal colonization and throughout the

co-infection (based on timeline in Figure 1A). We then confirmed that the cells were

depleted by staining with the RB6 antibody followed by flow cytometry (see materials and

methods). Following infection, we measured bacterial burden, weight loss, clinical score

and survival over time. Surprisingly, PMN depletion had no effect on bacterial burdens in

the nasopharynx or bacterial spread to the lungs or blood following co-infection (Fig 2A

and B). PMN-depletion was not associated with any significant differences in weight loss

as compared to the control group (Fig. 2C). Additionally, untreated controls and PMN-

depleted mice had comparable clinical signs of sickness at 18 hours and 48 hours post

IAV infection (Fig. 2D). Lower survival was observed in the depletion group, where only

25% survived to day 7 compared to ~43% survival from untreated controls; however, this

difference was not statistically significant (Fig. 2E). In conclusion, PMN depletion prior to

infection appeared to have a slight but not significant effect on the course of co-infection in

young mice in this model.

Aging increases susceptibility to IAV/*S. pneumoniae* co-infection.

We next tested if the new co-infection mouse model (Fig. 1A) recapitulates the age-

associated increase in susceptibility to secondary pneumococcal pneumonia. Old (20-24

months) B6 mice were inoculated i.n. with $5x10^6$ CFU of biofilm grown S. *pneumoniae*

TIGR4 and 48 hours later were infected with 500 PFU IAV i.n. plus 20 PFU i.t. When

compared to young co-infected controls, old mice displayed significantly more severe signs

of disease, as indicated by a higher average clinical score (Fig. 3A). While seven of 16

(43.75%) young mice showed clinical symptoms (i.e., clinical score greater than 1; Fig.

3A), all 13 old mice showed at least some degree of illness by day 2 post co-infection (p =

0.0012, by Fisher's exact test). Furthermore, whereas only 25% (4 out of 16) young mice

had a clinical score greater than 2, which is indicative of more severe disease, 92% (12 out

of 13) aged mice fell into this category (p = 0.005, by Fisher's exact test). In addition, old

mice died at a significantly accelerated rate. By day 2 post co-infection, 60% of old mice

had succumbed to the infection compared to only 25% of young mice. Differences in

survival were observed at each successive time point, and at the end of the experiment on

day 8, only 14% of old mice remained alive compared to 50% of young mice (Fig. 3B).

Importantly, the accelerated death observed in co-infected old mice was not observed in

old mice infected with *S. pneumoniae* alone or IAV alone (Fig S2).

We next tested whether these differences could be attributed to increased bacterial

loads in the nasopharynx, lungs, or blood. As old mice got sicker at earlier time points post

IAV co-infection, with the majority succumbing by day 4, we compared bacterial burden

across age groups at 18 and 48 hours post IAV co-infection. We found no significant

differences in the numbers of pneumococci in nasopharyngeal washes, pulmonary

homogenates, or blood at either time point (Fig 3C-E). Taken together, these findings

suggest that with aging there is an increased susceptibility to co-infection and an

accelerated course of disease that could not be attributed to a more rapid bacterial

dissemination or higher bacterial loads in the nasopharynx, lung, or bloodstream.

Aging is associated with more rapid lung inflammation.

We previously found that mice suffering from exacerbated PMN-mediated

pulmonary inflammation during pneumococcal pneumonia did not display higher bacterial

burdens in their lungs (27) despite a higher likelihood of severe disease (19, 20, 27). To

test whether PMN influx was higher in old co-infected mice, we measured the percentage

and number of pulmonary PMNs (Ly6G$^+$) by flow cytometry. We found that old mice had

significantly (6-fold) higher percentages and numbers of PMNs in their lungs as compared

to young controls at 18 hours post co-infection (Fig 4A). By 48 hours, although PMN

percentages and numbers appeared to be higher in young mice, the differences were not

statistically significant (Fig. 4A and B). Macrophages, which are important for host

resistance to *S. pneumoniae/IAV* co-infection (49) and display age-driven changes (60,

61), displayed no significant age-dependent differences in either percentage or number

(Fig. S3).

Next we tested the levels of inflammatory cytokines in co-infected young vs old

mice. No significant differences between age groups were detected in baseline

(uninfected) levels of any of the cytokines tested between the age groups (Fig 4B and not

shown). However, consistent with the enhanced PMN influx at 18 h post-infection in aged

mice, old mice had significantly higher levels of IL-10, IL-2, IL-1β and TNFα (Fig 4B)

compared with young mice. Levels of IL-12p70, IL-17, IL-6 and IFNγ, were slightly but not

significantly elevated 18 hours post co-infection (Fig S4A). By 48 hours there were no

significant differences between young and old mice in cytokine levels except for IFNγ,

which was higher in young mice (Fig S4B).

To determine if the pulmonary inflammation was associated with lung damage, we

analyzed H&E stained lung sections for alveolar congestion, hemorrhage, alveolar

thickness, neutrophils and lymphocytic infiltration (Fig. 4C). We found that the alveolar

spaces of both uninfected old and young mice were clear and free of inflammatory or red

blood cells (Fig. 4C). Similarly, at 18h hours post co-infection, the lungs of young mice did

not show any overt signs of disease (Fig. 4C). However, co-infected old mice had

significant pathology in the lungs by 18 hours post infection, including a loss of alveolar

architecture, and spotty inflammation consisting of infiltrates composed of neutrophils,

alveolar macrophages and mononuclear cells that were mixed with red blood cells (Fig.

4C). At 48 hours post-infection, there were clear signs of lung pathology in both young and

old co-infected mice (Fig. 4C). Taken together, these findings demonstrate that aging is

associated with earlier pulmonary inflammation and damage following co-infection, which

may account for the accelerated death observed in this mouse group.

Aging and IAV infection diminish the ability of PMNs to kill *S. pneumoniae ex vivo*.

Aged, co-infected mice experience an accelerated rate of pulmonary inflammation

but bacterial loads in the lungs of aged mice were not lower than in young mice, indicating

that PMN infiltration is not associated with bacterial clearance. Both aging (24) and IAV

infection (62) have been reported to diminish antibacterial function of PMNs. To assess the

ability of PMNs to kill *S. pneumoniae* in our co-infection model, we used a well-established

opsonophagocytic (OPH) killing assay (19, 63). We first compared the bactericidal activity

of bone marrow-derived PMNs from young or aged mice. The percentage of bacteria killed

upon incubation with PMNs was determined by comparing surviving CFU to no PMN

control reactions. We found that as previously reported for humans (23) and mice (24) the

ability of PMNs isolated from uninfected old mice to kill pneumococci was reduced 5-fold

compared to young controls (Fig. 5).

We next examined the bactericidal activity of PMNs isolated from young or old mice

2-days after i.t./i.n. IAV inoculation. As previously reported (47-50), PMNs from young IAV

infected mice had a slight (2-fold) but significant reduction in their ability to kill

pneumococci as compared to PMNs from uninfected controls (Fig. 5). Strikingly, IAV

infection abolished the ability of PMNs from old mice to kill *S. pneumoniae*; instead, PMNs

from IAV infected old mice promoted bacterial growth instead of death (Fig. 5). These

findings suggest that IAV infection completely abrogates the ability of PMNs from old mice

to kill *S. pneumoniae.*

DISCUSSION

S. pneumoniae remains a leading cause of secondary bacterial pneumonia

following influenza A virus infection and is associated with severe disease (30, 34-36),

particularly in the elderly (52). The majority of *S. pneumoniae/*IAV co-infection studies have

delivered bacteria into the lungs of mice pre-infected with IAV to reveal changes in the host

lungs and immune response that are crucial for priming invasive pneumococcal disease

(34, 41, 44-50). In this study, to investigate the transition of *S. pneumoniae* from colonizer

to pathogen upon IAV co-infection, a process that has just started to be elucidated (4, 37),

we have developed a modified murine infection model that recapitulates this transition and

results in severe clinical disease. In a previously established model, female BALB/cByJ

mice were first colonized intra-nasally with biofilm-grown pneumococci and then infected

with IAV by delivering the virus to the nasopharynx (4, 40). This model showed that

changes in the host environment in response to viral infection triggers the dispersal of

pneumococci from colonizing biofilms and their spread to the lower respiratory tract (4).

Importantly, the dispersed bacteria expressed higher levels of virulence factors required for

infection, thus, rendering them pathogenic (4). When we used this model to co-infect male

C57BL/6 (B6) mice, we observed a significant increase in dispersed bacteria in the

nasopharynx, but no disease and only a transient presence of *S. pneumoniae* in the lungs

of B6 mice. This could be due to the differences in host susceptibility to bacterial infection

between the different mouse strains, which has been well described for *S. pneumoniae*

(64, 65) and other infections (66) or due to gender (67): Here we performed experiments in

male instead of female mice due to both the easier availability of aged male animals and

the documented higher rate of pneumococcal pneumonia in men compared to women (68,

69).

Previous work of pneumococcal inoculation of IAV-infected lungs showed viral

infection to be crucial for creating an environment in the lungs that is more permissive for

bacterial infection (41, 44-50). The relatively low bacterial burden in co-infected male B6

mice observed here upon IAV inoculation only by the i.n. route suggested that the lung

environment was not sustaining the bacteria. Therefore, we co-infected *S. pneumoniae*

colonized B6 mice with IAV by delivering the virus not only i.n. to infect the nasopharynx,

but also i.t. to ensure infection of the lungs. This modified model recapitulated the increase

in non-adherent pneumococci in the nasopharynx observed upon viral co-infection (4, 70,

71), and resulted in bacterial spread into the lungs and circulation that increased over time.

Importantly, this mode of dual infection recapitulated both the increased colonization

burden (72, 73) as well as the clinical signs of severe disease observed in humans (30, 34-

36), and resulted in death of approximately half of co-infected young controls.

Although the elderly are at higher risk for secondary pneumococcal pneumonia

following IAV infection, animal studies exploring this age-driven susceptibility to co-

infection are few (51, 74, 75). Here, using the modified model, we found that old mice were

significantly more susceptible to *S. pneumoniae*/IAV co-infection. Old mice displayed more

severe signs of disease as compared to young controls and the majority (>85%) failed to

survive the co-infection. This increased susceptibility in old mice was not linked to higher

bacterial dissemination from the nasopharynx, or greater establishment of infection in the

lungs or systemic spread into the circulation at the time points tested. Although we did not

directly measure viral loads, susceptibility to viral infection alone as measured by weight

loss, within the first 5 days following IAV was also similar between the age groups. Rather,

the age-driven susceptibility to co-infection was associated with earlier and more severe

pulmonary inflammation. Production of cytokines in the lungs has been reported in young

mice infected with *S. pneumoniae* 7 days post IAV (76). Further, old mice displayed

changes in the expression of pattern recognition receptors in the lungs leading to altered

inflammatory responses (51). Similar to previous studies (51), we found here that co-

infected old mice had higher levels of TNFα as compared to young controls. However, in

contrast to previous reports that found reduced NLRP3 inflammasome expression in the

lungs and lower production of IL-1β, we found higher levels of IL-1β in old mice as

compared to young controls. This may be accounted for by differences in expression of

bacterial factors. The expression of pneumolysin, which was found to activate NLRP3

inflammasomes and lead to production of IL-1β (77), was elevated in pneumococci

dispersed from biofilms upon IAV infection (78) and therefore may have primed the IL-1β

production we observed in old mice.

We previously found that PMNs were key determinants of disease during primary

pneumococcal pneumonia and are required to initially control bacterial numbers (20, 79).

Therefore, we explored here the role of PMNs in *S. pneumoniae/*IAV co-infection. Similar

to other studies, we found that PMNs are recruited to the lungs of co-infected young mice

(49). Previous reports indicate that the PMN-mediated anti-pneumococcal function in IAV-

infected mice (62) and humans (73) is progressively reduced over time. For example, in

mice, PMNs demonstrably contribute to host defense at 3 days but not at 6 days post-

infection (62). Nevertheless, PMN depletion showed that these cells are important for

control of bacterial transmission (80) and control of pulmonary bacterial numbers (49) in

IAV co-infected mice. In contrast, here we found that PMN depletion starting prior to

bacterial colonization and continuing throughout viral co-infection had no significant effect

on the number of dispersed *S. pneumoniae* in the nasopharynx or their spread to the lungs

and blood. In this model IAV may rapidly impair PMN function, and in fact, we found that

within 2 days following IAV infection, the ability of bone marrow-derived PMNs to kill *S.*

pneumoniae was significantly blunted in young mice. Alternatively, the apparent inability of

PMNs to limit bacterial numbers in this model could be due to the enhanced virulence of

pneumococci dispersed from the nasopharyngeal environment (4) compared to broth-

grown bacteria typically used in other models (62).

In this study we found that aging was associated with earlier influx of PMNs into the

lungs of co-infected mice. We previously demonstrated that excessive PMN influx into the

lungs is detrimental for the ability of old mice to control invasive disease following primary

pneumococcal pneumonia (19) and that PMN depletion 18 hours after infection boosted

host survival (20). Similarly, exacerbated PMN influx into the lungs of old mice singly

infected with IAV was associated with host mortality, and depletion of these cells six days

following viral infection significantly boosted host survival (81). Uncontrolled PMN influx

can result in tissue damage, disruption of gaseous exchange and pulmonary failure. In

fact, it was reported that in IAV singly infected old mice, PMNs mediated lung inflammation

and damage and their depletion reduced the levels of inflammatory IL-1β and TNFα (81).

Therefore, the early increase in these inflammatory cytokines and lung damage we

observed here in *S. pneumoniae*/IAV co-infected old mice may be driven by the elevated

levels of pulmonary PMNs.

In this model, aging was associated with significantly increased levels of pulmonary

IL-10 during *S. pneumoniae*/IAV confection as compared to young controls, which may

account for their enhanced susceptibility to co-infection. IL-10 was previously shown to

impair host resistance to *S. pneumoniae* following IAV infection (82). IL-10 levels were

elevated in co-infected young mice as compared to those singly infected with *S.*

pneumoniae and blocking this cytokine resulted in reduced bacterial numbers and

extended host survival (82). The elevated IL-10 may also account for the complete

inability of PMNs from IAV infected old mice to kill *S. pneumoniae.* We previously found

that IL-10 impairs PMN anti-bacterial function by inhibiting ROS production in response to

infection (79). Blocking this cytokine boosted PMN anti-bacterial function and importantly

restored resistance of vulnerable hosts to primary pneumococcal pneumonia (79).

In summary, here we modified existing murine models to establish an experimental

system that reflects the transition of *S. pneumoniae* from asymptomatic colonizer to

invasive pulmonary pathogen upon IAV co-infection. In this model, IAV triggers the

transition of a pathobiont from a commensal to a pathogenic state through modification of

bacterial behavior in the nasopharynx, enhancement of bacterial colonization in the lung,

and compromise of PMN-mediated anti-bacterial immunity. Importantly this model

recapitulates the susceptibility of aging to co-infections. Moving forward, this model can be

used to dissect the multiple phases of pneumococcal disease progression from

commensals to pathogens, which can help inform specialized treatment options (70)

tailored to the susceptible elderly population.

MATERIALS AND METHODS

Mice. Young (8-10 weeks) and aged (18-24 months) male C57BL/6 mice were purchased

from Jackson Laboratories (Bar Harbor, ME) and the National Institute of Aging. Mice were

housed in a pathogen-free facility at Tufts University. All procedures were performed in

accordance with Institutional Animal Care and Use Committee guidelines.

Bacterial biofilms. NCI-H292 mucoepidermoid carcinoma cells (H292) were grown in 24-well plates in RPMI 1640 media with 10% FBS and 2mM L-glutamine until confluent. Cells were washed with 1x PBS and fixed in 4% paraformaldehyde for 1 hour on ice. *S. pneumoniae* TIGR4 (kind gift from Andrew Camilli) were grown on Tryptic Soy Agar plates supplemented with 5% sheep blood agar (blood agar plates) overnight, then diluted and grown in chemically defined liquid medium (CDM) (4, 40) supplemented with Oxyrase until OD_{600nm} of 0.2. Bacteria were diluted 1:1000 in CDM and seeded on the fixed H292 cells. The, bacteria/H292 cells were incubated at 34°C/ 5% CO_2 and media was changed every 12 hours. At 48 hours post-infection, the supernatant containing planktonic bacteria (non-adherent to NCI-H292 cells) cells was discarded, the cells gently washed with PBS and adherent biofilms collected in fresh CDM by vigorous pipetting. Biofilm aliquots were then frozen at -80°C in the CDM with 25% (v/v) glycerol.

Intranasal inoculation. Before use, biofilm aliquots were thawed on ice, washed once and diluted in PBS to the required concentration. The mice were restrained without anesthesia and infected i.n. with 10μl (5×10^6 CFU) of biofilm grown *S. pneumoniae*. The inoculum was equally distributed between the nostrils with a pipette. Bacterial titers were confirmed by serial dilution and plating on blood agar plates. To ensure stable colonization of the biofilm in the nasopharynx, groups of mice were euthanized at 18 and 48 hours post inoculation, and the nasal washes and tissue were collected and plated on blood agar plate for enumeration of *S. pneumoniae.*

Viral infection. The mouse-adapted H1N1 Influenza A virus PR8 (A/PR/8/34) was

obtained from Dr. Bruce Davidson (4, 40) and stored at -80℃. Before use, viral aliquots

were thawed on ice, diluted in PBS and used to inoculate mice. At 48 hours following

bacterial inoculation, mice were infected i.n. with 10µl of 500 plaque forming units (PFU) of

virus by pipetting the inoculum into the nostrils of non-anesthetized mice. Following i.n.

inoculation, mice were lightly anesthetized with isoflurane and challenged i.t. with 20 PFU

of virus in a 50µl volume pipetted directly into the trachea with the tongue pulled out to

facilitate delivery (20).

Clinical scoring and bacterial burden. Following co-infection, mice were monitored daily

and blindly scored for signs of sickness including weight loss, activity, posture and

breathing. Based on these criteria, the mice were given a clinical score of healthy [0] to

moribund [25] as previously described (57). Any mice displaying a score above 9 are

humanely euthanized in accordance with our protocol. Mice were euthanized at indicated

time points and the lung, nasal lavages and blood collected and plated on blood agar for

enumeration of bacterial loads, as previously described (20). For collection of sera, blood

was collected via cardiac puncture into Microtainer® tubes (BD Biosciences) and

centrifuged at 7607xg for 2 minutes to collect serum as per manufacturer's instructions.

Depletion of PMNs. Mice were intraperitoneally (i.p.) injected with 100µl (50µg/mouse) of

anti-Ly6G clone 1A8 (BD Biosciences) to deplete neutrophils. Mice were injected daily with

the depleting antibodies starting one day prior and ending two days after bacterial

inoculation, followed by every other day from day 1 post viral co-infection to the end of

each experiment. Treatment resulted in >90% neutrophil depletion as described below.

Cell isolation and Flow Cytometry. Mice lungs were harvested, washed in PBS, and

minced into small pieces. The sample was then digested for 45 minutes with RPMI 1640

1X supplemented with 10% FBS, 1 mg/ml Type II collagenase (Worthington, Lakewood,

NJ), and 50 U/ml deoxyribonuclease I to obtain a single-cell suspension as previously

described (20). The red blood cells were lysed using ACK lysis buffer (Gibco). Cells were

then stained with anti-mouse Ly6G clone 1A8 (BD Biosciences), F480 clone BM8

(BioLegend), CD11c clone N418 (eBioscience), and CD11b clone M1/70 (eBioscience).

For neutrophil depletion, cells isolated from the lungs at 18 and 48 h post co-infection were

also stained with either Ly6G clone 1A8 or RB6 clone RB6-8C5 (BioLegend) antibodies to

confirm cell depletion. The fluorescence intensities were measured on BD LSR II Flow

Cytometer at Tufts FACS Core Facility (Boston, MA) to capture at least 25,000 cells and

analyzed using FlowJo.

Isolation of PMNs and Opsonophagocytic Killing Assay (OPH). Femurs and tibias of

uninfected mice were collected and flushed with RPMI, supplemented with 10% FBS and 2

mM EDTA to obtain bone marrow cells. Neutrophils were isolated by density gradient

centrifugation, using Histopaque 1119 and Histopaque 1077 as previously described (83).

The neutrophils were resuspended in Hank's (Gibco) buffer/0.1% gelatin with no Ca+ or

Mg+ and tested for purity by flow cytometry using anti-Ly6G antibodies (eBioscience)

where 85-90% of enriched cells were Ly6G+. The ability of neutrophils to kill bacteria was

measured using a well-established opsonophagocytic (OPH) killing assay as previously

described (20). Briefly, 2.5×10^5 neutrophils were incubated in Hank's buffer/0.1% gelatin

with 10^3 CFU of *S. pneumoniae* pre-opsonized with 3% mouse sera. The reactions were

incubated in flat bottom 96-well non-binding plates for 45 minutes at 37°C. Each group

was plated on blood agar to enumerate viable CFU. Percent bacterial killing was

calculated in comparison to a no PMN control under the same conditions.

Histology. Whole lungs were harvested from groups of mice at 18 hours and at 48 hours

post co-infection and fixed in 10% neutral buffered formalin for 2 days. The tissues were

then embedded in paraffin, sectioned at 5 μm and stained with Hematoxylin and Eosin at

the Animal Histology Core at Tufts University. Sections of lung from three mice per group

were imaged using a Nikon Eclipse E400 microscope. Photomicrographs were captured

using a SPOT Idea 5.0-megapixel color digital camera and SPOT software.

Histopathologic scoring was performed by a board-certified anatomic pathologist

experienced in murine pathology, from 0 (no damage) to 4+ (maximal damage) for alveolar

congestion, hemorrhage, alveolar thickness, neutrophils, and lymphocytic infiltration (84).

Cytokine Analysis. Cytokines in the lungs and serum samples were measured using

Mouse Cytokine 8-Plex Array (Quanterix, Billerica, MA) following the manufacturer's

instructions. Frozen lung homogenates and serum samples were thawed on ice and mixed

by gentle vortexing. Levels in lung supernatants and serum samples were measured using

the Cirascan at the Imager at Tufts University Genomic Core (Boston, MA) and analyzed

by the Cirascan/Cirasoft program. Qlucore Omic Explorer (version 3.5) was used for the

generation of lung cytokine box plots. Concentrations of cytokines (IL-10, IL-2, IL-1β,

TNFα, IL-6, IFNγ, IL-17 and IL-12p70) were log-transformed, and displayed as Log$_2$ pg/ml.

Statistical Analysis. Statistical analysis was performed using Graph Pad Prism7. CFU

data were log-transformed to normalize distribution. Data are presented as mean values

+/- SEM. Significant differences ($p < 0.05$) were determined by Student's t-test. Differences

in fractions of mice that got sick were measured using Fisher's exact test. Survival analysis

was performed using the log rank (Mantel-Cox) test. Asterisks indicate significant

differences and p values are noted in the figures.

ACKNOWLEDGEMENTS

We would like to acknowledge James Nicholas Lee, Summer Schmaling, and Ognjen

Sekulovic for technical assistance with clinical score, virus preparation and nasal lavage

respectively. We would also like to thank Andrew Camilli for bacterial strains and Marta

Gaglia for protocols. They, along with Tim van Opijnen, Bharathi Sundaresh and Marcia

Osburne provided important feedback on the manuscript.

FUNDING

Research reported in this publication was supported by the National Institute On Aging of

the National Institutes of Health under Award Number R21 AG064215 to E.B.G. and F31

AI122615-01A1 to S.R.; King Abdullah Scholarship Program (KASP) implemented by the

Ministry of Higher Education (MOHE) under Award Number 7896504 to B.H.J.

REFERENCES

1.   Kadioglu A, Weiser JN, Paton JC, Andrew PW. 2008. The role of Streptococcus
     pneumoniae virulence factors in host respiratory colonization and disease. Nat Rev
     Microbiol 6:288-301.

2.   Chao Y, Marks LR, Pettigrew MM, Hakansson AP. 2014. Streptococcus
     pneumoniae biofilm formation and dispersion during colonization and disease. Front
     Cell Infect Microbiol 4:194.

3.   Marks LR, Parameswaran GI, Hakansson AP. 2012. Pneumococcal interactions
     with epithelial cells are crucial for optimal biofilm formation and colonization in vitro
     and in vivo. Infect Immun 80:2744-60.

4.   Marks LR, Davidson BA, Knight PR, Hakansson AP. 2013. Interkingdom signaling
     induces Streptococcus pneumoniae biofilm dispersion and transition from
     asymptomatic colonization to disease. MBio 4.

5.   Blanchette-Cain K, Hinojosa CA, Akula Suresh Babu R, Lizcano A, Gonzalez-
     Juarbe N, Munoz-Almagro C, Sanchez CJ, Bergman MA, Orihuela CJ. 2013.
     Streptococcus pneumoniae biofilm formation is strain dependent, multifactorial, and
     associated with reduced invasiveness and immunoreactivity during colonization.
     MBio 4:e00745-13.

6.   Boe DM, Boule LA, Kovacs EJ. 2017. Innate immune responses in the ageing lung.
     Clin Exp Immunol 187:16-25.

7.   Chong CP, Street PR. 2008. Pneumonia in the elderly: a review of the
     epidemiology, pathogenesis, microbiology, and clinical features. South Med J
     101:1141-5; quiz 1132, 1179.

8.   Bogaert D, De Groot R, Hermans PW. 2004. Streptococcus pneumoniae

colonisation: the key to pneumococcal disease. Lancet Infect Dis 4:144-54.

9.   Simell B, Auranen K, Kayhty H, Goldblatt D, Dagan R, O'Brien KL, Pneumococcal

Carriage G. 2012. The fundamental link between pneumococcal carriage and

disease. Expert Rev Vaccines 11:841-55.

10.   Orsi A, Ansaldi F, Trucchi C, Rosselli R, Icardi G. 2016. Pneumococcus and the

Elderly in Italy: A Summary of Available Evidence Regarding Carriage, Clinical

Burden of Lower Respiratory Tract Infections and On-Field Effectiveness of PCV13

Vaccination. Int J Mol Sci 17.

11.   Smith EL, Wheeler I, Adler H, Ferreira DM, Sa-Leao R, Abdullahi O, Adetifa I,

Becker-Dreps S, Esposito S, Farida H, Kandasamy R, Mackenzie GA, Nuorti JP,

Nzenze S, Madhi SA, Ortega O, Roca A, Safari D, Schaumburg F, Usuf E, Sanders

EAM, Grant LR, Hammitt LL, O'Brien KL, Gounder P, Bruden DJT, Stanton MC,

Rylance J. 2020. Upper airways colonisation of Streptococcus pneumoniae in adults

aged 60 years and older: A systematic review of prevalence and individual

participant data meta-analysis of risk factors. J Infect doi:10.1016/j.jinf.2020.06.028.

12.   Regev-Yochay G, Raz M, Dagan R, Porat N, Shainberg B, Pinco E, Keller N,

Rubinstein E. 2004. Nasopharyngeal carriage of Streptococcus pneumoniae by

adults and children in community and family settings. Clin Infect Dis 38:632-9.

13.   Flamaing J, Peetermans WE, Vandeven J, Verhaegen J. 2010. Pneumococcal

colonization in older persons in a nonoutbreak setting. J Am Geriatr Soc 58:396-8.

14.   Palmu AA, Kaijalainen T, Saukkoriipi A, Leinonen M, Kilpi TM. 2012.

Nasopharyngeal carriage of Streptococcus pneumoniae and pneumococcal urine

antigen test in healthy elderly subjects. Scand J Infect Dis 44:433-8.

15.   van Deursen AMM, van Houten MA, Webber C, Patton M, Scott D, Patterson S,

618   Jiang Q, Gruber WC, Schmoele-Thoma B, Grobbee DE, Bonten MJM, Sanders

619   EAM. 2018. The Impact of the 13-Valent Pneumococcal Conjugate Vaccine on

620   Pneumococcal Carriage in the Community Acquired Pneumonia Immunization Trial

621   in Adults (CAPiTA) Study. Clin Infect Dis 67:42-49.

622   16.   Esposito S, Mari D, Bergamaschini L, Orenti A, Terranova L, Ruggiero L, Ierardi V,

623   Gambino M, Croce FD, Principi N. 2016. Pneumococcal colonization in older adults.

624   Immun Ageing 13:2.

625   17.   van Deursen AM, van den Bergh MR, Sanders EA, Carriage Pilot Study G. 2016.

626   Carriage of Streptococcus pneumoniae in asymptomatic, community-dwelling

627   elderly in the Netherlands. Vaccine 34:4-6.

628   18.   Krone CL, van de Groep K, Trzcinski K, Sanders EA, Bogaert D. 2014.

629   Immunosenescence and pneumococcal disease: an imbalance in host-pathogen

630   interactions. Lancet Respir Med 2:141-53.

631   19.   Bou Ghanem EN, Clark S, Du X, Wu D, Camilli A, Leong JM, Meydani SN. 2015.

632   The alpha-tocopherol form of vitamin E reverses age-associated susceptibility to

633   streptococcus pneumoniae lung infection by modulating pulmonary neutrophil

634   recruitment. J Immunol 194:1090-9.

635   20.   Bou Ghanem EN, Clark S, Roggensack SE, McIver SR, Alcaide P, Haydon PG,

636   Leong JM. 2015. Extracellular Adenosine Protects against Streptococcus

637   pneumoniae Lung Infection by Regulating Pulmonary Neutrophil Recruitment. PLoS

638   Pathog 11:e1005126.

639   21.   Garvy BA, Harmsen AG. 1996. The importance of neutrophils in resistance to

640   pneumococcal pneumonia in adult and neonatal mice. Inflammation 20:499-512.

641   22.   Hahn I, Klaus A, Janze AK, Steinwede K, Ding N, Bohling J, Brumshagen C,

642   Serrano H, Gauthier F, Paton JC, Welte T, Maus UA. 2011. Cathepsin G and

643   neutrophil elastase play critical and nonredundant roles in lung-protective immunity

644   against Streptococcus pneumoniae in mice. Infect Immun 79:4893-901.

645   23.   Simell B, Vuorela A, Ekstrom N, Palmu A, Reunanen A, Meri S, Kayhty H,

646   Vakevainen M. 2011. Aging reduces the functionality of anti-pneumococcal

647   antibodies and the killing of Streptococcus pneumoniae by neutrophil phagocytosis.

648   Vaccine 29:1929-34.

649   24.   Bhalla M SS, Abamonte A, Herring SE, Roggensack SE, Bou Ghanem EN. 2020.

650   Extracellular adenosine signaling reverses the age-driven decline in the ability of

651   neutrophils to kill S. pneumoniae. doi:https://doi.org/10.1101/2020.04.14.041418.

652   25.   Pignatti P, Ragnoli B, Radaeli A, Moscato G, Malerba M. 2011. Age-related

653   increase of airway neutrophils in older healthy nonsmoking subjects. Rejuvenation

654   Res 14:365-70.

655   26.   Menter T, Giefing-Kroell C, Grubeck-Loebenstein B, Tzankov A. 2014.

656   Characterization of the inflammatory infiltrate in Streptococcus pneumoniae

657   pneumonia in young and elderly patients. Pathobiology 81:160-7.

658   27.   Bhowmick R, Tin Maung NH, Hurley BP, Ghanem EB, Gronert K, McCormick BA,

659   Leong JM. 2013. Systemic disease during Streptococcus pneumoniae acute lung

660   infection requires 12-lipoxygenase-dependent inflammation. J Immunol 191:5115-

661   23.

662   28.   Rudra Bhowmick NM, Bryan Hurley, Karsten Gronert, Beth McCormick and John

663   Leong. 2012. Systemic disease during Streprococcus pneumoniae lung infection is

664   facilitated by hepoxilin A3 mediated nutophil recruitment, Submitted, In revision.

29.   Bakaletz LO. 2017. Viral-bacterial co-infections in the respiratory tract. Curr Opin

Microbiol 35:30-35.

30.   McCullers JA. 2006. Insights into the interaction between influenza virus and

pneumococcus. Clin Microbiol Rev 19:571-82.

31.   Chertow DS, Memoli MJ. 2013. Bacterial coinfection in influenza: a grand rounds

review. JAMA 309:275-82.

32.   Centers for Disease C, Prevention. 2010. Estimates of deaths associated with

seasonal influenza --- United States, 1976-2007. MMWR Morb Mortal Wkly Rep

59:1057-62.

33.   Shrestha S, Foxman B, Weinberger DM, Steiner C, Viboud C, Rohani P. 2013.

Identifying the interaction between influenza and pneumococcal pneumonia using

incidence data. Sci Transl Med 5:191ra84.

34.   McCullers JA. 2014. The co-pathogenesis of influenza viruses with bacteria in the

lung. Nat Rev Microbiol 12:252-62.

35.   Palacios G, Hornig M, Cisterna D, Savji N, Bussetti AV, Kapoor V, Hui J, Tokarz R,

Briese T, Baumeister E, Lipkin WI. 2009. Streptococcus pneumoniae coinfection is

correlated with the severity of H1N1 pandemic influenza. PLoS One 4:e8540.

36.   Dhanoa A, Fang NC, Hassan SS, Kaniappan P, Rajasekaram G. 2011.

Epidemiology and clinical characteristics of hospitalized patients with pandemic

influenza A (H1N1) 2009 infections: the effects of bacterial coinfection. Virol J

8:501.

37.   Siegel SJ, Roche AM, Weiser JN. 2014. Influenza promotes pneumococcal growth

during coinfection by providing host sialylated substrates as a nutrient source. Cell

Host Microbe 16:55-67.

38.   Rowe HM, Meliopoulos VA, Iverson A, Bomme P, Schultz-Cherry S, Rosch JW.

2019. Direct interactions with influenza promote bacterial adherence during

respiratory infections. Nat Microbiol 4:1328-1336.

39.   Diavatopoulos DA, Short KR, Price JT, Wilksch JJ, Brown LE, Briles DE, Strugnell

RA, Wijburg OL. 2010. Influenza A virus facilitates Streptococcus pneumoniae

transmission and disease. FASEB J 24:1789-98.

40.   Reddinger RM, Luke-Marshall NR, Sauberan SL, Hakansson AP, Campagnari AA.

2018. Streptococcus pneumoniae Modulates Staphylococcus aureus Biofilm

Dispersion and the Transition from Colonization to Invasive Disease. MBio 9.

41.   McCullers JA, Rehg JE. 2002. Lethal synergism between influenza virus and

Streptococcus pneumoniae: characterization of a mouse model and the role of

platelet-activating factor receptor. J Infect Dis 186:341-50.

42.   Levandowski RA, Gerrity TR, Garrard CS. 1985. Modifications of lung clearance

mechanisms by acute influenza A infection. J Lab Clin Med 106:428-32.

43.   Pittet LA, Hall-Stoodley L, Rutkowski MR, Harmsen AG. 2010. Influenza virus

infection decreases tracheal mucociliary velocity and clearance of Streptococcus

pneumoniae. Am J Respir Cell Mol Biol 42:450-60.

44.   McCullers JA, Bartmess KC. 2003. Role of neuraminidase in lethal synergism

between influenza virus and Streptococcus pneumoniae. J Infect Dis 187:1000-9.

45.   Smith AM, McCullers JA. 2014. Secondary bacterial infections in influenza virus

infection pathogenesis. Curr Top Microbiol Immunol 385:327-56.

46.   Cundell DR, Gerard NP, Gerard C, Idanpaan-Heikkila I, Tuomanen EI. 1995.

Streptococcus pneumoniae anchor to activated human cells by the receptor for

platelet-activating factor. Nature 377:435-8.

47.   Ballinger MN, Standiford TJ. 2010. Postinfluenza bacterial pneumonia: host

defenses gone awry. J Interferon Cytokine Res 30:643-52.

48.   Metzger DW, Sun K. 2013. Immune dysfunction and bacterial coinfections following

influenza. J Immunol 191:2047-52.

49.   Sun K, Metzger DW. 2008. Inhibition of pulmonary antibacterial defense by

interferon-gamma during recovery from influenza infection. Nat Med 14:558-64.

50.   Nakamura S, Davis KM, Weiser JN. 2011. Synergistic stimulation of type I

interferons during influenza virus coinfection promotes Streptococcus pneumoniae

colonization in mice. J Clin Invest 121:3657-65.

51.   Cho SJ, Plataki M, Mitzel D, Lowry G, Rooney K, Stout-Delgado H. 2018.

Decreased NLRP3 inflammasome expression in aged lung may contribute to

increased susceptibility to secondary Streptococcus pneumoniae infection. Exp

Gerontol 105:40-46.

52.   https://www.cdc.gov/flu/about/burden/index.html 2018. Disease Burden of

Influenza. CENTERS FOR DISEASE CONTROL AND PREVENTION.

53.   Krone CL, Wyllie AL, van Beek J, Rots NY, Oja AE, Chu ML, Bruin JP, Bogaert D,

Sanders EA, Trzcinski K. 2015. Carriage of Streptococcus pneumoniae in aged

adults with influenza-like-illness. PLoS One 10:e0119875.

54.   Aaberge IS, Eng J, Lermark G, Lovik M. 1995. Virulence of Streptococcus

pneumoniae in mice: a standardized method for preparation and frozen storage of

the experimental bacterial inoculum. Microb Pathog 18:141-52.

55.   Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, Heidelberg J,

DeBoy RT, Haft DH, Dodson RJ, Durkin AS, Gwinn M, Kolonay JF, Nelson WC,

Peterson JD, Umayam LA, White O, Salzberg SL, Lewis MR, Radune D, Holtzapple

737  E, Khouri H, Wolf AM, Utterback TR, Hansen CL, McDonald LA, Feldblyum TV,

738  Angiuoli S, Dickinson T, Hickey EK, Holt IE, Loftus BJ, Yang F, Smith HO, Venter

739  JC, Dougherty BA, Morrison DA, Hollingshead SK, Fraser CM. 2001. Complete

740  genome sequence of a virulent isolate of Streptococcus pneumoniae. Science

741  293:498-506.

742 56. Bou Ghanem EN, Maung NHT, Siwapornchai N, Goodwin AE, Clark S, Munoz-Elias

743  EJ, Camilli A, Gerstein RM, Leong JM. 2018. Nasopharyngeal Exposure to

744  Streptococcus pneumoniae Induces Extended Age-Dependent Protection against

745  Pulmonary Infection Mediated by Antibodies and CD138(+) Cells. J Immunol

746  200:3739-3751.

747 57. Bhalla M, Hui Yeoh J, Lamneck C, Herring SE, Tchalla EYI, Heinzinger LR, Leong

748  JM, Bou Ghanem EN. 2020. A1 adenosine receptor signaling reduces

749  Streptococcus pneumoniae adherence to pulmonary epithelial cells by targeting

750  expression of platelet-activating factor receptor. Cell Microbiol 22:e13141.

751 58. Rello J, Pop-Vicas A. 2009. Clinical review: primary influenza viral pneumonia. Crit

752  Care 13:235.

753 59. Torres A, Loeches IM, Sligl W, Lee N. 2020. Severe flu management: a point of

754  view. Intensive Care Med 46:153-162.

755 60. Boyd AR, Shivshankar P, Jiang S, Berton MT, Orihuela CJ. 2012. Age-related

756  defects in TLR2 signaling diminish the cytokine response by alveolar macrophages

757  during murine pneumococcal pneumonia. Exp Gerontol 47:507-18.

758 61. Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, Loukov

759  D, Schenck LP, Jury J, Foley KP, Schertzer JD, Larche MJ, Davidson DJ, Verdu

760  EF, Surette MG, Bowdish DME. 2018. Age-Associated Microbial Dysbiosis

Promotes Intestinal Permeability, Systemic Inflammation, and Macrophage

Dysfunction. Cell Host Microbe 23:570.

62. McNamee LA, Harmsen AG. 2006. Both influenza-induced neutrophil dysfunction

and neutrophil-independent mechanisms contribute to increased susceptibility to a

secondary Streptococcus pneumoniae infection. Infect Immun 74:6707-21.

63. Bou Ghanem EN, Lee JN, Joma BH, Meydani SN, Leong JM, Panda A. 2017. The

Alpha-Tocopherol Form of Vitamin E Boosts Elastase Activity of Human PMNs and

Their Ability to Kill Streptococcus pneumoniae. Front Cell Infect Microbiol 7:161.

64. Gingles NA, Alexander JE, Kadioglu A, Andrew PW, Kerr A, Mitchell TJ, Hopes E,

Denny P, Brown S, Jones HB, Little S, Booth GC, McPheat WL. 2001. Role of

genetic resistance in invasive pneumococcal infection: identification and study of

susceptibility and resistance in inbred mouse strains. Infect Immun 69:426-34.

65. Kadioglu A, Andrew PW. 2005. Susceptibility and resistance to pneumococcal

disease in mice. Brief Funct Genomic Proteomic 4:241-7.

66. Bou Ghanem EN, Jones GS, Myers-Morales T, Patil PD, Hidayatullah AN, D'Orazio

SE. 2012. InlA promotes dissemination of Listeria monocytogenes to the mesenteric

lymph nodes during food borne infection of mice. PLoS Pathog 8:e1003015.

67. Kadioglu A, Cuppone AM, Trappetti C, List T, Spreafico A, Pozzi G, Andrew PW,

Oggioni MR. 2011. Sex-based differences in susceptibility to respiratory and

systemic pneumococcal disease in mice. J Infect Dis 204:1971-9.

68. Wagenvoort GH, Sanders EA, Vlaminckx BJ, de Melker HE, van der Ende A, Knol

MJ. 2017. Sex differences in invasive pneumococcal disease and the impact of

pneumococcal conjugate vaccination in the Netherlands, 2004 to 2015. Euro

Surveill 22.

69.   Gutierrez F, Masia M, Mirete C, Soldan B, Rodriguez JC, Padilla S, Hernandez I,

Royo G, Martin-Hidalgo A. 2006. The influence of age and gender on the

population-based incidence of community-acquired pneumonia caused by different

microbial pathogens. J Infect 53:166-74.

70.   Greene CJ, Marks LR, Hu JC, Reddinger R, Mandell L, Roche-Hakansson H, King-

Lyons ND, Connell TD, Hakansson AP. 2016. Novel Strategy To Protect against

Influenza Virus-Induced Pneumococcal Disease without Interfering with Commensal

Colonization. Infect Immun 84:1693-1703.

71.   Mina MJ, Klugman KP, McCullers JA. 2013. Live attenuated influenza vaccine, but

not pneumococcal conjugate vaccine, protects against increased density and

duration of pneumococcal carriage after influenza infection in pneumococcal

colonized mice. J Infect Dis 208:1281-5.

72.   Wadowsky RM, Mietzner SM, Skoner DP, Doyle WJ, Fireman P. 1995. Effect of

experimental influenza A virus infection on isolation of Streptococcus pneumoniae

and other aerobic bacteria from the oropharynges of allergic and nonallergic adult

subjects. Infect Immun 63:1153-7.

73.   Jochems SP, Marcon F, Carniel BF, Holloway M, Mitsi E, Smith E, Gritzfeld JF,

Solorzano C, Reine J, Pojar S, Nikolaou E, German EL, Hyder-Wright A, Hill H,

Hales C, de Steenhuijsen Piters WAA, Bogaert D, Adler H, Zaidi S, Connor V,

Gordon SB, Rylance J, Nakaya HI, Ferreira DM. 2018. Inflammation induced by

influenza virus impairs human innate immune control of pneumococcus. Nat

Immunol 19:1299-1308.

74. Stout-Delgado HW, Vaughan SE, Shirali AC, Jaramillo RJ, Harrod KS. 2012.

Impaired NLRP3 inflammasome function in elderly mice during influenza infection is

rescued by treatment with nigericin. J Immunol 188:2815-24.

75. Gay R, Han SN, Marko M, Belisle S, Bronson R, Meydani SN. 2004. The effect of

vitamin E on secondary bacterial infection after influenza infection in young and old

mice. Ann N Y Acad Sci 1031:418-21.

76. Smith MW, Schmidt JE, Rehg JE, Orihuela CJ, McCullers JA. 2007. Induction of

pro- and anti-inflammatory molecules in a mouse model of pneumococcal

pneumonia after influenza. Comp Med 57:82-9.

77. McNeela EA, Burke A, Neill DR, Baxter C, Fernandes VE, Ferreira D, Smeaton S,

El-Rachkidy R, McLoughlin RM, Mori A, Moran B, Fitzgerald KA, Tschopp J, Petrilli

V, Andrew PW, Kadioglu A, Lavelle EC. 2010. Pneumolysin activates the NLRP3

inflammasome and promotes proinflammatory cytokines independently of TLR4.

PLoS Pathog 6:e1001191.

78. Pettigrew MM, Marks LR, Kong Y, Gent JF, Roche-Hakansson H, Hakansson AP.

2014. Dynamic changes in the Streptococcus pneumoniae transcriptome during

transition from biofilm formation to invasive disease upon influenza A virus infection.

Infect Immun 82:4607-19.

79. Siwapornchai N, Lee JN, Tchalla EYI, Bhalla M, Yeoh JH, Roggensack SE, Leong

JM, Bou Ghanem EN. 2020. Extracellular adenosine enhances the ability of PMNs

to kill Streptococcus pneumoniae by inhibiting IL-10 production. J Leukoc Biol

doi:10.1002/JLB.4MA0120-115RR.

80.   Short KR, Reading PC, Wang N, Diavatopoulos DA, Wijburg OL. 2012. Increased

nasopharyngeal bacterial titers and local inflammation facilitate transmission of

Streptococcus pneumoniae. MBio 3.

81.   Kulkarni U, Zemans RL, Smith CA, Wood SC, Deng JC, Goldstein DR. 2019.

Excessive neutrophil levels in the lung underlie the age-associated increase in

influenza mortality. Mucosal Immunol 12:545-554.

82.   van der Sluijs KF, van Elden LJ, Nijhuis M, Schuurman R, Pater JM, Florquin S,

Goldman M, Jansen HM, Lutter R, van der Poll T. 2004. IL-10 is an important

mediator of the enhanced susceptibility to pneumococcal pneumonia after influenza

infection. J Immunol 172:7603-9.

83.   Swamydas M, Lionakis MS. 2013. Isolation, purification and labeling of mouse bone

marrow neutrophils for functional studies and adoptive transfer experiments. J Vis

Exp doi:10.3791/50586:e50586.

84.   Nishina K, Mikawa K, Takao Y, Shiga M, Maekawa N, Obara H. 1998. Intravenous

lidocaine attenuates acute lung injury induced by hydrochloric acid aspiration in

rabbits. Anesthesiology 88:1300-9.

FIGURES & LEGENDS

Figure 1. Combined intranasal/intratracheal IAV inoculation of *S. pneumoniae*-

colonized mice results in bacterial dissemination and disease. (A) Timeline of co-

infection; 8-10 weeks old male C57BL/6 (B6) mice were inoculated i.n. with 5×10^6 CFU of

biofilm grown S. *pneumoniae* TIGR4 to establish colonization in the nasopharynx. 48 hours

later, the mice were either mock treated (*Sp*) or received 500 PFU of Influenza A virus PR8

(IAV) i.n. and 20 PFU i.t. (B) Percent weight loss was monitored daily. (C) Blinded clinical

scoring was performed on day 2 and day 4 post IAV infection. The fractions denote the

number of sick mice observed over total number of mice. A score of 0 means no sign of

sickness observed and a score above 1 indicates observable sickness. (D) Survival was

monitored for 8 days post IAV infection with fractions denoting survivors over total number

of mice. (E) The bacterial burden in the nasopharynx and lung were determined at day 2

post IAV infection. Pooled data from three separate experiments, n = 6-12 mice per group

are shown. Statistically significant differences determined by Student's t-test for bacterial

burden and clinical score and the log-rank (Mantel-Cox) test for survival are indicated by

asterisks. #, indicates statistical significance ($p < 0.05$) between *Sp* and *Sp*+IAV groups by

Fisher's exact test.

Figure 2. PMN depletion does not significantly affect the course of disease during

IAV/*S. pneumoniae* co-infection. 8-10 weeks old C57BL/6 mice were intraperitoneally

(i.p.) injected with anti-Ly6G (clone 1A8) antibodies to deplete neutrophils or mock treated.

The antibodies were given daily from day -3 to day 1, and every other day from day 3 to

the end of each experiment (with respect to IAV-infection). (A) At 18 and 48 hours post

IAV infection, bacterial numbers in the nasal wash and lungs were determined. Bacteremia

(B) and weight loss (C) was monitored over time. (D) Mice were blindly scored for

symptoms of diseases at 18 and 48 hours post IAV-infection. The fractions indicate

number of mice with clinical sickness (clinical score above 1) over total number of mice.

(E) Survival was monitored over time where the fractions denote survivors over total

number of mice at 8 days post IAV infection. Data shown represent the means +/- SEM

and are pooled from three separate experiments with n=16 mice per group. Statistically

significant differences were determined by Student's t-test for bacterial burden and clinical

score and the log-rank (Mantel-Cox) and *ns* = not significant

Figure 3. Aging increases susceptibility to IAV/*S. pneumoniae* co-infection. Young

(8-10 weeks) and aged (18-24 months) C57BL/6 male mice were co-infected with *S.*

pneumoniae TIGR4 i.n. and Influenza A virus PR8 i.n. and i.t. (as in Figure 1A). (A) Clinical

score of co-infected young and aged mice at day 2 post IAV-infection is shown; the

number of mice with demonstrable illness over total number of mice is indicated. Data

shown are pooled from four experiments with n=13-16 mice per group. #, indicate

statistical significance by Fisher's exact test. (B) Survival of co-infected young and aged

mice were monitored over time with fractions denoting survivors over total of mice. Data

are pooled from four experiments with n=14 mice per group. Asterisks indicate statistical

significance by the log–rank (Mantel-Cox) test. (C-E) Bacterial burdens in the nasal wash,

lungs, and blood were determined at 18 and 48 hours post IAV inoculation. The mean +/-

SEM pooled from three separate experiments with n=7-10 mice per age group are shown.

LOD denotes the limit of detection.

Figure 4. Aging is associated with more rapid lung inflammation. Young and aged

C57BL/6 male mice were co-infected with S. *pneumoniae* and Influenza A virus PR8. 18

and 48 hours following IAV-infection (see experimental design in Fig. 1A), the lungs were

harvested. (A) The percentages and total number of PMNs (Ly6G+) in the lungs was

measured by flow cytometry. Young mice are represented by open bars and aged mice by

shaded bars. The mean +/- SEM pooled from three separate experiments with n = 6 mice

and n=12 mice per group at 18 and 48 hours respectively are shown. Statistically

significant differences determined by Student's t-test are indicated by asterisks. (B)

Cytokines in the supernatants of lung homogenates of young (n=4) or aged (n=5) mice at

18 hours post co-infection were measured by multiplex ELISA. Asterisks represent

statistical significance by Student's t-test. (C) Lungs were stained with Hematoxylin and

Eosin and shown are representative photographs at 100x and 400X are shown (inset)

Figure 5. Aging and IAV infection diminish the ability of PMNs to kill *S. pneumoniae*

***ex vivo*.** PMNs were isolated from bone marrow of young (8-10 weeks) and aged (18-24

months) C57BL/6 male mice that were mock-infected (uninf.) or singly infected with IAV

(i.n. + i.t.) for 2 days. PMNs were incubated with *S. pneumoniae* pre-opsonized with

homologous sera from the same mouse for 45 minutes at 37°C. The percentages of *S.*

pneumoniae killed upon incubation with PMNs were determined with respect to a no PMN

control. Data shown represent the means +/- SEM pooled from two experiments (n=3 mice

per group) where each condition was tested in quadruplicates per experiment. Asterisks

represent statistical significance as determined by Student's t-test.