1

A Murine Model for Enhancement of *Streptococcus pneumoniae* Pathogenicity Upon Viral Infection and Advanced Age

Basma H. Joma[*1, 2], Nalat Siwapornchai[*1,] Vijay K. Vanguri[3], Anishma Shrestha[1], Sara E. Roggensack[1, 4], Bruce A. Davidson[5], Albert K. Tai[6], Anders P. Hakansson[7], Simin N. Meydani[8], John M. Leong[#1,9] and Elsa N. Bou Ghanem[#10]

* Co-first authors; B.H.J. and N.S. contributed equally to this work; the order of their names was listed alphabetically

Co-corresponding authors

[1] Department of Molecular Biology and Microbiology at Tufts University School of Medicine, Boston, Massachusetts, USA

[2] Graduate Program in Immunology, Tufts Graduate School of Biomedical Sciences, Boston, USA

[3] UMass Memorial Health Care, University of Massachusetts Medical School, Worcester, Massachusetts, USA

[4] Graduate Program in Molecular Microbiology, Tufts Graduate School of Biomedical Sciences, Boston, USA

[5] Department of Anesthesiology, University at Buffalo School of Medicine, Buffalo, New York, USA

[6] Department of Immunology, Tufts University School of Medicine, Boston, Massachusetts, USA

[7] Department of Translational Medicine, Lund University, Malmö, Sweden

[8] Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts, USA

[9] Stuart B. Levy Center for Integrated Management of Antimicrobial Resistance at Tufts (Levy CIMAR)

[10] Department of Microbiology and Immunology, University at Buffalo School of Medicine, Buffalo, New York, USA

Running Title: Murine Model of Influenza-Pneumococcal Co-infection

Address correspondence and reprint requests to John M. Leong (J.M.L.), Department of Molecular Biology and Microbiology, Tufts University School of Medicine, 136 Harrison Avenue, Boston, MA 02114 and Elsa Bou Ghanem (E.N.B.G.), Department of Microbiology and Immunology, University at Buffalo School of Medicine, 955 Main Street, Buffalo, New York, USA. E-mail addresses: john.leong@tufts.edu (J.M.L.) and elsaboug@buffalo.edu (E.N.B.G.)

Keywords: Co-infection, secondary bacterial pneumonia, *Streptococcus pneumoniae*, Influenza A, colonization, aging, neutrophils, inflammation

ABSTRACT

Streptococcus pneumoniae (pneumococcus) resides asymptomatically in the nasopharynx but can progress from benign colonizer to lethal pulmonary or systemic pathogen. Both viral infection and aging are risk factors for serious pneumococcal infections. Previous work established a murine model that featured the movement of pneumococcus from the nasopharynx to the lung upon nasopharyngeal inoculation with influenza A virus (IAV) but did not fully recapitulate the severe disease associated with human co-infection. We built upon this model by first establishing pneumococcal nasopharyngeal colonization, then inoculating both the nasopharynx and lungs with IAV. In young (2 months) mice, co-infection triggered bacterial dispersal from the nasopharynx into the lungs, pulmonary inflammation, disease and mortality in a fraction of mice. In old mice (20-22 months), co-infection resulted in earlier and more severe disease. Aging was not associated with greater bacterial burdens but rather with more rapid pulmonary inflammation and damage. Both aging and IAV infection led to inefficient bacterial killing by neutrophils *ex vivo*. Conversely, aging and pneumococcal colonization also blunted IFN-α production and increased pulmonary IAV burden. Thus, in this multistep model, IAV promotes pneumococcal pathogenicity by modifying bacterial behavior in the nasopharynx, diminishing neutrophil function, and enhancing bacterial growth in the lung, while pneumococci increase IAV burden likely by compromising a key antiviral response. Thus, this model provides a means to elucidate factors, such as age and co-infection, that promote the evolution of *S. pneumoniae* from asymptomatic colonizer to invasive pathogen, as well as to investigate consequences of this transition on antiviral defense.

INTRODUCTION

Streptococcus pneumoniae (pneumococcus) is a Gram-positive pathobiont that typically resides asymptomatically in the nasopharynx of healthy individuals (1). It is hypothesized *S. pneumoniae* establishes an asymptomatic biofilm on the nasopharyngeal epithelium by attenuating the production of virulence factors and concomitant inflammation (2-5). However, when immunity is compromised, a common occurrence in aging (6), pneumococci can cause serious disease such as otitis media, pneumonia, meningitis and bacteremia (7). In humans, pneumococcal carriage is believed to be a prerequisite to invasive disease (8, 9). Bacterial isolates from invasive infections are genetically identical to those found in the nasopharynx of patients (9); this and other longitudinal studies have led to the suggestion that invasive disease often involves pneumococcal carriage in the upper respiratory tract (8, 9).

The rate of reported colonization is quite variable among adults and is confounded by differences in detection methods, but colonization may be more prevalent in the elderly (10). A meta-analysis of twenty nine published studies found that in individuals above 60 years of age, conventional culture showed 0-39% carriage, while 3-23% carriage was detected by molecular methods (11). Importantly, carriage was higher among nursing home residents (11). In older adults, conventional culture methods estimated carriage to be <5% (12-14) but more recent data from European studies using molecular detection methods indicated that carriage in the elderly ranges from 10-22% (10, 15-17). Thus, carriage rates may be much higher than what was previously estimated in elderly individuals (12-14).

Advanced age increases the risk of invasive pneumococcal disease and pneumococcal pneumonia (7). Aging is associated with immunosenescence, the overall

decline in immunity that accompanies aging, as well as inflammaging, a low-grade chronic

inflammation that render the elderly more susceptible to pulmonary infections (18).

Polymorphonuclear leukocytes (PMNs), also known as neutrophils, are a crucial

determinant of age-related susceptibility to primary pneumococcal infection (19). These

cells are required to control bacterial burden at the start of infection (20-22). However,

aging is accompanied by impaired PMN anti-bacterial function (23, 24). In addition, aged

hosts experience exacerbated PMN pulmonary influx during primary pneumococcal

pneumonia (19, 25, 26), and persistence of PMNs in the airways beyond the first few days

leads to tissue destruction and systemic spread of infection (27, 28).

In addition to advanced age, epidemiological and experimental data show that

invasive pneumococcal infections are strongly associated with viral infection (29-31). The

risk of pneumococcal pneumonia is enhanced 100-fold by influenza A virus (IAV) infection

(32, 33), resulting in the seasonal peak of pneumococcal disease during influenza

outbreaks (33). Further, *S. pneumoniae* is historically among the most common etiologies

of secondary bacterial pneumonia following influenza and associated with the most severe

outcomes (30, 34-36). Symptoms of secondary bacterial pneumonia include cough,

dyspnea, fever, muscle aches and when severe result in hospitalizations, respiratory

failure, mechanical ventilation, and can lead to death (30, 34-36).

IAV commonly infects the upper respiratory tract, and at this location viral infection

can enhance the nutritional environment for pneumococcus in the nasopharynx, leading to

greater bacterial loads and/or higher rates of bacterial acquisition (4, 37). IAV can also

directly bind the pneumococcal surface and enhance bacterial binding to the pulmonary

epithelium leading to increased colonization (38). In addition, viral infection of the

pulmonary epithelium induces the release of host components, such as adenosine

triphosphate and norepinephrine, that are sensed by biofilm-associated pneumococci,

triggering both the production of pneumococcal virulence factors and the dispersal of

bacteria (4, 37, 39, 40). This in turn facilitates bacterial spread to and colonization of the

lower respiratory tract (4).

IAV can infect not just the upper respiratory tract, but also the lungs, and murine

models featuring sequential pulmonary challenge with IAV first followed by pneumococcus

show the virus can compromise immune defense against pneumococcus (29, 34, 41). In

these models, IAV alters both the pulmonary environment and the immune response to

enhance subsequent bacterial colonization and tissue damage. For example, IAV pre-

infection increases mucus production and fibrosis and dysregulates ciliary function (34, 42,

43), thus impairing mechanical clearance of invading bacteria. Viral enzymes, along with

virus-elicited inflammation, result in the exposure of epithelial proteins that promote

pneumococcal adherence to and invasion of host cells (41, 44-46). Further, IAV triggers

type I and II interferon (IFN) responses that impair both the recruitment and antibacterial

function of phagocytes key to defense against pneumococcus (47-50). The combined

tissue damage and the compromise in immune function render the lung more permissive

for invasive *S. pneumoniae* infection (34, 45, 47). Less understood is how *S. pneumoniae*

infection may alter host antiviral responses and viral replication in the lung.

Notably, advanced age and IAV infection appear to synergistically enhance

susceptibility to pneumococcal lung infections (6, 18, 51, 52). Indeed, individuals ≥ 65

years old account for 70-85% of deaths due to pneumonia and influenza (52). Interestingly,

elderly individuals with influenza-like symptoms were reported to have an increased

pneumococcal carriage rate of 30% (53). In animal models in which bacteria are directly

instilled into the lungs following influenza infection, aging is associated with increase

susceptibility of secondary pneumococcal pneumonia (51). Age-dependent changes in the

expression of key components of innate immune signaling contribute to disease in this co-

infection model (51). However, the factors, including those that are age-dependent, that

trigger the transition of pneumococci from benign colonizer to pathogen are poorly defined,

in part because small animal models that recapitulate the transition from asymptomatic

colonization to overt clinical illness are lacking.

These above studies indicate that bacterial-viral synergy is multi-factorial and can

occur at different sites with the host. Insight into events that occur in both the nasopharynx

and lung and contribute to the heightened susceptibility of the aged to serious disease

upon pneumococcal/IAV coinfection is needed to develop better therapeutic and

preventative approaches. A current murine model for the spread of nasopharyngeal

pneumococci to the lung after viral infection of the upper respiratory tract relies on initial

bacterial colonization of the nasopharynx followed by viral infection, but does not

recapitulate the severe signs of human clinical disease (4). Influenza virus is capable of

infection not just of the upper respiratory tract, but also the lung (54, 55). To better

investigate the transition of *S. pneumoniae* from asymptomatic colonizers to invasive

pathogens following IAV infection, as well as the effect of host age on the disease process,

we built upon this mouse model of *S. pneumoniae/*IAV co-infection by incorporating viral

infection of the lower respiratory tract. The enhanced model recapitulates the severe and

age-exacerbated clinical disease observed in humans.

RESULTS

Intranasal IAV inoculation of mice pre-colonized with *S. pneumoniae* strain TIGR4

does not result in disease.

Biofilm-grown *S. pneumoniae* are relatively less virulent and thus adapted to host colonization rather than disease (2-4). A previously established murine model in BALB/cByJ mice utilizes biofilm-grown *S. pneumoniae* serotype 2 strain D39 and serotype 19F strain EF3030 to establish heavy carriage in the nasopharynx (4). Then, two days after bacterial inoculation, IAV is introduced into the nasal cavity and results in the spread of pneumococci from the nasopharynx (NP) to the lung (4). We first recapitulated this model with *S. pneumoniae* strain TIGR4, an invasive serotype 4 strain (56, 57) that we previously found to be highly virulent in aged C57BL/6 mice (19). We intranasally (i.n.) inoculated young (8-10 weeks) C57BL/6 (B6) mice with 1×10^6 colony forming units (CFU) of biofilm-generated *S. pneumoniae* TIGR4 by delivering the bacteria in 10 µl (a volume that is unlikely to inoculate the lung (58)) to the nares of non-anesthetized mice. Forty-eight hours later, mice were i.n. inoculated with 10 µl containing 20 PFU of Influenza A (IAV) virus PR8. (See Fig. S1A for general scheme). Control groups of mice were either infected with *S. pneumoniae* or IAV alone as controls. However, under these conditions, mice did not display signs of sickness, nor did bacteria spread into the lungs after seven days (not shown).

To increase the likelihood of clinical disease, we repeated the experiment with a five-fold higher (5×10^6 CFU) dose of bacteria and a 25-fold higher dose (500 PFU) of IAV. Similar to earlier reports for other *S. pneumoniae* strains (D39 and EF3030) (4), IAV co-infection resulted in a 10-fold increase in *S. pneumoniae* TIGR4 in the nasal lavage fluid at 2 days post-IAV infection (Fig. S1B). In addition, IAV co-infection was associated with the

detection of bacteria in the lungs of 40% of mice, compared to none in the control group

infected with *S. pneumoniae* TIGR4 alone. This trend is consistent with the previous

BALB/cByJ mice model of coinfection (4), but did not reach statistical significance.

Furthermore, co-infection was not associated with weight loss when assessed over the

course of 4 days post-infection (Fig. S1C). We also scored mice for clinical signs of the

disease based on weight loss, activity, posture and breathing and ranging from healthy

[score = 0] to moribund [score = 25] and requiring euthanasia if the score was >9 as

previously described (59). As secondary pneumonia can occur several days following IAV

(41), we monitored the disease course up to 7 days, but did not detect disease symptoms

or death in any of the co-infected mice (100% survival and 0 daily clinical score, including

weight loss for all mice). Therefore, despite promoting bacterial dispersal from the

nasopharynx into the lungs, similar to the previous work with other *S. pneumoniae* strains

and BALB/c mice (4), this model of *S. pneumoniae* TIGR4/IAV coinfection did not result in

overt clinical signs of disease.

Combined intranasal/intratracheal IAV inoculation of *S. pneumoniae*-colonized mice

results in bacterial dissemination and disease

IAV infection is not restricted to the upper respiratory tract, and can cause viral

pneumonia in a significant fraction of infected individuals (54, 55) that is likely to be crucial

for creating an environment in the lungs that is more permissive for bacterial infection (29,

34, 41). Indeed, viral lung infection diminishes pulmonary defenses against *S. pneumoniae*

and promotes secondary bacterial pneumonia (34, 41, 44-50). Delivery of IAV i.n. to

BALB/cByJ mice results in signs of viral pneumonia (4), but we found that pulmonary

access of inocula delivered via the nasopharynx is more restricted in B6 mice as compared

to BALB/c mice (unpublished observation), raising the possibility that the lack of disease

observed in co-infected B6 mice was due to the exclusive localization of virus in the nasal

cavity, with limited opportunity to alter systemic or pulmonary immunity. In fact, when we

measured pulmonary viral load two days following i.n. infection with 500 PFU IAV, we were

unable to detect any PFU in the lungs, while delivery of 20 PFU of IAV by intratracheal (i.t.)

inoculation was sufficient to establish lung infection (Fig. S2).

Therefore, to ensure delivery of IAV to both the nasopharynx and the lungs, we co-

infected *S. pneumoniae*-colonized B6 mice by delivering the virus by two routes. Mice

were inoculated i.n. with 5×10^6 CFU of biofilm grown S. *pneumoniae* TIGR4 and 48 hours

later infected not only with 500 PFU IAV i.n., but also 20 PFU i.t. to ensure pulmonary

infection (Fig. 1A). No mice in a control group inoculated i.n. with *S. pneumoniae* alone lost

weight (Fig 1B), displayed clinical signs of sickness (Fig 1C), or died (Fig 1D). A second

control group, inoculated i.n. and i.t. with IAV alone displayed no disease until after day 4,

when they exhibited weight loss (Fig 1B) and began succumbing to viral infection (Fig 1D).

In contrast, inoculation of IAV to animals pre-colonized with *S. pneumoniae* caused a

bacterial/viral co-infection that resulted in weight loss (Fig. 1B), clinical symptoms (Fig. 1C)

and death (Fig. 1D) that were detected starting day 2 post IAV introduction. At this time

point, a higher fraction of co-infected mice displayed signs of disease as compared to

controls infected with IAV only (55% vs 64%), and disease was more severe in mice that

displayed clinical symptoms, although this did not reach statistical significance. In addition,

the overall survival rate among co-infected mice was significantly lower than mice singly

infected with *S. pneumoniae* alone (Fig. 1D). These signs of exacerbated disease were

associated with significantly higher bacterial burdens in the nasopharynx as well as

translocation of *S. pneumoniae* into the lung in comparison to mice colonized with the

bacteria or mice infected with IAV alone (Fig. 1E). These findings demonstrated that the

new co-infection model leads to disease in a fraction of young healthy mice, which may

increase the likelihood of detecting enhanced susceptibility in vulnerable hosts.

PMN depletion may have a small effect on the course of disease during IAV/*S.*

***pneumoniae* co-infection**

We previously found that in primary pneumococcal pneumonia, PMNs are required

to control bacterial numbers early in the infection process; however, their persistence in

the lungs is detrimental to the host and can promote the infection at later time points (20).

To address the role of PMNs during co-infection, we treated young mice with PMN-

depleting anti-Ly6G antibody (1A8) one day prior to pneumococcal colonization and

throughout the co-infection (based on timeline in Figure 1A). We then confirmed that the

cells were depleted by staining with the RB6 antibody followed by flow cytometry (Fig S3).

Following infection, we measured bacterial burden, weight loss, clinical score and survival

over time. PMN depletion had no effect on bacterial burdens in the nasopharynx or

bacterial spread to the lungs or blood following co-infection (Fig 2A and B). However,

PMN-depleted mice appeared to lose more weight at day 3 and 4 post co-infection as

compared to the control group (Fig. 2C), and a greater proportion of PMN-depleted mice

displayed clinical signs of sickness as compared to the control group at both 18 hours and

48 hours post IAV infection (Fig. 2D). Additionally, lower survival was observed in the

PMN-depleted group, in which 25% survived to day 7 compared to ~43% in the untreated

control group (Fig. 2E). These differences did not reach statistical significance, but raise

the possibility that PMNs provide a measure of defense to co-infection in young mice.

Aging increases susceptibility to IAV/*S. pneumoniae* co-infection.

We next tested if the new co-infection mouse model (Fig. 1A) recapitulates the age-associated increase in susceptibility to secondary pneumococcal pneumonia. Old (20-24 months) B6 mice were inoculated i.n. with 5×10^6 CFU of biofilm grown S. *pneumoniae* TIGR4 and 48 hours later were infected with 500 PFU IAV i.n. plus 20 PFU i.t. When compared to young co-infected controls, old mice displayed significantly more severe signs of disease, as indicated by a higher average clinical score (Fig. 3A). While seven of 16 (~44%) young mice showed clinical symptoms (i.e., clinical score greater than 1; Fig. 3A), all 13 old mice showed at least some degree of illness by day 2 post co-infection (p = 0.0012, by Fisher's exact test). Furthermore, whereas only 25% (4 out of 16) young mice had a clinical score greater than 2, which is indicative of more severe disease, 92% (12 out of 13) aged mice fell into this category (p = 0.005, by Fisher's exact test). In addition, old mice died at a significantly accelerated rate. By day 2 post co-infection, 60% of old mice had succumbed to the infection compared to only 25% of young mice. Differences in survival were observed at each successive time point, and at the end of the experiment on day 8, only 14% of old mice remained alive compared to 50% of young mice (Fig. 3B). Importantly, the accelerated death observed in co-infected old mice was not observed in old mice infected with S. *pneumoniae* alone or IAV alone (Fig S4).

We next tested whether these differences could be attributed to increased bacterial loads in the nasopharynx, lungs, or blood. As old mice got sicker at earlier time points after IAV co-infection, with the majority succumbing by day 4, we compared bacterial burden across age groups at 18 and 48 hours after IAV co-infection. We found no significant differences in the numbers of pneumococci in nasopharyngeal washes, pulmonary homogenates, or blood at either time point (Fig 3C-E). Taken together, these findings

suggest that with aging there is an increased susceptibility to co-infection and an

accelerated course of disease that could not be attributed to a more rapid bacterial

dissemination or higher bacterial loads in the nasopharynx, lung, or bloodstream.

Aging is associated with more rapid lung inflammation.

To determine if the accelerated rate of death examined in co-infected old mice was

due to more lung damage, we analyzed H&E-stained lung sections for alveolar congestion,

hemorrhage, alveolar thickness, neutrophils and lymphocytic infiltration (Fig. 4A). We

found that the alveolar spaces of both uninfected old and young mice were clear and free

of inflammatory or red blood cells (Fig. 4A). At 18h hours post co-infection, the lungs of

young mice did not show any overt signs of disease (Fig. 4A). In contrast, co-infected old

mice had significant lung pathology by 18 hours post infection, including a loss of alveolar

architecture, and spotty inflammation consisting of infiltrates composed of neutrophils,

alveolar macrophages and mononuclear cells that were mixed with red blood cells (Fig.

4A). At 48 hours post-infection, there were clear signs of lung pathology in both young and

old co-infected mice (Fig. 4A).

Next, we tested the levels of inflammatory cytokines in co-infected young vs old

mice. No significant differences between age groups were detected in baseline

(uninfected) levels of any of the cytokines tested between the age groups (Fig 4B and not

shown). However, consistent with the enhanced PMN influx at 18 h post-infection in aged

mice, old mice had significantly higher levels of IL-10, IL-2, IL-1β and TNFα (Fig 4B)

compared with young mice. Levels of IL-12p70, IL-17, IL-6 and IFNγ, were slightly but not

significantly elevated 18 hours post co-infection (Fig S5A). By 48 hours there were no

significant differences between young and old mice in cytokine levels except for IFNγ,

which was higher in young mice (Fig S5B).

We previously found that mice suffering from exacerbated PMN-mediated

pulmonary inflammation during pneumococcal pneumonia did not display higher bacterial

burdens in their lungs (27) despite a higher likelihood of severe disease (19, 20, 27). To

test whether PMN influx was also higher in old co-infected mice, we measured the

percentage and number of pulmonary PMNs (Ly6G$^+$) by flow cytometry. We found that old

mice had significantly (6-fold) higher percentages and numbers of PMNs in their lungs as

compared to young controls at 18 hours post co-infection (Fig 4C). By 48 hours, most aged

mice had succumbed to the infection (Fig. 3B), confounding interpretation of PMN

numbers at this time point; PMN percentages and numbers appeared to be higher in

young mice, but the differences were not statistically significant (Fig. 4C). Macrophages,

which are important for host resistance to *S. pneumoniae/IAV* co-infection (49) and display

age-driven changes (60, 61), displayed no significant age-dependent differences in either

percentage or number at 18 or 48 hours after infection (Fig. S6). Taken together, these

findings demonstrate that aging is associated with earlier pulmonary inflammation and

damage following co-infection, which may contribute to the accelerated death observed in

this mouse group.

Aging and IAV infection diminish the ability of PMNs to kill *S. pneumoniae ex vivo*.

Aged, co-infected mice experience an accelerated rate of pulmonary inflammation

but bacterial loads in the lungs of aged mice were not lower than in young mice, indicating

that PMN infiltration is not associated with bacterial clearance. Both aging (24) and IAV

infection (62) have been reported to diminish antibacterial function of PMNs. To assess the

ability of PMNs to kill *S. pneumoniae* in our co-infection model, we used a well-established

opsonophagocytic (OPH) killing assay (19, 63). We first compared the bactericidal activity

of bone marrow-derived PMNs from young or aged mice. The percentage of bacteria killed

upon incubation with PMNs for 45 or 90 minutes was determined by comparing surviving

CFU to no PMN control reactions at the same timepoint. We found that as previously

reported for humans (23) and mice (24) the ability of PMNs isolated from uninfected old

mice to kill pneumococci was reduced 5-fold compared to young controls regardless of the

duration of infection (Fig. 5).

We next examined the bactericidal activity of PMNs isolated from young or old mice

2-days after i.t./i.n. IAV inoculation. As previously reported (47-50), PMNs from young IAV

infected mice had a slight (2-fold) but significant reduction in their ability to kill

pneumococci as compared to PMNs from uninfected controls (Fig. 5). Strikingly, IAV

infection diminished the ability of PMNs from old mice to kill *S. pneumoniae*; instead,

PMNs from IAV infected old mice promoted a slight increase in bacterial numbers (Fig. 5).

These findings suggest that IAV infection completely abrogates the ability of PMNs from

old mice to kill *S. pneumoniae.*

Aging and prior colonization with *S. pneumoniae* result in impaired IFN-α production

and higher viral burden in the lungs.

Finally, we investigated whether aging and/or bacterial co-infection had an impact

on antiviral responses. Old or young B6 mice were inoculated i.n. with 5×10^{6} CFU of

biofilm grown S. *pneumoniae* TIGR4 or mock challenged with PBS and 48 hours later were

infected with 500 PFU IAV i.n. and 20 PFU i.t. At 48 hours following viral infection, we

compared viral burden in the lung across age groups. We found that bacterial colonization

resulted in 10-fold (and statistically significant) higher pulmonary viral loads when

compared to mock-colonized controls, regardless of host age (Fig. 6A). Further, we found

that in co-infected hosts, aging was associated with significantly increased viral loads in

the lungs (Fig. 6A), suggesting that enhanced viral loads and impaired antiviral defenses

contribute to the differences in clinical manifestation across host age.

To test whether bacterial colonization and aging impair antiviral immune responses,

we measured the levels of IFN-α, a cytokine crucial for antiviral defense (64). We found

that in young mice, despite the higher viral burden (Fig. 6A), prior bacterial colonization

resulted in a 3-fold (but not statistically significant) decrease in IFN-α production in the

lungs in response to IAV challenge (Fig. 6B). Notably, aging was associated with

(statistically significant) 5- and 3-fold lower levels of IFN-α correspondingly in mice infected

with IAV alone or co-infected with IAV and *S. pneumoniae.* No differences in IFN-α

production were observed in the sera (Fig S7). These findings suggest that both

pneumococcal infection and aging blunt antiviral responses in this mouse model.

DISCUSSION

S. pneumoniae remains a leading cause of secondary bacterial pneumonia

following influenza A virus infection and is associated with severe disease (30, 34-36),

particularly in the elderly (52). The majority of *S. pneumoniae/*IAV co-infection

experimental studies have delivered bacteria into the lungs of mice pre-infected with IAV to

reveal changes in the host lungs and immune response that are crucial for priming

invasive pneumococcal disease (34, 41, 44-50). In this study, to investigate the transition

of *S. pneumoniae* from colonizer to pathogen upon IAV co-infection, a process that has

just started to be elucidated (4, 37), we have developed a modified murine infection model

that recapitulates this transition and results in severe clinical disease. In a previously

established model, female BALB/cByJ mice were first colonized intra-nasally with biofilm-

grown pneumococci and then infected with IAV by delivering the virus to the nasopharynx

(4, 40). This model showed that changes in the host environment in response to viral

infection triggers the dispersal of pneumococci from colonizing biofilms and their spread to

the lower respiratory tract (4). Importantly, the dispersed bacteria expressed higher levels

of virulence factors required for infection, thus, rendering them more highly pathogenic (4).

Nevertheless, upon dispersion following *in vivo* IAV infection, although bacterial migration

to the lung was detected, the burden was less than 100 CFU per lung and the mice did not

suffer overt disease. When we used this model to co-infect male C57BL/6 (B6) mice, we

also observed a significant increase in dispersed bacteria in the nasopharynx, but no

disease and only a transient presence of *S. pneumoniae* in the lungs of B6 mice. Here we

performed experiments in male instead of female mice due to both the easier availability of

aged male animals and the documented higher rate of pneumococcal pneumonia in men

compared to women (65, 66).

Previous work of pneumococcal inoculation of IAV-infected lungs showed viral

infection to be crucial for creating an environment in the lungs that is more permissive for

bacterial infection (41, 44-50). The relatively low bacterial burden in co-infected male B6

mice observed here upon IAV inoculation only by the i.n. route suggested that the lung

environment was not sustaining the bacteria. It is possible that human IAV/*S. pneumoniae*

co-infection involves viral infection of not only the upper respiratory tract, but the lower

respiratory tract as well (54, 55). Therefore, we co-infected *S. pneumoniae* colonized B6

mice with IAV by delivering the virus not only i.n. to infect the nasopharynx, but also i.t. to

ensure infection of the lungs. This modified model recapitulated the increase in non-

adherent pneumococci in the nasopharynx observed upon viral co-infection (4, 67, 68),

and resulted in bacterial spread into the lungs and circulation that increased over time.

Importantly, this mode of dual infection recapitulated both the increased colonization

burden (69, 70) as well as the clinical signs of severe disease observed in humans (30, 34-

36), and resulted in death of approximately half of co-infected young controls.

Although the elderly are at higher risk for secondary pneumococcal pneumonia

following IAV infection, animal studies exploring this age-driven susceptibility to co-

infection are few (51, 71, 72). Here, using the modified model, we found that old mice were

significantly more susceptible to *S. pneumoniae*/IAV co-infection. Old mice displayed more

severe signs of disease as compared to young controls and the majority (>85%) failed to

survive the co-infection. This increased susceptibility in old mice was not linked to higher

bacterial dissemination from the nasopharynx, greater establishment of infection in the

lungs, or systemic spread into the circulation at the time points tested. Susceptibility to viral

infection alone, as measured by weight loss within the first 5 days following IAV, was also

similar between the age groups. Rather, the age-driven susceptibility to co-infection was

associated with earlier and more severe pulmonary inflammation. Production of pulmonary

cytokines is elevated in young mice infected with *S. pneumoniae* at 7 days after IAV

inoculation (73). Further, old mice display changes in the expression of pattern recognition

receptors in the lungs leading to altered inflammatory responses (51). Similar to previous

studies (51), we found here that co-infected old mice had higher levels of TNFα compared

to young controls. However, in contrast to previous reports that found reduced NLRP3

inflammasome expression in the lungs and lower production of IL-1β, we found higher

levels of IL-1β in old vs. young mice. This may be accounted for by differences in

expression of bacterial factors. The expression of pneumolysin, which was found to

activate NLRP3 inflammasomes and lead to production of IL-1β (74), was elevated in

pneumococci dispersed from biofilms upon IAV infection (75) and therefore may have

primed the IL-1β production we observed in old mice.

We previously found that PMNs were key determinants of disease during primary

pneumococcal pneumonia and are required to initially control bacterial numbers (20, 76).

Therefore, we explored here the role of PMNs in *S. pneumoniae*/IAV co-infection. Similar

to other studies, we found that PMNs are recruited to the lungs of co-infected young mice

(49). Previous reports indicate that the PMN-mediated anti-pneumococcal function in IAV-

infected mice (62) and humans (70) is progressively reduced over time. For example, in

mice, PMNs demonstrably contribute to host defense at 3 days but not at 6 days post-

infection (62). Nevertheless, PMN depletion showed that these cells are important for

control of bacterial transmission (77) and control of pulmonary bacterial numbers (49) in

IAV co-infected mice. Similarly, here we found that PMN depletion starting prior to bacterial

colonization and continuing throughout viral co-infection appeared to slightly worsen

disease progression, with slightly greater average weight loss at days 3 and 4 after viral

inoculation and a greater proportion of PMN-depleted mice succumbing to co-infection.

However, PMN depletion had no significant effect on the number of dispersed *S.*

pneumoniae in the nasopharynx or their spread to the lungs and blood. It is possible that,

in this model, IAV may rapidly impair PMN function, limiting their efficacy even in PMN-

replete mice. In fact, we found that within 2 days following IAV infection, the ability of bone

marrow-derived PMNs to kill *S. pneumoniae* was significantly blunted in young mice.

Alternatively, the apparent inability of PMNs to limit bacterial numbers in this model could

be due to the enhanced virulence of pneumococci dispersed from the nasopharyngeal

environment (4) compared to broth-grown bacteria typically used in other models (62).

In this study we found that aging was associated with earlier influx of PMNs into the

lungs of co-infected mice. We previously demonstrated that excessive PMN influx into the

lungs is detrimental for the ability of old mice to control invasive disease following primary

pneumococcal pneumonia (19) and that PMN depletion 18 hours after infection boosted

host survival (20). Similarly, greater PMN influx into the lungs of old mice singly infected

with IAV was associated with host mortality, and depletion of these cells six days following

viral infection significantly boosted host survival (78). Uncontrolled PMN influx can result in

tissue damage, disruption of gaseous exchange and pulmonary failure. In fact, it was

reported that in IAV singly infected old mice, PMNs enhanced lung inflammation and

damage and their depletion reduced the levels of inflammatory IL-1β and TNFα (78).

Therefore, the early increase in these inflammatory cytokines and lung damage we

observed here in *S. pneumoniae*/IAV co-infected old mice may be driven by the elevated

levels of pulmonary PMNs.

The age-associated increased levels of pulmonary IL-10 during *S. pneumoniae*/IAV

confection observed here may also contribute to the enhanced susceptibility of old mice to

co-infection. IL-10 levels are elevated in co-infected young mice compared to those singly

infected with *S. pneumoniae* (79). We showed that blocking this cytokine boosts PMN anti-

bacterial function by inhibiting ROS production (76), and van der Poll and coworkers

demonstrated that inhibition of IL-10 restores resistance of vulnerable hosts to primary

pneumococcal pneumonia (79).

The more severe disease observed in old mice also correlated with higher viral lung

burden during both co-infection with *S. pneumoniae* and IAV and single infection with IAV,

although the difference reached statistical significance only in the former. Production of

type I IFN, which is crucial for control of viral replication, is dysregulated during aging (80)

and we found that pulmonary IFN-α levels of co-infected or singly infected old mice were

significantly lower compared their young counterparts. Hence, in this model, enhanced

disease associated with aging may be a reflection of a combination of an exuberant PMN

response and a muted type I interferon response.

While many studies have examined the effect of IAV on secondary pneumococcal

pneumonia, the effect of bacterial colonization on viral infection has been less explored.

We found that, regardless of host age, prior colonization with pneumococci resulted in

significantly higher viral pulmonary loads. In experiments involving two different hosts

(ferrets or cotton rats) and two different viruses (IAV and Human RSV, respectively),

nasopharyngeal colonization of donors with *S. pneumo*niae promoted viral transmission

(81, 82). In the latter (rat-RSV) model, prior pneumococcal colonization enhanced viral

infection of the upper respiratory tract but did not promote infection of the lungs (82). A

human experimental pneumococcal colonization study showed that prior colonization with

pneumococci did not increase the burden of a live attenuated influenza virus, but reduced

pro-inflammatory immune responses in the nasopharynx and significantly blunted antiviral

IgG production in the lungs (83). Hence, prior colonization with pneumococci may impair

antiviral defenses. Indeed, we found that in spite of their approximately one log higher viral

lung burden compared to singly infected mice, *S. pneumo*niae-pre-colonized young and

old mice respectively displayed 5 and 3-fold lower levels of pulmonary IFN-α compared to

non-colonized controls.

In summary, here we modified existing murine models to establish an experimental

system that reflects the transition of *S. pneumoniae* from asymptomatic colonizer to

invasive pulmonary pathogen upon IAV co-infection. In this model, IAV triggers the

transition of a pathobiont from a commensal to a pathogenic state through modification of

bacterial behavior in the nasopharynx, enhancement of bacterial colonization in the lung, and compromise of PMN-mediated anti-bacterial immunity. In turn, pneumococci modulate antiviral immune responses and promote IAV infection of the lower respiratory tract. Importantly, this model recapitulates the susceptibility of aging to co-infections. Moving forward, this multi-step model can be used to dissect both the multiple phases of pneumococcal disease progression from commensals to pathogens and the complexity of viral/bacterial interactions within different hosts, thus helping inform specialized treatment options (67) tailored to the susceptible elderly population.

MATERIALS AND METHODS

Mice. Young (8-10 weeks) and aged (18-24 months) male C57BL/6 mice were purchased from Jackson Laboratories (Bar Harbor, ME) and the National Institute of Aging. Mice were housed in a pathogen-free facility at Tufts University. All procedures were performed in accordance with Institutional Animal Care and Use Committee guidelines. The number of mice included in each experiment is based on power analysis. At least 6 mice per group for determination of bacterial burden and 12 mice per group for monitoring survival were included to sufficiently power our studies. In each experiment, an equal number of mice per group were planned. However, slight variations in the number of mice per group at later time points of several experiments occurred due to the required euthanasia of several mice in highly susceptible experimental groups.

Bacterial biofilms. NCI-H292 mucoepidermoid carcinoma cells (H292) were grown in 24-well plates in RPMI 1640 media with 10% FBS and 2mM L-glutamine until confluent. Cells were washed with 1x PBS and fixed in 4% paraformaldehyde for 1 hour on ice. *S.*

pneumoniae TIGR4 (kind gift from Andrew Camilli) were grown on Tryptic Soy Agar plates

supplemented with 5% sheep blood agar (blood agar plates) overnight, then diluted and

grown in chemically defined liquid medium (CDM) (4, 40) supplemented with Oxyrase until

OD_{600nm} of 0.2. Bacteria were diluted 1:1000 in CDM and seeded on the fixed H292 cells.

The, bacteria/H292 cells were incubated at 34°C/ 5% CO_2 and media was changed every

12 hours. At 48 hours post-infection, the supernatant containing planktonic bacteria (non-

adherent to NCI-H292 cells) cells was discarded, the cells gently washed with PBS and

adherent biofilms collected in fresh CDM by vigorous pipetting. Biofilm aliquots were then

frozen at -80°C in the CDM with 25% (v/v) glycerol.

Intranasal inoculation. Before use, biofilm aliquots were thawed on ice, washed once and

diluted in PBS to the required concentration. The mice were restrained without anesthesia

and infected i.n. with 10µl ($5x10^6$ CFU) of biofilm grown *S. pneumoniae*. The inoculum was

equally distributed between the nostrils with a pipette. This method of inoculation results in

pathogen delivery limited to the nasal cavity, without accessing the lower tract in C57BL/6

mice (58). Bacterial titers were confirmed by serial dilution and plating on blood agar

plates. To ensure stable colonization of the biofilm in the nasopharynx, groups of mice

were euthanized at 18 and 48 hours post inoculation, and the nasal washes and tissue

were collected and plated on blood agar plate for enumeration of *S. pneumoniae*.

Viral infection. The mouse-adapted H1N1 Influenza A virus PR8 (A/PR/8/34) was

obtained from Dr. Bruce Davidson (4, 40) and stored at -80 □C. Before use, viral aliquots

were thawed on ice, diluted in PBS and used to inoculate mice. At 48 hours following

bacterial inoculation, mice were infected i.n. with 10µl of 500 plaque forming units (PFU) of

virus by pipetting the inoculum into the nostrils of non-anesthetized mice. Following i.n.

inoculation, mice were lightly anesthetized with isoflurane and challenged i.t. with 20 PFU

of virus in a 50µl volume pipetted directly into the trachea with the tongue pulled out to

facilitate delivery (20).

Clinical scoring and bacterial burden. Following co-infection, mice were monitored daily

and blindly scored for signs of sickness including weight loss, activity, posture and

breathing. Based on these criteria, the mice were given a clinical score of healthy [0] to

moribund [25] modified from what was previously described (59). Any mice displaying a

score above 9 are humanely euthanized in accordance with our protocol. Mice were

euthanized at indicated time points and the lung, nasal lavages and blood collected and

plated on blood agar for enumeration of bacterial loads, as previously described (20). For

collection of sera, blood was collected via cardiac puncture into Microtainer[®] tubes (BD

Biosciences) and centrifuged at 7607xg for 2 minutes to collect serum as per

manufacturer's instructions.

Depletion of PMNs. Mice were intraperitoneally (i.p.) injected with 100µl (50µg/mouse) of

anti-Ly6G clone 1A8 (BD Biosciences) to deplete neutrophils. Mice were injected daily with

the depleting antibodies starting one day prior and ending two days after bacterial

inoculation, followed by every other day from day 1 post viral co-infection to the end of

each experiment. Treatment resulted in >90% neutrophil depletion as described below.

Cell isolation and Flow Cytometry. Mice lungs were harvested, washed in PBS, and

minced into small pieces. The sample was then digested for 45 minutes with RPMI 1640

1X supplemented with 10% FBS, 1 mg/ml Type II collagenase (Worthington, Lakewood,

NJ), and 50 U/ml deoxyribonuclease I to obtain a single-cell suspension as previously

described (20). The red blood cells were lysed using ACK lysis buffer (Gibco). Cells were

then stained with anti-mouse Ly6G clone 1A8 (BD Biosciences), F480 clone BM8

(BioLegend), CD11c clone N418 (eBioscience), and CD11b clone M1/70 (eBioscience).

For neutrophil depletion, cells isolated from the lungs at 18 and 48 h post co-infection were

also stained with either Ly6G clone 1A8 or RB6 clone RB6-8C5 (BioLegend) antibodies to

confirm cell depletion. The fluorescence intensities were measured on BD LSR II Flow

Cytometer at Tufts FACS Core Facility (Boston, MA) to capture at least 25,000 cells and

analyzed using FlowJo.

Isolation of PMNs and Opsonophagocytic Killing Assay (OPH). Femurs and tibias of

uninfected mice were collected and flushed with RPMI, supplemented with 10% FBS and 2

mM EDTA to obtain bone marrow cells. Neutrophils were isolated by density gradient

centrifugation, using Histopaque 1119 and Histopaque 1077 as previously described (84).

The neutrophils were resuspended in Hank's (Gibco) buffer/0.1% gelatin with no Ca+ or

Mg+ and tested for purity by flow cytometry using anti-Ly6G antibodies (eBioscience)

where 85-90% of enriched cells were Ly6G+. The ability of neutrophils to kill bacteria was

measured using a well-established opsonophagocytic (OPH) killing assay as previously

described (20). Briefly, 2.5×10^5 neutrophils were incubated in Hank's buffer/0.1% gelatin

with 10^3 CFU of *S. pneumoniae* pre-opsonized with 3% mouse sera. The reactions were

incubated in flat bottom 96-well non-binding plates for 45 minutes at 37°C. Each group

was plated on blood agar to enumerate viable CFU. Percent bacterial killing was

calculated in comparison to a no PMN control under the same conditions.

Histology. Whole lungs were harvested from groups of mice at 18 hours and at 48 hours

post co-infection and fixed in 10% neutral buffered formalin for 2 days. The tissues were

then embedded in paraffin, sectioned at 5 µm and stained with Hematoxylin and Eosin at

the Animal Histology Core at Tufts University. Sections of lung from three mice per group

were imaged using a Nikon Eclipse E400 microscope. Photomicrographs were captured

using a SPOT Idea 5.0-megapixel color digital camera and SPOT software.

Histopathologic scoring was performed by a board-certified anatomic pathologist

experienced in murine pathology, from 0 (no damage) to 4+ (maximal damage) for alveolar

congestion, hemorrhage, alveolar thickness, neutrophils, and lymphocytic infiltration (85).

Cytokine Analysis. Frozen lung homogenates and serum samples were thawed on ice

and mixed by gentle vortexing. Cytokines in the lungs and serum samples were measured

using Mouse Cytokine 8-Plex Array (Quanterix, Billerica, MA) following the manufacturer's

instructions. Levels in lung supernatants and serum samples were measured using the

Cirascan at the Imager at Tufts University Genomic Core (Boston, MA) and analyzed by

the Cirascan/Cirasoft program. Qlucore Omic Explorer (version 3.5) was used for the

generation of lung cytokine box plots. Concentrations of cytokines (IL-10, IL-2, IL-1β,

TNFα, IL-6, IFNγ, IL-17 and IL-12p70) were log-transformed, and displayed as Log_2 pg/ml.

IFNα was measured using Mouse IFN-alpha ELISA kit (R&D system, MN). following the

manufacturer's instructions.

Plaque assay. Madin-Darby Canine Kidney (MDCK) cells were grown overnight in 12 well

plates at 2×10^5 cells/well in DMEM media +10% FBS. Cells were washed twice with 1xPBS

and incubated with serial dilutions of viral inoculum or lung homogenates in DMEM

supplemented with 0.5% low endotoxin BSA (Sigma-Aldrich) for 50 minutes in 37 □C with

5% CO_2 incubator. The plates were shaken every 10 minutes during the incubation and

then washed twice with 1xPBS. 2mL of 2.4% of avicel overlay (FMC) was then added onto

the infected cells and were incubated for 3 days in 37 □C with 5% CO_2 incubator. After 3

days, avicel overlays were removed and cells were washed with 1x PBS and fixed in 4%

paraformaldehyde for 30 minutes at room temperature. 1% crystal violet were then added

for 5 minutes to count plaques.

Statistical Analysis. Statistical analysis was performed using Graph Pad Prism7. CFU

data were log-transformed to normalize distribution. Data are presented as mean values

+/- SEM. Significant differences ($p < 0.05$) were determined by Student's t-test. Differences

in fractions of mice that got sick were measured using Fisher's exact test. Survival

analysis, including the kinetics by which mice succumb to infection, was performed using

the log rank (Mantel-Cox) test. Asterisks indicate significant differences and p values are

noted in the figures.

ACKNOWLEDGEMENTS

We would like to acknowledge James Nicholas Lee, Summer Schmaling, and Ognjen

Sekulovic for technical assistance with clinical score, virus preparation and nasal lavage

respectively. We would also like to thank Andrew Camilli for bacterial strains and Marta

Gaglia for protocols. They, along with Tim van Opijnen, Bharathi Sundaresh and Marcia

Osburne provided important feedback on the manuscript.

FUNDING

Research reported in this publication was supported by the National Institute On Aging of the National Institutes of Health under Award Number R21 AG064215 to E.B.G. and F31 AI122615-01A1 to S.R.; King Abdullah Scholarship Program (KASP) implemented by the Ministry of Higher Education (MOHE) under Award Number 7896504 to B.H.J.

REFERENCES

1.   Kadioglu A, Weiser JN, Paton JC, Andrew PW. 2008. The role of Streptococcus

pneumoniae virulence factors in host respiratory colonization and disease. Nat Rev

Microbiol 6:288-301.

2.   Chao Y, Marks LR, Pettigrew MM, Hakansson AP. 2014. Streptococcus

pneumoniae biofilm formation and dispersion during colonization and disease. Front

Cell Infect Microbiol 4:194.

3.   Marks LR, Parameswaran GI, Hakansson AP. 2012. Pneumococcal interactions

with epithelial cells are crucial for optimal biofilm formation and colonization in vitro

and in vivo. Infect Immun 80:2744-60.

4.   Marks LR, Davidson BA, Knight PR, Hakansson AP. 2013. Interkingdom signaling

induces Streptococcus pneumoniae biofilm dispersion and transition from

asymptomatic colonization to disease. MBio 4.

5.   Blanchette-Cain K, Hinojosa CA, Akula Suresh Babu R, Lizcano A, Gonzalez-

Juarbe N, Munoz-Almagro C, Sanchez CJ, Bergman MA, Orihuela CJ. 2013.

Streptococcus pneumoniae biofilm formation is strain dependent, multifactorial, and

associated with reduced invasiveness and immunoreactivity during colonization.

MBio 4:e00745-13.

6.   Boe DM, Boule LA, Kovacs EJ. 2017. Innate immune responses in the ageing lung.

Clin Exp Immunol 187:16-25.

7.   Chong CP, Street PR. 2008. Pneumonia in the elderly: a review of the

epidemiology, pathogenesis, microbiology, and clinical features. South Med J

101:1141-5; quiz 1132, 1179.

8.   Bogaert D, De Groot R, Hermans PW. 2004. Streptococcus pneumoniae

677   colonisation: the key to pneumococcal disease. Lancet Infect Dis 4:144-54.

678   9.   Simell B, Auranen K, Kayhty H, Goldblatt D, Dagan R, O'Brien KL, Pneumococcal

679   Carriage G. 2012. The fundamental link between pneumococcal carriage and

680   disease. Expert Rev Vaccines 11:841-55.

681   10.   Orsi A, Ansaldi F, Trucchi C, Rosselli R, Icardi G. 2016. Pneumococcus and the

682   Elderly in Italy: A Summary of Available Evidence Regarding Carriage, Clinical

683   Burden of Lower Respiratory Tract Infections and On-Field Effectiveness of PCV13

684   Vaccination. Int J Mol Sci 17.

685   11.   Smith EL, Wheeler I, Adler H, Ferreira DM, Sa-Leao R, Abdullahi O, Adetifa I,

686   Becker-Dreps S, Esposito S, Farida H, Kandasamy R, Mackenzie GA, Nuorti JP,

687   Nzenze S, Madhi SA, Ortega O, Roca A, Safari D, Schaumburg F, Usuf E, Sanders

688   EAM, Grant LR, Hammitt LL, O'Brien KL, Gounder P, Bruden DJT, Stanton MC,

689   Rylance J. 2020. Upper airways colonisation of Streptococcus pneumoniae in adults

690   aged 60 years and older: A systematic review of prevalence and individual

691   participant data meta-analysis of risk factors. J Infect doi:10.1016/j.jinf.2020.06.028.

692   12.   Regev-Yochay G, Raz M, Dagan R, Porat N, Shainberg B, Pinco E, Keller N,

693   Rubinstein E. 2004. Nasopharyngeal carriage of Streptococcus pneumoniae by

694   adults and children in community and family settings. Clin Infect Dis 38:632-9.

695   13.   Flamaing J, Peetermans WE, Vandeven J, Verhaegen J. 2010. Pneumococcal

696   colonization in older persons in a nonoutbreak setting. J Am Geriatr Soc 58:396-8.

697   14.   Palmu AA, Kaijalainen T, Saukkoriipi A, Leinonen M, Kilpi TM. 2012.

698   Nasopharyngeal carriage of Streptococcus pneumoniae and pneumococcal urine

699   antigen test in healthy elderly subjects. Scand J Infect Dis 44:433-8.

15. van Deursen AMM, van Houten MA, Webber C, Patton M, Scott D, Patterson S,

Jiang Q, Gruber WC, Schmoele-Thoma B, Grobbee DE, Bonten MJM, Sanders

EAM. 2018. The Impact of the 13-Valent Pneumococcal Conjugate Vaccine on

Pneumococcal Carriage in the Community Acquired Pneumonia Immunization Trial

in Adults (CAPiTA) Study. Clin Infect Dis 67:42-49.

16. Esposito S, Mari D, Bergamaschini L, Orenti A, Terranova L, Ruggiero L, Ierardi V,

Gambino M, Croce FD, Principi N. 2016. Pneumococcal colonization in older adults.

Immun Ageing 13:2.

17. van Deursen AM, van den Bergh MR, Sanders EA, Carriage Pilot Study G. 2016.

Carriage of Streptococcus pneumoniae in asymptomatic, community-dwelling

elderly in the Netherlands. Vaccine 34:4-6.

18. Krone CL, van de Groep K, Trzcinski K, Sanders EA, Bogaert D. 2014.

Immunosenescence and pneumococcal disease: an imbalance in host-pathogen

interactions. Lancet Respir Med 2:141-53.

19. Bou Ghanem EN, Clark S, Du X, Wu D, Camilli A, Leong JM, Meydani SN. 2015.

The alpha-tocopherol form of vitamin E reverses age-associated susceptibility to

streptococcus pneumoniae lung infection by modulating pulmonary neutrophil

recruitment. J Immunol 194:1090-9.

20. Bou Ghanem EN, Clark S, Roggensack SE, McIver SR, Alcaide P, Haydon PG,

Leong JM. 2015. Extracellular Adenosine Protects against Streptococcus

pneumoniae Lung Infection by Regulating Pulmonary Neutrophil Recruitment. PLoS

Pathog 11:e1005126.

21. Garvy BA, Harmsen AG. 1996. The importance of neutrophils in resistance to

pneumococcal pneumonia in adult and neonatal mice. Inflammation 20:499-512.

22. Hahn I, Klaus A, Janze AK, Steinwede K, Ding N, Bohling J, Brumshagen C,

Serrano H, Gauthier F, Paton JC, Welte T, Maus UA. 2011. Cathepsin G and

neutrophil elastase play critical and nonredundant roles in lung-protective immunity

against Streptococcus pneumoniae in mice. Infect Immun 79:4893-901.

23. Simell B, Vuorela A, Ekstrom N, Palmu A, Reunanen A, Meri S, Kayhty H,

Vakevainen M. 2011. Aging reduces the functionality of anti-pneumococcal

antibodies and the killing of Streptococcus pneumoniae by neutrophil phagocytosis.

Vaccine 29:1929-34.

24. Bhalla M SS, Abamonte A, Herring SE, Roggensack SE, Bou Ghanem EN. 2020.

Extracellular adenosine signaling reverses the age-driven decline in the ability of

neutrophils to kill S. pneumoniae. doi:https://doi.org/10.1101/2020.04.14.041418.

25. Pignatti P, Ragnoli B, Radaeli A, Moscato G, Malerba M. 2011. Age-related

increase of airway neutrophils in older healthy nonsmoking subjects. Rejuvenation

Res 14:365-70.

26. Menter T, Giefing-Kroell C, Grubeck-Loebenstein B, Tzankov A. 2014.

Characterization of the inflammatory infiltrate in Streptococcus pneumoniae

pneumonia in young and elderly patients. Pathobiology 81:160-7.

27. Bhowmick R, Tin Maung NH, Hurley BP, Ghanem EB, Gronert K, McCormick BA,

Leong JM. 2013. Systemic disease during Streptococcus pneumoniae acute lung

infection requires 12-lipoxygenase-dependent inflammation. J Immunol 191:5115-

23.

28. Rudra Bhowmick NM, Bryan Hurley, Karsten Gronert, Beth McCormick and John

Leong. 2012. Systemic disease during Streprococcus pneumoniae lung infection is

facilitated by hepoxilin A3 mediated nutophil recruitment, Submitted, In revision.

29. Bakaletz LO. 2017. Viral-bacterial co-infections in the respiratory tract. Curr Opin
Microbiol 35:30-35.
30. McCullers JA. 2006. Insights into the interaction between influenza virus and
pneumococcus. Clin Microbiol Rev 19:571-82.
31. Chertow DS, Memoli MJ. 2013. Bacterial coinfection in influenza: a grand rounds
review. JAMA 309:275-82.
32. Centers for Disease C, Prevention. 2010. Estimates of deaths associated with
seasonal influenza --- United States, 1976-2007. MMWR Morb Mortal Wkly Rep
59:1057-62.
33. Shrestha S, Foxman B, Weinberger DM, Steiner C, Viboud C, Rohani P. 2013.
Identifying the interaction between influenza and pneumococcal pneumonia using
incidence data. Sci Transl Med 5:191ra84.
34. McCullers JA. 2014. The co-pathogenesis of influenza viruses with bacteria in the
lung. Nat Rev Microbiol 12:252-62.
35. Palacios G, Hornig M, Cisterna D, Savji N, Bussetti AV, Kapoor V, Hui J, Tokarz R,
Briese T, Baumeister E, Lipkin WI. 2009. Streptococcus pneumoniae coinfection is
correlated with the severity of H1N1 pandemic influenza. PLoS One 4:e8540.
36. Dhanoa A, Fang NC, Hassan SS, Kaniappan P, Rajasekaram G. 2011.
Epidemiology and clinical characteristics of hospitalized patients with pandemic
influenza A (H1N1) 2009 infections: the effects of bacterial coinfection. Virol J
8:501.
37. Siegel SJ, Roche AM, Weiser JN. 2014. Influenza promotes pneumococcal growth
during coinfection by providing host sialylated substrates as a nutrient source. Cell
Host Microbe 16:55-67.

38.   Rowe HM, Meliopoulos VA, Iverson A, Bomme P, Schultz-Cherry S, Rosch JW.

773   2019. Direct interactions with influenza promote bacterial adherence during

774   respiratory infections. Nat Microbiol 4:1328-1336.

775   39.   Diavatopoulos DA, Short KR, Price JT, Wilksch JJ, Brown LE, Briles DE, Strugnell

776   RA, Wijburg OL. 2010. Influenza A virus facilitates Streptococcus pneumoniae

777   transmission and disease. FASEB J 24:1789-98.

778   40.   Reddinger RM, Luke-Marshall NR, Sauberan SL, Hakansson AP, Campagnari AA.

779   2018. Streptococcus pneumoniae Modulates Staphylococcus aureus Biofilm

780   Dispersion and the Transition from Colonization to Invasive Disease. MBio 9.

781   41.   McCullers JA, Rehg JE. 2002. Lethal synergism between influenza virus and

782   Streptococcus pneumoniae: characterization of a mouse model and the role of

783   platelet-activating factor receptor. J Infect Dis 186:341-50.

784   42.   Levandowski RA, Gerrity TR, Garrard CS. 1985. Modifications of lung clearance

785   mechanisms by acute influenza A infection. J Lab Clin Med 106:428-32.

786   43.   Pittet LA, Hall-Stoodley L, Rutkowski MR, Harmsen AG. 2010. Influenza virus

787   infection decreases tracheal mucociliary velocity and clearance of Streptococcus

788   pneumoniae. Am J Respir Cell Mol Biol 42:450-60.

789   44.   McCullers JA, Bartmess KC. 2003. Role of neuraminidase in lethal synergism

790   between influenza virus and Streptococcus pneumoniae. J Infect Dis 187:1000-9.

791   45.   Smith AM, McCullers JA. 2014. Secondary bacterial infections in influenza virus

792   infection pathogenesis. Curr Top Microbiol Immunol 385:327-56.

793   46.   Cundell DR, Gerard NP, Gerard C, Idanpaan-Heikkila I, Tuomanen EI. 1995.

794   Streptococcus pneumoniae anchor to activated human cells by the receptor for

795   platelet-activating factor. Nature 377:435-8.

47.   Ballinger MN, Standiford TJ. 2010. Postinfluenza bacterial pneumonia: host

defenses gone awry. J Interferon Cytokine Res 30:643-52.

48.   Metzger DW, Sun K. 2013. Immune dysfunction and bacterial coinfections following

influenza. J Immunol 191:2047-52.

49.   Sun K, Metzger DW. 2008. Inhibition of pulmonary antibacterial defense by

interferon-gamma during recovery from influenza infection. Nat Med 14:558-64.

50.   Nakamura S, Davis KM, Weiser JN. 2011. Synergistic stimulation of type I

interferons during influenza virus coinfection promotes Streptococcus pneumoniae

colonization in mice. J Clin Invest 121:3657-65.

51.   Cho SJ, Plataki M, Mitzel D, Lowry G, Rooney K, Stout-Delgado H. 2018.

Decreased NLRP3 inflammasome expression in aged lung may contribute to

increased susceptibility to secondary Streptococcus pneumoniae infection. Exp

Gerontol 105:40-46.

52.   https://www.cdc.gov/flu/about/burden/index.html. 2018. Disease Burden of

Influenza https://www.cdc.gov/flu/about/burden/index.html, *on* CENTERS FOR

DISEASE CONTROL AND PREVENTION.

https://www.cdc.gov/flu/about/burden/index.html. Accessed

53.   Krone CL, Wyllie AL, van Beek J, Rots NY, Oja AE, Chu ML, Bruin JP, Bogaert D,

Sanders EA, Trzcinski K. 2015. Carriage of Streptococcus pneumoniae in aged

adults with influenza-like-illness. PLoS One 10:e0119875.

54.   Rello J, Pop-Vicas A. 2009. Clinical review: primary influenza viral pneumonia. Crit

Care 13:235.

55.   Torres A, Loeches IM, Sligl W, Lee N. 2020. Severe flu management: a point of

view. Intensive Care Med 46:153-162.

56.   Aaberge IS, Eng J, Lermark G, Lovik M. 1995. Virulence of Streptococcus

821   pneumoniae in mice: a standardized method for preparation and frozen storage of

822   the experimental bacterial inoculum. Microb Pathog 18:141-52.

823   57.   Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, Heidelberg J,

824   DeBoy RT, Haft DH, Dodson RJ, Durkin AS, Gwinn M, Kolonay JF, Nelson WC,

825   Peterson JD, Umayam LA, White O, Salzberg SL, Lewis MR, Radune D, Holtzapple

826   E, Khouri H, Wolf AM, Utterback TR, Hansen CL, McDonald LA, Feldblyum TV,

827   Angiuoli S, Dickinson T, Hickey EK, Holt IE, Loftus BJ, Yang F, Smith HO, Venter

828   JC, Dougherty BA, Morrison DA, Hollingshead SK, Fraser CM. 2001. Complete

829   genome sequence of a virulent isolate of Streptococcus pneumoniae. Science

830   293:498-506.

831   58.   Bou Ghanem EN, Maung NHT, Siwapornchai N, Goodwin AE, Clark S, Munoz-Elias

832   EJ, Camilli A, Gerstein RM, Leong JM. 2018. Nasopharyngeal Exposure to

833   Streptococcus pneumoniae Induces Extended Age-Dependent Protection against

834   Pulmonary Infection Mediated by Antibodies and CD138(+) Cells. J Immunol

835   200:3739-3751.

836   59.   Bhalla M, Hui Yeoh J, Lamneck C, Herring SE, Tchalla EYI, Heinzinger LR, Leong

837   JM, Bou Ghanem EN. 2020. A1 adenosine receptor signaling reduces

838   Streptococcus pneumoniae adherence to pulmonary epithelial cells by targeting

839   expression of platelet-activating factor receptor. Cell Microbiol 22:e13141.

840   60.   Boyd AR, Shivshankar P, Jiang S, Berton MT, Orihuela CJ. 2012. Age-related

841   defects in TLR2 signaling diminish the cytokine response by alveolar macrophages

842   during murine pneumococcal pneumonia. Exp Gerontol 47:507-18.

843   61.   Thevaranjan N, Puchta A, Schulz C, Naidoo A, Szamosi JC, Verschoor CP, Loukov

844   D, Schenck LP, Jury J, Foley KP, Schertzer JD, Larche MJ, Davidson DJ, Verdu

845   EF, Surette MG, Bowdish DME. 2018. Age-Associated Microbial Dysbiosis

846   Promotes Intestinal Permeability, Systemic Inflammation, and Macrophage

847   Dysfunction. Cell Host Microbe 23:570.

848   62.   McNamee LA, Harmsen AG. 2006. Both influenza-induced neutrophil dysfunction

849   and neutrophil-independent mechanisms contribute to increased susceptibility to a

850   secondary Streptococcus pneumoniae infection. Infect Immun 74:6707-21.

851   63.   Bou Ghanem EN, Lee JN, Joma BH, Meydani SN, Leong JM, Panda A. 2017. The

852   Alpha-Tocopherol Form of Vitamin E Boosts Elastase Activity of Human PMNs and

853   Their Ability to Kill Streptococcus pneumoniae. Front Cell Infect Microbiol 7:161.

854   64.   Wu W, Metcalf JP. 2020. The Role of Type I IFNs in Influenza: Antiviral

855   Superheroes or Immunopathogenic Villains? J Innate Immun 12:437-447.

856   65.   Wagenvoort GH, Sanders EA, Vlaminckx BJ, de Melker HE, van der Ende A, Knol

857   MJ. 2017. Sex differences in invasive pneumococcal disease and the impact of

858   pneumococcal conjugate vaccination in the Netherlands, 2004 to 2015. Euro

859   Surveill 22.

860   66.   Gutierrez F, Masia M, Mirete C, Soldan B, Rodriguez JC, Padilla S, Hernandez I,

861   Royo G, Martin-Hidalgo A. 2006. The influence of age and gender on the

862   population-based incidence of community-acquired pneumonia caused by different

863   microbial pathogens. J Infect 53:166-74.

864   67.   Greene CJ, Marks LR, Hu JC, Reddinger R, Mandell L, Roche-Hakansson H, King-

865   Lyons ND, Connell TD, Hakansson AP. 2016. Novel Strategy To Protect against

Influenza Virus-Induced Pneumococcal Disease without Interfering with Commensal

Colonization. Infect Immun 84:1693-1703.

68. Mina MJ, Klugman KP, McCullers JA. 2013. Live attenuated influenza vaccine, but

not pneumococcal conjugate vaccine, protects against increased density and

duration of pneumococcal carriage after influenza infection in pneumococcal

colonized mice. J Infect Dis 208:1281-5.

69. Wadowsky RM, Mietzner SM, Skoner DP, Doyle WJ, Fireman P. 1995. Effect of

experimental influenza A virus infection on isolation of Streptococcus pneumoniae

and other aerobic bacteria from the oropharynges of allergic and nonallergic adult

subjects. Infect Immun 63:1153-7.

70. Jochems SP, Marcon F, Carniel BF, Holloway M, Mitsi E, Smith E, Gritzfeld JF,

Solorzano C, Reine J, Pojar S, Nikolaou E, German EL, Hyder-Wright A, Hill H,

Hales C, de Steenhuijsen Piters WAA, Bogaert D, Adler H, Zaidi S, Connor V,

Gordon SB, Rylance J, Nakaya HI, Ferreira DM. 2018. Inflammation induced by

influenza virus impairs human innate immune control of pneumococcus. Nat

Immunol 19:1299-1308.

71. Stout-Delgado HW, Vaughan SE, Shirali AC, Jaramillo RJ, Harrod KS. 2012.

Impaired NLRP3 inflammasome function in elderly mice during influenza infection is

rescued by treatment with nigericin. J Immunol 188:2815-24.

72. Gay R, Han SN, Marko M, Belisle S, Bronson R, Meydani SN. 2004. The effect of

vitamin E on secondary bacterial infection after influenza infection in young and old

mice. Ann N Y Acad Sci 1031:418-21.

73.   Smith MW, Schmidt JE, Rehg JE, Orihuela CJ, McCullers JA. 2007. Induction of

pro- and anti-inflammatory molecules in a mouse model of pneumococcal

pneumonia after influenza. Comp Med 57:82-9.

74.   McNeela EA, Burke A, Neill DR, Baxter C, Fernandes VE, Ferreira D, Smeaton S,

El-Rachkidy R, McLoughlin RM, Mori A, Moran B, Fitzgerald KA, Tschopp J, Petrilli

V, Andrew PW, Kadioglu A, Lavelle EC. 2010. Pneumolysin activates the NLRP3

inflammasome and promotes proinflammatory cytokines independently of TLR4.

PLoS Pathog 6:e1001191.

75.   Pettigrew MM, Marks LR, Kong Y, Gent JF, Roche-Hakansson H, Hakansson AP.

2014. Dynamic changes in the Streptococcus pneumoniae transcriptome during

transition from biofilm formation to invasive disease upon influenza A virus infection.

Infect Immun 82:4607-19.

76.   Siwapornchai N, Lee JN, Tchalla EYI, Bhalla M, Yeoh JH, Roggensack SE, Leong

JM, Bou Ghanem EN. 2020. Extracellular adenosine enhances the ability of PMNs

to kill Streptococcus pneumoniae by inhibiting IL-10 production. J Leukoc Biol

doi:10.1002/JLB.4MA0120-115RR.

77.   Short KR, Reading PC, Wang N, Diavatopoulos DA, Wijburg OL. 2012. Increased

nasopharyngeal bacterial titers and local inflammation facilitate transmission of

Streptococcus pneumoniae. MBio 3.

78.   Kulkarni U, Zemans RL, Smith CA, Wood SC, Deng JC, Goldstein DR. 2019.

Excessive neutrophil levels in the lung underlie the age-associated increase in

influenza mortality. Mucosal Immunol 12:545-554.

79.   van der Sluijs KF, van Elden LJ, Nijhuis M, Schuurman R, Pater JM, Florquin S,

Goldman M, Jansen HM, Lutter R, van der Poll T. 2004. IL-10 is an important

mediator of the enhanced susceptibility to pneumococcal pneumonia after influenza

infection. J Immunol 172:7603-9.

80. Feng E, Balint E, Poznanski SM, Ashkar AA, Loeb M. 2021. Aging and Interferons:

Impacts on Inflammation and Viral Disease Outcomes. Cells 10.

81. Rowe HM, Livingston B, Margolis E, Davis A, Meliopoulos VA, Echlin H, Schultz-

Cherry S, Rosch JW. 2020. Respiratory Bacteria Stabilize and Promote Airborne

Transmission of Influenza A Virus. mSystems 5.

82. Nguyen DT, Louwen R, Elberse K, van Amerongen G, Yuksel S, Luijendijk A,

Osterhaus AD, Duprex WP, de Swart RL. 2015. Streptococcus pneumoniae

Enhances Human Respiratory Syncytial Virus Infection In Vitro and In Vivo. PLoS

One 10:e0127098.

83. Carniel BF, Marcon F, Rylance J, German EL, Zaidi S, Reine J, Negera E, Nikolaou

E, Pojar S, Solorzano C, Collins AM, Connor V, Bogaert D, Gordon SB, Nakaya HI,

Ferreira DM, Jochems SP, Mitsi E. 2021. Pneumococcal colonization impairs

mucosal immune responses to live attenuated influenza vaccine. JCI Insight 6.

84. Swamydas M, Lionakis MS. 2013. Isolation, purification and labeling of mouse bone

marrow neutrophils for functional studies and adoptive transfer experiments. J Vis

Exp doi:10.3791/50586:e50586.

85. Nishina K, Mikawa K, Takao Y, Shiga M, Maekawa N, Obara H. 1998. Intravenous

lidocaine attenuates acute lung injury induced by hydrochloric acid aspiration in

rabbits. Anesthesiology 88:1300-9.

FIGURE LEGENDS

Figure 1. Combined intranasal/intratracheal IAV inoculation of *S. pneumoniae*-colonized mice results in bacterial dissemination and disease. (A) Timeline of co-infection; 8-10 weeks old male C57BL/6 (B6) mice were inoculated i.n. with 5×10^6 CFU of biofilm grown S. *pneumoniae* TIGR4 to establish colonization in the nasopharynx. 48 hours later, the mice were either mock treated (*Sp*) or received 500 PFU of Influenza A virus PR8 (IAV) i.n. and 20 PFU i.t. (B) Percent weight loss was monitored daily. (C) Blinded clinical scoring was performed on day 2 and day 4 post IAV infection. The percentages denote the number of sick mice observed over total number of mice. A score of 0 means no sign of sickness observed and a score above 1 indicates observable sickness. (D) Survival was monitored for 8 days post IAV infection with fractions denoting survivors over total number of mice. (E) The bacterial burden in the nasopharynx and lung were determined at day 2 post IAV infection. Pooled data from four separate experiments are shown in which one group of twelve mice in each experimental condition were monitored over time and another group of 6 mice per experimental condition were used for measuring bacterial burden. Statistically significant differences determined by Student's t-test for bacterial burden and clinical score and the log-rank (Mantel-Cox) test for survival are indicated by asterisks. #, indicates statistical significance ($p < 0.05$) between *Sp* and *Sp*+IAV groups by Fisher's exact test.

Figure 2. PMN depletion may have a small effect on the course of disease during IAV/*S. pneumoniae* co-infection. 8-10 weeks old C57BL/6 mice were intraperitoneally (i.p.) injected with anti-Ly6G (clone 1A8) antibodies to deplete neutrophils or mock treated. The antibodies were given daily from day -3 to day 1, and every other day from day 3 to

the end of each experiment (with respect to IAV-infection). (A) At 18 and 48 hours post IAV

infection, bacterial numbers in the nasal wash and lungs were determined. Pooled data

from three separate experiments with a total of n=6 mice per experimental condition at 18h

and n=9 mice per condition at 48h are shown. Bacteremia (B) and weight loss (C) was

monitored over time. (D) Mice were blindly scored for symptoms of diseases at 18 and 48

hours post IAV-infection. The percentages indicate number of mice with clinical sickness

(clinical score above 1). (E) Survival was monitored over time where the fractions denote

survivors over total number of mice at 8 days post IAV infection. (C-E) Data are pooled

from three separate experiments with n=16 mice per group. Statistically significant

differences were determined by Student's t-test for bacterial burden and clinical score and

the log-rank (Mantel-Cox) and *ns* = not significant

Figure 3. Aging increases susceptibility to IAV/*S. pneumoniae* co-infection. Young

(8-10 weeks) and aged (18-24 months) C57BL/6 male mice were co-infected with *S.*

pneumoniae TIGR4 i.n. and Influenza A virus PR8 i.n. and i.t. (as in Figure 1A). (A) Clinical

score of co-infected young and aged mice at day 2 post IAV-infection is shown; the

percentage of mice with demonstrable illness is indicated. #, indicate statistical

significance by Fisher's exact test. (B) Survival of co-infected young and aged mice were

monitored over time with fractions denoting survivors over total of mice. Data are pooled

from four experiments with n=14 mice per group. Asterisks indicate statistical significance

by the log–rank (Mantel-Cox) test. (C-E) Bacterial burdens in the nasal wash, lungs, and

blood were determined at 18 and 48 hours post IAV inoculation. The mean +/- SEM pooled

from three separate experiments are shown with n=7 mice per condition at 18h and n=10

mice per condition at 48h. LOD denotes the limit of detection. For clinical score (A), data

shown are pooled from all mice from experiments shown in (B) and (C-E) that had survived

up to that timepoint (i.e., n= 13 for old and n= 16 for young).

Figure 4. Aging is associated with more rapid lung inflammation. Young and aged

C57BL/6 male mice were co-infected with S. *pneumoniae* and Influenza A virus PR8. 18

and 48 hours following IAV-infection (see experimental design in Fig. 1A), the lungs were

harvested. (A) Lungs were stained with Hematoxylin and Eosin and shown are

representative photographs at 100x or 400X (in inset). (B) Cytokines in the supernatants of

lung homogenates of young (n=4) or aged (n=5) mice at 18 hours post co-infection were

measured by multiplex ELISA. Asterisks represent statistical significance by Student's t-

test. (C) The percentages and total number of PMNs (Ly6G+) in the lungs was measured

by flow cytometry. Young mice are represented by open bars and aged mice by shaded

bars. The mean +/- SEM pooled from three separate experiments are shown where data

are pooled from twelve mice per age group, except for 48h post-co-infection, where due to

the kinetics of disease, data from only 7 surviving old mice are shown. Statistically

significant differences determined by Student's t-test are indicated by asterisks.

1000

Figure 5. Aging and IAV infection diminish the ability of PMNs to kill *S. pneumoniae*

ex vivo. PMNs were isolated from bone marrow of young (8-10 weeks) and aged (18-24

months) C57BL/6 male mice that were mock-infected (uninf.) or singly infected with IAV

(i.n. + i.t.) for 2 days. PMNs were incubated with *S. pneumoniae* pre-opsonized with

homologous sera from the same mouse for 45 or 90 minutes at 37°C. The percentages of

S. pneumoniae killed upon incubation with PMNs were determined with respect to a no

PMN control. Data shown represent the means +/- SEM pooled from two experiments (n=3

mice per group per timepoint) where each condition was tested in quadruplicates per

experiment. Asterisks represent statistical significance as determined by Student's t-test.

1010

Figure 6. Aging and prior colonization with *S. pneumoniae* result in impaired IFN-α

production and higher viral burden in the lungs. Young and aged C57BL/6 male mice

were either co-infected with S. *pneumoniae* and Influenza A virus PR8 (*Sp* + IAV),

challenged with virus alone (IAV) or mock challenged with PBS (uninfected) (as in Figure

1A). Two days post IAV-infection, (A) viral burdens in the lungs were determined and (B)

levels of IFN-α in the supernatants of lung homogenates were measured by ELISA. Data

from n=4 mice per group are shown. Statistically significant differences were determined

by Student's t-test.

A.

	Uninfected	18h	48h
Young	1	2	3
Aged	4	5	6

B.

IL-10

pg/mL (Log2)

ns

*, *p=0.0104*

Young Aged Young Aged

Uninfected   18h

IL-2

pg/mL (Log2)

ns

*, *p=0099*

Young Aged Young Aged

Uninfected   18h

IL-1β

pg/mL (Log2)

*, *p=0.0164*

Young Aged Young Aged

Uninfected   18h

TNFα

pg/mL (Log2)

ns

*, *p=0.0187*

Young Aged Young Aged

Uninfected   18h

C.

% PMN

% Ly6G+

*, *p=0.0003*

ns

Young

Aged

PMN Number

Ly6G+ 1x10⁵

*, *p=0.0006*

ns

Young

Aged

Hours post co-infection

A. Lung

B. Lung