Title:

Developmental constraints modulate reproductive fate and plasticity within the Arabidopsis ovule
primordium

Elvira Hernandez-Lagana[1]*, Gabriella Mosca[2]*, Ethel Mendocilla Sato[2]*, Nuno Pires[2], Anja Frey[2,4],
Alejandro Giraldo-Fonseca[2], Ueli Grossniklaus[2], Olivier Hamant[3], Christophe Godin[3], Arezki Boudaoud[3],
Daniel Grimanelli[1], Daphné Autran[1#], Célia Baroux[2#]

Affiliations :

1 Diversité, Adaptation et Développement des Plantes (DIADE), IRD, CIRAD, University of Montpellier,
Montpellier, France.

2 Department of Plant and Microbial Biology and Zurich-Basel Plant Science Center, University of Zurich,
Zurich, Switzerland.

3 Laboratoire Reproduction et Développement des Plantes, University of Lyon, ENS Lyon, UCB Lyon 1,
CNRS, INRAE, INRIA, Lyon, France.

4 Present address: Novartis Pharma Schweiz AG, Basel, Switzerland.

* equal contribution.

authors for correspondence: daphne.autran@ird.fr, cbaroux@botinst.uzh.ch

The authors declare no competing financial interests.

ORCID numbers

Name	Orcid #
Elvira Hernandez-Lagana	0000-0002-2645-3783
Gabriella Mosca	0000-0003-4509-498X
Ethel Mendocilla-Sato	0000-0003-0339-3535
Alejandro Giraldo-	0000-0003-2943-9810
Nuno Pires	0000-0002-7113-3519
Anja Frey	
Ueli Grossniklaus	0000-0002-0522-8974
Olivier Hamant	0000-0001-6906-6620
Arezki Boudaoud	0000-0002-2780-4717
Christophe Godin	0000-0002-1202-8460
Daniel Grimanelli	0000-0001-5424-114X
Daphné Autran	0000-0002-5227-8966
Célia Baroux	0000-0001-6307-2229

Summary

Sexual reproduction in multicellular organisms requires the separation of the germline from the soma. In flowering plants, the formation of spore mother cells (SMCs) marks the somatic-to-reproductive cell fate transition in floral organs. The female SMC differentiates at the apex of the ovule primordium, as a single, large, and elongated subepidermal cell. However, this singleness rule is not very robust: several SMC candidates develop in some *Arabidopsis* accessions and mutants. Yet, the basis of this plasticity is not understood. Here we investigate whether ovule primordium growth and overarching cellular architecture serve as an instructive cue to channel the formation of a single SMC. Combining quantitative analyses of 3D cell-based information, cell cycle reporters, and 2D growth simulations, we provide a comprehensive analysis of cell proliferation and growth events that shape ovule primordium development and SMC formation. This reveals that SMC characteristics emerge earlier than previously thought, and often for more than one cell. Thus, developmental constraints gradually canalize the fate of these cells, so that a single SMC enters meiosis ultimately. Among these constraints, we explore the contribution of dynamic cell division patterns and localized cell growth anisotropy. Consistent with 2D computational simulations, the reduction of anisotropy in *katanin* mutants leads to altered ovule shape and cell division patterns. Such disturbance of ovule morphology is sufficient to delay SMC canalization, as multiple cells express reproductive fate markers in *katanin* mutant ovules. In conclusion, we show that ovule growth and shape channel SMC fate for a single cell.

Highlights

1. SMC singleness emerges from a canalization process mediated by KATANIN.

2. Anisotropic cell growth and local, biphasic cell divisions patterns explain ovule primordium shape.

3. 2D-mechanical growth models relate ovule growth and SMC emergence.

4. A 3D image resource of developing ovule primordia at cellular resolution.

Introduction

A hallmark of sexual reproduction in multicellular organisms is the separation of the germline from the soma. In animals, primordial germ cells (PGCs) are set-aside during embryogenesis from a mass of pluripotent cells. The number of germ cells depends on the balance between proliferation (self-renewal) and differentiation, a process controlled by both intrinsic factors and the surrounding somatic tissue (Extavour and Akam, 2003; Kimble 2011). In flowering plants, by contrast, the spore mother cells (SMCs), representing the germline precursors, differentiate late in development. SMCs arise multiple times, i.e., in each flower during the formation of reproductive organs. In *Arabidopsis*, the female SMC differentiates in the nucellus of the ovule primordium, a digit-shaped organ outgrowing from the placental tissue of the gynoecium. The SMC is recognizable as a single, large, and elongated subepidermal cell, in central position, at the nucellus, and displaying a prominent nucleus and nucleolus (Bowman, 1993, Schneitz *et al.,* 1995, Bajon *et al.*, 1999, Schmidt *et al*., 2015).

Although SMC singleness may appear to be robust, more than one SMC candidate per primordium can develop at variable frequencies depending on the *Arabidopsis* accession (9,6% in *Columbia* (*Col-0*), 27% in *Monterrosso* (*Mr-0)*, Rodriguez-Leal *et al.,* 2015). Mutants in *Arabidopsis*, maize, and rice that compromise SMC singleness unveiled the role of regulatory pathways involving intercellular signaling, small RNAs, as well as DNA and histone methylation (Sheridan *et al.*, 1996; Sheridan *et al.*, 1999; Nonomura *et al.*, 2003; Zhao *et al.*, 2008; García-Aguilar *et al.,* 2010; Olmedo-Monfil *et al.*, 2010; Schmidt *et al.,* 2011; Su *et al.*, 2017; Su *et al.,* 2020). As the SMC forms, cell-cycle regulation contributes to the stabilization of its fate in a cell-autonomous manner through cyclin-dependent kinase (CDK) inhibitors and RETINOBLASTOMA-RELATED1 (RBR1) (Zhao *et al.,* 2017; Cao *et al.*, 2018). SMC singleness thus appears to result from a two-step control: first, restricting differentiation to only one cell and second, preventing self-renewal before meiosis (reviewed in Lora *et al.*, 2019, Pinto *et al.*, 2019).

However, the precise mechanisms underlying the plasticity in the number of SMC candidates and SMC fate specification are still poorly understood. In principle, SMC singleness may be controlled by successive molecular cues. However, even in that scenario, such cues must be positional, at least to some extent, and thus involve a spatial component. In the last decade, many different molecular cues defining spatial patterns in the ovule primordium were identified (Pinto *et al.*, 2019; Su *et al.,* 2020), however their coordination is unknown. Since SMCs emerge at the primordium apex concomitant with its elongation, we hypothesize that developmental constraints during ovule morphogenesis influence

SMC singleness and differentiation. Such a scenario could explain variation in the number of SMC

candidates, ultimately culminating in a single SMC entering meiosis, a process that was previously

discussed as a possible canalization (Rodriguez-Leal *et al.*, 2015). Canalization refers to the capacity of an

organism to follow a given developmental trajectory in spite of disturbances (Waddington, 1942;

Scharloo, 1991, Hallgrimsson *et al.*, 2019). Yet, canalization as proposed by Waddington embraces a

broader meaning, including the nature of the developmental process itself to explain phenotypic

variability (Waddington, 1942). Answering the questions of whether SMC formation follows a

stereotypical or plastic developmental process and whether it is intrinsically linked to or independent of

ovule primordium growth would unravel fundamental principles connecting cell fate establishment and

organ growth.

Such an analysis requires a high-resolution description of the ovule tissue topology and geometry in

time. Our current knowledge of ovule primordium growth in *Arabidopsis* is based on two-dimensional

(2D) micrographs from tissue sections or clearings. It is described in discrete developmental stages

capturing classes of primordia by their global shape and SMC appearance until meiosis and by the

presence of integument layers and ovule curvature later-on (Grossniklaus and Schneitz, 1998). In

addition, a 3D analysis of average cell volumes during primordium growth was recently provided (Lora *et*

al., 2017). Yet, we lack a view of the patterning processes shaping the early ovule primordium, the

relation of cell layers and proliferation domains in 3D, and a temporally resolved description of

primordium emergence. To address the question of SMC singleness, we thus developed a resource

where the growth of the ovule primordium is described and quantified at cellular resolution in 3D. We

combined 3D imaging, quantitative analysis of cell and tissue characteristics, reporter gene analyses, 2D

mechanical growth simulations, and genetic analyses. Using the *katanin* mutant affecting anisotropic cell

growth and cell division patterns (Luptovciak *et al.*, 2017a, Ovecka *et al.*, 2020), we show that an altered

ovule morphology is sufficient to disrupt SMC singleness. We also uncovered that differentiation of SMC

candidates initiates earlier than previously thought. Finally, we provide evidence for a developmental

canalization process leading to the specification of a single SMC prior to meiosis.

Results

Building a reference collection of 3D images capturing developing ovule primordia in 3D at cellular resolution

To generate 3D reconstructions of ovule primordia with cellular resolution at consecutive different stages we imaged primordia stained for cell boundaries in intact carpels by confocal microscopy. Carpels were cleared and stained using a modified PS-PI staining (Truernit *et al.*, 2008) and mounting procedure preserving their integrity (Mendocilla-Sato and Baroux, 2017; Figure 1A). We selected high signal-to-noise ratio images and segmented them based on cell boundary signals using a previously described pipeline (Mendocilla-Sato and Baroux, 2017; Figure 1B). We manually curated 92 ovules representing seven consecutive developmental stages (7-21 ovules per stage, Figure 1B, Table 1, Data S1), classified according to an extended nomenclature explained in the STAR Methods. In order to gain temporal resolution, we defined new stages (st0-I to st0-III) covering primordium emergence, which occurs prior to the straight digit-shape organ stage 1-I where the SMC becomes distinguishable by its apparent larger size in longitudinal views (Schneitz and Grossniklaus, 1998).

To evaluate the distinct contribution of domain-, layer- and cell-specific growth dynamics, we assigned a label to each segmented cell enabling data classification (Figure 1B, STAR Methods). We defined cell labels to distinguish *domains*, *layers* and *cell types*: the apical and basal domains, the L1 and L2,L3 layers (L1, epidermis; and the underlying L2 and L3 layers taken together), and seven cell types intersecting with the domains and layers. The latter include the self-explanatory cell labels "L1 apical", "L1 basal", "L2,L3 apical", "L2,L3 basal", "SMC", and specific labels for the "L1 dome", the upmost apical L1 cells in contact with the SMC, and the "CC" or companion cells, the elongated L2 cells adjacent to the SMC.

To quantitatively analyze the data with respect to cell labels, layers, domains, ovule stage and genotype, we developed an interactive, R-based interface named OvuleViz. The script reads cell descriptors exported from segmented image files and enables multiple plot generation from a user-based selection of (sub)datasets (Figure S1A-C, STAR Methods). Collectively, our work generated a reference collection of annotated, 3D images capturing ovule primordium development at cellular resolution from emergence until the onset of meiosis. The collection of 92 segmented images associated with a total of 7763 annotated cells and five morphological cell descriptors (volume, area, sphericity, prolate and oblate ellipticity) provides a unique resource for morphodynamic analyses of ovule primordium growth.

To identify correlations between growth patterns and differentiation, we performed a principle
component analysis (PCA) based on the aforementioned cell descriptors, *per* cell type and stage,
considered together or separately (Figure S1D-E). In this global analysis, the SMC appears distinct at late
stages (2-I and 2-II) but clear conclusions are hampered by a crowding effect that is not resolved by this
approach. This prompted us to investigate in detail the contribution of different layers, domains, and
cell types to ovule primordium growth and in relation to SMC differentiation.

Ovule primordium morphogenesis involves domain-specific cell proliferation and anisotropic cell
shape patterns

The ovule primordium emerges from the placenta as a small dome-shaped protrusion and grows into a
digit-shaped primordium with nearly cylindrical symmetry (stage 1-I) before enlarging at the base
(Figure 1B). Using our segmented images, we first quantified global changes in cell number, cell volume ,
and ovule primordium shape. Our analysis revealed two distinct phases of morphological events. Phase I
(stages 0-I to 0-III) is characterized by a 4.5-fold increase in total cell number but a moderate increase in
mean cell volume (10%, P=0.03). By contrast, Phase II (stages 1-I to 2-II) is characterized by a moderate
increase in cell number (50%) and the global mean cell volume is relatively constant (Figure 2A, S2A). To
quantify the resulting changes in organ shape, we extrapolated a continuous surface mesh of the ovule
outline and used it to compute its height and width at the base (Figure 2B-C, S2B, STAR Methods).
Anisotropic organ growth during Phase I was confirmed by a steady increase in height while primordium
width increased moderately (Figure 2C). This contrast in morphodynamic events between Phase I and II
is illustrated by the representation of fold-changes (FCs) in cell number and aspect ratio (Figure 2D)
which range between 1.5 and 2.0 in Phase I whereas drop to 1.4 and 1.2 in Phase II. These observations
confirmed that Phase I captures a distinct growth dynamic compared to Phase II.

Next, to capture possible specific patterns of growth, we analyzed cell number, cell size, and cell
shape using different viewpoints: one comparing the L1 and L2,L3 layers and one contrasting the apical
vs basal domains. Scoring cell number per viewpoint clearly showed a dominant contribution of the
epidermis (L1) relative to the subepidermal layers and of the basal relative to the apical domain (Figure
2E,F). To verify these findings with a cellular marker, we analyzed the M-phase-specific *CYCB1.1-dbGFP*
reporter (Ubeda-Tomas et al, 2009). We scored the number of GFP-expressing cells among 481 ovules
and plotted relative mitotic frequencies *per* cell layer and domain for each ovule stage to generate a
cell-based mitotic activity map (Figure 2G-H, Data S2, STAR Methods). In this approach, subepidermal

cells beneath the dome were distinguished from underlying L3 cells to gain resolution on the L2 apical

domain where the SMC differentiate. Consistent with our previous observation, in Phase I, a high

proliferation activity was scored in L1 cells at the primordium apex (scoring 64% of mitotic events). By

contrast, the L2 apical domain remains relatively quiescent (contributing only 3% of the mitotic events).

During Phase II, proliferation preferentially occurs in the basal domain with > 60% of mitotic events at all

stages. It is noteworthy that during this phase, L2 apical cells divide relatively rarely, with the exception

of SMC neighbors cells that show frequent divisions at stage 1-II. Thus, reporter analysis confirmed a

biphasic, temporal pattern of cell division with changing regional contributions, suggesting the L1 apical

and L2,L3 basal domains as consecutive sites of increased proliferation, contributing to morphological

changes in Phase I and II, respectively.

Cell size analysis, by contrast, did not reveal significant changes during primordium development

with the notable exception of the SMC (Figure 2A, S2C). The SMC gradually attains a distinct size starting

at stage 0-III when compared to other L2,L3 cells (Figure 2I) or even earlier, at stage 0-I, when compared

to all other cells (Figure 2I inset). Size differentiation of the SMC is not uniform among ovules,

demonstrating plasticity in the process (Figure S2D). In addition, different cell sizes between L1 and

L2,L3 layers are already discernible at primordium emergence (Figure 2J, S2E). The presence of larger

L2,L3 cells is consistent with the low division frequency observed previously - likely stemming from

longer growth phases. We propose that cell size asymmetry established at early stages favors SMC

emergence.

We then investigated cell shape changes during primordium elongation, using ellipticity and

sphericity indices computed by Imaris (Bitplane, Switzerland) following segmentation. The analysis did

not reveal significant differences between domains or layers (Figure S2F, STAR Methods). This could

indicate either a highly variable cell shape or more localized, cell type-specific shape changes. We thus

more specifically analyzed the subepidermal niche where the SMC differentiates. We found that

companion cells (CC) showed an increasing ellipticity (and decreasing sphericity), starting at stage 1-I

and culminating at stage 2-I (Figure S2F). By contrast, the SMC only showed a moderate decrease in

sphericity at late stages and no distinctive ellipticity at early stages (Figure S2F). In order to get raw

information on anisotropy of cells shapes, we developed an extension for MorphoMechanX

(www.morphomechanx.org) to (i) perform a semi-automatic labeling of cell layers and cell types from a

cellularized mesh and (ii) compute in each 3D cell the principal axes of shape anisotropy and the

corresponding indexes (Figure S2G, STAR Methods, Movie S1). The maximum anisotropy shape index of

the SMCs was, averaged *per* stage, consistently above the medium (and hence above the minimum)

anisotropy index (Figure 2K). This confirmed that the SMC has an anisotropic shape, with a
distinguishable major axis throughout ovule primordium development from early stages onwards
(Figure S2H). We measured the degree of alignment of the SMC major axis with the main growth axis of
the ovule during early stages (stage 0-II) and found a mean angle of 22° (+/- 11°, n=16) (Figure 2K, see
also Figure 2C showing main growth axis of the ovule along development). Thus, the SMC's anisotropic
shape emerges at the earliest stages and its orientation is aligned with the growth axis of the
primordium.
Taken together, these results suggest that anisotropic primordium growth is linked to domain-
specific cell proliferation combined with localized, anisotropic expansion.
213

214 2D mechanical simulations of ovule primordium development relate its coordinated growth to SMC
215 shape emergence

Organ shape is determined by the rate and direction of cell growth, the latter affected by signaling and
the mechanical state of the tissue and its geometry, with possible feedback interactions between them
(Basel *et al.*, 2014, Echevin *et al.*, 2019). Mechanical constraints arise from the growth process in form of
tensile and compressive forces that, in turn, influence cell and tissue growth depending on their
elastic/visco-plastic properties (Echevin *et al.*, 2019). To determine the role of ovule primordium growth
on SMC differentiation, which involves early differential cell growth, we sought to understand the
contributions of growth rate and anisotropy and their relation to signaling and mechanical constraints
(Coen *et al.*, 2004; Kenaway *et al.*, 2011). Resolving this question would define fundamental
developmental principles linking organ growth and SMC differentiation.
We first sought to determine morphogenetic principles of primordium growth at the tissue level by
developing a 2D Finite Element Method (FEM)-based mechanical model, which considered two
uncoupled but complementary modes of growth: (i) signal-based growth in which an abstract
morphogen captures the cumulative effects of biochemical signals without considering its explicit mode
of action (i.e., cell wall loosening, increase in turgor pressure) (Boudon *et al.*, 2015); and (ii) passive
strain-based growth in which relaxation accommodates excess of strain (and therefore also excess of
stress) that accumulated in the tissue by relaxing its reference configuration according to the amount of
strain. The mechanical equilibrium is computed to ensure compatibility with the tissue that is locally
growing at different rates and orientations. This introduces residual internal compressions and tensions
to which the action of turgor pressure is also added (Rodriguez *et al.,* 1994). A polarization field is used
to set the direction of anisotropic growth for both growth mechanisms (Coen *et al.,* 2004). The

simulation consists of iterations where the mechanical equilibrium is computed before each growth step

is specified by signal- and strain-based growth. This strategy extends previous tissue growth models

(Kuchen *et al.*, 2012, Borzog *et al.*, 2016, Boudon *et al.*, 2015, Mosca *et al.,* 2018, STAR Methods).

We designed a starting template consisting of an L1 layer distinct from the underlying L2, L3

tissue based on different growth and material properties (Jarrousse, 2014) (Table 2, Figure S3, STAR

Methods). We first set the model components to produce a realistic elongated, digit-shaped primordium

with a narrow dome and an L1 layer of stable thickness during development (Reference Model, FEM-

Model 1) (Figure 3A, S3A). This model combines the following hypotheses: a narrow domain of

anisotropic, signal-based growth, a broad domain competent for passive strain-based relaxation and

material anisotropy.

Then, using the versatility of the modeling framework to vary initial conditions, we asked

whether primordium morphogenesis is driven primarily by growth of the epidermis (L1) or if it requires

active growth of inner tissues. In the first scenario, L1 growth is responsible for creating a dome shape,

enabling underlying tissue growth by pulling on the inner layers and inducing cell divisions as observed

in Phase I. In another scenario, active growth of the inner tissues forces the L1 to grow and create a

protrusion. We ran several simulations with varying initial distribution patterns of the growth signal

(Table 2). The shape aberrations produced by FEM-Models 4, 5, 6 and 7 (Figure S3B) suggested that an

inner tissue-driven growth principle (*vs* L1-driven), involving a narrow strip of initial growth signal

diffusing in the inner layers, was more plausible (Reference Model, FEM-Model 1, Figure 3A, S3A). Next,

through modulation of passive strain-based growth, we determined that a broad tissue domain

uniformly competent for strain accommodation is necessary to resolve the high accumulation of strain,

limiting primordium elongation (FEM-Model 8, Figure S3C). Yet, material anisotropy influencing the

relaxation response is not necessarily required broadly and it is sufficient to specify it in the L1 only to

achieve a dome with a realistic shape and constant L1 thickness (FEM-Model 2, Figure S3C). Finally, we

asked whether growth anisotropy must be specified in the model or if geometrical and mechanical

constraints are sufficient to specify primordium shape. Clearly, removing this component abolished the

digit shape of the primordium and produced a hemi-spherical protrusion (FEM-Model 3, Figure 3B).

In summary, we identified parsimonious growth principles shaping the ovule primordium: an

inner tissue-driven process promoted by a narrow domain of anisotropic, signal-based growth is

complemented by passive tissue growth in response to accumulated strain. Furthermore, material

anisotropy in the L1 is predicted to play a role in constraining L1 thickness as observed experimentally.

Next, we aimed to analyze how cell division and cell shape, particularly that of the SMC, are

connected to tissue growth patterns at cellular scale. For this, we developed a complementary model

with cellular resolution. This vertex-based model features cells as 2D polygons whose walls are mass

spring (MS) segments (Smith, 2009; Amar, 2009). This model integrates mechanics (Hooke's law for MSs)

with signal- and strain-based growth as before (STAR Methods). We simulated growth starting with a

virtual grid of cells representing a flat placenta and focused on early stages of ovule primordium

formation until stage 1-I. We tested the same combination of growth hypotheses as for the continuous

model (Table 2), except for material anisotropy that cannot easily be embedded in a vertex-based

model. As we observed that L1 cells do not divide periclinally, nor change in volume on average, we

deduced that their growth occurs mainly in direction of the main axis of the primordium. For this reason,

we enforced this feature in our model, preventing growth of the anticlinal walls (both for signal-based

growth and passive strain-based growth, STAR Methods). This constraint limits the ways an organ can

grow in a digit-shaped fashion, as the L1 does not progressively populate the inner tissues by division.

Furthermore, our models included assumes a slower division rate for L2,L3 apical cells, consistent with

our previous observations (Figure S3D).

We identified MS-Model 2 to best capture the characteristic shape and aspect ratio of ovules

(Figure 3C, S3D). This reference model confirmed the growth hypotheses from FEM-based simulations

regarding the distribution and orientation of signal-based growth (Figure S3B) and passive strain-based

growth (Figure S3C).

Finally, we were interested to determine whether SMC morphology is an emerging property of

primordium growth and how anisotropic growth influences the process. Remarkably, in the cell-based

reference model including anisotropic cell growth, an SMC candidate emerged early: a central L2 apical

cell that, beside its larger size, adopted a trapezoidal shape and a principal anisotropy axis aligned with

the main direction of ovule growth (Figure 3E). Such features are qualitatively comparable what we

observed for stage 0-II primordia (Figure 3E, S3D, Figure 2K).

The MS-Model 3 corresponds to a virtual mutant with isotropic cell growth. Under these

conditions, ovule primordia displayed an altered shape with a wider surface and flatter dome as in the

FEM-Model 3 (Figure 3D). Interestingly, a larger, central L2 cell still emerged but the elongated shape

typical of SMCs arose later in the simulation progression as compared to the reference model (Figure 3F,

Movie S3). Thus, 2D simulations predict that anisotropic cell growth does not only affect ovule

primordium shape but also the timing of SMC candidate emergence, hence connecting the growth

process with SMC differentiation. In these models, the candidate SMC eventually divided as they missed

a causative rule to differentially regulate cell division. When we prevented the SMC candidate to divide,

its enlargement overrode primordium shape control in our simulations, indicating an additional level of

control in real primordia (Figure S3E).

Altogether, the two complementary growth models of ovule primordium development confirm

at different scales a role for regional cell growth compatible with previous observations. They predict

that tissue accommodation, anisotropy in mechanical properties, and a specific pattern of signal-based

growth collectively influence the directional growth of the primordium. We showed that altered ovule

primordium growth affects size and shape of the SMC candidate, linking organ architecture to SMC fate.

Reciprocally, unleashing SMC growth leads to deformation of the primordium, suggesting the existence

of a compensatory feedback mechanism that coordinates cell and organ growth.

katanin **mutants show a distinct ovule primordium architecture primarily through altered cell division**
patterns

To test the model predictions experimentally, we analyzed the consequences of cell growth anisotropy

perturbation on ovule primordium growth and SMC fate establishment. *Arabidopsis* plants carrying a

defective microtubule-severing protein KATANIN exhibit organ shape defects, including increased round-

and bumpiness, as a consequence of disorganized cortical microtubules and isotropic cellulose

microfibril orientations (Bichet et al 2001, Burk et al 2002). We thus analyzed the shape and cellular

organization of ovule primordia in *katanin* mutants using the *botero* (*bot1-7*) (*Ws* background), *lue1*, and

mad5 alleles (*Col* background) (Bichet *et al.*, 2001; Bouquin *et al.*, 2003; Brodensen *et al.*, 2008),

collectively referred to *katanin* (*kat*) mutants hereafter. We generated and analyzed a new dataset of 59

annotated 3D digital *kat* and corresponding wild-type ovule primordia at stages 0-III, 1-I, and 1-II (Data

S4a). *kat* mutant primordia have an increased size and a more isotropic shape (Figure 4A), being 1.5

times bigger in volume than wild-type primordia (*P*=0.007, stage 0-III, Figure 4B) with a smaller aspect

ratio (*P*<0.01 stages 1-I, 1-II, Figure 4C). The width-to-height aspect ratio does not inform on the shape

at the flanks, hence does not discriminate straight digit *versus* plump organ shapes. We estimated the

"plumpiness" of the primordia by calculating the volume fraction of a fitting, 3D parallelepiped

(bounding box) effectively occupied by the primordium (bounding box occupancy, Figure 4D, S4A). In *kat*

ovules, the relationship deviates from the wild type beginning at stage 1-I, when the primodium

normally starts elongating along the major growth axis. These measurements confirmed a marked

attenuation of anisotropic growth in *kat* primordia as was observed in roots, shoot organs, or seeds

(Bichet *et al.*, 2001; Uyttewaal *et al.,* 2012, Hervieux *et al.*, 2016; Ren *et al.*, 2017; Wigthman *et al.*, 2013;

Zhang et al 2013; Luptovciak et al 2017b). Yet, at a global level, the mean cell number, cell volume, and

sphericity are not significantly different between *kat* and wild-type primordia (Figure S4B). Thus, these

global approaches did not resolve the origin of primordia mis-shaping.

To refine the analysis, we contrasted different layers as done for wild-type primordia and found

local alterations that provide an explanation for the formation of broader and more isotropic primordia

in *kat* mutants. Notably, the epidermis of *kat* primordia contains 17.5% more cells (47 L1 cells in *kat* vs

40 in wild-type at stage 0-III, *P*=0.03). Also, *kat* epidermal cells have a 20% larger volume than in the wild

type (*P*<0.001). Early in development (stage 0-III), the apical domain of *kat* primordia is populated by

larger (17%, *P*<0.001), more spherical (2%, *P*<0.001) cells than in wild type (Figure 4E,SC), a

characteristic that extends to the basal domain at later stages (Figure S4D). To determine whether, in

kat primordia, the mitotic domains shifted in space and/or time, we scored *CYCB1.1-dbGFP* signals *per*

stage, layers, and equivalent domains as before, and expressed it as fold-changes relative to wild-type

primordia (Figure 4F, S4E). In Phase I (stage 0-II), we consistently scored 2.4 times less mitoses in the L1

dome domain, and 2.7 times more mitoses at the L1 basal domain of *kat* primordia. Hence, in *kat*

primordia, proliferation is mostly localized in the basal region and not higher in the early L1 dome as in

the wild type.

To elucidate whether these alterations affected the cellular domain where the SMC normally

differentiates, we interrogated L2 apical cells and specifically the L2 neighbors of the candidate SMC. At

stage 0-II, we scored 3.5 times more divisions in the L2 SMC neighbor cells in the apical domain, showing

that the relative mitotic quiescence is lost in this domain in *kat* primordia. Next, we focus on cells in

contact with the SMC (STAR methods). We found that in *kat* primordia, in contrast to wild-type and to

non-neighbor cells, SMC neighbors specifically enlarge as early as stage 0-III (Figure 4G, S4E) but without

a detectable increase in sphericity. Consistently, our *CYCB1.1-dbGFP* analysis showed that the mitotic

activity detected in SMC neighbors at stage 1-II in wild-type primordia significantly decreases in *ka*t

primordia (Figure 4F, S4E).

In conclusion, the absence of *KAT*-mediated cell growth anisotropy is associated with major

changes in the domains of cell division, which can be linked to geometric changes or to a slow-down of

mitosis as previously shown in *kat* mutants (Ovecka *et al.*, 2020; Komis *et al.*, 2017). These alterations

impact ovule primordium shape but also architecture of the subepidermal apical domain, where the

SMC differentiates.

Altered ovule primordium architecture in *katanin* mutants induces ectopic SMC fate

We next investigated whether *kat* ovule growth defects also affect cell identity and SMC singleness. In *kat* primordia, SMCs appear bigger in volume and more spherical than their wild-type counterparts at stage 0-III; yet, no significant differences were detected at stages 1-I and 1-II. However, we observed variability in volume at these stages (Figure 4G, S5A-B). When we analyzed the number of SMC neighbors (Figure 5A-B), we found seven or more in *kat* primordia compared to an average of five in the wild type. The *kat* SMC neighbors were 14% larger (P=0.006), 11% less ellipsoid (P=0.03), and 2% more spherical (P<0.01) than in the wild type (Figure 5B), reminiscent of mutants affecting SMC singleness (Pinto *et al.*, 2019).

To analyze the identity of these additional, enlarged cells, we quantified several SMC hallmarks. First, we found that ~ 35% *kat* primordia harbor more than one SMC candidate at stage 1-II, decreasing to 14.42% at stage 2-II (Figure 5C). Then, we assessed two complementary markers associated with SMC fate in *kat* primordia. The first marker is a GFP-tagged linker histone variant (H1.1-GFP) that marks the somatic-to-reproductive fate transition by its eviction in the SMC at stage 1-I (She *et al.*, 2013): H1.1-GFP eviction occurred in more than one cell in 28.57% *kat* primordia (n=43, Figure 5D). The second marker reports expression of the *KNUCKLES* transcription factor (KNU-YFP) expressed in the SMC (Tucker *et al.*, 2012). Detectable as early as stage 1-I in the wild type, it was ectopically expressed in 21% *kat* primordia at stage 1-II (n=43) (Figure 5E, S5B). To test whether the ectopic SMC candidates have a meiotic competence, we analyzed the AtDMC1-GUS reporter (Agashe *et al.*, 2002, Klimyuk and Jones, 1997), and indeed scored ectopic expression in 57% of *kat* primordia (n=56) (Figure 5F). We also analyzed the pWOX2-CenH3-GFP marker labeling centromeres starting at the functional megaspore stage (De Storme *et al.*, 2016) and found 20.7% *kat* primordia (n=29) with two labeled cells, as compared to 3,8% (n=26) in the wild type (Figure S5C).

Taken together, these results show that ectopic, abnormally enlarged SMC neighbors in *kat* primordia show at least some characteristics of SMC and/or functional megaspores and can thus adopt a reproductive fate.

The growth of a single SMC emerges from a canalization process

Based on our analysis of *kat* primordia, it appeared that the frequency of ectopic SMC candidates was high at early stages but decreased over time. This is reminiscent of the phenotypic plasticity in SMC differentiation observed, to a lesser degree, in different *Arabidopsis* accessions (Rodríguez-Leal *et al*,

2015). To characterize this plasticity during development we analyzed 1276 wild-type primordia from
stage 0-II to 2-II and scored the number of primordia with one or two enlarged, centrally positioned,
subepidermal cells (class A and B, respectively, Figure 6A-C). In the wild type, the majority of primordia
showed a single candidate SMC (class A) at the stage 0-II, but 27% primordia (n=289) had two SMC
candidates (class B), this frequency decreasing to 3% at stage 2-II (n=103). This finding is consistent with
~ 5% primordia at stage 1-II showing H1.1-GFP eviction (n=43) and KNU-YFP expression (n=88) in more
than one cell, respectively (Figure 5D-E).

Next, we quantified class A and B primordia in *kat* mutants by scoring 2587 ovules of three
alleles (Figure 6C, S6A). Clearly, the canalization process is delayed in *kat* mutant primordia with 33%
class B primordia at stage 1-II (n=113) compared to 12% in the wild type (n=281). Consistently, the two
previously used SMC markers are observed in ectopic cells in *kat* primordia already at stage 1-I (Figure
S6B-C), and more significantly at stage 1-II (28.5% ectopic eviction of H1.1-GFP and 21% ectopic
expression of KNU-YFP, Figure 5D-E). At stage 2-II, when 17% of cleared *kat* primordia (n=110) showed
ectopic SMCs, H1.1-GFP eviction and KNU-YFP expression was only found in 8% (n=25) and 6.7% (n=74)
of the primordia, respectively (Figure S5B-C). Therefore, developmental canalization is not only delayed
in *kat* ovules, but there is also a partial uncoupling of cellular morphology and marker expression.

Another outcome of this study is the early emergence of SMCs, confirming our former 3D cell-
based analysis that showed heterogeneity in cell volume of the L2,L3 apical cells already at stages 0-I/0-
II (Figure 2J). To corroborate this finding using a different criterion, we measured the nuclear area,
which is a distinctive feature of SMCs (She *et al.,* 2013; Rodriguez-Leal *et al.,* 2015). We compared the
nuclear area of SMC candidate to that of surrounding L1 and L2 cells in both class A and B primordia at
stages 0-II and 0-III. Wild-type class A and B primordia showed an enlarged nucleus in the candidate
SMCs from stage 0-II onwards. This correlation was also true in *kat* primordia, where SMC candidates
also showed a 1.37-fold increase in nuclear size compared to the wild type (Figure 6D, S6D-F).

The commitment to SMC fate is difficult to resolve at early stages due to the lack of molecular
markers. Building on our previous findings, we rationalized that cells from the L2 apical domain with a
low frequency of mitotic events could have started to transit toward S-phase. To learn more about cell
cycle progress in this domain, we analyzed a GFP-tagged PCNA variant marking the replication
machinery (Yokoyama *et al*., 2016). When engaged at active replication forks, PCNA-GFP shows nuclear
speckles characteristic of S-phase (Strzalka and Ziemienowicz, 2011). By contrast, it remains in the
nucleoplasm in G1- and G2-phase and is undetectable in M-phase. We quantified the nuclear
distribution patterns of PCNA-GFP in cells from the L2 apical domain in wild type and *kat* primordia

separately for class A and B ovules (Figure 6E-H). At stages 0-II and 0-III, PCNA-GFP was always
detectable in both classes, indicating that L2 apical cells are rarely in M-phase, consistent with the
CYCB1.1-dbGFP marker analysis. Although the majority of class A and B ovules at these stages showed a
nucleoplasmic pattern, a significant proportion exhibited the speckled S-phase pattern (Figure 6F). At
stage 0-II, 24% of class A ovules (n=112) showed the S-phase pattern in a single, centrally positioned
apical cell (candidate SMC), reaching nearly 40% at stage 0-III (n=79) (Figure 6F). Conversely, SMC
neighbors rarely showed the S-phase pattern (2 to 5%). In class B ovules, 30% (n=41) and 28% (n=60) at
stages 0-II and 0-III, respectively, exhibited the S-phase pattern in at least one of the cells in the apical
domain (Figure 6F). Together with our analysis of the *CYCB1.1-dbGFP* marker, these data suggest a
prolonged S-phase in SMC candidates in Phase I (stages 0-II, 0-III). Although it is not possible to resolve
mitotic *versus* meiotic S-phase, it is tempting to speculate that this is typical for the latter.

At the onset of Phase II, PCNA-GFP showed a nucleoplasmic pattern in the SMC of class A ovules
and in the L2 apical cells of class B ovules (Figure 6G-H, S6G-H). This is consistent with the detection of
newly replicated DNA in the SMC at stage 1-I (She *et al.,* 2013). In contrast to the SMC, the SMC
neighbors display the S-phase pattern (Figure 6G-H), consistent with the late divisions reported for these
cells in Phase II (Figure 2D). In *kat* primordia, we found an higher frequency of S-phase patterns in SMC
candidates in both class A and B ovules in Phase II (Figure 6H, S6F-G) and a lower frequency for SMC
neighbors in class A ovules at stages 1-I and 1-II, consistent with the deregulation of *CYCB1.1-dbGFP* in
these cells in *kat* primordia.

Collectively, our data indicate a cellular heterogeneity in the L2 apical domain that leads to the
emergence of one or several SMC candidates as early as stage 0-II. The gradual decrease in the number
of primordia with ambiguous SMC candidates suggests developmental canalization associated with a
cell-cycle progression, cellular elongation, and robust expression of SMC fate markers. In *kat* primordia
displaying architecture alterations, SMC singleness is largely disrupted: the selection of a single SMC is
significantly delayed and several SMC candidates persist in an ambiguous cell fate state.

Discussion

Organogenesis involves coordinated cell division and cell expansion, complex growth processes
orchestrated by biochemical and mechanical cues (Echevin *et al.*, 2019). How cell differentiation is
coordinated in space and time during organ growth and whether these processes are interrelated are
core questions for the elucidation of patterning principles (Whitewoods and Coen, 2017).

Here, we tested the hypothesis that ovule primordium growth and overarching organ
architecture determines the differentiation of a single SMC, the precursor cell of the female germline,
per primordium. We established a reference collection of images capturing ovule primordia in 3D at
cellular resolution to study how SMC cell fate relates to organogenesis in space and time. This reference
set populates a growing number of 3D-segmented images of plant tissues and organs, produced as
described here or with new tools for automated analyses (Wolny *et al.*, 2020). This is a pre-requisite to
generate a developmental atlas that will integrate organ morphology at cellular resolution and gene
expression (Hartmann *et al.*, 2020).

We report on an exhaustive morphodynamic analysis at cellular resolution, *per* stage, domain,
layer, and cell type of the ovule primordium. In combination with reporter gene analyses in wild-type
and mutant ovules and tissue growth simulations, this approach aims to elucidate fundamental
patterning principles. We found that cell division is not uniformly distributed in space and time and
follows a biphasic pattern where proliferation frequencies in L1 and L2,L3 domains alternate between
Phases I and II to shape the primordium. Strikingly, the L2 apical domain shows a low mitotic activity in
Phase I (stages 0-II, 0-III) with a prolonged S-phase in the SMC candidate. This is reminiscent of the
animal germline, where mitotic quiescence is a prerequisite to initiate meiosis (Kimble, 2011; Reik and
Surani, 2015) and suggests a functional analogy between the L2,L3 apical domain in ovule primordia and
the primordial germline niche in animals. Subsequently, the onset of Phase II coincides with the marked
elongation and mitotic quiescence of the SMC, contrasting with active divisions in the basal domain.
Thus, the fate of the SMC may be uncoupled early from that of the surrounding cells.

Tissue growth simulations allowed us to test the role of three main factors contributing to ovule
primordium patterning: i) anisotropic cell growth, ii) spatially confined specified growth signal along the
ovule axis, and iii) passive strain–based growth (*i.e.* the baseline competency of the tissue to
accommodate to accumulated strain, independently of growth signal). Omitting any of these factors, the
resulting models failed to shape a realistic primordium. In particular, isotropic growth produces
hemispheric instead of digit-shaped primordia with a flat dome resembling *kat* primordia, in which
anisotropic cell growth is altered (Luptovciak *et al.*, 2017a). Thus, the *kat* mutant phenotype provides an
in planta confirmation of our theoretical models. Furthermore, the observation that abnormally shaped
kat primordia contain cells with ectopic expression of SMC fate markers is consistent with the
hypothesis that organ growth and ovule primordium architecture control SMC differentiation.

Moreover, our results highlight the developmental constraints for differentiation of a single SMC
corresponding to a canalization process. The formation of a single SMC prevails in most angiosperms

(Huang and Russell, 1992). Yet, some *Arabidopsis* accessions develop ovule primordia with multiple SMC

candidates but ultimately only one SMC is committed to meiosis (Grossniklaus and Schneitz, 1998;

Rodriguez-Leal *et al.*, 2015). Our quantifications in two reference accessions (Col-0, Ws) suggest that

natural variation affects the timing of the canalization process. Canalization, referring to a progressive

constraint of cell fate choice, is well studied in animal systems where it is modulated by intercellular

signal-based feedback mechanisms (Heitzler and Simpson, 1991) and epigenetic regulation (Pujadas and

Feinberg, 2012). Possibly, the epigenetic and signal-based processes known to regulate SMC

differentiation (Lora *et al.*, 2019, Pinto *et al.*, 2019) also operate in SMC fate canalization. In addition,

the contribution of morphogenetic processes on cell fate specification was recently re-evaluated in the

light of tissue mechanics and patterning processes (Chan *et al.,* 2017). Our results are consistent with a

role of primordium growth constraints on SMC fate canalization. Possibly, the prolonged cell cycle of

subepidermal cells in Phase I leverages competitive cell growth on which canalization can operate.

Under this scenario, disturbance of growth homeostasis as in *kat* mutants would render this process

ineffective and lead to a higher frequency of primordia with unresolved SMC candidates, unveiling a

coupling of SMC fate canalization with organ growth (Figure 6I). De-canalization observed in *kat* mutants

is consistent with recent work in *Drosophila* embryos showing that organ shape contributes to

developmental canalization (Huang *et al.,* 2020). Interestingly, the formation of a broader dome in *kat*

ovule coincides with the formation of a broader inner domain with more cells with an unresolved SMC

fate. This is reminiscent of the shoot apical meristem where the shape of the dome correlates with the

size and position of domains expressing stem cell regulators, such as WUSCHEL and CLAVATA3 (Gruel *et*

al., 2016).

Epidermal growth controls shoot apical meristem and shoot growth through the modulation of

the expression domains of key regulators in the underlying layers (Savaldi-Goldstein *et al.*, 2007; Gruel

et al., 2016). Whether growth of the epidermis similarly controls the cell cycle and SMC fate regulators

in underlying layers of ovule primordia is an exciting question for future investigation, particularly

considering the role of mechanical constraints on gene regulation (Landrein *et al.*, 2015; Fal *et al.*, 2017).

Reciprocally, the role of the underlying layers for epidermis growth and dome shape remains to be

determined. Our ovule growth simulations predicted a role for inner tissue-driven growth, similar to

what has been proposed for the emergence of leaf primordia (Peaucelle *et al.,* 2011), although this

remains debated (Boudon *et al.*, 2015). Moreover, our simulations suggested that passive strain-based

growth in the L1 is necessary to achieve a normal primordium shape, and that anisotropic material

properties in this layer are required to constrain the relaxation along the main axis of the primordium. A

possible explanation for this is the requirement to resolve mechanical conflicts arising between tissue

layers that grow at different rates (Hervieux *et al.*, 2016). This would be consistent with the high

proliferation activity in the L1 dome at a stage of marked differential growth of the SMC. In this

scenario, divisions in L1 may release the mechanical stress generated by SMC growth, leading to

anisotropic organ growth. Consistent with this idea, we observed that reduced organ shape anisotropy

in *kat* ovules is associated with fewer divisions in the L1 dome. The higher division rate in the wild type

might confer higher flexibility for the L1 to bend and follow an anisotropic growth profile driven by the

inner layers. By contrast, in *katanin*, non-dividing and elongated cells in the L1 would be expected to

transmit higher tension preventing bending. Experimental alterations of properties in L1 or L2,L3 cells,

together with 3D modeling, would be required to explore such scenarios.

Cell-based models suggested that blocking divisions in the SMC enables it to grow and push on

the surrounding cells, affecting significantly the ovule morphology. Therefore, we speculate that, in

order to maintain ovule growth homeostasis, differential SMC growth must adapt to the constraints

provided by surrounding cells. A gradual reduction of SMC turgor pressure may provide a plausible

cause to limit SMC size and maintain correct ovule anisotropy. Due to the limitations of 2D MS-based

models, such predictions will need to be tested in 3D cell-based models able to properly reproduce a

mechanical scenario for a growing ovule primordium. Nonetheless, this prediction paves the way to

further investigate the potential role of turgor pressure gradients across a field of cells, which could

provide positional information through the directional movement of water and other molecules,

thereby linking organ growth homeostasis, cell growth, and cell fate (Long *et al.*, 2020; Beauzamy *et al.*,

2014). Such mechanisms could participate in defining the specific pattern of a growth signal predicted by

our simulations. However, the nature of the growth signal(s) and the mechanism(s) establishing its

pattern in the placenta prior to primordium emergence remain to be identified. Candidate signals may

be the many differentially localized hormones, gene products, and small RNAs known to affect ovule

primordium growth (Pinto *et al.*, 2019, Su *et al.*, 2020). For example, in Phase II, auxin responses are

restricted to the L1 dome while cytokinin signaling is confined to a region basal to the SMC (Bencivenga

et al., 2012). Whether this pattern is established already in Phase I and plays a causative role in

primordium patterning and SMC differentiation remains to be determined.

Author contributions

CB, DA, GM, EHL, DG, OH, AB, CG: Conceptualization; EHL, GM, CB, DA, OH, AB, CG, DG: Methodology; GM, NP, AG: Softwares; EHL, GM, EMS, DA, CB: Validation; EHL, GM, AG, DA, CB: Formal analysis; EHL, EMS, DA, CB: Investigation; EMS, AF, OH: Ressources; EHL, GM, NP, DA, CB: Data curation; EHL, GM, EMS, DG, DA, CB: Writing – Original Draft; EHL, GM, OH, AB, CG, UG, DG, DA, CB: Writing – Reviewing and Editing; EHL, GM, DA, CB: Visualization; CB, DA, DG, OH, AB, UG: Supervision; CB, DA, DG: Project administration; CB, DA, DG, OH, AB, CG, UG: Funding Acquisition. All authors reviewed and revised the manuscript.

Data availability

All materials, scripts and datasets generated and analyzed in the current study are available at repositories (see Star Methods) or from the corresponding authors upon reasonable request.

Acknowledgments

We thank S. Strauss (Max Planck Institute for Plant Breeding Research, Cologne, Germany) and B. Lane (John Innes Centre, Norwich, U.K.) for providing code and assessment for mesh generation and simulations; S. Guyer (Bitplane, Switzerland) and M. Lartaud (CIRAD, France) for their help with IMARIS analyses; O. Leblanc (IRD Montpellier, France) for his help with data visualization in R; Montpellier Rio Imaging and the Center for Microscopy and Image Analysis of the University of Zurich for microscopy imaging infrastructures; M. Ueda (Nagoya University, Japan), I. Siddiqi (Center for Cellular and Molecular Biology, India), N. de Storme (Katholieke Universiteit Leuven, Belgium), S. Matsunaga (Tokyo Science University, Japan), M. Tucker (University of Adelaide, Australia), and the NASC stock center for seed germplasms.

This work was funded by the University of Zurich, IRD, and a PRCI grant from the Agence Nationale de la Recherche/Swiss National Science Foundation (n°ANR-16-CE93-0002-01 to 04) to CB and DA; a grant from the Swiss National Science Foundation (n° 310030B_160336) to UG, a CONACYT fellowship (n°438277) to EHL, and a fellowship from the Forschungskredit of the University of Zurich (n°FK-74502-04-01) to GM.

References

References

Agashe, B., Prasad, C.K., and Siddiqi, I. (2002). Identification and analysis of DYAD: a gene required for meiotic
chromosome organisation and female meiotic progression in Arabidopsis. Development 129, 3935-3943.

Amar, M.B.G., A.; M$\sqrt{}$ ᵒller, M.M.; Cugliandolo, L. (2011). New Trends in the Physics and Mechanics of Biological
Systems, Volume 92, (Oxford: Oxford University Press).

Bajon, C.H., C.; Motomayor, J.C.; Sauvanet, A.; Robert D. (1999). Megasporogenesis in Arabidopsis thaliana L.: an
ultrastructural study. Sexual Plant Reproduction 12, 99-109.

Beauzamy, L., Nakayama, N., and Boudaoud, A. (2014). Flowers under pressure: ins and outs of turgor regulation in
development. Ann Bot 114, 1517-1533.

Bencivenga, S., Simonini, S., Benkova, E., and Colombo, L. (2012). The transcription factors BEL1 and SPL are
required for cytokinin and auxin signaling during ovule development in Arabidopsis. Plant Cell 24, 2886-2897.

Bichet, A., Desnos, T., Turner, S., Grandjean, O., and Hofte, H. (2001). BOTERO1 is required for normal orientation
of cortical microtubules and anisotropic cell expansion in Arabidopsis. Plant J 25, 137-148.

Boudon, F., Chopard, J., Ali, O., Gilles, B., Hamant, O., Boudaoud, A., Traas, J., and Godin, C. (2015). A
computational framework for 3D mechanical modeling of plant morphogenesis with cellular resolution. PLoS
Comput Biol 11, e1003950.

Bouquin, T., Mattsson, O., Naested, H., Foster, R., and Mundy, J. (2003). The Arabidopsis lue1 mutant defines a
katanin p60 ortholog involved in hormonal control of microtubule orientation during cell growth. J Cell Sci 116,
791-801.

Bowman, J.L.e. ed. (1993). Arabidopsis: an Atlas of Morphology and Development. (Berlin & New York: Springer-
Verlag).

Bozorg, B., Krupinski, P., and Jonsson, H. (2016). A continuous growth model for plant tissue. Phys Biol 13, 065002.

Brodersen, P., Sakvarelidze-Achard, L., Bruun-Rasmussen, M., Dunoyer, P., Yamamoto, Y.Y., Sieburth, L., and
Voinnet, O. (2008). Widespread translational inhibition by plant miRNAs and siRNAs. Science 320, 1185-1190.

Burk, D.H., and Ye, Z.H. (2002). Alteration of oriented deposition of cellulose microfibrils by mutation of a katanin-
like microtubule-severing protein. Plant Cell 14, 2145-2160.

Cao, L., Wang, S., Venglat, P., Zhao, L., Cheng, Y., Ye, S., Qin, Y., Datla, R., Zhou, Y., and Wang, H. (2018).
Arabidopsis ICK/KRP cyclin-dependent kinase inhibitors function to ensure the formation of one megaspore
mother cell and one functional megaspore per ovule. PLoS Genet 14, e1007230.

Chan, C.J., Heisenberg, C.P., and Hiiragi, T. (2017). Coordination of Morphogenesis and Cell-Fate Specification in
Development. Curr Biol 27, R1024-R1035.

Coen, E., Rolland-Lagan, A.G., Matthews, M., Bangham, J.A., and Prusinkiewicz, P. (2004). The genetics of
geometry. Proc Natl Acad Sci U S A 101, 4728-4735.

De Storme, N., Keceli, B.N., Zamariola, L., Angenon, G., and Geelen, D. (2016). CENH3-GFP: a visual marker for
gametophytic and somatic ploidy determination in Arabidopsis thaliana. BMC Plant Biol 16, 1.

Echevin, E., Le Gloanec, C., Skowronska, N., Routier-Kierzkowska, A.L., Burian, A., and Kierzkowski, D. (2019).
Growth and biomechanics of shoot organs. J Exp Bot 70, 3573-3585.

Extavour, C.G., and Akam, M. (2003). Mechanisms of germ cell specification across the metazoans: epigenesis and
preformation. Development 130, 5869-5884.

Fal, K., Asnacios, A., Chaboute, M.E., and Hamant, O. (2017). Nuclear envelope: a new frontier in plant
mechanosensing? Biophys Rev 9, 389-403.

Garcia-Aguilar, M., Michaud, C., Leblanc, O., and Grimanelli, D. (2010). Inactivation of a DNA methylation pathway
in maize reproductive organs results in apomixis-like phenotypes. Plant Cell 22, 3249-3267.

Grossniklaus, U., and Schneitz, K. (1998). The molecular and genetic basis of ovule and megagametophyte
development. Semin Cell Dev Biol 9, 227-238.

Hallgrimsson, B., Green, R.M., Katz, D.C., Fish, J.L., Bernier, F.P., Roseman, C.C., Young, N.M., Cheverud, J.M., and
Marcucio, R.S. (2019). The developmental-genetics of canalization. Semin Cell Dev Biol 88, 67-79.

Hartmann, J., Wong, M., Gallo, E., and Gilmour, D. (2020). An image-based data-driven analysis of cellular
architecture in a developing tissue. Elife 9.

Heitzler, P., and Simpson, P. (1991). The choice of cell fate in the epidermis in Drosophila. Cell 22, 1083-1092.

Hervieux, N., Dumond, M., Sapala, A., Routier-Kierzkowska, A.L., Kierzkowski, D., Roeder, A.H., Smith, R.S.,
Boudaoud, A., and Hamant, O. (2016). A Mechanical Feedback Restricts Sepal Growth and Shape in Arabidopsis.
Curr Biol.

Hong, L., Dumond, M., Zhu, M., Tsugawa, S., Li, C.B., Boudaoud, A., Hamant, O., and Roeder, A.H.K. (2018).
Heterogeneity and Robustness in Plant Morphogenesis: From Cells to Organs. Annu Rev Plant Biol 69, 469-495.

Huang, A., Rupprecht, J.F., and Saunders, T.E. (2020). Embryonic geometry underlies phenotypic variation in
decanalized conditions. Elife 9.

Huang, B.Q.R., S.D. (1992). Female Germ Unit: Organization, Isolation, and Function. International Review of
Cytology 140, 233-293.

Kennaway, R., Coen, E., Green, A., and Bangham, A. (2011). Generation of diverse biological forms through
combinatorial interactions between tissue polarity and growth. PLoS Comput Biol 7, e1002071.

Kimble, J. (2011). Molecular regulation of the mitosis/meiosis decision in multicellular organisms. Cold Spring Harb
Perspect Biol 3, a002683.

Klimyuk, V.I., and Jones, J.D. (1997). AtDMC1, the Arabidopsis homologue of the yeast DMC1 gene:
characterization, transposon-induced allelic variation and meiosis-associated expression. Plant J 11, 1-14.

Komis, G., Luptovciak, I., Ovecka, M., Samakovli, D., Samajova, O., and Samaj, J. (2017). Katanin Effects on
Dynamics of Cortical Microtubules and Mitotic Arrays in Arabidopsis thaliana Revealed by Advanced Live-Cell
Imaging. Front Plant Sci 8, 866.

Kuchen, E.E., Fox, S., de Reuille, P.B., Kennaway, R., Bensmihen, S., Avondo, J., Calder, G.M., Southam, P.,
Robinson, S., Bangham, A., *et al*. (2012). Generation of leaf shape through early patterns of growth and tissue
polarity. Science 335, 1092-1096.

Landrein, B., Kiss, A., Sassi, M., Chauvet, A., Das, P., Cortizo, M., Laufs, P., Takeda, S., Aida, M., Traas, J., et al.
(2015). Mechanical stress contributes to the expression of the STM homeobox gene in Arabidopsis shoot
meristems. Elife 4, e07811.

Long, Y., Cheddadi, I., Mosca, G., Mirabet, V., Dumond, M., Kiss, A., Traas, J., Godin, C., and Boudaoud, A. (2020).
Cellular Heterogeneity in Pressure and Growth Emerges from Tissue Topology and Geometry. Curr Biol 30, 1504-
1516 e1508.

Lora, J., Herrero, M., Tucker, M.R., and Hormaza, J.I. (2017). The transition from somatic to germline identity shows
conserved and specialized features during angiosperm evolution. New Phytol 216, 495-509.

Lora, J., Yang, X., and Tucker, M.R. (2019). Establishing a framework for female germline initiation in the plant
ovule. J Exp Bot 70, 2937-2949.

Luptovciak, I., Komis, G., Takac, T., Ovecka, M., and Samaj, J. (2017). Katanin: A Sword Cutting Microtubules for
Cellular, Developmental, and Physiological Purposes. Front Plant Sci 8, 1982.

Luptovciak, I., Samakovli, D., Komis, G., and Samaj, J. (2017). KATANIN 1 Is Essential for Embryogenesis and Seed
Formation in Arabidopsis. Front Plant Sci 8, 728.

Mendocilla-Sato, E.B., C. (2017). Analysis of 3D Cellular Organization of Fixed Plant Tissues Using a User-guided
Platform for Image Segmentation. Bio-protocols 7, e2355.

Mosca, G.A., M.; Strauss, S.; Runions, R.; Sapala, A.; Smith, R.S. (2018). Modeling plant tissue growth and cell
division. In Mathematical Modelling in Plant Biology. (Springer, Cham), pp. 107-138.

Nonomura, K., Miyoshi, K., Eiguchi, M., Suzuki, T., Miyao, A., Hirochika, H., and Kurata, N. (2003). The MSP1 gene is
necessary to restrict the number of cells entering into male and female sporogenesis and to initiate anther wall
formation in rice. Plant Cell 15, 1728-1739.

Olmedo-Monfil, V., Duran-Figueroa, N., Arteaga-Vazquez, M., Demesa-Arevalo, E., Autran, D., Grimanelli, D.,
Slotkin, R.K., Martienssen, R.A., and Vielle-Calzada, J.P. (2010). Control of female gamete formation by a small RNA
pathway in Arabidopsis. Nature 464, 628-632.

Ovefçka M, L.I., Komis G, ≈†amajov√⁻° O, Samakovli D, ≈†amaj J. (2020). Spatiotemporal Pattern of Ectopic Cell
Divisions Contribute to Mis-Shaped Phenotype of Primary and Lateral Roots of katanin1 Mutant. Front. Plant Sci.

Paluch, E., and Heisenberg, C.P. (2009). Biology and physics of cell shape changes in development. Curr Biol 19,
R790-799.

Panteris, E., Adamakis, I.D., Voulgari, G., and Papadopoulou, G. (2011). A role for katanin in plant cell division:
microtubule organization in dividing root cells of fra2 and lue1Arabidopsis thaliana mutants. Cytoskeleton
(Hoboken) 68, 401-413.

Peaucelle, A., Braybrook, S.A., Le Guillou, L., Bron, E., Kuhlemeier, C., and Hofte, H. (2011). Pectin-induced changes
in cell wall mechanics underlie organ initiation in Arabidopsis. Curr Biol 21, 1720-1726.

Pinto, S.C., Mendes, M.A., Coimbra, S., and Tucker, M.R. (2019). Revisiting the Female Germline and Its Expanding
Toolbox. Trends Plant Sci 24, 455-467.

Pujadas, E., and Feinberg, A.P. (2012). Regulated noise in the epigenetic landscape of development and disease.
Cell 148, 1123-1131.

Reik, W., and Surani, M.A. (2015). Germline and Pluripotent Stem Cells. Cold Spring Harb Perspect Biol 7.

Ren, H., Dang, X., Cai, X., Yu, P., Li, Y., Zhang, S., Liu, M., Chen, B., and Lin, D. (2017). Spatio-temporal orientation of
microtubules controls conical cell shape in Arabidopsis thaliana petals. PLoS Genet 13, e1006851.

Rodriguez, E.K., Hoger, A., and McCulloch, A.D. (1994). Stress-dependent finite growth in soft elastic tissues. J
Biomech 27, 455-467.

Rodriguez-Leal, D., Leon-Martinez, G., Abad-Vivero, U., and Vielle-Calzada, J.P. (2015). Natural variation in
epigenetic pathways affects the specification of female gamete precursors in Arabidopsis. Plant Cell 27, 1034-1045.

Savaldi-Goldstein, S., Peto, C., and Chory, J. (2007). The epidermis both drives and restricts plant shoot growth.
Nature 446, 199-202.

Scharloo (1991). Canalization: Genetic and Developmental Aspects. Annual Review of Ecology and Systematics 22,
65-93.

Schmidt, A., Schmid, M.W., and Grossniklaus, U. (2015). Plant germline formation: common concepts and
developmental flexibility in sexual and asexual reproduction. Development 142, 229-241.

Schmidt, A., Wuest, S.E., Vijverberg, K., Baroux, C., Kleen, D., and Grossniklaus, U. (2011). Transcriptome analysis of
the Arabidopsis megaspore mother cell uncovers the importance of RNA helicases for plant germline development.
PLoS Biol 9, e1001155.

Schneitz, K.; Hulskamp, M. and Pruitt, R.E. (1995). Wild-type ovule development in Arabidopsis thaliana: a light
microscope study of cleared whole-mount tissue. Plant Journal 7, 731-749.

She, W., Grimanelli, D., Rutowicz, K., Whitehead, M.W., Puzio, M., Kotlinski, M., Jerzmanowski, A., and Baroux, C.
(2013). Chromatin reprogramming during the somatic-to-reproductive cell fate transition in plants. Development
140, 4008-4019.

Sheridan, W.F., Avalkina, N.A., Shamrov, II, Batygina, T.B., and Golubovskaya, I.N. (1996). The mac1 gene:
controlling the commitment to the meiotic pathway in maize. Genetics 142, 1009-1020.

Sheridan, W.F., Golubeva, E.A., Abrhamova, L.I., and Golubovskaya, I.N. (1999). The mac1 mutation alters the
developmental fate of the hypodermal cells and their cellular progeny in the maize anther. Genetics 153, 933-941.

Sieber, P., Gheyselinck, J., Gross-Hardt, R., Laux, T., Grossniklaus, U., and Schneitz, K. (2004). Pattern formation
during early ovule development in Arabidopsis thaliana. Dev Biol 273, 321-334.

Skinner, D.J., Hill, T.A., and Gasser, C.S. (2004). Regulation of ovule development. Plant Cell 16 Suppl, S32-45.

Smith, R.S. (2009). Modeling plant morphogenesis and growth. In New Trends in the Physics and Mechanics of

Biological Systems. Lecture Notes of the Les Houches Summer School., Volume 22, M.G. Ben Amar, A.; M$\sqrt{}^{\circ}$ller,

M.M.; Cugliandolo, L., ed. (Oxford: Oxford University Press).

Strzalka, W., and Ziemienowicz, A. (2011). Proliferating cell nuclear antigen (PCNA): a key factor in DNA replication

and cell cycle regulation. Ann Bot 107, 1127-1140.

Su, Z., Zhao, L., Zhao, Y., Li, S., Won, S., Cai, H., Wang, L., Li, Z., Chen, P., Qin, Y., *et al.* (2017). The THO Complex

Non-Cell-Autonomously Represses Female Germline Specification through the TAS3-ARF3 Module. Curr Biol 27,

1597-1609 e1592.

Su, Z. ; Wang, N. ; Hou, Z. ; Li, B. ; Li, D. ; Liu, Y. ; Cai, H. ; Qin, Y. and Chen X. (2020). Regulation of Female Germline

Specification via Small RNA Mobility in Arabidopsis. Plant Cell. Jul 23:tpc.00126.2020. doi: 10.1105/tpc.20.00126.

Online ahead of print.

Truernit, E., Bauby, H., Dubreucq, B., Grandjean, O., Runions, J., Barthelemy, J., and Palauqui, J.C. (2008). High-

resolution whole-mount imaging of three-dimensional tissue organization and gene expression enables the study

of Phloem development and structure in Arabidopsis. Plant Cell 20, 1494-1503.

Tucker, M.R., Okada, T., Hu, Y., Scholefield, A., Taylor, J.M., and Koltunow, A.M. (2012). Somatic small RNA

pathways promote the mitotic events of megagametogenesis during female reproductive development in

Arabidopsis. Development 139, 1399-1404.

Ubeda-Tomas, S., Federici, F., Casimiro, I., Beemster, G.T., Bhalerao, R., Swarup, R., Doerner, P., Haseloff, J., and

Bennett, M.J. (2009). Gibberellin signaling in the endodermis controls Arabidopsis root meristem size. Curr Biol 19,

1194-1199.

Uyttewaal, M., Burian, A., Alim, K., Landrein, B., Borowska-Wykret, D., Dedieu, A., Peaucelle, A., Ludynia, M., Traas,

J., Boudaoud, A., *et al*. (2012). Mechanical stress acts via katanin to amplify differences in growth rate between

adjacent cells in Arabidopsis. Cell 149, 439-451.

Waddington, C.H. (1942). Canalization of development and the inheritance of acquired characters. Nature, 563-

565.

Whitewoods, C.D., and Coen, E. (2017). Growth and Development of Three-Dimensional Plant Form. Curr Biol 27,

R910-R918.

Wightman, R., Chomicki, G., Kumar, M., Carr, P., and Turner, S.R. (2013). SPIRAL2 determines plant microtubule

organization by modulating microtubule severing. Curr Biol 23, 1902-1907.

Wolny, A., Cerrone, L., Vijayan, A., Tofanelli, R., Vilches Barro, A., Louveaux, M., Wenzl, C., Steigleder, S., Pape, C.,

Bailoni, A., *et al.* (2020). Accurate and Versatile 3D Segmentation of Plant Tissues at Cellular Resolution. bioRxiv,

2020.2001.2017.910562.

Yokoyama, R., Hirakawa, T., Hayashi, S., Sakamoto, T., and Matsunaga, S. (2016). Dynamics of plant DNA

replication based on PCNA visualization. Sci Rep 6, 29657.

Zhang, Q., Fishel, E., Bertroche, T., and Dixit, R. (2013). Microtubule severing at crossover sites by katanin

generates ordered cortical microtubule arrays in Arabidopsis. Curr Biol 23, 2191-2195.

Zhao, X., Bramsiepe, J., Van Durme, M., Komaki, S., Prusicki, M.A., Maruyama, D., Forner, J., Medzihradszky, A.,

Wijnker, E., Harashima, H., *et al.* (2017). RETINOBLASTOMA RELATED1 mediates germline entry in Arabidopsis.

Science 356.

Zhao, X., de Palma, J., Oane, R., Gamuyao, R., Luo, M., Chaudhury, A., Herve, P., Xue, Q., and Bennett, J. (2008).

OsTDL1A binds to the LRR domain of rice receptor kinase MSP1, and is required to limit sporocyte numbers. Plant J

54, 375-387.

Figure 1

Figure 1. Reference set of 3D segmented images capturing Arabidopsis ovule primordium growth at cellular-resolution.

(A) 3D reconstruction of a whole gynoecium stained with PS-PI (cell wall dye) and visualized by CSLM. The cross section allows showing the nascent ovule primordia attached to the placenta.

(B) Ovule primordium developmental stages (0-I to 2-II) and organ viewpoints (domains) defined for the 3D quantitative analysis. All segmented data can be analyzed by an interactive interface named OvuleViz. See also Figure S1A-C.

n= number of ovules analyzed. Segmented images for all developmental stages are provided in Data S1. See also Figure S1 D-E, STAR Methods.

	0-I			0-II			0-III		
Cells above placenta*	2.5	(±3.5)	(n=21)	3.6	(±0.5)	(n=17)	5.9	(±1.0)	(n=11)
range (min-max)	2	-	3	3	-	4	5	-	8
total # cells	28	(±5.7)	(n=21)	38	(±8.0)	(n=17)	80	(±8.4)	(n=11)
width (µm) (W)	23	(±3.5)		26.3	(±2.4)		28.0	(±3.1)	
height (µm) (H)	5.3	(±1.2)		11.9	(±2.1)		22.5	(±3.6)	
H:W ratio	0.2	(±0.04)	(n=13)	0.45	(±0.06)	(n=14)	0.8	(±0.12)	(n=8)

	1-I			1-II		
cells above placenta*	7.3	(±1.3)	(n=15)	9.7	(±1.0)	(n=11)
range (min-max)	6	-	10	8	-	11
total # cells	105	(±10.6)	(n=15)	128	(±18.0)	(n=11)
width (µm) (W)	27	(±2.0)		27.8	(±3.5)	
height (µm) (H)	30	(±2.8)		39	(±5.9)	
H:W ratio	1.1	(±0.10)	(n=6)	1.4	(±0.22)	(n=8)

* scoring the length of L1 cell file above placenta

	2-I			2-II		
cells above placenta*	10	(±1.1)	(n=10)	12.6	(±1.0)	(n=7)
range (min-max)	9	-	12	11	-	14
total # cells	151	(±18.9)	(n=10)	165	(±19.8)	(n=7)

Table 1. Classification criteria of Arabidopsis ovule primordia

The table summarizes general characteristics of ovule primordia per stage: cell "layers" above the placenta scored as the number of L1 cells in a cell file drawn from the basis to the top, range thereof, total cell number, ovule shape including heigth, width, aspect ratio.

Figure 2

A Cell number and volume

B Surface curvature

C Ovule shape

D Fold change between stages

E Cell number – L1 *vs* L2,L3

F Cell number – apical *vs* basal

G *CYCB1.1-dbGFP*

H Mitotic activity map

I SMC *vs* nucellus

J L1 *vs* L2,L3

K SMC shape — SMC orientation relative to the ovule main axis

Figure 2. Ovule primordium morphogenesis involves domain-specific cell division and anisotropic cell growth patterns.

(A) Mean cell number *per* ovule increases mainly during stages 0-I to 0-III (Phase I), whereas cell volume *per* ovule remains constant on average across primordium development (stages 0-I to 2-II).

(B) Representative image of a continuous surface of an ovule primordium mesh and its projected median plane. Dashed lines indicate the minimal and maximal curvature points used to measure organ height and width. Color scale: minimal curvature mm-1 (see also Figure S2B).

(C) Anisotropic organ growth during Phase I and until stage 1-II. Mean width and height quantifications *per* stage.

(D) Phase I captures a distinct growth dynamic compared to Phase II. Fold-change of cell number and aspect ratio between stages are plotted.

(E) Mean cell number is increased at the L1 *versus* L2,L3 layers across developmental stages.

(F) Mean cell number is increased in the basal *versus* apical domain across developmental stages.

(G) Representative images of ovule primordia expressing the M-phase reporter *pCYCB1.1::db-GFP.* White arrows indicate dividing cells. Magenta signal: Renaissance SR2200 cell-wall label. Scale bar: 10μm.

(H) Domain-specific map of mitotic activity during ovule primordium development, scored using the *pCYCB1.1::db-GFP* reporter. The frequency of mitoses was calculated *per* ovule domain at each developmental stage and color-coded as indicated in the bar (right). n: total number of scored ovules.

(I) Mean SMC volume (dark blue) is significantly increased as compared to L2,L3 cells (pale blue) from stage 0-III onward, on even at earlier stages as compared to all other cells (inset).

(J) Mean cell volume is increased in L2,L3 cells as compared to L1 cells, at the two early developmental stages (st0-I, st0-III).

(K) The SMC consistently displays anisotropic shape with main axis of elongation aligned with ovule growth axis. The SMC anisotropy index (boxplot, left; stage 0-II, n=16 ovules) was calculated from the Maximum (dark blue), Minimum (light blue) and Medium (medium blue) covariance matrix eigenvalues, computed from 3D segmented cells (see Figure S2G). Image (middle): illustration of the SMC main anisotropy axis (orange arrow) related by an angle 'alpha' to the main axis of the ovule primordium (white arrow), stage 0-II. Radar plot (right): 'alpha' angle measured on z projections for n=16 ovule primordia at stage 0-II. See also Star Methods.

Error bars: Standard errors to the mean. Differences between cell types or primordium domains were assessed using a two-tailed Man Whitney U test in (A) and (I); a two-tailed Wilcoxon signed rank test in (J). *p≤0.05, **p≤0.01, ***p≤0.001. See also Figure S2, Data S2, STAR Methods.

Modeling hypotheses		Models							
		1	2	3	4	5	6	7	8
Growth anisotropy		●	●	-	●	●	●	●	●
Material anisotropy		●	-	○	○	○	○	○	○
Passive Strain-based relaxation		●	●	●	●	●	●	●	-
Signal-based growth									
Distribution	L1 only	-	-	-	-	-	●	-	-
	Inner tissue only (pit-shape)	-	-	-	-	-	-	●	-
	L1 + inner tissue (pit-shape)	●	●	●	-	●	-	-	●
	L1 + inner tissue (broad distribution)	-	-	-	●	-	-	-	-
Fixed high concentration	L1 only	-	-	-	-	●	●	-	-
	Inner tissue only	-	-	-	-	-	-	●	-
	L1 + inner tissue	●	●	●	●	-	-	-	●

Table 2. Hypotheses used to generate the mass-spring (MS) and continuous Finite Element Method (FEM) based simulations

Several growth and mechanical hypotheses were listed at start of modelling. To evaluate their effect on primordium growth each hypothesis was excluded (-) in at least one scenario. The FEM and mass-spring (MS) models presented in Figure 3 and S3 are numbered according to the scenarios (1-8) in the table. Since it is not possible for MS to simulate material anisotropy, Model 1 was only tested with FEM. Empty dots for "Material anisotropy" were considered only for the FEM models. See also STAR Methods.

Figure 3

Figure 3. Mechanical and cell-based 2D simulation models of ovule primordium development predict that ovule shape depends on cell growth anisotropy.

(A) Reference model for FEM-based simulation of ovule primordium growth. Growth stability is reached at T=24. The fine lines indicate principal strain directions (white: expansive strain, red: compressive strain).

(B) FEM-based simulation omitting the anisotropic growth parameter (FEM-model 3) showing abnormal ovule dome shape at T=24.

(C) Reference model for Mass Spring (MS) cell-based simulation of ovule primordium growth. Growth stability is reached at T=190.

(D) MS cell-based simulation with isotropic cell growth (MS-model 3), showing enlarged apex and dome shape and abnormal cell division planes in L1, at T=190.

(E) Reference model for MS cell-based simulation showing emergence of L2 cell anisotropy at T=58.

(F) MS cell-based simulation MS-Model 3 (see Table 2) showing reduced L2 cells anisotropy at T=58.

Heatmaps: strain levels (A-B), growth signal intensity (C-D), 2D anisotropy index (E-F). See also Figure S3, Table 2 for model hypotheses and STAR Methods.

Figure 4

Figure 4. *katanin* mutants show a distinct ovule primordium architecture, primarily through altered cell division patterns.

Comparison between wild-type (WT, Ws-4 accession) and *katanin (kat, bot1-7* allele*)* ovule primordia unless indicated otherwise.

(A) 3D segmented images at stage 1-II. External organ view (top) and longitudinal sections (bottom) are shown. Scale bar 10μm. See Data S2a for full datasets.

(B-D) Morphological difference between WT and *kat* primordia measured by their volume (B), the aspect ratio (C), and aspect ratio to bounding box occupancy relationship (D). D, Left: scheme representing the bounding box capturing the primordium's 3D surface. See also Figure S4A.

(E) Quantification of cell number, cell volume, and sphericity in comparisons of L1 *versus* L2,L3 and apical *versus* basal domains at stage 0-III. See also Figure S4C.

(F) Mitotic activity domains are altered in *kat* primordia. Top: Heatmap of Log2 fold change of mitotic activity in the *kat* mutant (*lue1* allele) *versus* WT (Col-0) *per* domain, at two developmental stages. The frequency of mitoses was measured as in Figure 2. Full maps of mitotic activity in different mutant alleles are shown Figure S4E. Bottom: representative images of *kat* primordia (*lue1* allele) expressing *pCYCB1.1::db-GFP*. Dashed lines mark L2 apical cells. Magenta signal: Renaissance SR2200 cell wall label. Scale bar 10μm.

(G) Mean cell volume and sphericity are increased in *katanin* L2 apical cells in contact with the SMC. SMC, cells in contact with the SMC, and cells not in direct contact with the SMC, are compared at stage 0-III.

Color code in all plots: Dark red: WT; Salmon: *kat* mutant. Error bars: standard error of the mean. Differences between WT and *kat* mutants in (B), (C), (E), and (G) were assessed using a Mann Whitney U test; a two-tailed Fischer's exact test was used in (F). P values: *p≤0.05, **p≤0.01, ***p≤0.001. See also Figure S4 and Data S4.

Figure 5

A Stage 1-II

WT / *kat*

top view

side 1

side 2

5 µm

B SMC L2 neighbors (st1-II)

number

volume

kat
WT

ellipticity

sphericity

C

Stage 1-II / Stage 2-II

WT / *kat* / *WT* / *kat*

85.60% n=132 / 33.94% n=109 / 95.78% n=95 / 14.42% n=104

D *H1.1-GFP*

WT / *kat*

1-II

% eviction in > 1 cell

WT *kat*

n= 43 42

E *KNU-nlsYFP*

WT / *kat*

1-II

% expression in > 1 cell

WT *kat*

n= 88 43

F *AtDMC1-GUS*

WT / *kat*

2-III 2-V

% expression in > 1 cell

WT *kat*

n= 169 99

Figure 5. Altered ovule primordium architecture in *katanin* mutants is associated with multiple SMCs.

(A) SMCs lose their typical pear shape in *katanin* (*kat*) mutants. 3D images of the apical-most cells according to top and side views as indicated, showing the SMC (yellow), SMC neighbors (purple), and the L1 dome (transparent red).

(B) Differential properties of L2,L3 apical domain cells (SMC and SMC neighbors) in terms of cell number, mean cell volume, ellipticity, and sphericity at stage 1-II. See also Figure S5A and Data S4b.

(C) Representative images of cleared wild-type (WT) and *kat* ovule primordia. The % indicate the frequency of ovules showing one SMC for WT primordia, or multiple SMC candidates (dashed lines) for *kat* primordia.

(D-E) Representative images and quantification of SMC fate markers in WT and *kat* primordia: eviction of the *H1.1::H1.1-GFP* marker (green, D) and ectopic expression of *KNU::nls-YFP* (yellow, E) in more than one cell *per* primordia, are increased in *kat* primordia. See also Figure S5B.

(F) The meiotic marker *AtDMC1::GUS* is ectopically expressed in *kat* ovules.

Mutant alleles: *bot1-7* (A-B), *mad5* (C-E), *lue1* (F). Additional *kat* alleles, stages and detailed quantifications are presented Figure S5 and Data S5. Magenta signal in (B) and (C): Renaissance SR2200 cell wall label. Scale bars for (C), (D) and (E): 10μm. n: number of ovules scored. Error bar: standard error of the mean. Differences between WT and *kat* genotype*s* were assessed using a Mann Whitney U test in (B), and a two-tailed Fischer's exact test in (D), (E), and (F). P values: *p≤0.05, **p≤0.01, ***p≤0.001. See also Figure S5 and Data S5.

Figure 6

Figure 6. The formation of a single SMC stems from a canalization process.

(A-B) In wild-type (WT) plants, ovule primordia harbour one (Class A) or occasionally two (Class B) SMC candidates with the frequency of class B gradually decreasing during development, reminiscent of developmental canalization. Typical images obtained by tissue clearing with SMC candidates highlighted in yellow (A) and plots showing the percentages of classes A and B (B). The frequency of class B ovules is significantly reduced from stage 1-I, suggesting that canalization (represented by the red dashed line) occur before that stage. The plot coloration is a visual aid only.

(C) Developmental canalization is delayed in *katanin* (*kat*) mutants (*mad5* allele). The proportion of class B ovules is significantly reduced from stage 2-II only. Quantifications for two additional mutants carrying *kat* alleles are presented Figure S6A. The plot coloration is a visual aid only.

(D) The candidate SMCs initially identified as enlarged cells consistently show an enlarged nucleus in both class A and class B primordia. Nuclear area is compared between the candidate SMCs (cSMC) and surrounding L2 and L1 cells (see Figure S6D and STAR Methods for further details). Box plots include jittered data points to visualize data variability. Red lines represent the median of cSMC nuclear area for comparison with other cell types. Equivalent quantifications for stage 0-III are presented in Figure S6F.

(E-F) During Phase I (stages 0-II, 0-III), S-phase is detected in candidate SMCs at higher frequency than in neighbors cells in class A ovules, and in both SMC candidates in class B ovules. Representative images of ovule primordia showing the speckled S-phase pattern of the *pPCNA1::PCNA:sGFP* reporter (green) (E). Magenta signal: Renaissance SR2200 cell wall label. Scale bar 10µm. Quantification of speckled S-phase pattern in the SMC candidate and L2 neighbors (class A ovules) or in both SMC candidates (class B) (F).

(G-H) During Phase II (stages 1-I to 2-II), in class A wild-type ovule primordia, SMC exits S-phase, and neighbors cells re-enter S-phase; while *katanin* primordia show the opposite pattern.. Representative images of class A ovule primordia primarily showing the nucleoplasmic G1/G2-phase pattern of *pPCNA1::PCNA:sGFP* in SMCs at stages 1-II (left panel) and 2-II (middle panel); and of the speckled S-phase pattern in neighbors cells (right panel). Quantification of speckled S-phase pattern in SMC candidates and neighbors in Phase II wild-type and *katanin* primordia (I). Representative images and quantifications of Phase I *katanin* primordia and of class B primodia are presented in Figure S6G-H. See also Figure 2H, S6E.

(I) Model for the role of *KATANIN* in primordium growth and SMC differentiation (graphical abstract). In WT primordia, SMC differentiation follows a developmental canalization process influenced by cell growth anisotropy that shapes the primordium apex. In *kat* mutants, reduced anisotropy modifies the cell proliferation pattern, enlarges the apex and the L2,L3 apical domain, leading to multiple SMC candidates (delayed canalization). We propose that ovule primordium shape, controlled by anisotropic cell growth, determines SMC singleness.

Images: scale bars : 10um. *Graphs*: n= total number of ovules. Error bar: standard error of the mean (F, I). Differences between cell types, domains, or genotypes as indicated in the graphs were assessed using Wilcoxon signed rank test in (D), and a two-tailed Fischer's exact test in B, C, F, and I. P values: *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$. Quantifications for additional alleles and detailed quantifications are provided in Figure S6 and Data S6.