1 **PHYTOCHROMES B1/B2 ARE MAJOR REGULATORS OF RIPENING-**
2 **ASSOCIATED EPIGENOME REPROGRAMMING IN TOMATO FRUITS**
3

Ricardo Bianchetti[1], Nicolas Bellora[2], Luis A de Haro[3], Rafael Zuccarelli[1], Daniele Rosado[1§],
Luciano Freschi[1], Magdalena Rossi[1], Luisa Bermudez[4,5*]
[1]Departamento de Botânica, Instituto de Biociências, Universidade de São Paulo, São Paulo,
Brasil.
[2]Instituto Andino Patagónico de Tecnologias Biológicas y Geoambientales - CCT-
CONICET, Patagonia Norte, Bariloche, Argentina.
[3]Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot,
Israel.
[4]Instituto de Agrobiotecnología y Biología Molecular (IABIMO), CICVyA, INTA-
CONICET, Argentina.
[5]Cátedra de Genética, Facultad de Agronomía, Universidad de Buenos Aires, Buenos Aires,
Argentina.
16
17 [§]Current address: Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, USA.
18
19 * Corresponding Author.
20 Luisa Bermúdez
21 Phone: +54 11 4621 1447
22 email: bermudez.luisa@inta.gob.ar
23

24 **Author contributions**
25 RB performed most of the experiments and analysed the data; LB, NB, LAH and RZ analysed
26 the data; DR performed the experiments; RB, LF, MR and LB conceived the project,
27 designed the experiments and wrote the paper, which was revised and approved by all
28 authors. LB agrees to serve as the author responsible for contact and ensures communication.
29

ACKNOWLEDGMENTS

This work was supported by FAPESP (Fundação de Amparo à Pesquisa do Estado de São

Paulo, Grant Number #2016/01128-9); RB was a recipient of a FAPESP fellowship

(#2017/24354-7). MR was a recipient of a CNPq fellowship. LB was a recipient of a CAPES-

PRINT scholarship (88887.370243/2019-00).

Abstract

Phytochrome-mediated light and temperature perception has been shown to be a major

regulator of fruit development. Furthermore, chromatin remodelling via DNA demethylation

has been described as a crucial mechanism behind the fruit ripening process; however, the

molecular basis underlying the triggering of this epigenetic modification remains largely

unknown. Here, through integrative analyses of the methylome, siRNAome and

transcriptome of tomato fruits from *phyA* and *phyB1B2* null mutants, we report that PHYB1

and PHYB2 control genome-wide DNA methylation during fruit development. The

experimental evidence indicates that PHYB1B2 signal transduction is mediated by the

coordinated expression of DNA methylases/demethylases, histone-modifying enzymes and

chromatin remodelling factors, resulting in the transcriptional regulation of photosynthetic

and ripening-associated genes. This new level of understanding provides insights into the

orchestration of epigenetic mechanisms in response to environmental cues affecting

agronomical traits.

Keywords: carotenoid, chlorophyll, DNA methylation, epigenetics, fleshy fruit,

phytochrome, RNA-seq, RdDM, tomato.

INTRODUCTION

As sessile organisms, plants must constantly monitor their environment and continuously tune their gene expression to enable adaptation and survival[1]. Light is one of the main environmental cues that controls plant growth and development from seed germination to senescence[2]. Plants employ different photoreceptors to detect and respond to changes in the incident spectral composition (from UV-B to far-red wavelengths), light direction and photoperiod. These photoreceptor families include (i) phytochromes (PHYs), which perceive red/far-red (R/FR) light; (ii) cryptochromes (CRYs), phototropins, and 'Zeitlupes', which sense blue/UV-A light; and (iii) the UV-B receptor UVR8[3].

After photoreceptor activation, complex signal transduction pathways control the expression of light-regulated genes via transcriptional, posttranscriptional, and posttranslational mechanisms[2]. Several hub proteins in the light signal transduction pathway triggered by PHYs, CRYs and UVR8 have been identified, including transcription factors (TFs) such as PHY-INTERACTING FACTORS (PIFs) and ELONGATED HYPOCOTYL5 (HY5) as well as the ubiquitin E3 ligase complex comprising CONSTITUTIVE PHOTOMORPHOGENIC1 (COP1)[2]. Additionally, the effect of light on alternative splicing (AS) has recently been described[4]. In *Arabidopsis thaliana,* 1,505 genes undergo changes in the AS levels of their transcripts within 1 h of exposure to red light in a PHY-dependent manner[5]. Similarly, in the moss *Physcomitrella patens*, 8.4% and 8.9% of AS events rapidly respond to PHY-mediated red and blue light perception[6]. Further studies showed that the interaction between PpPHY4 and a splicing factor coregulates 70% of intron retention (IR) events in response to red light in *P. patens*[7]. Moreover, light controls protein localization through PHY-mediated alternative promoter selection, allowing plants to metabolically respond to changing light conditions[8]. Finally, it has been extensively reported that activated PHYs induce post-translational changes in PIF proteins, including sequestration, phosphorylation, polyubiquitylation, and subsequent degradation through the 26S proteasome-mediated pathway[3]. Although the effect of light on plant phenotypes and the plant transcriptome has been studied for decades[9–11], the involvement of epigenetic regulatory mechanisms in light-dependent changes in the transcriptional landscape remains poorly addressed.

Posttranslational histone modifications, such as acetylation and methylation, have been
associated with the induction and repression of light-responsive genes[12,13]. Light-dependent
enrichment of the acetylation pattern of H3 and H4 in the enhancer and promoter regions of
the pea plastocyanin locus *PetE* has been reported[14], and the hyperacetylation of the *PetE*
promoter is linked to the transcriptional activity of this gene[15]. Moreover, a reduction in H3
acetylation is associated with a decrease in the expression of the *A. thaliana* light-responsive
genes CHLOROPHYLL a/b-BINDING PROTEIN GENE (CAB2) and the RIBULOSE
BISPHOSPHATE CARBOXYLASE/OXYGENASE small subunit (RBCS)[16]. Histone
methylation regulates PHY-mediated seed germination in *A. thaliana*. Upon R light
illumination, photoactivated PHYB (Pfr) targets PIF1 for proteasome-mediated degradation,
releasing the expression of the JUMONJI HISTONE DEMETHYLASES JMJ20 and JMJ22.
As a result, JMJ20 and JMJ22 reduce the levels of H4R3me2, which leads to the activation
of the gibberellic acid biosynthesis pathway to promote seed germination[17].

Together with histone modification, DNA methylation is a common epigenetic mark with a
direct impact on gene expression. Nevertheless, only a few reports have specifically
addressed the effect of light stimuli on DNA methylation. Light-dependent nuclear
organization dynamics during deetiolation are associated with a reduction in methylated
DNA[18]. In *Populus nigra*, 137 genes were shown to be regulated by methylation during the
day/night cycle[19]. Moreover, photoperiod-sensitive male sterility is regulated by RNA-
directed DNA methylation (RdDM) in rice[20]. Finally, using a methylation-sensitive amplified
polymorphism assay, DNA methylation remodelling was shown to be an active epigenetic
response to different light qualities in tomato seedlings[21].

Here, we investigate the impact of DNA methylation on gene expression regulation in
response to PHY-dependent light perception in tomato fruits. The *Solanum lycopersicum*
genome harbours five PHY-encoding genes: PHYA, PHYB1, PHYB2, PHYE and PHYF[22].
Previous studies have shown that PHYA, PHYB1 and PHYB2 are major regulators of fruit
chloroplast maturation and nutraceutical compound accumulation[23–26]. In this work, the
genome-wide transcriptome and methylome were comprehensively analysed in fruits from
phyA and *phyB1B2* null mutants, revealing that PHY-dependent DNA demethylation is a
crucial stimulus for triggering ripening-associated gene expression. Moreover, our results
showed that gene-body (GB) RdDM is positively correlated with gene expression during fruit

development.

RESULTS

Impact of light perception impairment on the fruit transcriptome

To investigate the role played by either PHYA or PHYB1/PHYB2 (hereafter PHYB1B2) in

overall gene expression during fruit development, the transcriptome of fruits at the immature

green (IG) and breaker (BK) stages from *phyA* and *phyB1B2* null mutants as well as their

wild-type (WT) counterpart, was determined by RNAseq. Among the approximately 20,000

transcriptionally active loci in each biological replicate (Supplementary Table 1), 1.2% and

2.4% at the IG stage and 9.1% and 11.2% at the BK stage were identified as differentially

expressed genes (DEGs) in *phyA* or *phyB1B2* mutants, respectively, compared to the WT

(Fig. 1a; Supplementary Table 2). For both genotypes, the number of exclusive DEGs was

significantly lower in the IG stage than in the BK stage; similarly, the number of genes that

were commonly regulated by PHYA and PHYB1B2 was 172 at the IG stage and 785 at the

BK stage (Fig. 1b). Subsequently, the altered expression of approximately 76% (23/30) of

the tested genes was validated by RT-qPCR (Supplementary Table 3). Comparison with

previously reported expression data for genes involved in ripening regulation, ethylene

biosynthesis and signalling, and carotenogenesis further validated our RNAseq data, as 90%

of the analysed genes on average showed the expected mRNA profile at IG and BK stages.

It is worth mentioning that most of the genes displayed the same transcript fluctuation in the

WT, *phyA* and *phyB1B2* genotypes, though this was somewhat attenuated in the mutants

(Supplementary Table 4). These results showed that PHY-meditated light perception

regulates more genes in BK than in the early stages of fruit development and that PHYB1B2

has a more substantial impact than PHYA in the fruit transcriptome in both analysed stages.

A closer look at DEGs function revealed a similar distribution of loci across MapMan

categories in response to *phyB1B2* and *phyA* mutations in both developmental stages,

although with distinct abundance levels (Fig. 1c). At the IG stage, eight categories were

mainly represented, including at least 2% of the DEGs identified in *phyA* and *phyB1B2*:

photosynthesis, lipid metabolism, phytohormone action, RNA biosynthesis, protein

modification, protein homeostasis, cell wall organization, and solute transport (Fig. 1c;

Supplementary Tables 5 and 6). It is worth highlighting the abundance of the DEGs within

the photosynthesis category in the *phyB1B2* mutant, among which 34 out of the 37 genes

were downregulated (Supplementary Table 6). In the BK stage, at least 2% of the DEGs were

related to the lipid metabolism, phytohormone action, RNA biosynthesis, protein

modification and homeostasis, cell wall organization and solute transport categories in both

genotypes (Fig. 1c; Supplementary Tables 7 and 8). However, while *phyA* deficiency also

affected carbohydrate metabolism and external stimuli (Supplementary Table 7), the

phyB1B2 mutant showed a large number of DEGs related to the cell cycle and chromatin

organization (Supplementary Table 8). Interestingly, the chromatin organization category

displayed 52 DEGs, 45 of which were upregulated. These genes encode nucleosome

constituent histones (H3, H4, H2A and H2B); DNA methylases/demethylases; histone post-

translational modifiers such as deacetylases, methylases/demethylases, histone

ubiquitination factors and histone chaperones; chromatin remodelling factors; and genes

involved in RNA-independent and RNA-directed DNA methylation (Supplementary Table

8). These results led us to further investigate the impact of DNA methylation on PHY-

mediated gene expression reprogramming.

PHYs regulate the epigenome profile

The global profile of methylated cytosines (mCs) in the epigenome of tomato fruits was

assessed by whole-genome bisulfite sequencing in the IG and BK stages for *phyA*, *phyB1B2*

and WT genotypes. In agreement with previous reports[27,28], regardless of the genotype and

fruit stage, the greatest total number of mCs was located in the CHH context, followed by

the CG and CHG contexts, while the methylation level was highest in the CG (80%) context

followed by the CHG (67%) and CHH (23%) contexts (Supplementary Table 9,

Supplementary Fig. 1). For further comparisons, we selected only cytosines with coverage

>10X, and except for chromosome 9 in the transposable elements TEs enriched region, all

samples met this cutoff. In all contexts, the highest cytosine density was associated with

gene-rich euchromatic regions located at chromosome arm ends (Supplementary Fig. 1).

Conversely, in symmetrical contexts (CG and CHG), the highest methylation rates were

found across pericentromeric regions enriched in TEs and in the CHH context in gene-rich

regions associated with a higher density of sRNAs (Supplementary Fig. 1). The comparison

of the methylation status between the two fruit stages showed that ripening-associated

demethylation[27] occurs mainly in the CG context, especially in gene-rich regions, and that it

is impaired in *phyB1B2* mutant BK fruits (Supplementary Fig. 1).

The subsequent comparison between genotypes revealed global epigenome alteration in *phy*

mutants in all contexts analysed. The most remarkable observation was the presence of

considerable hypermethylation in all contexts across gene-rich regions in BK-stage fruits

from *phyB1B2* (Fig. 2a). In contrast, *phyA* exhibited hypermethylation in CHG and CHH

contexts associated with TE-rich regions (Fig. 2a), suggesting that different PHYs control

DNA methylation across specific genomic regions through distinct regulatory mechanisms.

Interestingly, PHY-associated hypomethylation was exclusively detected in the CG context

of gene-rich regions in IG-stage fruits from *phyA* and in the CHH context of TE-rich regions

for BK-stage fruits from *phyB1B2*. In summary, these data revealed that both PHYA and

PHYB1B2 regulate the global methylome, but PHYB1B2 has a greater impact on ripening-

associated methylation reprogramming across gene-rich genomic regions in tomato fruits.

To investigate the relationship between PHY-regulated cytosine methylation and gene

expression, we first identified genes with differentially methylated promoters (DMPs, 2 kb

upstream of the transcription start site) in all three contexts. Interestingly, associated with the

massive alteration previously observed, the pattern of DMPs varied with the mC context,

stage and genotype (Fig. 2b, Supplementary Table 10 and 11). Regarding the CG context,

whereas the *phyA* mutant showed virtually the same frequency of hyper- and hypomethylated

promoters in the two stages, the status of hypermethylated promoters in *phyB1B2* increased

over 60% from the IG to BK stage, while the number of loci with hypomethylation decreased

50% (Fig. 2b, Supplementary Table 12). In contrast, *phyA* showed a greater number of

hypermethylated promoters in the CHG context in the IG stage than in the BK stage, while

the levels in the WT and *phyB1B2* mutant remained similar upon ripening (Fig. 2b,

Supplementary Table 13). In the CHH context, the number of hypermethylated promoters

decreased in both genotypes from the IG to BK stages (Fig. 2b, Supplementary Table 14).

These results indicate that PHY deficiency results in massive promoter hypermethylation in

both the IG and BK stages of tomato fruit development. Moreover, they reinforce the role of

PHYB1B2 in ripening-associated demethylation and its putative effect on gene expression.

Effect of PHY-mediated differential methylation on the transcriptome

To assess whether the differential methylation of gene promoters affects mRNA levels, we crossed data from DEGs and DMPs between genotypes. Supplementary Fig. 2 shows scatter plots of promoter methylation vs mRNA fold changes for comparisons of the two genotypes at the two examined developmental stages in the three mC contexts. The most evident result was that among the thousands of loci with identified DMPs (Fig. 2b), only hundreds of the loci were also differentially expressed (Supplementary Table 15) (0.7% for IG *phyA*, 1.6% for IG *phyB1B2*, 5.6% for BK *phyA* and 7.4% for BK *phyB1B2*), raising an intriguing question about the biological significance of the extensive change in the methylation pattern observed in the mutants. In contrast, the percentages of the DEGs showing DMPs were 73% for IG *phyA*, 76% for IG *phyB1B2*, 72% for BK *phyA* and 75% for BK *phyB1B2* (Supplementary Fig. 2). Many more DEGs with DMPs were observed in BK than in IG fruits and in *phyB1B2* than in the *phyA* genotype. The functional categorization of these genes revealed a similar category distribution to the DEGs (Fig. 1c, Supplementary Table 16-19). At the IG stage, there were seven categories in which at least 2% of the loci showed DMPs and differential expression in both genotypes: photosynthesis, phytohormone action, RNA biosynthesis, protein modification and homeostasis, cell wall organization and solute transport, whereas *phyB1B2* additionally impacted lipid metabolism (Fig. 1c). In the BK stage, the categories in which at least 2% of the DEGs showed DMPs were lipid metabolism, phytohormone action, RNA biosynthesis, protein modification and homeostasis, cell wall organization and solute transport-related functions in both genotypes, while only *phyA* impacted carbohydrate metabolism and external stimuli, and only *phyB1B2* affected photosynthesis, chromatin organization and cell cycle categories.

Interestingly, in the comparison of the IG and BK stages, 42.5%, 34.2% and 18.8% of the DMPs were associated with DEGs, while 79.5%, 76.6% and 71.5% of the DEGs showed differences in promoter methylation in WT, *phyA* and *phyB1B2*, respectively (Supplementary Fig. 3). All of these data demonstrated that the PHY-dependent mRNA profile is profoundly affected by promoter methylation; however, massive genome-wide PHY-induced methylation reprogramming has a still uncharacterized role beyond the regulation of mRNA accumulation. Moreover, promoter methylation has a greater effect on gene expression regulation during BK than in the IG stage. Additionally, the data showed that PHYB1B2 has

240   a more extensive influence on gene expression regulated via promoter methylation than

241   PHYA, reinforcing the above conclusion that PHYB1B2 affects CG ripening-associated

242   demethylation (Supplementary Fig. 2).

244   The sRNAome is altered by PHY deficiency

To assess the involvement of RdDM in PHY-mediated transcriptome regulation, the

sRNAome was analysed in fruits at the IG and BK stages from both mutants and the WT

genotype (Supplementary Table 20a). A total of 28,314 clusters of sRNAs were identified

across the whole genome in at least one of the samples, including 7,984 in gene bodies, 7,863

in promoter regions, 7,966 in TEs and the remaining 4,501 across intergenic regions

(Supplementary Fig. 1, Supplementary Table 20b). The methylation level was evaluated for

each sRNA cluster-targeted genomic region (sCTGR), and as previously observed for

promoter regions, a higher proportion of hypermethylation was observed in BK fruits from

phyB1B2 in the CG symmetrical context. Moreover, the greatest number of differentially

methylated sCTGRs was observed in the asymmetrical context CHH. (Fig. 3a,

Supplementary Table 20g-j).

sCTGR methylation levels and sRNA accumulation data were intersected, and among a total

of 154, 318, 267 and 257 differentially accumulated sRNA clusters (Supplementary Table

20c-f), 88, 154, 99 and 82 also showed differential methylation changes (>5%) in *phyA* IG,

phyB1B2 IG, *phyA* BK and *phyB1B2* BK fruits, respectively (Fig. 3b, (Supplementary Table

20g-j), showing a strong association ($P<0.005$) of the two datasets. Intriguingly, this positive

association was not observed in the transition from the IG to BK stages (Supplementary Fig.

4), suggesting that the global methylation changes via RdDM could be attributed to PHY

deficiency. Moreover, a clear disturbance in sRNA accumulation was observed in *phyB1B2*,

since almost no clusters with less sRNA accumulation were observed in BK compared to the

IG stage (Supplementary Fig. 4). We further analysed whether this positive correlation

observed between sRNA accumulation and sCTGR methylation impacted gene expression

levels. Notably, regardless of the fruit developmental stage, changes in the accumulation of

sRNA located in gene bodies (GBs), and not in the promoter (P) region, were positively

correlated with the mRNA level (Fig. 3c, Supplementary Table 20k-n). Among these loci,

two interesting examples were identified: the well-known ripening-associated genes

RIPENING INHIBITOR (*RIN*, Solyc05g012020[29]) and *FRUITFULL2* (*FUL2*,

Solyc03g114830[30]), which showed higher expression in *phyB1B2* at the IG stage (Fig. 4a)

and higher sRNA accumulation and sCTGR methylation across their GBs (Fig. 4b) compared

to WT. The premature expression of these TFs was in agreement with the previously reported

anticipation of ripening onset in the *phyB1B2* mutant[31]. Altogether, these findings revealed

(i) impaired RdDM in BK fruits of *phyB1B2*, indicated by the absence of clusters with less

sRNA accumulation, and (ii) that GB RdDM is an important mechanism that positively

regulates gene expression in a PHY-mediated manner during fruit development.

PHYB1B2-dependent methylation regulates fruit chlorophyll accumulation

The categorization of DEGs associated with differential promoter methylation revealed

prominent representation of the photosynthesis category in the fruits of the *phyB1B2* mutant

at the IG stage (Fig. 1c). Among the 32 genes, 22 were downregulated and hypermethylated

in the promoter region (Supplementary Tables 6 and 17). Most of these genes encode

chlorophyll-binding proteins, structural photosystem proteins and chlorophyll biosynthetic

enzymes. This might at least partly explain the reduction of 50% in the total chlorophyll level

observed in *phyB1B2* IG fruits (Fig. 5a). Detailed promoter analysis of the chlorophyll-

related enzymes *PROTOCHLOROPHYLLIDE OXIDOREDUCTASE 3* (*POR3,*

Solyc07g054210), and two CHLOROPHYLL A/B BINDING PROTEINs (*CBP*,

Solyc02g070990 and *CAB-3c*, Solyc03g005780) showed that the reduced mRNA levels of

these three genes in *phyB1B2* (Fig. 5b) were correlated with the presence of hypermethylated

regions in the promoters. Interestingly, several of these hypermethylated sequences

overlapped with HY5 and PIF photomorphogenic TF binding sites, such as G-box

(CACGTG), CA hybrid (GACGTA), CG hybrid (GACGTG) and PBE-box (CACATG)

motifs[32] (Fig. 5c). These results suggest that the transcription of genes involved in

chlorophyll metabolism and the photosynthetic machinery in tomato fruits is regulated by the

PHYB1B2-mediated methylation status of their promoter regions in addition to the PHY-

mediated post-translational regulation of HY5 and PIF protein levels[2].

The methylation-mediated regulation of fruit ripening is PHYB1B2-dependent

In their seminal study, Zhong *et al*. (2013)[27] revealed that the extensive methylation in the
promoter regions of ripening-associated genes gradually decreases during fruit development.
Interestingly, RNA biosynthesis, which includes transcription factors, was the most abundant
functional category among the DEGs that showed DMPs (Fig. 1c). Thus, we examined a set
of ripening-associated master transcription factors: *RIN*, Solyc05g012020, *NON-RIPENING*
(*NOR*, Solyc10g006880[33]), *COLORLESS NORIPENING* (*CNR*, Solyc02g077920[34]) and
APETALA2a (*AP2a*, Solyc03g044300[35]). The evaluation of the promoter regions clearly
showed that while their methylation level decreased from the IG to BK stage in the WT
genotype, they remained highly methylated in *phyB1B2* (Fig. 6a). As previously observed,
several hypermethylated regions are closely linked to HY5 and PIF binding sites. The
maintenance of high methylation levels in the promoters of these key regulatory genes at the
onset of fruit ripening was highly correlated with their transcriptional downregulation at the
BK stage (Fig. 6b).
Carotenoid accumulation is probably the most appealing and best investigated trait of tomato
fruits; in agreement with previous findings[26], ripe *phyB1B2* fruits showed a five-fold
reduction in carotenoid content compared to WT (Fig. 7a).
With the aim of evaluating whether this effect is a consequence of the methylation-mediated
regulation of carotenoid biosynthesis genes, we further analysed the promoters of
PHYTOENE SYNTHASE 1 (*PSY1,* Solyc03g031860), *PHYTOENE DESATURASE* (*PDS,*
Solyc03g123760), *15-CIS- ζ-CAROTENE* (*ZISO,* Solyc12g098710) and *ZETA-CAROTENE*
DESATURASE (*ZDS,* Solyc01g097810), which, with the exception of *PDS*, were
hypermethylated in *phyB1B2* BK fruits (Supplementary Table 11). The mC profile confirmed
the presence of hypermethylated regions in all four promoters, which were predominantly
located near light-dependent TF binding motifs in the *PSY1*, *PDS* and *ZISO* promoters (Fig.
7b), explaining the reduced mRNA levels of these genes observed in *phyB1B2* (Fig. 7c).
RIN is one of the main TFs controlling ripening-associated genes by directly binding to their
promoters. RIN binding occurs in concert with the demethylation of its targets[27]. To examine
whether RIN binding site methylation could be affected by the *phyB1B2* mutation in the
ripening-related master transcription factors and carotenoid biosynthesis gene promoters, we
mapped the available RIN ChIP-seq data[27] and performed *de novo* motif discovery
(Supplementary Fig. 5). Interestingly, the levels of mCs associated with RIN targets were

higher in the *NOR*, *CNR* and *AP2a* promoters in *phyB1B2* than in WT. Moreover, the *RIN*

promoter itself was hypermethylated across the RIN binding site in *phyB1B2* BK fruits,

suggesting a positive feedback regulatory mechanism (Fig. 6a). Finally, in the *phyB1B2*

mutant, the *PSY1*, *PDS*, *ZISO* and *ZDS* promoters showed higher methylation overlapping

with RIN target binding motifs, indicating that the upregulation of carotenoid biosynthesis

genes during tomato ripening is dependent on the PHYB1B2-mediated demethylation of RIN

target sites.

Altogether, our findings showed that PHYB1B2 is a major player in fruit ripening by

controlling the promoter demethylation of master transcriptional regulators and carotenoid

biosynthesis genes.

DISCUSSION

The dynamic methylation pattern during tomato fruit development has been demonstrated to

be a critical ripening regulation mechanism[27,28]. DNA demethylation, mainly in the CG

context, triggers the activation of genes involved in ripening and is required for pigment

accumulation and ethylene synthesis[27,36]. Simultaneously, the dynamic epigenome during

fruit development is strictly regulated by environmental cues[37]. The prevailing model

establishes PHYs as major components involved in the coordination of fruit physiology with

the ever-changing light and temperature environment[23,26]. Thus, we explored the link

between fruit epigenome reprogramming and these well-established light and temperature

sensors.

Our data clearly showed that *phyA* and *phyB1B2* deficiencies modified the epigenome profile

through methylome and sRNAome reprogramming. In particular, PHY-mediated DMPs and

GB methylation impacted the transcriptome pattern, affecting tomato fruit development and

demonstrating that a portion of the ripening-associated demethylation previously reported[27]

is dependent on active PHYs, especially PHYB1B2. However, the massive alteration of

methylation patterns observed in PHY mutants suggests the existence of a still unclear

genome regulatory mechanism. Whether it is related to light-induced AS[3] and/or PHY-

mediated alternative promoter usage to control protein localization[8], as mentioned above,

remains to be explored.

The *phyA* and *phyB1B2* mutants showed a positive correlation between cluster sRNA

accumulation, target methylation in GB and mRNA levels. In angiosperms, GB methylation

has been associated with constitutively expressed genes[38,39]; however, PHY deficiency

intriguingly seems to deregulate this mechanism, affecting temporally and spatially regulated

genes. The *RIN* and *FUL2* examples analysed here clearly showed that sRNA accumulation

and methylation were mainly located near transposable elements (TEs) (Fig. 4). It is known

that the insertion of TEs within GB can disrupt gene expression; thus, methylation-mediated

TE silencing and GB methylation are evolutionarily linked[38]. The enhancement of TE-

associated DNA methylation in GB (Fig. 3c) and the absence of clusters with less sRNA

accumulation in BK compared to the IG stage in *phyB1B2* (Supplementary Fig. 4) can be

explained by the overexpression of canonical RdDM genes. Solyc12g008420 and

Solyc06g050510 encode homologs of RNA-DEPENDENT RNA POLYMERASE (RDRP)

and the associated factor SNF2 DOMAIN-CONTAINING PROTEIN CLASSY 1 (CLSY1),

respectively, both of which were upregulated in BK fruits from *phyB1B2* plants

(Supplementary Table 2). These proteins are key players in canonical RdDM in *A. thaliana*

required for PolIV-dependent sRNA production[40]. Similarly, Solyc09g082480 and

Solyc03g083170, which were also upregulated in *phyB1B2* BK fruits, are homologs of *A.*

thaliana RNA-DIRECTED DNA METHYLATION 1 (RDM1) and DEFECTIVE IN

MERISTEM SILENCING 3 (DMS3), respectively. The protein products of these genes,

together with DEFECTIVE IN RNA- DIRECTED DNA METHYLATION 1 (DRD1), form

the DDR complex, which enables RNA Pol V transcription[41]. To our knowledge, this is the

first report to associate PHY-mediated sRNA accumulation and DNA methylation with

mRNA levels in plants.

Several pieces of evidence have shown that PHYB1B2 has a more substantial impact on

genome regulation than PHYA. For example, BK fruits from *phyB1B2* displayed (i) a large

number of DEGs associated with chromatin organization (Fig. 1c); (ii) overall promoter

hypermethylation in the CG context (Fig. 2b); (iii) the highest number of DEGs associated

with DMPs (Supplementary Fig. 2); and (iv) half the number of DMPs associated with DEGs

between the IG and BK stages compared to the WT (Supplementary Fig. 3). These massive

epigenomic alterations in *phyB1B2* led us to look more closely at genes related to chromatin

organization that showed alterations in mRNA levels.

The chromomethylase *SlMET1L* (Solyc01g006100) (also referred to as *SlCMT3*[42]) displays
the highest transcript abundance in immature fruits, which declines towards the fully ripe
stage[43]. Hence, in line with the higher level of DNA methylation, our transcriptome analysis
showed that *SlMET1L* was upregulated in *phyB1B2* BK fruits. Conversely, the upregulation
of the *SlROS1L* demethylase (Solyc09g009080[43]; also referred to as *SlDML1*[44]) was also
observed in *phyB1B2* BK fruits, which might seem contradictory at first glance. However, it
has been reported that the *Arabidopsis thaliana ROS1* gene promoter contains a DNA
methylation monitoring sequence (MEMS) associated with a Helitron transposon, which is
methylated by AtMET1, positively regulating *AtROS1* gene expression[45]. Similarly,
SlROS1L harbours two transposable elements within its promoter and showed a higher
methylation level in *phyB1B2* than in the WT genotype, suggesting a similar regulatory
mechanism in tomato (Supplementary Fig. 6, Supplementary Table 15).
The tomato homologue of *A. thaliana* DECREASED DNA METHYLATION 1 (DDM1,
Solyc02g085390) showed higher mRNA expression in *phyB1B2* mutant BK fruits than in
their WT counterparts. DDM1 is a chromatin remodelling protein required for maintaining
DNA methylation in the symmetric cytosine sequence[46], which can be associated with the
observed methylation differences in the CG context in *phyB1B2* (Fig. 2a).
Several histone modifiers showed altered expression in BK fruits from the *phyB1B2* mutant
(Supplementary Table 8). The methylation of lysine residues 9 and 27 on histone H3 is
associated with repressed genes. Histone lysine methyltransferases are classified into five
groups based on their domain architecture and/or differences in enzymatic activity[47]. The BK
fruits of the *phyB1B2* mutant displayed three differentially expressed lysine
methyltransferases: Solyc03g082860, an upregulated H3K27 Class IV homologue, and
Solyc06g008130 and Solyc06g083760, two H3K9 Class V homologs that showed lower and
higher expression than in WT fruits, respectively. Histone arginine methylation is catalysed
by a family of enzymes known as protein arginine methyltransferases (PRMTs).
Solyc12g099560, a PRMT4a/b homologue, was upregulated in *phyB1B2* BK fruits.
Interestingly, in *A. thaliana*, PRMT4s modulate key regulatory genes associated with the
light response[48], reinforcing the link between the PHYB1B2 photoreceptor and epigenetic
control. Finally, tomato histone demethylases have been recently identified. *SlJMJ6*, whose
expression peaks immediately after the BK stage, has been characterized as a positive

regulator of fruit ripening by removing the H3K27 methylation of ripening-related genes,

and *SlJMJ6*-overexpressing lines show increased carotenoid levels[49]. *SlJMJC1*

(Solyc01g006680), which exhibits the same expression pattern[49], is downregulated in the

phyB1B2 mutant, suggesting that this gene might exhibit a similar regulatory function to its

paralogue, inducing ripening in a PHYB1B2-dependent manner (Fig. 6 and 7).

Histone deacetylation plays a crucial role in the regulation of eukaryotic gene activity and is

associated with inactive chromatin[40]. Histone deacetylation is catalysed by histone

deacetylases (HDACs). Fifteen HDACs were identified in the tomato genome[50]. Among

these HDACs, *SlHDA10* (Solyc01g009120) and *SlHDT3* (Solyc11g066840) were found to

be downregulated and upregulated in *phyB1B2* BK fruits, respectively. SlHDA10 is localized

in the chloroplast, and its transcript is highly expressed in photosynthetic tissues[50]; whether

SlHDA10 deacetylates chloroplast proteins by silencing photosynthesis-related genes

remains to be determined. Although *SlHDT3* is mainly expressed in immature stages of fruit

development and its expression declines with ripening, its silencing results in delayed

ripening and reduced *RIN* expression and carotenogenesis. On the other hand, the expression

level of *SlHDT3* is increased in ripening-deficient mutants such as *Nr* or *rin*[51]. Our results

showed that *phyB1B2* mutant fruits displayed higher expression of *SlHDT3* and reduced *RIN*

transcript levels at the BK stage, suggesting reciprocal regulation between these two factors

(Fig. 8a). During the IG stage, *SlHDT3* is highly expressed, contributing to the epigenetic

inhibition of ripening. The reduction in *SlHDT3* expression towards BK releases DNA

methylation and both directly and indirectly upregulates *RIN*. Additionally, PHYB1B2

inhibits the expression of *SlHDT3* and induces that of *RIN*, thereby inducing or repressing

DNA methylation, respectively. Finally, RIN maintains *SlHDT3* downregulation in BK[51].

This mechanism precisely tunes ripening-related epigenetic reprogramming and contributes

to explaining the high methylation levels observed in the *phyB1B2* mutant (Fig. 2).

The link between DNA methylation levels and tomato fruit ripening-associated gene

expression has been previously reported[27,28], but the stimuli and the molecular mechanisms

underlying this relationship remained unknown. The integrated analysis of the experimental

evidence together with previous gene functional studies in tomato and *A. thaliana* allowed

us to propose that PHYB1B2 is an important triggering factor for chromatin remodelling and,

consequently, transcriptional regulation during fruit development. PHY signal transduction

is mediated by the coordinated expression of DNA methylases/demethylases, histone-

modifying enzymes and chromatin remodelling factors resulting in the induction of

photosynthesis and ripening-related genes in immature and breaker fruit stages, respectively

(Fig. 8b). The vast reservoir of data released here brings a new level of understanding about

how epigenetic mechanisms orchestrate the response to light and temperature fluctuations

affecting important agronomical traits in fleshy fruits.

MATERIAL AND METHODS

Plant material, growth conditions and sampling

phyA and *phyB1B2* phytochrome mutants in the *Solanum lycopersicum* (cv. MoneyMaker)

genetic background were previously characterized[52–54]. Tomato seeds were grown in 9L pots

containing a 1:1 mixture of commercial substrate and expanded vermiculite, supplemented

with 1 g L^{-1} of NPK 10:10:10, 4 g L^{-1} of dolomite limestone ($MgCO_3$ + $CaCO_3$) and 2 g L^{-1}

thermophosphate at 24/18 °C under a 16/8 h light/dark cycle under 230 - 250 µmol photons

m^{-2} s^{-1} irradiation and a relative humidity of 55%. Five replicates per genotype were

cultivated. Fruits were sampled at the immature green (15 mm diameter), mature green (when

the placenta displays a gelatinous aspect), breaker (beginning of ripening process when the

fruit shows the first yellowish colouration) and red ripe (7 days after the breaker stage) stages.

All fruits were harvested at the same time of day with four biological replicates (each

replicate was composed of a single fruit per plant). The columella, placenta, and seeds were

immediately removed, and the remaining tissues were frozen in liquid nitrogen, ground and

freeze-dried for subsequent analysis.

Transcriptional profile

Total RNA was extracted from immature green and breaker stage fruits with three

independent biological replicates of each genotype using a Promega ReliaPrep RNA tissue

kit according to the manufacturer's instructions. The RNA concentration was determined

with a spectrophotometer (Nanodrop ND-1000; NanoDrop Technologies, Wilmington, DE,

U.S.A.), RNA quality was assessed with a BioAnalyzer 2100 (Agilent Technologies), and

RNA libraries were constructed following the recommendations of an Illumina Kit

(Directional mRNA-Seq Sample Preparation) and sequenced using the Illumina NovaSeq

6000 System. Each library was sequenced, generating approximately 20 million 150 bp

paired end reads per sample. The raw sequencing reads that were generated were analysed

with FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and were filtered

and cleaned using Trimmomatic[55] (Parameters: ILLUMINACLIP: TruSeq3-

PE.fa:2:30:10LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:50). At least

95% (19.1-27.9 M) of the reads met the quality criteria and were mapped to the tomato

reference genome sequence SL3.0 with the ITAG3.2 annotation using STAR v2.4.2.

allowing one mismatch[56], approximately 84% of the reads were uniquely mapped

(Supplementary Table 1) and were used for statistical analysis.

497   Reverse transcription quantitative PCR (RT-qPCR)

Total RNA extraction was performed with the ReliaPrep™ RNA Cell and Tissue Miniprep

System (Promega), and cDNA synthesis was conducted with SuperScript™ IV Reverse

Transcriptase (Invitrogen). The primers used for qPCR are listed in Supplementary Table 21.

RT-qPCR was performed in a QStudio6 – A1769 PCR Real-Time thermocycler using 2X

Power SYBR Green Master Mix in a final volume of 10 µL. Absolute fluorescence data were

analysed using LinRegPCR software to obtain Ct and primer efficiency values. Relative

mRNA abundance was calculated and normalized according to the ΔΔCt method using

EXPRESSED and *CAC* as reference genes[57].

507   MethylC-Seq analysis

Methyl C sequencing was performed as described in a previous report[58]. gDNA (~5 g) was

extracted from a pool of the same three biological replicates used in the transcriptome

analyses, obtained from three IG and BK fruit samples per genotype, using the DNeasy Plant

maxi kit (Qiagen). The libraries were prepared with the EZ DNA Methylation-Gold Kit

(Zymo Research) and the Accel-NGS® Methyl-Seq DNA Library Kit (Swift Biosciences)

and further sequenced using the Illumina NovaSeq 6000 platform. Over 240 M reads were

sequenced from each genotype and stage. Raw reads were screened for quality using

Trimmomatic[55] (parameters: ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 LEADING:3

TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:50). Mapping to the tomato reference

genome sequence SL3.0 and the assessment of global methylation status were performed

using Bismark[59] (parameters: bismark -q --bowtie2 --non_directional -N 1 -p 4), and the

methylation status of DNA in the three possible contexts (CG, CHG and CHH) was

distinguished. At least 130 M reads were uniquely mapped (Supplementary Table 9). The

Bioconductor package methylKit[60] was used for the detection of methylation levels across

the analysed regions: promoters (2 kb upstream of transcription start site) and sRNA cluster-

targeted genome regions (sCTGRs). Only Cs with 10X coverage were considered.

Methylation differences with an FDR < 0.05 in each comparison (WT vs *phyA*; WT vs

phyB1B2) were recorded as differentially methylated promoters (DMPs) or differentially

methylated sCTGRs. Differential methylation in the CG, CHG and/or CHH context was

considered to exist if the region contained at least 10 differentially methylated Cs in the

corresponding context. Finally, for the comparison of global methylation levels between

genotypes, only common Cs with at least 10X coverage in all samples were analysed.

sRNAome profile

sRNA extraction and quality parameters were determined from three independent biological

replicates of green and breaker stage fruits from each genotype, as described above in the

"Transcriptional profile" section. After RNA integrity confirmation, libraries were prepared

using a TruSeq Small RNA Library Prep and sequenced using the Illumina HiSeq 4000

platform to generate a read length of 50 bp. The raw sequencing reads that were generated

were quality trimmed with Trimmomatic[55] to retain reads of 18-24 nt in length (parameters:

ILLUMINACLIP:TruSeq3-SE:2:30:10   LEADING:3   TRAILING:3

SLIDINGWINDOW:4:15   MINLEN:18   AVGQUAL:25). A   minimum   of   38%

(WT/breaker/A) and a maximum of 85% (WT/immature green/A) of the reads achieved the

quality criteria and were used for further analyses (Supplementary Table 20a). All libraries

were aligned to genome version SL3.0 using ShortStack v3.8.1[61] with default parameters

(allowing the distribution of multimapping reads according to the local genomic context).

Then, the *de novo* identification of clusters of sRNAs was performed for all libraries, and

individual counts for each library and cluster were obtained using the same software.

Statistical analysis for RNAseq and sRNAome

Genes/sRNA clusters with read/count numbers smaller than two per million were removed.

Read/count values were normalized according to the library size factors. Statistical analyses

were performed with edgeR from Bioconductor®[62,63] using a genewise negative binomial

generalized linear model with the quasi-likelihood test[64] and a cutoff of the false discovery

rate (FDR) ≤ 0.05.

Gene functional categorization

The DEGs were functionally categorized with MapMan application software[65] followed by

hand-curated annotation using MapMan categories.

***In silico* regulatory motif predictions and RIN ChIP-seq analyses**

To predict the transcription factor binding motifs, we used the complete collection of 530

plant transcription factor-binding sites (TFBS) modelled as position frequency matrices

(PFMs) from the JASPAR 2020 database[66]. Putative binding sites were obtained by scanning

the whole *S. lycopersicum* 3.00 genome with Fimo[67], p-value < 1e−5.

RIN ChIPseq reads were downloaded from the Sequence Read Archive (SRA) (accession

SRX15083[27]) and mapped to tomato genome version SL3.0 with STAR[56] (version 2.7.3X,

parameters: outFilterMismatchNmax 3, alignEndsType EndToEnd, alignIntronMax 5), and

peak calling was performed by using Macs2[68] (version 2.2.7.1, default parameters). Regions

of 200 bp centred on the top-scoring peaks (score>100, n=327) among transposable elements

were retrieved, and the binding motif was inferred *de novo* by using the MEME algorithm

(Supplemental Fig. 5) [67].

Carotenoid and chlorophyll analysis

Carotenoid and chlorophyll extraction was performed from aliquots of 20 mg dry mass,

sequentially with 100 µL of a saturated solution of NaCl, 200 µL of dichloromethane and

500 µL of hexane:diethyl ether 1:1 (v/v). The pellet was extracted three additional times with

500 µL of 1:1 (v/v) hexane:diethyl ether. All supernatant fractions were combined,

completely dried by vacuum and suspended in 200 µL of acetonitrile. Chlorophyll, phytoene,

phytofluene, lycopene, β-carotene and lutein levels were determined via HPLC with a

photodiode array detector[69].

Statistical analysis of RT-qPCR and metabolites

Statistical analyses of the RT-qPCR (Student's t-test, $p \leq 0.05$) and metabolic data (ANOVA, Tukey's test. $p \leq 0.05$) were performed with InfoStat/F software (http://www.infostat.com.ar).

Figure legends

Fig. 1. PHYA and PHYB1B2 modify the global transcriptomic profile of tomato fruit. (a) Number of differentially expressed genes (DEGs) in *phyA* and *phyB1B2* mutant fruits at immature green (IG) and breaker (BK) stages. (b) Venn diagram showing exclusive and common DEGs in *phyA* and *phyB1B2* mutants in both developmental stages. (c) Functional categorization of all DEGs and those DEGs with differentially methylated promoters (DMPs) in both analysed genotypes and stages. Only categories corresponding to at least 2% of the DEGs or DMPs in each comparison are shown (asterisks). Up- and downregulated genes are indicated in red and blue, respectively. Loci with hyper- and hypomethylated promoters are indicated in light red and light blue, respectively. DEGs and DMPs show statistically significant differences (FDR < 0.05) relative to WT.

Fig. 2. PHYA and PHYB1B2 regulate the tomato fruit epigenome. (a) Density plot of genes, transposable elements (TEs) and mC in all contexts (mCG, mCHG, mCHH) for the WT genotype. Global methylation changes for *phyA* and *phyB1B2* in comparison with the wild type (WT) at the immature green (IG) and breaker (BK) stages are shown (bin size, 1 Mb). Gene and TE densities were estimated according to the number of nucleotides covered per million. The methylation levels in the CG, CHG and CHH contexts are 40-90%, 25-80% and 10-30%, respectively. The mC difference was relative to the corresponding WT fruit stage within a -5% (hypomethylated) ≤ range ≤ +5% (hypermethylated). (b) Number of genes with differentially methylated promoters (DMPs, 2 kb upstream transcription start site) in *phyA*, *phyB1B2* and both mutants. Hyper- and hypomethylation are indicated by grey and darker-coloured bars, respectively. DMPs show statistically significant differences (FDR < 0.05) relative to WT.

Fig. 3. Phytochrome deficiency impacts the sRNAome profile. (a) Total number of differentially methylated sRNA cluster-targeted genome regions (sCTGRs). (b) Scatter plots

show the relationship between the differential accumulation of cluster sRNAs and a
minimum of 5% differential methylation of their sCTGRs. The result of Fischer´s test for the
association of the two datasets is shown ($p \leq 2.07e^{-5}$). (c) Boxplots show changes in the
accumulation of cluster sRNAs in promoter (P, 2 Kb upstream of the 5' UTR end) and gene
body (GB) regions for up- and downregulated DEGs. Asterisks indicate statistically
significant differences by the Wilcoxon–Mann–Whitney test (** $p<0.0001$). All results
represent the comparison of *phyA* and *phyB1B2* to the wild type in immature green (IG) and
breaker (BK) fruit stages.

Fig. 4. Methylation across promoter and gene body regions differentially affects gene
expression. (a) Relative expression of *RIPENING INHIBITOR* (*RIN*) and *FRUITFULL 2*
(*FUL2*) in immature green (IG) and mature green (MG) fruits from *phyB1B2* determined by
RT-qPCR. Red dots indicate data from RNA-seq in the same stage. Expression levels
represent the mean of at least three biological replicates and are relative to the wild type
(WT). Asterisks indicate statistically significant differences by the two-tailed Student´s *t* test
compared to WT (* $p<0.05$). (b) Differential gene body methylation (green bars) and sRNA
accumulation (black bars) within *RIN* and *FUL2* in IG fruits from the *phyB1B2* and WT
genotypes.

Fig. 5. PHYB1/B2-dependent methylation regulates fruit chlorophyll. (a) Relative content of
total chlorophyll in IG fruits from *phyB1B2* and WT genotypes. Values represent the mean
of at least three biological replicates. Asterisks indicate statistically significant differences by
the two-tailed Student's *t* test between genotypes (** $p<0.01$). (b) Relative expression of
CHLOROPHYLL A/B BINDING PROTEINs (*CBP* and *CAB-3c*) and
PROTOCHLOROPHYLLIDE OXIDOREDUCTASE 3 (*POR3*) in IG fruits from *phyB1B2*
determined by RNA-seq. Expression levels represent the mean of at least three biological
replicates and are relative to WT. Asterisks indicate statistically significant differences
compared to WT (* FDR$\leq$0.05). (c) Differential promoter methylation in *CBP*, *CAB-3c* and
POR3 in IG fruits from the *phyB1B2* and WT genotypes. HY5 and PIF transcription factor
binding motifs are denoted with arrows.

Fig. 6. PHYB1/B2-dependent methylation regulates master ripening-associated transcription
factors. (a) Differentially methylated promoters of the *RIPENING INHIBITOR* (*RIN*), *NON-*
RIPENING (*NOR*), *COLORLESS NORIPENING* (*CNR*) and *APETALA 2a* (*AP2a*) loci

between the *phyB1B2* and wild-type (WT) genotypes. Green and orange indicate total mC in

immature green (IG) and breaker (BK) fruits, respectively. HY5 and PIF transcription factor

binding motifs are denoted with arrows. Thick blue lines indicate RIN binding sites according

to ChIP-seq data[27]. (b) Relative expression from the RT-qPCR assay of genes encoding

master ripening transcription factors in BK and red ripe (RR) fruits from *phyB1B2*. Red dots

indicate data from RNA-seq in the same stage. Expression levels represent the mean of at

least three biological replicates and are relative to WT. Asterisks indicate statistically

significant differences by two-tailed Student's *t* test compared to WT (* $p<0.05$).

Fig. 7. PHYB1/B2-dependent methylation regulates fruit carotenogenesis. (a) Relative

contents of total carotenoids in red ripe (RR) fruits from phyB1B2 and wild-type (WT)

genotypes. Values represent the mean of at least three biological replicates. Asterisks indicate

statistically significant differences by the two-tailed Student's *t* test between genotypes (**

$p<0.01$). (b) Differentially methylated promoter sites of the *PHYTOENE SYNTHASE 1*

(*PSY1*), *PHYTOENE DESATURASE* (*PDS*), *15-CIS- ζ-CAROTENE* (*ZISO*) and *ZETA-*

CAROTENE DESATURASE (*ZDS*) loci between the *phyB1B2* and WT genotypes. Orange

colour indicates total mC in breaker (BK) fruits. Arrows denote HY5 and PIF transcription

factor binding motifs. Thick blue lines indicate RIN binding sites according to ChIP-seq

data[27]. (c) Relative expression of carotenoid biosynthetic enzyme-encoding genes in

immature green (IG), mature green (MG), BK and RR fruits from *phyB1B2* determined by

RT-qPCR. Red dots indicate data from RNA-seq in the same stage. The expression levels

represent the mean of at least three biological replicates and are relative to WT. Asterisks

indicate statistically significant differences by the two-tailed Student's *t* test compared to WT

(*$p<0.05$, ** $p<0.01$).

Fig. 8. Model of epigenetic mechanisms involved in fruit development regulation triggered

by PHYB1B2. (a) PHYB1B2 contributes to the reciprocal regulation between HDT3 and

RIN through the inhibition and induction of *HDT3* and *RIN* expression, respectively.

Increases in *RIN* expression during fruit maturation inhibit *HDT3* levels triggering DNA

demethylation and consequently fruit ripening. (b) Active PHYB1B2 regulates the

expression of genes associated with DNA methylation (*MET1L*), chromatin remodelling

(*DDM1*) and histone modifiers (*PRMT* and *HDT3*), promoting overall fruit DNA

demethylation during the ripening process. PHYB1B2-mediated methylation reprogramming

results in the upregulation of master TFs, which in turn trigger ripening and carotenogenesis.

Competing interests

The authors declare no competing interests.

Data availability

All high-throughput sequencing data reported in this paper have been uploaded to the

Sequence Read Archive (SRA) under NCBI Bioproject PRJNA646733, with accession

numbers SUB7763724, SUB7782168 and SUB7791358 for RNAseq, WGBS and small

RNAseq, respectively.

References

1.   Kaiserli, E., Perrella, G. & Davidson, M. L. Light and temperature shape nuclear

architecture and gene expression. *Curr. Opin. Plant Biol.* **45**, 103–111 (2018).

2.   Galvão, V. C. & Fankhauser, C. Sensing the light environment in plants:

photoreceptors and early signaling steps. *Curr. Opin. Neurobiol.* **34**, 46–53 (2015).

3.   Paik, I. & Huq, E. Plant photoreceptors: Multi-functional sensory proteins and their

signaling networks. *Semin. Cell Dev. Biol.* **92**, 114–121 (2019).

4.   Cheng, Y.-L. & Tu, S.-L. Alternative Splicing and Cross-Talk with Light Signaling.

Plant Cell Physiol. **59**, 1104–1110 (2018).

5.   Shikata, H. *et al.* Phytochrome controls alternative splicing to mediate light responses

in Arabidopsis. *Proc. Natl. Acad. Sci.* **111**, 18781–18786 (2014).

6.   Wu, H.-P. *et al.* Genome-wide analysis of light-regulated alternative splicing mediated

by photoreceptors in *Physcomitrella patens*. *Genome Biol.* **15**, R10 (2014).

7.   Lin, B.-Y., Shih, C.-J., Hsieh, H.-Y., Chen, H.-C. & Tu, S.-L. Phytochrome

Coordinates with a hnRNP to Regulate Alternative Splicing via an Exonic Splicing

Silencer. *Plant Physiol.* **182**, 243 LP – 254 (2020).

8.   Ushijima, T. *et al.* Light Controls Protein Localization through Phytochrome-

Mediated Alternative Promoter Selection. *Cell* **171**, 1316-1325.e12 (2017).

9.   Mazzella, M. A. *et al.* Phytochrome control of the Arabidopsis transcriptome

anticipates seedling exposure to light. *Plant Cell* **17**, 2507–2516 (2005).

10. Ibarra, S. E., Auge, G., Sánchez, R. A. & Botto, J. F. Transcriptional programs related
to phytochrome a function in arabidopsis seed germination. *Mol. Plant* **6**, 1261–1273
(2013).

11. Carlson, K. D., Bhogale, S., Anderson, D., Tomanek, L. & Madlung, A. Phytochrome
a regulates carbon flux in dark grown tomato seedlings. *Front. Plant Sci.* **10**, 1–20
(2019).

12. Perrella, G. & Kaiserli, E. Light behind the curtain: photoregulation of nuclear
architecture and chromatin dynamics in plants. *New Phytol.* **212**, 908–919 (2016).

13. Tessadori, F. *et al.* Phytochrome B and histone deacetylase 6 control light-induced
chromatin compaction in *Arabidopsis thaliana*. *PLoS Genet.* **5**, e1000638 (2009).

14. Chua, Y. L., Brown, A. P. & Gray, J. C. Targeted histone acetylation and altered
nuclease accessibility over short regions of the pea plastocyanin gene. *Plant Cell* **13**,
599–612 (2001).

15. Chua, Y. L., Watson, L. A. & Gray, J. C. The transcriptional enhancer of the pea
plastocyanin gene associates with the nuclear matrix and regulates gene expression
through histone acetylation. *Plant Cell* **15**, 1468–1479 (2003).

16. Bertrand, C. *et al.* Arabidopsis HAF2 gene encoding TATA-binding protein (TBP)-
associated factor TAF1, is required to integrate light signals to regulate gene
expression and growth. *J. Biol. Chem.* **280**, 1465–1473 (2005).

17. Cho, J.-N. *et al.* Control of seed germination by light-induced histone arginine
demethylation activity. *Dev. Cell* **22**, 736–748 (2012).

18. Bourbousse, C., Barneche, F. & Laloi, C. Plant Chromatin Catches the Sun. *Frontiers
in Plant Science* **10**, 1728 (2020).

19. Ding, C. J., Liang, L. X., Diao, S., Su, X. H. & Zhang, B. Y. Genome-wide analysis
of day/night DNA methylation differences in *Populus nigra*. *PLoS One* **13**, 1–14
(2018).

20. Ding, J. *et al.* RNA-directed DNA methylation is involved in regulating photoperiod-
sensitive male sterility in rice. *Mol. Plant* **5**, 1210–1216 (2012).

21. Omidvar, V. & Fellner, M. DNA Methylation and Transcriptomic Changes in
Response to Different Lights and Stresses in 7B-1 Male-Sterile Tomato. *PLoS One*

10, e0121864 (2015).

22. Alba, R., Kelmenson, P. M., Cordonnier-Pratt, M.-M. & Pratt, L. H. The Phytochrome
Gene Family in Tomato and the Rapid Differential Evolution of this Family in
Angiosperms. *Mol. Biol. Evol.* **17**, 362–373 (2000).

23. Alves, F. R. R. *et al.* Beyond the limits of photoperception: constitutively active
PHYTOCHROME B2 overexpression as a means of improving fruit nutritional
quality in tomato. *Plant Biotechnol. J.* **n/a**, (2020).

24. Gramegna, G. *et al.* PHYTOCHROME-INTERACTING FACTOR 3 mediates light-
dependent induction of tocopherol biosynthesis during tomato fruit ripening. *Plant.*
Cell Environ. **42**, 1328–1339 (2019).

25. Bianchetti, R. *et al.* Fruit-localized phytochromes regulate plastid biogenesis, starch
synthesis, and carotenoid metabolism in tomato. *J. Exp. Bot.* **69**, 3573–3586 (2018).

26. Bianchetti, R. *et al.* Phytochrome-Dependent Temperature Perception Modulates
Isoprenoid Metabolism. *Plant Physiol.* **183**, 869 LP – 882 (2020).

27. Zhong, S. *et al.* Single-base resolution methylomes of tomato fruit development reveal
epigenome modifications associated with ripening. *Nat. Biotechnol.* **31**, 154–159
(2013).

28. Zuo, J. *et al.* Relationships between genome methylation, levels of non-coding RNAs,
mRNAs and metabolites in ripening tomato fruit. *Plant J.* (2020).
doi:10.1111/tpj.14778

29. Vrebalov, J. *et al.* A MADS-Box Gene Necessary for Fruit Ripening at the Tomato
Ripening-Inhibitor Locus. *Science (80).* **296**, 343 LP – 346 (2002).

30. Bemer, M. *et al.* The Tomato FRUITFULL Homologs TDR4/FUL1 and MBP7/FUL2
Regulate Ethylene-Independent Aspects of Fruit Ripening. *Plant Cell* **24**, 4437 LP –
4451 (2012).

31. Gupta, S. K. *et al.* Complex and shifting interactions of phytochromes regulate fruit
development in tomato. *Plant. Cell Environ.* **37**, 1688–1702 (2014).

32. Zhang, Y. *et al.* A Quartet of PIF bHLH Factors Provides a Transcriptionally Centered
Signaling Hub That Regulates Seedling Morphogenesis through Differential
Expression-Patterning of Shared Target Genes in Arabidopsis. *PLOS Genet.* **9**,
e1003244 (2013).

33.   Giovannoni, J. J. Fruit ripening mutants yield insights into ripening control. *Curr.*
765   *Opin. Plant Biol.* **10**, 283–289 (2007).

766   34.   Manning, K. *et al.* A naturally occurring epigenetic mutation in a gene encoding an
767   SBP-box transcription factor inhibits tomato fruit ripening. *Nat. Genet.* **38**, 948–952
768   (2006).

769   35.   Karlova, R. *et al.* Transcriptome and Metabolite Profiling Show That APETALA2a Is
770   a Major Regulator of Tomato Fruit Ripening. *Plant Cell* **23**, 923 LP – 941 (2011).

771   36.   Lang, Z. *et al.* Critical roles of DNA demethylation in the activation of ripening-
772   induced genes and inhibition of ripening-repressed genes in tomato fruit. *Proc. Natl.*
773   *Acad. Sci.* **114**, E4511–E4519 (2017).

774   37.   Zhang, B. *et al.* Chilling-induced tomato flavor loss is associated with altered volatile
775   synthesis and transient changes in DNA methylation. *Proc. Natl. Acad. Sci.* **113**,
776   12580 LP – 12585 (2016).

777   38.   Bewick, A. J. & Schmitz, R. J. Gene body DNA methylation in plants. *Curr. Opin.*
778   *Plant Biol.* **36**, 103–110 (2017).

779   39.   Lu, X. *et al.* Genome-Wide Epigenetic Regulation of Gene Transcription in Maize
780   Seeds. *PLoS One* **10**, e0139582 (2015).

781   40.   Zhang, H., Lang, Z. & Zhu, J. K. Dynamics and function of DNA methylation in
782   plants. *Nat. Rev. Mol. Cell Biol.* **19**, 489–506 (2018).

783   41.   Pikaard, C. S. & Scheid, O. M. Epigenetic regulation in plants. *Cold Spring Harb.*
784   *Perspect. Biol.* **6**, 1–32 (2014).

785   42.   Gallusci, P., Hodgman, C., Teyssier, E. & Seymour, G. B. DNA Methylation and
786   Chromatin Regulation during Fleshy Fruit Development and Ripening. *Frontiers in*
787   *Plant Science*   **7**, 807 (2016).

788   43.   Cao, D. *et al.* Genome-wide identification of cytosine-5 DNA methyltransferases and
789   demethylases in *Solanum lycopersicum*. *Gene* **550**, 230–237 (2014).

790   44.   Liu, R. *et al.* A DEMETER-like DNA demethylase governs tomato fruit ripening.
791   *Proc. Natl. Acad. Sci.* **112**, 10804 LP – 10809 (2015).

792   45.   Lei, M. *et al.* Regulatory link between DNA methylation and active demethylation in
793   Arabidopsis. *Proc. Natl. Acad. Sci.* **112**, 3553 LP – 3557 (2015).

46.   Zemach, A. *et al.* The arabidopsis nucleosome remodeler DDM1 allows DNA

methyltransferases to access H1-containing heterochromatin. *Cell* **153**, 193–205
(2013).

47. Pontvianne, F., Blevins, T. & Pikaard, C. S. Arabidopsis Histone Lysine
Methyltransferases. *Adv Bot Res.* **53**, 1–22 (2010).

48. Hernando, C. E., Sanchez, S. E., Mancini, E. & Yanovsky, M. J. Genome wide
comparative analysis of the effects of PRMT5 and PRMT4/CARM1 arginine
methyltransferases on the *Arabidopsis thaliana* transcriptome. *BMC Genomics* **16**,
192 (2015).

49. Li, Z. *et al.* Histone demethylase SlJMJ6 promotes fruit ripening by removing H3K27
methylation of ripening-related genes in tomato. *New Phytol.* **227**, 1138–1156 (2020).

50. Zhao, L. *et al.* Identification and characterization of histone deacetylases in tomato
(*Solanum lycopersicum*). *Frontiers in Plant Science* **5**, 760 (2015).

51. Guo, J. E. *et al.* Silencing of histone deacetylase SlHDT3 delays fruit ripening and
suppresses carotenoid accumulation in tomato. *Plant Sci.* **265**, 29–38 (2017).

52. Kerckhoffs, L. H. J. *et al.* Molecular analysis of tri-mutant alleles in tomato indicates
the Tri locus is the gene encoding the apoprotein of phytochrome B1. *Planta* **199**, 152–
157 (1996).

53. Lazarova, G. I. *et al.* Molecular analysis of PHYA in wild-type and phytochrome A-
deficient mutants of tomato. *Plant J.* **14**, 653–662 (1998).

54. Kerckhoffs, L. H. J. *et al.* Characterization of the gene encoding the apoprotein of
phytochrome B2 in tomato, and identification of molecular lesions in two mutant
alleles. *Mol. Gen. Genet.* **261**, 901–907 (1999).

55. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

56. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–
21 (2013).

57. Expósito-Rodríguez, M., Borges, A. A., Borges-Pérez, A. & Pérez, J. A. Selection of
internal control genes for quantitative real-time RT-PCR studies during tomato
development process. *BMC Plant Biol.* **8**, 131 (2008).

58. Lister, R. *et al.* Highly Integrated Single-Base Resolution Maps of the Epigenome in
Arabidopsis. *Cell* **133**, 523–536 (2008).

59.   Krueger, F. & Andrews, S. R. Bismark: A flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics* **27**, 1571–1572 (2011).

60.   Akalin, A. *et al.* MethylKit: a comprehensive R package for the analysis of genome-
wide DNA methylation profiles. *Genome Biol.* **13**, R87 (2012).

61.   Axtell, M. J. ShortStack: Comprehensive annotation and quantification of small RNA
genes. *Rna* **19**, 740–751 (2013).

62.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139–140 (2009).

63.   McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of
multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids*
Res. **40**, 4288–4297 (2012).

64.   Chen, Y., Lun, A. T. L. & Smyth, G. K. From reads to genes to pathways: differential
expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-
likelihood pipeline. *F1000 Research* **5**, 1438 (2016).

65.   Thimm, O. *et al.* MapMan: a user-driven tool to display genomics data sets onto
diagrams of metabolic pathways and other biological processes. *Plant J.* **37**, 914–939
(2004).

66.   Fornes, O. *et al.* JASPAR 2020: update of the open-access database of transcription
factor binding profiles. *Nucleic Acids Res.* **48**, D87–D92 (2019).

67.   Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic*
Acids Res. **43**, W39–W49 (2015).

68.   Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137
(2008).

69.   Lira, B. S. *et al.* Manipulation of a senescence-associated gene improves fleshy fruit
yield. *Plant Physiol.* **175**, 77–91 (2017).

a

RIN

log2 FC — IG, MG

FUL2

log2 FC — IG, MG

b

Solyc05g012020 *RIN*

WT

phyB1B2

1 Kb

Solyc03g114830 *FUL2*

WT

phyB1B2

1 Kb

TE ▬ UTR ▬ Exon ■ Intron ←

a

Relative chlorophyll content

WT   *phyB1B2*

b

log2 FC

CBP   *CAB-3c*   *POR3*

c

Solyc02g070990 *CBP*

WT   0-100

phyB1B2   0-100

-1.5kb   +1

500 pb

Solyc03g005780 *CAB-3c*

WT   0-100

phyB1B2   0-100

-1.5kb   +1

500 pb

Solyc07g054210 *POR3*

WT   0-100

phyB1B2   0-100

+1   -2kb

500 pb

◆ G-box   ➤ PBE-box   ➤ CA-hybrid   ➤ CG-hybrid

a

Solyc05g012020 *RIN*

WT IG
phyB1B2 IG
WT BK
phyB1B2 BK

+1   -2kb
500 pb

Solyc10g006880 *NOR*

WT IG
phyB1B2 IG
WT BK
phyB1B2 BK

+1   -2kb
500 pb

Solyc02g077920 *CNR*

WT IG
phyB1B2 IG
WT BK
phyB1B2 BK

+1   -2kb
500pb

Solyc03g044300 *AP2a*

WT IG
phyB1B2 IG
WT BK
phyB1B2 BK

-2kb   +1
500 pb

b

RIN

BK   RR

NOR

BK   RR

CNR

BK   RR

AP2a

BK   RR

RIN-binding motif   G-box   PBE-box   CA-hybrid   CG-hybrid

a

b