Impact of low-frequency coding variants on human facial shape

Dongjing Liu[1], Nora Alhazmi[2,3], Harold Matthews[4,5], Myoung Keun Lee[6], Jiarui Li[7], Jacqueline T. Hecht[8], George L. Wehby[9], Lina M. Moreno[10], Carrie L. Heike[11], Jasmien Roosenboom[6], Eleanor Feingold[1,12], Mary L. Marazita[1,6], Peter Claes[4,7], Eric C. Liao[13], Seth M. Weinberg[1,6*], John R. Shaffer[1,6*]

[1]Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America

[2] Department of Oral Biology, Harvard School of Dental Medicine, Boston, Massachusetts, United States of America

[3] King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

[4] Department of Human Genetics, KU Leuven, Leuven, Belgium

[5] Medical Imaging Research Center, UZ Gasthuisberg, Leuven, Belgium

[6] Center for Craniofacial and Dental Genetics, Department of Oral Biology, School of Dental Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America

[7] Department of Electrical Engineering, ESAT/PSI, KU Leuven, Leuven, Belgium

[8] Department of Pediatrics, University of Texas McGovern Medical Center, Houston, Texas, United States of America

[9] Department of Health Management and Policy, University of Iowa, Iowa City, Iowa, United States of America

[10] Department of Orthodontics, University of Iowa, Iowa City, Iowa, United States of America

[11] Department of Pediatrics, Seattle Children's Craniofacial Center, University of Washington, Seattle, Washington, United States of America

[12] Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America

[13] Department of Surgery, Center for Regenerative Medicine, Massachusetts General Hospital, Shriners Hospital, Boston, Massachusetts, United States of America

[*] Corresponding author
E-mail: smwst46@pitt.edu (SMW)
E-mail: john.r.shaffer@pitt.edu (JRS)

Abstract

The contribution of low-frequency variants to the genetic architecture of normal-range facial traits is unknown. We studied the influence of low-frequency coding variants (MAF < 1%) on multi-dimensional facial shape phenotypes in 2329 healthy Europeans. We used MultiSKAT o scan the exome for face-associated low-frequency variants in a gene-based manner. Seven genes (*AR*, *CARS2*, *FTSJ1*, *HFE*, *LTB4R*, *TELO2*, *NECTIN1*) were significantly associated with shape variation of the cheek, chin, nose and mouth areas. These genes displayed a wide range of phenotypic effects, with some impacting the full face and others affecting localized regions. The missense variant rs142863092 in *NECTIN1* had a significant effect on chin morphology, and was predicted bioinformatically to be deleterious. *NECTIN1* is an established craniofacial gene that underlies a human syndrome that includes a mandibular phenotype. We further showed that *nectin1a* mutations can affect zebrafish craniofacial development, with the size and shape of the mandibular cartilage altered in mutant animals. These Findings highlighted the role of low-frequency coding variants in normal-range facial variation.

Introduction

Significant progress has been made in elucidating the genetic basis of human facial traits (Richmond, Howe, Lewis, Stergiakouli, & Zhurov, 2018; Weinberg, Cornell, & Leslie, 2018; Weinberg et al., 2019). Genome-wide association studies (GWASs) have identified and replicated numerous common genetic variants associated with normal-range facial morphology (Adhikari et al., 2016; Cha et al., 2018; Claes et al., 2018; Cole et al., 2016; Crouch et al., 2018; M. K. Lee et al., 2017; F. Liu et al., 2012; Paternoster et al., 2012; Shaffer et al., 2016); yet these variants cumulatively explain only a small fraction of the heritable phenotypic variation. Based on large-scale genomic studies of other complex morphological traits such as height (D. J. Liu et al., 2017; Lu et al., 2017; Marouli et al., 2017), we hypothesized that functional variants at hundreds or perhaps thousands of loci have yet to be discovered. While we expect that common variants, with a minor allele frequency (MAF) greater than 1%, account for most of the heritable variation in facial morphology, low frequency (MAF<1%) genetic variants may also play an important role. An exome-wide study of human height, for example, discovered 29 low-frequency coding variants with large effects of up to 2 centimeters per allele (Marouli et al., 2017).

Our previous GWAS in a modestly sized cohort of healthy individuals identified 1932 common genetic variants associated with facial variation at 38 loci, 15 of which were independently replicated (Claes et al., 2018). The success of this GWAS was attributed in part to an innovative data-driven phenotyping approach, in which the 3D facial surfaces were partitioned into hierarchically organized regions, each defined by multiple axes of shape variation. This approach allows for testing of genetic variants on facial morphology at multiple levels of scale – from the entire face (global) to highly localized facial regions (local). Extending this global-to-local analysis of facial traits to the analysis of low-frequency variants requires an appropriate and scalable statistical framework capable of accommodating the multivariate nature of the facial shape variables. A recently developed statistical approach, MultiSKAT (Dutta, Scott, Boehnke, & Lee, 2019), was designed for this purpose and showed desirable performance in its original development.

In this study, we evaluated the influence of low frequency coding variants, captured by the Illumina HumanExome BeadChip, on normal-range facial morphology in 2,329 individuals. We applied multivariate gene-based association testing methods to multi-dimensional facial shape phenotypes derived from 3D facial images. The results of our analyses pointed to novel genes, including at least one involved in orofacial clefts and several others with no previously described role in craniofacial development or disorders. We provided experimental evidence of our genetic association results through expression screening and knockout experiments in a zebrafish model. These results enhance our understanding of the genetic architecture of human facial variation.

Materials and Methods

Ethics statement

Institutional ethics (IRB) approval was obtained at each recruitment site and all subjects gave
their written informed consent prior to participation (University of Pittsburgh Institutional
Review Board #PRO09060553 and #RB0405013; UT Health Committee for the Protection of
Human Subjects #HSC-DB-09-0508; Seattle Children's Institutional Review Board #12107;
University of Iowa Human Subjects Office/Institutional Review Board #200912764 and
#200710721).

Sample and Phenotyping

The study cohort comprised 2,329 unrelated, healthy individuals of European ancestry aged 3 to
40 years. Participants were eligible if they had not experienced facial trauma, major surgery,
congenital facial anomalies that could potentially affect their natural facial structure. 3D images
of each participant's resting face were captured via digital stereophotogrammetry using the
3dMD face camera system. The data-driven phenotyping approached has been described in detail
in a previous work (Claes et al., 2018). Briefly, approximately 10,000 points—"quasi-
landmarks"—were automatically placed across the facial surface, by a non-rigid registration of a
standard facial template onto each surface. The result is that each quasi-landmark represents the
same facial position across all participants (White et al., 2019). The configurations were then co-
aligned to their mean using generalized Procrustes analysis (GPA). The quasi-landmarks were
then clustered into groups of co-varying component points in order to partition the full face into
two segments. GPA was repeated within each of the two segments, and the process was
continued for a total of four iterations to generate a hierarchy of 31 facial segments (which we
call modules) comprising overlapping groups of quasi-landmarks. The hierarchical structure is
illustrated in Fig 1, where modules formed successive levels representing the shift from more
globally integrated to more locally focused morphology. Shape variation within each module
was represented by the 3D coordinates of all quasi-landmarks contained therein. To reduce the
dimensionality, principal components analysis and parallel analysis were performed on the quasi-
landmarks. The result was a set of 31 multivariate phenotypes made up of 8 to 50 principal
components (PCs) that jointly captured near complete shape variance. The effects of sex, age,
height, weight, facial size and genetic ancestry were corrected for at the phenotyping stage.
These facial module phenotypes were successfully used in our previous GWAS of common
variants (Claes et al., 2018), which demonstrated a clear advantage of this data-driven
multivariate modeling approach for gene-mapping studies over the traditional utilization of *a
priori* (Shaffer et al., 2016) and univariate (M. K. Lee et al., 2017) facial traits.

Fig 1. Hierarchical clustering of facial shape.
Global-to-local facial segmentation obtained using hierarchical spectral clustering. Segments are colored in blue. The highest-level segment representing the full face was split into two sub-segments, and this bifurcation process was repeated until a five-level hierarchy comprising 31 segments was formed.

In addition to the phenotype quality control process described in (Claes et al., 2018), we further examined the phenotypic distribution of each module for extreme outlier faces, as phenotypic outliers may adversely impact low-frequency variant tests (Auer, Reiner, & Leal, 2016). To accomplish this, we looked at both the joint and the pairwise distribution of all PCs underlying each module. We visualized quantile-quantile (Q-Q) plots of chi-squared quantiles versus robust squared Mahalanobis distances to identify outliers that deviated from the rest of the sample. Mahalanobis distance is a metric measuring how far an observation is to the center of the joint distribution (centroid equivalent in a multivariate space). We identified one individual who was an outlier for several PCs in module 27 (chin), and revisited the associated facial images to confirm data validity and sample eligibility. This individual was excluded from any subsequent analysis involving module 27.

Genotyping
Participants were genotyped by the Illumina OmniExpress + Exome v1.2 array, which included approximately 245,000 coding variants in the exome panel. Standard data cleaning and

102 imputation procedures were implemented. Imputed genotypes with a certainty above 0.9 were
103 used to fill in any sporadic missingness among genotype calls of the directly genotyped variants.
104 We did not include any wholly unobserved, imputed SNPs in this analysis. Ancestry PCs based
105 on common LD-pruned SNPs were constructed and regressed out from the multivariate traits to
106 adjust for population structure.
107

MultiSKAT
MultiSKAT (Dutta et al., 2019) was specifically developed for testing sets of variants, in this
case coding variants within genes, for association with a multivariate trait. Testing low-
frequency variants in aggregate can improve power compared to individual tests of each variant.
The tool is flexible in relating multiple variants collectively to multiple phenotypes through the
use of several choices of kernels, and includes an omnibus test to obtain optimal association p-
values by integrating results across different kernels via Copula. This capability of
accommodating multivariate phenotypes fits well with our analysis of facial modules, as each
module was composed of several independent PCs. MultiSKAT can be applied to both common
and rare variants, although our analysis considered low-frequency variants exclusively.
118

MultiSKAT uses a phenotype kernel to model how one variant affects multiple traits, and a
separate genotype kernel to specify how multiple variants influence one trait. In reality, these
effects are often not known *a priori*, and the true relationship can be a mixture of different
effects. We used the heterogeneous and homogeneous phenotype kernels, which are appropriate
when the set of traits analyzed are orthogonal PCs. We used the Sequence Kernel Association
Test (SKAT) and burden test as the genotype kernel, and performed the omnibus test in
MultiSKAT to aggregate results across the 2×2 kernel combinations.
126

Gene-level analysis
Genome-wide coding variants with MAF<1% were aggregated into genes. Per the developer's
suggested practice for using the MultiSKAT method, we filtered out variants with three or fewer
minor alleles to ensure that there is no inflation in MultiSKAT test statistics. We excluded genes
with less than two qualified variants, leading to 31347 variants in 8091 genes being tested. When
grouping multiple variants into a gene, MultiSKAT assigns larger weights to rarer variants. We
applied a Bonferroni threshold to declare significance. To account for the correlation among
partially overlapping facial modules, we used a procedure based on eigenvalues as proposed by
Li and Ji (Li & Ji, 2005) and computed that the effective number of independent modules was 19.
The threshold for significance was therefore set as $p < 3.3 \times 10^{-7}$ (i.e., 0.05/(8091 $\times$ 19)). The
phenotypic effects of identified genes on face were visualized by creating and comparing the
average facial morphs in individuals who had variants in a certain gene and those who do not
carry any variants.
140

Gene-set enrichment analysis was carried out using GREAT (McLean et al., 2010), FUMA
(Watanabe, Taskesen, van Bochoven, & Posthuma, 2017) and ToppFun (Chen, Bardes, Aronow,
& Jegga, 2009). Expression of genes were looked up in the GTEx database (GTEx Consortium,
2013). Following our hypothesis that genes influencing typical facial presentation may also be
involved in facial anomalies, we examined whether any genes identified by MultiSKAT were
associated with non-syndromic cleft palate with or without cleft lip (NSCL/P) by retrieving
association p-values from a past study of our group, where we performed a gene-based low-
frequency variant association scan on NSCL/P (Leslie et al., 2017).

Variant-level analysis
For genes highlighted by MultiSKAT, we scrutinized the quality of genotype calls by inspecting
allele intensity cluster plots. We further performed association tests of individual SNPs using
MultiPhen (O'Reilly et al., 2012). MultiPhen works by finding the linear combination of PCs
that is mostly associated with the genotypes at each SNP, and is robust when variants with low
frequencies are tested against non-normal phenotypes. Variant level functional prediction was
performed using CADD (Rentzsch, Witten, Cooper, Shendure, & Kircher, 2019). CADD is a
comprehensive metric that weights and integrates diverse sources of annotation, by contrasting
variants that survived natural selection with simulated mutations. The scaled CADD score
expresses the deleteriousness rank in terms of order of magnitude. A score of 10, for instance, is
interpreted as ranking in the top 10% in terms of the damaging degree amongst reference
genome SNPs, and a score of 20 refers to 1%, 30 to 0.1%, etc. Variant identifiers and
chromosomal locations are indicated according to the hg19 genome build. Individual variants
were searched in literature and PhenoScanner (Kamat et al., 2019) for existing human phenotype
associations.

We quantified the magnitude of phenotypic effect of individual low-frequency variants by the
difference between averaged faces of variant carriers (those who were heterozygotes; there was
no homozygotes for the low-frequency variants tested) and non-carries, which was further
compared with the effects of significant common variants identified in the prior GWAS of the
same multidimensional traits (Claes et al., 2018). Specifically, the centroids of the
multidimensional space defined by PCs in a certain module were computed separately for people
carrying the variant and people who do not carry the variant. Then the Euclidean distance
between the two centroids was calculated as a measure of variant effect size.

Expression screen of candidate genes in zebrafish
The whole-mount RNA in situ hybridization (WISH) for *ar, cars2, ftsj1, hfe, ltb4r, telo2,*
nectin1a and *nectin1b* was performed on wild type zebrafish embryos at 24 hpf and 48 hpf as
described by Thisse et al. (C. Thisse & Thisse, 2008). All wild type embryos were collected
synchronously at the corresponding stages and fixed in 4% paraformaldehyde (PFA) overnight.

T7 RNA polymerase promoter was added to the reverse primers and was synthesized with
antisense DIG-labeled probe in order to generate antisense RNA probe. The probe primers for *ar*
are: forward 5'- GTCCTACAAGAACGCCAACG-3' and reverse 5'-
GGTCACAGACTTGGAAAGGG-3' at 59°C. The *cars2* probe primers are: forward 5'-
ATCTGGGTCATGCGTGTTCA-3' and reverse 5'- GGATTCCTGTGGTGCTTGGT at 59°C.
The *ftsj1* probe primers are: forward 5'- GGCGAGAAGTGCCTTCAAAC-3' and reverse 5'-
AGTCGTGCTTGTGTCTGGTT-3' and *hfe* probe primers are: forward 5'-
GGGGATGGATGCTTCTACGA-3' and reverse 5'- CGCGCACACAAAATCATCAC-3' at
59°C. The *ltb4r* probe primers are: forward 5'-GACGGTGCATTACCTGTGC-3' and reverse
5'-AGTCTTGTCCGCCAAGGTC-3' at 58°C. The primers for *telo2* are: forward 5'-
GCTCCACTGGTGAGAGTGAG-3' and reverse 5'-GTCAGCTGAGGAGAGTCTGCG-3'. The
primers for *nectin1a* probe are: forward 5'-AACACCCAGGAGATCAGCAA-3' and reverse 5'-
CCTCCACCTCAGATCCGTAC-3' at 57°C and the *nectin1b* probe primers are: forward 5'-
TGCTAACCCAGCATTGGGAG-3' and reverse 5'-GGTTCTTGGGCATTGGAGGA-3' at
59°C. Embryos were mounted using glycerol and imaged using Nikon AZ100 multizoom
microscope.
196

Phenotype of mutant zebrafish
Zebrafish adults and embryos were obtained and maintained as described by Kimmel et al. (C. B.
Kimmel, Ballard, Kimmel, Ullmann, & Schilling, 1995). Zebrafish nectin1a mutants were
generated by transgene insertion Tg(Nlacz-GTvirus) in Chr 21: 21731876 - 21731886 (Zv9), and
obtained from Zebrafish International Resource Center, allele Ia021885Tg (ZIRC catalog ID:
ZL6899.07). The retroviral-mediated insertional mutagenesis inserts a molecular tag in the DNA
and isolates the allele of interest. Therefore, this will induce a frameshift and probably causing
either nonsense-mediated mRNA decay or a truncated protein (Amsterdam & Hopkins, 2003;
Sivasubbu, Balciunas, Amsterdam, & Ekker, 2007). The PCR genotyping primers for *nectin1a*
are: forward 5'-TTAGACCAGCCCACCTCA-3' and reverse 5'-
AATATGAAATAGCGCCGTTGTG-3' at 62°C.
208

209 Alcian blue staining was performed as described by Walker et al. (Walker & Kimmel, 2007).
210 The craniofacial cartilages were dissected and flat-mounted and then imaged using Nikon AZ100
211 multizoom microscope. After imaging, each embryo tail was placed in a PCR tube for
212 genotyping. The protocol was used as described by (Westerfield, 1994) with modification of
213 using fresh embryos without fixation.
214

216 **Results**
217

In the gene-based test of exome-wide low-frequency variants, seven genes were significantly associated with one or more facial modules (*HFE, NECTIN1, CARS2, LTB4R, TELO2, AR,* and *FTSJ1;* Fig 2 and Table 1). Three of them showed associations with more than one module. Fig 3 and S1 Table show the results of these genes in multiple modules. Fig 3 shows the association signals propagating along the branching paths from the more global segments to the more local segments. Four genes (*HFE, CARS2, LTB4R,* and *TELO2*) were associated with nose-related modules, and the others were associated with the shape of chin, mouth, and cheek. *FTSJ1* had broad associations in the full face as well as in local regions, while the effects of other genes were more confined to only local modules. We observed well-calibrated test statistics and little evidence of inflation as shown in the Q-Q plots (Fig S1).

Fig 2. Composite Manhattan plot showing results across 31 facial modules.
Manhattan plot showing the position of genes on the x axis and MultiSKAT p-values on the y axis. A total of 31 points are plotted for each gene, representing their p-values in each of the 31 modules. The red horizontal line indicates the significance threshold (3.3×10^{-7}). The associated facial modules and the corresponding p-value for each gene that surpassed the threshold are shown above the Manhattan plot. The numbers to the bottom left of the facial images indicate the module identifiers in Fig 1.

236 Table 1. Single variant association and functional prediction for variants contributing to the gene-level significance

Chr	Gene	Gene Info	Gene-level association		Variant-level association							
			Module[a]	MultiSKAT P-value[a]	SNP	Pos (hg19)	Ref/Alt[b]	Function[c]	CADD score[d]	MAF	Module[e]	MultiPhen P-value[e]
6	*HFE*	Homeostatic iron regulator, binds to transferrin receptor (TFR) and reduces its affinity for iron-loaded transferrin	5, **22**	1.51E-10	rs149342416	26087686	G/C	Arg6Ser	15.3	0.09%	22	0.07
					rs143662783	26087718	C/G	Thr17Ile	13.4	0.09%	5	0.87
11	*NECTIN1*	Nectin 1, cell adhesion molecule	27	2.54E-07	rs142863092	119548369	G/A	Arg210His	25.2	0.09%	27	1.08E-03
					rs137991779	119549425	G/A	Gly44Ser	29.2	0.11%	27	0.15
13	*CARS2*	Cysteinyl-tRNA synthetase 2, mitochondrial	**10**, 20	5.22E-10	rs151097801	111296817	C/T	Pro138Leu	22.4	0.09%	20	0.12
					rs117788141	111357899	G/A	Val69Ile	28	0.09%	10	0.01
14	*LTB4R*	Leukotriene B4 receptor 1, receptor for extracellular ATP > UTP and ADP	20	6.88E-08	rs143666989	24780865	A/G	Gln332Arg	16.6	0.11%	20	0.11
					rs148153989	24780915	A/T	Met349Leu	12.5	0.09%	20	0.59
16	*TELO2*	Telomere length regulation protein homolog, regulate DNA damage response	10	1.19E-09	rs140903666	1544313	G/A	Ala11Thr	6.3	0.22%	10	8.21E-04
					rs144863771	1544314	C/A	Ala11Asp	10.7	0.22%	10	8.21E-04
					rs147858841	1555541	C/T	Ala132Val	9.4	0.11%	10	0.43
23	*AR*	Androgen receptor, steroid hormone receptors	18	1.77E-07	rs142280455	66905875	A/G	Ser598Gly	22.4	0.13%	18	0.81
					rs137852591	66941751	C/G	Gln267Glu	25	0.13%	18	3.91E-03
23	*FTSJ1*	Putative tRNA (cytidine(32)/guanosine(34)-2'-O)-methyltransferase	1, 6, 12, **28**	2.46E-10	rs142932029	48341118	G/A	Ser161Asn	7.4	0.08%	28	1.59E-14
					rs201095751	48341414	C/T	Splice site	0.1	0.11%	12	0.1

[a] For genes associated with multiple facial modules, the most significant module is in bold and only it's p-value is shown
[b] Alleles are listed as alternative/reference alleles on the forward strand of the reference genome
[c] For missense variant, amino acid substitution is given
[d] Bioinformatic prediction of variant effect, higher score indicates greater damaging effect
[e] Variants were tested against all module(s) with gene-level significance, and for genes associated with multiple modules, only the module yielding the smallest p-value in the variant-
level test is shown

Fig 3. Module-wide association results for significant genes.

For each gene, the –log10 p-value is shown as color shades ranging from min to max, for 31 facial segments arranged the same way as Fig 1. The global-to-local phenotyping enabled the discovery of genetic effects at different scales.

To visualize the effects of these genes on facial shape, we created the average module shape in non-carriers of the low-frequency variants for each gene, and a corresponding morph showing the change in shape from non-carriers to carriers (Fig 4). Blue and red indicate a local shape depression and protrusion, respectively, due to carrying any low-frequency variants. As an example, panel B in Fig 4 shows that *NECTIN1* variants shape the chin into a sharper and more protruding structure.

Fig 4. Phenotypic effect of the seven identified genes in their top associated module.
Blue and red indicate a local shape depression and protrusion, respectively, due to carrying the low-frequency variants in the gene. A) First column shows gene effect on a representative

module placing on the full face; middle column shows the lateral view of the average shape of
the corresponding module among people who do not carry any variant in the gene; right column
shows the change in the shape of the same module, from non-carrier to carrier, multiplied by a
constant (7), to make the changes more visibly distinctive. B) For *NECTIN1* gene, we show both
lateral (top) and frontal (bottom) view of its effect on chin shape. *NECTIN1* variant carriers on
average displayed a sharper, more protruding chin.
298
299
We employed various bioinformatics tools to explore the functions associated with the set of
identified genes. Enrichment was detected for a variety of biological processes (Fig S2),
especially ion-, metabolism-, transport- and regulation-related processes. Enriched gene ontology
(GO) molecular functions included signaling receptor and protein binding activity. Two genes
with relatively well characterized functions (*HFE* and *AR)* contributed a lot to these enrichment
results. In the GTEx database, these seven genes showed measurable expression level in adipose,
skin and muscle-skeletal tissue (Fig S3), among which the strongest expression was seen for
NECTIN1 in skin.
308
To explore whether facial genes also affect the risk of orofacial clefts, results of gene-based
associations of low-frequency (MAF<1%) variants with NSCL/P were retrieved from Leslie et al.
2017. Two out of the seven were not available from that study. S2 Table shows the SKAT and
CMC test results for the other five genes in the European, Asian, South American and the
combined samples. Two associations passed a Bonferroni corrected threshold for 40 tests (5
genes × 4 populations × 2 type of tests)— *TELO2* with a CMC p-value = 6.5 x 10^{-4}, and HFE
with a CMC p-value = 1.1×10^{-3}, both in the combined population of all ancestry groups.
316
317 Single variants in the genes showing significant associations in the gene-based tests were further
318 tested individually with the corresponding facial modules (Table 1). Six SNPs showed nominal
319 associations (p-value < 0.05) and the top association involved SNP rs142932029 in *FTSJ1* with
320 module 28 (p-value = 1.59×10^{-14}). As shown in S4 Fig, these low-frequency variants had larger
321 effects compared to previously reported common variants (Claes et al., 2018).
322
Most of the individual variants appeared at frequencies much lower than 1%, and all encode
nonsynonymous substitutions except one splice site SNP in *FTSJ1*. Variants in *NECTIN1,*
CARS2 and *AR* are predicted to be deleterious according to their CADD score (details in S3
Table). SNP rs137991779 in *NECTIN1* has a CADD score of 29.2, interpreted as ranking in the
top 0.12% in terms of deleteriousness among variants across the whole genome. PhenoScanner
linked those variants with a variety of human traits/disorders in previous studies (S4 Table,
mostly from UK Biobank), including height, vascular diseases, osteoporosis, neoplasms etc.,

suggesting that coding variants influencing facial shape may be pleiotropic and play roles in
other biological processes.
332
Zebrafish WISH was used to examine *ar, cars2, ftsj1, hfe, ltb4r, telo2, nectin1a* and *nectin1b*
expression pattern in the craniofacial region across key developmental stages (Fig 5). At 24
hours post fertilization (hpf), *ftsj1* was expressed in the hindbrain, and *hfe* and *ltb4r* were
expressed in the forebrain. We detected nectin1a and nectin1b transcripts in the eyes,
diencephalon, midbrain and hindbrain at 24 hpf. At 48 hpf, *ar* expression was detected in the
epiphysis*, cars2, nectin1a* and *nectin1b* were expressed in the palate (Fig 5 solid arrow), and
nectin1a was detected in the lower jaw (Fig 5 hollow arrow).
340

Fig 5. Whole-mount RNA in situ hybridization demonstrating genes expression in zebrafish.
Genes expression pattern in lateral and ventral views at the indicated embryonic stages as hours
per fertilization (hpf). *cars2, nectin1a* and *nectin1b* are expressed in zebrafish palate (solid
arrow). *nectin1a* is expressed in the lower jaw at 48 hpf (hollow arrow). cb: cerebellum, e:
epiphysis, ey: eye, h: heart, hb: hindbrain, op: olfactory placode, pq: palate quadrate, tel:
telencephalon.

To investigate if *nectin1a* is required for during normal craniofacial development, we analyzed
the *nectin1a* mutant allele Ia021885Tg. Breeding of *nectin1a*+/- intercross generated embryos

with Mendelian ratio (1 individual homozygous for the wild type allele: 2 heterozygous
individuals with one wild type and one mutant allele: 1 individual homozygous for the mutant
allele) demonstrating a mutant craniofacial phenotype, characterized by small head
structures (Fig 6). Using Alcain blue staining at 120 hpf, *nectin1a* mutants displayed dysmorphic
craniofacial development with smaller and distorted palate and abnormal Meckel's cartilage
compared to age-matched wild type zebrafish embryos from the same intercross. These results
show that *nectin1a* is genetically required for palate and mandible morphogenesis.

360
Fig 6. Alcian blue images for *nectin1a* zebrafish mutant comparted to wild type at day 5.
Top images: wild type alcian blue lateral view, palate and Meckel's cartilage. Middle images:
heterozygous *nectin1a* embryo alcian blue. Bottom images: homozygous *nectin1a* mutant lateral
view. The length of the palate was measured from the anterior midpoint to the posterior midpoint
of the palate. The width was measured as the maximum distance between the 2 lateral borders at
the anterior area. The length of the Meckel's cartilage was measured from the midline of the
Meckel's cartilage to the midline of an imaginary line drawn joining the joints between the
Meckel's cartilage and the palatoquadrate. The width was measured from the junction of the
Meckel's cartilage and the palatoquadrate of one side to the other side. Compared to wild type

animals. *nectin1a* mutants have smaller and shorter palate, and shorter and wider Meckel's
cartilage. L: length, W: width. Scale bar: 10 μm
372
373

374   **Discussion**
375

This study presented a discovery effort to identify low-frequency coding variants associated with
normal-range human facial shape, by undertaking gene-based association tests on a carefully
phenotyped human cohort followed by functional experiments of the association results. Overall,
we demonstrated that part of the morphological variation of facial shape is attributable to low-
frequency coding variants, and pinpointed putative functional genes involved. Seven genes (*AR*,
CARS2, *FTSJ1*, *HFE*, *LTB4R*, *TELO2* and *NECTIN1*) were identified, with phenotypic effects in
the area of cheek, chin, nose and mouth. Notably, *NECTIN1* is known to cause a syndrome
characterized by facial dysmorphology. Using a zebrafish model, we confirmed the expression of
nectin1a and *nectin1b* in the developing head and the abnormal craniofacial phenotype in
nectin1a mutants, with the affected structures being highly consistent with the associated facial
region in the human data analysis. Taken together, these findings support the contribution of
low-frequency coding variants to the genetic architecture of normal-range facial shape.
388

The seven genes identified by the multivariate approach are for the first time implicated in
normal facial morphology. Six of the seven genes (all but *cars2)* were expressed in embryonic
craniofacial tissues in zebrafish, demonstrating their potential involvement in craniofacial
development. Cellular processes/functions of these genes include metal ion transport (*HFE*),
signaling (*AR*, *LTB4R*), tRNA metabolism (*CARS2, FTSJ1*), DNA repair (*TELO2*) and cell
adhesion (*NECTIN1*). This diversity in their biological function led to a variety of enriched
functional pathways/categories in the gene-set enrichment analysis, yet without a strong signal in
any particular one, probably due to the small number of genes and the polygenic nature of facial
morphology. With the exception of *NECTIN1*, the role of these genes in patterning craniofacial
structures is unknown, and further investigation is needed to gain better understanding of how
these genes may affect the face.
400

Previous GWASs and studies of facial dysmorphology have demonstrated that there are common
genetic factors underlying normal-range facial variation and orofacial clefting (Claes et al., 2018;
F. Liu et al., 2012; Weinberg et al., 2009). Our findings suggest that low-frequency coding
variants may also help explain this relationship. Although none of the other genes implicated
here have been shown to be involved in craniofacial development, *NECTIN1* is an established
player that has been linked to both syndromic and isolated forms of orofacial clefting (Avila et
al., 2006; Sozen et al., 2001; Suzuki et al., 2000). Individuals with cleft lip/palate-ectodermal
dysplasia syndrome (OMIM:225060) have distinctive facial features including an

underdeveloped lower jaw (Zlotogora, 1994), which is consistent with the facial segment (chin)
where the *NECTIN1* association was observed. Although not passing the genome-wide threshold,
NECTIN1 also yielded some signals in modules representing the nose and cheek (Fig 3),
additional facial regions affected in this syndrome. Different variants in *NECTIN1* are likely
involved in normal-range variation and in craniofacial disorders, which may help explain
apparent differences in phenotypic severity. NECTIN1 protein belongs to the subfamily of
immunoglobulin-like adhesion molecules which are key components of cell adhesion junctions
and play critical roles in the development of many tissues, including in the fusion of palatal
shelves during palatogenesis (Cobourne, 2004). A handful of *NECTIN1* mutations that can
potentially disrupt gene function have been documented in non-syndromic cleft patients (Oner &
Tastan, 2016; Scapoli et al., 2006; Tongkobpetch, Suphapeetiporn, Siriwan, & Shotelersuk,
2008). In the current study, two coding variants in *NECTIN1* contributed to the gene-level
significance, both predicted to be deleterious. We performed lookups of the face-associated
genes in a previous exome scan of a NSCL/P cohort (Leslie et al., 2017). *NECTIN1* yielded a
small p-value of 0.004, although not passing the Bonferroni significance threshold. Two other
genes, *TELO2* and *HFE*, did pass that threshold. These results are in line with previous evidence
suggesting a role for same genes in normal and abnormal facial development.
426
Our zebrafish experiments provided a strong support for the relevance of *nectin1a* in palate and
mandible development. The mutants displayed changes in the shape and size of both the palate
and the Meckel's cartilage, from which the mandibles evolved. This affected cartilage structure
in zebrafish mutants aligns well with the associated human anatomical region (chin and
mandible), where the effects of *NECTIN1* were observed in the MultiSKAT test. These findings
for the first time demonstrate a role of *NECTIN1* in normal-range facial variation. We highlight
the approach of interrogating human candidate genes in a biological context using the zebrafish
model, where dynamic gene expression can be assayed in a high throuput fashion. Those
candidate genes with spatiotemporal gene expression in the craniofacial domains then can be
evaluated in functional studies, were mutants may already be available from large scale
mutagenesis projects, or can be generated by CRISPR mediated gene editing.
438
With the hierarchical facial segmentation, we were able to identify genetic effects at different
scales. For example, the effects of *FTSJ1* were observed globally in the full face, and also locally
in specific modules on the side of the face. By contrast, the effect of *NECTIN1* was confined to
localized facial parts only. These patterns may help with understanding the mechanisms by
which genes act along the growth of facial structure. Our multivariate data-driven phenotyping
approach eliminates the need of preselecting traits, captures more variation in the facial shape,
and is more effective for gene mapping.
446

The current study is an important extension and complement of our prior work on common SNPs
(Claes et al., 2018). Here we exclusively focused on coding variants with MAF below 1%, which
have been omitted based no standard QC procedure from previous facial GWAS attempts. We
compared results from this study to those from our prior GWAS (Claes et al., 2018), and noted
that common variants in or near (within 500kb) the seven associated genes showed no evidence
of association ($p > 0.001$ for all) with the same facial modules. This indicates that the current
study generated distinct, non-overlapping knowledge on facial genetics, although it is possible
that there are trans-acting common GWAS SNPs that regulate the expression of the seven
identified genes during facial morphogenesis. Low-frequency variants showed larger magnitude
of effects compared to common variants in (Claes et al., 2018). It is necessary, however, to point
out that this difference could partially or completely be a result of the drastically smaller groups
of variant carriers, and we therefore refrain from overinterpreting the comparison.

Our study demonstrated the power of applying gene-based tests of low-frequency variants that
are usually untestable individually. While some significant genes harbor variants with a small p-
value in our single-variant association test, others would have been missed if not tested in
aggregate. With a moderate sample size of 2329, it is highly desirable to collapse low-frequency
variants into putative functional units and perform burden-style tests. In addition to an increase
in power, another key benefit with analyzing low-frequency coding variants collectively is the
improved biological interpretability compared to GWASs. The gene-centered design of coding
variant tests facilities much clearer biological implications and options for experimental follow-
up. Our success with the functional validation of *NECTIN1* provides a practical example. We
expect future better-powered studies to discover more biological pathways emerging from
analyses of low-frequency coding variants.

Replication of rare and low-frequency variant association signals presents unique challenges.
The prominent barrier is the limited sample size. The low numbers or even absence of the
carriers in independent populations hindered the replication efforts of our findings. Six out of the
seven genes identified were not testable in a separate cohort of 664 participants due to a lack of
variant carriers. Given our sample size and the ExomeChip design, this study was not adequately
powered to identify genes harboring extra rare variants that may also contribute to facial traits.
Although complex traits are not expected to have a large fraction of the heritability explained by
rare and private variants, such variants may be influential, predictive, and actionable at the
individual level. In this regard, whole exome or whole genome sequencing of large samples
holds promise to give deeper insights into the role rare variants in facial morphology.

Like many other complex traits, research with a focus on uncovering the genetic architecture of
facial morphology is confronted with the challenge of missing heritability (Cole et al., 2017;
Tsagkrasoulis, Hysi, Spector, & Montana, 2017). Our study has extended the paradigm of

genetic factors involved in facial morphology from common to low frequency variants and highlighted novel candidate genes that may lead to encouraging follow-ups. Given that rare and low-frequency genetic variation might be highly specific to certain populations, and facial shapes have distinctive ancestry features, future studies may benefit from extending the discovery of influential low-frequency variants to other ethnic groups.

Acknowledgements

The authors thank all the dedicated staff, collaborators, and participants for their contribution to the study.

Funding

This study was supported by the National Institute for Dental and Craniofacial Research (NIDCR, (http://www.nidcr.nih. gov/) through the following grants: U01-DE020078 to SMW and MLM, R01-DE016148 to SMW and MLM, R01-DE027023 to SMW and JRS, and X01-HG007821 to MLM, SMW, and EF. Funding for initial genomic data cleaning by the University of Washington was provided by contract #HHSN268201200008I from the NIDCR awarded to the Center for Inherited Disease Research (CIDR, https://www.cidr.jhmi.edu/). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability

All of the genotypic markers are available to the research community through the dbGaP controlled-access repository (https://dbgap.ncbi.nlm.nih.gov/) at accession phs000949.v1.p1. The raw source data for the phenotypes – the 3D facial surface models – are available for the 3D Facial Norms dataset through the FaceBase Consortium (www.facebase.org). Access to these 3D facial surface models requires proper institutional ethics approval and approval from the FaceBase data access committee. KU Leuven provides the spatially dense facial mapping software, free to use for academic purposes: MeshMonk (https://github.com/TheWebMonks/meshmonk).

Reference

Adhikari, K., Fuentes-Guajardo, M., Quinto-Sánchez, M., Mendoza-Revilla, J., Camilo Chacón-Duque, J., Acuña-Alonzo, V., et al. (2016). A genome-wide association scan implicates DCHS2, RUNX2, GLI3, PAX1 and EDAR in human facial variation. *Nature Communications*, *7*(1), 11616. http://doi.org/10.1038/ncomms11616

Amsterdam, A., & Hopkins, N. (2003). Retroviral-Mediated Insertional Mutagenesis in
Zebrafish. *Methods in Cell Biology*, *77*, 3–20. http://doi.org/10.1016/S0091-679X(04)77001-
6
Auer, P. L., Reiner, A. P., & Leal, S. M. (2016). The effect of phenotypic outliers and non-
normality on rare-variant association testing. *European Journal of Human Genetics : EJHG*,
24(8), 1188–1194. http://doi.org/10.1038/ejhg.2015.270
Avila, J. R., Jezewski, P. A., Vieira, A. R., Orioli, I. M., Castilla, E. E., Christensen, K., et al.
(2006). PVRL1 variants contribute to non-syndromic cleft lip and palate in multiple
populations. *American Journal of Medical Genetics Part A*, *140*(23), 2562–2570.
http://doi.org/10.1002/ajmg.a.31367
Cha, S., Lim, J. E., Park, A. Y., Do, J.-H., Lee, S. W., Shin, C., et al. (2018). Identification of
five novel genetic loci related to facial morphology by genome-wide association studies.
Bmc Genomics, *19*(1), 481–17. http://doi.org/10.1186/s12864-018-4865-9
Chen, J., Bardes, E. E., Aronow, B. J., & Jegga, A. G. (2009). ToppGene Suite for gene list
enrichment analysis and candidate gene prioritization. *Nucleic Acids Research*, *37*(Web
Server), W305–W311. http://doi.org/10.1093/nar/gkp427
Claes, P., Roosenboom, J., White, J. D., Swigut, T., Sero, D., Li, J., et al. (2018). Genome-wide
mapping of global-to-local genetic effects on human facial shape. *Nature Genetics*, *50*(3), 1–
16. http://doi.org/10.1038/s41588-018-0057-4
Cobourne, M. T. (2004). The complex genetics of cleft lip and palate. *European Journal of*
Orthodontics, *26*(1), 7–16.
Cole, J. B., Manyama, M., Kimwaga, E., Mathayo, J., Larson, J. R., Liberton, D. K., et al.
(2016). Genomewide Association Study of African Children Identifies Association of
SCHIP1 and PDE8A with Facial Size and Shape. *PLoS Genetics*, *12*(8), e1006174.
http://doi.org/10.1371/journal.pgen.1006174
Cole, J. B., Manyama, M., Larson, J. R., Liberton, D. K., Ferrara, T. M., Riccardi, S. L., et al.
(2017). Human Facial Shape and Size Heritability and Genetic Correlations. *Genetics*,
205(2), 967–978. http://doi.org/10.1534/genetics.116.193185
Crouch, D. J. M., Winney, B., Koppen, W. P., Christmas, W. J., Hutnik, K., Day, T., et al.
(2018). Genetics of the human face: Identification of large-effect single gene variants.
Proceedings of the National Academy of Sciences of the United States of America, *115*(4),
E676–E685. http://doi.org/10.1073/pnas.1708207114
Dutta, D., Scott, L., Boehnke, M., & Lee, S. (2019). Multi-SKAT: General framework to test for
rare-variant association with multiple phenotypes. *Genetic Epidemiology*, *43*(1), 4–23.
http://doi.org/10.1002/gepi.22156
GTEx Consortium. (2013). The Genotype-Tissue Expression (GTEx) project. *Nature Genetics*,
45(6), 580–585. http://doi.org/10.1038/ng.2653

Kamat, M. A., Blackshaw, J. A., Young, R., Surendran, P., Burgess, S., Danesh, J., et al. (2019).
PhenoScanner V2: an expanded tool for searching human genotype-phenotype associations.
Bioinformatics, *35*(22), 4851–4853. http://doi.org/10.1093/bioinformatics/btz469

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B., & Schilling, T. F. (1995). Stages of
Embryonic-Development of the Zebrafish. *Developmental Dynamics*, *203*(3), 253–310.
http://doi.org/10.1002/aja.1002030302

Lee, M. K., Shaffer, J. R., Leslie, E. J., Orlova, E., Carlson, J. C., Feingold, E., et al. (2017).
Genome-wide association study of facial morphology reveals novel associations with
FREM1 and PARK2. *PLoS ONE*, *12*(4), e0176566–13.
http://doi.org/10.1371/journal.pone.0176566

Leslie, E. J., Carlson, J. C., Shaffer, J. R., Buxó, C. J., Castilla, E. E., Christensen, K., et al.
(2017). Association studies of low-frequency coding variants in nonsyndromic cleft lip with
or without cleft palate. *American Journal of Medical Genetics Part B: Neuropsychiatric
Genetics*, *173*(6), 1531–1538. http://doi.org/10.1002/ajmg.a.38210

Li, J., & Ji, L. (2005). Adjusting multiple testing in multilocus analyses using the eigenvalues of
a correlation matrix. *Heredity*, *95*(3), 221–227. http://doi.org/10.1038/sj.hdy.6800717

Liu, D. J., Peloso, G. M., Yu, H., Butterworth, A. S., Wang, X., Mahajan, A., et al. (2017).
Exome-wide association study of plasma lipids in >300,000 individuals. *Nature Genetics*,
49(12), 1758–1766. http://doi.org/10.1038/ng.3977

Liu, F., van der Lijn, F., Schurmann, C., Zhu, G., Chakravarty, M. M., Hysi, P. G., et al. (2012).
A Genome-Wide Association Study Identifies Five Loci Influencing Facial Morphology in
Europeans. *PLoS Genetics*, *8*(9), e1002932–13. http://doi.org/10.1371/journal.pgen.1002932

Lu, X., Peloso, G. M., Liu, D. J., Wu, Y., Zhang, H., Zhou, W., et al. (2017). Exome chip meta-
analysis identifies novel loci and East Asian–specific coding variants that contribute to lipid
levels and coronary artery disease. *Nature Genetics*, *49*(12), 1722–1730.
http://doi.org/10.1038/ng.3978

Marouli, E., Graff, M., Medina-Gomez, C., Lo, K. S., Wood, A. R., Kjaer, T. R., et al. (2017).
Rare and low-frequency coding variants alter human adult height. *Nature*, *542*(7640), 186–
190. http://doi.org/10.1038/nature21039

McLean, C. Y., Bristor, D., Hiller, M., Clarke, S. L., Schaar, B. T., Lowe, C. B., et al. (2010).
GREAT improves functional interpretation of cis-regulatory regions. *Nature Biotechnology*,
28(5), 495–501. http://doi.org/10.1038/nbt.1630

Oner, D. A., & Tastan, H. (2016). Identification of Novel Variants in the PVRL1 Gene in
Patients With Nonsyndromic Cleft Lip With or Without Cleft Palate. *Genetic Testing and
Molecular Biomarkers*, *20*(5), 269–272. http://doi.org/10.1089/gtmb.2015.0276

O'Reilly, P. F., Hoggart, C. J., Pomyen, Y., Calboli, F. C. F., Elliott, P., Jarvelin, M.-R., & Coin,
L. J. M. (2012). MultiPhen: Joint Model of Multiple Phenotypes Can Increase Discovery in
GWAS. *PLoS ONE*, *7*(5), e34861–12. http://doi.org/10.1371/journal.pone.0034861

Paternoster, L., Zhurov, A. I., Toma, A. M., Kemp, J. P., Pourcain, B. S., Timpson, N. J., et al.
(2012). Genome-wide Association Study of Three-Dimensional Facial Morphology
Identifies a Variant in PAX3 Associated with Nasion Position. *The American Journal of*
Human Genetics, *90*(3), 478–485. http://doi.org/10.1016/j.ajhg.2011.12.021
Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J., & Kircher, M. (2019). CADD: predicting
the deleteriousness of variants throughout the human genome. *Nucleic Acids Research*,
47(D1), D886–D894. http://doi.org/10.1093/nar/gky1016
Richmond, S., Howe, L. J., Lewis, S., Stergiakouli, E., & Zhurov, A. (2018). Facial Genetics: A
Brief Overview. *Frontiers in Genetics*, *9*, 462. http://doi.org/10.3389/fgene.2018.00462
Scapoli, L., Palmieri, A., Martinelli, M., Vaccari, C., Marchesini, J., Pezzetti, F., et al. (2006).
Study of the PVRL1 gene in Italian nonsyndromic cleft lip patients with or without cleft
palate. *Annals of Human Genetics*, *70*(Pt 3), 410–413. http://doi.org/10.1111/j.1529-
8817.2005.00237.x
Shaffer, J. R., Orlova, E., Lee, M. K., Leslie, E. J., Raffensperger, Z. D., Heike, C. L., et al.
(2016). Genome-Wide Association Study Reveals Multiple Loci Influencing Normal Human
Facial Morphology. *PLoS Genetics*, *12*(8), e1006149–21.
http://doi.org/10.1371/journal.pgen.1006149
Sivasubbu, S., Balciunas, D., Amsterdam, A., & Ekker, S. C. (2007). Insertional mutagenesis
strategies in zebrafish. *Genome Biology*, *8*(S1), S9. http://doi.org/10.1186/gb-2007-8-s1-s9
Sozen, M. A., Suzuki, K., Tolarova, M. M., Bustos, T., Fernández Iglesias, J. E., & Spritz, R. A.
(2001). Mutation of PVRL1 is associated with sporadic, non-syndromic cleft lip/palate in
northern Venezuela. *Nature Genetics*, *29*(2), 141–142. http://doi.org/10.1038/ng740
Suzuki, K., Hu, D., Bustos, T., Zlotogora, J., Richieri-Costa, A., Helms, J. A., & Spritz, R. A.
(2000). Mutations of PVRL1, encoding a cell-cell adhesion molecule/herpesvirus receptor,
in cleft lip/palate-ectodermal dysplasia. *Nature Genetics*, *25*(4), 427–430.
http://doi.org/10.1038/78119
Thisse, C., & Thisse, B. (2008). High-resolution in situ hybridization to whole-mount zebrafish
embryos. *Nature Protocols*, *3*(1), 59–69. http://doi.org/10.1038/nprot.2007.514
Tongkobpetch, S., Suphapeetiporn, K., Siriwan, P., & Shotelersuk, V. (2008). Study of the
poliovirus receptor related-1 gene in Thai patients with non-syndromic cleft lip with or
without cleft palate. *International Journal of Oral & Maxillofacial Surgery*, *37*(6), 550–553.
http://doi.org/10.1016/j.ijom.2008.01.024
Tsagkrasoulis, D., Hysi, P., Spector, T., & Montana, G. (2017). Heritability maps of human face
morphology through large-scale automated three-dimensional phenotyping. *Scientific*
Reports, *7*(1), 1–18. http://doi.org/10.1038/srep45885
Walker, M. B., & Kimmel, C. B. (2007). A two-color acid-free cartilage and bone stain for
zebrafish larvae. *Biotechnic & Histochemistry*, *82*(1), 23–28.
http://doi.org/10.1080/10520290701333558

Watanabe, K., Taskesen, E., van Bochoven, A., & Posthuma, D. (2017). Functional mapping and
annotation of genetic associations with FUMA. *Nature Communications*, *8*(1), 1826.
http://doi.org/10.1038/s41467-017-01261-5

Weinberg, S. M., Cornell, R., & Leslie, E. J. (2018). Craniofacial genetics: Where have we been
and where are we going? *PLoS Genetics*, *14*(6), e1007438.
http://doi.org/10.1371/journal.pgen.1007438

Weinberg, S. M., Naidoo, S. D., Bardi, K. M., Brandon, C. A., Neiswanger, K., Resick, J. M., et
al. (2009). Face shape of unaffected parents with cleft affected offspring: combining three-
dimensional surface imaging and geometric morphometrics. *Orthodontics & Craniofacial*
Research, *12*(4), 271–281. http://doi.org/10.1111/j.1601-6343.2009.01462.x

Weinberg, S. M., Roosenboom, J., Shaffer, J. R., Shriver, M. D., Wysocka, J., & Claes, P.
(2019). Hunting for genes that shape human faces: Initial successes and challenges for the
future. *Orthodontics & Craniofacial Research*, *22 Suppl 1*(S1), 207–212.
http://doi.org/10.1111/ocr.12268

Westerfield, M. (1994). A guide for the laboratory use of zebrafish Danio (Brachydanio) rerio.
In: University of Oregon Press, Eugene.

White, J. D., Ortega-Castrillón, A., Matthews, H., Zaidi, A. A., Ekrami, O., Snyders, J., et al.
(2019). MeshMonk: Open-source large-scale intensive 3D phenotyping. *Scientific Reports*,
9(1), 6085. http://doi.org/10.1038/s41598-019-42533-y

Zlotogora, J. (1994). Syndactyly, Ectodermal Dysplasia, and Cleft Lip/Palate. *Journal of*
Medical Genetics, *31*(12), 957–959. http://doi.org/10.1136/jmg.31.12.957

Supporting Information

S1 Fig. Q-Q plot of gene-based MultiSKAT tests by facial module

S2 Fig. FUMA enrichment results

S3 Fig. GTEx expression of MultiSKAT significant genes in tissues relevant to facial morphology. Dendrogram denotes similarity in expression level. TPM, transcripts per million

S4 Fig. Magnitude of variant effect on facial modules, quantified by the Euclidean distance between averaged faces of different genotype groups. The 95% confidence interval was obtained by 5000 bootstraps. The farther away the blue (common) or red (low-freq) rectangular boxes fall from line x=0, the larger the group distances and the greater the magnitude of effects. Common variants that yielded significant GWAS association in the same cohort with the same modules are used as a comparison to low-frequency variants. Genotype groups column indicates the two groups of people of whom the faces were averaged and distance was computed. For example, 0 vs 1/2 means minor allele homozygotes vs the remaining. The following two columns indicate sizes of the two groups in comparison. Low-frequency variants had large effects compared to previously reported common variants, although this could be a result from the much smaller size of carrier group and may not reflect genuine greater effects of low-frequency variants.

S1 Table. Module-wide association results of genes identified by MultiSKAT. Show modules with a p-value < 10E-4.

S2 Table. SKAT and CMC test results of the association between the seven facial genes and NSCL/P, retrieved from a previous exome-wide gene-based association study of NSCL/P

S3 Table. Functional prediction of individual variants in significant genes by CADD GRCh37-v1.4

S4 Table. PhenoScanner lookups for variants in seven significant genes. Show existing associations involving these variants with a p value < 10e-4.