A persistent giant algal virus, with a unique morphology, encodes an

unprecedented number of genes involved in energy metabolism

Romain Blanc-Mathieu[1,2], Håkon Dahle[3], Antje Hofgaard[4], David Brandt[5], Hiroki

Ban[1], Jörn Kalinowski[5], Hiroyuki Ogata[1] and Ruth-Anne Sandaa[6*]

1: Institute for Chemical Research, Kyoto University, Gokasho, Uji, 611-0011, Japan

2: Laboratoire de Physiologie Cellulaire & Végétale, CEA, Univ. Grenoble Alpes,

CNRS, INRA, IRIG, Grenoble, France

3: Department of Biological Sciences and K.G. Jebsen Center for Deep Sea Research,

University of Bergen, Bergen, Norway

4: Department of Biosciences, University of Oslo, Norway

5: Center for Biotechnology, Universität Bielefeld, Bielefeld, 33615, Germany

6: Department of Biological Sciences, University of Bergen, Bergen, Norway

*Corresponding author: Ruth-Anne Sandaa, +47 55584646, ruth.sandaa@uib.no

Abstract

Viruses have long been viewed as entities possessing extremely limited metabolic capacities. Over the last decade, however, this view has been challenged, as metabolic genes have been identified in viruses possessing large genomes and virions—the synthesis of which is energetically demanding. Here, we unveil peculiar phenotypic and genomic features of *Prymnesium kappa* virus RF01 (PkV RF01), a giant virus of the *Mimiviridae* family. We found that this virus encodes an unprecedented number of proteins involved in energy metabolism, such as all four succinate dehydrogenase (SDH) subunits (A–D) as well as key enzymes in the *β*-oxidation pathway. The *SDHA* gene was transcribed upon infection, indicating that the viral SDH is actively used by the virus— potentially to modulate its host's energy metabolism. We detected orthologous *SDHA* and *SDHB* genes in numerous genome fragments from uncultivated marine *Mimiviridae* viruses, which suggests that the viral SDH is widespread in oceans. PkV RF01 was less virulent compared with other cultured prymnesioviruses, a phenomenon possibly linked to the metabolic capacity of this virus and suggestive of relatively long co-evolution with its hosts. It also has a unique morphology, compared to other characterized viruses in the *Mimiviridae* family. Finally, we found that PkV RF01 is the only alga-infecting *Mimiviridae* virus encoding two aminoacyl-tRNA synthetases and enzymes corresponding to an entire base-excision repair pathway, as seen in heterotroph-infecting *Mimiviridae*. These *Mimiviridae* encoded-enzymes were found to be monophyletic and branching at the root of the eukaryotic tree of life. This placement suggests that the last common ancestor of *Mimiviridae* was endowed with a large, complex genome prior to the divergence of known extant eukaryotes.

Importance

Viruses on Earth are tremendously diverse in terms of morphology, functionality, and genomic composition. Over the last decade, the conceptual gap separating viruses and cellular life has tightened because of the detection of metabolic genes in viral genomes that express complex virus phenotypes upon infection. Here, we describe *Prymnesium kappa* virus RF01, a large alga-infecting virus with a unique morphology, an atypical infection profile, and an unprecedented number of genes involved in energy metabolism (such as the tricarboxylic (TCA) cycle and the β-oxidation pathway). Moreover, we show that the gene corresponding to one of these enzymes (the succinate dehydrogenase subunit A) is transcribed during infection and is widespread among marine viruses. This discovery provides evidence that a virus has the potential to actively regulate energy metabolism with its own gene.

Introduction

In their essay "Varieties of Living Things: Life at the Intersection of Lineage and Metabolism," Dupré and O'Malley proposed to address Schrödinger's question "What is Life?" by "*describing a spectrum of biological entities that illustrates why no sharp dividing line between living and non-living things is likely to be useful*" (1). Microbiologists have contributed considerably to this descriptive effort, both by reporting the existence of viruses endowed with genes coding for functions once thought to be exclusive to cellular life and by concomitantly proposing that actively infecting viruses are a "living form" (2–4). Genes encoding elements for photosynthesis (5, 6), carbon metabolism (7), and nitrogen- (8) and sulfur-cycling (9) have been found in bacterial viruses, where they are used to maintain or augment cellular processes during infection and to redirect energy and resources towards viral production (8, 10, 11). Genes for protein synthesis, including translation initiation, elongation, and termination, and a range of aminoacyl-tRNA synthetases have been found in *Mimiviridae*, a group of giant viruses infecting single-celled eukaryotes (12–14). *Mimiviridae* and other large DNA viruses, including some bacterial viruses, also have tRNA genes (15, 16). Ribosomal proteins have recently been reported in viral genomes derived from metagenomes (17). Genes involved in other metabolic processes, such as fermentation (18), glycosylation (19), photosynthesis (20), and rhodopsin (21), are encoded in *Mimiviridae* and other related large eukaryotic DNA viruses. Metabolic genes are frequently observed within virus genomes (20, 22, 23); although they represent a tiny fraction of the viral gene pool, these genes have the potential to dramatically modify the phenotype of an actively infected cell and alter the ecological role of the host (7, 24, 25). The infected host in this state has been referred to as a virocell (2). One might expect that the interplay between viral genes

and host genes in virocells would become increasingly fine-tuned and complex during

prolonged virus–host co-evolution, which also typically leads to lower virulence.

Much of the complexity of virocells may still be undetected, as most *Mimiviridae*

isolated with their natural host (mostly algae) are highly virulent, with several

involved in rapid algal bloom termination events (26).

Viruses of the *Mimiviridae* family are known to infect heterotrophic and

autotrophic microbial eukaryotes. This divide is also reflected in the phylogeny of

these viruses, some of which are classified into two proposed sub-families:

"Megavirinae" and "Mesomimivirinae" (27). The former contains viruses with

genomes larger than 1 Mbp, all isolated from Amoebozoa, while the latter includes

viruses with smaller genomes isolated from haptophyte algae of class

Prymnesiophyceae. Several *Mimiviridae* members outside these two groups have

been characterized to some extent as well, namely, viruses isolated from heterotrophs

(*Cafeteria roenbergensis* virus, CroV; *Bodo saltans* virus, BsV; Choano virus),

autotrophs (*Aureococcus anophagefferens* virus, AaV; Tetraselmis virus 1, TetV;

Pyramimonas orientalis virus, PoV; *Prymnesium kappa* virus RF01, PkV RF01), a

metazoan (Namao virus), and metagenomes (Klosneuviruses). The Mesomimivirinae

sub-family includes viruses infecting bloom-forming hosts, such as *Phaeocystis*

pouchetii, Phaeocystis globosa, and *Prymnesium parvum* (PpV, PgV Group I, and

PpDVAV, respectively) (28–30); it also includes several viruses infecting *Haptolina*

ericina and *Prymnesium kappa*, which normally do not form massive blooms but are

present at low densities in seawater year round (31). In marine environments, viruses

infecting low-density and non-bloom-forming algae may be the most common virus–

host systems—that is, low-density hosts (non-blooming) and viruses that appear to

have co-evolved in response to host growth strategy. Thus far, the only known

representatives of such viruses are *Prymnesium kappa* viruses RF01 (PkV RF01) and

RF02 (PkV RF02), *Haptolina ericina* virus RF02 (HeV RF02), and

Chrysochromulina ericina virus (CeV 01B, infecting *Haptolina ericina*) (32, 33).

Together with PgV, all of these viruses, except for PkV RF01, belong to the sub-

family Mesomimivirinae on the basis of their monophyletic relationship and, in the

case of PgV and CeV, a shared genomic similarity (27). In contrast, phylogenetic

analysis of two partially sequenced marker genes has placed PkV RF01 deep inside

the *Mimiviridae* clade, and characterization of its life cycle has revealed an atypical

infection profile (33). Here, we report new phenotypic features as well as new viral

functions inferred from analysis of the genome sequence of PkV RF01. We found that

this virus has a unique morphology, is less virulent than most other alga-infecting

viruses and possesses an unprecedented number of energy-generating genes. We

uncovered clues suggesting that members of *Mimiviridae* that potentially modulate

the metabolism of their hosts are widespread in the ocean. Our findings of peculiar

genomic features in a persistent virus provide new insights on virus–host coevolution

and may stimulate further advances in modeling the history of their interaction.

Results and Discussion

PkV RF01 has an atypical morphology

The icosahedral PkV RF01 particle is approximately 400 nm in diameter (Fig. 1A-B).

Beneath the capsid, several convoluted inner membranes fill approximately 66% of

the interior. Treatment of chloroform can be used to identify possible functions of

lipid membranes, as it acts to remove lipid molecules that might be essential for

successful infection (34). Some algal viruses in the NCLDV group are sensitive to

chloroform (30, 35, 36) with the suggestions that lipid containing inner or outer

membranes are involved in the infection process (35, 37). In our experiment,

chloroform treatment of PkV RF01 drastically reduced the infectivity of the virus.

(Fig. 1C). As no outer membrane was detected by cryo-electron tomography, the

sensitivity to chloroform might be linked to lipid components in either the capsid or

the inner convoluted membranes. Internal lipid-containing membranes have been

detected in several icosahedral-shaped double-stranded DNA viruses, including algal

viruses belonging to families *Phycodnaviridae* and *Mimiviridae*, mimiviruses, and

various bacteriophages (38–43). In all of these viruses, the inner membranes are

suggested to play a role in the release of the viral nucleoprotein core or genome by

fusing with the host plasma membrane (40, 42, 43). Inner membranes in currently

described NCLDVs more or less adopt the icosahedral morphology defined by the

outer layer of capsomers (44, 45). We detected several convoluted inner membranes

in PkV RF01 that do not follow the structure of the capsid. To our knowledge, this

structural inconsistency has not been previously detected in any double-stranded

DNA viruses, which calls for further investigation to understand the assembly process

of PkV RF01 and how it enters its host. Another striking feature of the PkV RF01

virion is an internal rod-shaped core (ca. 55 nm in diameter), which is filled with

dense material and positioned in the center of the virus particle. Similar features have

been observed in TEM images of large virus-like particles (VLPs) (300–700 nm)

occurring in waste vacuoles of phaeodarian radiolarians collected from different

oceans (46) and in zoospores of the green alga *Chlorococcus minutum* (47). To our

knowledge, however, these features have not been described in isolated viruses thus

far.

PkV RF01 has an atypical infection strategy

Only 2% of the total PkV RF01 viral particles produced during infection of *Haptolina*
ericina UiO028 (He UiO028) were infectious (able to produce progeny) (Table 1).
This infectivity was much lower than that of the other two prymnesioviruses, HeV
RF02 and PkV RF02, which produced 13% and 44% of infectious progeny
respectively (Table 1). The portion of infectious particles of PkV RF01 is low also
when compared to other algal viruses (48, 49). In addition, the latent period of PkV
RF01 was previously reported to be longer (ca. 24–32 h, (33)) in comparison with
other prymnesioviruses (28, 29, 32, 33) and it has been demonstrated that PkV RF01
is also able to infect multi-species (33), that is another unusual trait among algal
viruses (26).

The hosts of PkV RF01, PkV RF02, and HeV RF02 all belong to order the
Prymnesiales, whose members are normally present in low abundance but co-occur
year round (*K*-strategists) (50)*.* PkV RF01, PkV RF02, and HeV RF02 are less
virulent, as shown in the present study, and have longer latent periods compared with
viruses infecting bloom-forming haptophytes (*r*-strategists). Two of these viruses
(PkV RF01 and HeV RF02) are also able to infect multi species (generalists) (33).
Longer replication time and reduced virulence, as hosts becomes scarce, increases the
chances of vertical transmission rather than horizontal transmission of a virus. As
vertical parent-to-offspring transmission depends on host reproduction, it has been
argued that such transmission should select for reduced virulence because the virus
depend on host survival and reproduction for its transmission (51, 52). High
virulence, on the other hand, may be supported by large, dense host populations, as
e.g. algal blooms, because high host densities ensure successful horizontal
transmission of viral progeny to new hosts (51, 53)*.* Viruses infecting the recurrent

bloom-forming haptophytes, *Phaeocystis pouchetii* virus (PpV), and *Phaeocystis*

globosa virus (PgV), are indeed highly virulent with between 60%–100% of virus

particles produced being infectious, resulting in rapid lysis of their hosts (48, 54).

Broad host range might also increase the chance of transmission in an environment

with low host abundances (*K*-strategists). Such strategy requires a tradeoff whereby

the virus decreases its opportunity of transmission by evolving longer replication

times, higher decay rates and reduced infectivity (discussed in (55, 56)). This fits

well with our two multi-species infecting haptophyte viruses, PkV RF01 and

HeV RF02, that have reduced proportions of infectious particles and longer

replication times (33), relative to other haptophyte viruses with restricted host ranges

(specialists) like e.g. the *Emiliania huxleyi* virus (EhV), PpV and PgV.

The balance between fitness traits, such as virulence, latent period and host

range, and tradeoffs is the result of the adaptive evolution between viruses and their

hosts, resulting in relationships spanning from acute to stable coexistence

(persistence). In the ocean, persistent relationships—such as between PkV RF01 and

its hosts—seem to be most common among viruses infecting unicellular algae; this

has been demonstrated by several metabarcoding studies revealing the persistence of

dominance of viral OTUs over several months (57, 58). The atypical infection

strategy of PkV RF01 evokes a persistent nature, different than the vast majority of

other so far characterized algal viruses.

PkV RF01 has the largest genome among algal viruses

The genome of PkV RF01 was assembled as a linear DNA sequence of 1,421,182 bp.

This size is more than twice that of the genome of TetV, which means that PkV RF01

has the largest reported genome of any virus infecting a photosynthetic organism (Fig.

2A). Evidence for the linear structure of this genome is the presence of ~5-kbp

terminal inverted repeats. Despite being phylogenetically more closely related to alga-

infecting *Mimiviridae*, the genome size of PkV RF01 is in the range of heterotroph-

infecting *Mimiviridae*. The overall G+C content of PkV RF01 is 22.8%, which is low

compared with other *Mimiviridae* (23%–41%). Similar to other *Mimiviridae*, the

average G+C content of PkV RF01 in intergenic regions is relatively low, 17.8%.

This lower G+C content may reflect an ongoing loss of G and C nucleotides, more

prevalent in non-coding than coding regions because of weaker background selection

in non-coding regions. The genome of PkV RF01 is predicted to contain 1,161 genes

comprising 1,121 protein-coding DNA sequences (CDSs) and 40 tRNA genes

corresponding to 13 amino acids. Most tRNA genes (30 out of 40) are clustered in

three genomic regions that lack predicted CDSs, a feature also observed in other

Mimiviridae. For example, all tRNAs of TetV ($n = 10$) and CroV ($n = 22$) are encoded

consecutively on the same strand (18, 59). The average CDS length is 1,046 bp

(minimum: 297; maximum: 1,493). Intergenic regions average 217 bp in length, with

a cumulative sum of 244,005 bp, which corresponds to a gene density of 82.8%.

Of the 1,121 predicted CDSs, 641 (57%) exhibited sequence similarities

(BLASTP *E*-value conservative cutoff of 1×10^{-5}) to protein sequences in the

UniRef90 database (Fig. 2A). Among them, 165 were most similar to *Mimiviridae*.

Curiously, among the CDSs most similar to *Mimiviridae*, sixty were closest to

ChoanoVirus which was isolated from choanoflagellates cultures, followed by

Mesomimivirinae ($n = 49$) and Klosneuvirinae ($n = 30$) (Fig. 2B). Among the 181

closest homologs found in eukaryotic organisms 23 were haptophytes. A sequence-

based homology search of corrected nanopore reads and scaffolds composing the

initial assembly against *Lavidaviridae* proteomes (BLASTX; matrix: BLOSUM45, *E*-

value $< 1 \times 10^{-5}$) yielded no significant alignments against any major or minor

Lavidaviridae capsid proteins, which suggests that virophages were absent from the

sample used for sequencing.

A previous analysis of PkV RF01 family-B DNA polymerase (PolB) and the

major capsid protein (MCP) placed this virus in the family *Mimiviridae* (33). We also

recently reported that the PkV RF01 genome has additional NCLDV core genes, such

as A32-like virion packing ATPase (NCVOG0249) and RNApol (RNA pol subunit I

[NCVOG0274] and subunit II [NCVOG0271]), and orthologous genes that are

specific to *Mimiviridae*, namely, MutS7 (NCVOG2626) and asparagine synthase

(AsnS, NCVOG0061) (60). Phylogenetic reconstruction using five NCLDV core

genes confirmed the deep branching of PkV RF01 within the *Mimiviridae* family and

suggested that PkV RF01, along with ChoanoV1, TetV and AaV, is more closely

related to Mesomimivirinae than to Megavirinae (Fig. 3A). In support of this

evolutionary relationship, PkV RF01 has an additional copy of the second largest

RNA polymerase subunit gene (*rpb2*). This *rpb2* duplication is shared with all other

Mimiviridae infecting algae, including Mesomimivirinae members, AaV (whose

second copy is very short), and TetV and was previously proposed as a useful feature

to discriminate between the two main clades (autotroph versus heterotroph-infecting

viruses) within the *Mimiviridae* family (27). This additional *rpb2* copy is not found in

other *Mimiviridae* to the exception of ChoanoV1 whose genome was derived from a

single cell metagenome in choanoflagellates cultures. Phylogenetic analysis indicates

that these two *rbp2* copies were present in the ancestor of alga-infecting *Mimiviridae*

and ChoanoV1 (Fig. 3B). In agreement with the five NCLDV core genes phylogeny,

it suggests that PkV RF01 and ChoanoV1, although evolutionarily distant, are more

related with each other compared to any other *Mimiviridae*.

Out of 1,121 predicted protein-coding genes in the genome of PkV RF01, only

about a third could be annotated with some functional description based on their

sequence homology with characterized proteins. Such a small percentage is typical of

divergent eukaryotic viruses detected for the first time. A total of 339 proteins (30%)

showed significant sequence similarity with proteins in the Cluster of Orthologous

Gene (COG) database (61) (Fig. 4). The distribution of COG functions associated

with these hits was dominated by "Posttranslational modification, protein turnover,

chaperones" (43 proteins) and "Cell wall/membrane/envelope biogenesis" (42

proteins), which is approximately two times more proteins than in other *Mimiviridae*

members except for Tupanvirus. Among other well-represented categories, numbers

of proteins in "Replication, recombination and repair" (36 proteins) and

"Transcription" (23 proteins) were similar to those of other *Mimiviridae*, while the

categories of "Translation, ribosomal structure and biogenesis" (25 proteins) and

"Amino acid transport and metabolism" (20 proteins) were respectively in the same

range or higher than those of heterotroph-infecting *Mimiviridae* (mimiviruses, BsV,

and CroV). Interestingly, 24, 17, and 9 PkV RF01 proteins were respectively assigned

to the categories of "Lipid transport and metabolism", "Carbohydrates transport and

metabolism," and "Energy production and conservation," all much higher compared

with other *Mimiviridae* viruses (Fig. 5).

Similar to other *Mimiviridae*, PkV RF01 encodes several genes involved in

DNA repair, transcription, and translation. Notably, this virus has the full set of

enzymes required for the base excision repair (BER) pathway, which is also the case

for all *Mimiviridae* members except for those with smaller genomes (PgV, CeV, and

AaV). PkV RF01 BER enzymes are closer (i.e., have a greater alignment score) to

heterotroph-infecting *Mimiviridae* than to cellular homologs, thus suggesting that this

pathway was present in the last common ancestor of *Mimiviridae*. According to a

previous phylogenetic analysis, *Mimiviridae* BER enzymes are monophyletic with

regard to *Mimiviridae* and have not recently been acquired from eukaryotes (62).

Unlike alga-infecting *Mimiviridae*, PkV RF01 encodes two amino-acyl tRNA

synthetases (aaRS): an isoleucyl-tRNA synthetase (IleRS; ORF 480) and an

asparaginyl-tRNA synthetase (AsnRS; ORF 764). Both of these synthetases are found

in most lineages of heterotroph-infecting *Mimiviridae* (AsnRS is missing from CroV

and BsV, and IleRS is missing from *Mimivirus* lineage A). Phylogenetic analyses of

these two proteins revealed a deep branching of viral homologs, which formed a

monophyletic clade well separated from cellular homologs (Fig. 6).

A viral-encoded succinate dehydrogenase and energy production genes

We found six predicted protein-coding genes (ORFs 893 to 900) related to energy

production in an 8,026-bp region (Fig. 7A). Four ORFs (ORFs 893 and 898–900)

were predicted to code for all four subunits (SDHA, D, C, and B) of a functional

succinate dehydrogenase (SDH, or Electron Transport Chain Complex II) of the

oxidative phosphorylation pathway (Fig. 7B). In eukaryotes, all four subunits of this

enzyme are encoded in the nuclear genome. This enzyme acts in the mitochondrial

respiratory chain and participates in both the TCA cycle and the respiratory electron

transfer chain. In the TCA cycle, this succinate dehydrogenase oxidizes succinate to

fumarate, while its activity in the inner mitochondrial membrane involves the

reduction of a FAD cofactor followed by electron transfer through three Fe–S centers

to ubiquinone (Fig. 7C).

SDH genes have recently been reported in viral genomes assembled from

environmental samples for which functional experiments cannot be done (63). In a

RT-PCR experiment using primers specific for the PkV RF01 gene for SDHA

(hereafter, *vSDHA*), we detected transcripts of this gene in samples collected 24, 72,

and 96 h post infection (Fig. 8). The *vSDHA* primers were tested on an uninfected

culture to ensure that only the viral version of the *SDHA* gene was amplified (Fig. 9).

The MCP gene of PkV RF01 was used both for protocol optimization and later as an

internal positive control (Fig. 10). Although the transcription of the viral *SDHA*

suggests that the viral SDH is functional, we can only speculate on the possible role

of this enzyme during infection. One possibility is that the viral SDH sustains the

carbohydrate metabolism of infected cells (i.e., virocells) to supply building blocks of

viral particles such as amino acids and to support proper replication of this large virus.

Another possibility is that PkV RF01 uses its SDH as a part of an arms race with its

host to turn on the TCA cycle after the host had turned it off to counter viral

replication, or more simply to boost the energy metabolism of the virocells to

augment the fitness of the host and/or to maximize virus production efficiency.

The discovery of the viral SDH prompted us to search for other potential viral-

encoded SDHA and SDHB homologs in marine metagenomes. These two subunits

(SDHA and SDHB) form the catalytic core containing the redox cofactors that

participate in electron transfer to ubiquinone; they are thus more conserved than

SDHC and SDHD subunits. To test for the presence of this viral SDH in other viruses,

we searched for *vSDHA* and *B* in marine metagenomes of the *Tara* Oceans

expedition. The 50 most-similar and non-redundant SDHA and B sequences predicted

from 101 *Tara* Oceans genome fragments were most likely derived from *Mimiviridae*

viruses (Fig. 11). Indeed, out of 1,113 genes predicted from these 101 genome

fragments, 681 were annotated at some taxonomic level, of which 449 were predicted

to be cellular and 157 viral. Of the 157 viral genes, 146 and 130 had their last

common ancestor in *Mimiviridae* and Mesomimivirinae, respectively. A total of 32 of

the 101-genome fragments contained at least one gene predicted to be of *Mimiviridae*

origin, and the larger the genome fragment, the more *Mimiviridae* genes it was found

to encode (Fig. 11A). Functional analysis indicated that 12 of the 1,113 predicted

genes were NCLDV hallmark genes (encoding five VLTF3s, two capsid proteins, two

PCNAs, two helicases, and one PolB). The high proportion of unknown genes and

genes annotated as *Mimiviridae* in the 101 *Tara* Oceans genome fragments encoding

SDHA or SDHB strongly suggests that these fragments belong to *Mimiviridae*

viruses. This finding demonstrates that the presence of SDH is not restricted to PkV

RF01 and is arguably widespread among marine *Mimiviridae*. According to

phylogenetic analyses of cellular and viral SDHA and SDHB, the viral homologs

form a monophyletic group that branches deeply within eukaryotic lineages (Fig.

11B-C). Long-branch attraction bias could generate such topologies but, as explained

above for the IleRS and AsnRS, it is more likely that the viral SDHA and SDHB were

acquired at an early stage in the radiation of eukaryotic lineages. The transcription of

vSDHA and its occurrence in marine environments calls for further investigation to

understand the biological role and co-evolutionary significance of this viral SDH.

Other genes related to energy production were detected in the 8,026 bp-long

region. ORF 894 and ORF 896, respectively corresponding to cytochrome *c* (CytC)

and cytochrome b6-f complex iron-sulfur (Cyt b6-f) subunits, showed high sequence

conservation with *Chrysochromulina* sp. CCMP291 proteins (78% and 59% amino

acid [aa] identities, respectively). CytC is a short protein (~100 aa) involved in the

oxidative phosphorylation pathway, where it accommodates the transfer of electrons

between the coenzymes Q-cytochrome *c* reductase (complex III) and cytochrome *c*

oxidase (complex IV). The presence of Cyt b6-f between oxidative phosphorylation

genes is puzzling because the cytochrome b6-f complex is involved in photosynthesis.

The core of the chloroplast b6f complex, however, is similar to the analogous

respiratory cytochrome bc(1) complex. The other two predicted ORFs in this region

are similar to ubiquinone biosynthesis protein UbiB (ORF 895) or contain a NAD-

binding domain and a Fe-S cluster (ORF 897) and may thus be associated with

electron transport as well. ORF 897 has two distant (25%–31% aa identity) homologs

in the PkV RF01 genome (ORF 456 and ORF 625).

Some other genes were predicted to encode enzymes involved in pyruvate

metabolism. ORF 79 has sequence homology with L-lactate dehydrogenases; it might

thus catalyze the conversion of lactate to pyruvate, an intermediary compound serving

as a starting point for several major metabolic pathways, such as glycolysis,

gluconeogenesis, and the TCA cycle. ORF 727 was predicted to code for an

isochorismate hydrolase that also produces pyruvate from isochorismate. ORF 24 and

ORF 726 share sequence homology with phosphoenolpyruvate synthase and a partial

pyruvate kinase, respectively. The former catalyzes the conversion of pyruvate to

phosphoenolpyruvate (PEP), while the latter catalyzes the reverse reaction. Formation

of PEP is an initial step in gluconeogenesis.

A nearly complete viral-encoded β-oxidation pathway

In this study, 22 predicted genes were inferred to code for proteins involved in lipid

synthesis or degradation, including key enzymes of the β-oxidation pathway (Table

2). Several genes were predicted to code for lipase-like proteins (ORFs 386, 481, 635,

653, and 690), including a triacylglycerol lipase (ORF 386) that can break down

triacylglycerol into glycerol and fatty acids. Glycerol and fatty acids can be used as a

starting point for ATP production—by glycolysis and β-oxidation, respectively. In the

β-oxidation pathway, fatty acids are fully oxidized to produce acetyl-CoA, which can

then enter the TCA cycle to yield NADH and FADH2; these latter two products can

funnel through to the electron transport chain to produce ATP (Fig. 7C). Each *β*-

oxidation cycle itself also produces NADH and FADH2 cofactors. We found that PkV

RF01 encodes key *β*-oxidation enzymes. First, two distantly related ORFs (ORF 142

and ORF 904 sharing 22% aa identity) have sequence homology with a long-chain

fatty acyl-CoA synthetase. This enzyme catalyzes the formation of fatty acyl-CoA in

the cytosol. Fatty acyl-CoA can be imported to mitochondria using a (carnitine) CoA-

transferase also encoded in PkV RF01 (ORF 33). Once in the mitochondrial matrix,

fatty acyl-CoA serves as a substrate on which an acyl-CoA dehydrogenase (ORF

1046) oxidizes the fatty acyl-CoA and reduces a FAD cofactor to produce a FADH2

cofactor. We identified a 2,4-dienoyl-CoA reductase (ORF 30) that may facilitate the

next oxidation step to produce a NADH cofactor. FADH2 and NADH molecules

produced by a *β*-oxidation cycle can both be oxidized in the electron transport chain

to generate ATP. The enzymes involved in the two intermediate steps following each

oxidation, either an enoyl-CoA hydratase or a β-ketothiolase, were not detected in our

analysis.

Most of these genes have no homologs in reference viral genomes, and, to our

knowledge, this is the first report of a virus possessing proteins directly involved in

lipid-based energy production. By diverting host lipid machinery, interactions of

viruses with lipids or lipid based-structures have long been known to have structural

or signaling roles at different stages of the virus life cycle, such as entry, genome

replication, morphogenesis, and exit (64–66). More recently, several studies on

human viruses (two herpesviruses and one RNA virus) have shown that the metabolic

state of an infected cell can be shifted toward energy generation to support viral

replication (65). These studies have highlighted the increasing abundance—up to 48 h

after HCV infection—of enzymes involved in *β*-oxidation, amino acid catabolism,

and the TCA cycle (67) and an increase in cellular β-oxidation following the release

of free fatty acids caused by Dengue virus-induced autophagy (68). Among algal

viruses, EhV remodels the transcription of host lipid genes for fatty acid synthesis to

support viral assembly (69) and also to generate triacylglycerols stored in the virion

and available as an energy pool in later infection phases (70). Besides diverting the

host metabolism, EhV encodes seven proteins involved in the sphingolipid

biosynthesis pathway (71). This pathway produces a viral sphingolipid that is a

central component of EhV lipid membranes and that can also act as a signaling lipid

and induce programed cell death during the lytic infection phase (72). EhV also

encodes a triglyceride lipase (with detectable homology to predicted PkV RF01

lipases ORF 635 and ORF653) that is highly expressed during late infection

concomitantly with significant up-regulation of host β-oxidation genes (69). These

examples and our observations of several genes involved in β-oxidation clearly show

that viruses can introduce new metabolism-related genes, sometimes representing

entire pathways, into the host, most likely to satisfy the high metabolic requirement of

these giant viruses.

High representation of glycosyltransferases

Compared with other viruses, PkV RF01 was found to encode an unusually high

number of glycosyltransferases (GTs) as well as other carbohydrate-active enzymes.

Automated annotation of GTs (and other carbohydrate-active enzymes) in reference

viral proteomes using dbCAN2 (73) revealed that the largest number of GT domains

was encoded by PkV RF01 ($n = 48$), followed by CeV ($n = 13$), *Mimivirus* members,

and CroV and AaV ($n = 8$–10) (Fig. 12). We uncovered 48 GT domains encoded in

40 ORFs, 8 of which were predicted to encode more than one GT domain. These

domains correspond to 16 different GT families. Most domains were inferred to be

functional, as 31 out of 48 covered at least 70% of the dbCAN2 reference domain,

with coverage ranging from 44% to 99%. GTs were found scattered across the

genome of PkV RF01 but with some local clustering (Fig. 2A), the latter indicating

possible involvement in the same pathway. GT32 was the most represented domain,

with 11 proteins (as annotated by dbCAN2) and potentially three additional proteins

(ORFs 40, 84, and 861). Eight proteins possessed a GT25 domain that can catalyze

the transfer of various sugars onto a growing lipopolysaccharide chain during its

biosynthesis. Among these eight predicted ORFs, four contained an additional non-

overlapping GT domain (two GT2s, one GT6, and one GT60). Functional analyses of

GTs in mimiviruses (or in related *Paramecium bursaria* Chlorella viruses) have

demonstrated that some of these enzymes are functional, being able to modify viral

collagen-like proteins (74) and polymerize sugars (75). Conservation between PkV

RF01 GTs and functionally characterized GTs in viruses and cells is absent or

extremely low, which precludes any predictions as to the specific roles of these

enzymes in the PkV RF01 life cycle. Nevertheless, this putative glycosylation-

conducive autonomy possibly allows the virus to infect a variety of hosts, as the virus

can modify its own glycans, which are used for host recognition, independently of the

host system (76). In alpha-, flavi-, and herpes-viruses, fusion is mediated by viral

glycoproteins (40).

Other carbohydrate-active enzymes in the PkV RF01 genome include seven

glycoside hydrolases (GHs), four carbohydrate esterases (CEs), one polysaccharide

lyase (PL), one carbohydrate-binding module (CBM), and a putative sugar

fermentation stimulation protein A (ORF 1003) possibly involved in maltose

metabolism. These numbers are not excessively high compared with other viruses.

Other detected ORFs were homologous to enzymes involved in carbohydrate

transport and metabolism, notably a transketolase (ORF 528) involved in the pentose

phosphate pathway in all organisms and in the Calvin cycle of photosynthetic

organisms. Finally, we detected a 6-phosphofructo-2-kinase/fructose-2,6-

biphosphatase 2 (ORF 539) and a mannose-1-phosphate

guanylyltransferase/mannose-6-phosphate isomerase (ORF 836) respectively involved

in fructose and mannose metabolism.

Conclusions

The haptophyte virus PkV RF01 has been previously shown to have a longer

replication cycle and a broader host range compared with other prymnesioviruses and

most other algal viruses. Here, we revealed that PkV RF01 has atypical virion

morphology and that infections yield several orders of magnitude fewer infectious

particles than other tested prymnesioviruses. In-depth phylogenetic analysis using

genes conserved in NCLDVs confirmed that PkV RF01 belongs to *Mimiviridae* but is

deeply diverged from existing members, although closer to alga-infecting

Mimiviridae than heterotroph-infecting ones. Unlike other alga-infecting *Mimiviridae*,

however, PkV RF01 has a large genome (1.4 Mb) and contains genes coding for two

aminoacyl-tRNA synthetases and the complete BER pathway. All these features are

conserved in most heterotroph-infecting *Mimiviridae* and therefore must have been

lost in other alga-infecting *Mimiviridae*. This outlier virus features an

unprecedentedly high number of genes involved in energy metabolism and

glycosylation machinery that may enable its longer replication cycle and broader host

range compared with other algal viruses. These genomic and phenotypic features are

suggestive of a persistent infection behavior that probably evolved in response to the

host growth strategy. Because of nutrient limitations, these persistent systems of slow-

growing but ubiquitous hosts with less virulent viruses may represent the most

common type of virocells in oceans.

Materials and Methods

Culturing and infection

All algal host cultures were grown in liquid IMR/2 medium consisting of 70% aged

seawater, 30% distilled water (25 PSU), and additional selenite (10 nM final

concentration). The cultures were kept at 14°C and partially synchronized using a

14:10 h light: dark cycle with irradiance of 100 μmol photons m^{-2} s^{-2} supplied by

white fluorescent tubes. Viruses were produced by adding freshly produced viral

lysate (ca. 2 x 10^8 VLP/mL), propagated three time on the host before added to

exponentially growing host cultures (ca. $5x10^5$ cells/mL) in a ratio of 1:10 volume.

Infection was followed by flow cytometry (FCM) (77, 78) for 72 h by counting viral

particles and host cells, as described in (33). Burst size was calculated as the number

of viral particles released from each host cell, estimated from the total number of host

cells pre-infection and the total number of VLPs produced during the infection cycle

(33).

Infectious progeny

The percentage of viral infectious progeny was determined by comparing the most

probable number (MPN; endpoint dilution (78)) and flow cytometric total counts of

viral particles produced during infection. The number of infectious particles released

in a burst was determined based on the percentage of viral infectivity produced during

the infection cycle and the burst size. Infectivity was tested using *Haptolina ericin*a

UiO028 as a host, and also compared with two other prymnesioviruses, HeV RF02

and PkV RF02 (33), propagated on He UiO028 and *Prymnesium kappa* RCC3423,

respectively.

Briefly, 10× dilution were prepared from fresh viral lysate and added to

exponentially growing host cells in 96-well microtiter plates (eight replicates for each

dilution). The plates were incubated for 7 days under normal incubation conditions.

Cell lysis was measured by monitoring *in situ* fluorescence on a plate reader

(PerkinElmer EnSpire™ 2300 Multilabel Reader) at 460/680 nm. Numbers of

infectious particles were estimated from the proportion of lysed wells using the

MPN_ver4.xls excel spreadsheet from (79).

Sensitivity to chloroform

The effect of chloroform on infectivity, used to infer the presence of a lipid membrane

or lipid molecules in the capsid, was tested by adding 50% (v/v) chloroform to PkV

RF01 lysate. After mixing, the chloroform phase was separated from the solution by

centrifugation at 4,000 *g* for 5 min. The tubes were incubated at 37°C for 2 h with the

lids open to allow evaporation of any remaining chloroform.

Triplicates of exponentially growing He UiO028 cells (1,6 x 10^5 cells /mL)

were incubated with 1:10 volumes of chloroform-treated viruses (ca. 2 x 10^8

VLP/mL). The incubation was followed for 7 days by counting host cells by FCM

(78). Host cells in chloroform-treated or untreated medium at the same ratio used with

the viral lysate were used as controls. Virus propagation was confirmed in lysed

cultures by FCM.

Cryo-electron tomography

A small drop of concentrated PkV RF01 (8x109) was deposited on a glow-discharged,

200-mesh copper grid with holey carbon film (R2/1 Cu 200, Quantifoil Micro Tools

GmbH, Germany). The sample was blotted with filter paper and immediately plunge

frozen in liquid ethane. Grids were transferred under liquid nitrogen to a cryo-transfer

tomography holder (Fishione Instruments, USA) and inserted in a 200-kV

transmission electron microscope (Thermo Scientific Talos F200C) equipped with a

Ceta 16M camera. Tilt series were recorded at 45,000× magnification and −7 μm

defocus between −60° to 60° in 2° increments. Finally, reconstruction, segmentation,

and visualization of the tomograms was performed with IMOD v4.9 software (80).

Purification of viral particles and DNA isolation

Exponentially growing He UiO028 cultures (2 L) were infected with 20 mL of PkV

RF01 and inspected visually for lysis. An uninfected culture (100 mL) was used as a

control. Lysed algal cultures were checked for viruses by FCM counting. Lysed

cultures were first centrifuged to remove algal debris and some bacteria (5,500 rpm

for 15 min). Viruses were then pelleted by ultracentrifugation at 25,000 rpm in a

Beckman Coulter Optima L90K ultracentrifuge for 2 h. The pellets were resuspended

in SM buffer (0.1 M NaCl, 8 mM $MgSO_4 \cdot 7H_2O$, 50 mM Tris-HCl, and 0.005%

glycerin). Viral particles were further purified by Optiprep gradient centrifugation

(81). Fractions were checked for viruses by FCM and for infectivity by infection of

He UiO028.

Isolation of high-quality DNA for sequencing was done by following the

protocol of (82) with some modifications. Viral particles were disrupted by one

round of heating to 90°C for 2 min and then chilling on ice for 2 min. Disodium

ethylenediaminetetraacetic acid and proteinase K at a final concentration of 20 mM

and 100 μg mL^{-1}, respectively, were then added before incubation of the samples for

10 min at 55°C. Sodium dodecyl sulfate at a final concentration of 0.5% (w/v) was

subsequently added, and samples were incubated for an additional 1 h at 55°C.

Double-stranded DNA was then purified from the lysates using a Zymo Genomic

DNA Clean & Concentrator Kit-10 (Zymo Research, Irvine, CA, USA) according to

the manufacturer's protocols. To avoid shearing DNA, gentle pipetting and mixing

(accomplished by turning the tubes instead of vortexing) were performed in all steps.

Genome assembly

Isolated DNA from PkV RF01 was subjected to Illumina TruSeq PCR-free library

preparation (insert size 350 bp). The generated library was sequenced on an Illumina

MiSeq instrument in paired-end mode (2×300 bp) to yield approximately 1.9 million

reads, which corresponds to about 400× coverage. Reads were assembled into 2,498

contigs of 500 bp or more with a total assembly size of 4.75 Mb using Newbler (83).

In addition, a ligation-based $1D^2$ nanopore library (LSK-308) was constructed and

sequenced using an Oxford Nanopore MinION Mk1b device and a FLO-MIN107

flow cell, which resulted in 825 long reads with an N50 of 13.6 kb and a total of 9.89

Mb. To improve the assembly, short-read contigs were manually bridged with the

long reads. Manual assembly using Consed (84) yielded a linear genome sequence of

1.4 Mb with inverted terminal repeats. After assembly, the consensus was polished

using Nanopolish (85) and Pilon (86).

Phylogenetic analyses

Five core genes, SDHA, and SDHB

The phylogenetic position of PkV RF01 was inferred from concatenated protein

alignments of five core nucleocytoplasmic virus orthologous genes (NCVOGs) (87):

D5-like helicase-primase (NCVOG0023), DNA polymerase elongation subunit family

B (NCVOG0038), DNA or RNA helicases of superfamily II (NCVOG0076),

packaging ATPase (NCVOG0249), and Poxvirus Late Transcription Factor VLTF3-

like (NCVOG0262). Sequences were obtained from the NCVOG database

(ftp.ncbi.nlm.nih.gov/pub/wolf/COGs/NCVOG/) (88). Additional sequences were

obtained from genomes retrieved from GenBank and annotated with HMMER v3.12b

using the hmmsearch (89) command with hidden Markov models available in Schulz

et al. (2017) (13). Sequences from each NCVOG were aligned independently using

MAFFT L-INS-i (90). The alignments were trimmed with trimAl v1.2 in *gapyout*

mode (91) prior to concatenation using a custom Python script. Bayesian phylogenetic

trees were inferred with PhyloBayes 1.7 (92) using the CAT model and a GTR

substitution matrix. Four chains were run for 34,500–35,500 generations. The *bpcomp*

command was used to check for convergence and stop when *maxdiff* = 0.3. One chain

was discarded, and a consensus tree was constructed using the remaining three chains.

For phylogenetic analyses of succinate dehydrogenase subunits, top hits of PkV RF01

SDHA and SDHB were retrieved from UniProt (https://www.uniprot.org/) using

online PHMMR searches (https://www.ebi.ac.uk/Tools/hmmer/search/phmmer) and

also from the *Tara* Oceans project using online BLASTP searches (http://tara-

oceans.mio.osupytheas.fr/ocean-gene-atlas/) (Villar et al., 2018). Alignments

generated with MAFFT L-INS-i were filtered with trimAl in gapyout mode.

Maximum-likelihood phylogenies were inferred with RAxML 8.2.9 (93) using the

PROTCATALG model and automatic bootstrapping with the following options: '-N

autoMRE -f a -n autoresult'. Phylogenetic trees of PkV RF01, SDHA, and SDHB

were visualized using iTOL (94).

Rpb2, IleRS, and AsnRS

To reconstruct a phylogenetic tree based on the second largest RNA polymerase

subunit, homologs were recruited by comparing Mimivirus Rpb2 against all proteins

of viruses and selected organisms in the KEGG database using the GenomeNet

BLASTP tool (https://www.genome.jp/). Organisms were manually selected from the

KEGG list to ensure broad taxonomic coverage of the tree of life. The retrieved amino

acid sequences were aligned using MAFFT-LINSI (90) and then trimmed using

trimAl (91) with the following parameters: '-resoverlap 0.5 -seqoverlap 70 -gt 0.8 -st

0.001 -cons 50'. The tree was reconstructed using FastTree (95) as implemented in

the GenomeNet TREE tool (https://www.genome.jp/tools-bin/ete). Isoleucine tRNA

synthase and aspartyl tRNA synthetase viral and cellular homologs were retrieved and

aligned in the same way. Trees were searched using PhyloBayes MPI (96) with the

non-homogeneous CAT+GTR model (97). For each protein three chains were run

until *maxdiff* parameter reach < 0.3 (0.27 for AsnRS and 0.16 for IleRS). One chain

was discarded for IleRS, and a consensus tree was constructed using the remaining

chains.

Gene prediction and functional and taxonomic annotation

GeneMarkS with the option 'virus' (98) predicted 1,121 open reading frames (ORFs)

in the fully assembled genome sequence of PkV RF01, while tRNAscan-SE (99)

predicted 41 tRNAs. PkV RF01 CDS amino acid sequences were searched against

Virus-Host DB (100), RefSeq (101), UniRef90 (102), and COG (61) databases using

BLASTP with an *E*-value of 1×10^{-5} as the significant similarity threshold and

against the Conserved Domain Database (103) using RPS-BLAST with an *E*-value

threshold of 1×10^{-2}. The 10 best hits for each database were compiled in a single file

and manually inspected to transfer annotations of subject sequences to our query. In

ambiguous cases, such as distant homologs (often seen in viral genomes) or unclear or

contradictory annotations of subject sequences, the query was searched against KEGG

genes (104) to allow extensive manual checking using GenomeNet tools

(https://www.genome.jp/; alignment quality, length comparison to canonical genes,

and links with KEGG orthology). We automatically annotated glycosyltransferases

(GTs) and other carbohydrate-active enzymes (glycoside hydrolases, GHs;

polysaccharide lyases, PLs; carbohydrate esterases, CEs; and auxiliary activities,

AAs) in PkV RF01 and all viral genomes in Virus-Host DB (as of June 2018) using

the *hmm* option of the dbCAN2 pipeline and its profile database (73). We retained hits

with E-values $< 1 \times 10^{-5}$ and domain coverage $> 35\%$, which corresponded to default

settings.

Taxonomic and functional analysis of vSDHA homologs in OM-RGCv1

We searched PkV RF01 SDHA and SDHB against OM-RGCv1 (105) using the

Ocean Gene Atlas (106) BLAST-based tool and kept the top 50 hits with significant

E-values for further analysis. We then collected genome fragments (contigs) encoding

these 50 SDHAs and 50 SDHBs by searching via BLASTN for identical hits over full

SDHA or *SDHB* lengths against *Tara* ocean assemblies (downloaded from EBI) used

to construct OM-RGCv1. We predicted ORFs in these genome fragments using

GeneMarkS. The resulting 1,113 amino acid sequences were functionally annotated

by searching against Pfam protein families (107) using profile HMM scan (108) and

also taxonomically using a last common ancestor strategy as in (109); in brief, protein

sequences were searched against a database composed of UniRef cells, MMETSP

(110) and Virus-Host DB (100) data using DIAMOND (111). Selected hits were then

used to derive the last common ancestor of the query using a NCBI taxonomic tree re-

wired to reflect the taxonomy of NCLDVs.

PCR and RT-PCR optimization

We designed specific primers (Table 3) targeting a 256-bp region of the *mcp* gene to use both as an internal control in the RT-PCR and to confirm that our protocols were optimized. For each PCR, a negative control (sterile distilled H_2O) was included. PCR amplifications were carried out in 50-µL total volumes containing 1 µL of template using a DNA HotStarTaq Master Mix kit (Qiagen). The cycling protocol was as follows: 15 min at 95°C, followed by 35 cycles of 30 s at 94°C, 30 s at 59°C, and 30 s at 72°C, with a final extension of 12 min at 72°C.

RT-PCRs were performed using the SuperScript III One-Step RT-PCR with Platinum *Taq* DNA Polymerase system (Thermo Fisher). Cycling conditions were as follows: 16 min at 55°C and 2 min at 94°C, followed by 40 cycles of 15 s at 94°C, 30 s at 49°C, and 30 s at 68°C, and a final extension of 5 min at 68°C.

All PCR products were checked for the correct size on a 1.5% agarose gel stained with GelRed (Biotium). PCR products were further checked by sequencing using BigDye v3.1 (Thermo Fisher) for cycle sequencing (Sekvenseringslaboratoriet, UiB, Norway).

PCR amplification and RT-PCR analysis of *vSDHA*

To investigate whether the *vSDHA* gene is transcribed during infection, an infected culture of He_UiO028 plus PkV RF01 as well as an uninfected He UiO028 culture (control) were set up as described above. Samples were collected at 24, 72, and 96 h post infection from both cultures. RNA was extracted using an RNeasy Plus Universal Mini kit (Qiagen), with gDNA removed in an extra step using a TURBO DNA-free kit (Ambion).

Specific primers were designed to target a 150-bp region of the *vSDHA* gene (Table

3). For each PCR, two negative controls (sterile distilled H_2O and extracted DNA

from He028) were included. As positive controls for the transcription, we used

primers targeting the *mcp* gene (see above). As a positive PCR control, we used

genomic PkV RF01 DNA. PCR amplifications were conducted in 50-µL total

volumes containing 1 µL of template DNA using an ExTaq kit (Takara). The cycling

protocol was as follows: 5 min at 94°C, followed by 35 cycles of 30 s at 94°C, 30 s at

59°C, and 30 s at 72°C, with a final extension of 12 min extension at 72°C.

RT-PCRs were performed using a SuperScript III One-Step RT-PCR with Platinum

Taq DNA Polymerase system (Thermo Fisher). Cycling conditions were as follows:

16 min at 55°C and 2 min at 94°C, followed by 40 cycles of 15 s at 94°C, 30 s at

49°C, and 30 s at 68°C, with a final extension of 5 min at 68°C. PCR products were

checked as described above.

Data availability

684   Raw sequence reads and PkV RF01 genome sequence were deposited at the European
685   Bioinformatics Institute (EMBL-EBI) (https://www.ebi.ac.uk) under project name
686   PRJEB37450. The complete video records of a cryo-electron tomogram of a PkV
687   RF01 virion and sequence data as well as curated gene annotation table as reported in
688   this study are available at https://github.com/RomainBlancMathieu/PkV-RF01.

Acknowledgements

The recording of tilt series was performed with the help of Sebastian Schultz at the
Unit of Cellular Electron Microscopy, the Norwegian Radium Hospital. Initial
sequencing (MiSeq and Pacbio) of PkV RF01 total DNA was performed at the
Norwegian Sequencing Center (https://www.sequencing.uio.no/). We thank Hilde M.
K. Stabell and Solveig Siqveland, Department of Biological Sciences, University of
Bergen, Norway, for technical assistance with molecular biology experiments as well
as Christian Rückert, Bielefeld University, for support in manual finishing of genome
assembly and Minyue Fan, Kyoto University, for assistance in genes analysis. This

work was supported by the Research Council of Norway project entitled "Uncovering
the key players for regulation of phytoplankton function and structure: lesson to be
learned from algal virus-haptophyte coexistence" (VirVar, project number 294364 to
RAS). Additional funding was provided by the European Union Horizons 2020
research and innovation program, grant agreement no. 685778 ("Virus-X") to RAS
and DB. This work was also supported by the Future Development Funding Program
of the Kyoto University Research Coordination Alliance. HO was supported by
JSPS/KAKENHI (No. 18H02279), and Scientific Research on Innovative Areas from
the Ministry of Education, Culture, Science, Sports and Technology (MEXT) of Japan
(Nos. 16H06429, 16K21723, 16H06437). The Super Computer System, Institute for
Chemical Research, Kyoto University, provided computational time. We thank
Barbara Goodson, from Edanz Group (www.edanzediting.com/ac), for editing the
English text of a draft of this manuscript.

Competing interests

712 Authors declare having no competing interests.

References

1. Dupré JO. 2009. Varieties of Living Things: Life at the Intersection of Lineage
and Metabolism. Philosophy & Theory in Biology 1.

2. Forterre P. 2012. Virocell Concept, The, p. a0023264. *In* John Wiley & Sons,
Ltd (ed.), eLS. John Wiley & Sons, Ltd, Chichester, UK.

3. Raoult D, Forterre P. 2008. Redefining viruses: lessons from Mimivirus. Nat
Rev Microbiol 6:315–319.

4. Claverie J-M. 2006. Viruses take center stage in cellular evolution. Genome
Biology 7:110.

5. Mann NH, Cook A, Millard A, Bailey S, Clokie M. 2003. Bacterial
photosynthesis genes in a virus. Nature 424:741–741.

6. Lindell D, Sullivan MB, Johnson ZI, Tolonen AC, Rohwer F, Chisholm SW.
2004. Transfer of photosynthesis genes to and from Prochlorococcus viruses.
PNAS 101:11013–11018.

7. Hurwitz BL, Hallam SJ, Sullivan MB. 2013. Metabolic reprogramming by
viruses in the sunlit and dark ocean. Genome Biol 14:R123.

8. Thompson LR, Zeng Q, Kelly L, Huang KH, Singer AU, Stubbe J, Chisholm
SW. 2011. Phage auxiliary metabolic genes and the redirection of cyanobacterial
host carbon metabolism. Proc Natl Acad Sci USA 108:E757-764.

9.   Anantharaman K, Duhaime MB, Breier JA, Wendt KA, Toner BM, Dick GJ.
733   2014. Sulfur Oxidation Genes in Diverse Deep-Sea Viruses. Science 344:757–
734   760.

735   10.   Lindell D, Jaffe JD, Johnson ZI, Church GM, Chisholm SW. 2005.
736   Photosynthesis genes in marine viruses yield proteins during host infection.
737   Nature 438:86–89.

738   11.   Fridman S, Flores-Uribe J, Larom S, Alalouf O, Liran O, Yacoby I, Salama F,
739   Bailleul B, Rappaport F, Ziv T, Sharon I, Cornejo-Castillo FM, Philosof A,
740   Dupont CL, Sánchez P, Acinas SG, Rohwer FL, Lindell D, Béjà O. 2017. A
741   myovirus encoding both photosystem I and II proteins enhances cyclic electron
742   flow in infected Prochlorococcus cells. Nat Microbiol 2:1350–1357.

743   12.   Raoult D, Audic S, Robert C, Abergel C, Renesto P, Ogata H, La Scola B, Suzan
744   M, Claverie J-M. 2004. The 1.2-megabase genome sequence of Mimivirus.
745   Science 306:1344–1350.

746   13.   Schulz F, Yutin N, Ivanova NN, Ortega DR, Lee TK, Vierheilig J, Daims H,
747   Horn M, Wagner M, Jensen GJ, Kyrpides NC, Koonin EV, Woyke T. 2017.
748   Giant viruses with an expanded complement of translation system components.
749   Science 356:82–85.

750   14.   Abrahão J, Silva L, Silva LS, Khalil JYB, Rodrigues R, Arantes T, Assis F,
751   Boratto P, Andrade M, Kroon EG, Ribeiro B, Bergier I, Seligmann H, Ghigo E,
752   Colson P, Levasseur A, Kroemer G, Raoult D, Scola BL. 2018. Tailed giant
753   Tupanvirus possesses the most complete translational apparatus of the known
754   virosphere. Nat Commun 9:1–12.

755   15.   Miller ES, Heidelberg JF, Eisen JA, Nelson WC, Durkin AS, Ciecko A,
756   Feldblyum TV, White O, Paulsen IT, Nierman WC, Lee J, Szczypinski B, Fraser
757   CM. 2003. Complete Genome Sequence of the Broad-Host-Range Vibriophage
758   KVP40: Comparative Genomics of a T4-Related Bacteriophage. J Bacteriol
759   185:5220–5233.

760   16.   Yoshikawa G, Askora A, Blanc-Mathieu R, Kawasaki T, Li Y, Nakano M,
761   Ogata H, Yamada T. 2018. Xanthomonas citri jumbo phage XacN1 exhibits a
762   wide host range and high complement of tRNA genes. Sci Rep 8:4486.

763   17.   Mizuno CM, Guyomar C, Roux S, Lavigne R, Rodriguez-Valera F, Sullivan
764   MB, Gillet R, Forterre P, Krupovic M. 2019. Numerous cultivated and
765   uncultivated viruses encode ribosomal proteins. Nature Communications 10:752.

766   18.   Schvarcz CR, Steward GF. 2018. A giant virus infecting green algae encodes
767   key fermentation genes. Virology 518:423–433.

768   19.   Piacente F, Gaglianone M, Laugieri ME, Tonetti MG. 2015. The Autonomous
769   Glycosylation of Large DNA Viruses. International Journal of Molecular
770   Sciences 16:29315–29328.

20. Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh D, Denef VJ, McMahon
KD, Konstantinidis KT, Eloe-Fadrosh EA, Kyrpides N, Woyke T. 2020. Giant
virus diversity and host interactions through global metagenomics. Nature 1–7.

21. Needham DM, Yoshizawa S, Hosaka T, Poirier C, Choi CJ, Hehenberger E,
Irwin NAT, Wilken S, Yung C-M, Bachy C, Kurihara R, Nakajima Y, Kojima
K, Kimura-Someya T, Leonard G, Malmstrom RR, Mende DR, Olson DK, Sudo
Y, Sudek S, Richards TA, DeLong EF, Keeling PJ, Santoro AE, Shirouzu M,
Iwasaki W, Worden AZ. 2019. A distinct lineage of giant viruses brings a
rhodopsin photosystem to unicellular marine predators. PNAS 116:20574–
20583.

22. Roux S, Brum JR, Dutilh BE, Sunagawa S, Duhaime MB, Loy A, Poulos BT,
Solonenko N, Lara E, Poulain J, Pesant S, Kandels-Lewis S, Dimier C, Picheral
M, Searson S, Cruaud C, Alberti A, Duarte CM, Gasol JM, Vaqué D, Bork P,
Acinas SG, Wincker P, Sullivan MB. 2016. Ecogenomics and potential
biogeochemical impacts of globally abundant ocean viruses. Nature 537:689–
693.

23. Nishimura Y, Watai H, Honda T, Mihara T, Omae K, Roux S, Blanc-Mathieu R,
Yamamoto K, Hingamp P, Sako Y, Sullivan MB, Goto S, Ogata H, Yoshida T.
2017. Environmental Viral Genomes Shed New Light on Virus-Host
Interactions in the Ocean. mSphere 2.

24. Forterre P. 2013. The virocell concept and environmental microbiology. ISME J
7:233–236.

25. Rosenwasser S, Ziv C, Creveld SG van, Vardi A. 2016. Virocell Metabolism:
Metabolic Innovations During Host–Virus Interactions in the Ocean. Trends in
Microbiology 24:821–832.

26. Coy SR, Gann ER, Pound HL, Short SM, Wilhelm SW. 2018. Viruses of
Eukaryotic Algae: Diversity, Methods for Detection, and Future Directions.
Viruses 10:487.

27. Gallot-Lavallée L, Blanc G, Claverie J-M. 2017. Comparative Genomics of
Chrysochromulina Ericina Virus and Other Microalga-Infecting Large DNA
Viruses Highlights Their Intricate Evolutionary Relationship with the
Established Mimiviridae Family. Journal of Virology 91:e00230-17.

28. Jacobsen A, Bratbak G, Heldal M. 1996. Isolation and characterization of a virus
infecting phaeocystis pouchetii (prymnesiophyceae)1. Journal of Phycology
32:923–927.

29. Baudoux A-C, Brussaard CPD. 2005. Characterization of different viruses
infecting the marine harmful algal bloom species Phaeocystis globosa. Virology
341:80–90.

30. Wagstaff BA, Vladu IC, Barclay JE, Schroeder DC, Malin G, Field RA. 2017.
Isolation and Characterization of a Double Stranded DNA Megavirus Infecting
the Toxin-Producing Haptophyte Prymnesium parvum. Viruses 9.

31. Egge ES, Johannessen TV, Andersen T, Eikrem W, Bittner L, Larsen A, Sandaa
R-A, Edvardsen B. 2015. Seasonal diversity and dynamics of haptophytes in the
Skagerrak, Norway, explored by high-throughput sequencing. Mol Ecol
24:3026–3042.

32. Sandaa RA, Heldal M, Castberg T, Thyrhaug R, Bratbak G. 2001. Isolation and
characterization of two viruses with large genome size infecting
Chrysochromulina ericina (Prymnesiophyceae) and Pyramimonas orientalis
(Prasinophyceae). Virology 290:272–280.

33. Johannessen TV, Bratbak G, Larsen A, Ogata H, Egge ES, Edvardsen B, Eikrem
W, Sandaa R-A. 2015. Characterisation of three novel giant viruses reveals huge
diversity among viruses infecting Prymnesiales (Haptophyta). Virology
476:180–188.

34. Feldman HA, Wang SS. 1961. Sensitivity of Various Viruses to Chloroform.
Proceedings of the Society for Experimental Biology and Medicine 106:736–
738.

35. Martínez M, Boere A, Gilg I, van Lent J, Witte H, van Bleijswijk J, Brussaard C.
2015. New lipid envelope-containing dsDNA virus isolates infecting
Micromonas pusilla reveal a separate phylogenetic group. Aquatic Microbial
Ecology 74:17–28.

36. Mirza SF, Staniewski MA, Short CM, Long AM, Chaban YV, Short SM. 2015.
Isolation and characterization of a virus infecting the freshwater algae
Chrysochromulina parva. Virology 486:105–115.

37. Mackinder LCM, Worthy CA, Biggi G, Hall M, Ryan KP, Varsani A, Harper
GM, Wilson WH, Brownlee C, Schroeder DC. 2009. A unicellular algal virus,
Emiliania huxleyi virus 86, exploits an animal-like infection strategy. Journal of
General Virology, 90:2306–2316.

38. Huiskonen JT, Kivelä HM, Bamford DH, Butcher SJ. 2004. The PM2 virion has
a novel organization with an internal membrane and pentameric receptor binding
spikes. Nat Struct Mol Biol 11:850–856.

39. Yan X, Chipman PR, Castberg T, Bratbak G, Baker TS. 2005. The Marine Algal
Virus PpV01 Has an Icosahedral Capsid with T=219 Quasisymmetry. J Virol
79:9236–9243.

40. Huiskonen JT, Butcher SJ. 2007. Membrane-containing viruses with
icosahedrally symmetric capsids. Curr Opin Struct Biol 17:229–236.

41. King AM, Lefkowitz E, Adams MJ, Carstens EB. 2011. Virus Taxonomy: Ninth
Report of the International Committee on Taxonomy of Viruses. Elsevier.

42. Peralta B, Gil-Carton D, Castaño-Díez D, Bertin A, Boulogne C, Oksanen HM,
Bamford DH, Abrescia NGA. 2013. Mechanism of Membranous Tunnelling
Nanotube Formation in Viral Genome Delivery. PLOS Biology 11:e1001667.

43.   Philippe C, Krupovic M, Jaomanjaka F, Claisse O, Petrel M, le Marrec C. 2018.
Bacteriophage GC1, a Novel Tectivirus Infecting Gluconobacter Cerinus, an
Acetic Acid Bacterium Associated with Wine-Making. Viruses 10.

44.   Yan X, Yu Z, Zhang P, Battisti AJ, Chipman PR, Bajaj C, Bergoin M,
Rossmann MG, Baker TS. 2009. The Capsid Proteins of a Large, Icosahedral
dsDNA Virus. J Mol Biol 385:1287–1299.

45.   Klose T, Rossmann MG. 2014. Structure of large dsDNA viruses. Biol Chem
395:711–719.

46.   Gowing MM. 1993. Large virus-like particles from vacuoles of phaeodarian
radiolarians and from other marine samples. Marine Ecology Progress Series
101:33–43.

47.   Gromov BV, Mamkaeva KA. 1981. A virus infection in the synchronized
population of the *Chlorococcum minutum* zoospores. Algological Studies/Archiv
für Hydrobiologie, Supplement Volumes 252–259.

48.   Bratbak G, Jacobsen A, Heldal M, Nagasaki K, Thingstad F. 1998. Virus
production in Phaeocystis pouchetii and its relation to host cell growth and
nutrition. Aquatic Microbial Ecology 16:1–9.

49.   Zimmerman AE, Bachy C, Ma X, Roux S, Jang HB, Sullivan MB, Waldbauer
JR, Worden AZ. 2019. Closely related viruses of the marine picoeukaryotic alga
Ostreococcus lucimarinus exhibit different ecological strategies. Environ
Microbiol 21:2148–2170.

50.   Thomsen HA, Buck KR, Chavez FP. 1994. Haptophytes as components of
marine phytoplankton. DTU Research Database 187–208.

51.   Berngruber TW, Froissart R, Choisy M, Gandon S. 2013. Evolution of Virulence
in Emerging Epidemics. PLOS Pathogens 9:e1003209.

52.   Day T, Proulx SR. 2004. A General Theory for the Evolutionary Dynamics of
Virulence. The American Naturalist 163:E40–E63.

53.   King AA, Shrestha S, Harvill ET, Bjørnstad ON. 2009. Evolution of Acute
Infections and the Invasion-Persistence Trade-Off. The American Naturalist
173:446–455.

54.   Brussaard CPD, Bratbak G, Baudoux A-C, Ruardij P. 2007. Phaeocystis and its
interaction with viruses. Biogeochemistry 83:201–215.

55.   Leggett HC, Buckling A, Long GH, Boots M. 2013. Generalism and the
evolution of parasite virulence. Trends Ecol Evol (Amst) 28:592–596.

56.   Woolhouse MEJ, Taylor LH, Haydon DT. 2001. Population Biology of
Multihost Pathogens. Science 292:1109–1112.

57. Johannessen TV, Larsen A, Bratbak G, Pagarete A, Edvardsen B, Egge ED,
Sandaa R-A. 2017. Seasonal Dynamics of Haptophytes and dsDNA Algal
Viruses Suggest Complex Virus-Host Relationship. Viruses 9.

58. Gran-Stadniczeñko S, Krabberød AK, Sandaa R-A, Yau S, Egge E, Edvardsen
B. 2019. Seasonal Dynamics of Algae-Infecting Viruses and Their Inferred
Interactions with Protists. Viruses 11:1043.

59. Fischer MG, Allen MJ, Wilson WH, Suttle CA. 2010. Giant virus with a
remarkable complement of genes infects marine zooplankton. Proc Natl Acad
Sci USA 107:19508–19513.

60. Sandaa R-A, Dahle H, Brussaard CPD, Ogata H, Blanc-Mathieu R. Algal
viruses belonging to a subgroup within the Mimiviridae familyEncyclopedia of
Virology, 4th ed. Bamford, D., M. Zuckerman, Elsevier, Academic.

61. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a
tool for genome-scale analysis of protein functions and evolution. Nucleic Acids
Res 28:33–36.

62. Blanc-Mathieu R, Ogata H. 2016. DNA repair genes in the Megavirales
pangenome. Current Opinion in Microbiology 31:94–100.

63. Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO. 2020.
Dynamic genome evolution and complex virocell metabolism of globally-
distributed giant viruses. 1. Nature Communications 11:1710.

64. Ono A. 2010. Viruses and Lipids. Viruses 2:1236–1238.

65. Heaton NS, Randall G. 2011. Multifaceted roles for lipids in viral infection.
Trends Microbiol 19:368–375.

66. Lange PT, Lagunoff M, Tarakanova VL. 2019. Chewing the Fat: The Conserved
Ability of DNA Viruses to Hijack Cellular Lipid Metabolism. Viruses 11:119.

67. Diamond DL, Syder AJ, Jacobs JM, Sorensen CM, Walters K-A, Proll SC,
McDermott JE, Gritsenko MA, Zhang Q, Zhao R, Metz TO, Camp DG, Waters
KM, Smith RD, Rice CM, Katze MG. 2010. Temporal Proteome and Lipidome
Profiles Reveal Hepatitis C Virus-Associated Reprogramming of Hepatocellular
Metabolism and Bioenergetics. PLoS Pathog 6.

68. Heaton NS, Randall G. 2010. Dengue virus induced autophagy regulates lipid
metabolism. Cell Host Microbe 8:422–432.

69. Rosenwasser S, Mausz MA, Schatz D, Sheyn U, Malitsky S, Aharoni A,
Weinstock E, Tzfadia O, Ben-Dor S, Feldmesser E, Pohnert G, Vardi A. 2014.
Rewiring Host Lipid Metabolism by Large Viruses Determines the Fate of
Emiliania huxleyi, a Bloom-Forming Alga in the Ocean[C][W][OPEN]. Plant
Cell 26:2689–2707.

70. Malitsky S, Ziv C, Rosenwasser S, Zheng S, Schatz D, Porat Z, Ben-Dor S,
Aharoni A, Vardi A. 2016. Viral infection of the marine alga Emiliania huxleyi

triggers lipidome remodeling and induces the production of highly saturated
triacylglycerol. New Phytologist 210:88–96.

71. Wilson WH, Schroeder DC, Allen MJ, Holden MTG, Parkhill J, Barrell BG,
Churcher C, Hamlin N, Mungall K, Norbertczak H, Quail MA, Price C,
Rabbinowitsch E, Walker D, Craigon M, Roy D, Ghazal P. 2005. Complete
Genome Sequence and Lytic Phase Transcription Profile of a Coccolithovirus.
Science 309:1090–1092.

72. Vardi A, Van Mooy BAS, Fredricks HF, Popendorf KJ, Ossolinski JE,
Haramaty L, Bidle KD. 2009. Viral glycosphingolipids induce lytic infection
and cell death in marine phytoplankton. Science 326:861–865.

73. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y.
2018. dbCAN2: a meta server for automated carbohydrate-active enzyme
annotation. Nucleic Acids Res 46:W95–W101.

74. Luther KB, Hülsmeier AJ, Schegg B, Deuber SA, Raoult D, Hennet T. 2011.
Mimivirus collagen is modified by bifunctional lysyl hydroxylase and
glycosyltransferase enzyme. J Biol Chem 286:43701–43709.

75. Rommel AJ, Hülsmeier AJ, Jurt S, Hennet T. 2016. Giant mimivirus R707
encodes a glycogenin paralogue polymerizing glucose through α- and β-
glycosidic linkages. Biochem J 473:3451–3462.

76. Parakkottil Chothi M, Duncan GA, Armirotti A, Abergel C, Gurnon JR, Van
Etten JL, Bernardi C, Damonte G, Tonetti M. 2010. Identification of an l-
Rhamnose Synthetic Pathway in Two Nucleocytoplasmic Large DNA Viruses. J
Virol 84:8829–8838.

77. Marie D, Brussaard CPD, Thyrhaug R, Bratbak G, Vaulot D. 1999. Enumeration
of Marine Viruses in Culture and Natural Samples by Flow Cytometry. Appl
Environ Microbiol 65:45–52.

78. Brussaard CPD. 2004. Optimization of procedures for counting viruses by flow
cytometry. Appl Environ Microbiol 70:1506–1513.

79. Jarvis B, Wilrich C, Wilrich P-T. 2010. Reconsideration of the derivation of
Most Probable Numbers, their standard deviations, confidence bounds and rarity
values. J Appl Microbiol 109:1660–1667.

80. Kremer JR, Mastronarde DN, McIntosh JR. 1996. Computer visualization of
three-dimensional image data using IMOD. J Struct Biol 116:71–76.

81. Lawrence JE, Steward GF. 2010. Purification of viruses by centrifugation, p.
166–181. *In* Wilhelm, S, Weinbauer, M, Suttle, C (eds.), Manual of Aquatic
Viral Ecology. American Society of Limnology and Oceanography.

82. Sandaa R-A, E. Storesund J, Olesin E, Lund Paulsen M, Larsen A, Bratbak G,
Ray JL. 2018. Seasonality Drives Microbial Community Structure, Shaping both
Eukaryotic and Prokaryotic Host–Viral Relationships in an Arctic Marine
Ecosystem. Viruses 10.

83.   Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J,
967   Braverman MS, Chen Y-J, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV,
968   Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer MLI,
969   Jarvie TP, Jirage KB, Kim J-B, Knight JR, Lanza JR, Leamon JH, Lefkowitz
970   SM, Lei M, Li J, Lohman KL, Lu H, Makhijani VB, McDade KE, McKenna
971   MP, Myers EW, Nickerson E, Nobile JR, Plant R, Puc BP, Ronan MT, Roth GT,
972   Sarkis GJ, Simons JF, Simpson JW, Srinivasan M, Tartaro KR, Tomasz A, Vogt
973   KA, Volkmer GA, Wang SH, Wang Y, Weiner MP, Yu P, Begley RF, Rothberg
974   JM. 2005. Genome sequencing in microfabricated high-density picolitre
975   reactors. Nature 437:376–380.

976   84.   Gordon D, Green P. 2013. Consed: a graphical editor for next-generation
977   sequencing. Bioinformatics 29:2936–2937.

978   85.   Loman NJ, Quick J, Simpson JT. 2015. A complete bacterial genome assembled
979   de novo using only nanopore sequencing data. Nat Methods 12:733–735.

980   86.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo
981   CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: An Integrated Tool
982   for Comprehensive Microbial Variant Detection and Genome Assembly
983   Improvement. PLOS ONE 9:e112963.

984   87.   Yutin N, Wolf YI, Raoult D, Koonin EV. 2009. Eukaryotic large nucleo-
985   cytoplasmic DNA viruses: Clusters of orthologous genes and reconstruction of
986   viral genome evolution. Virology Journal 6:223.

987   88.   Yutin N, Wolf YI, Koonin EV. 2014. Origin of giant viruses from smaller DNA
988   viruses not from a fourth domain of cellular life. Virology 466–467:38–52.

989   89.   Eddy SR. 2011. Accelerated Profile HMM Searches. PLOS Computational
990   Biology 7:e1002195.

991   90.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software
992   version 7: improvements in performance and usability. Mol Biol Evol 30:772–
993   780.

994   91.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for
995   automated alignment trimming in large-scale phylogenetic analyses.
996   Bioinformatics 25:1972–1973.

997   92.   Lartillot N, Lepage T, Blanquart S. 2009. PhyloBayes 3: a Bayesian software
998   package for phylogenetic reconstruction and molecular dating. Bioinformatics
999   25:2286–2288.

1000   93.   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and
1001   post-analysis of large phylogenies. Bioinformatics 30:1312–1313.

1002   94.   Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the
1003   display and annotation of phylogenetic and other trees. Nucleic Acids Res
1004   44:W242–W245.

95.   Price MN, Dehal PS, Arkin AP. 2009. FastTree: Computing Large Minimum
Evolution Trees with Profiles instead of a Distance Matrix. Mol Biol Evol
26:1641–1650.

96.   Lartillot N, Rodrigue N, Stubbs D, Richer J. 2013. PhyloBayes MPI:
phylogenetic reconstruction with infinite mixtures of profiles in a parallel
environment. Syst Biol 62:611–615.

97.   Lartillot N, Philippe H. 2004. A Bayesian Mixture Model for Across-Site
Heterogeneities in the Amino-Acid Replacement Process. Mol Biol Evol
21:1095–1109.

98.   Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training
method for prediction of gene starts in microbial genomes. Implications for
finding sequence motifs in regulatory regions. Nucleic Acids Res 29:2607–2618.

99.   Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and
context for analysis of transfer RNA genes. Nucleic Acids Res 44:W54-57.

100. Mihara T, Nishimura Y, Shimizu Y, Nishiyama H, Yoshikawa G, Uehara H,
Hingamp P, Goto S, Ogata H. 2016. Linking Virus Genomes with Host
Taxonomy. Viruses 8:66.

101. Pruitt KD, Tatusova T, Maglott DR. 2007. NCBI reference sequences (RefSeq):
a curated non-redundant sequence database of genomes, transcripts and proteins.
Nucleic Acids Res 35:D61–D65.

102. Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH. 2015. UniRef clusters: a
comprehensive and scalable alternative for improving sequence similarity
searches. Bioinformatics 31:926–932.

103. Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY,
Geer RC, He J, Gwadz M, Hurwitz DI, Lanczycki CJ, Lu F, Marchler GH, Song
JS, Thanki N, Wang Z, Yamashita RA, Zhang D, Zheng C, Bryant SH. 2015.
CDD: NCBI's conserved domain database. Nucleic Acids Res 43:D222–D226.

104. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a
reference resource for gene and protein annotation. Nucleic Acids Res 44:D457–
D462.

105. Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G,
Djahanschiri B, Zeller G, Mende DR, Alberti A, Cornejo-Castillo FM, Costea
PI, Cruaud C, d'Ovidio F, Engelen S, Ferrera I, Gasol JM, Guidi L, Hildebrand
F, Kokoszka F, Lepoivre C, Lima-Mendez G, Poulain J, Poulos BT, Royo-
Llonch M, Sarmento H, Vieira-Silva S, Dimier C, Picheral M, Searson S,
Kandels-Lewis S, Tara Oceans coordinators, Bowler C, de Vargas C, Gorsky G,
Grimsley N, Hingamp P, Iudicone D, Jaillon O, Not F, Ogata H, Pesant S,
Speich S, Stemmann L, Sullivan MB, Weissenbach J, Wincker P, Karsenti E,
Raes J, Acinas SG, Bork P. 2015. Ocean plankton. Structure and function of the
global ocean microbiome. Science 348:1261359.

106. Villar E, Vannier T, Vernette C, Lescot M, Cuenca M, Alexandre A, Bachelerie
P, Rosnet T, Pelletier E, Sunagawa S, Hingamp P. 2018. The Ocean Gene Atlas:
exploring the biogeography of plankton genes online. Nucleic Acids Res
46:W289–W295.

107. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M,
Richardson LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L,
Piovesan D, Tosatto SCE, Finn RD. 2019. The Pfam protein families database in
2019. Nucleic Acids Res 47:D427–D432.

108. Eddy SR. 1998. Profile hidden Markov models. Bioinformatics 14:755–763.

109. Carradec Q, Pelletier E, Silva CD, Alberti A, Seeleuthner Y, Blanc-Mathieu R,
Lima-Mendez G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, Bertrand A,
Engelen S, Madoui M-A, Méheust R, Poulain J, Romac S, Richter DJ,
Yoshikawa G, Dimier C, Kandels-Lewis S, Picheral M, Searson S, Jaillon O,
Aury J-M, Karsenti E, Sullivan MB, Sunagawa S, Bork P, Not F, Hingamp P,
Raes J, Guidi L, Ogata H, Vargas C de, Iudicone D, Bowler C, Wincker P. 2018.
A global ocean atlas of eukaryotic genes. Nature Communications 9:373.

110. Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA,
Armbrust EV, Archibald JM, Bharti AK, Bell CJ, Beszteri B, Bidle KD,
Cameron CT, Campbell L, Caron DA, Cattolico RA, Collier JL, Coyne K, Davy
SK, Deschamps P, Dyhrman ST, Edvardsen B, Gates RD, Gobler CJ,
Greenwood SJ, Guida SM, Jacobi JL, Jakobsen KS, James ER, Jenkins B, John
U, Johnson MD, Juhl AR, Kamp A, Katz LA, Kiene R, Kudryavtsev A, Leander
BS, Lin S, Lovejoy C, Lynn D, Marchetti A, McManus G, Nedelcu AM,
Menden-Deuer S, Miceli C, Mock T, Montresor M, Moran MA, Murray S,
Nadathur G, Nagai S, Ngam PB, Palenik B, Pawlowski J, Petroni G, Piganeau
G, Posewitz MC, Rengefors K, Romano G, Rumpho ME, Rynearson T,
Schilling KB, Schroeder DC, Simpson AGB, Slamovits CH, Smith DR, Smith
GJ, Smith SR, Sosik HM, Stief P, Theriot E, Twary SN, Umale PE, Vaulot D,
Wawrik B, Wheeler GL, Wilson WH, Xu Y, Zingone A, Worden AZ. 2014. The
Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP):
Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through
Transcriptome Sequencing. PLOS Biology 12:e1001889.

111. Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using
DIAMOND. Nat Methods 12:59–60.

1079

Figure legends

FIG 1 PkV RF01 morphology and reduced viral infectivity under chloroform

treatment. (A) Screen shot of a cryo-electron tomogram of a PkV RF01 virion. (B)

Composite image of 61 cryo-electron tomograms (−60 to 60°, imaged every 2°).

Purple, capsid; green, inner membrane consisting of multiple irregular, convoluted

membranes; blue, internal rod-shaped core filled with dense material. The full set of

records is available on GitHub (see Data availability section). Scale bar, 100 nm. (C)

Reduction of PkV RF01 infectivity with chloroform. Experiments were set up in

triplicate, and host cells were counted by flow cytometry. Chloroform-treated PkV

RF01 was added to exponentially growing He UiO028 cells in a 1:10 volume ratio.

Controls were He UiO028 cells incubated with chloroform-treated medium (Control

A), untreated PkV RF01 (Control B), and untreated medium (Control C). SDs are

indicated with error bars.

FIG 2 Structure and gene taxonomic composition of the PkV RF01 genome sequence.

(A) Rhizome and genomic features of the PkV RF01 genome. As illustrated by the

rhizome (inner part of the figure), ORFans comprise the largest set of PkV RF01

genes, and a substantial portion (15%) have their best BLAST hits (in UniRef90)

against "*Mimiviridae*." Colors indicate taxonomic origin. Intergenic regions are white.

Percentage hits per taxonomic group higher than 5% of total genes are indicated. In

the outermost ring, rectangles indicate the positions of glycosyltransferases (white),

lipid-related enzymes (black), and succinate dehydrogenase genes (red), and the

numbers correspond to *Mimiviridae* key enzymes (1 and 3: DNA-directed RNA

polymerase II subunits 1 and 2, respectively; 2: DNA mismatch repair protein MutS7;

4: Packaging ATPase; 5: VLTF3, 6: Major capsid protein; 7: Eukaryotic translation

1105   initiation factor 4E; 8: Asparagine synthase; 9: DNA polymerase family B). The ring

1106   adjacent to the outermost ring shows GC skew over a 10-KB window. (B) Taxonomic

1107   breakdown of 180 genes with best hits to virus genes. Mega, Megavirinae; AaV,

1108   Aureococcus anophagefferens virus; TetV, Tetraselmis virus 1; PoV, Pyramimonas

1109   orientalis virus.

1110

FIG 3 Phylogenetic evidence for PkV RF01 as a distant relative of

"Mesomimivirinae." (A) Bayesian phylogenetic tree of NCLDVs reconstructed from

a concatenated alignment of five core nucleocytoplasmic virus orthologous genes.

Values at branches are posterior probabilities support. The tree was rooted using

Poxviridae as outgroup. The scale bar indicates substitutions per site. (B) Maximum

likelihood phylogenetic tree of cellular and NCLDV DNA-directed RNA polymerase

subunit beta (RPB2). Values at branches are Shimodaira-Hasegawa-like local support.

1118

1119   **FIG 4** COG functional distribution of 339 proteins encoded by PkV RF01.

1120

1121   **FIG 5** Comparative COG functional distribution among Mimiviridae members. COG

1122   sequences were automatically searched against the proteomes of each virus using

1123   BLASTP with an E-value of 1×10^{-5} as the significant similarity threshold.

1124

1125   **FIG 6** Bayesian phylogenetic trees of two viral amino-acyl tRNA synthetases and

1126   their cellular homologs. (A) Isoleucine tRNA synthases. (B) Aspartyl tRNA

1127   synthetases. Branches supported by posterior probability (PP) values >70% are

1128   indicated by circles whose diameters are proportional to the PP value.

1129

FIG 7 Genes in PkV RF01 predicted to encode enzymes of oxidative phosphorylation

and β-oxidation pathways. (A) Gene organization in the succinate dehydrogenase-

containing region. (B) Schematic representation of the canonical enzymatic complex

II in the mitochondrial membrane. (C) Location of succinate dehydrogenase in the

TCA cycle and electron transport chain as known in plants and a schematic

reconstruction of the PkV RF01-encoded β-oxidation metabolic pathway.

1136

FIG 8 The viral SDHA gene is transcribed during infection. Gels of PCR and RT-

PCR in combination with a TURBO DNA-free kit. Samples were taken 24, 72, and 96

h after infection. (A) PCR with *vSDHA*-specific primers was used to check for the

presence of genomic DNA after RNA isolation treated with 1x and 2x DNAse, in the

upper and lower panels respectively. P, positive control (PKV RF01 genomic DNA);

N, negative control (sdH$_2$0). (B) RT-PCR of RNA samples using *vSDHA*-specific

primers. M, DNA marker (MassRuler DNA Ladder Mix, Thermo Fisher, 80 to 10,000

bp).

1145

1146   **FIG 9** PCR optimization and confirmation of the SDHA gene in the PkV RF01

1147   genome. (A–B) Results of PCR with SDHA primers using genomic PkV RF01 DNA

1148   (A) and genomic He UiO028 DNA (B) as templates. Lanes 1 and 9, DNA ladder; 2–

1149   7, optimization of the PCR annealing temperature from 55°C (2) to 60°C (7); 8,

1150   negative control (sdH2O).

1151

FIG 10 PCR and RT-PCR optimization using an internal control gene (mcp).

PCR and RT-PCR were carried out after removal of genomic DNA using a TURBO

DNA-free kit. Samples were taken 24, 72, and 96 h after infection. Two different

protocols, both provided in the TURBO DNA-free kit manual, were used to optimize

the reactions. (A) PCR check for the presence of genomic DNA after RNA isolation

treated with 1x and 2x DNAse, in the upper and lower panels respectively. P, positive

control (PkV RF01 genomic DNA); N, negative control (sdH20). (B) Result of RT-

PCR of samples harvested 24, 72 and 96 h post infection. M, DNA marker

(MassRuler DNA Ladder Mix, Thermo Fisher, 80 to 10,000 bp).

1161

FIG 11 Origin of PkV RF01 SDHA and SDHB and their most similar homologs in

Tara Oceans metagenomes. (A) Taxonomy of genes predicted in *Tara* Oceans

metagenome assembled-genome fragments encoding the 50 SDHAs and SDHBs most

similar to PkV RF01 genes (for genome fragments having at least five predicted

genes). (B and C) Phylogenetic trees of viral and cellular SDHAs (B) and SDHBs (C).

Clades in green contain PkV RF01 SDHA or SDHB and their 50 most similar hits

identified in *Tara* Oceans metagenomes (predicted to be *Mimiviridae* homologs from

A). Red, eukaryotic phyla; black, unclassified eukaryotes. Trees are rooted with

Proteobacteria and Firmicutes homologs (not shown). Circles indicate branches with

posterior probability support ≥ 50%.

1172

1173 **FIG 12** Comparative distribution of glycosyltransferase domains among viruses.

1174

Tables

TABLE 1 Infection parameters of *Prymnesium kappa* viruses RF01 and RF02 and *Haptolina ericina* virus RF02.

Viral species and hosts	Infectious progeny/mL (MPN)	Host cells/mL (FCM)[a]	Total VLP/mL (FCM)	Burst size (VLP)[b]	Infectivity (%)[c]	Infectious particles in a burst[d]
PkV RF01 (He UiO028)	2.9×10^6 (± 0.2)	4.9×10^5	1.8×10^8 (±0.9)	363	2	6
PkV RF02 (Pk RCC3423)	2.2×10^8 (± 0.2)	4.6×10^5	5.0×10^8 (±0.1)	1093	44	483
HeV RF02 (He UiO028)	5.8×10^7 (±0.2)	4.9×10^5	4.4×10^8 (±0.0)	907	13	119

VLP, virus-like particle; MPN, most probable number; FCM, flow cytometry.
[a]Measurement performed in duplicates
[b]The number of viral particles released from each host cell, estimated from the total number of host cells pre-infection and the total number of VLPs produced during the infection cycle.
[c]Estimated as the percentage of infectious progeny of all VLPs produced during the infection cycle.
[d]Number of infectious particles released per host cell.

TABLE 2 Gene related to lipid metabolism.

ORF	Annotation	KEGG orthology	Pathway
30	2,4-dienoyl-CoA reductase, mitochondrial [EC:1.3.1.34]	K13236	Beta oxidation
33	Putative CoA-transferase	NS	Beta oxidation
121	glycerophosphoryl diester phosphodiesterase	K01126	Glycerophospholipids metabolisms
138	Fatty acid synthase (FASN)	K00665	Fatty acid biosynthesis
142	Long-chain-fatty-acid--CoA ligase ACSBG [EC:6.2.1.3]	K15013	Fatty acid degradation /biosynthesis / Beta Oxidation
175	Acetyl-CoA carboxylase / biotin carboxylase 1 [EC:6.4.1.2 6.3.4.14 2.1.3.15]	K11262	Fatty acid biosynthesis
236	Glutaryl-CoA dehydrogenase [EC:1.3.8.6]	K00252	Fatty acid degradation
293	Lysophospholipase like	NS	NS
357	Lysophospholipase like	NS	NS
386	Triacylglycerol lipase [EC:3.1.1.3]	K01046	Glycerolipid metabolism
481	Lipase like	NS	NS
635	Lipase-like	NS	NS
653	Lipase-like	NS	NS
690	Lipase-like	NS	NS
774	Lysophospholipid Acyltransferases [EC:2.3.1.22]	K14457	Glycerolipid metabolism
694	Lipase esterase (Carbohydrate esterase CE10)	NS	NS
695	Lipase esterase (Carbohydrate esterase CE10)	NS	NS
886	Stearoyl-CoA desaturase (Delta-9 desaturase) [EC:1.14.19.1]	K00507	Biosynthesis of unsaturated fatty acids
902	Fatty acid synthase (FASN)	K00665	Fatty acid biosynthesis

904	Long-chain-fatty-acid--CoA ligase ACSBG [EC:6.2.1.3]	K15013	Fatty acid degradation /biosynthesis / Beta Oxidation
1016	Cyclopropane-fatty-acyl-phospholipid synthase [EC:2.1.1.79]	k00574	NS
1046	Acyl-CoA dehydrogenase	K06445	Fatty acid degradation / Beta oxidation

1186

1187

1188 **TABLE 3** Forward and reverse PCR primers for amplification of vSDHA and MCP

1189

Primer name	Sequence (5´- 3´)	PCR product (bp)
vSDHA-F1	ATGTGCCGAGAAGCTCCTAA	154
vSDHA-R1	CTGCACAGGCTGTTCGATAA	
PkV-RF01-MCP-F	GATGAACCTTGCCCACAACT	256
PkV-RF01-MCP-F	GTGCATGGTACGTTTTCGTG	

1190

Figure 1

Figure 2

A

Archaea

Bacteria 23%

Eukaryotes 16%

Other viruses

Other NCLDV

15% *Mimiviridae*

43% ORFans

B

NCLDV

Mimiviridae

Mesomimivirinae (27%)

ChoanoV1 (33%)

Klosneuvirinae (17%)

AaV (6%)

Mega. (6%)

Phycodnaviridae (3%)

Marseilleviridae (3%)

Phages (2%)

Cafeteriavirus (2%)

TetV (<1%)

PoV (<1%)

Pithovirus (<1%)

Figure 3

A

Poxviridae
Asfarviridae
1
Iridoviridae
1
Marseilleviridae
1
Phycodnaviridae
1
0.95

Megavirinae
1
Hokovirus
0.7
Indivirus
1
Klosneuvirus
1
Catovirus
1
BodoSaltans
1
Klosneuvirinae

Cafeteria roenbergensis virus
1
NamaoVirus
0.75
Tetraselmis virus 1
Aureococcus anophagefferens virus
1
Prymnesium kappa RF01
0.88
ChoanoVirus
0.73
0.97

Organic Lake phycodnavirus 1
Organic Lake phycodnavirus 2
1
Phaeocystis globosa virus
1
Chrysochromulina ericina virus
1
Mesomimivirinae

Tree scale: 1

B

Bacterial / Mitochondrial
1.0
Poxviridae
1.0
Euk. RNAP-I
1.0
1.0
Iridoviridae
1.0
Orpheovirus
1.0
Pithovirus sibericum
1.0
Cedratvirus
0.9
African swine fever virus
1.0
Kaumoebavirus
0.3
Pacmanvirus
0.3
Euk. RNAP-II
0.9
1.0
Emiliania huxleyi virus
1.0
Mollivirus sibericum
1.0
Pandoraviruses
1.0
0.9
0.1
Euk. RNAP-IV/V
0.7
Noumeavirus
0.8
1.0
Marseillevirus marseillevirus
0.9
1.0
Euk. RNAP-III
1.0
1.0
Archaeal
0.9

Megavirinae
1.0
Namao virus
0.9
Cafeteria roenbergensis virus
0.4
Klosneuvirinae
1.0
1.0
Tetraselmis virus 1
ChoanoVirus
1.0
Prymnesium kappa virus RF01 ORF665
0.7
Chrysochromulina ericina virus
0.8
Phaeocystis globosa virus
1.0
Organic Lake phycodnavirus 1
0.9
Organic Lake phycodnavirus 2
1.0
0.4
1.0
Tetraselmis virus 1
Aureococcus anophagefferens virus
1.0
1.0
Prymnesium kappa virus RF01 ORF356
0.4
ChoanoVirus
0.7
Organic Lake phycodnavirus 2
Organic Lake phycodnavirus 1
1.0
Chrysochromulina ericina virus
1.0
Phaeocystis globosa virus

Tree scale: 1

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A — PCR

B — RT –PCR

1 x DNase treatment

2 x DNase treatment

Figure 9

Figure 10

Figure 11

Figure 12