Root electrotropism in Arabidopsis does not depend on auxin distribution but
requires cytokinin biosynthesis

Maddalena Salvalaio[1], Nicholas Oliver[1,2], Deniz Tiknaz[1], Maximillian Schwarze[1,3],
Nicolas Kral[1,4], Soo-Jeong Kim[1,5], Giovanni Sena[1*]

[1] Department of Life Sciences, Imperial College London, UK
[2] Present address: Department of Physics, Freie Universität Berlin, Germany
[3] Present address: School of Biosciences, University of Birmingham, UK
[4] Present address: Phytoform Labs Ltd., Lawes Open Innovation Hub, Harpenden, UK
[5] Present address: Department of Engineering, University of Cambridge, UK
12
[*] Corresponding author

AUTHOR CONTRIBUTIONS
M.Sa., N.O. and G.S. conceived and designed the experiments. M.Sa. contributed to the
development of the V-box, performed and analysed most of the electrotropism assays. N.O.
contributed to the development of the V-box, performed and analysed some of the WT
electrotropism assays. D.T. contributed to the development of the V-slide and performed some
of the electrotropism assays. M.Sc. performed the root excision experiments and the
electrotropism assays for *cyp735a1*, and analysed the data. N.K. contributed to the
development of the V-slide and performed the *R2D2* and *WAVE131Y* imaging. S.J.K.
performed the heat-treatment experiments. M.Sa. and G.S. prepared the manuscript, with all
the authors contributing to the Methods section.

ABSTRACT

An efficient foraging strategy for plant roots relies on the ability to sense multiple physical and
chemical cues in soil and to reorient growth accordingly (tropism). Root tropisms range from
sensing gravity (gravitropism), light (phototropism), water (hydrotropism), touch
(thigmotropism) and more. Electrotropism, also known as galvanotropism, is the phenomenon
of aligning growth with external electric fields and currents. Although observed in a few species
since the end of the 19th century, the molecular and physical mechanism of root electrotropism
remains elusive, limiting the comparison to more defined sensing pathways in plants.
Here we provide a first quantitative and molecular characterisation of root electrotropism
in the model system *Arabidopsis thaliana,* showing that it does not depend on an asymmetric
distribution of the plant hormone auxin, but that instead it requires the biosynthesis of a second

hormone, cytokinin. We also show that the dose-response kinetics of the early steps of root

electrotropism follows a power law analogous to the one observed in some physiological

reactions in animals.

A future full molecular and quantitative characterisation of root electrotropism would

represent a step forward towards a better understanding of signal integration in plants, and an

independent outgroup for comparative analysis of electroreception in animals and fungi.

INTRODUCTION

Plant roots navigate the complex soil environment in search of water and nutrients, through

various sensing mechanisms reorienting their growth towards or away from signal sources

(tropism) (Muthert et al., 2020). For example, hydrotropism redirects growth towards high

moisture (Miyazawa and Takahashi, 2019), (negative) phototropism redirects away from light

sources (Kutschera and Briggs, 2012), and gravitropism induces growth downwards, following

the gravity vector (Su et al., 2017). The integration of overlapping and frequently contradicting

molecular and physical signals is as critical for plant roots as it is for other soil organisms.

Although many molecular aspects of a few root tropisms have been understood (Muthert

et al., 2020), several key sensing mechanisms remain elusive. One of these is the capacity of

plant roots to sense electric fields (electrotropism or galvanotropism) (Navez, 1927). The local

physical and chemical properties of soil determine the presence of mobile electrical charges

and the generation of spontaneous electric fields (Pozdnyakov and Pozdnyakova, 2002). For

example, electrostatic fields can appear from charge separation in minerals like clay (Ward,

1990), by electrokinetic conversion caused by a conducting fluid like water flowing through

rocks (Revil et al., 2003) or by local accumulation of mineral ions important for plant

metabolism such as ammonium and nitrate ions. Local electric fields in soil can have biological

origin as well, for example from negatively charged bacteria (Olitzki, 1932) or from ions and

charged molecules released by microorganisms (Chabert et al., 2015) or plant roots

(Takamura, 2006). All this suggests that transient electrostatic fields in soil encode unique

information regarding the localisation of water, micronutrients and organisms, and it is

plausible that a sensing mechanism to detect such signal provides a selective advantage.

In fact, the reception of electric fields (electroreception) has been observed in vertebrates

(Crampton, 2019) and invertebrates (Clarke et al., 2013), including common model systems

such as *C. elegans* (Sukul and Croll, 1978) and *Dyctostylium* (Shanley et al., 2006), as well

as in fungi (McGillivray and Gow, 1986). Interestingly, electric sensing structures have been

identified only in a few aquatic animal species (Peters et al., 2007) and more recently in

bumble-bees (Sutton et al., 2016).

First recorded in 1882 (Elfving, 1882) and rediscovered at the start of the 20th century

(Ewart and Bayliss, 1905), root electrotropism has been studied sporadically in maize (*Zea*

mays), peas (*Pisum sativum*) and bean (*Vigna mungo*) but with contradicting results
(Wolverton et al., 2000). Crucially, the anatomical and molecular details of sensing electric
fields are still largely unknown in roots. A quantitative description of root electrotropism's
kinetics is also missing, preventing a comparative analysis with animal electroreception and
electrotaxis.
79
RESULTS
A novel root electrotropism assay
To study root electrotropism in the plant model system *Arabidopsis thaliana,* we developed
a new setup to stimulate, image and quantitate electrotropism in its primary root. Briefly, roots
were grown vertically in a transparent chamber (V-box) containing a buffered liquid medium
and two immersed electrodes connected to a power supply to generate a uniform electric field
and an ionic current perpendicular to the growing roots (Fig. 1a and Methods).
To maintain constant temperature and minimise the pH gradient generated by electrolysis,
the liquid medium was continuously circulated in a closed loop between the V-box, a cooled
water bath and a 2-litre reservoir bottle (Supplemental Fig. S1 and Methods). In each V-box,
the primary roots of five seedlings were imaged every 10 minutes with a camera mounted in
front of the V-box (Supplemental Fig. S1 and Methods). We measured the actual field
generated in the medium by immersing voltmeter probes in the two neighbouring positions of
each seedling: as expected, the field measured in the medium was lower than the nominal
imposed in air between the two electrodes, but still relatively uniform across the five positions
(Supplemental Fig. S2a). At the same time, we measured the current passing through the
circuit (Supplemental Fig. S2b and Methods).
To quantify the root response to the applied electric field, we took an image of the roots
every 10 minutes and measured the angle between the root tip and the vertical gravity vector
(Fig. 1b).
To confirm that the circulation of the liquid medium was effective in damping any pH
gradient created by electrolysis and in eliminating any chemotropic effect, we imposed a field
of 1.5 V/cm in a V-box without plants for one hour while maintaining liquid circulation. We then
turned off the field and the liquid circulation, immediately positioned the plants in the V-box
and imaged the roots for the next 80 minutes: we did not observe any significant deviation in
root growth direction (t-test between 1.5 V/cm and 0 V/cm at 80 m, p=0.859), compared to
roots in a V-box that never experienced the electric field (Supplemental Fig. S2c), indicating
that no significant pH gradient was left in the medium.
108

Root tip reorientation in external electric fields

We measured the response of wild-type (WT) *Arabidopsis* primary roots to a continuous

electric field. The distribution of reorientations to a range of fields intensities between 0.5 V/cm

and 2.5 V/cm shows a quick tropism towards the negative electrode, or cathode (Fig. 1c).

We wondered how much of this effect was due to trivial electrostatic, *i.e.* the physical pull

towards the negative electrode due to a hypothetical net positive charge accumulated on the

root tip, rather than a more complex biological response involving molecular signalling. To

address this, we deactivated the roots by immersing them in a 50 ℃ bath for 10 min, until

growth and gravitropism were suppressed (0/9 roots growing and bending to gravity after

exposure to 50 ℃ , *vs.* 10/10 after exposure to 23 ℃), transferred to the V-box and exposed

them to a 2.0 V/cm electric field: the root response shows that this simple treatment was

sufficient to completely inhibit electrotropism when compared to roots kept at a standard 23

℃ temperature (t-test between 50 ℃ and 23 ℃ at 5 h, $p<0.001$), strongly suggesting that

electrostatic alone could not explain the root response and that this is in fact a biological

phenomenon (Supplemental Fig. S3).

Response curves

The progressively sharper root tip reorientation as the field intensity was increased (Fig.

1c) suggests that the sensing mechanism is not acting as a simple on/off switch but that it can

distinguish electric fields of different strengths. To quantitate this, we plotted the orientation

angle at 5 h ("response") as a function of the electric field intensity ("stimulus"), and found a

best fit with a power function with exponent 0.45 (Fig. 2a, left), indicating that the resolution of

the sensor is higher at low-intensity stimuli (steep response curve) than at high-intensity stimuli

(shallow response curve). The analogous response curve as a function of the measured

current intensity is best fit with a power function with exponent 0.33 (Fig. 2a, right).

Since the root tip reorientations did not show any obvious overshoot (Fig. 1c), we looked

more closely at the angular velocity: the maximum average velocity in WT roots was reached

when the root tip was between 10 and 20 degrees orientation to the gravity vector and then

progressively decreased as the root tip approached its maximum reorientation (Supplemental

Fig. S4a). This indicates that the mechanism is able to sense and respond differently to the

changing relative orientation of the tip with the external electric field, and to progressively slow

down the root tip rotation as it approaches the target orientation. Moreover, the maximum

angular velocity appears to be roughly proportional to the electric field strength (Fig. 2b, left),

although this is much less evident as a function of the electric current (Fig. 2b, right).

Root tips are not damaged

Early reports noted that protracted exposure to external fields could cause physical

damage to plant root tips (Wawrecki and Zagórska-Marek, 2007). To control whether this was

the case in our experimental conditions, we developed a simple chambered slide (V-slide) to

be mounted on a standard confocal microscope stage (Supplemental Fig. S5a and Methods)

with electrodes on the chamber's sides and a circulating liquid medium for temperature control

similar to the one implemented in the V-box (Supplemental Fig. S5b). Seedlings from the

Arabidopsis transgenic line constitutively expressing the yellow fluorescent cell-membrane

marker WAVE 131Y (Geldner et al., 2009) were mounted on the V-slide and imaged at cellular

resolution while exposed to a 1.0 V/cm electric field (Methods). The field of view in our time-

lapse images comprises the meristem, the transition zone and the distal elongation zone, but

no cellular pattern perturbation was noticeable in any of these regions when compared to roots

not exposed to the field (Supplemental Fig. S6). Also, the time-lapse suggests that asymmetric

cell expansion in the elongation zone is causing the bending, as expected from other examples

of root tropism (Gilroy, 2008).

To further confirm that roots exposed to the electric field are not damaged and maintain

gravitropic response, we monitored root tips for 2 h after the electric field had been turned off,

observing a clear gravitropic behaviour (Supplemental Fig. S7).

Regions of competence

Since the root bending occurs in the elongation zone, we wanted to identify the region

responsible for sensing the electric field. We excised distal fragments of the root at 125 µm,

300 µm, 400 µm and 500 µm from the tip (Fig. 3a), and then exposed the cut root to 1.5 V/cm

(Methods).

In the first 2 h (Fig. 3b left), roots cut at 400 µm from the tip turned to angles

indistinguishable from uncut roots exposed to the same field (Wilcoxon between 400 µm cut

and uncut at 1.5 V/cm at 2 h, p=0.211), while roots cut at 500 µm from the tip did not turn and

at 2 h showed orientations indistinguishable from uncut roots not exposed to a field (t-test

between 500 µm cut at 1.5 V/cm and uncut at 0 V/cm at 2 h, p=0.164). These results indicate

that the 400 µm distal fragment is not necessary for the early (2 h) electrotropic response,

while the 500 µm distal fragment is; since the elongation zone involved in the bending spans

an extended region proximal to the 500 µm cut, we conclude that in *Arabidopsis* roots the

region between 400 µm and 500 µm from the tip is necessary for early sensing an external

electric field.

Between 2 h and 5 h of exposure (Fig. 3b right), while roots cut at 300µm on average

continue to turn like the uncut roots (t-test between 300 µm cut and uncut at 1.5 V/cm at 5 hrs,

p=0.071), roots cut at 400 µm quickly fail to sustain their response and at 5 h they show tip

orientations on average different than those of uncut roots exposed to the same field (t-test
between 400 μm cut and uncut at 1.5V/cm at 5 hrs, p<0.001). These results indicate that the
300 μm distal fragment is not necessary to maintain the electrotropic response up to 5hrs,
while the 400 μm distal fragment is; we conclude that in *Arabidopsis* roots the region between
300 μm and 400 μm from the tip is necessary for prolonged sensing of the imposed electric
field.

Interestingly, previous studies suggested that the movement of Ca++ ions accumulated in
the mucilage at the very tip of the root might be involved in the electrotropic sensing (Marcum
and Moore, 1990). Since any excision above 125 μm from the tip essentially removes all the
mucilage from the root, our results disprove this hypothesis for *Arabidopsis*.

Auxin distribution is not altered by the electric field
The fact that roots without tips could still respond to the electric field shows that even a
major disruption of the stereotypical auxin redistribution mechanism is not sufficient to inhibit
electrotropism.

On the other hand, an asymmetric accumulation of auxin is required for asymmetric cell
elongation and root bending in some tropisms, as suggested by the classic Cholodny-Went
model (Thimann and Went, 1937). To test whether this model applied to electrotropism, and
whether the external electric field is sufficient to induce an asymmetric distribution of auxin in
the root, we exposed roots expressing the auxin-sensitive fluorescent reporter *R2D2* (Liao et
al., 2015) to a field of 1.0 V/cm for 30 minutes, before quickly mounting them on a microscope
slide and imaging them with a confocal microscope (Methods). A ratiometric quantification of
R2D2 signal (Kral et al., 2016) in each epidermal cell (Fig. 4a and Methods) showed that the
average auxin response measured in the epidermal cells on the side facing the negative
electrode and on the side facing the positive electrode were statistically indistinguishable (Fig.
4b; Methods), both in the distal (Wilcoxon test, p=0.696) and the proximal (Wilcoxon test,
p=0.843) region of the root, indicating that auxin is not asymmetrically distributed after
exposure to the electric field.

Moreover, both sides were also indistinguishable from the average between the two sides
in roots not exposed to the field (mock), both in the distal (Wilcoxon test between the negative
exposed side and mock, p=0.775; and between the positive exposed side and mock, p=0.881)
and in the proximal (Wilcoxon test between the negative exposed side and mock, p=0.041;
and between the positive exposed side and mock, p=0.098) regions.

Auxin transport is not necessary for early response
Previous work indicated that auxin transport inhibitors can inhibit electrotropism response
in maize roots (Ishikawa and Evans, 1990). To verify this in *Arabidopsis,* and further explore

the role of auxin in root electrotropism, we used N-1-naphthylphthalamic acid (NPA) to inhibit

polar auxin transport (Sabatini et al., 1999) while exposing the roots to the electric field. We

tested 0.1 µM, 1.0 µM and 10 µM NPA and found that 10 µM NPA was the lowest

concentration of NPA to inhibit gravitropism (Methods), a known auxin-dependent tropism.

We pre-treated the seedlings for 3 hours in liquid medium containing 10 µM NPA

(Methods) and then transferred them to a V-box containing 10 µM NPA in the medium and the

reservoir. In the first 1 hour of field exposure (Fig. 4c left) the NPA-treated roots reoriented at

angles indistinguishable from those of untreated ones (Wilcoxon test between NPA-treated

and untreated at 1 h, $p=0.350$). From 1 to 5 hours of exposure (Fig. 4c right), NPA-treated

roots respond less than the untreated (t-test between NPA-treated and untreated at 5 h,

$p<0.001$), but still significantly more than the untreated and not exposed to the field (t-test

between untreated at 1.5 V/cm and 0 V/cm at 5 h, $p=0.001$). These results indicate that auxin

polar transport is not necessary for an early electrotropic response, but might play a role in

maintaining a long-term orientation.

To further explore the role of auxin transport, we tested the mutants of PIN2, an auxin

cellular exporter (Chen et al., 1998), and mutants of AUX1, an auxin cellular importer

(Marchant et al., 1999). Since *pin2* and *aux1* mutants do not respond to gravity, to obtain a

sufficient number of roots growing vertically in preparation for the electrotropic assay, we

wrapped the sides and bottom of the nursery boxes with aluminium foil to induce negative root

phototropism towards the bottom of the box (Methods). The same setup was used to

germinate WT plants to be compared with these to mutants.

Roots of *pin2* mutants (Fig. 4d) showed a significant response (paired Wilcoxon between

the time-points 0 h and 2 h, $p<0.01$; and between the time-points 0 h and 5 h, $p<0.01$), but

weaker than WT (t-test between WT and *pin2* at 2 h, $p<0.01$; and between WT and *pin2* at 5

h, $p<0.01$) in the same conditions. Their angular velocity did not show an obvious decrease

before reaching the target orientation, as with WT roots (Supplemental Fig. S4b).

Analogously, roots of *aux1* mutants (Fig. 4d) showed a significant early response (paired

t-test between the time-points 0 h and 2 h, $p<0.01$), but weaker than WT (t-test between WT

and *pin2* at 2 h, $p<0.01$; t-test between WT and *pin2* at 5 h, $p<0.01$) in the same conditions.

Interestingly, *aux1* roots on average failed to maintain their orientation for longer period (paired

t-test between the time-points 0 h and 5 h, $p=0.051$). Moreover, their angular velocity on

averaged decreased while approaching the final orientation, as with WT roots (Supplemental

Fig. S4b).

Taken together, these results suggest that although auxin transport seems to play a role

in maintaining a sustained response to the electric field, it is not necessary in triggering early

electrotropism.

Cytokinin biosynthesis is necessary for electrotropism

Given our results suggesting a limited role for auxin during electrotropism, we wondered

which other plant hormones might be involved instead. Root hydrotropism, or the growth

towards high concentration of water, was also previously shown to be largely independent

from auxin distribution in *Arabidopsis* (Shkolnik et al., 2016), while it requires biosynthesis and

asymmetric distribution of cytokinin (Chang et al., 2019). Drawing an analogy with

hydrotropism, we considered the possibility that electrotropism might act through cytokinin as

well.

To test this hypothesis, we analysed root electrotropism in triple mutants of *AtIPTs*, a

family of adenosine phosphate-isopentenyltransferases required for the first step of isoprenoid

cytokinin biosynthesis (Miyawaki et al., 2006; Kamada-Nobusada and Sakakibara, 2009).

Within the tested mutants, although a high degree of redundancy is expected among the

AtIPTs, we found two distinct behaviours: the triple mutants *atipt1,3,5* and *atipt3,5,7* both

responded strongly to the electric field (Fig. 5a) (t-test between 1.5 V/cm and 0 V/cm at 5 h,

p<0.001 in both cases), while the triple mutants *atipt1,3,7* and *atipt1,5,7* responded well in the

first 2 h of exposure (Fig. 5b, left) (t-test between 1.5 V/cm and 0 V/cm at 2 h, p<0.001 for

both mutants) but showed a much weaker, although still significant, response at 5 h (Fig. 5b

right) (t-test between 1.5 V/cm and 0 V/cm at 5 h, p<0.01 for *atipt1,5,7* and p<0.001 for

atipt1,3,7; t-test between WT and mutant both at 1.5 V/cm at 5 h, p<0.001 for *atipt1,5,7* and

p<0.01 for *atipt1,3,7*). These results suggest that cytokinin biosynthesis is in part required for

long-term root electrotropism, and that for this phenotype *AtIPT1 and AtIPT7* dominate their

family in a redundant way (the response is the weakest when both are mutated).

To confirm the role of cytokinin, we also tested the requirement for *CYP735A1,* a

cytochrome P450 monooxygenase enzyme acting downstream of *AtIPTs* and necessary for

the biosynthesis of the *trans*-zeatin (*tZ*) variation of cytokinin (Takei et al., 2004). Interestingly,

roots of *cyp735a1* mutants completely failed to respond to the electric field (Fig. 5c), with tip

orientations indistinguishable from that of WT roots not exposed to the field (t-test between

cyp735a1 exposed to 1.5 V/cm and WT not exposed at 5 h, p=0.748). Crucially, this phenotype

could be rescued with the addition of the cytokinin *trans*-Zeatin (*tZ*) to the medium (Fig. 5c) (t-

test between *cyp735a1* +10nM *tZ* at 1.5 V/cm and 0 V/cm, p<0.001), while the same treatment

did not affect WT response (Fig. 5c).

Taken together, these results indicate that cytokinin is necessary for electrotropism in

Arabidopsis roots.

To further investigate a possible parallel between the molecular pathways involved in

electrotropism and hydrotropism, we tested mutants of *MIZU-KUSSEIN1 (MIZ1),* which is

necessary for hydrotropism (Kobayashi et al., 2007) and its functional cytokinin asymmetric

distribution in roots (Chang et al., 2019). Translational fusion reporters have shown MIZ1

localisation in the lateral root cap and the cortex of meristem and elongation zone (Dietrich et
al., 2017): although the root tip is not necessary for hydrotropic response (Dietrich et al., 2017),
this requires MIZ1 in the transition zone (Dietrich et al., 2017).

When we exposed roots of *miz1* mutants to 1.5 V/cm (Fig. 6) they showed an unperturbed
electrotropic response in the first 2 h (Wilcoxon test between *miz1* and WT at 2 h, p=0.859)
and perhaps a weakened response at 5 h, although with only a weak statistical significance
(t-test between *miz1* and WT at 5 h, p=0.033), indicating that MIZ1 is not necessary for early
root electrotropism.

Taken together, these surprising results indicate that cytokinin plays an important role in
root electrotropism, but that the underlying molecular pathway differs early on from that of
hydrotropism.

DISCUSSION

The capability of plant roots to sense and combine numerous physical and chemical signals
in soil is quite extraordinary, notably in the absence of a centralised processing system. The
more we understand about the physical and molecular mechanism behind the various root
tropisms, the closer we will get to a complete understanding of signal integration in plants. In
this work we focused on the little-studied phenomenon of root electrotropism and present its
first quantitative characterisation in the flowering plant model system *Arabidopsis thaliana.*

A non-trivial observation from this work is that plant roots respond to weak (order of 1 V/cm
and 100 μA) external electric fields and currents in a progressive way, with an increasingly
stronger tendency to align with the field as the field and current intensities increase. This is
interesting, because it reveals a sophisticated way to discriminate between highly charged
particles (*e.g.* ions, micro-organisms, other plant roots) and weakly charged ones. At the same
time, the absence of overshooting also points to a mechanism that perhaps can be modelled
along the lines of a proportional–integral–derivative control system (Chevalier et al., 2019).
Moreover, we show that the kinetics (response curve) of root electrotropism follows a power-
law with exponents similar to those that have been traditionally measured in physiological
responses to external stimuli in animal systems (Stevens, 1970): our results suggest that a
power-law response to external stimuli might be a universal feature across life kingdoms, and
that it does not require a nervous system. Moreover, it might reveal constraints on the genetic
architecture of the underlying sensory system (Adler et al., 2014).

A second, perhaps unexpected, result from this work is the identification of two regions in
Arabidopsis roots required for the detection of the field: the section between roughly 400 μm
and 500 μm from the tip is necessary for the early detection (within 2 h from exposure), while
the section between roughly 300 μm and 400 μm is necessary for a prolonged detection (up
to 5 h from exposure). Interestingly, these sections seem to correspond to the well-

characterised anatomical transition zone, between the meristem and the elongation zone in
roots, which has been previously involved in root sensing (Muthert et al., 2020). In future
works, it will be crucial to discern whether the electrotropism mechanism depends on a
relatively large region like the transition zone or if it can be narrowed to a more specific cell
population therein.

Finally, we present evidence against the assumption that auxin asymmetric distribution is
required for electrotropism in *Arabidopsis*, similarly to what found for hydrotropism (Shkolnik
et al., 2016) and phototropism (Kimura et al., 2018). Instead, we show that cytokinin is required
for a full electrotypic response, although its response-specific regulation seems to act through
a different pathway than the one established for hydrotropism.

Overall, our results show that root electrotropism requires a sensing mechanism likely
localised in the transition zone, a limited role for auxin but an important role for cytokinin. The
latter could be involved in a signalling or regulatory mechanism in the transition zone, in an
actuator mechanism (tissue bending) in the elongation zone, or both.

Cytokinins have been shown to regulate root meristem activity and size by controlling cell
proliferation (Beemster and Baskin, 2000) and the developmental progression from
proliferation to elongation in the transition zone (Ioio et al., 2007). It has been suggested that
root bending in hydrotropism is based on asymmetric distribution of cytokinin in the meristem
to induce asymmetric cell proliferation (Chang et al., 2019). This seems an unlikely
mechanism for electrotropism, because roots without meristem (a 400μm segment cut from
the tip) still respond to the electric field, suggesting that root bending in this case does not
depend much on cell proliferation. An alternative mode of action for cytokinins during
electrotropism could be based on its regulatory action in the transition zone, where an
asymmetric delay in elongation would result in root bending. Although we have shown that
MIZ1 is not required for root electrotropism, it is still possible that a MIZ1-independent
mechanism could generate an asymmetry in cytokinin distribution in the transition zone.

More recently, cytokinins have also been involved in stress response through regulation
of downstream factors and through crosstalk with other hormones (Li et al., 2021; Wu et al.,
2021). This signalling role could be relevant during electrotropism, especially since the
transition zone has often been associated with signal integration in the root (Baluška et al.,
2010).

Future work on root electrotropism should focus on testing these hypotheses regarding
the role of cytokinin and on illuminating the still unknown molecular mechanism involved in
sensing an electric field or current.

METHODS

Plant material

Wild-type and mutant *Arabidopsis thaliana* plants were all from the Columbia (Col-0) ecotype; the following mutant alleles were used: *aux1-7* (NASC id 9583) for *aux1*; *eir1-1* (NASC id 8058) for *pin2;* SALK_093028C (NASC id N654306) for *cyp735a1*, *miz1-1* for *miz1* (courtesy of Prof. A. Kobayashi); *atipt* triple mutants as previously described with *atipt1-1, atipt3-2, atipt5-2, atipt7-1* (Miyawaki et al., 2006) (courtesy of Prof. O. Leyser).

The fluorescent line *WAVE131Y* is expressing *pUBQ10::WAVE131:YFP* (NASC id N781535); the fluorescent line *R2D2* is expressing *RPS5A-mDII-ntdTomato, RPS5A-DII-n3xVenus* (courtesy of Dr. Teva Vernoux).

Seeds were imbibed in water and kept in the dark for 2 days at 4 °C, in order to synchronise germination. All seeds were surface sterilised using 50% Haychlor bleach and 0.0005% Triton X-100 for 3 minutes and then rinsed 6 times with sterilised milliQ water. Seed germination protocols are described in the experiment-specific Methods sections below. Unless otherwise specified, all experiments were conducted with primary roots of seedlings 5-8 days post-germination, with roots approximately of the same length to be in the field of view of the V-box camera.

Electrotropism assay (V-box)

Seeds were sown individually inside PCR tubes filled with 1X MS gel medium: 0.44% Murashige and Skoog (MS) Basal medium (Sigma-Aldrich, M5519), 0.5% sucrose, 0.05% MES hydrate (Sigma-Aldrich M8250), 0.8% agar (Sigma-Aldrich 05040), pH adjusted to 5.7 with TRIS HCl (Fisher-Scientific 10205100). The PCR tubes had their end cut out to allow the root to grow through, and placed in a 3D-printed (Ultimaker 2+) holder inserted in a Magenta box (Sigma-Aldrich V8380). The Magenta box was filled with 150 ml of 1/500X MS liquid medium (0.00088% MS Basal medium, 0.5% sucrose, 0.05% MES hydrate, pH adjusted to 5.7 with TRIS HCl) to reach the end of the PCR tubes. These germination, or "nursery", boxes were placed in a growth chamber at 22 °C, with a 16 h/8 h light/dark photoperiod and light intensity 120 µmol m^{-2} s^{-1}

In preparation for the electrotropism assay, each PCR tube containing a single seedling was transferred to a modified 3D-printed holder in a Magenta box filled with 1/500X MS liquid medium. The modified module consisted of a main body with five holes for the PCR tubes containing the seedlings, and two side clips to position the electrodes consisting of platinum-iridium (Platinum:Iridium = 80:20; Alfa Aesar 41805.FF) foils (Fig. 1a), which were connected to an external power supply. In this paper, we refer to the Magenta box and its holder as the "V-box". In addition to the five holes for the plants, four extra holes were designed: two at each end of the front side to pump the medium out of the V-box right on top of the two electrodes,

and two on the back to pump the medium in, using a tubing system and peristaltic pumps

(Verdeflex AU R2550030 RS1) to circulate the medium at a speed of 1 ml/sec to and from a

2 l reservoir bottle (Extended Data Fig. 1). This configuration was designed and tested to

ensure a slow and symmetric flow of medium in the box, eliminating any biased effect of the

flow on the roots, which was confirmed by the control experiments at 0 V/cm, performed with

the pump circulating the medium at the same speed as in the exposure experiments. Crucially,

between the V-box and the reservoir bottle, the tubings were immersed in a cooled water bath

(Grant Instruments, LTDGG) maintained at the constant temperature of 19 ℃ (Extended Data

Fig. 1), which was enough to maintaining the medium inside the V-box at a constant 22 ℃, as

measured. All electrotropic experiments were performed at constant illumination.

The electric field was generated with a power supply attached to the platinum-iridium foil

electrodes that were immersed in the liquid medium in the V-box. Standard electric wires were

soldered on the top of two electrodes, always kept outside the liquid medium to avoid

contaminants from the solder. The voltage was set constant on the power supply, while the

current was measured independently with a multimeter in series.

In order to record the movement of the roots over time, a Raspberry Pi camera module V2

(913-2664) connected to a Raspberry Pi board module B+ (137-3331) was used. The

Raspberry Pi was programmed to take a picture every 10 minutes, using the command *crontab*

in the local Raspbian OS.

WT electrotropism assays (Fig. 1) were performed with the following sample sizes N and

number of replicates R: 0 V/cm, N=10, R=2; 0.5 V/cm, N=9, R=2; 1.0 V/cm, N=8, R=2; 1.5

V/cm, N=21, R=5; 2.0 V/cm, N=18, R=6; 2.5 V/cm, N=20, R=7; foil-wrapped and 1.5 V/cm

(control for *pin2* and *aux1*), N=15, R=3.

Mutants electrotropism assays (Fig. 4,5,6) were performed with the following sample sizes

N and number of replicates R: *aux1* at 1.5 V/cm, N=18, R=4; *aux1* at 0 V/cm, N=14, R=3; *pin2*

at 1.5 V/cm, , N=13, R=4; *pin2* at 0 V/cm, , N=13, R=3; *atipt3,5,7* at 1.5 V/cm, N=15, R=3;

atipt3,5,7 at 0 V/cm, N=10, R=2; *atipt1,3,5* at 1.5 V/cm, N=15, R=3; *atipt1,3,5* at 0 V/cm, N=10,

R=2; *atipt1,5,7* at 1.5 V/cm, N=14, R=3; *atipt1,5,7* at 0 V/cm, N=10, R=2; *atipt1,3,7* at 1.5

V/cm, N=14, R=5; *atipt1,3,7* at 0 V/cm, N=10, R=2; *cyp735a1* at 1.5 V/cm, N=20, R=4; *miz1*

at 1.5 V/cm, N=9, R=2; *miz1* at 0 V/cm, N=10, R=2

The control for medium circulation efficiency (Fig. S2) was performed with the following

sample sizes N and number of replicates R: 1.5 V/cm, N=10, R=2; 0 V/cm, N=10, R=2.

The experiment showing gravitropism after 2 hours of electric field (Fig. S7) was performed

with a sample size N=10 and replicates R=2.

High temperature treatment

Wild-type Col-0 seeds were germinated and grown in the nursery boxes as described in

the Electrotropism Assay section. The boxes undergoing treatment were then immersed in a

water bath set at 50 °C, for 10 minutes. The PCR tubes containing the seedlings were then

transferred to a V-box, exposed to an electric field of 2.0 V/cm for 5 hours and imaged, as

described in the Electrotropism Assay section.

Electrotropism assays for treated roots were performed with the following sample sizes N

and number of replicates R: 23℃, N=18, R=6; 50℃, N=10, R=2.

For the gravitropism control, seedlings were treated with high temperature as described

above and then transferred on 1X MS agar plates, to complete the assay as described in the

Gravitropic Assay section.

Gravitropic assays for treated roots were performed with the following sample sizes N and

number of replicates R: 23℃, N=10, R=2; 50℃, N=9, R=1).

Electrotropism on microscope (V-slide)

Seeds of the transgenic reporter line *WAVE131Y* (see Plant Material section) were

germinated on 1X MS agar medium (0.44% MS Basal medium, 0.5% sucrose, 0.05% MES

hydrate, pH adjusted to 5.7 with TRIS HCl, 0.8% agar) in square plates kept vertical in a

growth chamber at 22 °C, with a 16 h/8 h light/dark photoperiod.

At 2 days post-germination the seedlings were mounted on the V-slide with 1/500X MS

liquid medium (0.00088% MS Basal medium, 0.5% sucrose, 0.05% MES hydrate, pH adjusted

to 5.7 with TRIS HCl). The V-slide was then connected to the medium perfusion system

circulating the same 1/500X MS medium, and the V-slide's electrodes were connected to the

power supply (Voltcraft PS-1302-D).

Imaging was performed on a Leica SP5 laser scanning confocal microscope with 20X air

objective, while the liquid medium was circulated and the electric field maintained constant.

Fluorophore was excited with the 514 nm Argon laser line and the emission collected with

PMT detector at 524-570 nm. Images were collected at intervals of 2 minutes.

Root tip excisions

Seeds were germinated in nursery boxes as described.

3 days post-germination seedlings still in their PCR tubes, as previously described, were

transferred to hard 1X MS agar medium (0.44% MS Basal medium, 0.5% sucrose, 0.05%

MES hydrate, pH adjusted to 5.7 with TRIS HCl, 5.0% agar) where the roots were manually

dissected using a dental needle (Sterican, 27G) under a dissecting microscope (Nikon

SMZ1000) at 180X magnification, following the published protocol (Kral et al., 2016).

After root tip excision, the PCR tubes containing the seedlings were immediately moved

into the V-box for the electrotropism assay.

Electrotropism assays on excised roots were performed with the following sample sizes N

and number of replicates R: 125 µm, N=16, R=4; 300 µm, N=7, R=2; 400 µm, N=10, R=2; 500

µm, N=10, R=2; uncut at 0 V/cm, N=10, R=2; uncut at 1.5 V/cm, N=21, R=5.

R2D2 reporter

Seeds of the transgenic reporter line *R2D2* (see Plant Material section) were germinated

in nursery boxes and at 3days post-germination exposed to 1.0 V/cm for 30min in V-boxes,

as described in the Electrotropism Assay section. After exposure, *R2D2* roots were quickly

mounted on standard microscope slides with sterile deionised water and imaged using Leica

SP5 laser scanning confocal microscope, with 63X water immersion objective.

R2D2 expresses two versions of a protein that forms a complex with auxin: DII-

n3xVENUS, which is degraded within minutes upon binding with auxin; mDII-ntdTOMATO,

which contains a modified, non-degradable, version of DII. Both of the proteins are localised

in the nucleus. We followed the published protocol (Liao et al., 2015) to separately collect the

emission from mDII-ntdTOMATO (Extended Data Fig. 6a) and DII-n3xVENUS (Extended Data

Fig. 6b).

For each root, a mean background was defined as the average pixel intensity in the mDII

channel of a 80x80 pixels corner of the field of view not occupied by the root. The mean

background was then subtracted form all pixel intensities in both channels (mDII and DII). We

manually segmented with FIJI (Schindelin et al., 2012) the most visible cell nuclei in the

epidermis of each root, both in the distal (12 roots in mock conditions and 16 roots exposed

to the field) and in the proximal (11 roots in mock conditions and 13 roots exposed to the field)

regions of the root tip, and quantified the average pixel intensity (corrected after background

subtraction) for each segmented nucleus. Finally, we calculated the natural log of the ratio

between the average mDII (non-degradable, auxin-independent) and DII (degradable, auxin-

dependent) signals in each nucleus and mapped it on top of the root image (Extended Data

Fig. 6d).

For each root, we calculated the average and standard deviation of these ratios, for

epidermal nuclei facing the anode or the cathode (Fig. 4b).

Sample size as following: exposed to E field and imaged in distal region, N=16; exposed

to E field and imaged in the proximal region, N=13; not exposed to E field and imaged in the

distal region, N=24; not exposed to E field and imaged in the proximal region, N=22.

NPA treatment

To find the minimum concentration of N-1-naphthylphthalamic acid (NPA) that inhibits

gravitropism, seeds were germinated on 1X MS agar medium (0.44% MS Basal medium, 0.5%

sucrose, 0.05% MES hydrate, pH adjusted to 5.7 with TRIS HCl, 0.8% agar) and at 5-8 days

post-germination were transferred for 3 hrs into cell culture dishes containing 5 ml of 1/500X

MS liquid medium (0.00088% MS Basal medium, 0.5% sucrose, 0.05% MES hydrate, pH

adjusted to 5.7 with TRIS HCl) plus NPA at 0.1 µM, 1.0 µM and 10 µM.

To quantify the effect of NPA on electrotropism, after being treated with a concentration of

10 µM NPA for 3 hours as described above, the seedlings were transferred inside a V-box

and exposed to a 1.5 V/cm electric field as described in the Electrotropism Assay section.

Both the V-box and the reservoir bottle contained 10 µM NPA throughout the experiment.

Sample sizes N and number of replicates R were the following: untreated, N=21, R=5; 10

µM NPA treated, N=21, R=5.

Cytokinin treatment

Both the 1X MS agar medium (0.44% MS Basal medium, 0.5% sucrose, 0.05% MES

hydrate, pH adjusted to 5.7 with TRIS HCl, 0.8% agar) contained in the PCR tubes, and the

1/500X MS liquid medium (0.00088% MS Basal medium, 0.5% sucrose, 0.05% MES hydrate,

pH adjusted to 5.7 with TRIS HCl) in the nurseries, were supplemented with 10nM *trans*-zeatin

(Sigma-Aldrich Z0876) as previously suggested (Miyawaki et al., 2006). Still in their PCR

tubes, seedlings were transferred into V-boxes, filled with 1/500X MS liquid medium supplied

with 10nM *trans*-zeatin to conduct the electrotropism experiments.

Sample sizes N and number of replicates R were the following: WT + tZ, N=14, R=3;

cyp735a1 +tZ at 1.5 V/cm, N=13, R=3; *cyp735a1* +tZ at 0 V/cm, N=14, R=3.

Gravitropism assay

In the high-temperature experiment, after the treatment the seedlings were transferred

onto 1X MS square agar plates (0.44% MS Basal medium, 0.5% sucrose, 0.05% MES hydrate,

pH adjusted to 5.7 with TRIS HCl, 0.8% agar).

In the NPA experiment, after the treatment the seedlings were transferred onto 1/500X MS

square agar plates (0.00088% MS Basal medium, 0.5% sucrose, 0.05% MES hydrate, pH

adjusted to 5.7 with TRIS HCl, 0.8% agar) containing the desired concentration of NPA.

In both cases, the plates were moved to a growth room (22 °C, 16 h/8 h light/dark

photoperiod), rotated by 90 degrees to position the roots in a horizontal orientation, and

monitored for root gravitropic response.

Statistical analysis

When comparing two samples of measurements, each distribution was first checked for

normality with the Shapiro-Wilk test with alpha-level = 0.05. Normal distributions were tested

with the two-tails Student's t-test without assuming equal variances (Welch two sample t-test);

if one of the two distributions was not normal, the non-parametric Mann-Whitney U (Wilcoxon)

test was used. Unless stated otherwise, all comparisons were performed assuming

independence (unpaired test). All statistical tests were performed with R.

ACKNOWLEDGEMENTS

We thank T. Vernoux for providing *R2D2* seeds; O. Leyser for providing *atipt* triple mutant

seeds; A. Kobayashi and Tohoku University for providing *miz1-1* seeds; A. Guyon, O. Kranse,

C. Keyzor, C.D. Livesey-Clare for generous technical help; M. Levin and I. Bordeu for critical

discussions at various stages of the project. This work was partly supported by the BBSRC

Impact Acceleration Account BB/S506667/1 and the Allen Discovery Center at Tufts

University; N.K. was supported by the BBSRC DTP grant BB/M011178/1.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

Adler M, Mayo A, Alon U (2014) Logarithmic and Power Law Input-Output Relations in

Sensory Systems with Fold-Change Detection. Plos Comput Biol **10**: e1003781

Baluška F, Mancuso S, Volkmann D, Barlow PW (2010) Root apex transition zone: a

signalling–response nexus in the root. Trends in Plant Science **15**: 402–408

Beemster GTS, Baskin TI (2000) STUNTED PLANT 1 Mediates Effects of Cytokinin, But Not

of Auxin, on Cell Division and Expansion in the Root of Arabidopsis. Plant Physiol **124**: 1718–

1727

Chabert N, Ali OA, Achouak W (2015) All ecosystems potentially host electrogenic bacteria.

Bioelectrochemistry **106**: 88–96

Chang J, Li X, Fu W, Wang J, Yong Y, Shi H, Ding Z, Kui H, Gou X, He K, et al (2019)

Asymmetric distribution of cytokinins determines root hydrotropism in Arabidopsis thaliana.

Cell Res **29**: 984–993

Chen R, Hilson P, Sedbrook J, Rosen E, Caspar T, Masson PH (1998) The Arabidopsis

thaliana AGRAVITROPIC 1 gene encodes a component of the polar-auxin-transport efflux

carrier. Proc National Acad Sci **95**: 15112–15117

Chevalier M, Gómez-Schiavon M, Ng AH, El-Samad H (2019) Design and Analysis of a
Proportional-Integral-Derivative Controller with Biological Molecules. Cell Syst **9**: 338-
353.e10

Clarke D, Whitney H, Sutton G, Robert D (2013) Detection and learning of floral electric
fields by bumblebees. Science **340**: 66–69

Crampton WGR (2019) Electroreception, electrogenesis and electric signal evolution. J Fish
Biol **95**: 92–134

Dietrich D, Pang L, Kobayashi A, Fozard JA, Boudolf V, Bhosale R, Antoni R, Nguyen
T, Hiratsuka S, Fujii N, et al (2017) Root hydrotropism is controlled via a cortex-specific
growth mechanism. Nature Plants **3**: 17057

Elfving F (1882) Ueber eine Wirkung des galvanischen Stromes auf wachsende Wurzeln. Bot
Zeit 257–264

Ewart AJ, Bayliss JS (1905) On the nature of the galvanotropic irritability of roots. Proc Royal
Soc Lond Ser B Contain Pap Biological Character **77**: 63–66

Geldner N, Dénervaud-Tendon V, Hyman DL, Mayer U, Stierhof Y-D, Chory J (2009)
Rapid, combinatorial analysis of membrane compartments in intact plants with a multicolor
marker set. The Plant journal : for cell and molecular biology **59**: 169–178

Gilroy S (2008) Plant tropisms. Current Biology **18**: R275-7

Ioio RD, Linhares FS, Scacchi E, Casamitjana-Martinez E, Heidstra R, Costantino P,
Sabatini S (2007) Cytokinins determine Arabidopsis root-meristem size by controlling cell
differentiation. Current Biology **17**: 678–682

Ishikawa H, Evans ML (1990) Electrotropism of maize roots. Role of the root cap and
relationship to gravitropism. Plant physiology **94**: 913–918

Kamada-Nobusada T, Sakakibara H (2009) Molecular basis for cytokinin biosynthesis.
Phytochemistry **70**: 444–449

Kimura T, Haga K, Shimizu-Mitao Y, Takebayashi Y, Kasahara H, Hayashi K-I, Kakimoto
T, Sakai T (2018) Asymmetric Auxin Distribution is Not Required to Establish Root
Phototropism in Arabidopsis. Plant Cell Physiology **59**: 828–840

Kobayashi A, Takahashi A, Kakimoto Y, Miyazawa Y, Fujii N, Higashitani A, Takahashi
H (2007) A gene essential for hydrotropism in roots. Proc Natl Acad Sci U S A **104**: 4724–
4729

Kral N, Ougolnikova AH, Sena G (2016) Externally imposed electric field enhances plant
root tip regeneration. Regeneration **3**: 156–167

Kutschera U, Briggs WR (2012) Root phototropism: from dogma to the mechanism of blue
light perception. Planta **235**: 443–452

Li S-M, Zheng H-X, Zhang X-S, Sui N (2021) Cytokinins as central regulators during plant
growth and stress response. Plant Cell Rep **40**: 271–282

Liao C-Y, Smet W, Brunoud G, Yoshida S, Vernoux T, Weijers D (2015) Reporters for
sensitive and quantitative measurement of auxin response. Nature methods **12**: 207–210

Marchant A, Kargul J, May ST, Muller P, Delbarre A, Perrot-Rechenmann C, Bennett MJ
(1999) AUX1 regulates root gravitropism in Arabidopsis by facilitating auxin uptake within
root apical tissues. Embo J **18**: 2066–2073

Marcum H, Moore R (1990) Influence of Electrical Fields and Asymmetric Application of
Mucilage on Curvature of Primary Roots of Zea mays. American Journal of Botany **77**: 446

McGillivray AM, Gow NAR (1986) Applied Electrical Fields Polarize the Growth of Mycelial
Fungi. Microbiology **132**: 2515–2525

Miyawaki K, Tarkowski P, Matsumoto-Kitano M, Kato T, Sato S, Tarkowska D, Tabata S,
Sandberg G, Kakimoto T (2006) Roles of Arabidopsis ATP/ADP isopentenyltransferases
and tRNA isopentenyltransferases in cytokinin biosynthesis. Proc National Acad Sci **103**:
16598–16603

Miyazawa Y, Takahashi H (2019) Molecular mechanisms mediating root hydrotropism: what
we have observed since the rediscovery of hydrotropism. J Plant Res **133**: 3–14

Muthert LWF, Izzo LG, Zanten M van, Aronne G (2020) Root Tropisms: Investigations on
Earth and in Space to Unravel Plant Growth Direction. Front Plant Sci **10**: 1807

Navez AE (1927) Galvanotropism of roots. The Journal of general physiology **10**: 551–558

Olitzki L (1932) Electric Charge of Bacterial Antigens. J Immunol **4**: 251–256

Peters RC, Eeuwes LBM, Bretschneider F (2007) On the electrodetection threshold of
aquatic vertebrates with ampullary or mucous gland electroreceptor organs. Biol Rev **82**:
361–373

Pozdnyakov A, Pozdnyakova L (2002) Electrical fields and soil properties. Proceedings of
17th World Congress of Soil Science. p N. 1558

Revil A, Naudet V, Nouzaret J, Pessel M (2003) Principles of electrography applied to self-
potential electrokinetic sources and hydrogeological applications. Water Resources
Research **39**: SBH 3: 1-15

Sabatini S, Beis D, Wolkenfelt H, Murfett J, Guilfoyle T, Malamy J, Benfey P, Leyser O,
Bechtold N, Weisbeek P, et al (1999) An Auxin-Dependent Distal Organizer of Pattern and
Polarity in the Arabidopsis Root. Cell **99**: 463–472

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, et al (2012) Fiji: an open-source platform for biological-
image analysis. Nature methods **9**: 676–682

Shanley LJ, Walczysko P, Bain M, MacEwan DJ, Zhao M (2006) Influx of extracellular Ca2+
is necessary for electrotaxis in Dictyostelium. J Cell Sci **119**: 4741–4748

Shkolnik D, Krieger G, Nuriel R, Fromm H (2016) Hydrotropism: Root Bending Does Not
Require Auxin Redistribution. Mol Plant **9**: 757–759

Stevens SS (1970) Neural Events and the Psychophysical Law: Power functions like those
that govern subjective magnitude show themselves in neurelectric effects. Science **170**:
1043–1050
Su S-H, Gibbs NM, Jancewicz AL, Masson PH (2017) Molecular Mechanisms of Root
Gravitropism. Current Biology **27**: R964–R972
Sukul NC, Croll NA (1978) Influence of potential difference and current on the electrotaxis of
Caenorhaditis elegans. J Nematol **10**: 314–7
Sutton GP, Clarke D, Morley EL, Robert D (2016) Mechanosensory hairs in bumblebees
(Bombus terrestris) detect weak electric fields. Proceedings of the National Academy of
Sciences **113**: 7261–7265
Takamura T (2006) Electrochemical Potential around the Plant Root in Relation to Metabolism
and Growth Acceleration. *In* AG Volkov, ed, Plant Electrophysiology, Theory and Methods.
Springer, Berlin, pp 341–374
Takei K, Yamaya T, Sakakibara H (2004) Arabidopsis CYP735A1 and CYP735A2 Encode
Cytokinin Hydroxylases That Catalyze the Biosynthesis of trans-Zeatin. J Biol Chem **279**:
41866–41872
Thimann KV, Went FW (1937) Phytohormones. doi: 10.5962/bhl.title.5695
Ward SH (1990) Geotechnical and environmental geophysics. Soc. of Exploration
Geophysicists
Wawrecki W, Zagórska-Marek B (2007) Influence of a Weak DC Electric Field on Root
Meristem Architecture. Ann Bot-london **100**: 791–796
Wolverton C, Mullen JL, Ishikawam H, Evans ML (2000) Two distinct regions of response
drive differential growth in Vigna root electrotropism. Plant Cell Environ **23**: 1275–1280
Wu Y, Liu H, Wang Q, Zhang G (2021) Roles of cytokinins in root growth and abiotic stress
response of Arabidopsis thaliana. Plant Growth Regul 1–10
677

Figure 1 (a) Schematic of the 3D-printed module used in the V-box. **(b)** Representative time-lapse series of a single root as imaged by the Raspberry camera from the V-box; inset, schematic of the angle measured. **(c)** Polar plot of the average root tip orientations relative to the gravity vector, with time on the radial axis and orientation angle on the circumferential axis; error bars, s.e.m.

Figure 2. **(a)** Root electrotropism response curves: average root tip orientations after 5 h of exposure to a given electric field (left panel) or current (right panel); error bars, s.e.m.; R^2, coefficient of determination. **(b)** Average maximum angular velocity reached by the root tip after 5 h of exposure to a given electric field (left) or current (right); error bars, s.e.m.; R^2, coefficient of determination.

***Figure 3.* (a)** Points of excision overlapped to a representative microscope image of the

Arabidopsis primary root tip; arrow, transition point; scale bar, 100µm. (**b**) Polar plot of the

average root tip orientations with respect to the gravity vector, with time on the radial axis and

orientation angle on the circumferential axis; the same data is presented for the first 2 h (left

panel) and for the full 5 h (right panel); error bars, s.e.m.

Figure 4. **(a)** Relative response to auxin concentration in *Arabidopsis* primary root tips,
indicated by the ratiometric fluorescent reporter R2D2 (see Methods); left panels:
representative tip (distal) regions at 0 V/cm and 1 V/cm; right panels: representative higher
(proximal) regions at 0 V/cm and 1 V/cm; scale bar, 50µm. **(b)** Quantification of the average
relative response to auxin (as shown in a) in the epidermis cell layer facing the positive or
negative electrode, in the distal and proximal regions; the data for 0 V/cm is an average
between the two sides; error bars, s.e.m. **(c-d),** Polar plots of the average root tip orientation
relative to the gravity vector, with time on the radial axis and orientation angle on the
circumferential axis: NPA treatment **(c)**; *aux1* and *pin2* mutants **(d)**; error bars, s.e.m.

Figure 5. Polar plots of the average root tip orientation relative to the gravity vector, with time on the radial axis and orientation angle on the circumferential axis: **(a)** *atipt* triple mutants with strong electrotropic response; **(b)** *atipt* triple mutants with weak electrotropic response; **(c)** *cyp735a1* mutant; error bars, s.e.m.

725 **Figure 6**. Polar plots of the average root tip orientation in *miz1* mutant relative to the gravity

726 vector, with time on the radial axis and orientation angle on the circumferential axis.