Trophic niche of the invasive gregarious species *Crepidula fornicata*, in relation to ontogenic changes

Thibault Androuin [1, 2], Stanislas F. Dubois [2], Cédric Hubas [3], Gwendoline Lefebvre [4], Fabienne Le Grand [4], Gauthier Schaal [4], Antoine Carlier [2].

[1] ISMER, UQAR, Rimouski, Canada
[2] IFREMER, DYNECO-LEBCO, Technopole Brest-Iroise, CS10070, Plouzané, France
[3] UMR BOREA, Muséum National d'Histoire Naturelle, Concarneau, France
[4] UMR CNRS 6539, LEMAR-IUEM-UBO, Plouzané, France

Abstract

The slipper limpet *Crepidula fornicata* is a common and widespread invasive gregarious species along the European coast. Among its life-history traits, well documented ontogenic changes in behavior (i.e., motile male to sessile female) suggests a potential shift in feeding strategy across its life stages. Considering the ecological significance of this species in colonized areas, understanding how conspecifics share the trophic resource is crucial. Using fatty acids (FA) and stable isotopes (SI) as complementary trophic markers, we conducted a field survey between late winter and spring to investigate the trophic niche of three ontogenic stages of *C. fornicata* that bear different sexual (male/female) and motility (motile/sessile) traits. Potential trophic sources were characterized by their pigment, FA and SI compositions and discriminated well over the study period. We showed that the biofilm covering *C. fornicata* shells harbored a higher biomass of primary producers (i.e., chlorophytes and diatoms) than the surrounding sediment surface. Over the study period, we observed a covariation between the three ontogenic stages for both FA and SI compositions which suggest that the trophic niche of *C. fornicata* does not change significantly across its benthic life. During periods of low food availability, slipper limpets displayed an opportunistic suspension-feeding behaviour, relying on both fresh and detrital organic matter, likely coming from superficial sedimentary organic matter. During high food availability (i.e., spring phytoplankton bloom), all ontogenic stages largely benefited from this fresh supply of organic matter (pelagic diatoms in this case). However, the three ontogenic stages showed consistent differences in FA composition, and to a lesser extent in SI composition. These differences persist over time, as they originate more likely from ontogenic physiological changes (differential growth rates, metabolic rate or gametogenesis) rather than diet discrepancies.

Key words: *Crepidula fornicata*, trophic niche, ontogenic shift, fatty acids, stable isotopes, pigments, Bay of Brest

Corresponding authors: thibault.androuin01@gmail.com

1. Introduction

The slipper limpet *Crepidula fornicata* is a non-indigenous and invasive gastropod originating from the East coast of the US (Hoagland, 1985). This species lives in the infralittoral zone but can be found from the intertidal zone down to 60 m depth. It has extensively colonized shallow soft bottom habitats of European coasts, from Norway to the Mediterranean Sea (Blanchard, 1997). Because of its introduction in many parts of the world and its potential cascading effect on food web functioning (Arbach Leloup et al., 2008; Chauvaud et al., 2000; Cugier et al., 2010), several studies have closely investigated its diet and inferred potential trophic overlap with co-occurring benthic species (Blanchard et al., 2008; Decottignies et al., 2007;Decottignies et al., 2007; Lefebvre et al., 2009; Riera, 2007; Riera et al., 2002). *C. fornicata* was overall considered as an opportunistic suspension-feeder, able to feed on a large array of trophic sources (e.g., phytoplankton, microphytobenthos, macroalgae, bacteria), depending on their availability. Based on stable isotope ratios, it has been hypothesised that there are a potentially large contribution of microphytobenthos, and more specifically benthic diatoms, in the diet of *C. fornicata* (Decottignies et al., 2007; Guérin, 2004; Lefebvre et al., 2009; Riera, 2007). However, the unexpected presence of inorganic carbonates in *C. fornicata* soft tissues have led to overestimate $\delta^{13}C$ ratios in this consumer and then to overestimate the trophic role of microphytobenthos (Androuin et al., 2019).

C. fornicata is a hermaphroditic gregarious species, which begins its benthic life as a low-motile male and then becomes a sessile female. Sessile adults form stacks of several non-moving individuals, while juveniles and small males (~10 mm) are motile (Coe, 1936). Adult females are suspension-feeders, but contrary to bivalves, they lack labial palp and show no anatomical or functional potential for qualitative selection (Beninger et al., 2007). They form a food cord in a groove at the distal end of their gill filaments and ultimately catch their food with this cord with their radula before ingesting it (Shumway et al., 2014). For the related species *Crepidula fecunda*, which exhibits ontogenic behavior changes comparable to those of *C. fornicata*, it has been demonstrated that newly settled individuals first adopt a grazing feeding mechanism and gradually shift to a suspension-feeding behaviour once their gill is fully developed (Montiel et al., 2005). Young individuals are able to use both feeding mechanisms (i.e., grazing and suspension-feeding) during the motile phase of their life cycle (size < 28 mm), whereas females are exclusive suspension-feeders (Chaparro et al., 2002; Navarro and Chaparro, 2002). Such a shifting feeding mode has also been suggested for *C. fornicata* albeit without further behavioral evidence nor quantitative measurements. Using stomach content analysis, isolated adults of *C. fornicata* have been found to graze their surrounding habitat (Breton and Huriez, 2010). Younger individuals were also observed near grazing tracks but not analyzed. By comparing the radula to body length ratio, Yee and Padilla (2015) found a more important role of the radula in small individuals (< 12.5 mm) and suggest a potential grazing behavior for those individuals. So, the relative contributions of the radula to feeding in different life stages of *C. fornicata* remain unclear (Shumway et al., 2014).

Since *C. fornicata* often occurs in large densities (up to 2000 ind. m⁻²) on the seafloor with all ontogenic stages grouped in stacks (Guérin, 2004; Martin et al., 2006), one can expect strong intraspecific interactions for food. These interactions could be either facilitative or competitive depending on ontogenic feeding ecology. While purely suspension-feeding slipper limpets should compete for food among ontogenic stages, recent works suggested that younger individuals may be facilitated by adults, both via a higher substrate availability (de Montaudouin and Accolla, 2018) and through the grazing of microphytobenthic biofilm colonizing adult shells (Androuin et al., 2018). In winter, when food is less available in the bay of Brest for suspension-feeders (Lavaud et al., 2018; Chatterjee et al., 2013), grazing may help young slipper limpets to avoid intraspecific competition for food. Moreover, *C. fornicata* often proliferates on muddy and turbid habitats with high suspended inorganic load, thus grazing behavior could also prevent the overloading of their digestive tract with inert matter of low nutritional quality (Navarro and Chaparro, 2002). Investigating the relationship (facilitation *vs.* competition) between age classes in a fecund invasive species like *C. fornicata* is crucial to better understand its population dynamic and so the consequences for the surrounding habitat.

Different trophic markers have long been used to investigate the trophic niche of marine benthic invertebrates (e.g., Blanchet-Aurigny et al., 2015; Cresson et al., 2016; Dubois and Colombo, 2014) and to describe the origin of assimilated particulate organic matter (hereafter OM) (Ke et al., 2017; Lavaud et al., 2018; Liénart et al., 2017). Carbon and nitrogen stable isotopes (SI) are broadly used to infer trophic niche of consumers (Fry and Sherr, 1984; Layman et al., 2012). Classically, nitrogen isotope ratio informs about the trophic position of a species and carbon isotope ratio reflects the origin of assimilated food sources (e.g.,

continental *vs.* oceanic, benthic *vs.* planktonic). In coastal ecosystems, the diet of most of benthic primary
consumers is composed of a mixture of OM from various origins (phytoplankton, macroalgae, continental
detritus, zooplankton, etc) which are often difficult to disentangle with isotopes of only two elements, namely
carbon and nitrogen. This diversity of food sources implies that complementary trophic markers are relevant
to complement SI informations (Majdi et al., 2018). For instance, pigment analyses have been widely used
to study community composition of microscopic primary producers in the water column or in the sediment,
since some pigments are specific to clades of algae (Brotas and Plante-Cuny, 2003; Roy et al., 2011). To a
lesser extent, fatty acid (FA) compositions can be also specific of group of organisms, such as diatoms,
bacteria, copepods or vascular plants (Dalsgaard et al., 2003; Kelly and Scheibling, 2012). FA represents the
building stock of most lipid forms. They are energy reserves (neutral lipids) but also key structural component
of cell membranes (polar lipids). In molluscs as for *C. fornicata*, such lipids storages are essential for larval
development and during mature ontogenic stages for gonadal development (Desloup Paoli et Heral, 1986;
Leroy et al., 2013). Contrary to polar lipids, FA incorporated in the neutral fraction are directly mobilized
and reflect more closely the FA composition of the diet (Langdon and Waldock, 1981; Jezyk and Penicnak,
1966; Fernandez-Reiriz et al. 2015; Waldock and Nascimento, 1979). Extracting FA from this specific class
of lipids and from a tissue with a rapid turnover (e.g., digestive gland, gonad) allows assessing rapid changes
in the diet (McCutchan et al., 2003). Recently, the combined use of SI, FA and pigments improved our
understanding of trophic pathways from the sources of particulate OM to benthic primary consumers (Lavaud
et al., 2018; Majdi et al., 2018).
In this study, we investigated the extent of the trophic niche of *C. fornicata* and hypothesized that
intra-specific differences exist in diet, associated with ontogenic behavior changes (i.e., motile male to sessile
female). We expected that ontogenic trophic shift happens within stacks of *C. fornicata*, with a higher
contribution of biofilm to motile males than to sessile males and females. To test our hypothesis, we
conducted a field survey and characterized potential OM sources by their SI, FA and pigments compositions
and inferred their assimilation in *C. fornicata* tissues using both SI and FA trophic markers.
120
121 **2. Materials and methods**
122
123 **2.1. Sampling strategy**
124
The Bay of Brest (Brittany, France) is a 180 km² semi-enclosed marine ecosystem. The sampling
site is located near the Elorn estuary (48°23'N, 4°23', average depth: 10 m) in a dense *C. fornicata* beds
(~2000 ind. m^{-2}) (Guérin, 2004) dominated by gravelly mud sediment (Gregoire et al., 2016). Potential OM
sources and *C. fornicata* individuals were collected by SCUBA divers at five sampling dates (S1 = 26th
February, S2 = 21th March, S3 = 28th March, S4 = 12th April and S5 = 14th June) around mid and flood tide
to ensure homogeneous mixing between estuarine and oceanic water. The late winter - spring period was
chosen to encompass a period with potentially contrasting OM sources availability (e.g., spring blooms)
(Figure 1).
133

134
Figure 1: Sampling dates (S1 to S5) at the study site superimposed with weekly chlorophyll *a*
concentration at the entrance of the bay of Brest in 2018 (data from the French Coastal Monitoring
Network SOMLIT; http://somlit.epoc.u-bordeaux1.fr/fr/).

Suspended particulate organic matter (SPOM) was sampled at 50 cm above the sediment-water
interface using two 8-litersNiskin bottles, and immediately filtered on board onto a 200 µm nylon mesh to
remove large zooplankton and particles. In the laboratory, between 1 and 1.5 L was filtered on pre-combusted
(450°C for 5 hours) GF/F filters (0.7 µm). Three replicates for each of the three analyses (SI, FA and pigment)
were obtained. Superficial sedimentary organic matter (SSOM) was sampled from three cores of 15 cm
diameter and 15 cm depth. In the laboratory, the sediment-water interface was re-suspended by flushing
seawater with a 30 ml syringe following a standardized process: 60 ml of SSOM was pre-filtered on a 200
µm nylon mesh to be consistent with SPOM samples and filtered on pre-combusted (450°C during 5 hours)
GF/F filters (0.7 µm). Three replicates for each of the three analyses were obtained. Biofilm from one *C.*
fornicata stack was scraped off using a toothbrush and suspended in 600 ml of filtered seawater (0.7 µm).
200 ml of the suspended solution was filtered on pre-combusted (450°C for 5 hours) on GF/F filters (0.7
µm). Three replicates for each of the three analyses were obtained. Filters for FA analysis were put in glass
tubes containing 6 ml of chloroform-methanol (2:1, v:v) solution and stored at -80°C before analysis, whereas
filters for pigment and SI analysis were immediately stored at -80°C.
Females of *C. fornicata* were sampled at the bottom of the stacks (mean shell length 33 ± 6 mm),
attached to a dead *C. fornicata* shell. Sessile and motile males were sampled if they had a penis and a mean
shell length of 20 ± 8 mm and 10 ± 1 mm, respectively. Smallest males were not necessarily found at the
apex of the stack, fitting with the fact that they were fully motile and therefore potentially able to graze the
substrate around them. We used the digestive gland as a relevant trophic integrator tissue because it has a
higher turnover rate than muscle tissue and is an energy storage organ enriched in lipids (McCutchan et al.,
2003; Vander Zanden et al., 2015). However, since digestive gland and gonad are fused in a single organ in
C. fornicata, we analysed both tissues together for sessile males and females. Because gonads are
comparatively small in motile males, the whole body was used to ensure sufficient lipid concentration. At
each date and for each ontogenic stage, both stable isotope (SI) and fatty acid (FA) analyses of *C. fornicata*
were performed on subsamples originating from the same tissue sample. Samples were stored at -80°C for a
few days before freeze drying.
164
165 **2.2. Pigment analysis**
166
The photosynthetic communities of SSOM, biofilm and SPOM have been analyzed by the
quantification of pigments by High Performance Liquid Chromatography (HPLC) according to Brotas and
Plante-Cuny (2003). Filters were crushed and extracted in 3 ml of 95 % cold buffered methanol (2 %
ammonium acetate) for 20 min at -20°C in the dark. Samples were centrifugated for 3 minutes at 3000 g after
the extraction period. Extracts were then filtered with Whatman membrane filters (0.2 mm) immediately
before HPLC analysis. Pigment extracts were analysed using an Agilent 1260 Infinity HPLC composed of a
quaternary pump (VL 400 bar), a UV–VIS photodiode array detector (DAD 1260 VL, 190–950 nm), and a

100 µl sample manual injection loop (overfilled with 250 µl). Chromatographic separation was carried out
using a C18 column for reverse phase chromatography (Supelcosil, 25 cm long, 4.6 mm inner diameter). The
solvents used were A: 0.5 M ammonium acetate in methanol and water (85:15, v:v), B: acetonitrile and water
(90:10, v:v), and C: 100 % ethyl acetate. The solvent gradient followed the Brotas and Plante-Cuny method
(2003), with a flow rate of 0.5 mL min^{-1}. Identification and calibration of the HPLC peaks were performed
with chlorophyll *a*, ββ-carotene, chlorophyll c2, diatoxanthin, diadinoxanthin and fucoxanthin standards. All
peaks detected were identified by their absorption spectra and relative retention times using the Open Lab
CDS software (ChemStation Edition for LC/MS Systems, Agilent Technologies). Quantification was
performed by repeated injections of standards over a range of dilutions to establish a standard curve of
concentrations. Pigment percentages were expressed relatively to the surface/volume sampled (µg.cm^{-2} for
biofilm and SSOM, and µg L^{-1} for SPOM). We measured the mean surface of three stacks of *C. fornicata* to
standardize surfaces.

2.3. Stable isotope analysis

$\delta^{15}N$ and $\delta^{13}C$ analyses were carried out independently for both OM sources and *C. fornicata* tissues.
For OM sources, filters were freeze-dried and split in two equal parts. Half of the filter was scrapped off and
weighed in tin capsules for $\delta^{15}N$ analysis. The second half was decarbonated using acid-flume (10 N
hydrochloric acid solution) for 7 hours (Lorrain et al., 2003), dried at 40 °C for 12 h, scrapped off and weighed
in tin capsules for $\delta^{13}C$ analysis. *C. fornicata* samples were freeze-dried and ground into homogenous powder
using a mortar and pestle. Approximately 400 µg of powder was weighed in tin capsules for $\delta^{15}N$ analysis.
Because both lipids content and inorganic carbonates can influence $\delta^{13}C$ (Androuin et al., 2019; McCutchan
et al., 2003), approximately 400 µg of powder was added to 1 ml of cyclohexane in Eppendorf tubes. Tubes
were vortexed and centrifuged at 3000 g during 5 min. The supernatant was discarded, and the tubes dried at
40°C for a 12 h period. If the supernatant remained coloured, the sample was re-processed. Lipid-free tissues
were then weighed in silver capsules and in-cup decarbonated using 1N HCl. Each capsule was visually
checked, dried at 40°C for 1 h, and sealed. Samples were analysed for $\delta^{15}N$ and $\delta^{13}C$ by continuous flow on
a Thermo Scientific Flash EA 2000 elemental analyser coupled to a Delta V Plus mass spectrometer at the
Pôle de Spectrométrie Océan (PSO, Plouzané, France). Results are expressed in standard δ notation based on
international standards (Vienna Pee Dee Belemnite for $\delta^{13}C$ and atmospheric nitrogen for $\delta^{15}N$) following
the equation:

$$\delta = [(R_{sample}/R_{standard}) - 1] \times 10^3 \text{ (in ‰)}, \text{ where R is } ^{13}C/^{12}C \text{ or } ^{15}N/^{14}N.$$

International isotopic standards of known $\delta^{15}N$ and $\delta^{13}C$ values were used: IAEA-600 Caffeine, IAEA-CH-
6 Sucrose, IAEA-N-1 and IAEA-N-2 Ammonium Sulphate. The analytical precision was estimated using the
standard deviation of an internal standard (Thermo Acetanilide, n = 8), as ± 0.11 ‰ and ± 0.07 ‰ for $\delta^{13}C$
and $\delta^{15}N$ values, respectively.

2.4. Fatty acids analysis

Immediately after freeze-drying, tissues of *C. fornicata* were powdered and subsampled for FA
analyses: between 2 and 20 mg - depending on ontogenic stages - were put in glass tubes (previously heated
for 6 h at 450°C) containing 6 mL of a chloroform/methanol mixture (2:1, v:v), and extracted with a Dounce
homogenizer. OM sources and *C. fornicata* samples were sonicated during 10 min and kept at -20°C until
further analysis. The total lipid fractions were analyzed in OM sources, whereas only the neutral lipids were
analyzed in *C. fornicata* samples. The detailed analysis method for separation and methylation is detailed in
Le Grand et al. (2014). Fatty acid methyl esters (FAME) were analyzed in a Varian CP 8400 gas
chromatograph (GC) equipped with a split/splitless injector and a flame-ionization detector (FID). FAMEs
were identified using two different capillary columns (ZBWAX 30 m × 0.25 mm i.d., 0.25 µm thickness,
Phenomenex®; and ZB-5HT 30 m × 0.25 mm i.d., 0.25 µm thickness, Phenomenex®) by means of a standard
FAME mix (S37, PUFA1, PUFA3, BAME, Sigma Aldrich®) and other lab-made standard mixtures with
previously characterized and published FA composition (e.g., non-methylene interrupted FA, Kraffe et al.,
2004; Le Grand et al., 2013). FAs were expressed as the molar percentage of the total FA content.

2.5. Statistical analyses

230
Pigment and FA compositions of OM sources, and FA compositions of ontogenic stages of *C.*
fornicata were represented using a non-metric multidimensional Scaling (n-MDS). Homogeneity of the data
was tested using permutational analyses of multivariate dispersion (PERMDISP) (Anderson, 2001).
Statistical analyses on OM sources (pigments and FA) and *C. fornicata* (FA) were conducted using a non-
parametric distanced-based permutation multivariate analysis of variance (PERMANOVA) based on a Bray-
Curtis distance. Analyses were performed using two variables: OM sources or ontogenic stages (3 levels'
factors) and sampling dates (5 levels' factor). Each date was considered independent due to the relative high
turnover rate of both microorganisms found in the OM sources as well as the cells in the sampled tissues in
C. fornicata. Following significant PERMANOVA results, *post hoc* tests were carried out using multiple
pairwise comparisons with Bonferroni correction to identify differences among factors (Martinez Arbizu,
2017). However, the number of samples at each sampling date ($3 < n < 5$) was not sufficient to allow
significant differences among the two factor levels in interaction, because of lack of statistical power when
using Bonferroni correction in too many multiple comparisons. Therefore, *post hoc* comparisons of
interaction term were not investigated. Finally, a SIMPER analysis was used to identify the FA explaining
most of the dissimilarities between sampling dates and OM sources/ontogenic stages of *C. fornicata*.
Temporal variations and differences in SI ratios and pigment ratios/FA markers between OM
sources/ontogenic stages of *C. fornicata* and sampling dates were assessed using two-way factorial analyses
of variance (ANOVA). Because pigment and FA concentrations in SPOM were not comparable with biofilm
and SSOM (surface *vs.* volume), SPOM was not include in the analysis. Its temporal variation was assessed
using the non-parametric Kruskal-Wallis test. When ANOVA were significant, *post hoc* multiple
comparisons were carried out using Tukey HSD. Normality and homogeneity of residuals were graphically
assessed. When these conditions were not fulfilled, simple factor effect was verified using non-parametric
Kruskal-Wallis and Mann-Whitney test for sampling dates and OM sources, respectively. Statistical analyses
were performed in R version 3.3.0 (R Core Team, 2016) using packages 'vegan', 'plyr', 'FactoMiner', and
'ggplot'.
256
257 **3. Results**
258
259 **3.1. Organic matter sources**
260
261 3.1.1. Pigments and fatty acids compositions
262
Overall, OM sources were well discriminated by their pigment compositions (Figure 2a), and by
their FA compositions (Figure 2b).
Pigment compositions differed significantly between OM sources and sampling dates ($p < 0.001$ in both
cases) and the interaction between the two factors was significant ($p < 0.001$). SIMPER analysis revealed
that 80 % of the variability was explained by 7 pigments (Table S1). The biofilm was characterized by higher
percentages of chlorophyll *b* and neoxanthin, together with one unknown pigment. SSOM was characterized
by pheophytin *a*, pheophorbide *a*, and to a lesser extent lutein, whereas fucoxanthin and alloxanthin mainly
discriminated SPOM. Temporal variations were mainly driven by a constant increase in both pheophorbide
a and pheophytin *a* percentages for all OM sources. Fucoxanthin also showed an increase over time, except
for SPOM and SSOM at S5 (Table S1).
Similarly, FA compositions also significantly differed between OM sources and sampling dates ($p < 0.001$
in both cases), and the two factors showed a significant interaction ($p < 0.001$). According to the SIMPER
analysis, SSOM was characterized by higher percentages of 22:0, 16:1n-7, 18:1n-7 and a lower percentage
of 16:0 (Table S2). The FA 18:0 and 20:4n-6 mostly discriminated biofilm whereas SPOM had higher
percentages of 14:0 and 22:2n-6, and a lower percentage of 20:5n-3. In terms of temporal variations, both
biofilm and SPOM showed similar decrease in saturated FA (i.e., 16:0 and 18:0) and increase in 16:1n-7 and
20:5n-3, especially between S4 and S5 (Table S2). SSOM exhibited less variable FA composition over time.
280

Figure 2: n-MDS based on the total pigment (a) and fatty acid (b) compositions of organic matter sources (Biofilm, suspended particulate organic matter (SPOM), superficial sedimentary organic matter (SSOM)). S1 to S5 correspond to the sampling dates (S1 = 26th February, S2 = 21st March, S3 = 28th March, S4 = 12th April and S5 = 14th June).

Total FA concentration did not show significant temporal variations for any OM sources (Figure 3a), with the biofilm always exhibiting higher concentration of total FA than SSOM over the sampling period ($p < 0.001$). Chlorophyll (chl) a concentration increased over time in PPOM up to sampling date S4 (1.8 ± 0.2 µg L^{-1}) ($p < 0.05$) as well as in biofilm (reaching 3.9 ± 1 mg m^{-2} at S5) even if differences were not significant due to high between-samples variability (Figure 3b). Chl a concentration in SSOM slightly increased over time ($p < 0.05$) and was lower than in biofilm over the season (p < 0.001). Fucoxanthin concentration increased over time for all OM sources ($p < 0.05$ in all cases), followed by a decrease in S5 for SPOM and SSOM (Figure 3c). Fucoxanthin concentration was higher in biofilm than in SSOM ($p < 0.001$). The chl b: chl a ratio was 3 to 7-fold higher for biofilm than for SSOM ($p < 0.001$) and SPOM ($p < 0.001$) (Figure 3d). Fucoxanthin: chl a (Figure 3e) and chl c: chl a (Figure 3f) ratios did not show clear temporal patterns for any OM sources and. However, they were 5 to 50-fold higher in SSOM than in SPOM ($p < 0.001$) and biofilm ($p < 0.001$) over the studied period, respectively.

299
300 **Figure 3: Total fatty acids (a), chlorophyll *a* (b) and fucoxanthin (c) concentrations, and ratios between**
301 **chlorophyll *b* (d), fucoxanthin (e), and chlorophyll *c* (f) over chlorophyll *a* (mean ± SD, n ≥ 2) of organic**
302 **matter sources (Biofilm, suspended particulate organic matter (SPOM), superficial sedimentary**
303 **organic matter (SSOM)). S1 to S5 correspond to the sampling dates (S1 = 26ᵗʰ February, S2 = 21ˢᵗ**
304 **March, S3 = 28ᵗʰ March, S4 = 12ᵗʰ April and S5 = 14ᵗʰ June).**

305
306 3.1.2. Stable isotopes composition
307
The three OM sources were well discriminated by their $\delta^{13}C$ and $\delta^{15}N$ values over the studied period.
Their $\delta^{13}C$ signal varied significantly according to both OM sources and sampling dates, and the interaction
between the two factors was significant ($p < 0.001$, $p < 0.01$ and $p < 0.001$, respectively). PPOM was always
depleted in ^{13}C compared to RPOM ($p < 0.001$ at each date) and biofilm ($p < 0.01$ at each date) (Figure 4a,
Table S3). Significant temporal $\delta^{13}C$ variations were only observed in biofilm, with higher values at S5 than
at S2 ($p < 0.001$) or S4 ($p < 0.001$). Biofilm was significantly enriched in ^{15}N compared to both PPOM ($p <$
0.001) and RPOM ($p < 0.001$) (Figure 4b, Table S3). There was no interaction between OM sources and
sampling dates ($p = 0.26$).

316
Figure 4: δ¹⁵N (a) and δ¹³C (b) isotopic compositions (mean ± SD, n = 3), and overall isotopic biplot (c)
obtained of organic matter sources (Biofilm, suspended particulate organic matter (SPOM), superficial
sedimentary organic matter (SSOM)). S1 to S5 correspond to the sampling dates (S1 = 26ᵗʰ February,
S2 = 21ˢᵗ March, S3 = 28ᵗʰ March, S4 = 12ᵗʰ April and S5 = 14ᵗʰ June).

3.2. *Crepidula fornicata*

3.2.1. Fatty acids composition

FA composition of *C. fornicata* significantly differed between ontogenic stages and sampling dates
($p < 0.001$ in both cases) and the interaction between the two factors was significant ($p < 0.001$). The analysis
of multivariate dispersion was also significant ($p < 0.05$), indicating that multivariate dispersion was not
homogeneous. This was clearly illustrated by the n-MDS (when comparing the convex hull areas) where
motile males showed much higher variation than sessile females and sessile males exhibited an intermediate
level of variation (Figure 5). Pairwise SIMPER analyses between ontogenic stages revealed that sessile males
were mainly characterized by saturated FA 16:0 and 18:0, especially at the two first sampling dates (Table
S4). Sessile females differed from both motile and sessile males by higher percentages of C_{20} FA such as
20:5n-3, 22:6n-3 and 20:1n-11, but also higher percentages of odd branched FA as iso17:0 (Table S4). Sessile
males showed an overall comparable FA composition than sessile females but exhibited higher variability
between sampling dates, as shown by the n-MDS. The SIMPER analyses performed between dates revealed
that FA that most contributed to the observed temporal changes were the FA 16:0, 18:0 and 22:6n-3
decreasing over time, and the FA 20:5n-3 and 16:1n-7 increasing over time, all accounting for approximately
45 % of the dissimilarity over the 5 sampling dates (Table S4).

340
341 **Figure 5: n-MDS based on the total fatty acid compositions of ontogenic stages of *Crepidula fornicata***
342 **(motile males, sessile males, sessile females). S1 to S5 correspond to the sampling dates (S1 = 26th**
343 **February, S2 = 21st March, S3 = 28th March, S4 = 12th April and S5 = 14th June).**
344
All FA percentages and ratios changed significantly between sampling dates ($p < 0.001$ in all cases)
and between ontogenic stages ($p < 0.05$ in all cases, except for 16:1n-7 and 18:4n-3). The interaction terms
were always significant ($p \leq 0.05$). Overall, temporal variations in FA composition were the highest in motile
males, the lowest for females and intermediate for sessile males.
PUFA/SFA ratio increased over time for both motile and sessile males but remained constant in females
(Figure 6a). There was no significant difference between ontogenic stages at sampling date S5. The relative
abundance of branched FA was quite variable but significantly higher in sessile females than in motile males
($p > 0.05$) (Figure 6b). The highest values in branched FA was recorded in sessile females at S1 (11.2 ± 4.3)
and the lowest in sessile males at S5 (3.8 ± 1). The ratio between 20:5n-3 and 22:6n-3 exhibited temporal
variations for all ontogenic stages (Figure 6c), with a strong increase in S5 ($p < 0.001$ in all cases) where
values ranged from 2.5 ± 0.5 in sessile females to 4 ± 0.6 in motile males. The FA 16:1n-7 followed the same
trend as the 20:5n-3 / 22:6n-3 ratio with a more gradual increase over time for all ontogenic stages (Figure
6d). The highest values (from 3.6 ± 0.9 in sessile females to 6.4 ± 1.9 in motile males) were also recorded at
S5. The FA 18:4n-3 also showed strong temporal variations in all ontogenic stages (Figure 6e), with an
increase up to S4 followed by a decrease at S5. Finally, the n-3/n-6 ratio showed no temporal variation for
sessile females but increased significantly over time up to S4 in both sessile ($p < 0.01$ in all cases) and motile
males ($p < 0.05$ in all cases) (Figure 6f).

362
Figure 6: Relative abundance of fatty acids (FA) in the three ontogenic stages of *Crepidula fornicata*
(motile males, sessile males, sessile females) (mean ± SD, n = 5): (a) Polyunsaturated FA / Saturated
FA ratio, (b) Branched FA, (c) 20:5n-3 / 22:6n-3 ratio, (d) 16:1n-7, (e) 18:4n-3 and (f) n-3/n-6 ratio. S1
to S5 correspond to the sampling dates (S1 = 26th February, S2 = 21st March, S3 = 28th March, S4 =
12th April and S5 = 14th June).

3.2.2. Stable isotopes composition

Overall, the three ontogenic stages exhibited similar isotopic patterns over time (Figures 7a and 7b,
respectively). No interactions were found between ontogenic stage and sampling date, both for carbon (p =
0.6) and nitrogen ($p = 0.27$). Motile males were significantly depleted in ^{13}C compared to sessile females (p
< 0.05) and significantly depleted in ^{15}N compared to both sessile females ($p < 0.001$) and males ($p < 0.05$).
Significant temporal variations were observed for δ^{13}C ($p < 0.001$) with a marked ^{13}C enrichment at S5
compared to all other sampling dates ($p < 0.05$ in all cases), up to 2 ‰ when compared to S4. A significant
temporal decrease in δ^{15}N was found ($p < 0.001$) for all ontogenic stages.

Figure 7: δ¹⁵N (a) and δ¹³C (b) isotopic compositions (mean ± SD, n = 5), and corresponding isotopic biplot (c) for three ontogenic stages of *Crepidula fornicata* (motile males, sessile males, sessile females). S1 to S5 correspond to the sampling dates (S1 = 26th February, S2 = 21st March, S3 = 28th March, S4 = 12th April and S5 = 14th June).

4. Discussion

Contrary to our hypothesis, and while the OM sources were well discriminated by their pigments, FA and SI compositions, trophic markers measured in *Crepidula fornicata* suggested an overall similar trophic niche across its ontogenic stages. Our results confirmed that the slipper limpet is an opportunistic suspension-feeder that exploits both pelagic and benthic particulate OM in varying proportions according to the season and sources availability. Although differences in FA composition, and to a lesser extent in SI composition, were noticeable between ontogenic stages at each sampling date, we think they likely reflect ontogenic physiological changes linked to differential growth rate and energetic demand rather than profound changes in diet.

4.1. Composition and availability of potential food sources

δ¹³C in SPOM (-25.4 ± 0.7 ‰) in our study site was likely influenced by both marine (-22.3 ± 1.2 ‰, data from the French Coastal Monitoring Network SOMLIT; http://somlit.epoc.u-bordeaux1.fr/fr/) and terrestrial POM ((-27.5 ± 1.2 ‰, Mortillaro et al., 2014), as mentioned by Marchais et al. (2013). Over our study period, SPOM showed comparable chlorophyll *a* biomass (0.4 - 1.8 µg L⁻¹) than currently encountered in other areas of the bay of Brest (ca. 0.3 - 5 µg L⁻¹, Chatterjee et al. 2013). High percentages of fucoxanthin and alloxanthin suggested the presence of Bacillaryophyta (i.e., diatoms) and Cryptophyta in the water column, respectively (Brotas and Plante-Cuny, 2003; Roy et al., 2011).

Using the SOMLIT weekly monitoring of chlorophyll *a* and marine POM δ¹³C, a typical ¹³C enrichment (~3 ‰) was noticeable during a phytoplankton bloom that occurred between May 25th and June 1st in the bay. This bloom was mainly composed by three diatoms: *Cerataulina pelagica* (1.1 10⁶ cells L⁻¹), *Leptocylindricus danicus* (4.7 10⁴ cells L⁻¹) and *Rhizosolenia imbricata* (1.2 10⁴ cells L⁻¹) (data extracted from the REPHY network, IFREMER). While this bloom was not sampled in the PPOM sampling set, the ¹³C enrichment was reflected in the tissues of *C. fornicata* collected at S5 (14th June).

Among the three sources of organic matter, SSOM showed the most homogeneous isotopic composition over time, which is unexpected as SSOM is often considered as a mixture of pelagic and benthic OM sources and consequently highly variable according to OM sources proportions and isotopic compositions. SSOM appeared here as a complex mixture of low and high quality OM (Lefebvre et al., 2009; Rigolet et al., 2014). On one hand, it was characterized by i) pheophorbide *a* and pheophytin *a*, which are degradation products of chlorophyllide *a* and chlorophyll *a*, respectively (Brotas and Plante-Cuny, 1998; Cartaxana et al., 2003), ii) odd branched FA (such as ant15:0) and the 18:1n-7 indicating the presence of

bacteria (Hubas et al., 2017; Jaschinski et al., 2011; Meziane et al., 1997) and iii) the long chain saturated
fatty acids 22:0 which confirms the presence of a refractory terrestrial contribution in this environment
(Canuel, 2001). On the other hand, SSOM exhibited the highest PUFA/SFA ratio most of the time (Table
S1), suggesting higher quality/lability compared to SPOM and biofilm (Connelly et al., 2015, 2016; Parrish
et al., 2005). This was confirmed by high Fucoxanthin / Chl *a* and Chl *c* / Chl *a* ratios which indicated a
higher contribution of diatoms in SSOM than in SPOM and biofilm (Brotas and Plante-Cuny, 2003).

Biofilm scraped on shells of *C. fornicata* showed higher chlorophyll *a* concentration than the
surrounding sediment, suggesting higher biomass of primary producers on shells (Androuin et al., 2018). The
high percentages of chl *b* and neoxanthin, as well as the Chl *b* / Chl *a* ratio in the biofilm also suggest that
chlorophytes were abundant on shells (Brotas and Plante-Cuny, 2003). While these results were not supported
by FA (e.g., 16:4n-3, 18:3n-3 or 18:4n-3 characterizing chlorophytes, Fleurence et al. 1994; Kelly and
Scheibling, 2012), mollusc shells are known to harbour microchlorophytes or macrochlorophytes propagules
(Barillé et al., 2017; Mineur et al., 2007). FA characterizing the biofilm were 18:0 and 20:4n-6. While 18:0
is an ubiquitous FA in marine environment, the 20:4n-6 can be found in large proportion in red and brown
algae (Fleurence et al., 1994; Kelly and Scheibling, 2012). It is worth noting that *C. fornicata* shells were
partly covered with crustose red algae in our study site (pers. obs.). The high concentration of fucoxanthin
also suggests the presence of high biomass of diatoms on these shells, as already mentioned by Ní Longphuirt
et al. (2007).

435
436 **4.2. Trophic niche of *C. fornicata*, in relation to ontogenic changes**
437
Overall, SI ratios of *C. fornicata* showed that the three ontogenic stages had similar isotopic niches,
although the niche of sessile females does not fully overlap with those of motile males. According to the
respective SI ratios of potential food sources (Biofilm, SPOM, SSOM and marine POM) and those of *C.*
fornicata tissues, and considering classical diet to consumer trophic enrichment factor (~ 0.75 - 1 ‰ for
carbon and ~2.5 - 2.74 ‰ for nitrogen, (Caut et al., 2009; McCutchan et al., 2003), it is likely that *C. fornicata*
relies either on SSOM or marine POM depending of the season and food availability in the water column.
During lower food availability period (i.e., end of winter and early spring), it is difficult to disentangle SSOM
from marine POM since their SI signals do not differ. However marine POM was sampled close to the Bay
entrance with higher oceanic influence, whereas SPOM was sampled just above the *C. fornicata* beds around
mid and flood tide to ensure mixing with oceanic water. Therefore, SPOM was more readily available than
marine POM for the slipper limpet. It is then reasonable to assume that SI composition of *C. fornicata* refers
to SSOM rather than marine POM, since SSOM can be exploited through regular resuspension events linked
to tidal currents (Beudin, 2014). After the phytoplankton spring bloom that occurred in the bay of Brest at
the end of May, producing a ^{13}C enrichment in the water column (~3 ‰), a similar ^{13}C enrichment was found
for all stages of *C. fornicata* (~2 ‰). These results clearly showed that adults as well as young individuals of
C. fornicata benefited from the spring bloom. However, minor but consistent isotopic differences were also
found between ontogenic stages at each sampling date, which cannot necessarily be attributed to an ontogenic
diet shift. Indeed, inferring diet shift using SI ratios may be hampered by the effects of physiological changes
occurring during ontogeny such as gonadal maturation, metabolic rate or differential tissue growth between
youngs and adults (Blanchet-Aurigny et al., 2012; Hentschel, 1998; Rossi et al., 2004; Vander Zanden et al.,
2015). Lefebvre and Dubois (2016) analysed trophic enrichment factor and turnover rate in several marine
benthic invertebrates, including *C. fornicata*. They showed a clear negative relationship between growth and
enrichment factor values. Young motile males of . *fornicata* are growing more rapidly than larger sessile
males or even larger females (Hoagland, 1978; Walne, 1956). Enrichment factors are then expected to be
smaller for motile males than sessile males or females. This is consistent with small between-stage
differences found in SI ratios (Figure 7c). Even if motile males were analysed *in toto* (including muscle tissue
with longer turnover rate), we believe that this does not bias interpretations of between-stages differences
over the studied period. Analysing only tissues with fast turnover rate (as we did for sessile individuals)
would have increased these between-stages differences.

467
The FA profiles of the three ontogenic stages of *C. fornicata* showed a consistent temporal variation
over the study period, suggesting similar changes in food sources utilization. This temporal pattern resulted
from three distinct groups of sampling dates (Figure 5). The two first ones (S1 and S2) likely corresponded
to a period that integrated the trophic signal of winter season's food sources, whereas the last one (S5) clearly
showed the assimilation of the spring phytoplankton bloom. The two-intermediate sampling dates (S3 and

S4) corresponded to a transition reflecting an increase in food availability. As interpreted above with SI, all
ontogenic stages of *C. fornicata* may have probably exploited SSOM before the spring bloom when food in
the water column is less available. In this pool of OM, FA revealed that slipper limpets likely fed on benthic
diatoms (as suggested by 16:1n-7 and 20:5n-3; Dunstan et al. 1992; Napolitano et al. 1997; Passarelli et al.
2012), dinoflagellates (22:6n-3; Zhukova and Aizdacher 1995; Lavaud et al. 2018) and bacteria (Branched
FA and 18:1n-7; Perry et al. 1979; Zhukova et al. 1992; Haack et al. 1994), which is in agreement with
previous interpretations done in other coastal bays (Dubois et al., 2014; Leroy et al., 2013). In sediment,
bacteria are often associated with detritus and are therefore not considered as a high-quality food source
(Dalsgaard et al., 2003). This is confirmed by the PUFA/SFA ratio, a biomarker of fresh *vs.* detritic OM
(Connelly et al., 2015, 2016; Parrish et al., 2005), which was lower in the slipper limpet (~ 1.5, our study)
than in other suspension-feeding species of the bay of Brest, such as *Pecten maximus* (~ 2.8, Lavaud et al.
2018) or *Ophiotrix fragilis* (~ 2, Blanchet-Aurigny et al. 2015). The fact that *C. fornicata* lacks pre-ingestive
mechanisms for particle selection likely explains their opportunistic trophic behaviour based on both fresh
and detritic organic matter (Beninger et al., 2007). After spring bloom, the percentages of diatom's markers
16:1n-7 and 20:5n-3 drastically increased in *C. fornicata*'s tissue, as well as the 20:5n-3/22:6n-3 ratio,
confirming that the three ontogenic stages benefit from this food supply from the water column (Budge and
Parrish, 1998; Lavaud et al., 2018).Finally, the contribution of FA 18:4n-3 increased over time for all
ontogenic stages. According to the literature, this FA may originate from different primary producers such
as dinoflagellates (Budge and Parrish, 1998), brown or green macroalgae (Fleurence et al., 1994; Kelly and
Scheibling, 2012). Considering i) the absence or low temporal variation of other dinoflagellate biomarkers
(peridinin pigment and 22:6n-3 FA) in the OM sources, ii) the presence of brown algae on the rocky shore,
and iii) the frequent seasonal accumulation of green macroalgae in our study area (Study Centre for Algal
Promotion, http://www.ceva.fr; Ragueneau et al. 2018), we can expect a seasonal trophic role of these
macroalgae for *C. fornicata* at our study site, probably in the form of detrital particles. FA dietary biomarkers
in consumers are subject to physiological regulation (e.g., specific retention, *de novo* synthesis) during uptake
and trophic transfer, and can be species-specific (Galloway and Budge, 2020). Although the use of neutral
lipids theoretically limits this shortcoming, dedicated experimental studies on FA metabolism and FA trophic
transfer in marine limpet specifically are still needed (Zhukova 2019; Galloway and Budge, 2020).
Contrary to SI, FA compositions of the three ontogenic stages of *C. fornicata* showed low overlap,
especially between motile and sessile limpets. Considering that lipids and fatty acids profiles may be age and
sex-specific (Correia et al., 2003; Pernet et al., 2012), some of the differences in FA compositions are then
likely to originate from physiological changes between ontogenic stages. During winter period, motile males
were characterized by higher proportions of SFA such as 16:0 (25-34 %) and 18:0 (27-36 %). These FA are
very common in marine organisms and do not necessarily reflect a specific diet (Dalsgaard et al., 2003; Kelly
and Scheibling, 2012). Moreover, despite the dietary interest of short-chain SFA, where energy is more
efficiently released via beta-oxidation than for PUFA (Langdon and Waldock, 1981), temperature may also
influence the process of their utilization (Pernet et al., 2007). For example, it has been experimentally
demonstrated that cold-acclimated oysters (5-7°C) have a clear preference for PUFA (n-3) over SFA (16:0)
as fuel for energy compared to 'temperate' oysters (Chu and Greaves, 1991). The lower utilization of SFA
in cold-acclimated oysters has been attributed to the fact that SFA are not in the liquid phase under cold
temperature, thus making them less accessible for catabolic processes. In *Crepidula fornicata*, young
individuals have proportionally less energy reserves than adults (Guérin, 2004) and are more subjected to
low temperature effects due to a higher surface-to-volume ratio (Diederich et al., 2015). Hence, the lower
utilization of SFA could explain their higher SFA percentages during the winter period in the bay of Brest,
where temperatures fall down to 7°C (Figure S1). Moreover, the weight-specific metabolic rate, which is
higher in smaller organisms (Bayne and Newell, 1983; Bougrier et al., 1995), could be exacerbated in *C.*
fornicata because motile males are more active than adults through their motility (Coe, 1936; Hoagland,
1978; Walne, 1956). Together, this may explain the higher temporal and inter-individual variability in their
FA compositions. It also suggests that young motile individuals of *C. fornicata*, having less energetic storage
in winter while having more energetic needs, are probably in poor energetic condition during this period.
As we measured neutral FA in both digestive gland and gonad simultaneously (because the digestive
gland cannot be isolated from the gonad), the level of lipid storage and FA composition may also depend on
their sexual development stage. Indeed, females allocate more energy than males in the reproduction due to
maternal gametogenesis (Deslous-Paoli and Héral, 1986; Leroy et al., 2013). As an illustration of FA
composition changes, the n-3/n-6 ratio increased over time for both sessile and motile males whereas it

remains unchanged in females. This may be linked to a preferential allocation of n-3 to early embryos, which showed an increase over the reproductive period of the slipper limpet (Leroy et al., 2013).

5. Conclusion

Contrary to our hypothesis, *Crepidula fornicata* did not show ontogenic feeding shift, despite its peculiar way of life, and development from motile male to sessile female. However, we confirmed the opportunistic feeding behaviour of this suspension-feeder in our study site. Although we only found minor between-stage differences in both SI and FA compositions, it is likely that these differences originate from physiology rather than trophic behavior. These results suggest that young *C. fornicata* (i.e., one year old) may compete for food with adults. This intraspecific competition is expected to be higher in winter, when food is more limited and when young *C. fornicata* are probably in poorer condition. In a recent work, de Montaudouin and Accolla (2018) showed a recruitment facilitation through higher substrate availability in high density beds with a potential mitigation by the negative effect of intraspecific competition. Our study supports this idea of ontogenic trophic competition in *C. fornicata* when food is limiting.

6. Acknowledgements

We thank the PSO and LIPIDOCEAN analytical facilities for stable isotope and fatty acid facilities, respectively (Oanez Lebeau, Antoine Bideau and Rudolph Corvaisier). We are grateful to Aline Blanchet-Aurigny for commenting upon preliminary versions of this manuscript. We also thank the LEBCO diving team (Amelia Curd, Xavier Caisey and Aurélien Tancray) for providing biological samples. We finally thank two anonymous reviewers for valuable input for the manuscript. TA was funded by an IFREMER, LabexMER and Region Bretagne PhD grant. This work was funded by the TOTAL foundation for biodiversity.

7. Conflict of interest

The authors of this preprint declare that they have no financial conflict of interest with the content of this article.
Cédric Hubas is one of the PCI recommenders.

8. Supporting informations

Raw data and R-scripts are freely available at: https://www.seanoe.org/data/00646/75809/

9. References

Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. Austral Ecology 26, 32–46. https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x

Androuin, T., Dubois, S.F., Decottignies, P., Pelleter, E., Carlier, A., 2019. The dark side of soft tissues: Unexpected inorganic carbonate in the invasive slipper limpet *Crepidula fornicata* and its implications for stable isotope interpretations. Rapid Commun Mass Spectrom 33, 107–115. https://doi.org/10.1002/rcm.8322

Androuin, T., Polerecky, L., Decottignies, P., Dubois, S.F., Dupuy, C., Hubas, C., Jesus, B., Le Gall, E., Marzloff, M.P., Carlier, A., 2018. Subtidal microphytobenthos: a secret garden stimulated by the engineer species *Crepidula fornicata*. Front. Mar. Sci. 5, 475. https://doi.org/10.3389/fmars.2018.00475

Arbach Leloup, F., Desroy, N., Le Mao, P., Pauly, D., Le Pape, O., 2008. Interactions between a natural food web, shellfish farming and exotic species: The case of the Bay of Mont Saint Michel (France). Estuarine, Coastal and Shelf Science 76, 111–120. https://doi.org/10.1016/j.ecss.2007.06.011

Barillé, L., Le Bris, A., Méléder, V., Launeau, P., Robin, M., Louvrou, I., Ribeiro, L., 2017. Photosynthetic epibionts and endobionts of Pacific oyster shells from oyster reefs in rocky

versus mudflat shores. PLoS ONE 12, e0185187.
https://doi.org/10.1371/journal.pone.0185187

Bayne, B.L., Newell, R.C., 1983. Physiological energetics of marine molluscs, in: The Mollusca.
Elsevier, pp. 407–515. https://doi.org/10.1016/B978-0-12-751404-8.50017-7

Beninger, P., Decottignies, P., Guiheneuf, F., Barillé, L., Rincé, Y., 2007. Comparison of particle
processing by two introduced suspension feeders: selection in *Crepidula fornicata* and
Crassostrea gigas. Mar. Ecol. Prog. Ser. 334, 165–177.
https://doi.org/10.3354/meps334165

Beudin, A., 2014. Dynamique et échanges sédimentaires en rade de Brest impactés par l´invasion
de la crépidule. Université de Bretagne Occidentale.

Blanchard, M., 1997. Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe. Current
state and consequences. Scientia Marina 61, 10.

Blanchard, M., Pechenik, J.A., Giudicelli, E., Connan, J.-P., Robert, R., 2008. Competition for food
in the larvae of two marine molluscs, *Crepidula fornicata* and *Crassostrea gigas*. Aquat.
Living Resour. 21, 197–205. https://doi.org/10.1051/alr:2008025

Blanchet-Aurigny, A., Dubois, S., Quéré, C., Guillou, M., Pernet, F., 2015. Trophic niche of two
co-occurring ophiuroid species in impacted coastal systems, derived from fatty acid and
stable isotope analyses. Mar. Ecol. Prog. Ser. 525, 127–141.
https://doi.org/10.3354/meps11169

Blanchet-Aurigny, A., Guillou, M., Pernet, F., Gaffet, J.-D., Dubois, S.F., 2012. Tissue-diet
discrimination factors of isotopic ratios ($\Delta\delta^{13}$C and $\Delta\delta^{15}$N) in two brittle star species: Effect
of reproductive state, diet and tissue composition. Journal of Experimental Marine Biology
and Ecology 426–427, 68–77. https://doi.org/10.1016/j.jembe.2012.05.013

Bougrier, S., Geairon, P., Deslous-Paoli, J.M., Bacher, C., Jonquières, G., 1995. Allometric
relationships and effects of temperature on clearance and oxygen consumption rates of
Crassostrea gigas (Thunberg). Aquaculture 134, 143–154. https://doi.org/10.1016/0044-
8486(95)00036-2

Breton, G., Huriez, L., 2010. Les crépidules *Crepidula fornicata* (Linnaeus 1758) peuvent
temporairement redevenir brouteuses à l'âge adulte. Bulletin de la Société Linnéenne de
Normandie 120, 98–104.

Brotas, V., Plante-Cuny, M., 1998. Spatial and temporal patterns of microphytobenthic taxa of
estuarine tidal flats in the Tagus Estuary (Portugal) using pigment analysis by HPLC. Mar.
Ecol. Prog. Ser. 171, 43–57. https://doi.org/10.3354/meps171043

Brotas, V., Plante-Cuny, M.-R., 2003. The use of HPLC pigment analysis to study
microphytobenthos communities. Acta Oecologica 24, S109–S115.
https://doi.org/10.1016/S1146-609X(03)00013-4

Budge, S.M., Parrish, C.C., 1998. Lipid biogeochemistry of plankton, settling matter and sediments
in Trinity Bay, Newfoundland. II. Fatty acids. Organic Geochemistry 29, 1547–1559.
https://doi.org/10.1016/S0146-6380(98)00177-6

Canuel, E.A., 2001. Relations between river flow, primary production and fatty acid composition
of particulate organic matter in San Francisco and Chesapeake Bays: a multivariate
approach. Organic Geochemistry 32, 563–583.

Cartaxana, P., Jesus, B., Brotas, V., 2003. Pheophorbide and pheophytin a-like pigments as useful
markers for intertidal microphytobenthos grazing by *Hydrobia ulvae*. Estuarine, Coastal
and Shelf Science 58, 293–297. https://doi.org/10.1016/S0272-7714(03)00081-7

Caut, S., Angulo, E., Courchamp, F., 2009. Variation in discrimination factors (Δ^{15}N and Δ^{13}C):
the effect of diet isotopic values and applications for diet reconstruction. Journal of Applied
Ecology 46, 443–453. https://doi.org/10.1111/j.1365-2664.2009.01620.x

Chaparro, O., Thompson, R., Pereda, S., 2002. Feeding mechanisms in the gastropod *Crepidula*
fecunda. Mar. Ecol. Prog. Ser. 234, 171–181. https://doi.org/10.3354/meps234171

Chatterjee, A., Klein, C., Naegelen, A., Claquin, P., Masson, A., Legoff, M., Amice, E., L'Helguen, S., Chauvaud, L., Leynaert, A., 2013. Comparative dynamics of pelagic and benthic micro-algae in a coastal ecosystem. Estuarine, Coastal and Shelf Science 133, 67–77. https://doi.org/10.1016/j.ecss.2013.08.015

Chauvaud, L., Jean, F., Ragueneau, O., Thouzeau, G., 2000. Long-term variation of the Bay of Brest ecosystem: benthic-pelagic coupling revisited. Mar. Ecol. Prog. Ser. 200, 35–48. https://doi.org/10.3354/meps200035

Chu, F.-L.E., Greaves, J., 1991. Metabolism of palmitic, linoleic, and linolenic acids in adult oysters, *Crassostrea virginica*. Mar. Biol. 110, 229–236. https://doi.org/10.1007/BF01313708

Coe, W.R., 1936. Sexual phases in Crepidula. J. Exp. Zool. 72, 455–477. https://doi.org/10.1002/jez.1400720306

Connelly, T., McClelland, J., Crump, B., Kellogg, C., Dunton, K., 2015. Seasonal changes in quantity and composition of suspended particulate organic matter in lagoons of the Alaskan Beaufort Sea. Mar. Ecol. Prog. Ser. 527, 31–45. https://doi.org/10.3354/meps11207

Connelly, T.L., Businski, T.N., Deibel, D., Parrish, C.C., Trela, P., 2016. Annual cycle and spatial trends in fatty acid composition of suspended particulate organic matter across the Beaufort Sea shelf. Estuarine, Coastal and Shelf Science 181, 170–181. https://doi.org/10.1016/j.ecss.2016.08.016

Correia, A.D., Costa, M.H., Luis, O.J., Livingstone, D.R., 2003. Age-related changes in antioxidant enzyme activities, fatty acid composition and lipid peroxidation in whole body *Gammarus locusta* (Crustacea: Amphipoda). Journal of Experimental Marine Biology and Ecology 289, 83–101. https://doi.org/10.1016/S0022-0981(03)00040-6

Cresson, P., Ruitton, S., Harmelin-Vivien, M., 2016. Feeding strategies of co-occurring suspension feeders in an oligotrophic environment. Food Webs 6, 19–28. https://doi.org/10.1016/j.fooweb.2015.12.002

Cugier, P., Struski, C., Blanchard, M., Mazurié, J., Pouvreau, S., Olivier, F., Trigui, J.R., Thiébaut, E., 2010. Assessing the role of benthic filter feeders on phytoplankton production in a shellfish farming site: Mont Saint Michel Bay, France. Journal of Marine Systems 82, 21–34. https://doi.org/10.1016/j.jmarsys.2010.02.013

Dalsgaard, J., St. John, M., Kattner, G., Müller-Navarra, D., Hagen, W., 2003. Fatty acid trophic markers in the pelagic marine environment. Advances in Marine Biology 46, 225–340. https://doi.org/10.1016/S0065-2881(03)46005-7

de Montaudouin, X., Accolla, C., 2018. Intraspecific trade-offs between facilitation and competition in the invasive mollusc *Crepidula fornicata*. Mar. Ecol. Prog. Ser. 604, 163–171. https://doi.org/10.3354/meps12726

Decottignies, P, Beninger, P., Rincé, Y., Robins, R., Riera, P., 2007. Exploitation of natural food sources by two sympatric, invasive suspension-feeders: *Crassostrea gigas* and *Crepidula fornicata*. Mar. Ecol. Prog. Ser. 334, 179–192. https://doi.org/10.3354/meps334179

Decottignies, Priscilla, Beninger, P.G., Rincé, Y., Riera, P., 2007. Trophic interactions between two introduced suspension-feeders, *Crepidula fornicata* and *Crassostrea gigas*, are influenced by seasonal effects and qualitative selection capacity. Journal of Experimental Marine Biology and Ecology 342, 231–241. https://doi.org/10.1016/j.jembe.2006.10.005

Deslous-Paoli, J., Héral, M., 1986. *Crepidula fornicata* L. (Gastéropode, Calyptraeidae) dans le bassin de Marennes-Oléron : composition et valeur énergétique des individus et des pontes. Oceanologica acta 9, 305–311.

Diederich, C., Bashevkin, S., Chaparro, O., Pechenik, J., 2015. Desiccation tolerance and lifting behavior in *Crepidula fornicata* (Gastropoda). Mar. Ecol. Prog. Ser. 528, 235–243. https://doi.org/10.3354/meps11284

Dubois, S., Blanchet, H., Garcia, A., Massé, M., Galois, R., Grémare, A., Charlier, K., Guillou, G., Richard, P., Savoye, N., 2014. Trophic resource use by macrozoobenthic primary

consumers within a semi-enclosed coastal ecosystem: Stable isotope and fatty acid
assessment. Journal of Sea Research 88, 87–99.
https://doi.org/10.1016/j.seares.2014.01.004

Dubois, S.F., Colombo, F., 2014. How picky can you be? Temporal variations in trophic niches of
co-occurring suspension-feeding species. Food Webs 1, 1–9.
https://doi.org/10.1016/j.fooweb.2014.07.001

Dunstan, G.A., Volkman, J.K., Jeffrey, S.W., Barrett, S.M., 1992. Biochemical composition of
microalgae from the green algal classes Chlorophyceae and Prasinophyceae. 2. Lipid
classes and fatty acids. Journal of Experimental Marine Biology and Ecology 161, 115–
134. https://doi.org/10.1016/0022-0981(92)90193-E

Fleurence, J., Gutbier, G., Mabeau, S., Leray, C., 1994. Fatty acids from 11 marine macroalgae of
the French Brittany coast. J Appl Phycol 6, 527–532. https://doi.org/10.1007/BF02182406

Fry, B., Sherr, E.B., 1984. $\delta^{13}C$ measurements as indicators of carbon flow in marine and freshwater
ecosystems. Contributions in Marine Science 27, 13–47.

Guérin, L., 2004. La crépidule en rade de Brest : un modèle biologique d´espèce introduite
proliférante en réponse aux fluctuations de l´environnement. Université de Bretagne
Occidentale, Brest.

Haack, S.K., Garchow, H., Odelson, D.A., Forney, L.J., Klug, M.J., 1994. Accuracy,
reproducibility, and interpretation of fatty acid methyl ester profiles of kodel bacterial
communities. Applied and Environmental Microbiology 60, 2483–2493.

Hentschel, B.T., 1998. Intraspecific variation in $\delta^{13}C$ indicate ontogenic diet changes in deposit-
feeding polychaetes. Ecology 79, 1357–1370. https://doi.org/10.1890/0012-
9658(1998)079[1357:IVICIO]2.0.CO;2

Hoagland, K.E., 1978. Protandry and the evolution of environmentally mediated sex change: study
of the Mollusca. Malacogia 17, 365–391.

Hubas, C., Boeuf, D., Jesus, B., Thiney, N., Bozec, Y., Jeanthon, C., 2017. A nanoscale study of
carbon and nitrogen fluxes in mats of purple sulfur bacteria: Implications for carbon
cycling at the surface of coastal sediments. Front. Microbiol. 8, 1995.
https://doi.org/10.3389/fmicb.2017.01995

Jaschinski, S., Brepohl, D., Sommer, U., 2011. Seasonal variation in carbon sources of
-mesograzers and small predators in an eelgrass community: stable isotope and fatty acid
analyses. Mar. Ecol. Prog. Ser. 431, 69–82. https://doi.org/10.3354/meps09143

Ke, Z., Tan, Y., Huang, L., Zhao, C., Jiang, X., 2017. Spatial distributions of $\delta^{13}C$, $\delta^{15}N$ and C/N
ratios in suspended particulate organic matter of a bay under serious anthropogenic
influences: Daya Bay, China. Marine Pollution Bulletin 114, 183–191.

Kelly, J.R., Scheibling, R.E., 2012. Fatty acids as dietary tracers in benthic food webs. Marine
Ecology Progress Series 446, 1–22. https://doi.org/10.3354/meps09559

Kraffe, E., Soudant, P., Marty, Y., 2004. Fatty acids of serine, ethanolamine, and choline
plasmalogens in some marine bivalves. Lipids 39, 59–66. https://doi.org/10.1007/s11745-
004-1202-x

Langdon, C.J., Waldock, M.J., 1981. The effect of algal and artificial diets on the growth and fatty
acid composition of *Crassostrea gigas* Spat. J. Mar. Biol. Ass. 61, 431–448.
https://doi.org/10.1017/S0025315400047056

Lavaud, R., Artigaud, S., Le Grand, F., Donval, A., Soudant, P., Flye-Sainte-Marie, J., Strohmeier,
T., Strand, Ø., Leynaert, A., Beker, B., Chatterjee, A., Jean, F., 2018. New insights into the
seasonal feeding ecology of *Pecten maximus* using pigments, fatty acids and sterols
analyses. Mar. Ecol. Prog. Ser. 590, 109–129. https://doi.org/10.3354/meps12476

Layman, C.A., Araujo, M.S., Boucek, R., Hammerschlag-Peyer, C.M., Harrison, E., Jud, Z.R.,
Matich, P., Rosenblatt, A.E., Vaudo, J.J., Yeager, L.A., Post, D.M., Bearhop, S., 2012.
Applying stable isotopes to examine food-web structure: an overview of analytical tools.
Biological Reviews 87, 545–562. https://doi.org/10.1111/j.1469-185X.2011.00208.x

Le Grand, F., Soudant, P., Marty, Y., Le Goïc, N., Kraffe, E., 2013. Altered membrane lipid composition and functional parameters of circulating cells in cockles (*Cerastoderma edule*) affected by disseminated neoplasia. Chemistry and Physics of Lipids 167–168, 9–20. https://doi.org/10.1016/j.chemphyslip.2013.01.004

Le Grand, F., Soudant, P., Siah, A., Tremblay, R., Marty, Y., Kraffe, E., 2014. Disseminated neoplasia in the soft-shell clam *Mya arenaria*: membrane lipid composition and functional parameters of circulating cells. Lipids 49, 807–818. https://doi.org/10.1007/s11745-014-3917-4

Lefebvre, S., Dubois, S.F., 2016. The stony road to understand isotopic enrichment and turnover rates: insight into the metabolic part. Vie et Milieu 66, 306–315.

Lefebvre, S., Marín Leal, J.C., Dubois, S., Orvain, F., Blin, J.-L., Bataillé, M.-P., Ourry, A., Galois, R., 2009. Seasonal dynamics of trophic relationships among co-occurring suspension-feeders in two shellfish culture dominated ecosystems. Estuarine, Coastal and Shelf Science 82, 415–425. https://doi.org/10.1016/j.ecss.2009.02.002

Leroy, F., Meziane, T., Riera, P., Comtet, T., 2013. Seasonal variations in maternal provisioning of *Crepidula fornicata* (Gastropoda): fatty acid composition of females, embryos and larvae. PLoS ONE 8, e75316. https://doi.org/10.1371/journal.pone.0075316

Liénart, C., Savoye, N., Bozec, Y., Breton, E., Conan, P., David, V., Feunteun, E., Grangeré, K., Kerhervé, P., Lebreton, B., Lefebvre, S., L'Helguen, S., Mousseau, L., Raimbault, P., Richard, P., Riera, P., Sauriau, P.-G., Schaal, G., Aubert, F., Aubin, S., Bichon, S., Boinet, C., Bourasseau, L., Bréret, M., Caparros, J., Cariou, T., Charlier, K., Claquin, P., Cornille, V., Corre, A.-M., Costes, L., Crispi, O., Crouvoisier, M., Czamanski, M., Del Amo, Y., Derriennic, H., Dindinaud, F., Durozier, M., Hanquiez, V., Nowaczyk, A., Devesa, J., Ferreira, S., Fornier, M., Garcia, F., Garcia, N., Geslin, S., Grossteffan, E., Gueux, A., Guillaudeau, J., Guillou, G., Joly, O., Lachaussée, N., Lafont, M., Lamoureux, J., Lecuyer, E., Lehodey, J.-P., Lemeille, D., Leroux, C., Macé, E., Maria, E., Pineau, P., Petit, F., Pujo-Pay, M., Rimelin-Maury, P., Sultan, E., 2017. Dynamics of particulate organic matter composition in coastal systems: A spatio-temporal study at multi-systems scale. Progress in Oceanography 156, 221–239. https://doi.org/10.1016/j.pocean.2017.03.001

Lorrain, A., Savoye, N., Chauvaud, L., Paulet, Y.-M., Naulet, N., 2003. Decarbonation and preservation method for the analysis of organic C and N contents and stable isotope ratios of low-carbonated suspended particulate material. Analytica Chimica Acta 491, 125–133. https://doi.org/10.1016/S0003-2670(03)00815-8

Majdi, N., Hette-Tronquart, N., Auclair, E., Bec, A., Chouvelon, T., Cognie, B., Danger, M., Decottignies, P., Dessier, A., Desvilettes, C., Dubois, S., Dupuy, C., Fritsch, C., Gaucherel, C., Hedde, M., Jabot, F., Lefebvre, S., Marzloff, M.P., Pey, B., Peyrard, N., Powolny, T., Sabbadin, R., Thébault, E., Perga, M.-E., 2018. There's no harm in having too much: A comprehensive toolbox of methods in trophic ecology. Food Webs 17, e00100. https://doi.org/10.1016/j.fooweb.2018.e00100

Martin, S., Thouzeau, G., Chauvaud, L., Jean, F., Guérin, L., Clavier, J., 2006. Respiration, calcification, and excretion of the invasive slipper limpet, *Crepidula fornicata* L.: Implications for carbon, carbonate, and nitrogen fluxes in affected areas. Limnol. Oceanogr. 51, 1996–2007. https://doi.org/10.4319/lo.2006.51.5.1996

Martinez Arbizu, P., 2017. pairwiseAdonis: Pairwise multilevel comparison using adonis. R package version 0.3.

McCutchan, J.H.J., Lewis, W.M.J., Kendall, C., McGrath, C.C., 2003. Variation in trophic shift for stable isotope ratios of carbon, nitrogen, and sulfur. Oikos 102, 378–390.

Meziane, T., Bodineau, L., Retiere, C., Thoumelin, G., 1997. The use of lipid markers to define sources of organic matter in sediment and food web of the intertidal salt-marsh-flat ecosystem of Mont-Saint-Michel Bay, France. Journal of Sea Research 38, 47–58. https://doi.org/10.1016/S1385-1101(97)00035-X

Mineur, F., Belsher, T., Johnson, M.P., Maggs, C.A., Verlaque, M., 2007. Experimental assessment of oyster transfers as a vector for macroalgal introductions. Biological Conservation 137, 237–247. https://doi.org/10.1016/j.biocon.2007.02.001

Montiel, Y.A., Chaparro, O.R., Segura, C.J., 2005. Changes in feeding mechanisms during early ontogeny in juveniles of *Crepidula fecunda* (Gastropoda, Calyptraeidae). Marine Biology 147, 1333–1342. https://doi.org/10.1007/s00227-005-0050-1

Napolitano, G.E., Pollero, R.J., Gayoso, A.M., Macdonald, B.A., Thompson, R.J., 1997. Fatty acids as trophic markers of phytoplankton blooms in the Bahía Blanca estuary (Buenos Aires, Argentina) and in Trinity Bay (Newfoundland, Canada). Biochemical Systematics and Ecology 25, 739–755. https://doi.org/10.1016/S0305-1978(97)00053-7

Navarro, J.M., Chaparro, O.R., 2002. Grazing–filtration as feeding mechanisms in motile specimens of *Crepidula fecunda* (Gastropoda: Calyptraeidae). Journal of Experimental Marine Biology and Ecology 270, 111–122. https://doi.org/10.1016/S0022-0981(02)00013-8

Ní Longphuirt, S., Clavier, J., Grall, J., Chauvaud, L., Le Loc'h, F., Le Berre, I., Flye-Sainte-Marie, J., Richard, J., Leynaert, A., 2007. Primary production and spatial distribution of subtidal microphytobenthos in a temperate coastal system, the Bay of Brest, France. Estuarine, Coastal and Shelf Science 74, 367–380. https://doi.org/10.1016/j.ecss.2007.04.025

Parrish, C., Thompson, R., Deibel, D., 2005. Lipid classes and fatty acids in plankton and settling matter during the spring bloom in a cold ocean coastal environment. Mar. Ecol. Prog. Ser. 286, 57–68. https://doi.org/10.3354/meps286057

Passarelli, C., Hubas, C., Nicolas Segui, A., Grange, J., Meziane, T., 2012. Surface adhesion of microphytobenthic biofilms is enhanced under *Hediste diversicolor* (O.F. Müller) trophic pressure. Journal of Experimental Marine Biology and Ecology 438, 52–60. https://doi.org/10.1016/j.jembe.2012.10.005

Pernet, F., Malet, N., Pastoureaud, A., Vaquer, A., Quéré, C., Dubroca, L., 2012. Marine diatoms sustain growth of bivalves in a Mediterranean lagoon. Journal of Sea Research 68, 20–32. https://doi.org/10.1016/j.seares.2011.11.004

Pernet, F., Tremblay, R., Comeau, L., Guderley, H., 2007. Temperature adaptation in two bivalve species from different thermal habitats: energetics and remodelling of membrane lipids. Journal of Experimental Biology 210, 2999–3014. https://doi.org/10.1242/jeb.006007

Perry, G.J., Volkman, J.K., Johns, R.B., Bavor, H.J., 1979. Fatty acids of bacterial origin in contemporary marine sediments. Geochimica et Cosmochimica Acta 43, 1715–1725. https://doi.org/10.1016/0016-7037(79)90020-6

R Core Team, 2012. R: A language and environment for statistical computing.

Ragueneau, O., Raimonet, M., Mazé, C., Coston-Guarini, J., Chauvaud, L., Danto, A., Grall, J., Jean, F., Paulet, Y.-M., Thouzeau, G., 2018. The impossible sustainability of the Bay of Brest? Fifty years of ecosystem changes, Interdisciplinary knowledge construction and key questions at the science-policy-community interface. Front. Mar. Sci. 5. https://doi.org/10.3389/fmars.2018.00124

Riera, P., 2007. Trophic subsidies of *Crassostrea gigas*, *Mytilus edulis* and *Crepidula fornicata* in the Bay of Mont Saint Michel (France): A δ^{13}C and δ^{15}N investigation. Estuarine, Coastal and Shelf Science 72, 33–41. https://doi.org/10.1016/j.ecss.2006.10.002

Riera, P., Stal, L., Nieuwenhuize, J., 2002. δ^{13}C versus δ^{15}N of co-occurring molluscs within a community dominated by *Crassostrea gigas* and *Crepidula fornicata* (Oosterschelde, The Netherlands). Mar. Ecol. Prog. Ser. 240, 291–295. https://doi.org/10.3354/meps240291

Rigolet, C., Thiébaut, E., Dubois, S., 2014. Food web structures of subtidal benthic muddy habitats: evidence of microphytobenthos contribution supported by an engineer species. Mar. Ecol. Prog. Ser. 500, 25–41. https://doi.org/10.3354/meps10685

Rossi, F., Herman, P.M.J., Middelburg, J.J., 2004. Interspecific and intraspecific variation of δ^{13}C and δ^{15}N in deposit- and suspension-feeding bivalves (*Macoma balthica* and *Cerastoderma*

edule): Evidence of ontogenetic changes in feeding mode of *Macoma balthica*. Limnol.
Oceanogr. 49, 408–414. https://doi.org/10.4319/lo.2004.49.2.0408
Roy, S., Llewellyn, C.A., Egeland, E.S., Johnsen, G., 2011. Phytoplankton Pigments,
Characterization, Chemotaxonomy, and Applications in Oceanography, Cambridge
University Press. ed. Cambridge, UK.
Shumway, S.E., Ward, J.E., Heupel, E., Holohan, B.A., Heupel, J., Heupel, T., Padilla, D.K., 2014.
Observations of feeding in the common Atlantic slippersnail *Crepidula fornicata* L., with
special reference to the "mucus net." Journal of Shellfish Research 33, 279–291.
https://doi.org/10.2983/035.033.0127
Vander Zanden, M.J., Clayton, M.K., Moody, E.K., Solomon, C.T., Weidel, B.C., 2015. Stable
isotope turnover and half-life in animal tissues: a literature synthesis. PLoS ONE 10,
e0116182. https://doi.org/10.1371/journal.pone.0116182
Walne, P.R., 1956. The biology and distribution of the slipper limpet *Crepidula fornicata* in Essex
rivers. Fishery Investigations Series II 20, 1–50.
Yee, A.K., Padilla, D.K., 2015. Allometric scaling of the radula in the Atlantic slippersnail
Crepidula fornicata. Journal of Shellfish Research 34, 903–907.
https://doi.org/10.2983/035.034.0320
Zhukova, N., Aizdacher, N., 1995. Fatty acid composition of 15 species of marine microalgae.
Phytochemistry 39, 351–356. https://doi.org/10.1016/0031-9422(94)00913-E
Zhukova, N.V., Kharlamenko, V.I., Svetashev, V.I., Rodionov, I.A., 1992. Fatty acids as markers
of bacterial symbionts of marine bivalve molluscs. Journal of Experimental Marine
Biology and Ecology 162, 253–263. https://doi.org/10.1016/0022-0981(92)90205-O
858

7. Supplementary materials

861 **Table S1: Pigment (%, mean ± SD, n ≥ 2) composition of organic matter sources (Biofilm, suspended particulate organic matter (SPOM), superficial**
862 **sedimentary organic matter (SSOM)) over the sampling survey. UK: Unknown pigments.**

Pigments	S1 = 26th February Biofilm	SPOM	SSOM	S2 = 21th March Biofilm	SPOM	SSOM	S3 = 28th March Biofilm	SPOM	SSOM	S4 = 12th April Biofilm	SPOM	SSOM	S5 = 14th June Biofilm	SPOM	SSOM
Alloxanthin	4 ± 2.2	11.7 ± 1.3	3.9 ± 0.1	1.2 ± 0.8	7.1 ± 1.2	3 ± 0.8	1.7 ± 1	9.3 ± 0.3	3.3 ± 0.3	2 ± 0.2	5.1 ± 0.8	4.1 ± 0.3	1.7 ± 0.6	15.1 ± 1.3	2.1 ± 0.4
β caroten	1.8 ± 0.6	0 ± 0	0.7 ± 0.2	0.2 ± 0.1	0.3 ± 0.2	0 ± 0	0.4 ± 0.1	0.6 ± 0.1	0 ± 0	1.5 ± 0.9	3.2 ± 3.4	0.1 ± 0.1	1.5 ± 0.5	2.8 ± 0.5	1.9 ± 1.6
Chlorophyll *a*	4.7 ± 0.6	7.7 ± 1.4	0.8 ± 0.2	4.1 ± 0.4	6.1 ± 0.7	0.3 ± 0.1	4.1 ± 0.4	7.2 ± 0.1	0.4 ± 0.1	4.7 ± 0.6	6.3 ± 0.6	0.3 ± 0.1	5 ± 0.4	4.7 ± 0.1	0.7 ± 0.1
Chlorophyll *b*	19.8 ± 1.7	7.8 ± 0.7	0.6 ± 0.1	25.8 ± 2.8	4.8 ± 0.5	0.3 ± 0.1	21.7 ± 0.1	7.1 ± 0.1	0.3 ± 0.1	19.6 ± 2.9	4.3 ± 0.6	0.2 ± 0.1	16 ± 0.6	5.1 ± 0	0.3 ± 0.2
Chlorophyllide	0.3 ± 0.1	0 ± 0	0.1 ± 0	0.3 ± 0.3	0.1 ± 0.1	0.1 ± 0	0.4 ± 0.1	0.2 ± 0	0.1 ± 0	0.5 ± 0.4	0.3 ± 0.1	0.1 ± 0	0.4 ± 0.1	0.2 ± 0	0 ± 0
Diadinoxanthin	0.8 ± 0.1	2.7 ± 0.4	1.1 ± 0.1	0.6 ± 0.1	3.4 ± 0.2	1 ± 0.1	0.7 ± 0.1	3.4 ± 0	0.9 ± 0.3	0.8 ± 0.1	2.1 ± 0.1	1.2 ± 0	1 ± 0	1.5 ± 0	1 ± 0.2
Xanthophyll	1.1 ± 0	2.5 ± 0.1	3.5 ± 0.2	0 ± 0	1.8 ± 1.6	2.9 ± 0.7	0.6 ± 0.1	2 ± 0.1	3.3 ± 0.3	0.7 ± 0.2	2.1 ± 0.2	3.1 ± 0.1	0.6 ± 0.1	3 ± 0.2	2.1 ± 0.5
Fucoxanthin-like	1.1 ± 0.1	2.7 ± 0.2	4.2 ± 0.4	1.2 ± 0.1	3.5 ± 0.1	5.6 ± 0.8	1.4 ± 0.4	4 ± 0.3	5.1 ± 0.2	2.8 ± 1.9	3 ± 0.1	5.7 ± 0.6	1.7 ± 0.2	3.1 ± 0.1	4.5 ± 0.5
Fucoxanthin	12.5 ± 2.8	22.5 ± 1.8	5.8 ± 0.9	10.8 ± 2.4	21.5 ± 1.4	5 ± 1.2	12.2 ± 3.4	19.2 ± 0.1	6.1 ± 1.3	12.3 ± 1.3	22.1 ± 0.9	6.8 ± 0.3	16.1 ± 2.3	11.3 ± 0.1	4.3 ± 1.1
Pheophorbide	14.8 ± 1.2	20 ± 2.7	32.8 ± 1.7	6.7 ± 2.4	23.7 ± 3	25.3 ± 1.6	8.8 ± 1.2	15.8 ± 0.5	26.9 ± 1.4	13.2 ± 3.4	22.3 ± 1.7	24.3 ± 0.6	12 ± 1.1	18.8 ± 2.8	20.8 ± 1.1
Lutein	3 ± 0.1	8.5 ± 0.8	13.9 ± 1.6	1.5 ± 0.5	7.5 ± 0.7	11.8 ± 2.3	2.2 ± 0.3	2.9 ± 0.4	10.6 ± 1	1.2 ± 0.3	4.3 ± 0.6	8.3 ± 1.6	1 ± 0.1	2 ± 0	3.8 ± 1.5
Neoxanthin	7.7 ± 1.2	1.7 ± 0.2	0 ± 0	14.8 ± 0.7	2.7 ± 0.4	0 ± 0	12.9 ± 0.5	5.1 ± 0.2	0 ± 0	6.1 ± 4.3	2.1 ± 0.1	2.9 ± 0.1	7.2 ± 0.9	2 ± 0.2	0.7 ± 1.2
Pheophytin	4.1 ± 3.3	0 ± 0	23.2 ± 1.1	1 ± 0.7	0.8 ± 1.4	32.9 ± 2.4	2.9 ± 0.6	1.3 ± 0.1	31.1 ± 1.8	7.9 ± 1.4	9.5 ± 5.8	30.1 ± 1.9	15 ± 1.5	17.9 ± 1.8	51.4 ± 2.6
Chlorophyll *c*	1.5 ± 0.5	4.8 ± 0.5	3.7 ± 0.2	0 ± 0	12.5 ± 1.5	6.4 ± 0.2	1.3 ± 0.2	15.2 ± 0.1	7.2 ± 0.7	1.3 ± 0.3	6.6 ± 1	8 ± 1.4	1.9 ± 0.1	4.1 ± 0.3	3.9 ± 0.4
UK 3	1.1 ± 0.3	0.9 ± 0	0.8 ± 0.1	1.6 ± 0.2	0.9 ± 0.4	0.7 ± 0.1	1.1 ± 0	1.6 ± 0.1	0.6 ± 0	1.1 ± 0.3	0.7 ± 0.4	1 ± 0.2	0.7 ± 0	0.7 ± 0.2	0.4 ± 0.1
UK 6	17.3 ± 1.3	0 ± 0	0 ± 0	26.7 ± 4.5	0 ± 0	0 ± 0	23 ± 1.3	0 ± 0	0 ± 0	20.4 ± 3.8	0 ± 0	0 ± 0	14.5 ± 0.9	0 ± 0	0 ± 0
UK 7	1.2 ± 0.1	0 ± 0	0 ± 0	1.7 ± 0.1	0 ± 0	0 ± 0	1.1 ± 0.3	0 ± 0	0 ± 0	1.6 ± 0.1	0 ± 0	0 ± 0	0.9 ± 0	0 ± 0	0 ± 0
UK 8	0.6 ± 0.1	0 ± 0	0 ± 0	1.1 ± 0.6	0 ± 0	0 ± 0	1 ± 0.8	0 ± 0	0 ± 0	0.7 ± 0.6	0 ± 0	0 ± 0	1.3 ± 0.5	0 ± 0	0 ± 0
UK 9	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.8 ± 0.3	0 ± 0	0 ± 0	0.8 ± 0	0.2 ± 0.2
Violaxanthin	0.6 ± 0.2	1.3 ± 0.3	0 ± 0	0 ± 0	0.7 ± 0.7	0 ± 0	0.9 ± 0.1	3.3 ± 0.4	0 ± 0	0.3 ± 0.5	1.8 ± 1.7	0 ± 0	0.2 ± 0.4	4.5 ± 0.1	0 ± 0
Zeaxanthin	2 ± 0.1	5.2 ± 0.2	5 ± 0.5	0.8 ± 0.4	2.5 ± 0.4	4.8 ± 1.1	1.6 ± 0	1.7 ± 0.1	4.2 ± 0.3	1.1 ± 0	3.2 ± 0.3	3.7 ± 0.6	0.9 ± 0.1	2.3 ± 0.1	1.9 ± 0.6
Chl *c* / Chl *a* ratio	0.3 ± 0.1	0.6 ± 0.2	4.5 ± 0.7	0 ± 0	2.1 ± 0.5	27.3 ± 17.1	0.3 ± 0.1	2.1 ± 0	17.1 ± 1.2	0.3 ± 0.1	1.1 ± 0.1	28 ± 14.9	0.4 ± 0.1	0.9 ± 0	5.8 ± 1.1
Chl *b* / Chl *a* ratio	4.2 ± 0.2	1 ± 0.1	0.7 ± 0.1	6.3 ± 0.7	0.8 ± 0	1.2 ± 0.5	5.3 ± 0.5	1 ± 0	0.7 ± 0.2	4.2 ± 0.6	0.7 ± 0	0.6 ± 0	3.2 ± 0.4	1.1 ± 0	0.5 ± 0.3
Fuco / Chl *a* ratio	2.6 ± 0.3	3 ± 0.8	7 ± 1.7	2.6 ± 0.6	3.6 ± 0.6	19.3 ± 6.7	2.9 ± 0.6	2.7 ± 0	14.5 ± 3.2	2.7 ± 0.5	3.5 ± 0.4	23 ± 9.8	3.2 ± 0.2	2.4 ± 0.1	6.3 ± 1.4

	S1 = 26th February			S2 = 21th March			S3 = 28th March			S4 = 12th April			S5 = 14th June		
Fatty acids	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM
TMTD	0.1 ± 0.1	0 ± 0	0 ± 0	0.1 ± 0.1	0.4 ± 0	0.6 ± 0	0.1 ± 0.1	0 ± 0	0.2 ± 0.2	0 ± 0	0 ± 0	0.3 ± 0	0 ± 0	0 ± 0	0 ± 0
15:0iso	1.8 ± 0.7	1 ± 0.2	2.7 ± 0.4	0.6 ± 0.5	1 ± 0.1	1.9 ± 0.8	1.3 ± 0.9	1.1 ± 0.1	2 ± 0.1	1.2 ± 0.4	1.3 ± 0.2	2.5 ± 0.2	1.4 ± 0.3	1 ± 0.2	2 ± 0.2
15:0ant	0.6 ± 0.1	1.2 ± 0.1	3.7 ± 0.7	0.3 ± 0.1	0.9 ± 0.1	2.9 ± 0.6	0.8 ± 0.2	1 ± 0.1	3 ± 0.4	0.5 ± 0.2	1.1 ± 0.2	3.2 ± 0.3	0.2 ± 0.2	0.4 ± 0.3	2.3 ± 0.8
16:0iso	0.8 ± 0.2	0.9 ± 0.2	2.3 ± 0.4	0.2 ± 0	0.5 ± 0	0.9 ± 0.1	0.6 ± 0.2	0.5 ± 0	0.8 ± 0.2	0.5 ± 0.2	0.5 ± 0.1	0.9 ± 0.1	0.6 ± 0.1	0.2 ± 0.1	0.5 ± 0.4
17:0iso	0.9 ± 0.3	1.7 ± 0.2	1.4 ± 0.2	0.1 ± 0.1	0.3 ± 0.1	1.2 ± 0.1	0.5 ± 0.4	2.8 ± 0.1	1.2 ± 0.1	0.8 ± 0.5	2.7 ± 0.2	1.6 ± 0.1	1.8 ± 0.6	0.8 ± 0.5	0.7 ± 0.6
17:0ant	2.5 ± 2.4	0.3 ± 0.3	0.9 ± 0.1	0.9 ± 0.6	1.2 ± 0	1.1 ± 0.1	2.6 ± 0.8	0 ± 0	0.6 ± 0	1.6 ± 0.5	0 ± 0	0.2 ± 0.2	1.2 ± 0.2	0.6 ± 0.1	0.9 ± 0.8
18:0iso	0.1 ± 0.1	1.2 ± 0.9	1.1 ± 0.2	0.1 ± 0.1	0.4 ± 0	0.6 ± 0.1	1.1 ± 0.1	0.9 ± 0.2	0.4 ± 0.1	0.4 ± 0.3	0.9 ± 0.3	0.6 ± 0.1	1.7 ± 0.4	1.2 ± 0	1.7 ± 0.4
∑ BFA	6.8 ± 2.4	6.3 ± 0.3	12.1 ± 1.4	2.2 ± 0.2	4.4 ± 0.1	8.5 ± 1.6	6.8 ± 0.7	6.2 ± 0.4	8 ± 0.7	5 ± 1.3	6.6 ± 0.9	9 ± 0.4	7 ± 0.6	4.2 ± 0.3	8.1 ± 1.5
14:0	3.1 ± 0.3	7.3 ± 1.3	4.9 ± 0.8	1.5 ± 0.4	5.2 ± 0.5	3.5 ± 0	4.2 ± 0.6	12 ± 0.8	4.1 ± 0.2	6.1 ± 2.1	8.6 ± 1	4.1 ± 0.3	4.9 ± 0.2	5.4 ± 1.1	5.3 ± 0.6
15:0	0.9 ± 0.3	2.6 ± 0.4	1.5 ± 0.2	0.2 ± 0.1	1.9 ± 0.1	1.2 ± 0.1	0.9 ± 0.2	1.7 ± 0.1	1 ± 0.1	0.8 ± 0.1	1.4 ± 0.2	1.2 ± 0.1	0.7 ± 0.1	0.8 ± 0.2	0.7 ± 0.6
16:0	21.5 ± 3.4	33 ± 4.9	22.4 ± 3.1	38.6 ± 0.7	26.3 ± 2.3	17.6 ± 1.6	29.5 ± 4.3	32.7 ± 1.9	14.8 ± 1.4	30.5 ± 4.4	34.1 ± 2.3	18.2 ± 0.2	19.9 ± 4.6	25 ± 4.7	19.3 ± 4.3
17:0	1.6 ± 0.7	1.6 ± 0.3	1.3 ± 0.2	0.8 ± 0.8	0.7 ± 0	0.9 ± 0.1	0.8 ± 0.2	0.9 ± 0.1	0.8 ± 0.1	0.7 ± 0.1	0.9 ± 0.1	1 ± 0.1	1.6 ± 0.4	0.2 ± 0.2	1.8 ± 0.5
18:0	16.5 ± 1.3	11.8 ± 1.7	9.6 ± 1.3	42.3 ± 2.6	13.3 ± 1.4	10.9 ± 2.5	21.5 ± 4.8	9.2 ± 0.7	6.8 ± 1	21.9 ± 3.8	14.9 ± 5.3	7.6 ± 0.3	6.7 ± 2.5	6.5 ± 1.3	8 ± 2.1
20:0	0.6 ± 0.1	1.7 ± 0.1	1.9 ± 1.7	0.6 ± 0	1.3 ± 0	2.8 ± 0.3	0.8 ± 0.2	0.9 ± 0	2.6 ± 0.3	0.9 ± 0.1	0.4 ± 0.7	3 ± 0.2	1.1 ± 0.7	0.9 ± 0.2	2.2 ± 0.3
22:0	1 ± 0.1	1.4 ± 1.2	3.2 ± 3	0.3 ± 0.2	1.6 ± 0	4.4 ± 0.7	1.1 ± 0.3	0.7 ± 0.6	4.5 ± 0.5	1.4 ± 0.4	1.4 ± 0.3	5.4 ± 0.3	1.2 ± 0.9	0.7 ± 0.1	3.5 ± 0.8
24:0	0.1 ± 0	0 ± 0	0.5 ± 0.4	0 ± 0	0 ± 0	0.1 ± 0.3	0.7 ± 0.3	4.4 ± 2	6.1 ± 5.2	0.1 ± 0.1	0.1 ± 0.2	0.3 ± 0.3	1.9 ± 1.5	1.1 ± 0.2	4.9 ± 1.2
∑ SFA	45.3 ± 4.7	59.4 ± 7.6	45.4 ± 3.3	84.2 ± 2.1	50.3 ± 3.8	41.4 ± 3.2	59.6 ± 9.9	62.4 ± 1.1	40.7 ± 7.3	62.3 ± 7.4	61.8 ± 5.9	40.7 ± 0.5	38.1 ± 10	40.5 ± 7.4	45.7 ± 8.3
14:1n-5	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0.5 ± 0.1	0.2 ± 0.2	0 ± 0	0 ± 0	0.2 ± 0.3	0.1 ± 0.1	0 ± 0
16:1n-11	0.3 ± 0.3	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0.6 ± 0.1	0 ± 0	0 ± 0	0 ± 0
16:1n-9	0.9 ± 0.1	1.5 ± 0.9	0.3 ± 0.5	0.2 ± 0.2	2 ± 0.2	1.2 ± 0.2	0.6 ± 0.3	0.4 ± 0.4	0.7 ± 0.2	0.9 ± 0.2	0.5 ± 0.1	0.9 ± 0.1	0.5 ± 0.4	0.8 ± 0.1	0.5 ± 0.4
16:1n-7	3.8 ± 0.1	1.9 ± 1.5	5.6 ± 3	0.8 ± 0.4	4.5 ± 0.5	8.8 ± 0.1	2.7 ± 1.8	0.9 ± 0.2	7.3 ± 2.4	2.5 ± 1.1	1.6 ± 0.7	0 ± 0	7.9 ± 0.6	3.4 ± 0.8	9 ± 1.9

16:1n-5	0.5 ± 0.1	0.8 ± 0.1	0.4 ± 0.4	0.1 ± 0	0.6 ± 0.1	0.7 ± 0.1	0.4 ± 0	0.5 ± 0.4	0.8 ± 0.2	0.3 ± 0.1	0.4 ± 0.1	1.2 ± 0.3	0.5 ± 0.4	0.1 ± 0.2	0.8 ± 0.7
17:1n-7	0.1 ± 0.2	0.4 ± 0.1	0.8 ± 0.3	0.2 ± 0.1	0.4 ± 0.1	0.6 ± 0.1	0.6 ± 0.3	0.8 ± 0.1	0.3 ± 0.2	0.7 ± 0.3	0.8 ± 0.1	0.7 ± 0.2	0.4 ± 0.3	1.2 ± 0.2	0.1 ± 0.3
18:1n-11	0.5 ± 0.3	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0.3 ± 0	0 ± 0	0 ± 0	0 ± 0
18:1n-9	1.6 ± 0.7	0.3 ± 0.5	1.3 ± 2.3	1 ± 0.7	2.5 ± 0.7	2.9 ± 0.2	0.9 ± 0.8	0 ± 0	1.9 ± 1	2.2 ± 0.9	0.2 ± 0.3	3.1 ± 1.1	1.8 ± 0.3	5.8 ± 1.4	2.3 ± 0.2
18:1n-7	2 ± 0.2	0 ± 0	0 ± 0	1.1 ± 0.3	2.7 ± 0.5	5.8 ± 0.3	2.1 ± 1.3	0.5 ± 0.5	4.2 ± 1.5	2 ± 0.8	0.6 ± 0.7	6.9 ± 1.4	4.3 ± 0.6	3.7 ± 1	4.6 ± 1.1
18:1n-5	0.6 ± 0.1	0.1 ± 0.2	0 ± 0	0.1 ± 0.1	0.1 ± 0.2	0 ± 0	0.4 ± 0	0 ± 0	0.3 ± 0.2	0.3 ± 0.2	0 ± 0	0.3 ± 0.3	0.6 ± 0.3	0 ± 0	0 ± 0
20:1n-11	1.2 ± 0.4	1.1 ± 1.5	0.3 ± 0.5	0.2 ± 0.1	3.6 ± 0.6	0.5 ± 0.4	0.5 ± 0.4	1.2 ± 1	0.6 ± 0.6	0.6 ± 0.3	0.4 ± 0.7	0.8 ± 0	0.5 ± 0.4	3.6 ± 0.8	0.1 ± 0.3
20:1n-9	0.3 ± 0	0.3 ± 0.5	0.3 ± 0.4	0.1 ± 0	0 ± 0	0.3 ± 0.3	0.3 ± 0.2	0 ± 0	0.4 ± 0.8	0.3 ± 0	0.5 ± 0.4	0.1 ± 0.2	0.3 ± 0.3	0 ± 0	0 ± 0
20:1n-7	0.6 ± 0.5	0 ± 0	0 ± 0	0.1 ± 0	0 ± 0	0 ± 0	0.2 ± 0.1	0 ± 0	0 ± 0	0.2 ± 0.1	0 ± 0	0.1 ± 0.2	0.3 ± 0.3	0 ± 0	0 ± 0
22:1n-9	0.2 ± 0.1	0 ± 0	0.3 ± 0.6	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
∑ MUFA	12.7 ± 1.3	6.6 ± 4.2	9.2 ± 6.4	3.9 ± 0.5	16.5 ± 2.7	20.7 ± 0.6	8.7 ± 4.8	4.3 ± 2.4	17 ± 6.6	10.4 ± 2.5	5.2 ± 1.9	15.1 ± 3.8	17.3 ± 3.4	18.6 ± 4.3	17.5 ± 4.7
16:2n-4	0.2 ± 0.2	0 ± 0	0.2 ± 0.4	0.2 ± 0.2	0.1 ± 0.1	0.7 ± 0.1	2.1 ± 0.7	0 ± 0	0.4 ± 0.5	0.9 ± 1.1	0.3 ± 0.3	0.2 ± 0.2	0.9 ± 0.2	1.5 ± 0.3	0.8 ± 0.7
16:3n-4	0.6 ± 0.1	0.3 ± 0.5	0.7 ± 0.6	0.1 ± 0.1	1.2 ± 0.1	1.3 ± 0.1	0 ± 0	0 ± 0	0 ± 0	0.5 ± 0.1	0 ± 0	1 ± 0.9	0.5 ± 0.4	0.3 ± 0.3	0.5 ± 0.4
16:3n-6	0.4 ± 0.2	0.2 ± 0.3	0 ± 0	0.1 ± 0.1	0 ± 0	0.3 ± 0.3	0.4 ± 0.3	0 ± 0	0.2 ± 0.4	0.3 ± 0.1	0.1 ± 0.2	1.1 ± 0.2	0.3 ± 0.3	0 ± 0	0 ± 0
16:3n-3	0.2 ± 0.1	0 ± 0	0.2 ± 0.3	0.1 ± 0.1	1 ± 0.1	1.2 ± 0.3	0.6 ± 0.5	0 ± 0	1.2 ± 0.9	0.4 ± 0.1	0.3 ± 0.3	0.9 ± 0.2	0.1 ± 0.1	0.1 ± 0.2	0.2 ± 0.3
16:4n-3	0.3 ± 0.1	1.3 ± 1.1	1.2 ± 0.1	0.2 ± 0.2	1.3 ± 0.1	0.7 ± 0.1	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
18:2n-6	4.2 ± 2.1	0.9 ± 0.8	0.7 ± 1	1.9 ± 0.4	1.2 ± 0.2	1.1 ± 0.2	3 ± 1.8	0.9 ± 0.1	1.2 ± 0.3	2.3 ± 0.9	0.9 ± 0.2	1.4 ± 0.3	3.1 ± 0.6	2.5 ± 0.5	1.3 ± 0.4
18:2n-4	0.2 ± 0.1	0.2 ± 0.4	0 ± 0	0 ± 0.1	0.2 ± 0.2	0.4 ± 0.1	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
18:3n-6	0.2 ± 0	0.1 ± 0.3	0.4 ± 0.7	0.1 ± 0	0.3 ± 0	0 ± 0	0.3 ± 0	0 ± 0	0.2 ± 0.2	0.2 ± 0.2	0.1 ± 0.2	0.4 ± 0.1	0.2 ± 0.1	0.2 ± 0.3	0 ± 0
18:3n-4	0.6 ± 0.1	1.4 ± 0.5	5.9 ± 1.7	0.1 ± 0	1.2 ± 0.4	1.6 ± 0.1	0.4 ± 0	0.8 ± 0.3	2.5 ± 1.7	0.2 ± 0	0.7 ± 0.3	0.7 ± 0.1	0 ± 0.1	0.2 ± 0.1	3.2 ± 0.4
18:3n-3	0.7 ± 0.3	0.6 ± 0.7	0.3 ± 0.5	0.2 ± 0.1	1.3 ± 0.2	0.7 ± 0	0.5 ± 0.2	0.2 ± 0.3	0.4 ± 0.4	0.3 ± 0.2	0.4 ± 0.4	0.7 ± 0.1	0.5 ± 0.4	2.8 ± 0.8	0.3 ± 0.5
18:4n-3	1.1 ± 0.9	2.7 ± 0.3	1.2 ± 0.2	0.2 ± 0.1	3.8 ± 0.5	1.6 ± 0.2	1 ± 0.4	2.5 ± 0.5	2.2 ± 0.6	0.8 ± 0.3	1.9 ± 0.6	2.5 ± 0.5	1.1 ± 0.4	4.3 ± 1.2	0.9 ± 0.9
20:2n-6	0.6 ± 0.2	1.3 ± 0.2	0.9 ± 0.2	0.1 ± 0	0.1 ± 0.2	0.2 ± 0.3	0.3 ± 0	0 ± 0	0.4 ± 0.7	0.4 ± 0.4	1.1 ± 0.2	1.3 ± 0.1	0.5 ± 0.4	1.6 ± 0.5	1.3 ± 0.6
20:3n-6	0.2 ± 0	0.5 ± 0	0 ± 0	0.1 ± 0	0.5 ± 0	0.5 ± 0.1	0.2 ± 0.1	0 ± 0	0.4 ± 0.1	0.3 ± 0.2	0.3 ± 0.6	0.4 ± 0	0.2 ± 0.2	0 ± 0	0 ± 0
20:3n-3	0.2 ± 0	0.5 ± 0.9	0 ± 0	0.1 ± 0	0 ± 0	0.3 ± 0.2	0 ± 0	0 ± 0	0 ± 0	0.2 ± 0.1	0 ± 0	0.4 ± 0	1.7 ± 1.5	0.4 ± 0.2	0.2 ± 0.3
20:4n-6	6.3 ± 0.9	0 ± 0	0.7 ± 0.6	1.6 ± 0.2	0.2 ± 0.2	1.5 ± 0.1	2.8 ± 1.7	0.2 ± 0.3	1.2 ± 0.4	2 ± 0.7	0.1 ± 0.2	2.2 ± 0.4	5.9 ± 1.6	0.2 ± 0.2	1.8 ± 0.7
20:4n-3	0.4 ± 0.2	1.6 ± 0.2	1.1 ± 0.2	0.1 ± 0	0.7 ± 0	0.7 ± 0.1	0.8 ± 0.3	2.3 ± 0.5	0.6 ± 0.1	1.3 ± 0.5	2.2 ± 0.9	0.5 ± 0.4	0.2 ± 0.2	1.7 ± 1	0 ± 0
20:5n-3	8.2 ± 2.1	2.2 ± 1.2	4.6 ± 1	1.4 ± 0.7	4.5 ± 0.6	6.3 ± 1.2	3.4 ± 2.5	0.8 ± 0.3	3 ± 3	2.2 ± 0.6	1.7 ± 1	9.7 ± 2.3	11.2 ± 4.4	6.7 ± 1.8	8.5 ± 3.1
21:5n-3	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	3 ± 5.2	2.3 ± 3.4	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0

22:2n-6	0.3 ± 0.1	0.4 ± 0	0.9 ± 0.1	0.3 ± 0	0.9 ± 0	1.3 ± 0.2	2.1 ± 1.5	7.3 ± 1	2.8 ± 2	1.5 ± 0.8	9.1 ± 3.2	3 ± 2.5	3 ± 2.2	2.7 ± 1.2	2.9 ± 1.6
22:4n-6	0.6 ± 0.5	0.5 ± 0.1	0 ± 0	0.1 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
22:5n-6	0.7 ± 0.1	2.1 ± 1.3	1.7 ± 0.5	0.1 ± 0	0.4 ± 0	0.6 ± 0.1	1.9 ± 0.2	4.4 ± 0.5	2.5 ± 0.9	1.6 ± 0.6	2.3 ± 0.8	0.9 ± 0.4	1 ± 1	1.7 ± 0.7	0.2 ± 0.3
22:5n-3	0.7 ± 0.8	0 ± 0	0 ± 0	0.1 ± 0	0.1 ± 0.2	0 ± 0	0.7 ± 0.1	2.1 ± 0.6	2.1 ± 1.7	0.6 ± 0.5	2.2 ± 0.5	2.3 ± 0.7	0.5 ± 0.3	1.3 ± 1.2	0.6 ± 0.6
22:6n-3	4.3 ± 2.8	0.9 ± 1.1	0.7 ± 0.7	0.5 ± 0.2	2.9 ± 0.4	1.9 ± 0.2	1.7 ± 0.3	4 ± 0.5	4.5 ± 0.8	1.5 ± 2.1	0 ± 0	3.6 ± 1.7	3.7 ± 1.7	6 ± 1.8	2.9 ± 1.1
∑ PUFA	31.2 ± 6.5	17.6 ± 4.7	21.5 ± 3.2	7.8 ± 1.6	22 ± 1.8	22.9 ± 1.8	22.3 ± 6.1	25.6 ± 1.6	28.7 ± 2.3	19.8 ± 4.7	23.9 ± 4.9	33.3 ± 3.1	34.5 ± 7	33.9 ± 4.9	25.6 ± 5.6
A	0.5 ± 0.4	0 ± 0	0 ± 0	0.1 ± 0	0.1 ± 0.1	0 ± 0	0.2 ± 0	0 ± 0	0 ± 0	0.3 ± 0	0 ± 0	0 ± 0	1.2 ± 1.4	0 ± 0	0 ± 0
M	0.8 ± 0.2	0.4 ± 0	0.2 ± 0.4	0.2 ± 0.1	0.4 ± 0	0 ± 0	0.5 ± 0.2	0 ± 0	0 ± 0	0.4 ± 0.2	0 ± 0	0 ± 0	0.4 ± 0.3	0 ± 0	0 ± 0
N	0.4 ± 0.1	0.5 ± 0	1.4 ± 1.8	0.2 ± 0.1	0.5 ± 0	0.6 ± 0.1	0.5 ± 0.2	0 ± 0	0.1 ± 0.2	0.4 ± 0	0.7 ± 0.6	0.4 ± 0	0 ± 0	0 ± 0	0 ± 0
O	0.8 ± 0	5.2 ± 2.1	5.5 ± 3.9	0.8 ± 0.6	3.1 ± 0.6	0.5 ± 0.5	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1 ± 0.6	2.2 ± 1.5	2.7 ± 2.1
P	1.4 ± 0	4 ± 0.6	4.9 ± 1.1	0.5 ± 0.2	1.9 ± 0.1	3.9 ± 0.5	1.1 ± 0.5	0.9 ± 0.8	4.8 ± 1.3	1.4 ± 1.5	1.2 ± 0.3	0.6 ± 0.6	0.4 ± 0.3	0.3 ± 0.3	0.5 ± 0.5
W	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0	0.6 ± 0	0.8 ± 0.1	0.1 ± 0.1	0.6 ± 0	0.5 ± 0	0.2 ± 0	0.7 ± 0.6	0.6 ± 0.1	0.2 ± 0.2	0.2 ± 0.2	0 ± 0
∑ UK FA	4 ± 0.7	10.1 ± 1.5	11.9 ± 5.2	1.7 ± 1.1	6.5 ± 0.6	5.8 ± 1.1	2.4 ± 0.6	1.5 ± 0.8	5.4 ± 1.3	2.6 ± 1.4	2.6 ± 1.3	1.6 ± 0.6	3.2 ± 1.2	2.7 ± 1.5	3.2 ± 1.9
PUFA/SFA	0.7 ± 0.2	0.3 ± 0.1	0.5 ± 0.1	0.1 ± 0	0.4 ± 0.1	0.6 ± 0.1	0.4 ± 0.2	0.4 ± 0	0.7 ± 0.2	0.3 ± 0.1	0.4 ± 0.1	0.8 ± 0.1	1 ± 0.4	0.9 ± 0.3	0.6 ± 0.2
EPA/DHA	2.3 ± 1	NA	NA	2.8 ± 0.5	1.6 ± 0	3.4 ± 0.4	1.9 ± 1.3	0.2 ± 0.1	0.8 ± 0.8	NA	NA	3.1 ± 1.6	3.1 ± 0.6	1.1 ± 0	2.9 ± 0.1
∑ n-3/ ∑ n-6	1.2 ± 0.6	1.7 ± 0.4	1.8 ± 0.5	0.7 ± 0.3	4.4 ± 0	2.5 ± 0.4	0.8 ± 0.2	0.9 ± 0.1	2.1 ± 1.1	1.2 ± 0.4	0.6 ± 0.1	1.9 ± 0.4	1.3 ± 0.3	2.7 ± 0.7	2 ± 1.1

867
868

Table S3: Stable isotope (mean ± SD, n = 3) composition of organic matter sources (Biofilm, suspended particulate organic matter (SPOM), superficial sedimentary organic matter (SSOM)) over the sampling survey.

	S1 = 26th February			S2 = 21th March			S3 = 28th March			S4 = 12th April			S5 = 14th June		
	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM	Biofilm	SPOM	SSOM
$\delta^{13}C$	-22.6 ± 0.1	-25.2 ± 0.2	-23 ± 0.2	-23.4 ± 0.5	-25.3 ± 0.1	-22.7 ± 0.3	-22.6 ± 0.4	-25.3 ± 0.1	-22.6 ± 0.1	-23.4 ± 1	-24.6 ± 0.4	-22.5 ± 0.1	-21.8 ± 0.2	-25.1 ± 0.3	-22.5 ± 0.1
$\delta^{15}N$	9.4 ± 0.6	7.3 ± 0	7 ± 0.7	10.3 ± 0.5	6.7 ± 0.3	7.1 ± 0.1	9.2 ± 0.7	6.6 ± 0.1	7 ± 0.1	9.2 ± 0.7	6.9 ± 0.9	7.1 ± 0.1	9.4 ± 0.7	6.4 ± 0.5	6.9 ± 0.2
C/N	3.9 ± 0.3	6.6 ± 1.1	5.1 ± 1.3	5.8 ± 3	5.9 ± 0.6	5.3 ± 0.6	5.3 ± 0.4	6 ± 0.5	5.5 ± 0.6	5 ± 0.5	5.6 ± 0.9	5.4 ± 0.5	3.6 ± 0.6	3.9 ± 0.4	5.9 ± 0.8

Table S4: Fatty acid (FA) (%, mean ± SD, n = 5) composition of the three ontogenic stages of *Crepidula fornicata* (motile males, sessile males, sessile females) over the sampling survey. Only FA accounting for more than 0.5 % of total FA in at least one sample was shown. BFA: Branched FA; SFA: saturated FA; MUFA: monounsaturated FA; PUFA: polyunsaturated FA; NMI FA: non-methyl-interrupted FA; DMA: Dimethyl acetals FA; EPA: 20:5n-3; DHA: 22:6n-3.

Fatty acids	S1 = 26th February			S2 = 21th March			S3 = 28th March			S4 = 12th April			S5 = 14th June		
	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male
TMTD	1.3 ± 0.3	2.1 ± 1.1	0.8 ± 0.6	1.3 ± 0.2	0 ± 0.1	0.5 ± 0.4	1.3 ± 0.7	2.5 ± 0.4	0.9 ± 1.2	2.1 ± 1	2.9 ± 0.8	2.5 ± 1	2.1 ± 1.7	2 ± 0.7	1 ± 0.6
15:0iso	0.2 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.2	0.1 ± 0.1	0 ± 0	0 ± 0	0.1 ± 0.1	0.1 ± 0.1	0.3 ± 0.4	0.1 ± 0.1	0.5 ± 0.3	1 ± 0.2
15:0ant	0.1 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0.1	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0.3 ± 0.5	0 ± 0	0.4 ± 0.3	0.7 ± 0.1
16:0iso	1.3 ± 0.5	1.1 ± 0.3	1.1 ± 0.1	0.7 ± 0.1	0.9 ± 0.2	0.1 ± 0.2	0.9 ± 0.1	1 ± 0.1	0.6 ± 1	0.8 ± 0.2	0.5 ± 0.1	0.6 ± 0.4	0.7 ± 0.2	0.3 ± 0	0.4 ± 0.2
17:0iso	5.4 ± 1.8	4.1 ± 1	3.4 ± 0.5	3.1 ± 1.8	1.7 ± 2.4	1.1 ± 1	3.5 ± 2	2.8 ± 1.4	2.5 ± 1.1	3.8 ± 1	2.4 ± 0.1	3 ± 0.3	3 ± 1	1.3 ± 0.2	1.4 ± 0.3
17:0ant	3.6 ± 1.6	3.2 ± 0.7	2.4 ± 0.6	2.1 ± 0.5	3.5 ± 1	1.5 ± 0.3	2.8 ± 0.7	3.1 ± 0.4	3 ± 1	2.4 ± 0.7	1.8 ± 0.2	2.5 ± 0.3	2.4 ± 0.9	1.2 ± 0.2	1.5 ± 0.3
18:0iso	0.7 ± 0.4	0.6 ± 0.4	0.3 ± 0.4	0.3 ± 0.3	0.2 ± 0.3	0 ± 0	0.5 ± 0.5	0.9 ± 0.2	0.2 ± 0.4	0.4 ± 0.6	0.7 ± 0.1	2.1 ± 2.9	0.5 ± 0.4	0.1 ± 0.1	0 ± 0
∑ BFA	11.2 ± 4.3	8.9 ± 2.1	7.1 ± 1.3	6.3 ± 2.3	6.4 ± 2.6	2.8 ± 1	7.8 ± 2.6	7.8 ± 1.8	6.3 ± 2.2	7.5 ± 2.2	5.6 ± 0.3	8.9 ± 4	6.6 ± 2.6	3.8 ± 1	5 ± 0.9
14:0	1.5 ± 0.3	1.3 ± 0.7	1.4 ± 0.4	1.3 ± 0.2	3 ± 0.9	1.6 ± 0.1	1.3 ± 0.4	1.4 ± 0.3	0.8 ± 0.8	2 ± 0.5	2.3 ± 0.4	1.9 ± 0.7	2.4 ± 0.6	2.7 ± 0.6	2.2 ± 0.2
15:0	0.6 ± 0.1	0.6 ± 0.2	0.8 ± 0.2	0.4 ± 0	0.5 ± 0.1	0.6 ± 0.4	0.5 ± 0.1	0.4 ± 0	0.1 ± 0.3	0.5 ± 0.1	0.5 ± 0.1	0.6 ± 0	0.4 ± 0	0.7 ± 0.4	1.5 ± 0.3
16:0	11.9 ± 0.8	16.7 ± 3.1	25.2 ± 3.5	11.7 ± 1	11.6 ± 3.3	33.7 ± 1.7	10.9 ± 1	13 ± 0.9	12.3 ± 2.3	12 ± 0.4	12.6 ± 3.1	14.2 ± 2.1	11.2 ± 0.9	9.7 ± 1.4	7.6 ± 0.9
17:0	1.5 ± 0.1	1.2 ± 0.3	1 ± 0.1	1.4 ± 0.1	1.2 ± 0.1	0.7 ± 0.4	1 ± 0.5	0.8 ± 0.1	0.4 ± 0.5	0.9 ± 0.5	0.8 ± 0.1	1 ± 0.1	1.1 ± 0.2	0.7 ± 0.1	0.7 ± 0.1
18:0	6.3 ± 0.9	17.7 ± 7.3	27.5 ± 4.2	6.8 ± 0.7	9.2 ± 3.9	35.6 ± 0.9	6.6 ± 0.9	10.4 ± 1.3	14.8 ± 4.5	6.9 ± 0.4	9.1 ± 5.6	14.3 ± 3.1	6.9 ± 1.2	5.6 ± 1.1	7.5 ± 1.1
20:0	0.5 ± 0.1	0.7 ± 0.2	0.5 ± 0.3	0.5 ± 0.3	0.2 ± 0.2	0.6 ± 0.3	0.2 ± 0	0.2 ± 0.3	0 ± 0	0.2 ± 0	0.2 ± 0.1	0.1 ± 0.2	0.4 ± 0.1	0.2 ± 0.1	0.3 ± 0
22:0	0.5 ± 0.1	0.3 ± 0.2	0.1 ± 0.2	0.4 ± 0.1	0.2 ± 0	0 ± 0	0.4 ± 0.1	0.1 ± 0.1	0 ± 0	0.3 ± 0.1	0.2 ± 0.1	0 ± 0	0.3 ± 0.1	0 ± 0	0 ± 0
24:0	0.4 ± 0.1	0.6 ± 0.4	0.6 ± 0.4	0.2 ± 0	0.2 ± 0.1	0.7 ± 0.2	0.3 ± 0.2	0.1 ± 0.1	1.6 ± 1.8	0.4 ± 0.1	0.3 ± 0.1	1.2 ± 0.5	0.3 ± 0.1	0.1 ± 0.1	0 ± 0
∑ SFA	23.2 ± 1.5	39.2 ± 9.6	57 ± 8.2	22.9 ± 1.5	26 ± 7.1	73.5 ± 2.4	21.4 ± 1.1	26.4 ± 1.7	30.1 ± 5.9	23.3 ± 0.9	25.8 ± 8.6	33.3 ± 4.6	23.1 ± 1.7	19.9 ± 1.2	19.7 ± 1.6
14:1n-5	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0.2 ± 0.5	0.1 ± 0.1	0 ± 0	0 ± 0
16:1n-11	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.4 ± 0.1	0.7 ± 0.4	0 ± 0	0.1 ± 0.1	0.2 ± 0.1	1.1 ± 1	0 ± 0	0.1 ± 0.1	0.2 ± 0.1
16:1n-9	0.7 ± 0.2	0.5 ± 0.3	0.3 ± 0.3	0.7 ± 0.2	0.1 ± 0.2	0.1 ± 0.2	0.5 ± 0.1	0.7 ± 0.5	0.4 ± 1	0.5 ± 0	0.6 ± 0.1	0.7 ± 0.3	0.3 ± 0	0.5 ± 0.4	4.1 ± 2.9
16:1n-7	1.4 ± 0.6	0.9 ± 0.2	0.8 ± 0.3	1.7 ± 0.2	1.3 ± 0.4	0.3 ± 0.4	1.5 ± 0.3	1.2 ± 0.4	2.6 ± 1.3	2.3 ± 0.6	2.9 ± 0.8	2.1 ± 0.3	3.6 ± 0.9	5.8 ± 1.9	6.4 ± 1.9
16:1n-5	0.4 ± 0.4	0 ± 0	0 ± 0	0.3 ± 0.1	0.1 ± 0.1	0 ± 0	0.3 ± 0.1	0.8 ± 0.4	1.4 ± 1.4	0.5 ± 0.2	0.4 ± 0.2	1.8 ± 1.2	0.9 ± 0.4	3.4 ± 2.1	2.2 ± 4.3

17:1n-7	0.1 ± 0.1	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0 ± 0	0.4 ± 0.5	0.1 ± 0.3	0 ± 0	0.3 ± 0.4	0 ± 0.1	0.2 ± 0.3	0 ± 0	0 ± 0	0 ± 0
18:1n-11	0.5 ± 0.2	0.4 ± 0.1	0.1 ± 0.3	0.5 ± 0.1	0.4 ± 0.1	0 ± 0	0.5 ± 0.2	0.9 ± 0.2	1.6 ± 1	0.8 ± 0.2	0.5 ± 0.1	1.4 ± 1	0.7 ± 0.2	2.1 ± 1.4	6 ± 1
18:1n-9	0 ± 0.1	0 ± 0	5.1 ± 2.3	2.1 ± 0.3	2.1 ± 0.2	4 ± 0.8	2 ± 0.2	2.3 ± 0.1	1.6 ± 1	2 ± 0.2	2 ± 0.1	1.8 ± 0.4	1.5 ± 0.2	1.5 ± 0.2	1.2 ± 0.2
18:1n-7	2.6 ± 0.2	1.8 ± 0.3	1.4 ± 0.2	2.8 ± 0.3	2.1 ± 0.3	1 ± 0.1	2.5 ± 0.2	1.7 ± 0.3	0.7 ± 1.1	2.3 ± 0.3	2.3 ± 0.5	2 ± 0.9	3.3 ± 0.4	4.4 ± 1	2.6 ± 0.8
18:1n-5	0.4 ± 0	0.2 ± 0.2	0.1 ± 0.2	0.4 ± 0	0.4 ± 0	0.2 ± 0.4	0.4 ± 0	0.2 ± 0.2	0.3 ± 0.4	0.3 ± 0	0.3 ± 0.1	0.5 ± 0.3	0.5 ± 0.1	0.6 ± 0.1	0.6 ± 0.1
20:1n-11	5.8 ± 0.5	5.3 ± 0.7	3.2 ± 0.7	6.1 ± 0.8	6.4 ± 0.9	1.9 ± 0.5	5.6 ± 0.6	4 ± 0.9	2.3 ± 1.5	5.2 ± 0.9	4.1 ± 1	2.3 ± 0.6	5.9 ± 0.6	4.3 ± 0.8	2.5 ± 0.3
20:1n-9	0.9 ± 0.1	1 ± 0.2	0.5 ± 0.4	0.8 ± 0	1.2 ± 0.1	0.1 ± 0.2	0.8 ± 0.2	1 ± 0.2	0.4 ± 0.5	0.8 ± 0.1	0.9 ± 0.1	0.7 ± 0.1	0.6 ± 0.2	0.7 ± 0.1	0.4 ± 0.1
20:1n-7	4.7 ± 0.5	3.9 ± 0.6	2.4 ± 0.8	4.6 ± 0.2	5.2 ± 0.9	1.3 ± 0.2	4.6 ± 0.4	3.5 ± 0.8	5.9 ± 4.3	4.2 ± 0.3	4.2 ± 1.1	4 ± 1	5.5 ± 0.9	5.5 ± 0.7	5 ± 0.8
∑ MUFA	17.8 ± 1.1	14 ± 2.1	13.9 ± 2.5	20.4 ± 0.6	19.4 ± 1.6	8.8 ± 0.7	19.9 ± 1	17.2 ± 1.5	17.2 ± 1.5	19.9 ± 0.9	18.8 ± 2.8	18.9 ± 2.8	22.9 ± 1.1	28.7 ± 1.4	31.3 ± 0.5
16:2n-7	0 ± 0	0 ± 0	0 ± 0	0.7 ± 1.5	2.6 ± 2.5	0.7 ± 1	0.8 ± 1.7	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
16:2n-4	0.4 ± 0.5	0.1 ± 0.1	0 ± 0	0.2 ± 0	0.5 ± 0.4	0.2 ± 0.5	0.7 ± 0.3	1 ± 0.9	0.2 ± 0.4	0.4 ± 0.2	0.7 ± 0.3	0.1 ± 0.3	0.5 ± 0.2	0.7 ± 0.4	0.6 ± 0.2
16:3n-4	0.1 ± 0.1	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0.1	0 ± 0	0 ± 0.1	0.1 ± 0.1	0 ± 0	0.1 ± 0.2	0.3 ± 0.2	0.3 ± 0.3	0.2 ± 0.1	0.3 ± 0.1	0.1 ± 0.1
16:3n-6	0.4 ± 0.3	0.6 ± 0.2	0.2 ± 0.3	0.5 ± 0.1	0.1 ± 0.2	0 ± 0	0.1 ± 0.2	0.1 ± 0.3	0 ± 0	0.5 ± 0.2	0.2 ± 0.1	0 ± 0	0.5 ± 0.2	0 ± 0	0 ± 0
16:3n-3	0.1 ± 0.1	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0.1	0 ± 0	0.1 ± 0.1	0.2 ± 0.2	0.5 ± 0.7	0.3 ± 0.1	0.2 ± 0.1	1.8 ± 2.7	1.6 ± 0.9	0.6 ± 0.2	0.8 ± 0.5
16:4n-3	0.3 ± 0.2	0.2 ± 0.2	0 ± 0	0.3 ± 0.1	0.6 ± 0.5	0.1 ± 0.2	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.2 ± 0.1	0.4 ± 0.1	0.6 ± 0.1
18:2n-6	1.3 ± 0.2	1.4 ± 0.3	1.6 ± 0.3	1.2 ± 0.2	0.9 ± 0.2	0.7 ± 0.4	1.1 ± 0.1	1.5 ± 0.2	0.9 ± 0.6	1.3 ± 0.1	1.4 ± 0.2	1.6 ± 0.5	0.9 ± 0.1	3 ± 1.3	6.3 ± 1.4
18:2n-4	0.4 ± 0.1	0.2 ± 0.2	0 ± 0	0.5 ± 0.1	0.3 ± 0.1	0 ± 0	0.5 ± 0	0.4 ± 0	0 ± 0	0.4 ± 0.1	0.3 ± 0.1	0 ± 0	0.8 ± 0.2	0.7 ± 0.2	0.4 ± 0.3
18:3n-6	0.1 ± 0	0 ± 0	0 ± 0	0.1 ± 0	0.1 ± 0.2	0 ± 0	0.2 ± 0	0 ± 0.1	0 ± 0	0.1 ± 0.1	0.1 ± 0.1	0 ± 0	0.3 ± 0.2	0.2 ± 0.1	0.2 ± 0.3
18:3n-4	1 ± 0.5	2.7 ± 1.2	3.1 ± 1.7	0.4 ± 0.3	0.4 ± 0.5	0 ± 0	1.1 ± 0.3	3.5 ± 1.4	1.3 ± 1.3	1 ± 0.4	0.6 ± 0.3	2.4 ± 0.8	0.3 ± 0.1	0.3 ± 0	0 ± 0
18:3n-3	0.9 ± 0.2	0.2 ± 0.2	0 ± 0	0.9 ± 0.2	1 ± 0.8	1.7 ± 1	0.9 ± 0.1	1.2 ± 0.3	2.1 ± 3.3	1.1 ± 0.2	1.6 ± 0.4	0.7 ± 0.4	0.5 ± 0.1	0.9 ± 0.2	0.5 ± 0.2
18:4n-3	0.7 ± 0.6	0 ± 0	0.6 ± 0.6	1.8 ± 0.4	1.4 ± 0.5	2.3 ± 0.2	1.7 ± 0.3	3 ± 0.6	2.9 ± 0.7	3.1 ± 1	4.4 ± 0.8	3.2 ± 0.8	2.7 ± 0.5	2 ± 0.3	1.5 ± 0.2
18:4n-1	0.1 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0.1 ± 0.1	0 ± 0	0 ± 0.1	0 ± 0	0.4 ± 1	0 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0.1 ± 0.1	0 ± 0
20:2n-6	1.2 ± 0.1	2.2 ± 0.5	1.1 ± 0.2	1.2 ± 0.2	2.1 ± 0.6	0.9 ± 0.2	1.1 ± 0.2	1.1 ± 0.2	0.4 ± 0.5	1 ± 0.2	1 ± 0.1	0.5 ± 0.4	0.9 ± 0.1	0.6 ± 0.1	0.4 ± 0
20:3n-6	0.1 ± 0	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0.1	0 ± 0	0.2 ± 0.1	0 ± 0	0.3 ± 0.6	0.1 ± 0.1	0 ± 0.1	0.1 ± 0.3	0.2 ± 0.1	0.1 ± 0.1	0.1 ± 0.1
20:3n-3	0.4 ± 0.1	0.1 ± 0.1	0 ± 0	0.4 ± 0.1	0.2 ± 0.1	0 ± 0	0.4 ± 0.1	0.3 ± 0.3	0 ± 0	0.4 ± 0.1	0.4 ± 0.1	0 ± 0	0.3 ± 0.1	0.2 ± 0	0.1 ± 0.1
20:4n-6	2.7 ± 0.3	3.5 ± 0.5	2.3 ± 0.6	2.7 ± 0.4	3.5 ± 0.3	1.2 ± 0.1	2.6 ± 0.6	2.6 ± 0.5	0.8 ± 1.2	2.5 ± 0.4	1.7 ± 0.4	1.4 ± 0.4	1.7 ± 0.2	1.2 ± 0.1	0.7 ± 0.1
20:4n-3	0.6 ± 0.1	0.3 ± 0.2	0.3 ± 0.3	0.6 ± 0.2	0.4 ± 0.1	0 ± 0	0.5 ± 0.1	0.6 ± 0.1	0.5 ± 0.8	0.6 ± 0.1	0.8 ± 0.2	0.3 ± 0.3	0.4 ± 0.1	0.5 ± 0.1	0.4 ± 0.1
20:5n-3	10.5 ± 2.6	5.7 ± 1.2	3.3 ± 0.7	12.1 ± 1.3	8.3 ± 1.3	2.5 ± 0.5	11.1 ± 1.8	6.9 ± 0.9	3.4 ± 1.7	11.4 ± 1.8	11.1 ± 2.6	5.8 ± 1.9	14.6 ± 2.4	16.3 ± 3.4	9.4 ± 2.1
21:5n-3	0.4 ± 0	0.3 ± 0.2	0.1 ± 0.3	0.5 ± 0.1	0.6 ± 0.1	0 ± 0	1.5 ± 0.6	0.7 ± 0.2	0 ± 0	0.5 ± 0.1	0.8 ± 0.2	0.1 ± 0.2	0.9 ± 0.2	0.8 ± 0.1	0.5 ± 0.1

22:2n-6	0.2 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.4 ± 0.1	0.2 ± 0.3	0 ± 0	0 ± 0	0.1 ± 0.1	0.1 ± 0.1	0 ± 0	0.3 ± 0.1	0.1 ± 0	0.4 ± 0.2
22:4n-6	0.6 ± 0.2	0.4 ± 0.1	0.1 ± 0.2	0.5 ± 0.3	0.4 ± 0.1	0 ± 0	0.6 ± 0.1	0.2 ± 0.2	0 ± 0	0.6 ± 0.2	0.2 ± 0.1	0.4 ± 0.4	0.3 ± 0.1	0.2 ± 0	0 ± 0.1
22:5n-6	0.4 ± 0.1	0.3 ± 0.2	0.1 ± 0.2	0.5 ± 0.1	0.4 ± 0.1	0 ± 0	0.4 ± 0.1	0.4 ± 0.4	0.5 ± 1	0.4 ± 0.1	0.4 ± 0.1	0.3 ± 0.3	0.2 ± 0.1	0.2 ± 0	0 ± 0.1
22:5n-3	2 ± 0.5	1.2 ± 0.3	0.5 ± 0.5	2.4 ± 0.3	1.2 ± 0.2	0.4 ± 0.9	2.2 ± 1.3	1.4 ± 0.4	2.1 ± 2.2	2.2 ± 0.5	2.8 ± 2.2	2.8 ± 1.8	1.6 ± 0.3	1.4 ± 0.1	0.9 ± 0.1
22:6n-3	9.7 ± 2.1	4.9 ± 0.9	2.6 ± 0.5	11.1 ± 1.4	6 ± 0.6	1.9 ± 0.5	10.3 ± 1.6	7 ± 0.7	4.5 ± 2.4	9.4 ± 2	7.8 ± 1.5	5.4 ± 1.5	6.1 ± 1.4	4.3 ± 0.7	2.3 ± 0.3
∑ PUFA	34.7 ± 5.1	24.5 ± 4.5	15.7 ± 2.5	38.8 ± 4.6	31.1 ± 2.5	13 ± 1.8	38.4 ± 3.5	32.4 ± 2.4	21.1 ± 10.3	37.9 ± 3.4	37.3 ± 3.7	29 ± 4.2	36.5 ± 2.9	38 ± 2.3	33.8 ± 2.6
20:2i	1.1 ± 0.3	1.1 ± 0.2	0.7 ± 0.5	0.9 ± 0.2	1.6 ± 0.6	0 ± 0	0.9 ± 0.2	0.8 ± 0.3	1.4 ± 2.8	0.9 ± 0.2	1 ± 0.3	0.1 ± 0.3	1.1 ± 0.1	1.3 ± 0.5	0.8 ± 0.2
20:2j	0.5 ± 0.1	0.4 ± 0.1	0.2 ± 0.2	0.4 ± 0	0.6 ± 0.1	0 ± 0	0.4 ± 0.1	0.3 ± 0.3	0.2 ± 0.5	0.3 ± 0.1	0.4 ± 0.1	0 ± 0	0.4 ± 0.1	0.4 ± 0.1	0.3 ± 0.1
22:2i	1.9 ± 0.3	2.6 ± 0.5	1.3 ± 0.7	1.6 ± 0.5	3.8 ± 1.5	0.1 ± 0.2	1.7 ± 0.4	1.9 ± 0.6	0.3 ± 0.6	1.5 ± 0.3	1.7 ± 0.8	0.8 ± 0.3	1 ± 0.1	0.7 ± 0.2	0.3 ± 0.2
22:2j	5.4 ± 0.8	6 ± 0.6	3.2 ± 2	5.1 ± 1.1	8.7 ± 3.2	1.4 ± 0.3	4.2 ± 0.7	4.4 ± 1.4	1.8 ± 1.7	4.6 ± 0.8	4.5 ± 1.9	2 ± 0.7	5.3 ± 1	3.6 ± 0.6	1.7 ± 0.9
22:3i	0.5 ± 0.2	1.1 ± 0.3	0.1 ± 0.2	0.5 ± 0.2	1.4 ± 0.6	0 ± 0	0.5 ± 0.2	0 ± 0	0 ± 0	0.5 ± 0.1	0.6 ± 0.3	0 ± 0	0 ± 0	0 ± 0	0 ± 0
∑ NMI FA	9.4 ± 1.4	11.3 ± 1.2	5.4 ± 3.3	8.4 ± 1.8	16 ± 5.6	1.5 ± 0.3	7.8 ± 1.3	7.3 ± 2.4	3.7 ± 3.4	7.8 ± 1.4	8.2 ± 3.1	3 ± 1.1	7.9 ± 1.2	6 ± 0.5	3 ± 1
16:0DMA	0.1 ± 0	0.3 ± 0.2	0 ± 0	0 ± 0	0.1 ± 0.1	0 ± 0	0.2 ± 0.2	0.4 ± 0.2	0.4 ± 0.5	0.5 ± 0.2	0.5 ± 0.2	1.8 ± 2.1	0.5 ± 0.4	3.2 ± 2.3	7.6 ± 1.9
18:0DMA	1 ± 0.5	0 ± 0	0 ± 0	1.2 ± 0.4	0.1 ± 0.1	0 ± 0	1.7 ± 1	4 ± 1.9	13.5 ± 5.8	0.8 ± 0.4	0.8 ± 0.6	2.7 ± 2	0.5 ± 0.1	1.5 ± 0.6	4.9 ± 1.1
20:1n-7DMA	0.6 ± 0.3	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	1.2 ± 0.5	1.8 ± 1	7.1 ± 3	0.2 ± 0.1	0.2 ± 0.2	1.5 ± 0.8	0 ± 0	0.1 ± 0.1	1.2 ± 0.2
∑ DMA FA	1.7 ± 0.7	0.3 ± 0.2	0 ± 0	1.3 ± 0.5	0.2 ± 0.1	0 ± 0	3.1 ± 1.4	6.2 ± 3	20.9 ± 8.9	1.5 ± 0.7	1.4 ± 1	6 ± 1.8	1.1 ± 0.5	4.9 ± 2.9	13.7 ± 3
A	0.3 ± 0.1	0.2 ± 0.2	0.1 ± 0.2	0.2 ± 0	0.4 ± 0.1	0 ± 0	0.2 ± 0	0.3 ± 0.4	0 ± 0	0.2 ± 0.1	0.3 ± 0.1	0.1 ± 0.2	0.1 ± 0.1	0 ± 0	0 ± 0
B	0.2 ± 0	0 ± 0	0 ± 0	0.2 ± 0	0.1 ± 0.1	0 ± 0	0.1 ± 0.1	0 ± 0	0.2 ± 0.3	0 ± 0.1	0.1 ± 0.1	0.2 ± 0.4	0.1 ± 0	0 ± 0	0 ± 0
W	0.3 ± 0.1	0.6 ± 0.1	0.1 ± 0.3	0.3 ± 0.1	0.7 ± 0.2	0 ± 0	0.2 ± 0.2	0.4 ± 0.2	0 ± 0	0.3 ± 0.1	0.3 ± 0.1	0.1 ± 0.2	0.2 ± 0	0.2 ± 0	0.1 ± 0.1
∑ UK FA	0.8 ± 0.1	0.8 ± 0.2	0.2 ± 0.4	0.6 ± 0.1	1.1 ± 0.3	0 ± 0	0.5 ± 0.2	0.7 ± 0.6	0.2 ± 0.3	0.5 ± 0.1	0.6 ± 0.1	0.4 ± 0.4	0.3 ± 0.1	0.2 ± 0.1	0.1 ± 0.1
PUFA/SFA	1.5 ± 0.3	0.7 ± 0.2	0.3 ± 0.1	1.7 ± 0.3	1.3 ± 0.3	0.2 ± 0	1.8 ± 0.2	1.2 ± 0.1	0.8 ± 0.5	1.6 ± 0.2	1.6 ± 0.6	0.9 ± 0.2	1.6 ± 0.2	1.9 ± 0.2	1.7 ± 0.2
EPA/DHA	1.1 ± 0.1	1.2 ± 0.2	1.3 ± 0.1	1.1 ± 0.1	1.4 ± 0.2	1.3 ± 0.1	1.1 ± 0.1	1 ± 0.2	0.8 ± 0.3	1.2 ± 0.2	1.4 ± 0.1	1.1 ± 0.2	2.5 ± 0.5	3.8 ± 0.3	4 ± 0.6
∑ n-3/ ∑ n-6	3.7 ± 1	1.5 ± 0.1	1.4 ± 0.4	4.5 ± 0.6	2.6 ± 0.5	2.8 ± 0.7	4.5 ± 0.5	3.6 ± 0.3	Inf ± NA	4.6 ± 0.9	5.9 ± 0.4	5.5 ± 2.8	5.6 ± 1.1	5.3 ± 1.9	2.2 ± 0.5

878 **Table S5: Stable isotope (mean ± SD, n = 5) composition of the three ontogenic stages of *Crepidula fornicata* (motile males, sessile males, sessile females)**
879 **over the sampling survey.**

	S1 = 26th February			S2 = 21th March			S3 = 28th March			S4 = 12th April			S5 = 14th June		
	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male	Sessile female	Sessile male	Motile male
$\delta^{13}C$	-19.5 ± 0.3	-19.4 ± 0.4	-19.4 ± 0.5	-19.8 ± 0.3	-19.7 ± 0.3	-20.1 ± 0.5	-19.2 ± 0.6	-19.6 ± 0.4	-19.9 ± 0.3	-20.1 ± 0.4	-20.3 ± 0.8	-20.6 ± 0.5	-18.1 ± 0.3	-18.5 ± 0.1	-18.5 ± 0.3
$\delta^{15}N$	9.3 ± 0.4	8.9 ± 0.8	8.9 ± 0.3	9.6 ± 0.5	8.9 ± 0.5	8.4 ± 1.1	9.7 ± 0.3	9.4 ± 0.6	8.5 ± 0.4	8.4 ± 0.7	8.4 ± 0.6	8.3 ± 0.5	8.6 ± 0.6	8.3 ± 0.5	7.5 ± 0.3
C/N	4 ± 0.3	3.8 ± 0.8	4 ± 0.2	3.2 ± 0.8	3.8 ± 0.6	3.5 ± 0.7	3.6 ± 0.4	3.3 ± 0.4	3.8 ± 0.2	3.6 ± 0.4	4.3 ± 0.2	3.8 ± 0.2	3.5 ± 0.6	4.1 ± 1.2	3.4 ± 0.3

882
883

884
Figure S1: Mean daily temperature ($\pm$ SD) at the bottom in our study site
886