Thalamic encoding of lexical status is lateralized during reading aloud

Dengyu Wang[1,2], Witold J. Lipski[1]*, Alan Bush[3,7], Anna Chrabaszcz[4], Christina Dastolfo-Hromack[5], Michael W. Dickey[5], Julie A. Fiez[4,6] & R. Mark Richardson[3,7,8]*

1. Department of Neurological Surgery, University of Pittsburgh School of Medicine, Pittsburgh, PA, 15213, USA
2. School of Medicine, Tsinghua University, Beijing, 100084, China
3. Brain Modulation Lab, Massachusetts General Hospital, Boston, MA, 02114, USA
4. Department of Psychology, University of Pittsburgh, Pittsburgh, PA, 15213, USA
5. Department of Communication Science and Disorders, University of Pittsburgh, Pittsburgh, PA, 15213, USA
6. University of Pittsburgh Brain Institute, Pittsburgh, PA, 15213, USA
7. Harvard Medical School, Boston, MA, 02115, USA

*For correspondence:
lipskiw@upmc.edu
mark.richardson@mgh.harvard.edu

Abstract

To explore whether the thalamus participates in lexical status encoding, local field potentials were recorded in patients undergoing deep brain stimulation lead implantation while they read aloud single-syllable words and nonwords. Bilateral decreases in thalamic beta (12-30Hz) activity were preferentially locked to stimulus presentation, and these decreases were greater when nonwords were read. Increased broadband gamma (70-150Hz) activity also was locked preferentially to speech onset bilaterally, but greater nonword-related increases in this activity were observed only on the left, demonstrating lateralization of thalamic gamma selectivity for lexical status. In addition, this lexical status effect was strongest in more anterior thalamic locations, regions which are more likely to receive pallidal than cerebellar afferents. These results provide evidence from intracranial thalamic recordings for the lateralization and topography of subcortical lexical status processing.

Introduction

Reading words aloud depends on our ability to transform information about letter combinations into plans for producing speech sounds. Determining whether a group of letters represents a word is one important component of this process. While word reading can be supported by processes that permit mapping entire word forms to lexical representations, nonword reading depends upon sublexical processes that map spelling and sound (Coltheart et al., 2001). Functional neuroimaging has allowed for increasingly detailed study of the cortical regions that participate in these phonological processes, for instance demonstrating that a region of the inferior frontal gyrus (Brodmann's areas 44 and 45) is significantly more active for nonword reading than for word reading (Fiez et al., 1999; Hagoort et al., 1999; Heim et al., 2005; Herbster et al., 1997). The role of subcortical regions in spoken word production remains elusive, however, due to the low resolution of neuroimaging techniques for measuring subcortical activity. Resolving this knowledge gap is important, given that cortical activity is modulated by thalamic outflow through basal ganglia-thalamo-cortical and cerebello-thalamo-cortical circuits (Behrens et al., 2003; Bosch-Bouju et al., 2013; Hwang et al., 2017; Zhang et al., 2010). Despite the cortico-centric focus of most experimental work and accompanying models, there is increasing recognition of the role of subcortical processes in language production (Llano, 2015).

Neurosurgical procedures involving invasive recording and stimulation in epilepsy patients undergoing electrode implantation for seizure mapping traditionally have provided the only direct means to test hypotheses related to cortical function during reading (Juphard et al., 2011). Epilepsy surgery, however, rarely provides access to the thalamus and basal ganglia. Movement disorders surgery, on the other hand, routinely provides direct access to the thalamus and basal ganglia in awake patients. Regionalization of language function within the left thalamus was established in surgery for movement disorders in the 1970s, in seminal studies by Ojemann and colleagues that employed electrical stimulation principles borrowed from traditional cortical language mapping protocols (Johnson & Ojemann, 2000). More recently, event-related potential recordings in patients undergoing deep brain stimulation (DBS) have suggested that thalamic structures are engaged in the analysis of syntactic, semantic and lexical information during acoustically presented language tasks (Tiedt et al., 2017; Wahl et al., 2008).

We recently developed a protocol to study subcortical activity during single-syllable

word/nonword reading, in patients undergoing DBS lead implantation (Chrabaszcz et al., 2019;

Lipski et al., 2018). Here, we explore whether the thalamus participates in the encoding of

lexical status, by recording local field potentials (LFP) in patients undergoing DBS lead

implantation for essential tremor targeting the ventral intermediate nucleus of the thalamus

(Vim, which corresponds to the ventral portion of the ventral lateral posterior nucleus (VLp)

(Macchi & Jones, 1997)). Subjects performed a reading aloud task where they were asked to

read aloud single-syllable words or nonwords that appeared on a computer screen (Figure 1).

We assessed thalamic participation in lexical status encoding by comparing task-related neural

responses when participants spoke nonwords vs. words.

Figure 1: Schematic of experiment. ITI: inter-trial interval.

Results

Eleven subjects read aloud single-syllable words alternating with nonwords, intraoperatively

during implantation of DBS leads targeting the Vim nucleus of the thalamus. LFP recordings in

four subjects were obtained unilaterally (3 left, 1 right) during microelectrode mapping, where

each of these subjects performed up to four sessions of 120 trials. Recordings from seven

subjects were obtained from DBS lead contacts, where each of these subjects performed two

sessions: in the first session unilateral recordings were made from the left thalamus, and in the

second session bilateral recordings were preformed simultaneously. A total of 117 recordings

(data recorded in one location in one session) from 89 recording sites pooled across subjects

were collected. Recording locations were determined in MNI (Montreal Neurological Institute)

space (Figure 2) and comprised locations within or bordering (within 1mm) the ventral anterior

nucleus (VA) and the ventral lateral anterior nucleus (VLa) (38/89), or the VLp (51/89).

a **b**

Figure 2: Localization and MNI-transformation of recording sites. a Sagittal view of the recording locations of all the subjects relative to the thalamus (yellow), VA and VLa (green), VLp (blue), and Vim (orange), superimposed on a T2-weighted image. **b** A plot of all the recording sites in MNI space with their MNI-defined coordinates. Recording contacts of different subjects are color-coded. In both **a** and **b**, right hemisphere recording locations are flipped to the left hemisphere: left side contacts are marked with circles, and right side contacts are marked with triangles.

Behavioral performance

Across all subjects, the mean speech production latency (interval between stimulus presentation and onset of speech) was 1.07 ±0.29s, and the mean duration of speech was 0.70 ±0.09s. Nonword production latency (1.23 ±0.37s) was significantly longer than word production latency (0.98 ±0.24s) across subjects (two-tailed paired t-test, t(10)=4.47, *P*=0.0012; Figure 3a). Nonword production duration across subjects (0.68 ±0.07s) was not significantly different from word production duration (0.72 ±0.13s; two-tailed paired t-test, t(10)=1.99, *P*=0.074; Figure 3b).

a **b**

Figure 3: Behavioral outcomes. a Mean and SEM of speech production latency for each
subject across all trials (first column), word trials (second column), or nonword trials (third
column), superimposed with the mean and SEM of speech production latency across subjects.
b Mean and SEM of speech duration for each subject across all trials (first column), word trials
(second column), or nonword trials (third column), superimposed with the mean and SEM of
speech duration across subjects. Two-tailed paired t-test, **P<0.01, ns: not significant.
83

Thalamic neural activity is modulated during reading aloud

Thalamic LFP activity exhibited significant time-frequency modulation during the reading aloud

task (Figure 4). Compared to baseline (a period of 1000ms preceding stimulus presentation),

there was a significant decrease in spectral power in the beta frequency band (12-30Hz) that

occurred at stimulus presentation and lasted until the end of speech (-1.08-0.62s relative to

speech onset, two-tailed Wilcoxon signed-rank test, n=117, *P*<0.05, Bonferroni corrected). In

contrast, a significant increase in broadband gamma (70-150Hz) activity occurred shortly before

the onset of speech and persisted throughout the utterance (-0.15-0.59s relative to speech

onset, two-tailed Wilcoxon signed-rank test, n=117, *P*<0.05, Bonferroni corrected). Average *z*-

scored task-related beta and broadband gamma response amplitudes of each trial were then

calculated over the respective significant time windows for all the recordings. As a result, 66/117

(56%) of the recordings showed significant beta activity decreases during the task compared to

baseline (one-tailed one-sample t-test, *P*<0.05, Bonferroni corrected), and significant task-

related broadband gamma activity increases were observed in 91/117 (78%) of the recordings

(one-tailed one-sample t-test, *P*<0.05, Bonferroni corrected).

Figure 4: Thalamic neural activity shows task-related modulations. A spectrogram of thalamic neural activity during the reading aloud task, averaged across all trials and all recordings. Trials are aligned to speech onset. Significant changes compared to baseline are marked in white contours (-1.08-0.62s for beta activity, and -0.15-0.59s for broadband gamma activity; Wilcoxon signed-rank test, n=117, *P*<0.05, Bonferroni corrected). Average time points of stimulus presentation, speech onset and offset are marked with black dashed lines.

Beta and broadband gamma responses differ in timing properties

The average interval between stimulus presentation and onset of a significant change from baseline in the spectral power of a particular frequency band, i.e. the mean band response latency, was shorter for significant beta decrease responses than for significant broadband gamma increase responses (0.79 ±0.18s vs. 0.99 ±0.18s, two-tailed two-sample t-test, t(155)=-6.97, *P*<10^{-5}). The mean band response onset to speech onset interval was also greater for significant beta decrease responses than for significant broadband gamma increase responses (0.31 ±0.15s vs. 0.13 ±0.14s, two-tailed two-sample t-test, t(155)=7.67, *P*<10^{-5}). To characterize the temporal properties of these responses, we examined their trial-to-trial relationships to stimulus presentation versus speech onset (Figure 5a-d). Of the 66 recordings that showed significant task-related beta power decreases, 43 (65.2%) had beta responses time-locked to stimulus presentation, whereas only 19 (28.8%) had beta responses time-locked to speech onset (Figure 5e). In contrast, the majority (70/91, 77.0%) of significant broadband gamma power increases were time-locked to speech onset, with a minority (19/91, 21.0%) time-locked to stimulus presentation (Figure 5f). These relationships were dissociated (χ^2 test, α=0.05; Supplementary Table 1), with beta decreases more likely to be stimulus-locked (χ^2(1)=31.4, *P*<10^{-5}) and broadband gamma increases more likely to be speech onset-locked (χ^2(1)=36.1, *P*<10^{-5}). These temporal correlations did not differ between recording sides (χ^2 test, α=0.05; Supplementary Table 2).

Figure 5: Beta decrease response is locked to stimulus presentation while broadband gamma increase response is locked to speech onset. a, **b** Raster plots for beta band responses (**a**) and broadband gamma responses (**b**) across trials of two representative recordings. Trials are aligned to stimulus presentation (indicated with black dashed lines) and sorted by speech production latency. Speech onsets are denoted with bold black lines. Onsets of significant beta activity decreases are marked with red asterisks in **a** and onsets of significant

broadband gamma activity increases are marked with blue asterisks in **b**. **c**, **d** Band response

onset to speech onset interval (red asterisks) and band response latency (blue asterisks) are

correlated (Pearson's correlation, α=0.05) with speech production latency respectively, for the

two representative recordings. **e**, **f** The same correlation analysis is performed for all the

recordings with significant beta decrease responses and all the recordings with significant

broadband gamma increase responses, and the results are summarized in **e** and **f**, respectively.

Recordings locked to stimulus presentation are shown in red, recordings locked to speech onset

are shown in blue, and recordings not locked to either stimulus presentation or speech onset

are shown in gray.

Thalamic beta activity is selective to lexical status

To investigate the involvement and lateralization of the thalamus in lexical processing, only

recordings that were from subjects with bilateral lead recordings (n=7) and that showed

significant task-related modulation were included for lexicality-related analyses. As a result, 55

recordings (21 unilateral session left-side recordings, 34 from bilateral session left-side (20) and

right-side (14) recordings) were included for beta lexical selectivity analysis. Nonword

production was associated with a greater suppression of beta power compared with reading

words. These differential beta responses were observed in both hemispheres (Figure 6a, c). In

the left thalamus, significant word vs. nonword beta amplitude differences occurred between

0.8s and 1.8s after stimulus presentation (two-tailed paired t-test, n=41, *P*<0.05, Bonferroni

corrected). Similarly, significant beta amplitude differences occurred in the right thalamus at 0.2-

1.7s relative to stimulus presentation (two-tailed paired t-test, n=14, *P*<0.05, Bonferroni

corrected).

To quantify the lexical selectivity of thalamic beta activity, a lexical selectivity index was

determined for each of these recordings. We utilized the time window of significant task-related

beta activity decrease (-1.08-0.62s relative to speech onset (see Figure 4)) to calculate a mean

task-related beta response power value for each trial, and then the values were compared

between nonword trials vs. word trials in each recording by performing a two-sample t-test. The

resulting t-statistic was the beta lexical selectivity index for this recording: a beta lexical

selectivity index less than zero indicated stronger beta activity decrease (thus having more

negative z-scored power) during nonword trials than during word trials, and vice versa. The

mean beta lexical selectivity indexes for unilateral session left-side recordings (-0.49 ±0.76),

bilateral session left-side (-0.52 ±0.82) and right-side (-0.82 ±0.89) recordings were all

significantly less than zero (two-tailed one-sample t-test, t(20)=-2.97, *P*=0.0075; t(19)=-2.83,

P=0.011; t(13)=-3.48, *P*=0.0041; Figure 6e), suggesting that the magnitude of the beta

decrease was significantly nonword-selective in each case. There were no significant

differences in beta lexical selectivity between recordings in the left and right thalamus, or

between recordings from the unilateral and bilateral sessions on the left side (two-tailed two-

sample t-test, unilateral session left-side recordings vs. bilateral session right-side recordings:

t(33)=1.18, *P*=0.25; bilateral session left-side recordings vs. bilateral session right-side

recordings: t(32)=1.05, *P*=0.30; unilateral session left-side recordings vs. bilateral session left-

side recordings: t(39)=0.093, *P*=0.93).

Lexical selectivity of thalamic broadband gamma activity is lateralized

74 recordings (26 unilateral session left-side recordings, 22 bilateral session left-side

recordings, and 26 bilateral session right-side recordings) were included for broadband gamma

lexical selectivity analysis. Significantly greater broadband gamma activity increases during

nonword production were observed in the left thalamus, starting 0.1s after speech onset and

persisting throughout the following 0.4s (two-tailed paired t-test, n=48, *P*<0.05, Bonferroni

corrected; Figure 6b). In the right thalamus, however, the broadband gamma response curves

observed during nonword and word reading were similar, without a significant difference in

response amplitudes (two-tailed paired t-test, n=26, α=0.05, Bonferroni corrected; Figure 6d).

We next calculated broadband gamma lexical selectivity indexes, using the significant time

window of -0.15-0.59s relative to speech onset (see Figure 4). A broadband gamma lexical

selectivity index greater than zero indicated stronger broadband gamma activity increase during

nonword trials than during word trials, and vice versa. The mean broadband gamma lexical

selectivity indexes of both unilateral session left-side recordings (0.53 ±0.96) and bilateral

session left-side recordings (0.46 ±1.00) were significantly greater than zero (two-tailed one-

sample t-test, t(25)=2.80, *P*=0.0097; t(21)=2.14, *P*=0.044), indicating significant correlation of

the magnitude of the gamma response to lexical status (Figure 6f). In contrast, broadband

gamma responses in right side recordings did not show significant lexical selectivity (-0.22

±0.68, two-tailed one-sample t-test, t(25)=-1.67, *P*=0.11). The differences in broadband gamma

lexical selectivity between the left and right thalamus were demonstrated with two-tailed two-

sample t-test (unilateral session left-side recordings vs. bilateral session right-side recordings:

t(50)=3.25, *P*=0.0021; bilateral session left-side recordings vs. bilateral session right-side

recordings: t(46)=2.79, *P*=0.0077), further suggesting that selectivity of thalamic broadband

gamma activity to lexical status is lateralized to the left. Recordings in the unilateral and bilateral

sessions on the left side did not differ in broadband gamma lexical selectivity (two-tailed two-

sample t-test, t(46)=0.25, *P*=0.80), indicating the consistency of broadband gamma lexical selectivity between task sessions.

Figure 6: Differential band responses during word vs. nonword reading aloud in the left and right thalamus. a-d Comparisons of average band response amplitudes during word (blue) vs. nonword (red) reading aloud for beta band (**a**, **c**) and broadband gamma (**b**, **d**), in the left (**a**, **b**) and right thalamus (**c**, **d**). Band responses are averaged across trials of respective trial types and across recordings that showed significant task-related band responses in each side, aligned

to stimulus presentation for beta band (**a**, **c**) and speech onset for broadband gamma (**b**, **d**).
Average time points of speech onset and end of speech are marked with dashed lines in **a** and
c, and average time points of stimulus presentation and end of speech are marked with dashed
lines in **b** and **d**, for word trials (blue) and nonword trials (red) respectively. Black bars indicate
significant differences of response amplitudes between word and nonword trials (two-tailed
paired t-test, *P*<0.05, Bonferroni corrected). Standard errors are shaded in light colors. **e** Dot
plot of beta lexical selectivity indexes of lead recordings that showed significant task-related
beta decrease responses, grouped by recording sides and recording sessions (unilateral
session left-side recordings in the first column, bilateral session left-side recordings in the
second column, and bilateral session right-side recordings in the third column). Mean and SEM
of beta lexical selectivity index across recordings of each group is superimposed on each
column, respectively. **f** Dot plot of broadband gamma lexical selectivity indexes of lead
recordings that had significant task-related broadband gamma increase responses, grouped by
recording sides and recording sessions (unilateral session left-side recordings in the first
column, bilateral session left-side recordings in the second column, and bilateral session right-
side recordings in the third column). Mean and SEM of broadband gamma lexical selectivity
index across recordings of each group is superimposed on each column, respectively. Two-
tailed one-sample t-test, †*P*<0.05, ††*P*<0.01; two-tailed two-sample t-test, *******P*<0.01, ns: not
significant.
231
Left side broadband gamma lexical selectivity is topographically organized
We observed that the recording sites in the left thalamus that were significantly nonword-
selective in terms of broadband gamma response activity (broadband gamma lexical selectivity
index>1.645 based on normal approximation of t-distribution) appeared to comprise more
anterior locations in MNI space (Figure 7a). Pearson's correlation tests demonstrated that the
broadband gamma lexical selectivity index significantly correlated with recording location along
the anterior-posterior axis (MNI-defined y coordinate; n=48, ρ=0.49, *P*=0.00041; Figure 7b) and
the ventral-dorsal axis (MNI-defined z coordinate; n=48, ρ=0.30, *P*=0.041; Supplementary
Figure 1a), but not the lateral-medial axis (MNI-defined x coordinate; n=48, ρ=-0.14, *P*=0.34;
Supplementary Figure 1b) within the left thalamus. To avoid multicollinearity and to test out
possible interactions between variables, a stepwise linear regression model was applied, which
determined that the anterior-posterior location of the recording was the only significant predictor
(estimated coefficient=0.17, SE=0.044, *P*=0.00041), while neither the ventral-dorsal location
(*P*=0.72) nor the interaction between ventral-dorsal location and anterior-posterior location
(*P*=0.14) explained the result. In order to control for subject differences and session differences,
we further fitted linear mixed effects models to the data, entering subject and session as random
effects. The results indicate that even after accounting for subject and session variability, the
broadband gamma lexical selectivity index in the left thalamus had a significant gradient along

the anterior-posterior axis, with greater broadband gamma lexical selectivity index more likely to

be observed anteriorly (estimated coefficient=0.17, SE=0.043, *P*=0.00032). Based on the linear

mixed effects modeling results, neither ventral-dorsal location (estimated coefficient=0.061,

SE=0.035, *P*=0.087) nor lateral-medial location (estimated coefficient=-0.11, SE=0.093, *P*=0.24)

of the recording had significant effect on broadband gamma lexical selectivity in the left

thalamus. Taken together, these results suggest that broadband gamma lexical selectivity is

dependent on the anterior-posterior location of the recording in the left ventral lateral thalamus,

with greater nonword selectivity more likely to occur anteriorly.

Figure 7: Broadband gamma lexical selectivity depends on anterior-posterior location of
the recording in the left thalamus. a Left side lead recording sites are plotted together with
anatomical structures (thalamus in yellow, VA and VLa in green, VLp in blue, and Vim in
orange), viewed from a lateral direction. Recording sites where significantly nonword-selective
recordings in terms of broadband gamma response amplitudes (broadband gamma lexical
selectivity index>1.645 based on normal approximation of t-distribution) were observed in either
session are filled with black, and remaining recording sites are shown in open circles. **b**
Broadband gamma lexical selectivity indexes of left side lead recordings that showed significant
task-related broadband gamma responses are correlated (Pearson's correlation, α=0.05) with
recording locations along the anterior-posterior axis (y coordinates in MNI space). Recordings
inside VA or VLa are green-colored and recordings inside VLp are blue-colored.

No significant correlation (Pearson's correlation, α=0.05) between broadband gamma lexical

selectivity index and recording location was observed for right side recordings (Supplementary

Figure 2 and 3), or for beta lexical selectivity on either side (Supplementary Figure 4 and 5), by

any statistical modeling. These results suggest that the topography of lexical selectivity is

unique to broadband gamma responses in the left thalamus.

Discussion

Subcortical contributions to language production have been hypothesized largely from correlations of focal brain damage with aphasic syndromes and from language studies using functional magnetic resonance imaging (fMRI) or scalp electroencephalography (EEG) (Hebb & Ojemann, 2013). Our results are the first to demonstrate thalamic neural activity during reading aloud. We discovered that the encoding of lexical status is differentially represented in thalamic neural activity. Whereas greater beta activity decrease occurred during nonword trials as compared to word trials in both hemispheres, greater gamma increases in the left but not right thalamus were associated with the spoken production of nonwords as compared to words. Importantly, the level of broadband gamma lexical selectivity was greater in more anterior thalamic recording locations, regions more likely to receive basal ganglia than cerebellar afferents (Alexander et al., 1986).

We first observed that thalamic beta activity showed a task-related decrease, locked to stimulus presentation, consistent with the expected beta desynchronization that accompanies motor action. Beta oscillations are proposed to signal the maintenance of current sensorimotor and cognitive states (Engel & Fries, 2010), and event-related beta activity decreases during the preparation and the execution of voluntary movements have been observed across relevant brain regions, including the thalamus (Klostermann et al., 2007; Kühn et al., 2004; Paradiso, 2004; Tzagarakis et al., 2010). Beta activity was reduced across the entirety of each task event, which requires a series of underlying neural dynamics (processing the visual stimulus, retrieval of lexical information, encoding of the phonological information, and execution of articulatory movement), suggesting a state change that facilitated the process of speech production. Significant increases in broadband gamma activity began before but were predominantly locked to speech onset. Given that broadband gamma activity is thought to index synchronized local neuronal firing (Ray et al., 2008; Ray & Maunsell, 2011) and is associated with the functional activation of relevant brain regions during a wide range of sensorimotor and cognitive tasks, including speech and language processing (Bouchard et al., 2013; Chrabaszcz et al., 2019; Crone et al., 2006; Juphard et al., 2011; Uhlhaas et al., 2011), our data suggest that the ventral lateral thalamus tracks speech production.

Studies of the brain networks involved in lexical processing during reading primarily have been confined to cortical regions (Dietz et al., 2005; Fiez et al., 1999; Hagoort et al., 1999; Heim et al., 2005; Herbster et al., 1997; Juphard et al., 2011; Mechelli et al., 2003; Xu, 2001). In this

study, we utilized the balanced design of words and nonwords in the stimulus sets and
simultaneous bilateral recordings in the seven participants to determine whether lexical
processing is differentially represented in the thalamus during reading aloud. Thalamic
broadband gamma oscillations showed significantly stronger activation during nonword reading
aloud than during word reading aloud, which evolved around speech onset and was present
throughout the entire utterance. Notably, this broadband gamma lexical selectivity was found
only in recordings from the left side. According to classic speech production models (Levelt et
al., 1999), the time course of the observed broadband gamma activity difference between word
and nonword trials corresponded to the articulatory stage of reading aloud. It is unlikely that this
lexicality effect was motoric, considering that the motor complexity was balanced between word
and nonword stimuli, and that this differential broadband gamma modulation was not
equivalently observed in the right hemisphere. In fact, the dual-route theory of reading aloud has
suggested that words and nonwords are read aloud differently: while real words can be read
aloud via either grapheme-to-phoneme conversions or direct word-to-sound mapping, nonwords
can only be read aloud via grapheme-to-phoneme conversions (Coltheart et al., 2001). It is
possible that phonological encoding was still ongoing during nonword production, after a failed
internal lexicon lookup procedure. Indeed, stronger gamma (50-150Hz) responses for pseudo-
words than for words have been reported in Broca's area (Brodmann's areas 44 and 45) during
silent word reading, where the length of this differential response increased with the length of
the stimuli (Juphard et al., 2011). In addition, it is possible that reading aloud nonwords created
a learning or error correction demand (Hickok, 2014). We propose that the lexicality effect on
thalamic broadband gamma activity during reading may reflect left thalamic participation in
phonological encoding, learning, and feedback monitoring during speech and language
processing. Although previous studies have supported a lateralized thalamic role in language
(Johnson & Ojemann, 2000), our results from simultaneous bilateral direct recordings in the
thalamus are the first to provide direct supporting evidence.
337
A topography for broadband gamma lexical selectivity was observed in the left thalamus. The
lexical selectivity of the broadband gamma response was significantly correlated with the
anterior-posterior location of recording sites in the left thalamus, with a higher lexical selectivity
more likely to appear anteriorly. In contrast, a gradient for broadband gamma lexical selectivity
was not observed in the right thalamus. This finding further supports a unique language role
lateralized to the left thalamus, and suggests functional heterogeneity in the left ventral lateral
thalamus during speech and language processing. The anterior portion (VA and VLa) receives

input primarily from the internal globus pallidus and substantia nigra pars reticulata, and has

strong connections with frontal cortex, including Broca's area (Alexander et al., 1986; Behrens

et al., 2003; Bosch-Bouju et al., 2013; Hintzen et al., 2018; Hwang et al., 2017; Zhang et al.,

2010). Broca's area has consistently been associated with lexico-phonological processing (Dietz

et al., 2005; Fiez et al., 1999; Hagoort et al., 1999; Heim et al., 2005; Herbster et al., 1997;

Juphard et al., 2011; Mechelli et al., 2003; Xu, 2001). Thus, this thalamic region may participate

in differential lexical processing during nonword vs. word reading aloud in concert with Broca's

area. In contrast, the posterior region (VLp), which receives cerebellar projections and

preferentially sends output to primary motor cortex (Behrens et al., 2003; Bosch-Bouju et al.,

2013; Hintzen et al., 2018; Hwang et al., 2017; Zhang et al., 2010), might be more related to

motor control of a selected motor plan. This idea is supported by previous stimulation studies

that have reported location-dependent effects of thalamic stimulation on speech and language:

stimulation of VLa could cause acceleration of language processes, while stimulation of VLp

often affected motor aspects of speech, such as perseveration and stuttering speech (Hebb &

Ojemann, 2013). We note that VA has been proposed to participate in selection of a language

unit during speech production, via basal ganglia-thalamo-cortical loop interactions (Crosson,

2013); it also is included in the "planning loop" in the GODIVA model of speech production

(Bohland et al., 2010).

In contrast, although significantly stronger nonword-related than word-related beta activity

decreases accompanied the reading aloud task bilaterally, neither lateralization nor topography

was observed for thalamic beta lexical selectivity. Therefore, beta activity decreases likely

represent nonspecific changes of cognitive and sensorimotor states that prepare the entire

thalamo-cortical network for a behavioral change. Note that in the current work we did not try to

compare task-related band response strength between different recording locations and

different recording sides among subjects and sessions, because of a number of uncontrollable

factors that affect the band oscillatory power (e.g. recording impedances and baseline neural

activity are variable across subjects, sessions, and recording sides). In contrast, the lexical

selectivity index, which was calculated by comparing word-related and nonword-related band

response strength within each recording, is mostly independent of those factors and thus

comparable at the group level. This idea is supported by the fact that lexical selectivity indexes

in both beta and broadband gamma bands remained consistent across sessions.

In summary, our results are the first demonstration of time-frequency modulations of thalamic
neural activity during reading aloud. These data suggest that lateralized and topographically
organized thalamic pathways participate in speech production differently, based on whether a
word or nonword is being read.
382

Materials and methods

Subjects
Eleven human subjects (3 females, 68.4 ±8.0 years) with essential tremor undergoing awake
DBS implantation surgery targeting the Vim nucleus of the thalamus were studied. All
participants were right-handed native English speakers. None had significant cognitive
impairment based on a detailed neuropsychological evaluation performed during clinical
evaluation for DBS surgery. All but one underwent bilateral (left side first) DBS lead implantation
(one subject had one lead implanted in the left hemisphere previously and underwent right side
lead implantation in the current study). Full demographic description of subjects is provided in
Table 1. All protocols were approved by the Institutional Review Board of the University of
Pittsburgh (IRB Protocol #PRO13110420), and all participants gave written informed consent.

Table 1: Subject characteristics.

Subject	Age	Gender	Handedness	MMSE	LFP recording side	LFP recording type	Number of recordings	Mean number of trials per session	Trials rejected, %
1	61	Male	Right	NR	Left	Mapping electrodes	12	111	7
2	70	Female	Right	30/30	Right	Mapping electrodes	6	107	11
3	66	Male	Right	NR	Left	Mapping electrodes	6	116	3
4	75	Male	Right	26/30	Left	Mapping electrodes	9	112	5
5	64	Male	Right	30/30	Both	DBS leads	12	104	13
6	53	Male	Right	NR	Both	DBS leads	12	113	5
7	67	Male	Right	26/30	Both	DBS leads	12	111	7
8	71	Male	Right	26/30	Both	DBS leads	12	108	9
9	84	Female	Right	30/30	Both	DBS leads	12	118	2
10	73	Female	Right	30/30	Both	DBS leads	12	109	9
11	68	Male	Right	28/30	Both	DBS leads	12	114	5

MMSE: Mini-Mental State Examination; NR: not recorded.

Stimuli and experimental paradigm

Subjects performed a reading aloud task either in the course of subcortical mapping procedure during the surgery (4/11 subjects with mapping electrode recordings) or after the placement of DBS lead in each side (7/11 subjects with lead electrode recordings). For the 4 subjects with mapping electrode recordings, each subject performed up to 4 task sessions, while the 7 subjects with lead electrode recordings each performed 2 task sessions (the first session occurred after left lead was implanted, and the second session took place after bilateral implantation was completed). Each session included 120 trials. The stimuli were consonant-vowel-consonant (CVC) words and nonwords that were presented on a computer screen. Four lists of 120 stimuli (Supplementary Table 3) were constructed based on a previous work (Moore et al., 2017). The first 60 stimuli of each list alternated between unique words and nonwords, and those words and nonwords were balanced along a number of psycholinguistic features, including phoneme probability, phonological neighborhood density, bigram frequency, and

biphone probability (see Supplementary Table 4 for the results of statistical comparisons (two-
tailed two-sample t-test, n=60, α=0.05) between the two conditions in terms of these
psycholinguistic parameters, which were calculated using the English Lexicon Project database
(Balota et al., 2007)). The nonwords were duplicated twice to construct the last 60 stimuli of
each list. One of the four stimulus lists was presented to the subjects during each task session.
For lexicality-related analyses, only the first 60 trials of each session were used.
416
The experimental paradigm was programmed using MATLAB (MathWorks, Natick, MA) and
Psychophysics Toolbox extensions (Brainard, 1997). A schematic of the experiment is shown in
Figure 1. Before each trial, a white fixation cross was presented on the screen. Each trial was
initiated manually by the experimenter, with the appearance of a green fixation cross on the
screen. The green fixation cross lasted 250ms, and was followed by a variable time delay (500-
1000ms) during which the screen remained black. Then the CVC syllable stimulus was
presented in white on the screen, and subjects were instructed to read it aloud. The text
remained on the screen until the subjects finished speaking. A white fixation cross was
presented on the screen during the inter-trial interval (ITI).
426

Electrophysiological recordings

For the 4/11 subjects where physiological subcortical mapping was administered, LFP
recordings were performed using the Neuro-Omega recording system (Alpha Omega, Nazareth,
Israel) and mapping electrodes that have a stainless steel macroelectrode ring (0.55mm in
diameter, 1.4mm in length) 3mm above the tip, while in the other 7/11 subjects LFP signal was
recorded from Medtronic Model 3387 DBS leads (Medtronic, Minneapolis, MN) with four
platinum-iridium electrodes (1.27mm in diameter, 1.5mm in length) that are spaced 1.5mm
apart, using Grapevine Neural Interface Processor (Ripple LLC, Salt Lake City, UT). The
mapping electrodes and the DBS leads targeted the Vim. For subjects with mapping electrode
recordings, three mapping electrodes were placed in three trajectories (anterior, central, and
posterior or central, posterior, and medial) of a standard cross-shaped Ben-Gun array with a
2mm center-to-center spacing, and made simultaneous recordings starting at 15mm above the
surgical target with manual advance in 0.1mm steps. The reading aloud task was carried out in
pauses during subcortical mapping procedure and subjects performed in up to four recording
sessions, with each session corresponding to a different recording depth. Mapping electrode
recordings were performed only during the left side implantation except for one subject who
underwent unilateral implantation in the right side, and thus got mapping electrode recordings

only in the right side. For subjects with DBS lead electrode recordings, the task was

administered in two recording sessions, one after the implantation of the DBS lead in the left

side, receiving recordings from only the left DBS lead electrodes, and the other after bilateral

DBS lead implantation was completed, receiving simultaneous recordings from bilateral lead

electrodes. The LFP signal recorded from mapping electrodes was sampled at 44kHz and

bandpass filtered from 0.075Hz to 10kHz, and data recorded from DBS lead electrodes were

collected at 30kHz. Signal collected in one recording site in one session counted as one

recording. Subject recording characteristics are summarized in Table 1.

Audio recordings

Subjects' speech signal was recorded using an omnidirectional microphone (Audio-Technica

model ATR3350iS Mic, frequency response 50-18,000Hz (Audio-Technica, Machida, Japan) for

6 subjects, and PreSonus model PRM1 Precision Flat Frequency Mic, frequency response 20-

20,000Hz (PreSonus, Baton Rouge, LA) for 5 subjects) placed approximately 8cm away from

the subject's mouth and oriented at an angle of about 45°. The audio signal was collected by

Grapevine Neural Interface Processor at a sampling frequency of 30kHz. For subjects with

mapping electrode recordings, the audio signal was then synchronized with neural signal

recorded by the Neuro-Omega system using digital pulses delivered to both recording systems

via a USB data acquisition unit (model USB-1208FS, Measurement Computing, Norton, MA).

Electrode localization

DBS lead electrodes and mapping electrodes were localized using LEAD-DBS toolbox (Horn et

al., 2019; Horn & Kühn, 2015). Postoperative brain scans were coregistered to preoperative

brain scans using open-source Advanced Normalization Tools (ANTs). Pre- and postoperative

acquisitions were then normalized into MNI ICBM152 NLIN 2009b stereotactic space (Fonov et

al., 2011). Both coregistration and normalization results underwent manual quality check. Semi-

automatic reconstruction of electrodes in MNI space was performed in LEAD-DBS and MNI-

defined coordinates were determined for all the electrode contacts (Figure 2). A digitized and

normalized to MNI space version of the Ewert atlas (Ewert et al., 2018) was used to categorize

the electrode contacts. A contact was assigned to a nucleus if it was within or in the vicinity of

the nucleus (1mm cut-off) based on the minimum Euclidean distance between the contact and

the voxels of the nucleus.

Data pre-processing

The audio signal was segmented into trials and the speech sound was coded by communication
science students trained in phonetics in a custom-designed graphical user interface
implemented in MATLAB. The coding results were manually checked by a speech-language
pathologist. For each trial, (1) the onset of speech was identified, (2) the end of speech was
identified, and (3) the speech content was identified. Trials were considered to have correct
speech responses and were included in further analyses if they met all the following criteria: (1)
the subject's speech response could be clearly identified by the coder, (2) the subject's
response was a CVC syllable consisting of the targeted phonemes, and (3) the response did not
make nonword into a word or word into a nonword.

Electrophysiological data were pre-processed using custom code based on FieldTrip toolbox
(Oostenveld et al., 2011) in MATLAB. The data were resampled at 1kHz and band-pass filtered
from 2 to 400Hz. The data were also notch-filtered at 60Hz and its' harmonics to remove line
noise. Time series data from all recording sites were visually and quantitatively inspected for
quality control. The data were then segmented into trials, each spanning 2s before stimulus
presentation to 2s after the end of speech. Trials with artefacts or excessive noise were
identified both manually and quantitatively, and were excluded for subsequent analyses.
Combined with trials that did not meet the criteria for correct speech responses, an average 6.9
±3.4% of trials per subject were rejected (Table 1). The remaining data were common-average
referenced to minimize noise. For spectral-temporal analysis, the data were decomposed using
Morlet wavelet transformation (width=7) over frequencies of 2Hz to 200Hz in increment steps of
2Hz. For band activity analyses, instantaneous analytic amplitudes of beta and broadband
gamma frequency bands were extracted from respective bandpass filters using Hilbert
transform (MATLAB function *hilbert*). The resulting signal of each trial was z-scored relative to
the baseline (a period of 1000ms preceding stimulus presentation).

Task-related responses
Time-frequency data were averaged across all trials centering on speech onset (from 2s before
speech onset to 2s after speech onset), z-scored to baseline, and then averaged across all
recordings. Nonparametric two-tailed Wilcoxon signed-rank test was performed to determine
significant time-frequency modulations compared to baseline (n=117, α=0.05, Bonferroni
corrected). Significant time windows for beta and broadband gamma determined by the
statistical results were used to calculate a mean beta response strength and mean broadband
gamma response strength for each trial in each recording. Then one-tailed one-sample t-test

was performed on each recording to determine recordings that had significant beta activity

decreases and recordings that had significant broadband gamma activity increases, respectively

(α=0.05, Bonferroni corrected).

Locking analysis

In an effort to characterize the timing properties of beta decrease response and broadband

gamma increase response for recordings with significant task-related changes in either band

activity, we examined the trial-to-trial relationships of significant band response onsets to

stimulus presentation versus speech onset in these recordings. First, beta or broadband gamma

time series data of each trial were smoothed using a moving average kernel of 200ms (MATLAB

function *smoothdata*) and z-scored to baseline, in order to minimize single-trial noise. For

recordings with significant beta decrease responses, a thresholding method with a critical value

of z=-1.645 was applied to determine the onset of beta response for each trial. Specifically, for

each trial, the period with band response power below the threshold that gave minimum

summed activity was considered as activation period and the beginning of the period was

determined as onset of beta response. For recordings with significant broadband gamma

increase responses, the onset of broadband gamma increase was determined in a similar way,

except that a critical value of z=1.645 was used and period of maximum summed activity above

the threshold was considered as activation period. For each band, a trial of one recording was

discarded for locking analysis if no beyond-threshold period was present throughout the trial.

Two intervals were calculated for each trial: (1) the interval between stimulus presentation and

the onset of significant band response (band response latency), and (2) the interval between the

onset of significant band response and the onset of speech. Then the two intervals were

correlated (Pearson's correlation, α=0.05) with speech production latency (interval between

stimulus presentation and onset of speech) across trials for each recording, respectively. The

band response of a recording was considered to be more time-locked to stimulus presentation,

if (1) band response onset to speech onset interval was significantly correlated with speech

production latency, and (2) the correlation coefficient (Pearson's ρ) between band response

onset to speech onset interval and speech production latency was greater than the correlation

coefficient between band response latency and speech production latency. On the contrary, the

band response was considered to be more time-locked to speech onset, if (1) band response

latency was significantly correlated with speech production latency, and (2) the correlation

coefficient between band response latency and speech production latency was greater than the

correlation coefficient between band response onset to speech onset interval and speech

production latency. If a band response did not meet any of the two criteria, it was considered not

locked to either stimulus presentation or speech onset. χ^2 tests were performed to differentiate

the locking properties of broadband gamma increase response and beta decrease response,

and to test if the locking properties differed between recording sides (α=0.05).

Lexical selectivity analysis

We utilized the first 60 trials of recordings that were from the seven subjects with bilateral data

and that showed significant task-related modulations in beta or broadband gamma activity to

make comparisons between the left and right thalamus in terms of band lexical selectivity. A

mean z-scored band response curve averaged across trials and across recordings was

obtained for each frequency band, each trial type (word/nonword), in each side. Trials were

aligned to stimulus presentation for beta and aligned to speech onset for broadband gamma,

based on their respective locking properties. Periods of significant differences between word-

related responses and nonword-related responses were determined with two-tailed paired t-test

using a sliding window of 100ms (α=0.05, Bonferroni corrected).

To quantify the extent of band response difference between word and nonword trials, beta

lexical selectivity indexes and broadband gamma lexical selectivity indexes were calculated for

respective recordings. For each recording, the mean response power values of a particular

frequency band over the corresponding significant time window determined before (see **Task-**

related responses) were calculated for word trials and nonword trials. Two-sample t-test was

then performed between nonword-related power values and word-related power values, and the

resulting t-statistic was the lexical selectivity index of that frequency band for that recording. For

beta, a lexical selectivity index smaller than zero meant that the recording showed stronger beta

activity decrease (thus having more negative z-scored power) during nonword trials than during

word trials, and vice versa. For broadband gamma, a lexical selectivity index greater than zero

indicated that the recording showed stronger broadband gamma activity increase during

nonword reading aloud than during word reading aloud, and vice versa. The beta or broadband

gamma activity of a recording was considered significantly nonword-selective, if lexical

selectivity index was smaller than -1.645 for beta or greater than 1.645 for broadband gamma

(normal approximation of t-distribution). Two-tailed one-sample t-test was performed to test if

lexical selectivity indexes of a selected group of recordings were significantly different from zero

(α=0.05). Two-tailed two-sample t-test was performed to test if lexical selectivity indexes

significantly differed between two groups of recordings (α=0.05).

Analysis of location dependency of lexical selectivity

We further sought to examine possible dependence of band lexical selectivity on recording location, for both beta band and broadband gamma and in both left and right recording sides. Several regression models were applied to the data. First, simple linear regression (Pearson's correlation, α=0.05) was performed to correlate lexical selectivity index with recording location: anterior-posterior location (MNI-defined y coordinate), ventral-dorsal location (MNI-defined z coordinate), and lateral-medial location (MNI-defined x coordinate), respectively. A series of stepwise linear regression models (MATLAB function *stepwiselm*, response variable: lexical selectivity index, predictor variables: MNI-defined x, y, z coordinates and their interactions, α=0.05) were then carried out to test out possible variable interactions, minimize multicollinearity, and determine the final location dependency model of band lexical selectivity. Finally, in order to account for possible lexical selectivity differences that might exist due to subject- and session-specific variations, linear mixed effects models (MATLAB function *fitglme*) were also applied to the data, setting subject and session as random effects.

Acknowledgments

This work was supported by NIH grant U01NS098969 to R.M.R., the Hamot Health foundation to R.M.R., and a University of Pittsburgh Brain Institute NeuroDiscovery Pilot Research Award to R.M.R. This work was supported by the University of Pittsburgh-Tsinghua University Scholars Program to D.W.

Competing interests

The authors declare no competing interests.

References

Alexander, G. E., DeLong, M. R., & Strick, P. L. (1986). Parallel Organization of Functionally Segregated Circuits Linking Basal Ganglia and Cortex. *Annual Review of Neuroscience*, *9*(1), 357–381. https://doi.org/10.1146/annurev.ne.09.030186.002041

Balota, D. A., Yap, M. J., Cortese, M. J., Hutchison, K. A., Kessler, B., Loftis, B., Neely, J. H., Nelson, D. L., Simpson, G. B., & Treiman, R. (2007). The english lexicon project. In *Behavior Research Methods* (Vol. 39, Issue 3, pp. 445–459). Springer New York LLC. https://doi.org/10.3758/BF03193014

Behrens, T. E. J., Johansen-Berg, H., Woolrich, M. W., Smith, S. M., Wheeler-Kingshott, C. A.
M., Boulby, P. A., Barker, G. J., Sillery, E. L., Sheehan, K., Ciccarelli, O., Thompson, A. J.,
Brady, J. M., & Matthews, P. M. (2003). Non-invasive mapping of connections between
human thalamus and cortex using diffusion imaging. *Nature Neuroscience*, *6*(7), 750–757.
https://doi.org/10.1038/nn1075

Bohland, J. W., Bullock, D., & Guenther, F. H. (2010). Neural representations and mechanisms
for the performance of simple speech sequences. *Journal of Cognitive Neuroscience*,
22(7), 1504–1529. https://doi.org/10.1162/jocn.2009.21306

Bosch-Bouju, C., Hyland, B. I., & Parr-Brownlie, L. C. (2013). Motor thalamus integration of
cortical, cerebellar and basal ganglia information: Implications for normal and parkinsonian
conditions. In *Frontiers in Computational Neuroscience* (Issue NOV).
https://doi.org/10.3389/fncom.2013.00163

Bouchard, K. E., Mesgarani, N., Johnson, K., & Chang, E. F. (2013). Functional organization of
human sensorimotor cortex for speech articulation. *Nature*, *495*(7441), 327–332.
https://doi.org/10.1038/nature11911

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.
http://www.ncbi.nlm.nih.gov/pubmed/9176952

Chrabaszcz, A., Neumann, W. J., Stretcu, O., Lipski, W. J., Bush, A., Dastolfo-Hromack, C. A.,
Wang, D., Crammond, D. J., Shaiman, S., Dickey, M. W., Holt, L. L., Turner, R. S., Fiez, J.
A., & Richardson, R. M. (2019). Subthalamic nucleus and sensorimotor cortex activity
during speech production. *Journal of Neuroscience*, *39*(14), 2698–2708.
https://doi.org/10.1523/JNEUROSCI.2842-18.2019

Coltheart, M., Rastle, K., Perry, C., Langdon, R., & Ziegler, J. (2001). DRC: A dual route
cascaded model of visual word recognition and reading aloud. *Psychological Review*,
108(1), 204–256. https://doi.org/10.1037/0033-295X.108.1.204

Crone, N. E., Sinai, A., & Korzeniewska, A. (2006). High-frequency gamma oscillations and
human brain mapping with electrocorticography. *Progress in Brain Research*, *159*, 275–
295. https://doi.org/10.1016/S0079-6123(06)59019-3

Crosson, B. (2013). Thalamic mechanisms in language: a reconsideration based on recent
findings and concepts. *Brain and Language*, *126*(1), 73–88.
https://doi.org/10.1016/j.bandl.2012.06.011

Dietz, N. A. E., Jones, K. M., Gareau, L., Zeffiro, T. A., & Eden, G. F. (2005). Phonological
decoding involves left posterior fusiform gyrus. *Human Brain Mapping*, *26*(2), 81–93.
https://doi.org/10.1002/hbm.20122

Engel, A. K., & Fries, P. (2010). Beta-band oscillations—signalling the status quo? *Current*

Opinion in Neurobiology, *20*(2), 156–165. https://doi.org/10.1016/j.conb.2010.02.015

Ewert, S., Plettig, P., Li, N., Chakravarty, M. M., Collins, D. L., Herrington, T. M., Kühn, A. A., &

Horn, A. (2018). Toward defining deep brain stimulation targets in MNI space: A subcortical

atlas based on multimodal MRI, histology and structural connectivity. *NeuroImage*, *170*,

271–282. https://doi.org/10.1016/j.neuroimage.2017.05.015

Fiez, J. A., Balota, D. A., Raichle, M. E., & Petersent, S. E. (1999). Effects of lexicality,

frequency, and spelling-to-sound consistency on the functional anatomy of reading.

Neuron, *24*(1), 205–218. https://doi.org/10.1016/S0896-6273(00)80833-8

Fonov, V., Evans, A. C., Botteron, K., Almli, C. R., McKinstry, R. C., & Collins, D. L. (2011).

Unbiased average age-appropriate atlases for pediatric studies. *NeuroImage*, *54*(1), 313–

327. https://doi.org/10.1016/j.neuroimage.2010.07.033

Hagoort, P., Indefrey, P., Brown, C., Herzog, H., Steinmetz, H., & Seitz, R. J. (1999). The neural

circuitry involved in the reading of German words and pseudowords: A PET study. *Journal*

of Cognitive Neuroscience, *11*(4), 383–398. https://doi.org/10.1162/089892999563490

Hebb, A. O., & Ojemann, G. A. (2013). The thalamus and language revisited. *Brain and*

Language, *126*(1), 99–108. https://doi.org/10.1016/j.bandl.2012.06.010

Heim, S., Alter, K., Ischebeck, A. K., Amunts, K., Eickhoff, S. B., Mohlberg, H., Zilles, K., von

Cramon, D. Y., & Friederici, A. D. (2005). The role of the left Brodmann's areas 44 and 45

in reading words and pseudowords. *Brain Research. Cognitive Brain Research*, *25*(3),

982–993. https://doi.org/10.1016/j.cogbrainres.2005.09.022

Herbster, A. N., Mintun, M. A., Nebes, R. D., & Becker, J. T. (1997). Regional cerebral blood

flow during word and nonword reading. *Human Brain Mapping*, *5*(2), 84–92.

https://doi.org/10.1002/(sici)1097-0193(1997)5:2<84::aid-hbm2>3.0.co;2-i

Hickok, G. (2014). The architecture of speech production and the role of the phoneme in speech

processing. *Language, Cognition and Neuroscience*, *29*(1), 2–20.

https://doi.org/10.1080/01690965.2013.834370

Hintzen, A., Pelzer, E. A., & Tittgemeyer, M. (2018). Thalamic interactions of cerebellum and

basal ganglia. *Brain Structure & Function*, *223*(2), 569–587.

https://doi.org/10.1007/s00429-017-1584-y

Horn, A., & Kühn, A. A. (2015). Lead-DBS: A toolbox for deep brain stimulation electrode

localizations and visualizations. *NeuroImage*, *107*, 127–135.

https://doi.org/10.1016/j.neuroimage.2014.12.002

Horn, A., Li, N., Dembek, T. A., Kappel, A., Boulay, C., Ewert, S., Tietze, A., Husch, A., Perera,

T., Neumann, W.-J., Reisert, M., Si, H., Oostenveld, R., Rorden, C., Yeh, F.-C., Fang, Q.,

Herrington, T. M., Vorwerk, J., & Kühn, A. A. (2019). Lead-DBS v2: Towards a

comprehensive pipeline for deep brain stimulation imaging. *NeuroImage*, *184*, 293–316.

https://doi.org/10.1016/j.neuroimage.2018.08.068

Hwang, K., Bertolero, M. A., Liu, W. B., & D'Esposito, M. (2017). The Human Thalamus Is an

Integrative Hub for Functional Brain Networks. *The Journal of Neuroscience*, *37*(23), 5594–

5607. https://doi.org/10.1523/JNEUROSCI.0067-17.2017

Johnson, M. D., & Ojemann, G. A. (2000). The role of the human thalamus in language and

memory: Evidence from electrophysiological studies. *Brain and Cognition*, *42*(2), 218–230.

https://doi.org/10.1006/brcg.1999.1101

Juphard, A., Vidal, J. R., Perrone-Bertolotti, M., Minotti, L., Kahane, P., Lachaux, J.-P., & Baciu,

M. (2011). Direct Evidence for Two Different Neural Mechanisms for Reading Familiar and

Unfamiliar Words: An Intra-Cerebral EEG Study. *Frontiers in Human Neuroscience*, *5*, 101.

https://doi.org/10.3389/fnhum.2011.00101

Klostermann, F., Nikulin, V. V, Kühn, A. A., Marzinzik, F., Wahl, M., Pogosyan, A., Kupsch, A.,

Schneider, G.-H., Brown, P., & Curio, G. (2007). Task-related differential dynamics of EEG

alpha- and beta-band synchronization in cortico-basal motor structures. *European Journal*

of Neuroscience, *25*(5), 1604–1615. https://doi.org/10.1111/j.1460-9568.2007.05417.x

Kühn, A. A., Williams, D., Kupsch, A., Limousin, P., Hariz, M., Schneider, G.-H., Yarrow, K., &

Brown, P. (2004). Event-related beta desynchronization in human subthalamic nucleus

correlates with motor performance. *Brain*, *127*(4), 735–746.

https://doi.org/10.1093/brain/awh106

Levelt, W. J. M., Roelofs, A., & Meyer, A. S. (1999). A theory of lexical access in speech

production. In *Behavioral and Brain Sciences* (Vol. 22, Issue 1, pp. 1–75).

https://doi.org/10.1017/S0140525X99001776

Lipski, W. J., Alhourani, A., Pirnia, T., Jones, P. W., Dastolfo-Hromack, C., Helou, L. B.,

Crammond, D. J., Shaiman, S., Dickey, M. W., Holt, L. L., Turner, R. S., Fiez, J. A., &

Richardson, R. M. (2018). Subthalamic nucleus neurons differentially encode early and late

aspects of speech production. *Journal of Neuroscience*, *38*(24), 5620–5631.

https://doi.org/10.1523/JNEUROSCI.3480-17.2018

Llano, D. A. (2015). The Thalamus and Language. In G. Hickok & S. L. Small (Eds.).

Neurobiology of Language (pp. 95-114). Cambridge, MA: Academic Press.

Macchi, G., & Jones, E. G. (1997). Toward an agreement on terminology of nuclear and

subnuclear divisions of the motor thalamus. *Journal of Neurosurgery*, *86*(1), 77–92.

https://doi.org/10.3171/jns.1997.86.1.0077

Mechelli, A., Gorno-Tempini, M. L., & Price, C. J. (2003). Neuroimaging studies of word and

pseudoword reading: Consistencies, inconsistencies, and limitations. *Journal of Cognitive*

Neuroscience, *15*(2), 260–271. https://doi.org/10.1162/089892903321208196

Moore, M. W., Fiez, J. A., & Tompkins, C. A. (2017). Consonant Age-of-Acquisition Effects in

Nonword Repetition Are Not Articulatory in Nature. *Journal of Speech, Language, and*

Hearing Research, *60*(11), 3198–3212. https://doi.org/10.1044/2017_JSLHR-L-16-0359

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data.

Computational Intelligence and Neuroscience, *2011*, 1–9.

https://doi.org/10.1155/2011/156869

Paradiso, G. (2004). Involvement of human thalamus in the preparation of self-paced

movement. *Brain*, *127*(12), 2717–2731. https://doi.org/10.1093/brain/awh288

Ray, S., Crone, N. E., Niebur, E., Franaszczuk, P. J., & Hsiao, S. S. (2008). Neural correlates of

high-gamma oscillations (60-200 Hz) in macaque local field potentials and their potential

implications in electrocorticography. *The Journal of Neuroscience : The Official Journal of*

the Society for Neuroscience, *28*(45), 11526–11536.

https://doi.org/10.1523/JNEUROSCI.2848-08.2008

Ray, S., & Maunsell, J. H. R. (2011). Different Origins of Gamma Rhythm and High-Gamma

Activity in Macaque Visual Cortex. *PLoS Biology*, *9*(4), e1000610.

https://doi.org/10.1371/journal.pbio.1000610

Tiedt, H. O., Ehlen, F., Krugel, L. K., Horn, A., Kühn, A. A., & Klostermann, F. (2017).

Subcortical roles in lexical task processing: Inferences from thalamic and subthalamic

event-related potentials. *Human Brain Mapping*, *38*(1), 370–383.

https://doi.org/10.1002/hbm.23366

Tzagarakis, C., Ince, N. F., Leuthold, A. C., & Pellizzer, G. (2010). Beta-band activity during

motor planning reflects response uncertainty. *The Journal of Neuroscience : The Official*

Journal of the Society for Neuroscience, *30*(34), 11270–11277.

https://doi.org/10.1523/JNEUROSCI.6026-09.2010

Uhlhaas, P. J., Pipa, G., Neuenschwander, S., Wibral, M., & Singer, W. (2011). A new look at

gamma? High- (>60 Hz) γ-band activity in cortical networks: function, mechanisms and

impairment. *Progress in Biophysics and Molecular Biology*, *105*(1–2), 14–28.

https://doi.org/10.1016/j.pbiomolbio.2010.10.004

Wahl, M., Marzinzik, F., Friederici, A. D., Hahne, A., Kupsch, A., Schneider, G. H., Saddy, D.,

Curio, G., & Klostermann, F. (2008). The Human Thalamus Processes Syntactic and

Semantic Language Violations. *Neuron*, *59*(5), 695–707.

https://doi.org/10.1016/j.neuron.2008.07.011

Xu, B. (2001). Conjoint and Extended Neural Networks for the Computation of Speech Codes:

The Neural Basis of Selective Impairment in Reading Words and Pseudowords. *Cerebral*

Cortex, *11*(3), 267–277. https://doi.org/10.1093/cercor/11.3.267

Zhang, D., Snyder, A. Z., Shimony, J. S., Fox, M. D., & Raichle, M. E. (2010). Noninvasive

Functional and Structural Connectivity Mapping of the Human Thalamocortical System.

Cerebral Cortex, *20*(5), 1187–1194. https://doi.org/10.1093/cercor/bhp182

Thalamic encoding of lexical status is lateralized during reading aloud

Wang et al., 2020

Supplementary Information

Supplementary Figure 1: Correlation between left thalamic broadband gamma lexical
selectivity and recording locations along ventral-dorsal and lateral-medial axes.
Broadband gamma lexical selectivity indexes of left side lead recordings that showed significant
task-related broadband gamma increase responses are correlated (Pearson's correlation,
α=0.05) with ventral-dorsal locations (**a**) and lateral-medial locations (**b**) of the recordings,
respectively.

Supplementary Figure 2: Broadband gamma lexical selectivity does not depend on anterior-posterior location of the recording in the right thalamus. a Right side lead recording sites are flipped to the left and plotted together with anatomical structures (thalamus in yellow, VA and VLa in green, VLp in blue, and Vim in orange), viewed from a lateral direction. Recording sites where significantly nonword-selective recordings in terms of broadband gamma response amplitudes (broadband gamma lexical selectivity index>1.645 based on normal approximation of t-distribution) were observed are filled with black (n=0), and remaining recording sites are shown in open circles. **b** Broadband gamma lexical selectivity indexes of right side lead recordings that showed significant task-related broadband gamma increase responses are correlated (Pearson's correlation, α=0.05) with recording locations along the anterior-posterior axis. Recordings inside VA or VLa are green-colored and recordings inside VLp are blue-colored.

Supplementary Figure 3: Correlation between right thalamic broadband gamma lexical selectivity and recording locations along ventral-dorsal and lateral-medial axes.
Broadband gamma lexical selectivity indexes of right side lead recordings that showed significant task-related broadband gamma increase responses are correlated (Pearson's correlation, α=0.05) with ventral-dorsal locations (**a**) and lateral-medial locations (**b**) of the recordings, respectively.

Supplementary Figure 4: Beta lexical selectivity does not depend on anterior-posterior location of the recording in either side. a Left side lead recording sites are plotted together with anatomical structures (thalamus in yellow, VA and VLa in green, VLp in blue, and Vim in orange), viewed from a lateral direction. Recording sites where significantly nonword-selective recordings in terms of beta response amplitudes (beta lexical selectivity index<-1.645 based on normal approximation of t-distribution) were observed in either session are filled with black, and remaining recording sites are shown in open circles. **b** Right side lead recording sites are flipped to the left and plotted in the same way. **c** Beta lexical selectivity indexes of left side lead recordings that showed significant task-related beta decrease responses are correlated (Pearson's correlation, α=0.05) with recording locations along the anterior-posterior axis. **d** Beta lexical selectivity indexes of right side lead recordings that showed significant task-related beta decrease responses are correlated (Pearson's correlation, α=0.05) with recording locations along the anterior-posterior axis. In **c** and **d**, recordings inside VA or VLa are green-colored and recordings inside VLp are blue-colored.

Supplementary Figure 5: Correlation between thalamic beta lexical selectivity and recording locations along ventral-dorsal and lateral-medial axes. a, **c** Beta lexical selectivity indexes of left side lead recordings that showed significant task-related beta decrease responses are correlated (Pearson's correlation, α=0.05) with ventral-dorsal locations (**a**) and lateral-medial locations (**c**) of the recordings, respectively. **b**, **d** Beta lexical selectivity indexes of right side lead recordings that showed significant task-related beta decrease responses are correlated (Pearson's correlation, α=0.05) with ventral-dorsal locations (**b**) and lateral-medial locations (**d**) of the recordings, respectively.

819 **Supplementary Table 1: Broadband gamma increase response and beta decrease**

820 **response differ in locking properties.**

Response type	Number of recordings	Stimulus-locked (%)	Speech-locked (%)	Not locked (%)
Broadband gamma increase response	91	19 (21)	70 (77)	2 (2)
Beta decrease response	66	43 (65)	19 (29)	4 (6)
χ^2	-	31.4	36.1	1.55
P	-	$< 10^{-5}$	$< 10^{-5}$	0.21

822 **Supplementary Table 2: Locking properties of broadband gamma increase response and**

823 **beta decrease response do not differ between recording sides.**

Response type	Number of recordings	Stimulus-locked (%)	Speech-locked (%)	Not locked (%)
Broadband gamma increase response				
Left	59	10 (17)	47 (80)	2 (3)
Right	32	9 (28)	23 (72)	0 (0)
χ^2	2.49			
P	0.29			
Beta decrease response				
Left	49	31 (63)	14 (29)	4 (8)
Right	17	12 (71)	5 (29)	0 (0)
χ^2	1.50			
P	0.47			

825 **Supplementary Table 3: Stimulus lists.**

Stimulus list 1		Stimulus list 2		Stimulus list 3		Stimulus list 4	
First 60	Last 60	First 60	Last 60	First 60	Last 60	First 60	Last 60
daif	daif	dight	dight	deave	deave	dipe	dipe
van	dutt	void	dute	vowed	doif	vote	doan
dutt	foat	dute	fape	doif	feam	doan	fode
vied	fute	vet	foon	vat	feeve	vain	fett
foat	meef	fape	mide	feam	vune	fode	meave
town	mipe	type	tep	tied	mafe	towed	mep
fute	neave	foon	veem	feeve	mupe	fett	neam
teed	nime	tune	naid	team	nipe	time	noit
meef	paim	mide	noop	vune	noove	meave	peff
putt	pote	pain	pime	poof	pode	pine	poom
mipe	teeve	tep	peaf	mafe	poun	mep	taid
pout	toff	peeve	taff	pave	teep	phone	tife
neave	vapp	veem	toove	mupe	tem	neam	vadd
note	vife	node	vade	need	vight	paid	vate
nime	vean	naid	vome	nipe	ved	noit	veet
moon	laz	mood	zool	moat	soith	neat	leeth
paim	saith	noop	loath	noove	luth	peff	sish
mauve	lesh	mean	luss	main	losh	mode	luthe
pote	rel	pime	rall	pode	rith	poom	ral
knife	reez	knot	suzz	knight	ruzz	might	riz
teeve	shuss	peaf	saze	poun	sar	taid	soush
fought	shule	fight	seash	foot	shoss	knit	shar
toff	sosh	taff	shez	teep	shiz	tife	shoth
five	sithe	feed	shile	foam	sile	food	thaze
vapp	thace	toove	thar	tem	thale	vadd	theal
fade	thauz	fame	thil	fain	thall	fate	sule
vife	thole	vade	thure	vight	thice	vate	sazz
deep	thoss	dine	thush	doubt	thel	dive	thosh
vean	zal	vome	zel	ved	zoice	veet	zoil
dawn	zole	date	soth	dime	zus	dean	zil
laz	daif	zool	dight	soith	deave	leeth	dipe

zoom	dutt	zone	dute	zip	doif	zap	doan
saith	foat	loath	fape	luth	feam	sish	fode
thought	fute	thud	foon	thumb	feeve	thief	fett
lesh	meef	luss	mide	losh	vune	luthe	meave
thin	mipe	theme	tep	that	mafe	them	mep
rel	neave	rall	veem	rith	mupe	ral	neam
this	nime	wrath	naid	zeal	nipe	thus	noit
reez	paim	suzz	noop	ruzz	noove	riz	peff
shoes	pote	these	pime	those	pode	thaws	poom
shuss	teeve	saze	peaf	sar	poun	soush	taid
shawl	toff	sure	taff	south	teep	shies	tife
shule	vapp	seash	toove	shoss	tem	shar	vadd
sell	vife	soar	vade	sill	vight	sews	vate
sosh	vean	shez	vome	shiz	ved	shoth	veet
sear	laz	size	zool	shore	soith	sale	leeth
sithe	saith	shile	loath	sile	luth	thaze	sish
sash	lesh	shell	luss	shale	losh	share	luthe
thace	rel	thar	rall	thale	rith	theal	ral
rush	reez	seal	suzz	role	ruzz	lace	riz
thauz	shuss	thil	saze	thall	sar	sule	soush
rise	shule	rose	seash	rash	shoss	lash	shar
thole	sosh	thure	shez	thice	shiz	sazz	shoth
race	sithe	rail	shile	raise	sile	lore	thaze
thoss	thace	thush	thar	thel	thale	thosh	theal
loathe	thauz	loss	thil	lure	thall	lose	sule
zal	thole	zel	thure	zoice	thice	zoil	sazz
laze	thoss	lies	thush	lows	thel	rule	thosh
zole	zal	soth	zel	zus	zoice	zil	zoil
laws	zole	leash	soth	lice	zus	rice	zil

828 **Supplementary Table 4: *P* values of statistical tests comparing psycholinguistic**

829 **parameters between word and nonword stimuli.**

	Stimulus list 1	Stimulus list 2	Stimulus list 3	Stimulus list 4
Weighted phonological neighborhood density	0.23	0.43	0.88	0.86
Summed bigram frequency	0.97	0.51	0.39	0.32
Summed phoneme probability	0.47	0.33	0.37	0.51
Summed biphone probability	0.65	0.65	0.65	0.90

830 Two-tailed two-sample t-test, n=60, α=0.05.