1 Midcell localization of PBP4 of *Escherichia coli* modulates the timing of
2 divisome assembly
3
4

5 Jolanda Verheul[1#], Adam Lodge[2,7#], Hamish C.L. Yau[2,8#], Xiaolong Liu[1,9#], Xinwei Liu[1],
6 Alexandra S. Solovyova[3], Athanasios Typas[4,5] Manuel Banzhaf[6*], Waldemar Vollmer[2**], and
7 Tanneke den Blaauwen[1***]
8
[1]Bacterial Cell Biology, Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam;
Science Park 904, 1098 XH Amsterdam, The Netherlands.
[2]Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle upon Tyne, NE2 4AX,
United Kingdom.
[3]NUPPA, Leech Building, Framlington Place, Newcastle upon Tyne NE2 4HH, UK
[4]European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, Germany
[5]European Molecular Biology Laboratory, Structural & Computational Unit, Heidelberg, Germany
[6]Institute of Microbiology & Infection and School of Biosciences, University of Birmingham, Edgbaston,
Birmingham, UK. B15 2TT
[7]Present address: Iksuda Therapeutics, The Biosphere, Draymans Way, Newcastle Helix, Newcastle upon Tyne,
NE4 5BX, UK
[8]Present address: Procter & Gamble, Newcastle Innovation Centre, Whitley Road, Newcastle-Upon Tyne,
NE12 9TS, UK.
[9]Department of Biochemistry, University of Oxford, OX1 3QU, Oxford, United Kingdom.
23
24 [#]contributed equally.
25 *Corresponding author. Tel: +44 121 4145586; Email: m.banzhaf@bham.ac.uk
26 **Corresponding author. Tel: +44 191 208 3216; Email: w.vollmer@ncl.ac.uk
27 ***Corresponding author. Tel: +31 20 525 3852; Email: t.denblaauwen@uva.nl
28
29 **Running title:** PBP4 modulates divisome assembly timing

30
31
ABSTRACT (176 words)

Insertion of new material into the *Escherichia coli* peptidoglycan (PG) sacculus between the

cytoplasmic membrane and the outer membrane requires a well-organized balance between

synthetic and hydrolytic activities to maintain cell shape and avoid lysis. The hydrolytic

enzymes outnumber the enzymes that insert new PG by far and very little is known about

their specific function. Here we show that the DD-carboxy/endopeptidase PBP4 localizes in a

PBP1A/LpoA and FtsEX dependent fashion at midcell during septal PG synthesis. Midcell

localization of PBP4 requires its non-catalytic domain 3 of unknown function, but not the

activity of PBP4 or FtsE. Microscale thermophoresis with isolated proteins shows that domain

3 is needed for the interaction with NlpI, but not PBP1A or LpoA. *In vivo* crosslinking

experiments confirm the interaction of PBP4 with PBP1A and LpoA. We propose that PBP4

functions together with the amidases AmiA and B to create denuded glycan strands to attract

the initiator of septal PG synthesis, FtsN. Consistent with this model, we found that the

divisome assembly at midcell was significantly affected in cells lacking PBP4.

IMPORTANCE (140 words)

Peptidoglycan biosynthesis is a major target for antibacterials. The covalently closed

peptidoglycan mesh, called sacculus, protects the bacterium from lysis due to its turgor.

Sacculus growth is facilitated by the balanced activities of synthases and hydrolases, and

disturbing this balance leads to cell lysis and bacterial death. Because of the large number

and possible redundant functions of peptidoglycan hydrolases, it has been difficult to

decipher their individual functions. In this paper we show that the DD-endopeptidase PBP4

localizes at midcell during septal peptidoglycan synthesis in *Escherichia coli* and is important

for the timing of the assembly of the division machinery. This shows that inhibition of certain

hydrolases could weaken the cells and might enhance antibiotic action.

KEYWORDS: bacterial cell division, endopeptidase, immunolabeling, LpoA, NlpI, PBP1A,

PBP4, peptidoglycan, preseptal, protein interactions

INTRODUCTION

The peptidoglycan (PG) layer is sandwiched between the cytoplasmic membrane (CM) and

the outer membrane (OM) of the Gram-negative bacterium *Escherichia coli* forming a

covalently closed network of glycan strands that are interconnected by short peptides [1].

The PG layer maintains the shape of the bacterium by stabilizing the cell against its high

internal osmotic pressure. To proliferate, the rod-shaped bacterium grows in length and then

divides by binary fission into two equally sized daughter cells. The balanced activities of PG

synthases and hydrolases ensures a safe extension of the PG layer without defects that would

cause cell lysis [2]. Beta-lactam antibiotics disturb this balance by inactivating the

transpeptidase (TPase) activity of the penicillin binding proteins (PBP)s causing lysis of

growing cells. We currently know more than twenty PG hydrolases that can cleave either in

the glycan chains or peptides within the PG layer, but how their activities are regulated is

poorly understood. This is particularly true for the PG endopeptidases [2]. Recently, the OM

bound lipoprotein NlpI was shown to interact with similar affinity with several

endopeptidases, suggesting that NlpI acts as an interaction hub for endopeptidases to

regulate their activities by competing protein-protein interactions. [3].

The basic precursor (lipid II) for PG synthesis is the β-(1,4) linked disaccharide *N*-

acetylglucosamine-*N*-acetylmuramic acid (Glc*N*Ac-Mur*N*Ac) with an undecaprenol-

pyrophosphate linked to C1 of Mur*N*Ac and a stem peptide with the sequence L-alanine-D-

iso-glutamate-diaminopimelic acid (Dap)-D-alanine-D-alanine coupled to the C3 of Mur*N*Ac.

Lipid II is used by glycosyltransferases (GTases), which polymerize the glycan chains, and

TPase, which crosslink the stem peptides. The main PG synthases in *E. coli* are the class A PBPs

(PBP1A and PBP1B) with GTase and TPase activities [4–8], the integral membrane proteins

RodA and FtsW with GTase activity [9–11] and their partner TPases PBP2 and PBP3,

respectively [12–15].

The hydrolases are present in greater redundancy than the synthases [2,16]. The

carboxypeptidases (CPase) PBP5, PBP6A and PBP6B remove the terminal D-Ala from nascent

pentapeptides [17–20]. The endopeptidases (EPases) PBP4, PBP7, MepA, MepM, MepH, and

MepS cleave the crosslinks in the PG [21–23]. The amidases AmiA-D hydrolyze the bond

between MurNAc and L-alanine at position 1 of the stem peptide [24–27]. Finally, the lytic

transglycosylases Slt70 and MltA-G cleave the glycan strands to release 1,6-anhydroMurNAc

containing turnover products [28–30]. The large number of hydrolases reflects the

adaptability to various environmental conditions {Pazos; Muellersw} but begs the question of

how their potentially dangerous activities are controlled to avoid cell lysis.

Binary fission in *E. coli* is initiated by the assembly of the Z-ring at midcell, which

consist of a network of dynamic FtsZ filaments that are anchored to the cytoplasmic

membrane by FtsA and ZipA [31,32]. The Z-ring also recruits the PG synthases PBP1A and

PBP1B to midcell [4,33]. A number of Z-associated proteins play a role in the organization and

dynamics of the ring [34–39]. Of these, the integral membrane protein FtsX interacts with

FtsA [40] and its cytoplasmic ATPase partner protein FtsE interacts with FtsZ [41]. Both localize

early to midcell, together with the Z-ring [24]. As a complex FtsEX recruits EnvC, which

activates AmiA and AmiB that are needed for septum cleavage [26,42]. FtsE versions without

ATPase activity still localize at midcell and recruit as FtsEX-EnvC complex but the latter is not

able to activate the amidases [43]. Presumably an ATPase dependent conformational change

in FtsEX induces a conformational change in EnvC, which is then able to relieve an

autoinhibitory alpha-helix from the active site of AmiA and AmiB, activating them [44]. The

third amidase AmiC is activated by the OM-anchored lipoprotein NlpD [24] and all three

amidases AmiA-C are activated under certain stress conditions by the recently identified ActS

[46,47]. The recruitment of the other cell division proteins occurs with a time delay of about

20% of the cell division cycle, depending on the growth condition [48–50]. FtsK, FtsBLQ, FtsW,

PBP3 and FtsN localize in an interdependent fashion. FtsBLQ inhibit the PG synthase activity

of PBP3, FtsW and PBP1B until they are outcompeted by the accumulation of FtsN that

relieves this inhibition and initiates septum synthesis [51–54].

While investigating the specific function of various PG hydrolases, we found that PBP4

localizes specifically at midcell as part of the division machinery. PBP4 is a periplasmic

endopeptidase [55] with a C-terminal amphipathic alpha-helix that associates with

membranes [56]. Deletion of *dacB*, which encodes PBP4, does not have morphological

consequences but causes a decrease in the percentage of PG crosslinks and of PG-attached

lipoprotein [57] and an increased sensitivity to bile salts [58]. Overproduction of PBP4 is toxic

[59] presumably because the enhanced cleavage of peptide cross-links weakens the PG mesh

and eventually causes lysis. PBP4 has three domains that are assembled in an unusual way

[60] (Fig. 1A). A non-catalytical domain of unknown function (domain 2) is inserted into the

transpeptidase domain 1, and a third domain (domain 3) is inserted into domain 2. Domain 3

is positioned above the active site with its catalytic serine 62, which resides in domain 1, and

might be involved in substrate binding or regulation [61] (Fig. 1A). Site directed mutagenesis

suggested that residues from domain 1 (R361) and 2 (D155) are important for the

endopeptidase activity [62].

Here we determined the localization of PBP4 and different variants during the cell

cycle of *E. coli.* The timing of localization at midcell and interactions of PBP4 support a role in

remodeling of PG synthesized by the divisome.

RESULTS

PBP4 localizes to the lateral wall and at midcell. To determine the localization of PBP4 in *E.*

coli cells as a function of the bacterial cell cycle, we generated an antiserum against PBP4 and

removed unspecific antibodies by adsorption to cells of a *dacB* deletion mutant that lacks

PBP4. The purified antibodies were specific for PBP4 (Fig. S1) and used to immunolabel the

wild-type strain MC4100 grown in rich (TY) medium at 37°C. PBP4 localized strongly at midcell

and to a lesser extend in the lateral wall in wild-type cells (Fig. 1B).

PBP4 localizes between the peptidoglycan layer and the outer membrane. Although

it was reported that PBP4 has a stretch of weakly amphipathic amino acids at its C-terminus

[56], it was not known whether PBP4 freely diffuses in the periplasm or associates with the

membrane or other proteins. It has been previously observed that chemically fixing freely

diffusing periplasmic proteins can shock them into the cell poles [12,63]. The fact that we did

not observe any polar localization of PBP4 after fixation (Fig. 1B and S1), suggests that it is

either associated with the membrane or other proteins. Next, we investigated whether PBP4

was accessible to anti-PBP4 in cells with an intact PG layer. For this, wild-type cells were grown

in TY at 37°C and immunolabeled without the typically used Triton X-100-mediated

permeabilization of the OM or without digesting the PG mesh with lysozyme. PBP4 was not

accessible without permeabilization of the OM but was fully accessible in cells with an intact

PG (Fig. 1C). In contrast, proteins like FtsN and PBP3 were not accessible without degradation

of the PG layer by lysozyme (Fig. 1C). The OM pore inserting BAM complex is also not

accessible without lysozyme treatment [64]. This suggests that PBP4 resides between the OM

and the PG layer.

Activity of PBP4 is not needed for midcell localization. Substrate binding can be a key

determinant in the localization of proteins [18,65,66]. To test whether PBP4 substrate binding

is needed to localize at division sites we expressed the active site mutants PBP4(S62G) and

(S62A) [61] that fails to bind β-lactams and therefore is not able to hydrolyze the PG stem

peptide (Fig. S2 and S3). A plasmid encoding the mutant protein was constitutively expressed

in a Δ*dacB* background by the leakage of a weakened p*Trc99*A promoter. Interestingly, the

wild-type and active site mutant when expressed from plasmid were both able to localize at

midcell (Fig. 2). This indicates that although PBP4(S62G) cannot hydrolyze its substrate or bind

β-lactams covalently (Fig. S2C Fig. S3), it is either still able to interact with its substrate or it

localizes through protein interactions.

Since PBP4(S62G) could still interact with its PG substrate, we mutagenized the gene

to express PBP4 versions in which residues D155 or R361 involved in the binding of the DAP

residue in the stem peptide by alanine [62] were replaced. The genes encoding PBP4(D155A)

or PBP4(R361A) were expressed by promoter leakage from plasmid in a ΔdacB background,

and both PBP4 versions localized at midcell (Fig. 2). An immunoblot analysis showed that all

mutants were expressed to similar levels and that PBP4(R361A) and PBP4(D155A) were able

to bind the fluorescent β-lactam Bocilin-FL (Fig. S2). Therefore, all three active site protein

variants are potentially able to interact with substrate as an inactive protein.

A PBP4 version lacking domain 3 (PBP4ΔD3) was still able to bind β-lactams (Fig. S2B

and S3) but did not have DD-carboxypeptidase or DD-endopeptidase activity (Fig. S3).

Analytical ultracentrifugation (Fig. S4A) and circular dichroism analysis (Fig. S4B) of isolated

PBP4ΔD3, showed that the removal of domain 3 did not affect the structure, the dimerization

of the rest of the protein or its interaction with PG sacculi (Fig. S4C), which suggests a folded

active site domain. However, although the inactive PBP4ΔD3 was capable of binding substrate,

it did not localize at midcell (Fig. 2). Hence, we conclude that the midcell localization of PBP4

is largely driven by protein-protein interactions and to a lesser extent by substrate interaction

but is independent from its own activity.

PBP4 localizes at inactive divisomes. To further dissect whether the localization of

PBP4 at the division site is dependent on the availability of its substrate or the presence of

other proteins at the division site, cell division was inhibited through the specific inactivation

of PBP3 by aztreonam [67]. Aztreonam treatment for 1-3 mass doublings stalls the division

machinery at midcell resulting in filamentous growth. In the longer filaments new division

machineries are formed and localize at future division sites [5]. In these filaments PBP4 still

localized albeit somewhat weaker than in untreated cells at all possible division sites (Fig. S5),

indicating that the activity of PBP3 was not required for the recruitment of PBP4. While PBP3

is inhibited, preseptal PG synthesis [68] still occurs at the potential division sites in the

filamenting cells [18,33,69]. The pre-septal localization of PBP4 mimics the localization of

PBP1A and PBP1B, which were previously shown to be recruited to preseptal PG synthesis

sites by early cell division proteins [4,33].

PBP1A/LpoA assist in midcell localization of PBP4. PBP4 was recently reported to

interact with PBP1A, LpoA and the EPase adaptor protein NlpI [3]. To investigate whether

PBP1A and/or LpoA proteins were needed for midcell localization of PBP4, strains deleted for

mrcA (PBP1A), *lpoA*, *mrcB* (PBP1B) or *lpoB* were grown in rich medium at 37°C and

immunolabeled with anti-PBP4. The amount of PBP4 that localized at midcell was identical

for the parental wild-type strain BW25113, Δ*mrcB* and Δ*lpoB,* but reduced in the Δ*mrcA* and

Δ*lpoA* strains (even after correction for the smaller diameter of these cells, [70], whereas the

PBP4 concentration was similar for all strains (Fig. 3). This could indicate that PBP1A/LpoA

assist in the initial localization of PBP4 at midcell.

NlpI seems not to be involved in midcell localization of PBP4. We next tested if the

interaction with the OM anchored NlpI affected the localization of PBP4. In the Δ*nlpI* cells the

amount of PBP4 at midcell per μm average cell circumference was reduced by 42 ± 6.7% (n =

3) of the wild type and the protein localized later at midcell than in the wild-type cells (Fig.

S6). However, the PBP4 concentration in the Δ*nlpI* strain was also lower than in the wild-type

cells (64.8 ± 17.6%, n = 3). This could indicate that the cells down-regulate PBP4 expression

or that PBP4 becomes unstable in the Δ*nlpI* strain. Labeling of PBP4 in a Δ*nlpI* strain did not

require lysozyme (Fig. S6), suggesting that NlpI is not needed to maintain PBP4 localized

outside the PG layer, which could be facilitated by additional interactions of PBP4.

PBP4 domain 3 facilitates interaction with NlpI but not PBP1A/LpoA. Microscale

thermophoresis (MST) in which one protein is labelled with a fluorophore and titrated with

an unlabeled partner protein was used to analyze the interactions between PBP4 and its

known interaction partners further. The difference in thermophoresis of the complex versus

the unbound fluorescently labeled proteins is then used to calculate the affinity of the binding

partners. Different versions of PBP4 and its interaction partners were purified and assayed

for direct protein-protein interactions. Wild-type PBP4 interacted with PBP1A, LpoA and NlpI

[3]. PBP4 lacking domain 3 (PBP4ΔD3), which did not localize at midcell (Fig. 2), still interacted

with PBP1A and LpoA but not with NlpI (Fig. 4). These results suggest that domain 3 of PBP4

facilitates the interaction with NlpI, but not PBP1A.

NlpI and PBP1A/LpoA have different binding sites in PBP4. PBP4 can form a multi-

enzyme complex with NlpI and PBP1A/LpoA [3]. PBP1A interacted with PBP4 and could be

cross-linked to PBP4 in cells (Fig. 4C). We found that full length LpoA interacted with both

PBP4 and PBP4ΔD3, with apparent K_D values of 315 ± 38 nM and 153 ± 31 nM, respectively

(Fig. 4A; Fig. S7). This suggests that domains 1 or 2 of PBP4 are sufficient for the interaction

with LpoA. MST assays of PBP4 or PBP4ΔD3 with LpoA's C-terminal (residues N257-S678;

$LpoA^C$) or N-terminal (LpoA residues G28-T256; $LpoA^N$) domains also yielded positive binding

curves (Fig. 4A; Fig. S7) [71–73]. We found that $LpoA^C$ and the full length LpoA version had

comparable affinities for both PBP4 versions, suggesting that the C-terminal domain of LpoA

is sufficient for interaction with PBP4 (Fig. 4A). Interestingly, the removal of domain 3 from

PBP4 resulted in an increased affinity of $LpoA^N$ for PBP4 (app. K_D decreased by ~50 fold) (Fig.

4A Fig. S7). Hence, the interaction between the N-terminal domain of LpoA and PBP4 possibly

occurs via a different mechanism dependent on a conformational rearrangement of domain

3 of PBP4. This suggests that the two domains of LpoA, which likely serve different primary

functions, are both sufficient for an interaction with PBP4, albeit with varying affinities. In

conclusion PBP4 robustly interacts with PBP1A/LpoA independent of its domain 3.

LpoA has a small effect on PBP4 activity. We found previously that NlpI does not

affect PBP4 activity [3] but considered it possible that PBP4 interaction with PBP1A/LpoA may

affect its activity. We first verified that NlpI or PBP4 did not affect the activity of PBP1A in the

presence or absence of LpoA using an *in vitro* glycosyltransferase assay (Fig. 5). We next

tested if PBP1A or LpoA affected the activity of PBP4 using *in vitro* PG degradation assays.

LpoA decreased the activity of PBP4 by about 30% in a continuous DD-CPase assay with

soluble substrate and an endpoint PG digestion assay (Fig 5B and C). To assess whether LpoA

also affects PBP4 activity *in vivo*, a Δ*dacB*Δ*lpoA* strain and Δ*dacB* strain were transformed

with the plasmids expressing PBP4 or the active site mutant PBP4(S62G) and the toxicity of

their expression was compared. Although the double mutant was less fit than the single

mutant, it was equally sensitive to the overexpression of PBP4 (Fig. 5D), leaving no indication

that LpoA also inhibits the activity of PBP4 *in vivo*.

PBP4 localizes as early divisome protein. Since PBP4 localizes at midcell it could be

part of or associated with the divisome. The division machinery is assembled in two successive

steps [48,74]. First, FtsZ and its membrane bound partners ZipA and FtsA in conjunction with

other Z-ring associated proteins form the proto-ring that localizes at midcell

{denBlaauwen:2017dq}. Second, after approximately 20% of the cell division cycle the

remaining proteins assemble subsequently at the division site. To obtain more details on PBP4

localization in relation to other cell division proteins, MC4100 cells were grown to steady state

in minimal glucose medium at 28°C. The cells were immunolabeled with antibodies against

PBP4, FtsZ (early localizing), or FtsN) late localizing). PBP4 localized strongly at midcell and

hardly in the lateral cell wall (Fig. 6). To determine whether PBP4 belongs to the early or late

divisome localizing proteins, we analyzed the timing of its midcell localization. To this end, we

plotted the extra fluorescence (FCPlus) present at midcell in comparison to the amount of

fluorescence in the rest of the cell (Fig. 6). PBP4 localization coincided with that of FtsZ,

indicating that PBP4 localizes relatively early.

Midcell localization of PBP4 depends on FtsE. To investigate if PBP4 localization is

dependent on the presence of the Z-ring or the later localizing division associated proteins

we localized PBP4 in various division defective strains. First, the localization of PBP4 was

assessed in a strain that expressed Tre1 that ADP-ribosylates FtsZ at R174, which renders it

unable to participate in protofilament formation [75,76]. The inactive E415Q variant of Tre1

was used as negative control. PBP4 was unable to localize in cells with insufficient FtsZ

protofilaments to form division Z-rings (Fig. 7A), whereas it localized as in WT cells in the

presence of the TreE415Q variant, proving that a Z-ring is needed for PBP4 localization. Next,

strains harboring temperature sensitive alleles of FtsE, FtsQ and PBP3 (FtsI) were grown in

minimal glucose medium at 28°C to an OD_{450} of 0.2, diluted 1:4 in prewarmed medium of 28°C

or 42°C and grown for two mass doublings before PBP4 immunolabelling. PBP4 localized in

filamentous cells with thermo-labile FtsQ or PBP3 indicating that these proteins are not

needed to recruit PBP4 (Fig. 7B). However, PBP4 localized very diffuse around potential

division sites in the filamentous *ftsE*(ts) cells (Fig. 7B). This suggests that a functional FtsE is

important for PBP4 localization. The FtsEX/EnvC complex assists in the assembly of the

divisome [77,78] and is essential for amidase function to cleave PG for cell separation [43]. To

investigate a possible link between PBP4 and this complex we further dissected the

localization requirement of PBP4. PBP4 localized at division sites in a Δ*envC* strain, indicating

that EnvC is not needed for PBP4 localization. However, PBP4 failed to localize in a Δ*ftsEX*

strain (Fig. 8). To verify whether FtsE and FtsX were required for PBP4 localization we

constructed a Δ*ftsE* strain. The *ftsE* and *fts*X genes reside in an operon and the expression of

*fts*X requires the *fts*E gene [79]. We therefore constructed a Δ*ftsE* strain with a weakened

p*Trc*99A promoter placed upstream of *fts*X to ensure sufficient expression of FtsX. The

resulting strain, XL36 grew with a mild filamentous phenotype in rich medium that could be

complemented by the expression of FtsE(wt) (Fig S9). PBP4 did not localized at septal

positions in the Δ*fts*E strain XL36 (Fig. 8A) but localized normally when FtsE was expressed to

complement this strain (Fig 8B). This suggests that FtsE and FtsX enhance the septal

localization of PBP4. Interestingly, PBP4 localized at division sites in a strain lacking the three

amidases AmiA, AmiB and AmiC (Fig. S10) and in a strain lacking essential components of the

Tol-Pal system, which is involved in the coordination of OM constriction and PG invagination

during cell division (Fig. S10).

To determine whether the activity of FtsE was needed for the localization of PBP4, we

expressed three inactive FtsE variants that had been shown to localize to midcell

{Arends:2009id}. The K41Q and D162A versions of FtsE cannot bind ATP and FtsE(E163A)

binds ATP but cannot hydrolyze it. PBP4 was able to localize at the division site in cells that

expressed one of these inactive variants of FtsE as the only copy (Fig. 8B) showing that the

activity of FtsE is not required for the localization of PBP4.

PBP4 affects the timing of the divisome assembly. To measure the timing of divisome

in various strain backgrounds, we grew Δ*dacB* cells carrying either an empty plasmid or

plasmids constitutively expressing PBP4 or PBP4 versions to steady state in minimal glucose

medium at 28°C. Cells were fixed, divided into three aliquots and labeled with antibodies

against FtsZ, PBP4 or FtsN. We determined the timing of the arrival of these proteins at

midcell by analyzing fluorescence signals of several thousands of cells [80]. In the parental

strain FtsZ and FtsN started to localize at midcell at about 26 % and 49%, respectively, of the

cell cycle age. Similarly, in the Δ*dacB* strain expressing wild-type PBP4 from plasmid FtsZ and

FtsN started to localize at midcell at about 28% and 49%, respectively, of the cell division cycle

age (Fig. 9E). Expression of inactive PBP4 versions affected the timing of the divisome

assembly differently. Especially the Δ*dacB* cells expressing PBP4(D155A) from plasmid

became slightly filamentous (Fig. 9B), assembled the divisome very early and showed a longer

cell division period (Fig. 9E). Next, we measured the timing of the divisome assembly in the

absence of PBP4 in the ΔdacB strain and in its parental BW25113. A difference was observed

between the two strains, which was not significant due to the large variation in the

localization timing of FtsZ and FtsN in both strains (Fig 9F). We inferred that BW25113 is not

able to reach a reproducible steady state, possibly due to a known frameshift in the *rph* gene,

which leads to pyrimidine starvation and hence irregular DNA replication [81]. The uracil and

thymine that we add as standard to the minimal medium for this strain is evidently unable to

fully compensate the deficiency. To solve this possible issue, the *dacB* gene was deletion from

the wild-type strain MC4100 and the cells were grown to steady state in minimal medium in

the absence of an antibiotic. The results reproduced the difference between the Δ*dacB* and

wild-type strains but now with a significance of P = 0.0001 (Fig. 9F). No significant difference

between the Δ*dacB* and its parental in other parameters that could account for a change in

cell cycle such as mass doubling time, cell length, diameter or the percentage of constricting

cells was detected (Table S1). However, FtsZ started to localize (t_0) 8 min earlier in the Δ*dacB*

strain and reached 50% of its maximum intensity ($t_{1/2}$) 9 min earlier compared to the parental

strain. FtsN localized 8.4 min (t_0) and 5.7 min ($t_{1/2}$) earlier (see for an explanation of t_0 and

$t_{1/2}$ Fig. 9D). These results show that PBP4 profoundly affects the timing of divisome assembly.

DISCUSSION

E. coli contains a large number of seemingly redundant PG hydrolases, which possibly fine tune peptidoglycan biogenesis and remodelling in response to environmental parameters. Apart from their enzymatic activities, little is known about the actual function of most PG hydrolases. Here we report that the DD-endopeptidase PBP4 localizes at midcell during septal PG synthesis in *E. coli* and is important for the timing of the assembly of the division machinery.

PBP4 localises at midcell during cell division in an FtsEX dependent manner. PBP4 localizes predominantly at midcell during cell division in an FtsEX dependent manner. Inactive PBP4 variants were able to localize at midcell indicating that substrate hydrolysis is not a requirement for localization. However, a version of PBP4 lacking the non-catalytic domain 3 (PBP4ΔD3), which is required for activity but not β-lactam binding, did not localize to mid-cell. From these results we infer that PBP4 localizes to midcell either by recognizing its substrate (without the need to hydrolyse it) and/or by protein-protein interactions. PBP4 molecules appear to arrive early and concomitantly with the assemblage of the proto-ring and may associate to preseptal PG synthesis, which is consistent with our observation that PBP4 affects the timing of divisome assembly.

PBP4 might be involved in preseptal PG synthesis. PBP4 interacts with NlpI, PBP1A and LpoA and these proteins are able to form a ternary complex ([3] and (Fig. 4). PBP4ΔD3 interacted with PBP1A and LpoA but not with NlpI, confirming differential PBP4 binding sites of PBP1A, LpoA and NlpI. The interaction with PBP1A and/or LpoA may contribute to the recruitment of PBP4 at midcell, as the amount of PBP4 at midcell was slightly reduced in Δ*mrcA* and Δ*lpoA* strains. PBP1A and its partner LpoA are involved in preseptal PG synthesis [33,80]. Although somewhat weaker, PBP4 still localizes at putative division sites in cell

filaments generated with aztreonam that have lost septal synthesis but not preseptal

synthesis activity. Possibly, some PBP4 enriches at preseptal PG synthesis sites together with

PBP1A/LpoA, which localize through the interaction of PBP1A with ZipA [33], and preseptal

PG synthesis provides substrates for PBP4. The interaction between NlpI and PBP4 could

function to sequester PBP4 in the lateral wall to prevent its interaction with potential

substrate (Fig. 10).

Why does the loss of PBP4 affects the timing of divisome assembly? The absence of

PBP4 interferes with the timing of the divisome assembly. In most bacteria studied thus far,

the Z-ring with its associated early division proteins localizes first to prepare the future cell

division site, perhaps by generating the border between the side wall and the future septum

[33,68,82]. After a time-delay, the proteins that synthesize the bulk of the septal PG arrive

and septation starts [48–50]. In the absence of PBP4, the assembly of the divisome starts

earlier in the cell cycle and it takes more time. This change of divisome assembly could be

caused by direct or indirect effects. PBP4 could be directly needed to open the PG layer to

allow insertion of new septal material or to remodel the nascent PG to allow new insertions.

Alternatively, the absence of PBP4 might enable NlpI to bind other EPases, coordinating their

activity for remodelling the PG layer during septation. However, the robust localization of

PBP4 at midcell suggests a direct function for PBP4 in cleavage of existing glycan crosslinks to

allow the insertion of new material. Since deletion of PBP4 does not have dramatic effects,

another endopeptidase must be involved as well or be able to replace PBP4 to a large extend.

We hypothesise that during elongation sufficient PBP4 molecules are retained by NlpI in the

lateral wall above the PG layer to avoid promiscuous PBP4 activity (Fig. 10). This is consistent

with the observations that the absence of NlpI resulted in a reduced amount of PBP4 present

in the cells (Fig. S6 and [3]). NlpI is not enhanced at midcell [3] and therefore we do not

anticipate a role for NlpI in the retention of PBP4 at midcell. PBP4 might be initially recruited

to midcell through interactions with PBP1A and LpoA, and the FtsEX-EnvC complex, likely

directing PBP4's activity already to preseptal sites (Fig. 10). Once septal peptidoglycan

synthesis starts by the mature divisome, more substrate should become available for PBP4

facilitating its accumulation at midcell, which we observed in the immunolocalization

experiments. Hence, PBP4 could initially remove crosslinks to help AmiA and AmiB to

generate denuded glycan strands. These would facilitate the accumulation of FtsN, which is

required to initiate septal PG synthesis. Alternatively, PBP4 could simply remove old glycan

strands during preseptal PG synthesis. Because of its impact on the timing of the divisome

assembly, the first option seems to be more likely. Whether the function of PBP4 remains the

same during septation will need further investigation.

MATERIALS AND METHODS
391
Culturing conditions. *E. coli* K12 cells were grown to steady state [80] in glucose minimal

medium containing 6.33 g of $K_2HPO_4.3H_2O$, 2.95 g of KH_2PO_4, 1.05 g of $(NH_4)_2$, 0.10 g of

$MgSO_4.7 H_2O$, 0.28 mg of $FeSO_4.7 H_2O$, 7.1 mg of $Ca(NO_3)_2.4 H_2O$, 4 mg of thiamine and 4 g of

glucose per L. For strain MC4100 and its derivatives 50 µg lysine (Gb1) and for BW25114 and

MG1655 derivatives 2 µg uracil, 20 µg thymidine, 50 µg arginine and glutamine (Gb4) per liter

pH 7.0 at 28°C or 37°C were added. Absorbance was measured at 450 nm with a 300-T-1

spectrophotometer (Gilford Instrument Laboratories Inc.). Alternatively, cells were grown in

TY (5g yeast extract, 10 g bacto trypton and 85 mM NaCl, pH 7.0), or LB, which is TY with 170

mM NaCl and absorbance was measured at 600 nm.

Strain construction. Strain XL35, the FtsE partial deletion strain, was constructed as

follows: Primers priXL221 and priXL222 were used to amplify the upstream homologous

sequences (*UHS*) from the *ftsE* start codon, and primers priXL223 and priXL224 were used to

amplify the downstream homologous sequence after the FtsE 151[th] amino acid (the potential

promoter of *ftsX*), from the MC4100 genome. Primers priXL51 and priXL54 were used to

amplify the kanamycin resistance cassette from plasmid pKD4 [83]. The amplified overlap PCR

product (*UHS-FRT-KAN-FRT-DHS*) with primers priXL221 and priXL224 was used to construct

the recombinant strain XL34 in which the first 150 amino acids encoding sequence of *ftsE* was

deleted [83]. Subsequently, the kanamycin selection cassette was removed with plasmid

pCP20 [83] yielding XL35.

Strain XL36, the *ftsE* clean deletion strain, was constructed similarly as XL35. Primers

priXL221 and priXL222 were used to amplify the upstream homologous sequences from the

ftsE start codon, and primers priXL225 and priXL226 were used to amplify the downstream

homologous sequences after the *ftsE* stop codon, from the MC4100 genome. In order to

restore the FtsX expression, a *pTrc99Adown* promoter [84] was amplified from plasmid

pSAV057 [85] with primers priXL77 and priXL227 and a chloramphenicol selection cassette

was amplified from plasmid pKD3 with primers priXL52 and priXL54. The amplified overlap

PCR product (*UHS-FRT-CAM-FRT-DHS*) with primers priXL221 and priXL226 was used to

construct the recombinant strain XL33 in which the entire *ftsE* coding sequence was deleted

and FtsX was expressed from the chromosomally encoded p*Trc99Adown* promoter.

Subsequently, the kanamycin selection cassette was removed with plasmid pCP20 yielding

XL36 (Table 1).

Plasmid construction. The *ftsE* gene was amplified from the chromosome of LMC500

[86] using the forward primer priXL187 and the reverse primer priXL188, and sequenced to

characterize the mutation. The wild-type *ftsE* gene was cloned into pSAV057, p15A origin and

Cam[R] [85] to produce the plasmid pXL135. *FtsE* was expressed under the control of the

p*Tcr*99A down promoter. QuickChange site directed mutagenesis (Agilent technologies, Santa

Clara, CA) and Gibson assembly [87] approaches were applied afterwards to construct the

inactive *ftsE* mutant plasmids pXL137, pXL138 and pXL139, using the primers listed in Table 2.

To construct the PBP4 expression plasmids, the wild-type *dacB* gene was firstly

amplified from the *E. coli MG1655* genome using primers priXL244 and priXL245, and

subsequently cloned into plasmid pSAV057 and pSAV057-dsba^{ss}-mCherry-PBP5 (pNM010 [20])

with *EcoRI* and *HindIII*, to generate the PBP4 plasmids without and with mCherry fusion.

Similarly, QuickChange site directed mutagenesis and Gibson assembly approaches were

applied afterwards to construct the inactive *dacB* mutant plasmids from these two wild-type

plasmids, using the primers listed in Table 2. Tre1 and Tre1E415Q were expressed from the

arabinose inducible promoter of pBAD18 as described [75].

Immunolabeling. After reaching steady state for minimal glucose grown cells or the

desired OD for rich medium grown cells, the cells were fixed for 15 min by addition of a

mixture of formaldehyde (f. c. 2.8%) and glutaraldehyde (f. c. 0.04%) to the cultures in the

shaking water bath and immunolabeled as described [88] with Rabbit polyclonal antibodies

against PBP4, NlpI [3] preabsorbed against Δ*dacB* or Δ*nlpI*, strains, respectively, or against

FtsZ or FtsN [48]. As secondary antibody, donkey anti-rabbit conjugated to Cy3 or to Alexa488

(Jackson Immunochemistry, USA) diluted 1:300 in blocking buffer (0.5% (wt/vol) blocking

reagents (Boehringer, Mannheim, Germany) in PBS) was used, and the samples were

incubated for 30 minutes at 37°C. For immunolocalization, cells were immobilized on 1%

agarose in PBS slabs coated object glasses as described [76] and photographed with an Orca

Flash 4.0 (Hamamatsu, Japan) CCD camera mounted on an Olympus BX-60 (Japan)

fluorescence microscope through a 100x/*N.A.* 1.35 oil objective. Images were taken using the

program ImageJ with MicroManager (https://www.micro-manager.org).

SIM images were obtained with a Nikon Ti Eclipse microscope (Japan) and captured

using a Hamamatsu Orca-Flash 4.0 LT camera. Phase contrast images were acquired with a

Plan APO 100x/1.45 Ph3 oil objective. SIM images were obtained with a SR APO TIRF

100x/1.49 oil objective, using 3D-SIM illumination with a 561nm laser, and were

reconstructed with Nikon-SIM software using the values 0.20-0.25-0.20 for the parameters

Illumination Modulation Contrast (IMC), High Resolution Noise suppression (HNS) and Out of

focus Blur Suppression (OBS), respectively.

Image analysis. Phase contrast and fluorescence images were combined into

hyperstacks using ImageJ (http://imagej.nih.gov/ij/) and these were linked to the project file

of Coli-Inspector running in combination with the plugin ObjectJ

(https://sils.fnwi.uva.nl/bcb/objectj/). The images were scaled to 15.28 pixel per μm. The

fluorescence background has been subtracted using the modal values from the fluorescence

images before analysis. Slight misalignment of fluorescence with respect to the cell contours

as found in phase contrast was corrected using Fast-Fourier techniques as described [80].

Data analysis was performed as described [80]. In brief, midcell was defined as the central

part of the cell comprising 0.8 μm of the axis. From either cell part, midcell and remaining cell,

the volume, the integrated fluorescence, and thus the concentration of fluorophores was

calculated. The difference of the two concentrations is multiplied with the volume of midcell.

It yields FCPlus (surplus of fluorescence) at midcell. For age calculation, all cell lengths are

sorted in ascending order. Then the equation

$age = \ln(1 - 0.5 * rank / (nCells - 1)) / \ln(0.5)$

is used, where *rank* is a cell's index in the sorted array, *nCells* is the total number of cells, and

age is the cell's age expressed in the range 0 .. 1.

Ni^{2+}-NTA pulldown assays. Nickel-Nitrilotriacetic acid (Ni-NTA) beads (100μl) were

pre-equilibrated with dH$_2$O and binding buffer (10 mM HEPES/NaOH, 10 mM MgCl$_2$, 150 mM

NaCl, 0.05% Triton X-100, pH 7.5) by centrifugation at 4000 × g, 4 min at 4°C, and then

incubated with proteins of interest. Equimolar amounts (1-2 μM) of His$_6$-tagged and untagged

proteins were incubated alone (control) and in combination, for 10 min at 4°C in a 200 μl

reaction volume, containing 10 mM binding buffer. An aliquot of this mixture is taken as an

'applied' sample. The protein samples were then added to pre-equilibrated Ni-NTA beads and

incubated overnight on a spinning plate at 4°C with 1.3 ml of binding buffer.

Beads were centrifuged at 4000 × g, 4 min, 4°C and washed a further 4-6 times with 1

ml of washing buffer (binding buffer with 30 mM Imidazole) before re-suspending in 250 μl

washing buffer and transferring beads to Proteus spin columns (Generon), centrifuging as

described above. Bound proteins were eluted by the addition of 50 μl of SDS-PAGE loading

buffer and boiling at 100°C for 5 min. Spin columns were centrifuged a final time at 1500 × g,

5 min, RT, to collect 'eluted' protein before separation by SDS-PAGE. Eluted samples were run

alongside applied samples for comparison.

Microscale thermophoresis. Microscale thermophoresis (MST) is an immobilisation-

free method which allows the detection of biomolecular interactions in solution. This

technique is based on the specific directed movement of a protein along a heat gradient

(thermophoresis), an observation first reported by Carl Ludwig in 1856 [89]. The

thermophoresis of a protein changes upon ligand binding due to one or more changes to size,

charge and/or hydration shell. In MST, a localized heat gradient is initiated by an IR-laser

(wavelength 1470 nm). A protein of interest is fluorescently labelled and the change in its

thermophoretic mobility, in the presence of an unlabelled ligand, is measured and expressed

as a change in the normalised fluorescence (FNorm).

Proteins of interest (10-20 µM) were fluorescently labelled with an amine

reactive dye (NT-647- N-hydroxysuccinimide (NHS)), cysteine reactive dye (NT-647 maleimide)

or a histidine reactive dye (NT-647-Tris-NTA), according to manufacturer's instructions

(Nanotemper). Fluorescently labelled proteins were diluted to an appropriate concentration

and a fixed concentration was titrated against a two-fold serial dilution of unlabelled ligand,

across 16 samples, in MST running buffer (25 mM HEPES/NaOH, 150 mM NaCl, 0.05% Triton

X-100, pH 7.5).

MST measurements were carried out as described in [90], using standard or premium

capilaries on a Monolith NT.115^{TM} MST machine (Nanotemper). Binding curves and Kinetic

parameters were plotted and estimated using manufacturer provided software (NT Analysis

1.5.41 and MO. Affinity Analysis (x64)). Capillary scans were carried out prior to all

measurements to check for consistent fluorescence counts, confirming that any subsequent

change in fluorescence was due to ligand binding and not due to inaccurate pipetting or

adsorption and dilution effects.

SDS-denaturation (SD) test. In instances where capillary scans showed a ligand

concentration dependant change in raw fluorescence, we investigated whether this was a

property of the binding interaction by carrying out an SD-test.

Samples (10 µL) with the highest and lowest concentration of unlabelled ligand were

centrifuged 10 000 × g, 5 min, RT and mixed 1:1 volume ratio with SD-test buffer (40 mM DTT,

4% SDS). Mixtures were boiled at 100 °C for 10 min, to abolish ligand binding, before being

spun down and subjected to another capillary scan. If the fluorescence between samples that

contained the highest and lowest concentration of ligand, after SDS treatment, were now

back to ± 10% of each other, the initial observations were a property of ligand binding, and a

binding curve was plotted from the raw fluorescence data. If the ligand concentration

dependent change in fluorescence was still observed, then this indicated that the

fluorescently labelled protein was aggregated, and assay and buffer conditions were

optimised accordingly

***In vivo* cross-linking / co-immunoprecipitation assays.** Method is described in, and

adapted from [91]. An overnight culture of *E. coli* BW25113 cells and an appropriate mutant

strain was used to inoculate 150 ml of Lennox LB (Fisher Scientific) and was cultivated to an

OD_{578} of 0.5-0.6 at 37°C before harvesting by centrifugation (4500 × g, 4°C, 25 min). Cells were

resuspended in 6 ml of CL buffer 1 (50 mM NaH_2PO_4, 20% sucrose, pH 7.4). The amine reactive

cross-linker, DTSSP (3,3'-dithiobis (sulfosuccinimidylpropionate) (ThermoFisher), was freshly

dissolved (20 mg/ml in dH_2O) and added to the isolated cell suspension and incubated at 4°C

with agitation for 1 h. Cross-linked cells were then harvested by centrifugation (4500 × g, 4°C,

25 min) and resuspended in 6 ml CL buffer 2 (100 mM Tris/HCl, 10 mM $MgCl_2$, 1 M NaCl, pH

7.5). DNase, protease inhibitor cocktail and phenylmethylsulfonly fluoride were added prior

to sonication at low levels before ultracentrifugation of the lysate (140,000 × g, 4°C, 1 h). The

membrane pellet was resuspended in 2.5 ml of CL buffer 3 (25 mM Tris/HCl, 10 mM $MgCl_2$, 1

M NaCl, 1% Triton X-100, 20% glycerol, pH 7.5) and the solubilised membrane extracted o/n

with stirring at 4°C.

Samples were ultracentrifuged (140,000 × *g*, 4°C, 1 h) to remove debris, before

removing 2 × 1.2 ml of each supernatant to be subsequently diluted with 0.6 ml of CL buffer

4 (75 mM Tris/HCl, 10 mM $MgCl_2$, 1 M NaCl, pH 7.5). One sample was incubated with an

optimised concentration of specific antibody with the other used as a negative control. Both

samples were incubated at 4°C with agitation for 5 h. For the isolation of antibodies, and thus

cross-linked interaction partners, 100 µl of protein G-coupled agarose bead resin (Roche)

were washed (2 × CL buffer 4, 2 × CL wash buffer [2:1 CL buffer 3 and CL buffer 4]) and added

to each sample, and incubated o/n at 4°C with agitation.

Samples were centrifuged and the supernatant retained before washing the beads 10

× 1 ml with CL wash buffer. After the final wash, beads were resuspended in 250 µl CL wash

buffer and transferred to 2 ml spin dry columns and centrifuged to isolate the beads. These

were then resuspended in 50 µl of fresh SDS-loading buffer and boiled to elute bound proteins,

and reverse cross-linkage, and were collected by centrifugation (10,000 × g, RT, 5 min).

Supernatant and elution samples were resolved by SDS-PAGE and transferred to a

nitrocellulose membrane by Western blotting to detect for specific interaction partners using

purified antibodies. The secondary antibody used here is Trueblot Anti-Rabbit IgG-HRP

specific for native antibodies.

Peptidoglycan digestion assays. To test for hydrolase activity on PG, 10 µl of sacculi

isolated from strains of interest (usually *E. coli* strains MC1061, BW25113 or D456) were

incubated with 1-10 µM of respective enzymes, 37°C for between 1 h and overnight, as

indicated for respective proteins. The standard reaction conditions were 10 mM HEPES/NaOH,

10 mM $MgCl_2$, 150 mM NaCl, 0.05% Triton X-100, pH 7.5, in 100 µl reaction volume. Following

incubation, samples were boiled at 100°C, 10 min, to terminate reactions before digesting

remaining PG overnight at 37°C, with 1 µM cellosyl. The samples were centrifuged at 10,000

× g for 5 min, RT, to obtain digested muropeptide products in the supernatant.

To test for activity against soluble muropeptides; first, 100 µl of intact sacculi were

incubated overnight at 37°C with 1 µM cellosyl and cellosyl digestion buffer (20 mM $NaPO_4$,

pH 4.8). Next, samples were boiled at 100 °C, 10 min, before centrifugation at 10,000 × g for

5 min, RT, to obtain the soluble muropeptides in the supernatant [92]. Ten µl of the

supernatant was then used as muropeptide substrate for incubating with enzymes of interest,

for the appropriate incubation time, at 37°C. Reactions were then terminated by boiling at

100°C for 10 min.

After digestion of respective substrates, products were reduced with NaBH$_4$, adjusted

to pH 4-5 and separated for analysis by reversed-phase HPLC as described below in HPLC

analysis.

Reduction of muropeptides with sodium borohydride. PG digestion samples were

transferred to 2 ml vials following centrifugation. Muropeptides were reduced in a 1:1 volume

ratio of sodium borohydride buffer (0.5 M sodium borate, pH 9.0) and a small spatula of

sodium borohydride pellets, centrifuging at 3000 × g, 30 min, RT as described in [92]. The

samples were adjusted to pH 4-5 using HPLC grade phosphoric acid before separation and

analysis by reversed-phase HPLC.

Reversed-phase HPLC analysis of muropeptides. Following the protocol of [92],

reduced muropeptides were separated for analysis on reversed-phase HPLC systems with a

Prontosil 120-3-C18-AQ 3 µm reversed-phase HPLC column (Bischoff). A linear gradient of

solvent A (50 mM sodium phosphate, pH 4.31 supplemented with 0.2% NaN$_3$) to 100 %

solvent B (75 mM sodium phosphate, pH 4.95, 15% methanol) over 90 or 180 min was used

to separate muropeptides at 52°C. Unlabelled muropeptides were detected at UV absorbance

205 nm. In assays where [14]C-radiolabelled muropeptides were used, detection was achieved

by flowing scintillation cocktail along with standard buffers to give a radioisotope scintillation

count (radioactivity CPM). Muropeptide profiles were recorded and analysed using Laura

V4.2.11.129 (LabLogic Systems Ltd.).

Spectrophotometric D-alanine release assay (DD-carboxypeptidase assay). This

protocol was adapted from [62]. The carboxypeptidase (CPase) activity of PG hydrolases

results in the release of the terminal D-Ala residue from the pentapeptide stem of PG

precursors. Using UDP-Mur*N*Ac pentapeptide as a substrate, and in this case PBP4, it was

possible to spectrophotometrically measure the release of D-Ala.

Each reaction sample consisted of 200 μl of CPase buffer (50 mM HEPES/NaOH, 10

mM $MgCl_2$, pH 7.6), 3 units of D-amino acid oxidase (Sigma), 6 units of horseradish peroxidase

(HRP) (Sigma), Amplex Red (Sigma), and an optimised concentration of protein.

All constituents of the reaction were added and mixed directly in a quartz cuvette

(Hellma, 10 mM light path, 15 mM centre), before the addition, and brief mixing by pipette,

of purified UDP-Mur*N*Ac pentapeptide (BACWAN, Warwick University) to begin the reaction.

The released D-Ala residues from the CPase activity of PBP4 are oxidatively deaminated by

the action of D-amino acid oxidase to produce pyruvate and hydrogen peroxide (H_2O_2). The

released H_2O_2 is reduced to H_2O by HRP using Amplex Red as an electron donor. Oxidised

Amplex Red produces resorufin, which has an intense pink colour and the production of which

was measured spectrophotometrically using a Cary 100 Bio UV-visible spectrophotometer

(wavelength 555 nm). The change in absorption over 10 min was measured and analysed

using the complementing software.

Circular dichroism. Proteins were dialysed overnight against 10 mM $NaPO_4$, pH 7.5

and concentrated/diluted to 0.4 mg/ml. CD measurements were taken using a Jasco J-810

spectropolarimeter using a wavelength range of 180-250 nm. The average of 10 runs was

taken for each protein with a buffer control subtraction. For a direct comparison, correcting

for the differing amino acid sequences, the collected data were converted to molecular CD

and plotted against wavelength (nm). The resulting CD spectra are compared in Fig. S4 and

show that PBP4 lacking domain 3 is folded, consisting of both α-helices (~190, 208 and 222

nm) and β-sheets (~210 nm).

Analytical ultracentrifugation. Purified PBP4 and PBP4ΔD3 were dialyzed over night

against 25 mM HEPES/NaOH, 150 mM NaCl, pH 7.5, in preparation for AUC experiments. AUC

sedimentation velocity (SV) experiments were carried out in a Beckman Coulter (Palo Alto,

CA, USA) ProteomeLab XL-I analytical ultracentrifuge using absorbance at 280 nm and

interference detection. The AUC runs were executed at the rotation speed of 45,000 rpm and

the temperature of 20°C using an 8-hole AnTi50 rotor and double-sector aluminium-Epon

centerpieces. The sample volume was 400 µl and the sample concentrations ranged between

approximately 0.25 and 1.4 mg/ml. The partial specific volumes ($\bar{v}$) of the proteins were

calculated from their amino acid sequence, using the program SEDNTERP [93]. Sedimentation

velocity boundaries were analyzed using the size-distribution c(s) model implemented in the

program SEDFIT http://www.analyticalultracentrifugation.com [94]. The experimental values

of the sedimentation coefficient were converted to the standard conditions (s20,w), which is

the value of sedimentation coefficient in water at 20°C. The size-distribution peaks were

integrated to obtain the weight-averaged values for sedimentation coefficient and molecular

mass.

The atomic coordinates from the published high-resolution structure of [60] (pdb

accession code 2EX2) were used to calculate the sedimentation coefficient values for the

monomeric and dimeric forms of PBP4 /PBP4ΔD3 using the program SoMo [95]. PBP4

/PBP4ΔD3 crystallographic dimer was built using program PyMol (https://pymol.org/2/).

PG binding assay. To assay for interactions of respective proteins with purified PG, we

used a PG binding protocol as described in [96]. Briefly, ~100 µg purified PG was pelleted by

centrifugation (10,000 × g, 10 min, 4°C) and resuspended in binding buffer (10 mM

Tris/Maleate, 10 mM MgCl$_2$, 50 mM NaCl, pH 6.8). PG was incubated with 10 µg of desired

protein before incubating on ice for 30 min and centrifuging at 10,000 × g, 10 min, 4°C. An

aliquot of the supernatant (S) was taken for SDS-PAGE. The pelleted PG was washed with 200

μl binding buffer by centrifugation as before and the wash supernatant was taken as sample

W for SDS-PAGE. The pelleted PG was then resuspended in 100 μl of 2% SDS and incubated

for 1 h on a stirring plate, 4°C. Mixtures were centrifuged a final time at 10,000 × g, 10 min,

4°C, with the supernatant, now containing any initially bound protein, taken as the pellet

sample (P). A negative control sample with no PG was assayed in parallel to determine that

binding of protein was specific. Samples, along with corresponding controls, were analysed

by SDS-PAGE and visualised by Coomassie staining. Presence of protein band in pellet sample

(P) indicated binding to PG, whilst presence in supernatant sample (S) indicated no binding.

Bocillin binding assay. A fluorescent-bocillin binding assay was used to determine

whether a PBP was purified with a correctly folded active site. Bocillin FL is a commercially

available dye-β-lactam conjugate, synthesised from penicillin V and BODIPY (Molecular

Probes, In., Eugene, Oreg.). Ten μg of purified PBP was incubated for 10 min at 37°C, with 20

ng of Bocillin FL in a 50 μl volume with 10 mM HEPES/NaOH, 10 mM $MgCl_2$, pH 7.5. A negative

control sample was pre-incubated with 1 mM ampicillin for 30 min, 37°C, to block the PBP

active site, prior to incubation with Bocillin FL. After incubation, samples were boiled at 100

°C for 10 min with 40 μl of SDS-PAGE loading buffer and resolved by SDS-PAGE. Fluorescent

signal was observed using Typhoon Fluorescence-imager (Excitation; 488 nm, emission; 520

nm, BP20 PMT 400-600 V), followed by visualisation of gels by Coomassie staining. Proteins

incubated with ampicillin should have no detectable binding of Bocillin FL. For whole cells, a

culture was grown in 5 mL TY and induced with 10 μM IPTG for 3 mass doublings to an OD_{600}

of 0.5 and the cells were pelleted and washed once with PBS (8 g NaCl, 1.8 g Na_2HPO_4, 0.24 g

KH_2PO_4 per liter, adjusted to pH 7.4) at 4°C and resuspended in 100 μl PBS with 5μg/ml Bocilin

FL and incubated for 20 min at RT. To remove the excess Bocillin FL, the samples were pelleted

and resuspended in 100 µl PBS with cOmplete™ Mini Protease Inhibitor Cocktail (Sigma) and

2 U DNAse I (NEB). Then sonicated on ice for 10 sec and boiled 15' at 99°C after adding 25 µL

of SDS-PAGE 5x loading buffer and resolved by SDS-PAGE. Fluorescent signal was observed

using a home-made imager for Midori green, followed by visualisation of gels by Coomassie

staining.

***In vitro* PG synthesis assay with radiolabelled lipid II.** The assay was published by

Bertsche et al., [97]. Lipid II (1.2 nmol, 11,000 dpm) was vacuum dried and dissolved in 5 µl of

methanol. The reaction was performed in a total volume of 50 µl in 25 mM Tris/HCl, 10 mM

$MgCl_2$, 100 mM NaCl, 0.05% Triton X-100, pH 7.5 (buffer B), with 0.5 µM PBP1A, 1 µM LpoA,

1 µM PBP4(S62A), 2 µM NlpI. The reaction was performed for 1 h at 30°C. The pH of the

samples was adjusted to 4.8 prior to boiling for 5 min followed by a digestion with 50 g/ml

cellosyl for 1 h at 37°C and boiling for 5 min.

ACKNOWLEDGEMENTS

We thank David Roper and Smita Chauhan (University of Warwick) for help with the

carboxypeptidase assay, Helen Waller (Newcastle University) for help with circular dichroism

methodology, Lisa Atkinson and Daniela Vollmer (Newcastle University) for preparation of PG

sacculi, and Thomas van Wingen for with help the Tre1 assay. We thank See-Yeun Ting for the

plasmids expressing Tre1 and Tre1E415Q (University of Washington School of Medicine,

Seattle, USA). The work was supported by funding from the Wellcome Trust (101824/Z/13/Z),

to WV). X.L. was supported by the China Scholarship Council fellowship (File

No.201406220123).

J.V., A.L., H.C.L.Y, X.L., X.L., A.S.S., and M.B did most experiments. A.T., M.B., V.W., and T.dB.

designed most experiments. M.B, W.V and T.dB analyzed the data and wrote the manuscript.

All authors critically read the manuscript and provided feedback.

References

691 1. Vollmer W, Blanot D, Pedro MAD. Peptidoglycan structure and architecture. FEMS microbiol. revs. 2008;32: 149–167. doi:10.1111/j.1574-6976.2007.00094.x

693 2. Egan AJF, Errington J, Vollmer W. Regulation of peptidoglycan synthesis and remodelling. Nat. Rev. Microbiol. 2020;18: 446–460. doi:10.1038/s41579-020-0366-3

695 696 3. Banzhaf M, Yau HCL, Verheul J, Lodge A, Kritikos G, Mateus A, et al. Outer membrane lipoprotein NlpI scaffolds peptidoglycan hydrolases within multi-enzyme complexes in Escherichia coli. The EMBO Journal. 2020; e102246. doi:10.15252/embj.2019102246

698 699 4. Gray AN, Egan AJF, Veer ILV, Verheul J, Colavin A, Koumoutsi A, et al. Coordination of peptidoglycan synthesis and outer membrane constriction during *Escherichia coli* cell division. Storz G, editor. eLife. 2015;4: e07118. doi:10.7554/elife.07118

701 702 5. Bertsche U, Kast T, Wolf B, Fraipont C, Aarsman MEG, Kannenberg K, et al. Interaction between two murein (peptidoglycan) synthases, PBP3 and PBP1B, in Escherichia coli. Molecular Microbiology. 2006;61: 675–690. doi:10.1111/j.1365-2958.2006.05280.x

704 6. Born P, Breukink E, Vollmer W. In vitro synthesis of cross-linked murein and its attachment to sacculi by PBP1A from *Escherichia coli*. J. Biol. Chem. 2006;281: 26985–26993. doi:10.1074/jbc.m604083200

706 707 7. Egan AJF, Biboy J, Veer I van't, Breukink E, Vollmer W. Activities and regulation of peptidoglycan synthases. Philosophical transactions of the R.Soc. London Series B, Biolog. Sci. 2015;370. doi:10.1098/rstb.2015.0031

709 8. Pazos M, Vollmer W. Regulation and function of class A Penicillin-binding proteins. Curr. Opin. Microbiol. 2021;60: 80–87. doi:10.1016/j.mib.2021.01.008

711 712 9. Rohs PDA, Buss J, Sim SI, Squyres GR, Srisuknimit V, Smith M, et al. A central role for PBP2 in the activation of peptidoglycan polymerization by the bacterial cell elongation machinery. PLoS Genetics. 2018;14: e1007726. doi:10.1371/journal.pgen.1007726

714 10. Meeske AJ, Riley EP, Robins WP, Uehara T, Mekalanos JJ, Kahne D, et al. SEDS proteins are a widespread family of bacterial cell wall polymerases. Nature. 2016;537: 634–638. doi:10.1038/nature19331

716 717 11. Taguchi A, Welsh MA, Marmont LS, Lee W, Sjodt M, Kruse AC, et al. FtsW is a peptidoglycan polymerase that is functional only in complex with its cognate penicillin-binding protein. Nat. Microbiol. 2019;4: 587–594. doi:10.1038/s41564-018-0345-x

719 720 12. Ploeg R van der, Verheul J, Vischer NOE, Alexeeva S, Hoogendoorn E, Postma M, et al. Colocalization and interaction between elongasome and divisome during a preparative cell division phase in *Escherichia coli.* Mol. Microbiol. 2013;87: 1074–1087. doi:10.1111/mmi.12150

722 13. Datta P, Dasgupta A, Singh AK, Mukherjee P, Kundu M, Basu J. Interaction between FtsW and penicillin-binding protein 3 (PBP3) directs PBP3 to mid-cell, controls cell septation and mediates the formation of

a trimeric complex involving FtsZ, FtsW and PBP3 in mycobacteria. Mol. Microbiol. 2006;62: 1655–1673.
doi:10.1111/j.1365-2958.2006.05491.x

14. Fraipont C, Alexeeva S, Wolf B, Ploeg R van der, Schloesser M, Blaauwen T den, et al. The integral membrane
FtsW protein and peptidoglycan synthase PBP3 form a subcomplex in *Escherichia coli*. Microbiol.
2011;157: 251–259. doi:10.1099/mic.0.040071-0

15. Liu X, Biboy J, Consoli E, Vollmer W, Blaauwen den, T. MreC and MreD balance the interaction between the
elongasome proteins PBP2 and RodA. PLoS Genetics. 2020;16: e1009276.
doi:10.1371/journal.pgen.1009276

16. Pazos M, Peters K, Vollmer W. Robust peptidoglycan growth by dynamic and variable multi-protein
complexes. Curr. Opin. Microbiol. 2017;36: 55–61. doi:10.1016/j.mib.2017.01.006

17. Sarkar SK, Dutta M, Chowdhury C, Kumar A, Ghosh AS. PBP5, PBP6 and DacD play different roles in intrinsic -
lactam resistance of *Escherichia coli.* Microbiology (Reading, England). 2011;157: 2702–2707.
doi:10.1099/mic.0.046227-0

18. Potluri L, Karczmarek A, Verheul J, Piette A, Wilkin J-M, Werth N, et al. Septal and lateral wall localization of
PBP5, the major D,D-carboxypeptidase of *Escherichia coli*, requires substrate recognition and membrane
attachment. Mol. Microbiol. 2010;77: 300–323. doi:10.1111/j.1365-2958.2010.07205.x

19. Hugonnet J-E, Mengin-Lecreulx D, Monton A, Blaauwen T den, Carbonnelle E, Veckerlé C, et al. Factors
essential for L,D-transpeptidase-mediated peptidoglycan cross-linking and β-lactam resistance in
Escherichia coli. eLife. 2016;5: 2006. doi:10.7554/elife.19469

20. Meiresonne NY, Ploeg R van der, Hink MA, Blaauwen T den. Activity-Related Conformational Changes in d,d-
Carboxypeptidases Revealed by In Vivo Periplasmic Förster Resonance Energy Transfer Assay in
Escherichia coli. Søgaard-Andersen L, editor. mBio. 2017;8: e01089-17. doi:10.1128/mbio.01089-17

21. Singh SK, Saisree L, Amrutha RN, Reddy M. Three redundant murein endopeptidases catalyse an essential
cleavage step in peptidoglycan synthesis of *Escherichia coli* K12. Mol. Microbiol. 2012;86: 1036–1051.
doi:10.1111/mmi.12058

22. Meberg BM, Paulson AL, Priyadarshini R, Young KD. Endopeptidase penicillin-binding proteins 4 and 7 play
auxiliary roles in determining uniform morphology of *Escherichia coli.* J. Bacteriol. 2004;186: 8326–8336.
doi:10.1128/jb.186.24.8326-8336.2004

23. Peters K, Kannan S, Rao VA, Biboy J, Vollmer D, Erickson SW, et al. The Redundancy of Peptidoglycan
Carboxypeptidases Ensures Robust Cell Shape Maintenance in *Escherichia coli*. mBio. 2016;7.
doi:10.1128/mbio.00819-16

24. Peters NT, Dinh T, Bernhardt TG. A fail-safe mechanism in the septal ring assembly pathway generated by the
sequential recruitment of cell separation amidases and their activators. J. Bacteriol. 2011;193: 4973–
4983. doi:10.1128/jb.00316-11

25. Heidrich C, Templin MF, Ursinus A, Merdanovic M, Berger J, Schwarz H, et al. Involvement of N-
acetylmuramyl-L-alanine amidases in cell separation and antibiotic-induced autolysis of *Escherichia coli.*
Mol. Microbiol. 2001;41: 167–178.

26. Bernhardt TG, Boer PAJD. The Escherichia coli amidase AmiC is a periplasmic septal ring component exported
via the twin-arginine transport pathway. Mol. Microbiol. 2003;48: 1171–1182.

27. Uehara T, Parzych KR, Dinh T, Bernhardt TG. Daughter cell separation is controlled by cytokinetic ring-
activated cell wall hydrolysis. EMBO J. 2010;29: 1412–1422. doi:10.1038/emboj.2010.36

28. Straaten KE van, Dijkstra BW, Vollmer W, Thunnissen A-MWH. Crystal structure of MltA from *Escherichia coli*
reveals a unique lytic transglycosylase fold. J. Mol. Biol. 2005;352: 1068–1080.
doi:10.1016/j.jmb.2005.07.067

29. Heijenoort J van. Peptidoglycan hydrolases of *Escherichia coli.* Microbiol and Mol. Biol. Revs.: MMBR.
2011;75: 636–663. doi:10.1128/mmbr.00022-11

30. Yunck R, Cho H, Bernhardt TG. Identification of MltG as a potential terminase for peptidoglycan
polymerization in bacteria. Mol.Microbiol. 2015. doi:10.1111/mmi.13258

31. Addinall SG, Lutkenhaus J. FtsZ-spirals and -arcs determine the shape of the invaginating septa in some
mutants of *Escherichia coli*. Mol. Microbiol. 1996;22: 231–237.

32. Raychaudhuri D. ZipA is a MAP-Tau homolog and is essential for structural integrity of the cytokinetic FtsZ
ring during bacterial cell division. EMBO J. 1999;18: 2372–2383. doi:10.1093/emboj/18.9.2372

33. Pazos M, Peters K, Casanova M, Palacios P, VanNieuwenhze M, Breukink E, et al. Z-ring membrane anchors
associate with cell wall synthases to initiate bacterial cell division. Nature Commun. 2018;9: 5090–12.
doi:10.1038/s41467-018-07559-2

34. Galli E, Gerdes K. FtsZ-ZapA-ZapB Interactome of Escherichia coli. J.Bacteriol. 2012;194: 292–302.
doi:10.1128/jb.05821-11

35. Durand-Heredia JM, Yu HH, Carlo SD, Lesser CF, Janakiraman A. Identification and Characterization of ZapC, a
Stabilizer of the FtsZ Ring in *Escherichia coli.* J. Bacteriol. 2011;193: 1405–1413. doi:10.1128/jb.01258-10

36. Hale CA, Shiomi D, Liu B, Bernhardt TG, Margolin W, Niki H, et al. Identification of *Escherichia coli* ZapC
(YcbW) as a component of the division apparatus that binds and bundles FtsZ polymers. J. Bacteriol.
2011;193: 1393–1404. doi:10.1128/jb.01245-10

37. Roach EJ, Wroblewski C, Seidel L, Berezuk AM, Brewer D, Kimber MS, et al. Structure and mutational analyses
of *Escherichia coli* ZapD reveal charged residues involved in FtsZ filament bundling. DiRita VJ, editor. J.
Bacteriol. 2016;198: 1683–1693. doi:10.1128/jb.00969-15

38. Marteyn BS, Karimova G, Fenton AK, Gazi AD, West N, Touqui L, et al. ZapE is a novel cell division protein
interacting with FtsZ and modulating the Z-ring dynamics. mBio. 2014;5: e00022-14.
doi:10.1128/mbio.00022-14

39. Mehla J, Liechti G, Morgenstein RM, Caufield JH, Hosseinnia A, Gagarinova A, et al. ZapG (YhcB/DUF1043), a
novel cell division protein in gamma-proteobacteria linking the Z-ring to septal peptidoglycan synthesis.
J Biol Chem. 2021;296: 100700. doi:10.1016/j.jbc.2021.100700

40. Du S, Pichoff S, Lutkenhaus J. FtsEX acts on FtsA to regulate divisome assembly and activity. Proc. Natl Acad
Sci. USA. 2016;113: E5052-61. doi:10.1073/pnas.1606656113

41. Corbin BD, Wang Y, Beuria TK, Margolin W. Interaction between Cell Division Proteins FtsE and FtsZ. J.
Bacteriol. 2007;189: 3026–3035. doi:10.1128/jb.01581-06

42. Priyadarshini R, Popham DL, Young KD. Daughter Cell Separation by Penicillini-Binding Proteins and
Peptidoglycan Amidases in *Escherichia coli.* J. Bacteriol. 2006;188: 5345–5355. doi:10.1128/jb.00476-06

43. Yang DC, Peters NT, Parzych KR, Uehara T, Markovski M, Bernhardt TG. An ATP-binding cassette transporter-
like complex governs cell-wall hydrolysis at the bacterial cytokinetic ring. Proc. Natl Acad Sci. USA.
2011;108: E1052-60. doi:10.1073/pnas.1107780108

44. Yang DC, Tan K, Joachimiak A, Bernhardt TG. A conformational switch controls cell wall-remodelling enzymes
required for bacterial cell division. Mol. Microbiol. 2012;85: 768–781. doi:10.1111/j.1365-
2958.2012.08138.x

45. Tsang M-J, Yakhnina AA, Bernhardt TG. NlpD links cell wall remodeling and outer membrane invagination
during cytokinesis in Escherichia coli. PLoS Genetics. 2017;13: e1006888.
doi:10.1371/journal.pgen.1006888

46. Serrano CKG, Winkle M, Martorana AM, Biboy J, Morè N, Moynihan P, et al. ActS activates peptidoglycan
amidases during outer membrane stress in *Escherichia coli.* Mol Microbiol. 2021.
doi:10.1111/mmi.14712

47. Mueller EA, Iken AG, Öztürk MA, Winkle M, Schmitz M, Vollmer W, et al. The active repertoire of *Escherichia*
coli peptidoglycan amidases varies with physiochemical environment. Mol Microbiol. 2021;116: 311–
328. doi:10.1111/mmi.14711

48. Aarsman MEG, Piette A, Fraipont C, Vinkenvleugel TMF, Nguyen-Distèche M, Blaauwen T den. Maturation of
the *Escherichia coli* divisome occurs in two steps. Mol. Microbiol. 2005;55: 1631–1645.
doi:10.1111/j.1365-2958.2005.04502.x

49. Perez AJ, Cesbron Y, Shaw SL, Villicana JB, Tsui H-CT, Boersma MJ, et al. Movement dynamics of divisome
proteins and PBP2x:FtsW in cells of *Streptococcus pneumoniae*. Proc. Natl Acad Sci. USA. 2019;116:
3211–3220. doi:10.1073/pnas.1816018116

50. Monteiro JM, Pereira AR, Reichmann NT, Saraiva BM, Fernandes PB, Veiga H, et al. Peptidoglycan synthesis
drives an FtsZ-treadmilling-independent step of cytokinesis. Nature. 2018;554: 528–532.
doi:10.1038/nature25506

51. Tsang M-J, Bernhardt TG. A role for the FtsQLB complex in cytokinetic ring activation revealed by an ftsL allele
that accelerates division. Mol. Microbiol. 2014. doi:10.1111/mmi.12905

52. Bernard CS, Sadasivam M, Shiomi D, Margolin W. An altered FtsA can compensate for the loss of essential cell
division protein FtsN in *Escherichia coli*. Mol. Microbiol. 2007;64: 1289–1305. doi:10.1111/j.1365-
2958.2007.05738.x

53. Boes A, Olatunji S, Breukink E, Terrak M. Regulation of the Peptidoglycan Polymerase Activity of PBP1b by
Antagonist Actions of the Core Divisome Proteins FtsBLQ and FtsN. Blaauwen T den, Salama NR, editors.
mBio. 2019;10: e01912-18. doi:10.1128/mbio.01912-18

54. Blaauwen T den, Luirink J. Checks and Balances in Bacterial Cell Division. Mbio. 2019;10.
doi:10.1128/mbio.00149-19

55. Korat B, Mottl H, Keck W. Penicillin-binding protein 4 of *Escherichia coli*: molecular cloning of the *dacB* gene,
controlled overexpression, and alterations in murein composition. Mol. Microbiol. 1991;5: 675–684.

56. Harris F, Brandenburg K, Seydel U. Investigations into the mechanisms used by the C-terminal anchors of
Escherichia coli penicillin-binding proteins 4, 5, 6 and 6b for membrane interaction. Eur.J. Biochem.
2002. doi:10.1046/j.1432-1033.2002.03295.x

57. Pedro MAD, Schwarz U. Heterogeneity of newly inserted and preexisting murein in the sacculus of *Escherichia*
coli. Proceedings of the Proc. Natl Acad Sci. USA. 1981;78: 5856–5860.

58. Iwaya M, Strominger JL. Simultaneous deletion of D-alanine carboxypeptidase IB-C and penicillin-binding
component IV in a mutant of E*scherichia coli* K12. Proc. Natl Acad Sci. USA. 1977;74: 2980–2984.
doi:10.1073/pnas.74.7.2980

59. Nelson DE, Young KD. Contributions of PBP 5 and DD-carboxypeptidase penicillin binding proteins to
maintenance of cell shape in *Escherichia coli*. J. Bacteriol. 2001;183: 3055–3064.
doi:10.1128/jb.183.10.3055-3064.2001

60. Kishida H, Unzai S, Roper DI, Lloyd A, Park S-Y, Tame JRH. Crystal Structure of Penicillin Binding Protein 4
(*dacB)* from *Escherichia coli*, both in the Native Form and Covalently Linked to Various Antibiotics.
Biochemistry. 2006;45: 783–792. doi:10.1021/bi051533t

61. Thunnissen MMGM, Fusetti F, Boer BD, Dijkstra BW. Purification, crystallisation and preliminary X-ray analysis
of penicillin binding protein 4 from *Escherichia coli,* a protein related to class a β-lactamases. J. Mol. Biol.
1995;247: 149–153. doi:10.1006/jmbi.1994.0128

62. Clarke TB, Kawai F, Park S-Y, Tame JRH, Dowson CG, Roper DI. Mutational analysis of the substrate specificity
of *Escherichia coli* penicillin binding protein 4. Biochemistry. 2009;48: 2675–2683.
doi:10.1021/bi801993x

63. Hocking J, Priyadarshini R, Takacs CN, Costa T, Dye NA, Shapiro L, et al. Osmolality-Dependent Relocation of
Penicillin-Binding Protein PBP2 to the Division Site in *Caulobacter crescentus*. J.Bacteriol. 2012;194:
3116–3127. doi:10.1128/jb.00260-12

64. Consoli E, Collet J-F, Blaauwen T den. The *Escherichia coli* Outer Membrane β-Barrel Assembly Machinery
(BAM) Anchors the Peptidoglycan Layer by Spanning It with All Subunits. Int. J. Mol. Sci. 2021;22: 1853.
doi:10.3390/ijms22041853

65. Liu X, Meiresonne NY, Bouhss A, Blaauwen T den. FtsW activity and lipid II synthesis are required for
recruitment of MurJ to midcell during cell division in *Escherichia coli.* Mol. Microbiol. 2018;109: 855–
884. doi:10.1111/mmi.14104

66. Alcorlo M, Dik DA, Benedetti SD, Mahasenan KV, Lee M, Domínguez-Gil T, et al. Structural basis of denuded
glycan recognition by SPOR domains in bacterial cell division. Nature Commun.. 2019;10: 5567–13.
doi:10.1038/s41467-019-13354-4

67. Sykes RB, Bonner DP, Bush K, Georgopapadakou NH. Azthreonam (SQ 26,776), a synthetic monobactam
specifically active against aerobic gram-negative bacteria. Antimicrob Agents Ch. 1982;21: 85–92.
doi:10.1128/aac.21.1.85

68. Nanninga N. Cell division and peptidoglycan assembly in *Eschenchia coli.* Mol. Microbiol. 1991;5: 791–795.
doi:10.1111/j.1365-2958.1991.tb00751.x

69. Pedro MAD, Quintela JC, Höltje JV, Schwarz H. Murein segregation in *Escherichia coli*. J. Bacteriol. 1997;179:
2823–2834.

70. Banzhaf M, Saparoea B van den B van, Terrak M, Fraipont C, Egan A, Philippe J, et al. Cooperativity of
peptidoglycan synthases active in bacterial cell elongation. Mol. Microbiol. 2012;85: 179–194.
doi:10.1111/j.1365-2958.2012.08103.x

71. Jean NL, Bougault CM, Lodge A, Derouaux A, Callens G, Egan AJF, et al. Elongated structure of the outer-
membrane activator of peptidoglycan synthesis LpoA: implications for PBP1A stimulation. Structure
(London, England : 1993). 2014;22: 1047–1054. doi:10.1016/j.str.2014.04.017

72. Jean NL, Bougault C, Derouaux A, Callens G, Vollmer W, Simorre J-P. Backbone and side-chain (1)H, (13)C, and
(15)N NMR assignments of the N-terminal domain of *Escherichia coli* LpoA. Biomolecular NMR
assignments. 2015;9: 65–69. doi:10.1007/s12104-014-9546-2

73. Typas A, Banzhaf M, Saparoea B van den B van, Verheul J, Biboy J, Nichols RJ, et al. Regulation of
peptidoglycan synthesis by outer-membrane proteins. Cell. 2010;143: 1097–1109.
doi:10.1016/j.cell.2010.11.038

74. Goehring NW, Beckwith J. Diverse paths to midcell: assembly of the bacterial cell division machinery. Current
Biology : CB. 2005;15: R514-26. doi:10.1016/j.cub.2005.06.038

75. Ting S-Y, Bosch DE, Mangiameli SM, Radey MC, Huang S, Park Y-J, et al. Bifunctional Immunity Proteins
Protect Bacteria against FtsZ-Targeting ADP-Ribosylating Toxins. Cell. 2018;175: 1380-1392.e14.
doi:10.1016/j.cell.2018.09.037

76. Koppelman C-M, Aarsman MEG, Postmus J, Pas E, Muijsers AO, Scheffers D-J, et al. R174 of *Escherichia coli*
FtsZ is involved in membrane interaction and protofilament bundling, and is essential for cell division.
Mol. Microbiol. 2004;51: 645–657.

77. Pichoff S, Du S, Lutkenhaus J. Disruption of divisome assembly rescued by FtsN-FtsA interaction in *Escherichia*
coli. Proc. Natl Acad Sci. USA. 2018;115: E6855–E6862. doi:10.1073/pnas.1806450115

78. Du S, Henke W, Pichoff S, Lutkenhaus J. How FtsEX localizes to the Z ring and interacts with FtsA to regulate
cell division. Mol. Microbiol. 2019. doi:10.1111/mmi.14324

79. Arends SJR, Kustusch RJ, Weiss DS. ATP-binding site lesions in FtsE impair cell division. J. Bacteriol. 2009;191:
3772–3784. doi:10.1128/jb.00179-09

80. Vischer NOE, Verheul J, Postma M, Saparoea B van den B van, Galli E, Natale P, et al. Cell age dependent
concentration of *Escherichia coli* divisome proteins analyzed with ImageJ and ObjectJ. Front. Microbiol.
2015;6: 586. doi:10.3389/fmicb.2015.00586

81. Jensen KF. The Escherichia coli K-12 "wild types" W3110 and MG1655 have an rph frameshift mutation that
leads to pyrimidine starvation due to low pyrE expression levels. J Bacteriol. 1993;175: 3401–3407.
doi:10.1128/jb.175.11.3401-3407.1993

82. Potluri L-P, Kannan S, Young KD. ZipA is required for FtsZ-dependent preseptal peptidoglycan synthesis prior
to invagination during cell division. J. Bacteriol. 2012;194: 5334–5342. doi:10.1128/jb.00859-12

83. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR
products. Proc. Natl Acad Sci. USA. 2000;97: 6640–6645. doi:10.1073/pnas.120163297

84. Chen JC, Weiss DS, Ghigo J-M, Beckwith J. Septal Localization of FtsQ, an Essential Cell Division Protein in
Escherichia coli. J Bacteriol. 1999;181: 521–530. doi:10.1128/jb.181.2.521-530.1999

85. Alexeeva S, Gadella TWJ, Verheul J, Verhoeven GS, Blaauwen T den. Direct interactions of early and late
assembling division proteins in *Escherichia coli* cells resolved by FRET. Mol. Microbiol. 2010;77: 384–398.
doi:10.1111/j.1365-2958.2010.07211.x

86. Taschner PE, Huls PG, Pas E, Woldringh CL. Division behavior and shape changes in isogenic *ftsZ, ftsQ, ftsA,*
pbpB, and *ftsE* cell division mutants of *Escherichia coli* during temperature shift experiments. J Bacteriol.
1988;170: 1533–1540. doi:10.1128/jb.170.4.1533-1540.1988

87. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. Enzymatic assembly of DNA molecules
up to several hundred kilobases. Nat. Chem.Biol. 2009;6: 343–345. doi:10.1038/nmeth.1318

88. Buddelmeijer N, Aarsman M, Blaauwen T den. Immunolabeling of proteins in situ in *Escherichia coli* K12
strains. Bio-protocol. 2013;3: 1–5.

89. Ludwig C. Diffusion between unequally heated regions of initially uniform solutions. Sitzungsber Akad Wiss
Wien Math-Naturwiss K1. 1856;20: 593.

90. Jerabek-Willemsen M, Wienken CJ, Braun D, Baaske P, Duhr S. Molecular interaction studies using microscale
thermophoresis. Assay and drug develop. technol. 2011;9: 342–353. doi:10.1089/adt.2011.0380

91. Müller P, Ewers C, Bertsche U, Anstett M, Kallis T, Breukink E, et al. The essential cell division protein FtsN
interacts with the murein (peptidoglycan) synthase PBP1B in *Escherichia coli.* J. Biol. Chem. 2007;282:
36394–36402. doi:10.1074/jbc.m706390200

92. Glauner B. Separation and quantification of muropeptides with high-performance liquid chromatography.
Anal. Biochem. 1988;172: 451–464. doi:10.1016/0003-2697(88)90468-x

93. laue T, Shah B, Ridgeway T, Pelletier S. Computer-aided interpretation of analytical sedimentation data for
proteins. Royal Soc. Chem. 1992;90: 125.

94. Schuck P. Size-distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and
lamm equation modeling. Biophys. J. 2000;78: 1606–1619. doi:10.1016/s0006-3495(00)76713-0

95. Brookes E, Demeler B, Rosano C, Rocco M. The implementation of SOMO (SOlution MOdeller) in the
UltraScan analytical ultracentrifugation data analysis suite: enhanced capabilities allow the reliable
hydrodynamic modeling of virtually any kind of biomacromolecule. Eur. Biophys. J.: EBJ. 2010;39: 423–
435. doi:10.1007/s00249-009-0418-0

96. Ursinus A, Ent F van den, Brechtel S, Pedro M de, Höltje J-V, Löwe J, et al. Murein (peptidoglycan) binding
property of the essential cell division protein FtsN from *Escherichia coli.* J. Bacteriol. 2004;186: 6728–
6737. doi:10.1128/jb.186.20.6728-6737.2004

97. Bertsche U, Breukink E, Kast T, Vollmer W. In vitro murein peptidoglycan synthesis by dimers of the
bifunctional transglycosylase-transpeptidase PBP1B from *Escherichia coli*. J. Biol. Chem. 2005;280:
38096–38101. doi:10.1074/jbc.m508646200

98. Casadaban MJ, Cohen SN. Analysis of gene control signals by DNA fusion and cloning in *Escherichia coli*. J.
Mol. Biol. 1980;138: 179–207. doi:10.1016/0022-2836(80)90283-1

99. Ishino F, Jung HK, Ikeda M, Doi M, Wachi M, Matsuhashi M. New mutations fts-36, lts-33, and ftsW clustered
in the mra region of the *Escherichia coli* chromosome induce thermosensitive cell growth and division. J.
Bacteriol. 1989;171: 5523–5530.

100. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of Escherichia coli K-12 in-
frame, single-gene knockout mutants: the Keio collection. Mol. Syst. Biol.. 2006;2: 2006.0008.
doi:10.1038/msb4100050

962 **TABLE 1 Strain and plasmids used in this study.**

Strain	Mutation	Genotype	Source
LMC500 (MC4100)	Wild-type	F⁻, *araD139, Δ(argF-lac)U169, deoC1, flbB5301, lysA1, ptsF25, rbsR, relA1, rpsL150*	[86]
BL21(DE3)	Overexpression strain	F⁻ *ompT, dcm lon hsdS hsdSB(rB-mB-)* λ(DE3)	Novagen
BW25113	Keio background strain	F⁻, *Δ(araD-araB)567, lacZ4787(del)::rrnB-3, LAM-, rph-1, Δ(rhaD-rhaB)568, hsdR514*	[83]
DH5α	Storage strain	F⁻ *endA1 glnV44 thi-1 recA1* φ80*ΔlacZ*ΔM15, λ⁻	Invitrogen
MC1061	Laboratory strain	Δ(*araA-leu*)7697 Δ(*lac*)X74 *galK*16 *galE*15(GalS) λ⁻e14- *mcrA*0 *relA1 rpsL*150(str^R) *spoT1 mcrB1 hsdR2*	[98]
LMC509	FtsZ (ts)	MC4100 *ftsZ*184	[86]
LMC511	FtsA (ts)	MC4100 *ftsA*10	[86]
LMC512	FtsA (ts)	MC4100 *ftsA*1882	[86]
LMC515	FtsE (ts)	MC4100 *ftsE*1181	[86]
LMC531	FtsQ (ts)	MC4100 *ftsQ*(ts)	[86]
LMC2487	FtsW (ts)	JLB17; F⁻ *thr, trp, his, thy, ara, lac, gal, xyl, mtl, rspL, tonA*, *ftsW* amino acid replacement G311D	[99]
LMC510	PBP3 (ts)	MC4100 *ftsI* 2158	[86]
BCB677	ΔPBP1A	BW25113Δ*mrcA*	[100]
BCB678	ΔPBP1B	BW25113Δ*mrcB*	[100]
BCB192	ΔLpoA	BW25113Δ*lpoA*	[100]
LMC2217	ΔLpoB	BW25113Δ*lpoA*	[100]
BCB655	ΔPBP4	BW25113Δ*dacB*	[100]
BCB1433	ΔPBP4	XWL010 MC4100Δ*dacB*	This work
BCB904	ΔNlpI	BW25113Δ*nlpI*	[100]
BCB1363	ΔPBP4ΔLpoA	BW25113Δ*dacB:kan*^R, Δ*lpoA*	This work
BCB744	ΔFtsE	BW25113Δ*ftsE* kan^R CAG60397	This work
BCB746	ΔFtsEX	BW25113Δ*ftsE*X cat^R CAG60399	This work
BCB1081	ΔFtsE 1-150	XL34 LMC500Δ*ftsE* 1-150::*kan-fts*X	This work
BCB849	ΔFtsE 1-150	XL35 LMC500Δ*ftsE* 1-150::*fts*X	This work
BCB1082	ΔFtsE	XL33 LMC500Δ*ftsE*::-cam-p*Trc99A*down *fts*X	This work
BCB850	ΔFtsE	XL36 LMC500Δ*ftsE*::p*Trc99A*down *fts*X	This work

Plasmid	Name	Characteristics	Source
pET21b	pET21b-PBP4(S62A)Δ1-60	Inactive PBP4 lacking residues 1-60, Amp^R	[3]

pET21b	pET21b-His-PBP4(S62A)Δ1-60	For purification of His-PBP4 (active site mutant), Amp[R]	[3]
pET21b	pET21b-PBP4Δ1-60	Native PBP4 lacking residues 1-60, Amp[R]	[3]
pET28a	pET28a-His-LpoA(sol)	Soluble LpoA (LpoAΔ1-27) construct, N-terminal His-tag, Kan[R]	[71]
pET28a	pET28-His-LpoA[N]	Purification of His-LpoA[N], Kan[R]	[71]
pET28a	pET28-His-LpoA[C]	Purification of His-LpoA[C], Kan[R]	[71]
pTK1A-His	pTK1A-His	Full length His-PBP1A, Kan[R]	[71]
pET28a-	pET28a-His-NlpI	Soluble NlpI construct, N-terminal His-tag, KanR	[6]
pBAD18	pBAD18-His-PBP4ΔD3	for purification of His- PBP4 lacking domain 3, Amp[R]	This work
pBAD18	Tre1	Arabinose inducible Tre1, Amp[R]	[75]
pBad18	Tre1 E415Q	TreE415Q inactive Tre1 variant, Amp[R]	[75]
pXL86	pTHV mCh-FtsE wt	p*Trc* 99A down expressing mCh-FtsE(wt) colE1 ori, Amp[R]	This work
pXL89	pTHV mCh-FtsE ts	p*Trc* 99A down expressing mCh-FtsE(ts) (P135S) colE1 ori, Amp[R]	This work
pXL110	pSAV mNG-NNN-FtsE	p*Trc* 99A down expressing mNG-FtsE, p15A origin, Cam[R]	This work
pXL133	pSAV-PBP4	p*Trc* 99A down expressing PBP4, p15A origin, Cam[R]	This work
pXL134	pSAV-PBP4 S62G	p*Trc* 99A down expressing PBP4 S62G, p15A origin, Cam[R]	This work
pXL135	pSAV-FtsE	p*Trc* 99A down expressing FtsE, p15A origin, Cam[R]	This work
pXL137	pSAV-FtsE K41Q	p*Trc* 99A down expressing FtsE K41Q, p15A origin, Cam[R]	This work
pXL138	pSAV-FtsE D162A	p*Trc* 99A down expressing FtsE D162A, p15A origin, Cam[R]	This work
pXL139	pSAV-FtsE E163A	p*Trc* 99A down expressing FtsE E163A, p15A origin, Cam[R]	This work
pXL140	pSAV-PBP4-ΔD3	p*Trc* 99A down expressing PBP4Δ173-247 (domain 3), p15A origin, Cam[R]	This work
pXL150	pSAV-PBP4-D155A	p*Trc* 99A down expressing PBP4 D155A, p15A origin, Cam[R]	This work
pXL151	pSAV-PBP4-R361A	p*Trc* 99A down expressing PBP4 R361A, p15A origin, Cam[R]	This work

963 Amp[R], Kan[R] and Cam[R] are ampicillin, kananamycin and chloramphenicol resistance,

964 respectively

967 **TABLE 2 Primers used in this study.**

Primer	Sequence 5'-3'	Purpose
priXL51	ACCATGGCTAATTCCCATGTCAG	GA_pkd3_F
priXL54	GTGTAGGCTGGAGCTGCTTC	PCR_pkd3_R
priXL187	CCGGCCGAATTCAACAACAACTCGATGATTCGCTTTGAACATGTCAGC	FtsE-EcoRI fw
priXL188	GGCCAAGCTTTTATTCATGGCCCACGCCTCC	FtsE-EcoRI rv
priXL221	CATGAAGCCGTTGGCTTAACC	ΔFtsE-FA-F
priXL222	TAGGAACTTCGAAGCAGCTCCAGCCTACACTGTTAATCCTCTCGGGCAAAAAG	ΔFtsE -FA-R
priXL223	GGATATTCATATGGACCATGGCTAATTCCCATGTCAGGCGGTGGTGAACAAGCCCGC	ΔFtsE- 1-150
priXL224	GGTGAGCATGCGATAGGAACG	ΔFtsE-1:150
priXL225	TGAGCGGATAACAATTTCACACAGGAAACAGACCATGAATAAGCGCGATGCAATC	ΔFtsE-clean
priXL226	CTTTTCAGATCCTGCAATGCG	ΔFtsE-clean
priXL230	ATTCCGGCGCAGGGCAAAGTACTCTCCTGAAGCTG	FtsE-K41Q-F
priXL231	GATCAGCTTCAGGAGAGTACTTTGCCCTGCGCCGGA	FtsE-K41Q-R
priXL232	CGGTACTGCTAGCGGCTGAACCGACTGGTAACCTG	FtsE-D162A-F
priXL233	AGGTTACCAGTCGGTTCAGCCGCTAGCAGTACCGC	FtsE-D162A-R
priXL234	GTACTGCTAGCGGACGCTCCGACTGGTAACCT	FtsE-E163A-F
priXL235	AGGTTACCAGTCGGAGCGTCCGCTAGCAGTACCGC	FtsE-E163A-R
priXL236	AGATGGCGCTGCCTGCCGGCACCCAGAAAG	PBP4-S62G-F
priXL244	GCGCGAATTCatgCGATTTTCCAGATTTATCA	EcoRI-PBP4-F
priXL245	GCGCAAGCTTctaATTGTTCTGATAAATATC	HindIII-PBP4-R
priXL250	GTTGACCGCAACGGTGGCGGAAGCGGGTCTGTGCAGGATGGAGCCAGCTATG	PBP4ΔD3-F
priXL251	TCCATCCTGCACAGACCCGCTTCCGCCACCGTTGCGGTCAACTATGGCGGC	PBPΔD3-R
priXL266	CCGGCTGGCCATGGAATGCCATGACACAATGCTTTAGCGCTC	PBP4-D155A-F
priXL267	GGAGCGCTAAAGCATTGTGTCATGGCATTCCATGGCCAGCCGG	PBP4-D155A-R
priXL268	GCCGATGGTTCAGGGCTTTCGGCGCATAACCTGATTGCCCCCGCC	PBP4-R361A-F
priXL269	GGCGGGGGCAATCAGGTTATGCGCCGAAAGCCCTGAACCATCGGC	PBP4-R361A-R
prXW5-F	TAGTATGACGGCTCGATTCCAGGTTGTTAGCGCGAGATTgtgtaggctggagctgcttc	LMC500ΔdacB
priXW6-R	AATCTGAAGCCCCGGCCATGTGCCGGGGTTTCTTTTTGAatgggaattagccatggtcc	LMC500ΔdacB

968
969

Midcell localization of PBP4 of *Escherichia coli* modulates the timing divisome assembly

Jolanda Verheul[1#], Adam Lodge[2,7#], Hamish C.L. Yau[2,8#], Xiaolong Liu[1,9#], Xinwei Liu[1], Alexandra S. Solovyova[3], Athanasios Typas[4,5] Manuel Banzhaf[6*], Waldemar Vollmer[2**], and Tanneke den Blaauwen[1***]

[1]Bacterial Cell Biology, Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam; Science Park 904, 1098 XH Amsterdam, The Netherlands.
[2]Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle upon Tyne, NE2 4AX, United Kingdom.
[3]NUPPA, Leech Building, Framlington Place, Newcastle upon Tyne NE2 4HH, UK
[4]European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, Germany
[5]European Molecular Biology Laboratory, Structural & Computational Unit, Heidelberg, Germany
[6]Institute of Microbiology & Infection and School of Biosciences, University of Birmingham, Edgbaston, Birmingham, UK. B15 2TT
[7]Present address: Iksuda Therapeutics, The Biosphere, Draymans Way, Newcastle Helix, Newcastle upon Tyne, NE4 5BX, UK
[8]Present address: Procter & Gamble, Newcastle Innovation Centre, Whitley Road, Newcastle-Upon Tyne, NE12 9TS, UK.
[9]Department of Biochemistry, University of Oxford, OX1 3QU, Oxford, United Kingdom.

[#]contributed equally.
*Corresponding author. Tel: +44 121 4145586; Email: m.banzhaf@bham.ac.uk
**Corresponding author. Tel: +44 191 208 3216; Email: w.vollmer@ncl.ac.uk
***Corresponding author. Tel: +31 20 525 3852; Email: t.denblaauwen@uva.nl

FIGURES AND LEGENDS

Fig 1. Structure of PBP4 and its localization at midcell and in the lateral wall during fast

growth. A. Crystal structure of PBP4 in which the three domains (numbers 1-3) and some of

the residues that are essential for its endopeptidase activity are indicated. B. MC4100 cells

were grown exponentially in TY medium at 37°C to an $OD600$ of 0.3 and fixed and

immunolabeled with BW25113Δ*dacB* pre-adsorpted anti-PBP4. Upper panel phase contrast

image corresponding to the SIM fluorescence image in the lower panel. Scale bar equals 2 μm

C. BW25113 wild-type cells were grown in TY at 37°C and harvested in the exponential phase

at an OD600 of 0.3 and fixed. Samples were divided into three aliquots and cells were

immunolabeled with antibodies against PBP3, FtsN and PBP4. The first aliquot was

immunolabeled without any treatment, the second after permeabilizing the cells with

TritonX-100 (right: phase contrast and fluorescence images) and the third after

permeabilizing cells with Triton-X100 and lysozyme (left: phase contrast and fluorescence

images). The scale bar equals 5 μm. PBP4, but not PBP3 or FtsN, is accessible without

degrading the peptidoglycan layer.

Fig 2. Activity of PBP4 is not required for localization. Cells were grown in minimal glucose

medium to steady state at 28°C, fixed and immunolabeled with antibodies against PBP4. (A)

The extra fluorescence at midcell (FCPlus) in the Δ*dacB* strain transformed with the empty

plasmid (EP, yellow), or plasmids expressing wild-type PBP4 (black), PBP4S63G (blue),

PBP4D155A, (purple) PBP4R361A (red), PBP4ΔD3 (green) was determined and plotted as

function of the cell division cycle age as in bins of 5% age classes with the error bar indicating

the 95% confidence interval. (B) Demographs of the localization fluorescence pattern of the

PBP4 variants shown in (A) with the cells sorted according to their cell length. The white line

indicates the length of the cells. Intensity scaling is identical for all maps. Number of cells

analyzed for each immunolabeling was at least 2000 cells.

Fig 3. PBP1A and LpoA contribute to PBP4 midcell localization. Cells were grown

exponentially to an OD of 0.3 in TY at 37°C, then fixed and immunolabeled with antibodies

against PBP4. Because fluorescence imaging by microscopes is usually not directly

comparable between different experiments, all results in each experiment were normalized

to the parental strain BW25113. (A) PBP4 fluorescence at midcell per μm circumference of

the cell. (B) Concentration of PBP4 in the cells. (C) length of the cells. (D) Diameter of the cells.

Δ*lpoA* (n=5), Δ*lpoB* (n=5), Δ*mrcA* (PBP1A, n=2), Δ*mrcB* (PBP1B, n=2). Each point is the average

of 1000-2000 cells. Based on the one-way Anova the difference in midcell localization is

69   significant (P=0.009) while the difference in concentration of PBP4 is not significant

70   (P=0.0484).

Fig 4. PBP4 interacts with PBP1A, LpoA and NlpI. (A) Summary of apparent K_D values of the

interactions of PBP4 or a version lacking domain 3 (PBP4ΔD3) with LpoA, the C-terminal

domain of LpoA (LpoA^C), the N-terminal domain of LpoA (LpoA^N), NlpI and PBP1A determined

by microscale thermophoresis (MST). Corresponding binding curves are shown in Suppl.

Figure S7. (B) Pulldown experiment showing that domain 3 of PBP4 is required for interaction

with NlpI. Oligo-histidine tagged NlpI (His-NlpI) pulled down PBP4 to Ni-NTA beads. His-tagged

PBP3ΔD3 did not pull down untagged NlpI. A, applied sample; E, eluted sample. (C) *In vivo*

cross-linking/co-immunoprecipitation showing interaction between PBP1A and PBP4.

Growing cells of BW25113 (wt) or BW25113Δ*dacB* (ΔPBP4) were chemically cross-linked with

DTSSP and cell extract was immunoprecipitated with purified anti-PBP4 antibody. Control

samples did not receive anti-PBP4 antibody. The cross-linker was cleaved by reducing agent

and proteins were separated by SDS-PAGE and transferred to a membrane, followed by

detection of PBP1A with specific antibodies. PBP1A was detected in the elution of the sample

from wild-type and not from ΔPBP4. (D) *In vivo* cross-linking/co-immunoprecipitation (as in

panel C) showing that PBP4 and be cross-linked with LpoA in cells.

Fig 5 PBP4 or NlpI do not affect the activity of PBP1A whereas LpoA reduces the activity of

PBP4. (A) Summary of the results from an *in vitro* PG synthesis assay with PBP1A (in the

presence or absence of LpoA) and radio-labelled lipid II substrate. The presence of NlpI,

catalytically inactive PBP4(S62A) or both together, does not affect the cross-linkage of the PG

product of a PBP1A or PBP1A/LpoA reaction (left side), and NlpI, PBP4(S62A) or both together,

don't induce enhanced carboxypeptidase activity of PBP1A, which would be seen as a higher

content of tetrapeptides (right side). Values are mean ± variation of two independent repeats.

Example chromatograms are shown in Figure S8. (B) The presence of LpoA modestly reduces

the activity of PBP4 in a DD-carboxypeptidase assay using the substrate *N*-acetyl-L-Lys-D-Ala-

D-Ala. (C) LpoA also reduces slightly the activity of PBP4 against PG sacculi from BW25113, as

seen by the reduced digestion of TetraTetra upon a 30 min incubation period.

(D The Δ*dacB* and Δ*dacB*Δ*lpoA* mutant strains are equally sensitive to overexpression of PBP4.

The average growth over time as measured by OD650 nm is plotted against the added IPTG

concentration. The culture grown in TY with glucose to suppress the expression is represented

by -5 on the X-axe. All other data points are from cell grown in TY without glucose (n=4).

Fig 6 Cell division cycle timing of the localization of PBP4. (A) MC4100 cells were grown to

steady state in minimal glucose medium at 28°C and immunolabeled with antibodies specific

for PBP4. Phase contrast (top) and fluorescence is shown (bottom). Scale bar equals 2 µm. (B)

The extra fluorescence at midcell compared to the rest of the cell (FCPlus) is plotted as

function of the normalized cell division cycle age. The dots (grey for MC4100 and light pink

for the BW25113 Δ*dacB strain*) are the values measured for the individual cells and the

markers with bars (black for the wild-type MC4100, red for the BW25113 Δ*dacB*) are 5% age

bins with 95% confidence. (C) Demographs of the localization of FtsZ, PBP4 and FtsN in the

cells sorted according to cell length. The white line indicates the length of the cells. (D)

Comparison of the timing of the localization at midcell of FtsZ (black), PBP4 (green) and FtsN

(blue) during the cell division cycle age. The FCPlus values are min-max normalized to enable

timescale comparison, despite differences in molecule number and antibody affinities. (E)

BW25113 wild type cells were grown to steady state in minimal glucose medium at 28°C and

immunolabeled with antibodies specific for FtsZ, PBP4 and FtsN. The FCPlus of three

independent biological experiments in 5% age bins was determined. The binned FCPlus values

were min-max normalized. The average min-max normalized FCPlus values of the three

experiments were subsequently plotted as function of the cell division cycle.

Fig 7 PBP4 localization is dependent on the presence of the proto-ring. A. MC4100 was

transform with a plasmid that expressed either Tre1 that abolished the ability of FtsZ to

polymerize by ADP-ribosylation of residue R174 or with the inactive variant Tre1(E415Q). Cells were grown in Gb1 medium at 28°C and expression of the inhibitor was induced for 2 mass doublings (MD) with 0.15% arabinose. Subsequently, the cells were fixed and immunolabeled with antibodies against FtsZ or PBP4. From left to right: the phase contrast (PC), corresponding fluorescence image of the PBP4 labeling and the merged images ate shown. In the demographs the cells are sorted according to cell length (contours in white). The number of cells analyzed were 3694 and 3756 for Tre1E415Q and 1275 and1621 for Tre1 for anti FtsZ and anti PBP4, respectively. **B.** Isogenic strains of MC4100 producing different temperature sensitive (ts) versions of cell division proteins were grown to steady state in Gb1 medium at 28°C and split in two parts. These were 1:4 diluted in prewarned medium and grown for two MD at either 28°C or 42°C. The cells were fixed, labeled with antibodies specific for PBP4 (a4). From left to right: the phase contrast (PC), corresponding fluorescence image of the PBP4 labeling and the merged images at the permissive temperature, demograph of PBP4 fluorescence localization where cells are sorted according to their cell length (contours in white). This is followed by the same series from the non-permissive temperature samples. The number of cells analyzed were 2809 and 1576 for LMC515 FtsE(ts), 3833 and 1203 for LMC531 FtsQ (ts) and 3876 and 926 for LMC510 PBP3(ts), for the cells grown at 28°C and 42°C, respectively. The scale bar equals 5 μm. Within one antibody staining the brightness and contrast of the samples is identical and therefore directly comparable.

Fig 8 PBP4 is recruited by the septal cleavage complex FtsEX. (A) Cells of BW25113 expressing

the proteins indicated were grown in LB at 37°C to OD_{600} = 0.3, fixed and labeled with specific

antibodies against PBP4. From left to right, the phase contrast, corresponding fluorescence

image of the PBP4 labeling, and the merged former two images, map of fluorescence with the

cell lengths outlined (with the same contrast and brightness for all maps) of PBP4 localization

where cells are sorted according to their cell length are shown. The number of cells analyzed

were 687 for Δ*envC*, 434 for Δ*ftsE*, and 346 for Δ*ftsEX*. The scale bar equals 5 μm. (B) Cells of

LMC500 or XL36 (LMC500Δ*ftsE*::p*Trc99A*down *ftsX*) expressing wild-type or active site

versions of FtsE were immunolabeled with anti-PBP4 antibodies. Cells were grown in TY at

30°C and FtsE expression from plasmid was induced for 2 mass doubling with 30 μM IPTG,

and cell were fixed and harvested at an OD$_{600}$ of 0.3. The maps of PBP4 fluorescence with the

cell lengths outlined is shown for the indicated strains. Number of cells analyzed: LMC500

(2462), (wt) (2588), K41Q (728), D162A (554), and E163E (1052).

Fig 9 Localization of PBP4 depends on the presence of domain 3 but not on activity. Cells

were grown in minimal glucose medium to steady state at 28°C, fixed and immunolabeled

with the indicated antibodies. (A and B) Graphs with the percentage of constricting cells and

the average cell length in μm for the various mutants expressed from plasmid in the Δ*dacB*

strain and also of the parental strain BW24113 (n = 4). (C) The concentration of PBP4 in

fluorescent units per μm^3 in these cells for a representative experiment (out of the four

repeats).

(D) Graphical illustration of the meaning of t_0 and $t_{1/2}$. (E) Cell division cycle age timing of FtsZ,

PBP4, and FtsN for the parental strain BW25113, the Δ*dacB* strain carrying plasmids for the

expression of PBP4WT, PBP4S63G, PBP4D155A, or PBP4R361A. Plasmids were not induced.

(F). Cell division cycle age timing of FtsZ and FtsN for the Δ*dacB* strain and its parental

BW25113 (left) and the Δ*dacB* strain and its parental LMC500. P 0.0001 = ***, P 0.001 = **.

Number of points correspond to the number of independent biological replicates.

Fig 10 Model of the organization of PBP4 in the periplasm of *E. coli*. In a minimal medium

grown cell of average size of 3 by 0.86 μm 230000 Lpp trimeric proteins are present of which

1/3 is bound to PG. (A) During elongation PBP4 (or a fraction of the PBP4 molecules) is kept

away from its substrate through its interaction with NlpI and possibly LpoA. (B) PBP1A and

LpoA associate with ZipA and the Z-ring to assist in preseptal PG synthesis. Because of their

presence, the absence of NlpI and the newly synthesized PG, PBP4 is attracted by PBP1A and

LpoA. More PBP4 is accumulating at midcell and becoming clearly visible by immunolabeling

upon maturation of the divisome and synthesis of septal PG. All proteins apart from the FtsZ

polymer have been drawn according to their crystal structure shape and hydrated crystal

structure sizes (scale bar equals 4 nm). The membranes are assumed to be 2 nm and the

distance between the outer and the inner membrane 21 nm.

Midcell localization of PBP4 of *Escherichia coli* modulates the timing of
divisome assembly
3

4 Jolanda Verheul[1#], Adam Lodge[2,7#], Hamish C.L. Yau[2,8#], Xiaolong Liu[1,9#], Xinwei Liu[1],
5 Alexandra S. Solovyova[3], Athanasios Typas[4,5] Manuel Banzhaf[6*], Waldemar Vollmer[2**], and
6 Tanneke den Blaauwen[1***]
7
[1]Bacterial Cell Biology, Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam;
Science Park 904, 1098 XH Amsterdam, The Netherlands.
[2]Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle upon Tyne, NE2 4AX,
United Kingdom.
[3]NUPPA, Leech Building, Framlington Place, Newcastle upon Tyne NE2 4HH, UK
[4]European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, Germany
[5]European Molecular Biology Laboratory, Structural & Computational Unit, Heidelberg, Germany
[6]Institute of Microbiology & Infection and School of Biosciences, University of Birmingham, Edgbaston,
Birmingham, UK. B15 2TT
[7]Present address: Iksuda Therapeutics, The Biosphere, Draymans Way, Newcastle Helix, Newcastle upon Tyne,
NE4 5BX, UK
[8]Present address: Procter & Gamble, Newcastle Innovation Centre, Whitley Road, Newcastle-Upon Tyne,
NE12 9TS, UK.
[9]Department of Biochemistry, University of Oxford, OX1 3QU, Oxford, United Kingdom.
22
23 [#]contributed equally.
24 *Corresponding author. Tel: +44 121 4145586; Email: m.banzhaf@bham.ac.uk
25 **Corresponding author. Tel: +44 191 208 3216; Email: w.vollmer@ncl.ac.uk
26 ***Corresponding author. Tel: +31 20 525 3852; Email: t.denblaauwen@uva.nl
27

28 **Supplementary figures:**

FIG S1 Pre-adsorption of antibodies against Δ*dacB* BW25113 strain results in PBP4 specific

antibodies. (A) Demograph with cells sorted according to length of PBP4 fluorescence of the

BW25113Δ*dacB* (PBP4) strain, BW245113 (wt) strain before purification of the antibody and

the BW25113Δ*dacB* strain and BW245113 strain after purification of the antibody. (B)

Absolute fluorescence average profiles of cells from the BW25113 Δ*dacB* strain and the

BW245113 strain before purification of the antibody and from the BW245113 strain after

purification of the antibody (C) Phase contrast and corresponding fluorescence image of anti

PBP4 immunolabeled BW25113Δ*dacB* (left) and wild-type cells using the supernatant of

BW25113Δ*dacB* pre-adsorbed antibodies. The scale bar equals 5 μm.

FIG S2 All PBP4 mutants are equally well expressed and only S62G is not able to bind

Bocillin-FL. (A) An immunoblot of membranes of PBP4 wild-type and variants. *This extra

band is due to non-specificity of the primary antibody, which was for the purpose of

immunoblotting not affinity purified. (B) Immunoblot of supernatant after pelleting the

membrane of PBP4 wild-type and variants. (C) The corresponding gel where binding of the

fluorescent β-lactam Bocillin-FL is visible for all mutants apart from S62G. EP is empty plasmid,

WT is wild-type. The other samples are PBP4 variants.

FIG S3 PBP4ΔD3 binds beta-lactam but is inactive against PG substrates. (A) PBP4 versions

were incubated with the fluorescent β-lactam Bocillin FL with or without pre-incubation with

Penicillin G (PenG), followed by SDS-PAGE analysis and detection of covalent Bocillin-PBP4

adducts by fluorescence scanner. PBP4 and PBPP4ΔD3, but not catalytically inactive

PBP4(S62A), bound Bocillin FL. (B) Only wild-type PBP4, but not PBPP4ΔD3 or PBP4(S62A),

was active in a DD-carboxypeptidase assay. (C) PBP4 versions were incubated with

muropeptides from BW25113 prior to their separation by high-performance liquid

chromatography (HPLC). A control sample contained no enzyme. PBP4 DD-endopeptidase

activity is demonstrated by the reduction in the dimer (TetraTetra) substrate peak and

increase in the monomer (Tetra) product peak. PBPP4ΔD3 and PBP4(S62G) were inactive. (D)

Quantification of the Tetra and TetraTetra peaks shown in panel C (top) and quantification of

a similar analysis with PG sacculi (incubated with PBP4 versions or no enzyme), followed by

generation of muropeptides and HPLC analysis (bottom).

FIG S4 The absence of domain 3 does not affect the dimerization, secondary structure or PG

binding of PBP4. (A) Analytical ultracentrifugation sedimentation velocity experiment of PBP4

and PBP4ΔD3 shows that both proteins are mainly dimers. The determined sedimentation

coefficients (s_{exp}) for the dimers fit well with the theoretical values (s_{calc}) that were calculated

from atomic coordinates of both protein dimers (pdb accession code for monomer is 2EX2).

(B) PBP4 and PBP4ΔD3 show similar far UV circular dichroism spectra. (C) PBP4 and PBP4ΔD3

co-sediment with PG sacculi from BW25113 cells, but did not sediment in control samples

without sacculi, demonstrating binding to PG. S, supernatant; W, wash fraction; P, pellet

fraction.

FIG S5 PBP4 localizes at site of inactive divisomes. MC4100 cells were grown to steady state

in minimal glucose medium at 28°C and split in two parts. One part was 1:4 diluted in

prewarmed medium without aztreonam and the other part was 1:4 diluted in medium with

10 μg/ml aztreonam. The cells continued to grow for 0, 1, 2 or 3 mass doublings (MD) and

were fixed and immunolabeled with antibodies specific for FtsZ or PBP4. Demographs with

identical brightness and contrast of the FtsZ or the PBP4 fluorescence of cells grown in the

absence or presence of aztreonam sorted according to length. The white line shows the

borders of the cells. Number of analyzed cells per demograph for FtsZ and PBP4 were, 5084

and 3448 (0), 1430 and 1858 (1), 857 and1630 (2) and 659 and 595 (3), respectively.

FIG S6 PBP4 localizes between the outer membrane and the peptidoglycan layer in a Δ*nlpI*

strain. (A) BW 25113 Δ*nlpI* cells were grown exponentially in TY medium at 37°C and fixed

while shaking when at an OD600 of 0.3. The cells were harvested and split in three portions

of which one was directly immunolabeled with to Δ*dacB* cells pre-adsorbed anti-PBP4, the

second was first treated with Triton X-100 and the last was treated with Triton X-100 and

lysozyme and then immunolabeled. The scale bar equals 2 μm. (B) Map of PBP4 fluorescence

sorted according to cell length of wild-type cells and Δ*nlpI* cells displayed at the same

brightness and contrast values.

FIG S7 Interactions of PBP4 and PBPP4ΔD3 measured by microscale thermophoresis (MST).

MST binding curves for interactions using different fluorescently-labelled proteins (LpoA,

LpoA[C], Lpo[N], NlpI or PBP1A) titrated against fixed concentrations of PBP4 or PBPP4ΔD3.

Apparent K_D values are mean ± SD of three independent experiments and summarized in Fig.

4A.

FiG S8 PBP4 or NlpI do not affect the activity of PBP1A or PBP1A/LpoA. Representative HPLC

chromatograms of samples from an *in vitro* PG synthesis assay of PBP1A and radiolabelled

lipid II, in the presence of proteins indicated. The main PG products upon digestion with the

muramidase cellosyl are the disaccharide pentapeptide (Penta) and the bis-disaccharide

tetrapentapeptide (TetraPenta). The % peptides present in cross-links is quantified on the

right side and presented as mean ± variation of two independent experiments.

FIG S9 NG-FtsE effectively localizes and recruits PBP4. Strains were grown in LB at 37°C to an

OD of 0.3 in the presence of 30 µM IPTG to induce expression of mNG-FtsE then fixed and

immunolabeled with antibodies specific for PBP4. (A) LMC500 (1108 cells analyzed). (B) XL36

(LMC500::p*Trc*99down*ftsX*, Δ*ftsE*) with pXL110 mNG-FtsE(wt) (1189 cells analyzed). From left

to right, phase contrast image, anti-PBP4 immunolabeling, mNG-FtsE, and the corresponding

map of diameter with the fluorescence maps of cells sorted according to length. Scale bar

equals 5 µm.

	Phase contrast	Anti PBP4	Merged	Dia	Fluor
wt					
Δ*tolA*					
Δ*pal*					
Δ*nplD*					
Δ*amiABC*					

FIG S10 PBP4 localization is not dependent on the presence of TolA, Pal, NlpD or AmiABC.

Isogenic strains of the wild-type strain BW25113 were grown in LB at 37°C to a OD600 of 0.3,

fixed and labeled with specific antibodies against PBP4. From left to right, the phase contrast,

corresponding fluorescence image of the PBP4 labeling, and the merged former two images,

map of diameters (Dia) and map of fluorescence (Fluor) PBP4 localization where cells are

sorted according to their cell length are shown. The number of cells analyzed were 1053 for

BW25113 (*wt*), 918 for Δ*tolA*, 1314 for Δ*pal*, 1600 for Δ*nlpD*, and 1410 for Δ*amiABC*. The scale

bar equals 5 μm.

Table S1. Morphological parameters LMC500 and LMC500Δ*dacB*

Parameter	LMC500	n	LMC500Δ*dacB*	n	p
Length (μm)	2.57 ± 0.07	8	2.58 ± 0.09	12	0.79
Diameter (μm)	0.89 ± 0.09	8	0.91 ± 0.19	12	0.73
Constrictions (%)	16.9 ± 1.8	8	17.4 ± 1.3	12	0.53

148   n = number of biological replicates, P from a two tailed T-test to verify whether that the means

149   are significant different.