Title: The critical ecological role of an endemic, large-bodied frugivore on a small tropical

island

Authors: Rohit Naniwadekar[1,*], Abhishek Gopal[1], Navendu Page[2], Sartaj Ghuman[1], Vivek

Ramachandran[3] and Jahnavi Joshi[4]

Present Address of Authors:

[1]Nature Conservation Foundation, 1311, "Amritha", 12[th] Main, Vijayanagar 1[st] Stage, Mysuru,

Karnataka, India 570017.

[2]Wildlife Institute of India, Post Box no. 18, Chandrabani, Dehradun, Uttarakhand, India

248001.

[3]National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary Road,

Bengaluru, Karnataka, India 560065.

[4]Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad, Telangana, India 500007.

* **Corresponding Author** – Rohit Naniwadekar. Email: rohit@ncf-india.org. ORCiD -

https://orcid.org/0000-0002-9188-6083.

ABSTRACT

Mutualistic interactions, like seed dispersal, are important for maintaining plant diversity in

tropics. Islands, which have a smaller subset of plants and frugivores, when compared with

mainland communities offer an interesting setting to understand the relative role of ecological

and evolutionary processes in influencing the interactions between plants and frugivores. In this

study, we examined the relative influence of functional traits and phylogenetic relationships on

the plant-seed disperser interactions on an island and a mainland site. The island site allowed us

to investigate the organization of the plant-seed disperser community in the natural absence of

key frugivore groups (bulbuls and barbets) of Asian tropics. The largest frugivore on the island,

the point endemic Narcondam Hornbill, was also the most abundant frugivore on the island and

played a significant quantitative and central role in the plant-seed disperser community. Species

strength of frugivores was positively associated with frugivore abundance. Among plants, figs

had the highest species strength and played a central role in the community during the study

period. We found that the island had among the highest reported densities of hornbills and figs in

the world. Island-mainland comparison revealed that the island plant-seed disperser community

was more asymmetric, connected and nested as compared to the mainland community with

which it shared certain plant and frugivore species. Neither functional traits nor phylogenetic

relationships were able to explain the patterns of interactions between plants and frugivores on

the island or the mainland. Diffused nature of interactions between plants and frugivores, trait

convergence and plasticity in foraging behavior likely contribute to shaping the interactions

between plants and frugivores. This study underscores the need to study plant-seed disperser

communities on tropical islands, particularly in the Indian Ocean, which have inadequate

representation in the literature.

INTRODUCTION

Endozoochory, which is a mutualistic interaction between plants and frugivorous animals, is a critical stage in plant regeneration and contributes to the maintenance of tree diversity in tropics (Terborgh et al. 2002). The community of food plants and seed dispersers form a network of bipartite interactions. These networks could be influenced by different ecological (e.g. functional traits, phenological overlap) and evolutionary (e.g. co-evolution) processes (Vázquez et al. 2009; Guimarães Jr et al. 2011). There are limited studies that have investigated the relative role of eco-evolutionary processes on plant-seed disperser interactions.

Interactions could be influenced by trait complementarity which could be modulated by trait conservatism or trait convergence (Rezende et al. 2007; Bastazini et al. 2017). Trait conservatism will result in closely related species interacting with similar partners. On the other hand, interactions could be influenced by trait convergence wherein a similar trait is exhibited by unrelated species. Interacting communities often exhibit organize themselves into modules. These modules could be organized based on species' shared evolutionary history coupled with similar traits (Donatti et al. 2011) or through trait convergence (Peralta 2016) or abundance (Schleuning et al. 2014b). Traits relevant to the seed dispersal services, and the number of partners that a frugivore has, have been demonstrated to exhibit phylogenetic signal (Rezende et al. 2007). Some studies have also detected phylogenetic signal in species that act as connectors between modules that enable network integration (Schleuning et al. 2014b). The ecophylogenetic approach outlines four scenarios based on the interactions that are mediated by phylogeny or functional traits, which allows the assessment of relative contribution of traits and phylogenetic relationships on plant-animal interactions (Bastazini et al. 2017). The influence of phylogenetic component would be indicative of the role of evolutionary processes like co-evolution in the

organization of the community, while the influence of functional component controlling for

phylogenetic history would imply trait complementarity. This enables an understanding of the

relative roles of these factors in community assembly. Despite substantial availability and

progress in the use of phylogenetic data in understanding community structure, systematic

examination of the role of evolutionary processes in tandem with ecological processes in

assembly of plant-frugivore communities is limited.

Bulk of the information on plant-disperser communities are from mainland sites with few studies

from oceanic islands and even fewer in an island-mainland comparative framework (Schleuning

et al. 2014*a*). Despite zoochory being a pervasive interaction in Asian wet tropics, there is

limited information on plant-disperser communities from the mainland Asian wet tropics (Corlett

2017) and very poor information from oceanic islands in the Indian Ocean (Escribano-Avila et

al. 2018). Island systems have experienced a disproportionately higher percentage of species

extinctions as compared to mainland systems (Heinen et al. 2018). Given the naturally

asymmetric nature of plant-frugivore communities on islands (Schleuning et al. 2014*a*),

frugivore loss on islands can be expected to have significant cascading impacts on plant

communities. Oceanic barrier restricts the species that can colonize and persist on the island.

Therefore, they offer a natural laboratory to evaluate the influence of the absence of key

mainland frugivore groups on plant-seed disperser interactions. Comparing distribution of

interactions on island and mainland can provide insights into the organization of plant-seed

disperser communities.

Given this background, we examined the organization of the plant-disperser community on a

remote and small tropical island (Narcondam Island) in South Asia using the network and

ecophylogenetic approaches (Fig. 1A). The Narcondam island, which is the only place where the

Narcondam Hornbill *Rhyticeros undulatus* is found (Fig. 1B), is characterized by a species-poor

frugivore assemblage, enabled us to examine the relative quantitative role of different species in

seed dispersal. In the Asian tropics, small-bodied frugivores like bulbuls and barbets are known

to play key roles in seed dispersal (Naniwadekar et al. 2019). However, the Narcondam Island

does not have any bulbuls and barbets and most of the dominant frugivores on the island are

medium- and large-bodied avian frugivores (> 100 g) (Fig. 1C). This offers an interesting system

to understand the distribution of interactions in absence of key frugivore group. We compared

this island community with a mainland community from north-east India with which it shares

several frugivore and plant species. The specific aims of the study were to 1) identify key seed

dispersers and plant groups on an oceanic island community by comparing species-level

properties (degree, species strength and weighted betweenness) across different species, 2)

examine the influence of abundance and morphological traits on species strength of plants and

frugivore species, 3) compare the network-level properties (web asymmetry, connectance,

specialization and weighted nestedness) of the island community with a mainland community,

and 4) examine the influence of phylogenetic and functional component on the distribution of

interactions between plants and frugivores on the island and mainland communities. We

expected that large-bodied frugivores will not be restricted by morphological constraints and will

demonstrate dietary shifts on islands in the absence of small-bodied frugivore. This will alter the

organization of the plant-seed disperser community on the island as compared to mainland.

These questions scale up from species-level to community-level and across spatial scales from

island to mainland. On the temporal axis, we examine the influence of ecological processes on

species roles in community and ecological and evolutionary processes on distribution of

interactions in the community. This system allowed us to explicitly examine the change in the

role of large-bodied frugivores in plant-seed disperser communities in the absence of small

frugivores (Fig. 1C).

MATERIALS AND METHODS

Study area: The study was carried out on a small (area: 6.8 km^2), volcanic island, the

Narcondam Island (13º30'N, 94º38'E), in the Andaman Sea in India. The island is part of the

Indo-Myanmar Biodiversity Hotspot (Myers et al. 2000). Three distinct vegetation types on the

island include the narrow strip of littoral forests, forests dominated by deciduous species in the

north-eastern part of the island and evergreen forests in most of the island (Page et al. in prep.).

The Endangered Narcondam Hornbill *Rhyticeros narcondami* is found only on the Narcondam

Island (BirdLife International 2017). Around 50 species of birds have been reported from the

island (Raman et al. 2013). There are two terrestrial mammals on the island, including a shrew

(*Corcidura* sp.) and an invasive rat (*Rattus* sp.). There are at least four species of bats on the

island including a fruit bat species (*Pteropus hypomelanus*). Narcondam Hornbill (body mass:

600-750 g) is the largest frugivore on the island. Additional details of study area are in Appendix

S1.

Frugivore and plant abundance: We conducted the study between December 2019 – February

2020. We used variable-width line-transect surveys to estimate the density and abundance of

birds on Narcondam Island (Thomas et al. 2010). We walked 20 unique trails on the island

(Appendix S2: Fig. S1). We divided the entire elevation gradient into three elevation zones: low

(0-200 m), mid (200-400 m) and high (400-700 m) based on topography, vegetation structure

and composition. One to three observers walked trails in the mornings and afternoons. During

the trail walks, we recorded species identity, the number of individuals and the perpendicular

distance of the sighting to the trail. The trail lengths varied between 0.22–1.26 km and the total

sampling effort was 51.4 km (Appendix S2: Table S1).

We laid 49 belt transects (50 m × 10 m) across the three different elevation zones (low: 18, mid:

14, high: 17) to estimate the richness and abundance of woody plants on the Narcondam Island.

Within the belt, we enumerated all wood plants that were ≥ 10 cm GBH (girth at breast height)

along with their girth and taxonomic identity.

Plant-frugivore interactions: To systematically document plant-frugivore interactions, we

performed spot-sampling during our trail walks, and opportunistically outside our trail walks

following Palacio *et al*. (2016). Whenever we encountered a frugivore on a fruiting tree, we

recorded the species identity, number of frugivores, and frugivore foraging behaviour (fruits

swallowed, pecked or dropped) to ascertain whether the frugivore was a seed disperser or not.

Plant-frugivore interaction information was systematically collected outside trail sampling

whenever frugivore was detected on fruiting trees. To determine if we were missing

documenting interactions of frugivore species with low detection probability through spot

sampling, we systematically observed fruiting trees. We observed 40 fruiting individuals of 12

species following Naniwadekar et al. (2019a) (Appendix 2: Table S2). Additional details of the

tree watches are in Appendix S1. We measured widths of at least five fruits each of 23 species of

fleshy-fruited plants following Naniwadekar et al. (2019*a*) (Appendix S2: Table S3). Plants were

classified as small-seeded (seed width: < 5 mm), medium-seeded (seed width: 5-15 mm) and

large-seeded (> 15 mm) following Naniwadekar et al. (2019a) (Appendix S2: Table S3).

Analysis

Frugivore and plant abundance: We estimated densities of perched birds with more than 40

visual detections using the R package 'Distance' (Thomas et al. 2010; Miller et al. 2019). We

estimated combined density of two Imperial-pigeon species (Pied Imperial-Pigeon *Ducula*

bicolor and Green Imperial-Pigeon *Ducula aenea*), since we had 45 detections overall.

Additional details of the analysis are provided in the Appendix S1. We estimated the overall tree

density and basal area ($m^2 ha^{-1}$), and density of 27 species of fleshy-fruited plants that were

fruiting during the study period.

Island plant-seed disperser community: We created a matrix of plants and frugivores with the

frequency of interactions documented for each plant-frugivore combination through spot

sampling. We only considered those animals as seed dispersers which we had seen swallowing

the seeds. We didn't consider Alexandrine Parakeet *Psittacula eupatria* in the analysis as we

documented the parakeets predating on seeds or feeding on unripe fruits as has also been

documented elsewhere (Shanahan et al. 2001). We plotted the interaction accumulation curve

using the 'random' method and 1000 permutations as implemented in the R package 'vegan' to

determine if we had adequately documented the diversity of unique plant-seed disperser

interactions (Oksanen et al. 2019).

To determine the relative roles of species in the organization of plant-seed disperser

communities, we examined species-level properties like the degree (number of mutualistic

partners), species strength (importance of species from partner's perspective) and weighted

betweenness centrality (connector species connecting multiple guides in community)

(Bascompte et al. 2006; Mello et al. 2015). We used the R package 'bipartite' for this analysis

(Dormann et al. 2008).

We used the general linear model to examine the influence of abundance (estimated using the

vegetation plots) and fruit type (small-seeded, figs, medium-seeded and large-seeded) on species

strength of the 27 plant species that were fruiting during the study period. During exploratory

analysis, we noticed that figs consistently had higher species strength values. Since certain fig

groups are known to be keystone species in tropical forests and provide key nutrients (Shanahan

et al. 2001), we performed the analysis with figs as a separate category. Species strength was

$\log_{10}$ transformed to approximate normality. In the case of frugivores, since there were only

seven species and species strength values were not normally distributed, we used Spearman's ρ

to examine the correlation between abundance (encounter rate km^{-1}) of frugivores and body mass

with species strength. We obtained body mass information from Wilman et al. (2014). To

determine if the island plant-seed disperser community was organized into distinct sub-

communities, we used the program MODULAR (Marquitti et al. 2014) following Naniwadekar

et al. (2019). Additional details in Appendix S1.

Comparing the island and the mainland plant-seed disperser communities: To determine the

differences in the organization of plant-seed disperser communities between island and

mainland, we compared plant-disperser communities between the Narcondam Island with that of

Pakke Tiger Reserve in north-east India. Information on plant-seed disperser communities is

limited from the Asian tropics. However, there is detailed information on interactions between

48 avian frugivores and 43 plant species is available from Pakke Tiger Reserve in north-east

India (Naniwadekar et al. 2019). Pakke is part of the Eastern Himalaya Biodiversity Hotspot. It is

about 1500 km straight line distance north from the Narcondam Island. Representative species of

all the frugivore genera found on the Narcondam Island are also found in Pakke with species like

the Green Imperial-Pigeon, Asian Koel *Eudynamys scolopaceus* and Common Hill Myna

Gracula religiosa being reported from both sites. Several genera of fleshy-fruited plants found

on Narcondam Island are also found in Pakke. Unlike the island community which is relatively

young, the mainland site is several million years old. We examined network-level properties like

web asymmetry, connectance (proportion of realized interactions), specialization (H_2') and

nestedness (weighted NODF) and compared it with the mainland site (Pakke). Values of web

asymmetry range between -1 to 1 with negative values indicating higher diversity of plant

species as compared to animal species and vice versa (Blüthgen et al. 2007). Specialization is a

measure of complementary specialization of the entire network with high values indicating that

the species are more selective. Weighted NODF is a quantitative measure of nestedness with

high values indicating higher nestedness.

Community assembly on island and mainland: To determine the influence of phylogeny and

functional traits on plant-frugivore interactions on the mainland and island site, we used the

following approach. The plant phylogeny was pruned from the megaphylogeny (Qian and Jin

2016) for both the island and mainland communities individually. This is a time-calibrated

phylogeny with largest representation from 98.6% of families and 51.6% of all genera of seed

plants. Scientific names of plants were standardized by referring to 'The Plant List'

(www.theplantlist.org). However, some species from Pakke and Narcondam were missing from

the megaphylogeny. In those cases, we chose the closest species from the Oriental region. When

we failed to find a congener, we chose a genus close to the focal genus (Appendix S2: Table S3).

We used fruit width and seed size class information for the analysis (Appendix S2: Table S3). In

the case of the birds, we referred to the comprehensive and time-calibrated bird phylogeny

available at www.birdtree.org (Jetz et al. 2012). We found all the focal species for the two sites

in the database except *Pycnonotus flaviventris*, *Chrysocolaptes guttacristatus* and *Sitta*

cinnamoventris. We used congeners from the Oriental region for these species for the analysis

(Appendix S2: Table S4). For the frugivores, we used body mass and percentage fruits in the diet

of frugivores (degree of frugivory) obtained from Wilman et al. (2014) (Appendix S2: Table S4).

We obtained 100 trees from posterior distribution of 10000 trees based on Ericson Sequenced

Species backbone phylogeny with 6670 OTUs for seven bird species for Narcondam and 48

species for Pakke found in the network (Jetz et al. 2012). The maximum clade credibility tree

was used for further analyses as these trees did not vary substantially in topology and branch

lengths. We calculated the phylogenetic distances (pair-wise distances between species using

their branch lengths) using 'cophenetic.phylo' in the R package 'ape' (Paradis and Schliep

2018). We used Gower method to estimate pair-wise distances among traits of both plants and

frugivores using "vegdist" in the R package 'vegan'. We then followed the novel approach

outlined by Bastazini et al. (2017) to determine whether phylogenetic proximity influenced the

evolution of species trait and their interactions in the network or whether species traits influence

the observed pattern of interactions after removing the potential confounding effect of the

phylogeny. We applied this method for Narcondam and Pakke separately. We used the R

packages 'SYNCSA' for this analysis (Debastiani and Pillar 2012; Oksanen et al. 2019). We

used R (ver. 3.5.3) for all the analysis (R Core Team 2019).

In addition, we assessed the phylogenetic signal separately in trait data for birds and plants in

both sites, Narcondam and Pakke. We calculated the Blomberg's *K* for each trait using the R

package "phytools" (Blomberg et al. 2003; Revell 2012).

RESULTS

Frugivore and plant abundance: We detected six avian frugivores during the trail walks. We

had 207 visual detections of Narcondam Hornbill, 71 detections of Asian Koel, and 45 detections

of the two Imperial-Pigeons. We had only two detections each of the Common Hill Myna and

Eye-browed Thrush *Turdus obscurus*. Information on detection probability and mean flock size

is summarized in Appendix S2: Table S5.

Narcondam Hornbill was the most common frugivore on the island and its abundance (mean

(95% CI): 1026 birds (751-1402)) was almost four times that of the Asian Koel (Fig. 2A;

Appendix S1: Table S5). The estimated overall mean (95% CI) density of the Narcondam

Hornbill was 151 (110-206) birds km^{-2}. The mean densities of the Narcondam Hornbill varied

between 101-168 birds km^{-2} across the three elevation zones with the lowest mean densities in

the highest elevation band (Fig. 2B) (Appendix S2: Table S5).

The mean ($\pm$ SE) density and basal area of all the woody plants ($\geq$ 10 cm GBH) was highest in

the 0-200 m elevation band (density: 2091.1 $\pm$ 124.4 plants ha^{-1}; basal area (m^2 ha^{-1}): 58.3 $\pm$ 5.3),

followed by 200-400 m elevation band (density: 1505.7 $\pm$ 101.1 plants ha^{-1}; basal area (m^2 ha^{-1}):

49.1 $\pm$ 3.0) and least in the 400-700 m elevation band (density: 1311.8 $\pm$ 82.7 plants ha^{-1}; basal

area (m^2 ha^{-1}): 39.2 $\pm$ 4.2). However, the mean abundance of the different categories of the

frugivore food plant species (figs, small-seeded non-fig plants, medium-seeded plants and large-

seeded plants), which were fruiting during the study period, was highest in the high elevation

band (400-700 m), followed by middle elevation band and lowest in the low elevation band (0-

200 m) (Fig. 2C).

Island plant-seed disperser community: We documented 752 interactions between seven

frugivore species and 27 plant species during the spot scans performed on trail walks and

opportunistically. We documented all the frugivore species (except Andaman Green Pigeon

Treron chloropterus) previously reported from the island. We documented the migrant Daurian

Starling *Agropsar sturninus*, for the first time. During spot scans on trail walks, we documented

262 interactions between six frugivore species and 19 plant species. During opportunistic spot

scans, we documented 490 interactions between seven frugivore species and 26 plant species.

Narcondam Hornbill was documented feeding on 22 plant species, followed by Asian Koel,

which was documented feeding on 16 plant species (Appendix S2: Table S6). Of the 752

interactions, 533 were of the Narcondam Hornbill, 111 were of the Asian Koel, 36 of the Green

Imperial-Pigeon, 34 of the Pied Imperial Pigeon, 25 of the Common Hill Myna, 12 of the Eye-

browed Thrush and one of the Daurian Starling. Among the 27 plant species, eight species were

represented by *Ficus*. We observed 282 interactions on *Ficus rumphii*, 80 on *Ficus glaberrima*,

54 on the small-seeded *Aidia densiflora* and 50 on the large-seeded *Caryota mitis*. We

documented 64 unique interactions between plants and frugivores on the island during the study

period. The interaction accumulation curve indicated that we had adequately documented

interaction diversity (Appendix S2: Fig. S2).

During tree watches, we documented 684 interactions (Narcondam Hornbill: 481, Asian Koel:

167, Pied Imperial-Pigeon: 23, Green Imperial-Pigeon: 7, Eye-browed Thrush: 5, Common Hill

Myna: 1), between 11 plant species and six frugivore species. We did not document any

interaction during tree watches which we had not documented during the spot scans. For eight of

the 11 tree species, we observed greater diversity of frugivores foraging on the particular plant

species during spot sampling as compared to tree watches.

Ficus rumphii (among plants) and Narcondam Hornbill (among frugivores) had the highest

degree, species strength and weighted betweenness centrality highlighting the central role they

played in the community during the study period (Appendix S2: Table S6). General linear model

indicated that the species strength in plants was influenced by plant type (figs and non-fig small-,

medium- and large-seeded plants) and not by plant abundance (Adj. $R^2 = 0.40$, $F_{4,22} = 5.322$, $p =$

0.004) (Fig. 3A; Appendix S2: Table S7). *Ficus* were different from other small-seeded plants

and had higher species strength values (mean ($\pm$ SE) = 0.739 ($\pm$ 0.101), n = 8 species) as

compared to small-seeded (mean ($\pm$ SE) = 0.0816 ($\pm$ 0.01), n = 9 species), medium-seeded (mean

($\pm$ SE) = 0.0376 ($\pm$ 0.01), n = 6 species) and large-seeded plant species (mean ($\pm$ SE) = 0.0328 ($\pm$

0.01); n = 4 species) (Fig. 3A). Among frugivores, species strength was significantly and

positively correlated to their abundance (Spearman's $\rho = 0.89$, $p = 0.0123$, n = 7 species; Fig.

3B) and not significantly correlated to their body mass (Spearman's $\rho = 0.71$, $p = 0.088$, n= 7).

The observed modularity of the island plant-disperser community was 0.33, and it was not

significantly different from that obtained through the null models ($p = 0.45$). The observed

community was subdivided into four modules and all the species were consistently allocated to

the same modules across the 25 runs (Table S8). One of the four modules comprised of the

Narcondam Hornbill and several medium and all the four large-seeded plants including

Endocomia macrocomia, *Caryota mitis*, *Pycnarrhaena lucida* and *Planchonella longipetiolata*

highlighting hornbill's importance for large-seeded plants (Table S8).

Comparing the island and the mainland plant-seed disperser communities: The island

network was more asymmetric (-0.588) as compared to the mainland site (0.044). The observed

values of connectance and nestedness on Narcondam were almost twice that of the mainland site

and specialization (H_2') was almost half that of the mainland site (Table 1). The observed

network on Narcondam was less connected and less weighted than the null model predictions but

it was more specialized than expected by random (Table 1).

Community assembly on island and mainland: Among woody plants, we detected a strong

phylogenetic signal for seed size and fruit width in Pakke Tiger Reserve (seed size: Blomberg's

$K = 0.73$, $p = 0.001$; fruit width: Blomberg's $K = 0.43$, $p = 0.005$) and Narcondam (seed size:

Blomberg's $K = 1.48$, $p = 0.001$; fruit width: Blomberg's $K = 0.45$, $p = 0.027$). Among

frugivores, we detected a phylogenetic signal for the body mass of frugivores in Pakke

(Blomberg's $K=1.22$, $p = 0.012$) and Narcondam Island (Blomberg's $K = 1.28$, $p = 0.043$). We

did not detect phylogenetic signal in the percentage of fruits in the diet of the frugivores of Pakke

(Blomberg's $K = 0.23$, $p = 0.38$) and Narcondam (Blomberg's $K = 0.90$, $p = 0.08$). We did not

find any significant correlation for the four models indicating that neither the functional or the

phylogenetic component was able to explain the observed interaction patterns between plants and

frugivores in both island and mainland sites (Table 2; Fig. 4).

DISCUSSION

In this study, we demonstrate the role of large-bodied frugivores, especially the endemic

Narcondam Hornbill, in contributing disproportionately to seed dispersal in the absence of key

small-bodied frugivore groups on the island. Narcondam Hornbill plays a central role in the

plant-dispersal community as it is super-abundant, has very high species strength and

betweenness centrality values and is critical for the select large-seeded plants fruiting. This study

also demonstrates the key role played by figs in the plant-seed disperser community as

demonstrated by its very high species strength values as compared to other plant types. In

comparison to the mainland community, the island community was more asymmetric, connected,

nested and less-specialized. We have documented trait convergence and dietary shifts in

frugivores which potentially contribute to the diffused nature of plant-seed disperser interactions.

This also likely contributes to the absence of the role of phylogenetic or functional component in

the assembly of plant-seed disperser communities on mainland or the island. This is one of the

few studies from oceanic islands of the Indian Ocean and demonstrates how the largest

frugivore, which is a point-endemic, plays a critical quantitative role in seed dispersal. This is

also one of the few studies that performed a paired comparison between an island and mainland

community.

Narcondam for the hornbill and by the hornbill

There is no other place in the world where hornbills are reported to occur in such high densities

as on Narcondam Island (151 birds km^{-2}). On Sulawesi, the Knobbed Hornbills *Rhyticeros*

cassidix occur in very high densities of up to 84 birds km^{-2} (Kinnaird et al. 1996). In north-east

India, Wreathed Hornbill *Rhyticeros undulatus*, a close relative of the Narcondam Hornbill,

seasonally occur in densities of up to 68 birds km^{-2} (Naniwadekar and Datta 2013). Our density

estimates are similar to densities estimated by previous studies (Raman et al. 2013; Manchi

2017). Reduction in hunting pressures, following the suggestions by Sankaran (2000), may have

resulted in the recovery of Narcondam Hornbill population. Among other frugivores, there is

limited information on the density of the Asian Koel from other sites, but comparison with

existing studies show that Narcondam has very high densities of as compared to other sites

(Riley 2001). The mean Imperial-Pigeon densities are slightly lower than those reported from a

mainland site in north-east India (Dasgupta and Hilaluddin 2012).

In this study, multiple measures (abundance and different network metrics – species strength,

degree, betweenness and modularity) indicate that Narcondam Hornbills are keystone seed

dispersers on the island. Generally, as the body mass increases, the abundance of the frugivore

and the frequency of visitation is found to decline (Godínez-Alvarez et al. 2020). Interestingly,

the Narcondam Island is unique where the largest frugivore, the Narcondam Hornbill, is also the

most abundant frugivore on the island. Abundant species have been reported to influence their

partners strongly and Narcondam Hornbill had the highest species strength values highlighting

its importance for plants (Vázquez et al. 2007). Larger frugivores, like hornbills, are known to

remove more fruits per visit as compared to other smaller frugivores (Naniwadekar et al. 2019).

Highest number of interactions of the Narcondam Hornbill coupled with potentially higher fruit

removal rates point towards the key quantitative role played by the hornbill on the island.

Modularity analysis revealed that the hornbill was the key seed disperser for the large-seeded

plants as has been demonstrated in previous studies (Naniwadekar et al. 2019; Gopal et al. 2020).

The hornbill was the only frugivore that has high betweenness centrality value pointing towards

its vital role as a connector between different guilds within the community. The Narcondam

Hornbill was found to be playing a key role in seed dispersal in this asymmetric system. This

high reliance on a frugivore and the potentially low redundancy highlights the fragility of the

island ecosystem because of strong reliance on a single frugivore.

Narcondam has very high densities of hornbill food plants, which likely explains the high

hornbill densities. The overall fig density on Narcondam Island was 27 trees ha^{-1}. The densities

of the six fig species which were consumed by frugivores (including hornbills) was 24 trees ha^{-1}.

This is very high as compared to other sites in the world where the densities of hemi-epiphytic

figs are typically less than one tree ha^{-1} (Harrison 2005). *Ficus* (species consumed by hornbills)

density was 2.7 trees ha^{-1} at a site in north-east India (Datta 2001). *Ficus* (overall) densities were

1.4 trees ha^{-1} at a site in the southern Western Ghats (Page and Shanker 2020). *Ficus* densities of

up to 10 trees ha^{-1} have been reported in Sulawesi (Kinnaird et al. 1996). Consistently across the

mainland site and the island site, certain species of *Ficus* attracted the highest diversity of

frugivores. On the island, *Ficus* represented the top six species in species strength, highlighting

the key role they potentially played in the plant-disperser community at least during the study

period if not throughout the year. Association between canopy, hemi-epiphytic figs and hornbills

has been documented (Harrison et al. 2003). Most dominant fig species during our study on

Narcondam Island, which are important hornbill food plants are mostly canopy, hemi-epiphytic

figs. Being small-seeded, they often play the pioneering role in attracting frugivores on oceanic

islands and accelerating plant colonization on the island (Harrison 2005). Figs and hornbills may

have played a critical role in setting up the plant community on the island which needs further

investigation.

Based on published literature on hornbill food plants from north-east (Datta 2001; Naniwadekar

et al. 2015, 2019) and Thailand (Poonswad et al. 1998; Kanwatanakid-Savini et al. 2009;

Chaisuriyanun et al. 2011), we identified at least 35 species of Narcondam Hornbill food plants.

Overall density of the 35 hornbill food plant species (GBH $\geq$ 30 cm) on Narcondam Island is 327

trees ha^{-1}. In Pakke Tiger Reserve, the density of 45 hornbill food plant species was 163 trees ha^{-}

^{1} (Datta 2001), highlighting the hyper abundance of hornbill food plants on Narcondam Island.

Interestingly, the abundance of food plants is consistently distributed across the entire elevation

gradient. The hyper abundance of frugivores and their food plants makes this island a unique

ecosystem.

Plant-seed disperser communities on island and mainland

Island communities have been reported to be more asymmetric as compared to mainland systems

due to higher immigration rates of plants or human-caused frugivore extinctions on the island

(Schleuning et al. 2014*a*; Nogales et al. 2016). There are no known human-driven extinctions on

the Narcondam island. We are unlikely to find additional species of frugivores that can play an

important functional role in seed dispersal on the island as our list of frugivores is comparable to

the frugivores reported by the previous studies (Raman et al. 2013). We feel that dispersal

limitation of frugivores could also be a key factor in the web asymmetry. Andaman and Nicobar

archipelago of which Narcondam Island is part of, is an interesting example of this. Bulbuls and

barbets, which are the key frugivores on the mainland (Naniwadekar et al. 2019), are absent on

the remote and tiny Narcondam Island. Endemic species of bulbuls are found on other islands in

the Andaman and Nicobar archipelago, but barbets are absent from the archipelago. Thus, the

variation in the dispersal ability of different frugivore groups might also play an important role in

contributing towards the web asymmetry on oceanic islands. Among other network properties,

higher connectance on islands has been reported elsewhere (González-Castro et al. 2012).

Forbidden links are thought to contribute towards lower connectance (Olesen et al. 2011).

Additionally, connectance and nestedness are negatively correlated (Rezende et al. 2007). In our

case, higher connectance is associated with higher nestedness on the island site as compared to

the mainland site. Given that most of the important frugivores on the island are medium (100-

500 g) or large-bodied (> 500 g) with ability to handle small and large fruits, it has likely

resulted in higher connectance and nestedness.

Organization of plant-seed disperser communities

We detected a phylogenetic signal in bird body mass and seed and fruit traits on the island and

the mainland. However, functional and phylogenetic component did not influence the network

interactions in the plant-seed disperser communities on the island and mainland. The role of

evolutionary history in shaping local networks of interacting species is thought to be limited

(Segar et al. 2020). Unlike the plant-pollinator networks, the seed dispersal networks can be

expected to be less specialized and diffused as unrelated species may exhibit trait convergence

and forage on similar plant species (Jordano 2000). We found that distantly related groups of

frugivores dispersed seeds of a common assemblage of plants, many of which are distantly

related to each other. A diverse array of frugivores disperse the super-generalist *Ficus*, and even

the large-seeded plants, which otherwise have a smaller disperser assemblage, were

predominantly dispersed by distant frugivore clades, like hornbills and *Ducula* pigeons, on the

mainland site. Unlike Bastazini et al. (2017), we did not detect signal in the functional

component of the plant-seed disperser communities. This is a likely consequence of the absence

of trait complementarity among interacting frugivores and plants. While the small-bodied

frugivores may not be able to disperse large seeds due to morphological constraints, large-bodied

frugivores like hornbills, fed on plants that were large- and small-seeded resulting in lack of trait

complementarity.

Another important reason could be the behavioral plasticity shown by frugivorous birds while

foraging. In the mainland site, where there are almost 50 frugivore species with a higher

preponderance of small-bodied frugivores, we rarely encountered *Ducula* pigeons on figs that

attracted more than half of the frugivore species found in the area. On the mainland site, *Ducula*

pigeons were mostly observed foraging on medium-seeded and large-seeded plants. On the

island, where there are hardly any small frugivores, we mostly saw the *Ducula* pigeons foraging

on *Ficus*. This behavior of *Ducula* pigeons feeding on figs on the island but less frequently on

the mainland demonstrates phenotypic plasticity in *Ducula* pigeons, likely as a response to lack

of competitive interactions, an aspect that has been seldom documented elsewhere. Additionally,

on Narcondam Island, we documented hornbills feeding on *Callicarpa arborea*, a small-seeded

species which is mostly dispersed by bulbuls and barbets on the mainland site (Naniwadekar et

al. 2019). Asian Koel, which is a rare frugivore in the mainland site, was seen only once feeding

on fruit there. However, on the island, it is the second-most important frugivore. It remains to be

determined, whether exploitative competition on the mainland site drives larger frugivores to

feed on different resources in the mainland forested system. While habitat fragmentation often

results in loss of large-bodied frugivores thereby affecting large-seeded plants, it will be

interesting to investigate the implications of the absence of small-bodied frugivore on the

structuring of plant communities.

Bats are known to play an important role in seed dispersal on oceanic islands (McConkey and

Drake 2015). We recorded spat out wads of pulp of three *Ficus* spp., *Balakata*, and

Planchonella. Direct detections of foraging bats were mostly near our camp on fruiting

Manilkara zapota (planted), *Terminalia catappa*, *Ficus rumphii* and *Ardisia*. We failed to detect

foraging bats during the six night-walks that lasted for at least 90 minutes each. Role of bats in

seed dispersal on the island needs to be evaluated in the future.

One may argue about the lack of phenological completeness of the study and its influence on the

inferences drawn. The abundances of the frugivores, which is an independent measure used to

demonstrate the relative importance of frugivores, is unlikely to change since it is a tiny and

remote island in the middle of the Andaman Sea. Narcondam Hornbill is likely to remain the

most abundant frugivore on the island year-round and thereby play a key role in seed dispersal

throughout the year. Additionally, the frugivore species composition on the island is unlikely to

change as previous studies have documented similar assemblage of frugivores. The frugivore

data is relatively robust, and we have had a diverse representation of plant families in our study

with varying fruit (figs, drupes and arillate dehiscent capsular fruits) and seed traits thereby

capturing the functional diversity in traits of fruits and seeds making the data amenable for

ecophylogenetic analysis. Additionally, this is a remote island making it extremely challenging

to conduct long bouts of fieldwork, thereby posing challenges to collect long-term data.

Only those plants and animals that are able to cross the oceanic barriers and establish contribute

to structuring of communities on islands. Degree of isolation, area and age of islands thereby

offers an interesting gradient to understand the relative importance of processes in structuring of

communities. As demonstrated in this study, ;arge-bodied frugivores, particularly the Narcondam

Hornbill, play a pivotal role on Narcondam Island in the absence of small frugivores on a

relatively young and moderately isolated island. Several threatened hornbill species are found on

oceanic islands (IUCN 2019). Narcondam Island and its endemic hornbill give insights into the

potential role of hornbills, which are among the largest frugivores on islands, in shaping and

potentially maintaining the island tree communities by playing a key role as a seed disperser. The

ecological role of hornbills and their contribution to mutualism and maintenance of plant species

diversity is poorly understood from other oceanic islands. Narcondam Island through the hyper

abundance of hornbills and its food plants and the asymmetric nature of the plant-disperser

community with heavy reliance on a single frugivore demonstrates the unique ecological and

evolutionary setting of oceanic islands. The disproportionate and singular role that species can

play on oceanic islands points towards the extreme vulnerability of oceanic islands to any

external perturbations. The perturbations can disrupt the delicate and intricate relationships

between plants and seed dispersers, with dramatic cascading impacts on the entire community.

Unfortunately, studies like this might not be possible in many other oceanic islands that harbor

hornbills as they have been already modified extensively due to anthropogenic activities. It is

indeed fortunate that Narcondam Island has been preserved as a Wildlife Sanctuary thereby

safeguarding not only a unique hornbill species but also a unique ecosystem.

ACKNOWLEDGEMENTS

We are grateful to the Andaman and Nicobar Forest Department for giving us the permit (No:

WII/NVP/NARCONDAM/2019). We thank Mr. Dependra Pathak, DGP (A&N) for giving us the

necessary permissions. We thank Commandant A. K. Bhama and Captain Kundan Singh from

the Indian Coast Guard for giving us permission and support. We thank Mr. Abhishek Dey, DC

(South Andamans) for giving us permission.We thank Kulbhushansingh Suryawanshi, Divya

Mudappa and T. R. Shankar Raman for providing us field equipment and for useful discussions.

We thank Mr. D. M. Shukla (PCCF, Wildlife), Mr. A. K. Paul and Mr. Soundra Pandian for their

support. We are grateful to the Dean and the then Director WII, Dr. G. S. Rawat for facilitating

the research permit application and supporting the project. We are indebted to the Special Armed

Police unit led by Ms. Usha Rangnani (SP) for providing us logistic support at Narcondam

Island. We thank Elrika D'Souza, Evan Nazareth, Rachana Rao and Rohan Arthur for providing

us logistic support in Port Blair. We thank Prasenjeet Yadav, Adarsh Raju, Suri Venkatachalam,

Hari Sridhar, Shashank Dalvi, Anand Osuri, Narayan Sharma and Aparajita Datta for discussions

and support.

Funding - We thank Wildlife Conservation Trust, IDEAWILD, Nature Conservation

Foundation, Mr. Uday Kumar, MMMRF, Mr. Rohit and Deepa Sobti, and Mr. Arvind Datar for

providing funding support.

Author contributions - RN conceived the idea and developed with VR, NP, SG, AG and JJ;

RN, AG, SG and NP collected the data; RN and JJ analyzed the data with inputs from AG, NP

and VR; RN wrote the manuscript with inputs from JJ, AG, NP, SG and VR. All authors gave

final approval for publication.

Conflicts of interest – No conflicts of interest to declare.

LITERATURE CITED

Abdulali, H. 1971. Narcondam Island and notes on some birds from the Andaman Islands.

Journal of Bombay Natural History Society 68:385–411.

Barber, M. J. 2007. Modularity and community detection in bipartite networks. Physical Review

E 76:066102.

Bascompte, J., P. Jordano, and J. M. Olesen. 2006. Asymmetric coevolutionary networks

facilitate biodiversity maintenance. Science 312:431–433.

Bastazini, V. A. G., P. M. A. Ferreira, B. O. Azambuja, G. Casas, V. J. Debastiani, P. R.

Guimarães, and V. D. Pillar. 2017. Untangling the tangled bank: a novel method for partitioning

the effects of phylogenies and traits on ecological networks. Evolutionary Biology 44:312–324.

BirdLife International. 2017. Rhyticeros narcondami (amended version of 2016 assessment). The

IUCN Red List of Threatened Species 2017: e.T22682531A110038017.

https://dx.doi.org/10.2305/IUCN.UK.2017-1.RLTS.T22682531A110038017.en.

Blomberg, S. P., T. Garland, and A. R. Ives. 2003. Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. Evolution 57:717–745.

Blüthgen, N., F. Menzel, T. Hovestadt, B. Fiala, and N. Blüthgen. 2007. Specialization,

constraints, and conflicting interests in mutualistic networks. Current Biology 17:341–346.

Chaisuriyanun, S., G. A. Gale, S. Madsri, and P. Poonswad. 2011. Food consumed by Great

Hornbill and Rhinoceros Hornbill in tropical rainforest, Budo Su-Ngai Padi National Park,

Thailand. The Raffles Bulletin of Zoology 24:123–135.

Corlett, R. T. 2017. Frugivory and seed dispersal by vertebrates in tropical and subtropical Asia:

An update. Global Ecology and Conservation 11:1–22.

Dasgupta, S., and Hilaluddin. 2012. Differential effects of hunting on populations of hornbills

and imperial pigeons in the rainforests of the Eastern Indian Himalaya. Indian Forester 138:902–

909.

Datta, A. 2001. *An ecological study of sympatric hornbills and fruiting patterns in a tropical*

forest in Arunachal Pradesh (PhD Dissertation). Saurashtra University, Gujarat, India, Gujarat,

India.

Debastiani, V. J., and V. D. Pillar. 2012. SYNCSA: R tool for analysis of metacommunities

based on functional traits and phylogeny of the community components. Bioinformatics 28.

Donatti, C. I., P. R. Guimarães, M. Galetti, M. A. Pizo, F. M. D. Marquitti, and R. Dirzo. 2011.

Analysis of a hyper-diverse seed dispersal network: modularity and underlying mechanisms:

Modularity in a seed dispersal network. Ecology Letters 14:773–781.

Dormann, C. F., B. Gruber, and J. Fründ. 2008. Introducing the bipartite package: analysing

ecological networks. R News 8:8–11.

Escribano-Avila, G., C. Lara-Romero, A. Traveset, and R. Heleno. 2018. Tropical seed dispersal

networks: emerging patterns, biases, and keystone species traits. *in* W. Dattilo and V. Rico-Gray,

eds. Ecological Networks in the Tropics. Springer, Cham.

Godínez-Alvarez, H., L. Ríos-Casanova, and B. Peco. 2020. Are large frugivorous birds better

seed dispersers than medium- and small-sized ones? Effect of body mass on seed dispersal

effectiveness. Ecology and Evolution accepted.

González-Castro, A., A. Traveset, and M. Nogales. 2012. Seed dispersal interactions in the

Mediterranean Region: contrasting patterns between islands and mainland. Journal of

Biogeography 39:1938–1947.

Gopal, A., D. Mudappa, T. R. S. Raman, and R. Naniwadekar. 2020. Forest cover and fruit crop

size differentially influence frugivory of select rainforest tree species in Western Ghats, India.

Biotropica in press.

Guimarães Jr, P. R., P. Jordano, and J. N. Thompson. 2011. Evolution and coevolution in

mutualistic networks. Ecology Letters 14:877–885.

Harrison, R. D. 2005. Figs and the diversity of tropical rainforests. BioScience 55:1053–1064.

Harrison, R. D., A. A. Hamid, T. Kenta, J. Lafrankie, H.-S. Lee, H. Nagamasu, T. Nakashizuka,

et al. 2003. The diversity of hemi-epiphytic figs (Ficus; Moraceae) in a Bornean lowland rain

forest. Biological Journal of the Linnean Society 78:439–455.

Heinen, J. H., E. E. van Loon, D. M. Hansen, and W. D. Kissling. 2018. Extinction-driven

changes in frugivore communities on oceanic islands. Ecography 41:1245–1255.

IUCN. 2019. The IUCN Red List of Threatened Species. Version 2019-1.

http://www.iucnredlist.org.

Jetz, W., G. H. Thomas, J. B. Joy, K. Hartmann, and A. O. Mooers. 2012. The global diversity of

birds in space and time. Nature 491:444–448.

Jordano, P. 2000. Fruits and frugivory. Pages 125–166 *in* M. Fener, ed. Seeds: The Ecology of

Regeneration in Plant Communities. CABI Publishing, Wallingford.

Kaiser-Bunbury, C. N., A. Traveset, and D. M. Hansen. 2010. Conservation and restoration of

plant–animal mutualisms on oceanic islands. Perspectives in Plant Ecology, Evolution and

Systematics, Comparative ecological research on oceanic islands 12:131–143.

Kanwatanakid-Savini, C., P. Poonswad, and T. Savini. 2009. An assessment of food overlap

between gibbons and hornbills. The Raffles Bulletin of Zoology 57:189–198.

Kinnaird, M. F., T. G. O'Brien, and S. Suryadi. 1996. Population fluctuation in Sulawesi Red-

Knobbed Hornbills: tracking figs in space and time. The Auk 113:431–440.

Manchi, S. 2017. *Status, ecology and conservation of Narcondam Hornbill Aceros narcondami*

on the Narcondam Island, India. Salim Ali Centre for Ornithology and Natural History,

Coimbatore, Tamil Nadu, India.

Marquitti, F. M. D., P. R. Guimarães, M. M. Pires, and L. F. Bittencourt. 2014. MODULAR:

software for the autonomous computation of modularity in large network sets. Ecography

37:221–224.

McConkey, K. R., and D. R. Drake. 2015. Low redundancy in seed dispersal within an island

frugivore community. AoB PLANTS 7.

Mello, M. A. R., F. A. Rodrigues, L. da F. Costa, W. D. Kissling, Ç. H. Şekercioğlu, F. M. D.

Marquitti, and E. K. V. Kalko. 2015. Keystone species in seed dispersal networks are mainly

determined by dietary specialization. Oikos 124:1031–1039.

Miller, D. L., E. Rexstad, L. Thomas, L. Marshall, and J. Laake. 2019. Distance sampling in R.

Journal of Statistical Software 89:1–28.

Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca, and J. Kent. 2000.

Biodiversity hotspots for conservation priorities. Nature 403:853–858.

Naniwadekar, R., S. Chaplod, A. Datta, A. Rathore, and H. Sridhar. 2019. Large frugivores

matter: insights from network and seed dispersal effectiveness approaches. Journal of Animal

Ecology 88:1250–1262.

Naniwadekar, R., and A. Datta. 2013. Spatial and temporal variation in hornbill densities in

Namdapha Tiger Reserve, Arunachal Pradesh, north-east India. Tropical Conservation Science

6:734–748.

Naniwadekar, R., C. Mishra, and A. Datta. 2015. Fruit resource tracking by hornbill species at

multiple scales in a tropical forest in India. Journal of Tropical Ecology 31:477–490.

Nogales, M., R. Heleno, B. Rumeu, A. González-Castro, A. Traveset, P. Vargas, and J. M.

Olesen. 2016. Seed-dispersal networks on the Canaries and the Galápagos archipelagos:

interaction modules as biogeographical entities: Modularity of seed-dispersal networks on

oceanic islands. Global Ecology and Biogeography 25:912–922.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, R. B. O'Hara, et

al. 2019. vegan: Community Ecology Package.

Olesen, J. M., J. Bascompte, Y. L. Dupont, H. Elberling, C. Rasmussen, and P. Jordano. 2011.

Missing and forbidden links in mutualistic networks. Proceedings of the Royal Society B:

Biological Sciences 278:725–732.

Page, N. V., and K. Shanker. 2020. Climatic stability drives latitudinal trends in range size and

richness of woody plants in Western Ghats, India. PLOS ONE 15:e0235733.

Palacio, R. D., C. Valderrama-Ardila, and G. H. Kattan. 2016. Generalist species have a central

role in a highly diverse plant-frugivore network. Biotropica 48:349–355.

Paradis, E., and K. Schliep. 2018. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in {R}. Bioinformatics 35:526–528.

Peralta, G. 2016. Merging evolutionary history into species interaction networks. Functional

Ecology 30:1917–1925.

Poonswad, P., A. Tsuji, N. Jirawatkavi, and V. Chimchome. 1998. Some aspects of food and

feeding ecology of sympatric hornbill species in Khao Yai National Park, Thailand. Pages 137–

157 *in* P. Poonswad, ed. The Asian Hornbills: Ecology and Conservation., Thai studies in

biodiversity no. 2. BIOTEC, NSTDA.

Qian, H., and Y. Jin. 2016. An updated megaphylogeny of plants, a tool for generating plant

phylogenies and an analysis of phylogenetic community structure. Journal of Plant Ecology

9:233–239.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Raman, T. R. S., D. Mudappa, T. Khan, U. Mistry, A. Saxena, K. Varma, N. Ekka, et al. 2013.

An expedition to the Narcondams: observations of the marine and terrestrial fauna including the

island-endemic hornbill. Current Science 105:346–360.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217–223.

Rezende, E. L., P. Jordano, and J. Bascompte. 2007. Effects of phenotypic complementarity and

phylogeny on the nested structure of mutualistic networks. Oikos 116:1919–1929.

Riley, J. 2001. Population sizes and the status of endemic and restricted-range bird species on

Sangihe Island, Indonesia. Bird Conservation International 12:53–78.

Sankaran, R. 2000. Narcondam Hornbill Aceros narcondami. Pages 57–66 *in* L. Vijayan, R.

Sankaran, K. Sivakumar, and V. Murugan, eds. A study on the ecology, status and conservation

perspectives of certain rare endemic avifauna of the Andaman and Nicobar Islands. Report

submitted to Salim Ali Centre for ornithology and natural History, Coimbatore, India.

Coimbatore, India.

Schleuning, M., K. Böhning-Gaese, D. M. Dehling, and K. C. Burns. 2014*a*. At a loss for birds:

insularity increases asymmetry in seed-dispersal networks. Global Ecology and Biogeography

23:385–394.

Schleuning, M., L. Ingmann, R. Strauß, S. A. Fritz, B. Dalsgaard, D. Matthias Dehling, M. Plein,

et al. 2014*b*. Ecological, historical and evolutionary determinants of modularity in weighted

seed-dispersal networks. (J. Gomez, ed.)Ecology Letters 17:454–463.

Segar, S. T., T. M. Fayle, D. S. Srivastava, T. M. Lewinsohn, O. T. Lewis, V. Novotny, R. L.

Kitching, et al. 2020. The role of evolution in shaping ecological networks. Trends in Ecology &

Evolution 35:454–466.

Shanahan, M., S. So, S. G. Compton, and R. Corlett. 2001. Fig-eating by vertebrate frugivores: a

global review. Biological Reviews 76:529–572.

Terborgh, J., N. Pitman, M. Silman, H. Schichter, and P. Nunez V. 2002. Maintenance of tree

diversity in tropical forests. Seed dispersal and frugivory: ecology, evolution and conservation.

CABI Publishing, Wallingford.

Thomas, L., S. T. Buckland, E. A. Rexstad, J. L. Laake, S. Strindberg, S. L. Hedley, J. R. B.

Bishop, et al. 2010. Distance software: design and analysis of distance sampling surveys for

estimating population size. Journal of Applied Ecology 47:5–14.

Vázquez, D. P., N. Blüthgen, L. Cagnolo, and N. P. Chacoff. 2009. Uniting pattern and process

in plant–animal mutualistic networks: a review. Annals of Botany 103:1445–1457.

Vázquez, D. P., C. J. Melián, N. M. Williams, N. Blüthgen, B. R. Krasnov, and R. Poulin. 2007.

Species abundance and asymmetric interaction strength in ecological networks. Oikos 116:1120–

1127.

Wilman, H., J. Belmaker, J. Simpson, C. de la Rosa, M. M. Rivadeneira, and W. Jetz. 2014.

EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. Ecology

95:2027–2027.

Table 1. Network-level properties (observed and null model values) for the Narcondam Island,

and observed values for the mainland site, Pakke Tiger Reserve. Null model values for the

mainland site can be found in Naniwadekar et al. (2019a).

	Observed value: Narcondam	**Null mean: Narcondam**	**Null SD: Narcondam**	**Observed value: Mainland**
Connectance	0.339	0.463	0.018	0.154
Specialization (H_2')	0.233	0.144	0.009	0.500
Weighted NODF	45.178	59.76	3.834	24.885

683 **Table 2.** Correlation coefficients for the two models that evaluate the influence of the functional

684 (after removing phylogenetic signal) and phylogenetic components on the observed patterns of

685 interactions between plants and frugivores for the mainland site (Pakke Tiger Reserve) and the

686 island site (Narcondam Wildlife Sanctuary). None of the values were significant at $p = 0.05$.

Parameter	ρ (Mainland)	ρ (Island)
Functional component (after removing phylogenetic signal)	0.04	0.03
Phylogenetic component	0.023	-0.17

 FIGURES

Figure 1. Schematic illustrating mainland and island plant and seed-dispersal communities. A

map of the South and Southeast Asia showing mainland and island site (a). The island endemic

large-bodied frugivore Narcondam Hornbill *Rhyticeros narcondami* on *Caryota mitis* (b).

Organization of plant-seed disperser community on mainland and island. Large-bodied

frugivores, many of which predominantly feed on large-seeded plants, can exhibit ecological

release on the island (in the absence of small frugivores) resulting in dietary shifts. Dietary

changes will in turn result in changes in interaction frequencies with corresponding changes in

network structure. Painting and sketches by Sartaj Ghuman. Map source: Google Earth.

Figure 2. Abundance of Narcondam Hornbill *Rhyticeros narcondami*, Asian Koel *Eudynamys*

scolopaceus and Imperial-Pigeons (Pied Imperial-Pigeon *Ducula bicolor* and Green Imperial-

Pigeon *Ducula aenea*) on Narcondam Island (A). Density (km^{-1}) of Narcondam Hornbill on the

island and across the three elevation zones on the island (B). Density (ha^{-1}) of 27 species of

fleshy-fruited plants that were fruiting across the three elevation zones during the study period

(C).

Figure 3. Relationship between species strength ($\log_{10}$ transformed in the case of plants) and

plant type (A) and frugivore abundance (B). We separated figs and small-seeded plants for they

consistently had higher species strength values. Frugivore codes - ASKO: Asian Koel, CHMY:

Common Hill Myna, DAST: Daurian Starling, EBTH: Eye-browed Thrush, GIPI: Green

Imperial-Pigeon, NAHO: Narcondam Hornbill and PIPI: Pied Imperial-Pigeon.

Figure 4. Seed dispersal interactions in Narcondam Island (A) and the mainland site, Pakke

Tiger Reserve (B). Also shown is the pruned phylogeny for the plants and the frugivores and

respective traits for plants (seed size and fruit width) and frugivores (body mass and percentage

diet that is comprised of fruits).

Figure 1.

Figure 2.

Figure 3.