Title: Novel therapeutic targets to mitigate early neuronal dysfunction and cognitive deficits in tauopathy

Authors: Shon A. Koren[1], Matthew J. Hamm[1], Ryan Cloyd[2], Sarah N. Fontaine[2], Emad Chishti[2], Chiara Lanzillotta[3], Fabio Di Domenico[3], Nicholas T. Seyfried[4], Duc Duong[4], David Powell[5], Moriel Vandsburger[6], Anika M.S. Hartz[2], John Koren 3rd, [1], Jeffrey M. Axten[7], Nicholas J. Laping[7], Jose F. Abisambra[1],*

Affiliations:

[1]Department of Neuroscience & Center for Translational Research in Neurodegenerative Disease, University of Florida, Gainesville, FL, USA.

[2]Sanders Brown Center on Aging & Department of Physiology, University of Kentucky, Lexington, KY, USA.

[3]Department of Biochemical Sciences, Sapienza University of Rome, Rome, Italy.

[4]Department of Biochemistry and Emory Integrated Proteomics Core, Emory University School of Medicine, Atlanta, GA, USA.

[5]Magnetic Resonance Imaging and Spectroscopy Center, University of Kentucky, Lexington, KY, USA.

[6]Department of Bioengineering, University of California, Berkeley, CA, USA.

[7]GlaxoSmithKline, Research and Development, Collegeville, PA, USA.

*To whom correspondence should be addressed: j.abisambra@ufl.edu

One Sentence Summary: Multi-target kinase inhibition rescues cognitive function in early stage tauopathy mice and reverses proteomic shifts common to Alzheimer's disease in humans.

Abstract: Tauopathies are a group of more than twenty known disorders that involve progressive neurodegeneration, cognitive decline, and pathological tau accumulation. Current therapeutic strategies provide only limited, late-stage symptomatic treatment. This is partly due to lack of understanding of the molecular mechanisms linking tau and cellular dysfunction, especially during the early stages of disease progression. In this study, we treated early stage tau transgenic mice with a multi-target kinase inhibitor to identify novel substrates that contribute to cognitive impairment and exhibit therapeutic potential. Drug treatment significantly ameliorated brain atrophy and cognitive function as determined by behavioral testing and a sensitive imaging technique called manganese-enhanced magnetic resonance imaging (MEMRI) with quantitative R1 mapping. Surprisingly, these benefits occurred despite unchanged hyperphosphorylated tau levels. To elucidate the mechanism behind these improved cognitive outcomes, we performed quantitative proteomics to determine the altered protein network during this early stage in tauopathy and compare this model with the human AD proteome. We identified a cluster of preserved pathways shared with human tauopathy with striking potential for broad multi-target kinase intervention. We further report high confidence candidate proteins as novel therapeutically relevant targets for the treatment of tauopathy.

Introduction

Tauopathies, the most common of which is Alzheimer's disease (AD), are a group of

neurological disorders defined by the neuropathological accumulation of tau protein that present

progressive cognitive dysfunction and brain atrophy. No cure for tauopathies exists, and current

treatment strategies are palliative *(1, 2)*. Several clinical trials have targeted toxic tau species

through immunotherapies or by inhibiting tau post-translational modifications and fibrillization

with mixed results *(1–4)*. However, the disease etiology of tauopathies, AD in particular, are

almost certainly multifactorial. A promising strategy to mitigate the complex nature of tauopathies,

similar to established approaches for other chronic, progressive diseases, encompass multi-target

or combinatorial treatments *(3, 5)*. However, these strategies are limited by requiring a thorough

understanding of the cellular perturbations that underly the disease.

Modern systems biology approaches provide unparalleled power in investigating the

cellular alterations in disease. These tools highlight promising candidate targets for novel

therapeutics while elucidating the pathophysiology of the disease. Recent studies focused on the

molecular network changes in AD suggest concurrent changes in energy metabolism, immune

response, synapse activity, cytoskeletal stability, and RNA metabolism pathways present early in

disease progression *(6–8)*. These reports validate decades of research utilizing *in vitro* and *in vivo*

disease models. Using these tools, the pathways most closely associated with early cognitive

decline and neurodegeneration can be determined and used to assess novel therapeutics.

Importantly, these resources have not yet been applied to establish the efficacy of multi-target

drugs in mitigating cognitive decline in animal models of tauopathy.

Aberrant phosphorylation and kinase signaling is a hallmark of AD and other tauopathies

(8, 9). Here, we report the use of a multi-target kinase inhibitor, GSK2606414 or 414, to treat tau

transgenic mice in a proof-of-concept study to evaluate whether such strategies can mitigate the

early negative outcomes of tauopathy. This compound potently inhibits multiple tyrosine and

serine/threonine kinases at low micromolar concentrations *(10)* and ameliorates phenotypes in a

variety of neurological disorder and neurodegenerative models *(11–16)*. The targets of 414 include

kinases involved in tauopathy pathogenesis, including protein kinase R-like endoplasmic

reticulum kinase, or PERK *(10–12)*, the MAPK cascade *(10)*, receptor-interacting

serine/threonine-protein kinase 1, or RIPK1 *(17)*, and KIT *(18)*. We show that brain atrophy and

phenotypes of cognitive dysfunction evident in this mouse model of tauopathy are substantially

reduced by treatment with 414. Using manganese-enhanced MRI (MEMRI) with parametric

mapping, we show that multi-target kinase inhibition rescues deficits in hippocampal calcium

activity. The rescue of tauopathy occurs in the absence of changes to toxic, hyper-phosphorylated

tau levels and without inhibition of PERK, the target canonically associated with the

neuroprotective benefits of 414 *(10, 11)*. We use quantitative proteomics to investigate both the

cellular alterations and the networks responsive to multi-target kinase inhibition associated with

cognitive rescue in this early stage tauopathic mouse model. Finally, we identify novel candidate

proteins that are rescued by drug treatment and are consistent between the tauopathy mouse and

human proteome. These analyses identify novel candidate targets for therapeutic intervention in

future studies.

Results

Tau transgenic mice exhibit common neurodegenerative features ameliorated by multi-target
kinase inhibition

To investigate the pathways responsible for early cognitive impairment in tauopathy and

their amenability to treatment with a multi-target kinase inhibitor, we used the rTg4510 tau

transgenic mouse model of frontotemporal lobar degeneration (FTLD). The rTg4510 (Tg) mice

exhibit well-characterized disease progression driven by the overexpression of P301L human tau

primarily in forebrain neurons *(19–21)*. Neurofibrillary tau pathology appears as early as two

months, with progressively worsening cognitive impairment detectable as early as two and a half

months. Additionally, significant brain atrophy is detectable in this model by 5 months of age (Fig.

1A).

We treated four-month-old Tg and non-transgenic control (NTg) mice with the multi-target

kinase inhibitor GSK2606414 (414) for 30 days. Four-month-old Tg mice display significant tau

burden, but they do not present with severe cognitive impairment and therefore provide a

therapeutic window amenable to treatment strategies. T2-weighted MRI (Fig. 1B-F) revealed that

Tg mice treated with vehicle (Tg + Veh) exhibited an approximately 20% reduction in total brain

volume compared with non-transgenic control mice treated with either vehicle (NTg + Veh, **p

= 0.0063) or 414 (NTg + 414, **p = 0.0012). Surprisingly, tau transgenic mice treated with 414

(Tg + 414) exhibited only a 10% reduction in total brain volume that lacked statistical significance

from NTg + Veh control animals (ns, p > 0.15).

Having observed that 414 treatment decreased brain atrophy in the Tg mice, we then

examined if treatment with 414 also rescued cognitive deficits. To analyze changes in neuronal

activity *in vivo*, we performed an imaging approach that measures brain function using manganese-

enhanced MRI (MEMRI) with parametric mapping *(22, 23)* . MEMRI utilizes manganese as a

contrast agent that can flow through calcium channels such as voltage-gated calcium channels *(24–*

26) and N-methyl-d-aspartate (NMDA) receptors *(27)* that lengthens R1 relaxation rates in tissue

(28). When MEMRI is combined with an imaging protocol to quantify the change in R1 relaxation

rates over time, or ΔR1, the acquired data reflects calcium dynamics as a measure of neuronal

activity in the living brain *(23, 28, 29)*.

We previously established that this tau transgenic mouse line exhibits altered calcium

homeostasis detected by MEMRI as early as 3mo of age which continues as the mice age *(22)*.

Here, we corroborated those findings, identifying altered ΔR1 in 5mo Tg mice (Fig. 2A-D). The

strongest differences in ΔR1 observed were found in hippocampal regions such as the dentate

gyrus (DG, Fig. 2E) and Cornu ammonis area 1 (CA1, Fig. 2F). In both regions, our experimental

Tg + Veh mice exhibited significantly decreased ΔR1 when compared to the control NTg + Veh

mice (DG, ***p = 0.0003; CA1, *p = 0.0226). Tg + 414 treated mice displayed no apparent

impairment in calcium activity measured by MEMRI in both regions relative to NTg + Veh mice

(DG and CA1, ns) and displayed significant improvement in calcium activity relative to the Tg +

Veh animals (DG, **p = 0.0019; CA1, *p = 0.0211). These findings demonstrate that, in tandem

with the changes in total brain volume, treatment with the multi-target kinase inhibitor 414 rescued

hippocampal activity deficits in this model.

Since calcium dyshomeostasis underlies cognitive impairments in tauopathy *(30)*, we

evaluated the effects of 414 in two behavioral paradigms: open field as a measure of anxiety-like

phenotype, and Y-maze as a measure of cognition. The open field behavioral paradigm

demonstrated a Tg phenotype wherein Tg + Veh mice spent more time in the field periphery than

either NTg group (NTg + Veh: **p = 0.0012; NTg + 414: ***p = 0.0001), suggesting a transgene-

dependent anxiety-like behavior (Fig. 3G). Concurrent with our brain volume and calcium

signaling findings, Tg mice treated with 414 spent significantly less time in the periphery than Veh

treated Tg mice (*p=0.0260) and were statistically indistinguishable from either NTg treatment

group (ns, p > 0.15). In the Y-maze, Tg + Veh mice exhibited decreased novel arm entries (*p =

0.0365) compared to NTg control mice. Treatment of Tg animals with 414, though statistically

indistinguishable from the NTg groups (p > 0.15, ns) only partially rescued the number of entries

(ns, p = 0.1095) compared to vehicle-treated Tg mice (Fig. 2H). Additional Y-maze analysis

parameters, such as the total time spent in the novel arm, indicated Tg + Veh mice had reduced

performance compared to NTg + Veh mice (*p = 0.0433). Again, the Tg + 414 treatment group

was statistically indistinguishable from either NTg group (ns, p > 0.15) but displayed an

incomplete rescue of this phenotype as compared to Tg + Veh mice (ns, p = 0.0882) (Supplemental

Fig. 1). Consistent with our MEMRI and brain volume data, these data demonstrated that treatment

with 414 mitigates the cognitive deficits of early-stage tauopathy in this model.

GSK2606414 rescues functional deficits without altering tau hyper-phosphorylation

140   Reduction of tau pathology typically correlates and is interpreted as the reason why novel

therapeutic strategies improve tauopathy outcomes. Previously, GSK2606414 had been reported

to reduce tau pathology by preventing PERK activation and consequent enhancement of GSK3β

activity *(11)*. GSK3β phosphorylates tau at sites that are associated with disease *(31)*. To test

whether the reduction of tau was responsible for the cognitive rescue in Tg mice, we measured

changes in the levels of disease-associated, hyper-phosphorylated tau at the S396/S404 (PHF1)

epitope along with total tau levels (Fig. 3A). Quantification of immunoblots indicated that 414 did

not modify the levels of PHF1, total tau, nor the relative phosphorylation at the PHF1 epitope (Fig.

3B-D). We further measured changes in the levels of active, Y216-phosphorylated and total

GSK3β (Fig. 3E) and found that 414 similarly did not modify the levels of activated GSK3β (Fig.

3F). These data suggest that the neuroprotective benefits conferred by 414 in these Tg mice are

not directly related to changes in tau or phosphorylated tau, but rather through alternative targets.

Functional deficits of early stage tau transgenic mice do not depend on PERK/UPR activation

GSK2606414 was developed as a first-in-class inhibitor of PERK *(10)*. PERK, an ER-

associated kinase, functions as part of the unfolded protein response (UPR) by attenuating global

translation and dampening the level of incoming nascent proteins in the ER under conditions of

ER stress *(32)*. However, long term PERK activation leads to sustained suppression of translation

and activation of pro-apoptotic signaling cascades. Previous studies have used 414 to inhibit PERK

to rescue or prevent neurodegeneration by limiting the deleterious chronic activation of the UPR

(11, 12, 32). We evaluated whether PERK inhibition was a primary factor in the neuroprotective

effects of 414 treatment in these early stage Tg mice. Consistent with several reports showing

PERK activity does not begin until late-stage tauopathy progression in this model *(11, 33, 34)*,

PERK activation (as measured by the phosphorylation of T980) was minimally detected by

immunofluorescent stains of the hippocampus and cortex of 5mo Tg mice (Fig. 3G-H). At 8mo,

Tg mice had several thousand-fold greater levels of phosphorylated PERK (Fig. 3I-J) which was

inhibited by 414 treatment as previously reported (Fig. 3K) *(11)*. We used quantitative real-time

PCR gene card arrays to verify our immunofluorescence findings by monitoring transcript levels

of genes directly altered by PERK activation. The expression levels of genes downstream of PERK

were largely unchanged in the 5mo Tg animals compared to NTg controls, and they were

unmodified by 414. These data confirm the absence of strong PERK activity (Fig. 3L). We also

measured other mediators of the UPR distinct from the PERK pathway such as ATF6, IRE1, and

other eIF2α kinases. Only the transcriptional regulator ATF6 exhibited a significant change in

transcript levels in Tg mice, with a reduction of ATF6 mRNA levels in Tg + Veh mice which was

Tg mice treated with 414 (Supplemental Fig. 2). PERK reportedly switches preferential substrate

phosphorylation in tauopathies between eIF2α and other targets such NRF2, a transcription factor

which undergoes nuclear translocation and induces the expression of redox response proteins when

phosphorylated by PERK *(35)*. Consistent with our other data, 414 did not alter NRF2 nuclear

translocation in Tg model mice (Fig. 3M-N). Considering the off-target effects of kinase inhibitors,

and those targeted by 414 beyond PERK at nanomolar and low micromolar concentrations (Table

S1), we reasoned that the neuroprotective effect of 414 is likely due to a multi-target response

distinct from inhibiting PERK kinase activity.

Quantitative hippocampal proteomics reveals pathways responsible for cognitive decline in early

stage tauopathy mice

To assess the effects of a compound impacting multiple cellular pathways simultaneously,

we turned to quantitative proteomics to identify proteins and pathways most impacted by

transgenicity and our test agent, 414. Despite the prevalence of the rTg4510 model in the study of

neurodegeneration, no quantitative analysis of the brain proteome in these mice has been reported.

To this end, we evaluated the hippocampal proteome of NTg and Tg mice treated with either Veh

or 414 (n = 4 per group, all female) using a multiplexed tandem mass tagging (TMT) approach

(Fig. 4A). Following protein quantification, batch normalization, and statistical analyses, 337

proteins were identified as significantly altered across all group comparisons (FDR adjusted p

value < 0.05, Fig. 4B). First, we identified the impact of 414 treatment on proteins in NTg animals

(termed: "Drug Effect") and removed these proteins from further analysis. Next, we determined

which proteins were altered due to Tg genotype (termed: "Tg Effect") by comparing NTg + Veh

vs Tg + Veh samples and specifically excluded any genes known to be dysregulated due to

transgene insertion in this model *(36, 37)*. We then identified the proteins that were rescued by 414

treatment in the Tg mice: proteins in the "Tg Effect" group that were restored to statistically normal

levels in Tg mice (no significant difference from NTg + Veh mice) following treatment with 414

(termed: "Drug Rescued"). These groupings are summarized in (Fig. 4C) and representative

proteins belonging to each group are shown in (Fig. 4D). Interestingly, ≈67% of the proteins altered

in Tg mice were rescued after treatment with 414 (186 out of 276). Importantly and as expected,

the human *MAPT* P301L transgene was significantly upregulated in Tg mice and unaffected by

414 (Fig. 4E) thus confirming the tau immunoblots and validating the proteomic results.

Next, we analyzed the pathway level changes in the 5mo Tg brain proteome. To collapse

the altered protein into pathways, we used g.Profiler *(38)* to assess the changes in Reactome *(39)*

annotated pathways (FDR adjusted p value < 0.1, Table S2). Cytoscape *(40)* was used to cluster

each pathway into larger umbrella networks that were down-regulated (Fig. 4F) or up-regulated

(Fig. 4E) in Tg mice. The down-regulated protein network clustered into nine discrete groups,

namely: mTOR pathway activity, synaptic activity and long-term potentiation, reactive oxygen

species defense, calcium dynamics, vesicle-mediated transport and endocytosis, rho GTPase

activity, muscle contraction, RNA metabolism and splicing, as well as aerobic respiration and

mitochondrial activity. The number of up-regulated protein networks was considerably larger

despite having fewer significantly altered proteins compared to down-regulated proteins. Up-

regulated pathways clustered into the following groups: Golgi/ER transport, ion conductance,

immune response kinase activity, aerobic respiration and mitochondrial function, axonal and

neuritic morphogenesis, neurotransmitter release, amino acid metabolism, heat shock stress

response, insulin signaling, rho GTPase activity, cell cycle, Wnt and Hedgehog signaling, as well

as gap junctions and plasma membrane.

Multi-target kinase inhibition via GSK2606414 treatment rescues broad proteomic shifts in

tauopathy

Having identified and mapped the pathway changes in Tg mice, we sought to determine

the pathways that were affected by 414 that associated with cognitive rescue. To that end, we

repeated clustering analysis of Reactome pathways (FDR adjusted p-value < 0.1) representing

"Drug Rescued" proteins that were either initially down-regulated (Fig. 5A) or up-regulated (Fig.

5B) in Tg + Veh mice and rescued back toward NTg control levels with 414 treatment. Pathway

groups initially down-regulated in Tg + Veh mice and rescued up toward NTg control levels

included: aerobic respiration, long-term potentiation, muscle contraction, ion conductance, and rho

GTPase activity. Other pathways initially up-regulated in Tg + Veh mice and restored to NTg

control levels included: amino acid metabolism, aerobic respiration and mitochondrial function,

heat shock stress response, Golgi-ER membrane trafficking, axonal and neuritic morphogenesis,

apoptotic protein degradation, cell cycle, gap junction and plasma membrane, immune response

kinase activity, ion conductance, NMDA post-synaptic activity, Wnt and Hedgehog signaling, and

RNA metabolism and translation.

To validate these results, we focused on two distinct cellular processes that were rescued

by 414: RNA translation and mitochondrial redox defense. RNA translation was assessed by a

novel metabolic labeling method developed to investigate the translational dysregulation of

tauopathy *in vivo (34)*. This method involved an intraperitoneal injection of puromycin, a tRNA

analog, shortly before death to rapidly label nascent polypeptide chains *(41)*. Anti-puromycin

antibodies were then used to probe for proteins that incorporated the puromycin as a direct measure

of polypeptide elongation and RNA translation. Anti-puromycin staining of the CA1 region

revealed significantly increased translation activity in Tg + 414 (*p < 0.05) compared to Tg + Veh

mice (Fig. 5C-D). Secondly, since nitroxidative stress is a marker of mitochondrial dysregulation,

and it is implicated in the pathogenesis of AD and other tauopathies *(42, 43)*, we measured changes

in protein nitration between our animal groups. Immunoblot results identified a significant increase

in protein nitration (3-NT) in Tg + Veh animals which was rescued with 414 treatment (Fig. 5E-

F). Together, these results validated the pathway level changes that were rescued in Tg mice by

treatment with 414.

Early stage tauopathy exhibits common protein alterations targetable by multi-target kinase

inhibition

Finally, we investigated how the Tg hippocampal proteome compared with human

tauopathy to provide a context for our findings in AD. We correlated the 276 proteins altered by

"Tg Effect" with two independent proteomic analyses of the human AD brain to identify both

model-specific alterations and novel candidate proteins (Table S3). The majority of significantly

altered mouse proteins mapped to human homologs (>90%) in both human tauopathy datasets, as

expected. First, to account for the disparity in pathology between the tauopathy Tg mouse model

and AD, we assessed the consistency of proteomic alterations between Tg mouse and preclinical

(or asymptomatic) AD and late-stage AD samples (Fig. 6A). Individuals with asymptomatic AD

(Asym AD) included in this analysis exhibited minimal cognitive impairment with moderate to

high amyloid plaque frequency, but with low to no cortical tau pathology (Table S4) *(44)*. Late

AD samples, however, had significant cognitive impairment and high degrees of amyloid and tau

pathology. By identifying the proteins consistent between Tg mice and Asym AD distinct from

late AD, we classified which proteins are considered early markers of AD- and tauopathy-specific

progression, respectively. We found that 15 of 20 human homologs significantly altered in Asym

AD vs healthy control brains were consistently altered in Tg animals (75%; hypergeometric test,

****$p < 0.0001$). Of these 15, nine homolog proteins were rescued by 414 in Tg mice: NUCB1,

SNCA, FKBP1A, PKM, YWHAZ, HSPE1, GRPEL1, NDUFB10, and HOMER1. These nine

proteins are implicated in other neurodegenerative diseases or cellular processes known to be

dysregulated early in tauopathy pathogenesis such as synaptic plasticity, proteostasis, glucose

metabolism, and mitochondrial function. Tg mice matched to a greater extent with the late AD

proteome, where 30 out of 48 significantly altered proteins were inter-specially consistent (≈63%;

hypergeometric test, ****p < 0.0001). Of these 30 proteins, 23 homologs were rescued by 414 in

Tg mice: NUCB1, NTM, ATOX1, SNCA, ENO1, GJA1, FKBP1A, ATP6V1E1, PKM, STMN1,

VSNL1, YWHAZ, HINT1, HPCA, PEPD, ROGDI, PPP1R7, PACSIN1, HSPE1, RTN1,

NDUFB5, DMTN, and HOMER1. These proteins function in pathways rescued by 414 in Tg mice,

namely aerobic respiration/glucose metabolism, mitochondrial function, endocytosis and

trafficking, immune system activation, signaling cascades. Interestingly, several proteins were

found consistent across disease staging: NUCB1, SNCA, FKBP1A, PKM, YWHAZ, HSPE1,

HOMER1. Since each of these were rescued by 414 in Tg mice, they may be important candidate

proteins to consider for future studies.

Considering that the proteome in late-stage neurodegenerative diseases may relate more to

widespread brain organ failure and may not be a specific response to any one disease *(45)*, we

investigated how the Tg proteome matched the protein alterations in AD and another chronic,

degenerative neurodegenerative disease, Parkinson's disease (PD) *(46)*. We applied a similar

approach used with asymptomatic and late stage AD, wherein we analyzed whether proteins found

altered in the Tg hippocampal proteome were consistently altered in AD or PD (Fig. 6B).

Interestingly, Tg animals and PD brains did not share consistently altered homologous proteins,

suggesting a tauopathy specific effect. Of the 39 significantly altered proteins found in AD, 26 had

homologs consistently changed in Tg animals (≈67%; hypergeometric test ****p<0.0001), and 16

were rescued by 414: AKAP5, STMN3, EEF1B2, FKBP1A, ATP6V1E1, NDUFS6, CYCS,

CPLX1, HINT1, PACSIN1, CIAPIN1, RHEB, DDAH1, NDUFB10, ALDH6A1, PIN1. These

proteins function in aerobic respiration/glucose metabolism, RNA translation, apoptosis, immune

system activation, reactive oxygen and nitrogen species defense, membrane trafficking, and

neurotransmitter release.

Four proteins were consistently shared across both late AD datasets with mouse homologs rescued

by 414 in Tg mice (Fig. 6C). These high-confidence, novel candidate proteins are PACSIN1,

FKBP1A, ATP6V1E1, and HINT1. Future studies aimed at investigating the therapeutic impact

of targeting one or all these proteins is currently underway.

Discussion

Tauopathy brains are characterized by aberrant aggregation of a variety of tau conformers,

which include hyperphosphorylated tau species. To reduce tau hyperphosphorylation with the goal

of limiting tau pathogenesis, clinical trials using small molecule kinase inhibitors have

predominantly targeted either individual kinases or whole kinase families with some success *(47–*

50). Nilotinib, a multi-family kinase inhibitor, was used in a recently completed Phase II clinical

trial in which it reduced hippocampal volume loss, cognitive decline, and phosphorylated tau levels

in early to moderate AD patients *(51)*. The compound targets multiple tyrosine kinase families

(52), strongly supporting the therapeutic potential of broad kinase inhibitors in treating

tauopathies. Despite these encouraging outcomes, the precise pathways targeted by Nilotinib and

other kinase inhibitors are unknown. Given the numerous roles of kinases in biological processes,

identification of discrete targets that can be chemically regulated, as well as the mechanisms that

contribute to neuronal dysfunction in tauopathies, will likely improve the chances for success in

the clinic.

Here, we uncover critical biological processes that contribute to neurotoxic processes

driving tauopathies using a multi-family kinase inhibitor, GSK2606414. In doing so, we

demonstrate the potential to mitigate the negative functional consequences of tauopathy without

altering tau hyper-phosphorylation. We treated early stage rTg4510 tau transgenic mice and,

compared to previous studies, we firmly established cognitive and molecular benefits of broad-

based kinase inhibition. With TMT quantitative proteomics, we identified biological processes that

are affected in the tauopathy mice and rescued by the compound. Finally, we cross-reference our

findings to human AD and PD brain proteomic data to narrow our results to high confidence unique

targets for future study. This approach revealed four distinct proteins associated with cognitive

decline in human tauopathy with strong therapeutic potential.

GSK2606414 ameliorates molecular underpinnings of neurodegeneration including

neuronal loss, cognitive dysfunction, and pathological protein accumulation in a variety of disease

models *(11–16, 18)*. These studies established that the benefits of GSK2606414 were conferred by

inhibition of its primary target, PERK, which was upregulated in disease. While PERK is active

in the rTg4510 tau transgenic model, this does not occur until after 6mo *(11, 33)*. By treating

rTg4510 mice from 4-5mo, a time point in which PERK levels not activated, our study identified

numerous PERK-independent pathways targeted by GSK2606414 that elicited rescue from toxic

outcomes in tauopathy. This has important implications for data interpretation of this and previous

studies.

The mechanism by which GSK2606414 confers benefits implicates multiple but

identifiable and targetable pathways. While GSK2606414 is a potent PERK inhibitor, the

concentration needed to reduce PERK activity *in vivo* also inhibits many other significant kinases

(12). In fact, GSK2606414 targets are numerous and have substantial implications in promoting

neurodegeneration *(10, 17, 18)*. While GSK2606414 inhibits many tyrosine kinases shared with

Nilotinib and dozens of others involved in pathways mediating neurodegenerative diseases *(10)*,

our broad approach to also inhibit serine and threonine kinases casts a wider net to identify

phosphorylation-dependent events linked to tau pathogenesis.

We found that measures of early neurodegeneration and cognitive decline in rTg4510 mice

were consistent with other studies. Volumetric T2-weighted MR imaging revealed brain atrophy

in line with other studies in the same transgenic model *(19, 20, 53, 54)*. Neuronal functional deficits

were assessed using MEMRI, a functional imaging approach that uses manganese to approximate

calcium homeostasis *(23)*. Hippocampal decreases in $\Delta R1$, a measurement of manganese retention

over time, supported other studies identifying tau-associated suppression of neuronal calcium

activity in the same model of tauopathy *(55)*. Behavioral phenotypes involving hippocampal

activity were also impaired at this stage in rTg4510 mice, supporting other studies in this model

(20, 56, 57) and MR imaging indications of brain dysfunction and degeneration present in this

study. Importantly, the functional deficits in this model appear to be tau-dependent since transgenic

mice fed doxycycline to suppress tau transgene expression exhibited rescued cognitive and

behavioral activity *(19, 20, 55, 56)*. While transgenicity in rTg4510 mice disrupts the genetic

expression of several genes *(37)*, these studies suggest that the functional deficits in this model are

largely tau dependent.

To evaluate the protein network changes behind this functional rescue, we completed the

first deep proteomic profiling of the brain in rTg4510 mice by quantitatively measuring the

hippocampal proteome. Many of the pathways affected by disease in this early stage of rTg4510

mice overlap with altered pathways in AD brain such as glucose metabolism, immune response,

synaptic activity, and RNA metabolism, highlighting the similarity between this model and human

tauopathy *(6–8, 44)*. The largest alterations in kinase pathway activities were the MAPK cascade,

which is consistent with the involvement of MAPK in human tauopathies *(8)*. Pathways involving

regulation of calcium dynamics were also down-regulated which supports the deficits in calcium-

related activity measured by MEMRI. Pathways involved in muscle contraction were also

dysregulated in the rTg4510 hippocampus, which is consistent with locomotor alterations reported

in this model *(56)* and other mouse models of tauopathy *(58)*. Interestingly, pathways relating to

cell cycle re-entry were altered in rTg4510 mice. These data support the evidence that cell cycle

dysregulation occurs as a result of, and a potential promoter for, tau pathogenesis as seen in human

AD brain and mouse models of tauopathy *(59–61)*. Comparing our proteomic results with a study

of proteostatic dysfunction and protein aggregation in rTg4510 reveals no apparent correlation of

protein abundance between soluble and insoluble fractions *(62)*. These studies suggest that our

proteomic findings reflect alterations in expression and degradation rather than a protein shift

toward insolubility.

Treatment with GSK2606414 broadly affected the hippocampal proteome of rTg4510 mice

and only minimally affected non-transgenic controls. Many pathways critical for neuronal health

altered in human tauopathies were rescued in 414-treated tau transgenic mice. This provides

further support for the multi-target engagement of GSK2606414. Multiple kinase-related pathways

were rescued in transgenic animals treated with GSK2606414, such the MAPK and Wnt signaling

cascades responsible for downstream kinase regulation and tau hyper-phosphorylation *(63)*. These

results were comparable with a recent report on the phospho-proteome changes in AD suggesting

the broad activation of the MAPK cascade in tauopathy *(8)*. We verified the proteomic alterations

found in this study by confirming both translation elongation and mitochondrial redox

accumulation were altered in tau transgenic mice and reduced with 414 treatment (Fig. 5C-F).

Furthermore, mass spectrometry quantification of human P301L tau revealed no significant

alteration of total tau levels with 414 treatment which matched our immunoblot results.

Given the congruent pathway-level results between rTg4510 mice and human tauopathy

studies, we searched for novel candidate targets that were associated with rescued cognitive

function with GSK2606414 treatment. We compared our results with quantitative proteomic cases

of AD as no other deep, high sample number proteomic investigation of another tauopathy has

been reported. To focus on the impact of tau pathology, we assessed the correlation between

cortical proteomic results from cases of asymptomatic AD which exhibit amyloid pathology, but

no tau pathology in the cortex, along with late stage AD cases *(44)*. As expected, more proteins

were consistent between rTg4510 mice and late AD than asymptomatic cases. However, a

considerable number of proteins matched with asymptomatic AD and late stage AD, suggesting

that these targets may relate to an early vulnerability which progressively declines over disease

course. These changes were specific to tauopathy as no protein was consistent with the PD

proteome. Interestingly, alpha-synuclein was down-regulated in transgenic mice and rescued with

GSK2606414 treatment earlier in this model than reported elsewhere *(64, 65)*. Since decreased

alpha-synuclein monomer levels are associated with their pathological oligomerization and

pathological accumulation *(66, 67)*, this early involvement supports a role of for alpha-synuclein

in the pathophysiology of tauopathy. Overlapping the proteins consistent in both AD datasets with

data from transgenic mice revealed four novel, high-confidence candidates that were associated

with cognitive decline and drug-mediated multi-target kinase inhibition: PACSIN1, FKBP1A,

ATP6V1E1, and HINT1.

The neuronal protein PACSIN1 participates in endocytosis *(68–70)* and directly interacts

with tubulin to promote microtubule formation *(71)*. There is evidence for direct tau-PACSIN1

binding, suggesting that this interaction impinges upon tau's capacity to bind microtubules *(72)*.

Network analyses of AD brain identified PACSIN1 as a hub microtubule regulator and a

cerebrospinal fluid marker altered in disease *(6, 44)*. Thus, PACSIN1 represents a protein with a

direct link to tauopathy that could perturb cellular trafficking and tau/microtubule stability.

The peptidyl-prolyl *cis-trans* isomerase (PPIase) FKBP1A promotes protein folding and

trafficking *(73)*. Along with other related PPIases such as FKBP51 *(74–79)*, FKBP1A modifies

tau's structure at the proline-rich region. Perturbation of protein quality control and trafficking

mechanisms is a nearly ubiquitous feature of proteinopathy neurodegenerative diseases, with these

deficiencies often cited as a crucial step in disease cascades that lead to protein mis-folding and

aggregation *(80, 81)*. In addition, FKBP1A activity enhances alpha-synuclein aggregation into

Lewy body inclusions in PD *(82–86)*. FKBP1A associates with tau pathology in neurofibrillary

tangles, suggesting that the protein may play a role in AD pathogenesis *(87)*. Of note, PIN1,

another PPI, was similarly down-regulated in late AD brain and was rescued by GSK2606414 in

Tg animals in this study. FKBP1A has also been implicated in calcium regulation in myocytes and

neuronal signaling *(88, 89)*. Importantly, FKBP1A was the only protein found altered in early AD

and late AD in our analyses, suggesting that it may be a useful target to disrupt disease

pathogenesis. While the full involvement of FKBP1A in proteostasis and calcium activity is

unresolved, there is increasing evidence of its importance in neurodegenerative diseases.

ATP6V1E1 encodes the E1 subunit of the V1 domain of the V-type proton ATPase (V-

ATPase), or ATP6VE1. V-ATPase participates in numerous physiological processes involved in

tauopathy including glucose metabolism, membrane trafficking, protein degradation, and vesicular

acidification *(90)*. Our data indicate that murine ATP6VE1 is reduced in Tg mice and its human

homolog down-regulated in both AD tissue sets (Fig. 6). Thus, ATP6V1E1 function may

deteriorate with disease progression, despite a clear role for the protein in neurodegenerative

disease.

HINT1, or histidine triad nucleotide-binding protein 1, was originally described as a

protein kinase regulator, but has more recently been implicated in regulation of G protein coupled

receptors, or GPCRs, as well as transcription factors that influence Wnt/beta-catenin signaling

(91–94). HINT1 has been implicated in many neuropsychiatric disorders including schizophrenia,

mood disorders, and drug addiction *(95)*. In Schizophrenia, HINT1 is downregulated in both the

dorsolateral prefrontal cortex and prefrontal cortex *(96, 97)*, which mirror its abundance in AD.

While HINT1 is not classically associated with neurodegenerative disease, it is possible that

HINT1 plays a large role in the molecular mechanisms of neuropsychiatric symptoms present in

dementia and in other neurological disorders.

Taken together, our pre-clinical study contributes two major findings. First, we identify

discrete pathways contributing to cognitive changes in tauopathy and highlight four unique

proteins associated with cognitive rescue with strong therapeutic potential. Further work

investigating the novel targets identified in this study could discover better targets amenable to

pharmacological intervention and novel involvement in tauopathy. Second, we demonstrate that

broad, multi-family kinase inhibition can be a useful tool to mitigate the molecular and functional

decline in tauopathy. Since extensively multi-targeted approaches meet safety challenges in the

clinic, the development of compounds that selectively target key kinase cascades at the same time

could greatly benefit combinatorial therapies for the treatment of tauopathy. Studies combining

FDA-approved selective kinase inhibitors may be beneficial toward advancing possible

therapeutic strategies.

Materials and Methods

Animals

All animal studies were approved by the University of Kentucky's Institutional Animal Care and

Use Committee (IACUC) and abided by that committee's policies on animal care and use in

accordance with the Guide for the Care and Use of Laboratory Animals, the Animal Welfare

Regulations Title 9 Code of Federal Regulations Subchapter A, "Animal Welfare," Parts 1–3, and

the Public Health Service Policy on Humane Care and Use of Laboratory Animals. Mice were kept

in standard housing on a 12h light/dark cycle and received food and water *ad libitum*. The tau

transgenic (rTg4510) and parental mice were maintained and genotyped as described previously

(20) and were maintained on mixed FVB and 129S6 backgrounds. Mice of both sexes were used

in experiments unless otherwise stated.

Treatment with GSK2606414

The multi-target inhibitor GSK2606414 (GlaxoSmithKline) was suspended in vehicle (0.5%

hydroxypropylmethyl cellulose + 0.1% Tween-80 in water at pH 4.0) to a dosage of 50 mg/kg, as

previously described *(12)*. Prior to treatment, all mice were handled for 10 minutes twice a day to

acclimate the animals for gavage. Unless otherwise stated, four-month-old rTg4510 and non-

transgenic mice were treated with GSK2606414 or vehicle for 30 days.

A total of 100 mg/kg GSK2606414 was delivered by oral gavage once (100 mg/kg) or twice a day

(50 mg/kg each separated by 10-12 hours). Mice were weighed to ensure precise dosing prior to

each treatment. Daily weight records were carefully monitored due to the potential of irritating the

esophagus and previous reports of GSK2606414-mediated weight loss *(12)*. Animals that lost more

than 20% of starting body weight were excluded from the study (Supplemental Fig. 3). These

animals corresponded to mice in both vehicle and GSK2606414 groups, indicating that weight loss

was not due to the compound.

Tissue Preparation

Mouse brains were harvested as previously described *(33)*. Briefly, mice were irreversibly

anesthetized using isoflurane and transcardially perfused with 0.9% saline. The right brain

hemisphere was drop-fixed in 4% para-formaldehyde for immunohistochemistry. The left

hemisphere was microdissected, snap frozen in liquid nitrogen, and stored at -80°C until

homogenization. For Western blot analysis, brains were weighed, and homogenization buffer

(1:100 Protease Inhibitor, Calbiochem; 1:100 Phosphatase Arrest II, G Biosciences; 1:100

Phosphatase Arrest III G Biosciences; 100μM PMSF; in 1x RIPA, Thermo) was added to a final

concentration of 10% (w/v). Samples were homogenized using a using a small electronic pestle

(VWR). Samples were then centrifuged at 4°C at 13,000xg for 15 minutes. Protein concentration

was established using the bicinchoninic acid (BCA) assay (ThermoScientific).

Protein analysis of brain homogenates

Western blots were performed as previously described *(33)*. Briefly, proteins were separated via

SDS-PAGE, transferred to PVDF membranes, blocked in 5% (w/v) milk-TBS for 1 hour, and then

incubated in primary antibody overnight at 4°C. Primary antibodies diluted in 5% (w/v) milk-TBS

were as follows: PHF1 1:1000 (Peter Davies), total tau 1:1000 (H150, SantaCruz), pY216-GSK3β

1:1000 (Abcam), total GSK3β 1:1000 (Cell Signaling Technologies), 3-NT (1:500, Santa Cruz),

4-HNE 1:1000 (EMD Millipore) and GAPDH and actin for loading controls (1:1000, Cell

Signaling Technologies). Different loading controls were used throughout experiments to ensure

optimal normalization despite non-specific antibody binding and species reactivity. HRP-

conjugated secondary antibodies to mouse or rabbit IgG (Southern Biotech and ThermoScientific)

were used at a 1:1000 dilution where appropriate. Bands were visualized using ECL

(ThermoScientific). Image quantification was performed using ImageJ, normalized to loading

controls.

Open Field Test

Open field-testing was performed in the University of Kentucky Rodent Behavior Core. All testing

was performed at the same time of day for each cohort. Animals were acclimated to testing room

30 minutes prior to testing. Open field-testing was done with the automated Photobeam Activity

System (PAS) with Flexfield Animal Activity System (San Diego Instruments, Inc. San Diego)

coupled to a computer used to eliminate human interaction and bias. Each animal was placed into

the middle of the open field, facing away from the experimenter and exploratory behavior recorded

for 30min. The chamber was cleaned with 10% ethanol between tests to eliminate olfactory cues

from previous animals. The time mice spent in the peripheral areas were recorded in six, 5-min

intervals and is reported by normalizing to time spent in the first 5-min window. Recordings were

taken using EthoVision XT and reviewed by two blinded investigators. Several cohorts were tested

independently over periods of months, but each animal was tested on the 21st day of treatment.

Y-maze

Forced alternation test was performed according to a previously published protocol (30). A

symmetrical Y-maze (Stoelting Co, Wood Dale, IL, USA) with grey Perspex® walls and a grey

steel bottom (arms: 35 cm long, 5 cm wide, 10 cm high) was used for the test. The light in the

testing area was kept at 30±5 lux, the test was performed at room temperature (23± 2°C). Mice

were handled for 5 days (5min/day/mouse) before testing. During the 5 min sample trial (T1), mice

were allowed to explore two arms of the Y-maze, the third arm was blocked. The mice were placed

into the end of the start arm facing the wall, away from the center. After completion of the sample

trial, each mouse was returned to its home cage for 30 min. During the test trial (T2), the block in

arm 3 was removed and the mouse was allowed to explore the entire maze for 5 min. In between

animals and trials, the maze was cleaned with Quatricide® solution to prevent odor cues. Total

duration and number of entries into open arms is reported.

Manganese-enhanced magnetic resonance imaging (MEMRI)

All MRI procedures were performed in accordance with the University of Kentucky MRISC

IACUC Protocol. Brain volume was calculated from T2 images based on manual tracing of the

brain. Manganese chloride (30mM) prepared in saline was delivered to mice via intraperitoneal

injection (66 mg/kg). All MR imaging was performed on a horizontal bore 7T Bruker ClinScan

magnet (Bruker, Ettlingen, Germany) using a cylindrical volume coil for excitation and a cryocoil

for detection. In preparation for imaging, animals are anesthetized using isoflurane (3-5%) in

oxygen at a rate of 0.5-1.0 L/min and maintained using 1-3% isoflurane in oxygen. Body

temperature was maintained using circulating water, and vital signs (core temperature and

respirations) were monitored using a physiological monitoring system SA Instruments Inc. (SAI,

Stony Brook, New York). When applicable, MEMRI was performed after behavioral testing to

reduce stress-related confounds in behavioral test results.

Look-locker imaging was performed following non-selective spin inversion in one slice of the

hippocampus. Fifty images were acquired following inversion with image spacing of 100ms (total

sequence repetition time of 5s) to fully sample the T1-relaxation curve. Additional image

parameters included TR/TE = 5500/1.9, Matrix = 128 x 128, number of averages = 3, field of view

= 17mm x 17mm x 0.7mm. T2-weighted images were acquired covering the entire brain

(excluding the cerebellum and olfactory bulb) using a turbo-spin echo sequence with TR/TE =

3360/42, Slices = 21, Matrix = 448 x 336, number of averages = 2, field of view = 25mm x 25mm

x 0.5mm. The imaging procedures for scanning one mouse were completed in 45min. Imaging was

performed prior to the injection of MnCl$_2$ (baseline) and repeated at 6h post-injection. Image

mapping and analysis was performed in MATLAB (Mathworks, Nattic, MA, USA). Images from

the Look-Locker series were used to reconstruct voxel-by-voxel signal relaxation curves which

were fit to the equation S(t) = So(1-2*exp(-R1*T1)), where S(t) represents the signal at a given

inversion time (TI), So represents the steady state signal at maximal T1, and R1 represents the

longitudinal relaxation rate. Regions of interest (DG, CA1, CA1, CA3, and superior medial cortex)

were identified using the Allen Brain Mouse Atlas. Within each the change in R1 relaxation rates

(ΔR1) before and after MnCl$_2$ exposure was calculated as:

550
$$\Delta R_1 = R1_{6h} - R1_{baseline}$$

Quantitative real-time RT-PCR gene card arrays

Total RNA was extracted from snap frozen hippocampal tissues using TRI reagent solution

(Ambion) following the manufacturer's protocol. RNA was quantified using a BioTech

spectrophotometer and cDNA was generated using SuperScript IV reverse transcriptase

(Invitrogen). Real-time RT-PCR was performed using a TaqMan custom Array (Life

Technologies) according to manufacturer's instructions. Gene card array was designed to evaluate

changes in the expression of genes delineated in Figure 4. Gene expression was evaluated by

normalizing to GAPDH and 18S expression as internal controls. The real-time values for each

sample were averaged and evaluated using the comparative CT (cycle time) method.

Protein Extraction and Quantification for Proteomics

Microdissected hippocampal mouse tissue were processed for protein extraction which were

prepared according to *(98)* with the following modifications. Briefly, samples were ground in

liquid nitrogen into fine powder and incubated in extraction buffer (0.1 M Tris-HCl pH 8.8, 10

mM EDTA, 0.2 M DTT, 0.9 M sucrose), continued by grinding in a fume hood, and then the

extract was agitated for 2 hrs at room temp. After washing twice with 0.1 M ammonium acetate in

methanol and twice with 80% acetone, the dried pellet was dissolved with 50mM ammonium

bicarbonate buffer. The mixtures were cooled on ice before centrifugation (4°C) for 20 min at

12,000 g. The soluble proteins were performed using an EZQ Protein Quantitation Kit (Invitrogen,

Carlsbad, CA, USA) with the SoftMax Pro Software v5.3 (Molecular Devices, Downingtown, PA,

USA).

Protein Extraction, Digestion, TMT Labeling, and LC-MS/MS

Proteins were dissolved in protein buffer (8M Urea, 50 mM Tris-HCl (pH 8.0), 50 mM

triethylammonium bicarbonate, 0.1% SDS (w/v), 1% Triton-100 (w/v), 1 mM PMSF, 10 μg/ml

leupeptin, 1% phosphatase inhibitor cocktail 2 and 3 (v/v)). For each sample, a total of 50 μg of

protein were reduced, alkylated, trypsin-digested, and labeled according to the manufacturer's

instructions (Thermo Scientific). For each mass-spectrometer batch, late-stage 8mo transgenic

tissue were labeled with TMT tags 126 and 127N, non-transgenic mice treated with 414 were

labeled with TMT tags 128C and 129N, non-transgenic mice treated with vehicle were labeled

with TMT tags 128C and 129N, transgenic mice treated with 414 were labeled with 129C and

130N, transgenic mice treated with vehicle were labeled with 130C and 131N, and a mixed-sample

internal batch was labeled with 131C. Labeled peptides were desalted with C18-solid phase

extraction and dissolved in strong cation exchange (SCX) solvent A (25% (v/v) acetonitrile, 10

mM ammonium formate, and 0.1% (v/v) formic acid, pH 2.8). The peptides were fractionated

using an Agilent HPLC 1260 with a polysulfoethyl A column (2.1 × 100 mm, 5 μm, 300 Å;

PolyLC, Columbia, MD, USA). Peptides were eluted with a linear gradient of 0–20% solvent B

(25% (v/v) acetonitrile and 500 mM ammonium formate, pH 6.8) over 50 min followed by ramping

up to 100% solvent B in 5 min. The absorbance at 280 nm was monitored and a total of 19 fractions

were collected. The fractions were lyophilized, desalted, and resuspended in LC solvent A (0.1%

formic acid in 97% water (v/v), 3% acetonitrile (v/v)). A hybrid quadrupole Orbitrap (Q Exactive)

MS system (Thermo Fisher Scientific, Bremen, Germany) was used with high energy collision

dissociation (HCD) in each MS and MS/MS cycle. The MS system was interfaced with an

automated Easy-nLC 1000 system (Thermo Fisher Scientific, Bremen, Germany). Each sample

fraction was loaded onto an Acclaim Pepmap 100 pre-column (20 mm × 75 μm; 3 μm-C18) and

separated on a PepMap RSLC analytical column (250 mm × 75 μm; 2 μm-C18) at a flow rate at

350 nl/min during a linear gradient from solvent A (0.1% formic acid (v/v)) to 30% solvent B

(0.1% formic acid (v/v) and 99.9% acetonitrile (v/v)) for 95 min, to 98% solvent B for 15 min, and

hold 98% solvent B for additional 30 min. Full MS scans were acquired in the Orbitrap mass

analyzer over m/z 400–2000 range with resolution 70,000 at 200 m/z. The top ten most intense

peaks with charge state ≥ 3 were fragmented in the HCD collision cell normalized collision energy

of 28%, (the isolation window was 2 m/z). The maximum ion injection times for the survey scan

and the MS/MS scans were 250 ms, respectively and the ion target values were set to 3e6 and 1e6,

respectively. Selected sequenced ions were dynamically excluded for 60 sec.

Raw data files obtained from the Orbitrap Fusion were processed using Proteome Discoverer™

(version 2.3). MS/MS spectra were searched against the UniProt mouse proteome database

downloaded Dec 4, 2019 with 63369 total sequences including the sequence for 0N4R P301L tau

mutant cDNA. The SEQUEST HT search engine was used and parameters were the following:

fully-tryptic specificity; maximum of two missed cleavages; minimum peptide length of 6; fixed

modifications for TMT tags on lysine residues and peptide N-termini (+ 229.162932 Da) and

carbamidomethylation of cysteine residues (+ 57.02146 Da); variable modifications for oxidation

of methionine residues (+ 15.99492 Da); phosphorylation of serine, threonine and tyrosine (+

79.9663 Da); precursor mass tolerance of 20 ppm; fragment mass tolerance of 0.05 Daltons.

Percolator was used to filter peptide spectral matches (PSM) and peptides to a false discovery rate

(FDR) of less than 1% using target-decoy strategy. Following spectral assignment, peptides were

assembled into proteins and were further filtered based on the combined probabilities of their

constituent peptides to a final FDR of 1%. In cases of redundancy, shared peptides were assigned

to the protein sequence in adherence with the principles of parsimony. Reporter ions were

quantified from MS2 scans using an integration tolerance of 20 ppm with the most confident

centroid setting.

Each MS batch protein abundance value was normalized to the internal standard (131C)

abundance, enabling batch-corrected analysis across groups. The criteria for protein inclusion in

statistical analysis were that each protein must i) be present in all samples, ii) have a global FDR

< 0.05, iii) contain at least one unique peptide, and iv) be an annotated master protein. These

filtering criteria reduced the total amount of unique protein hits to 24,149 peptides and 1099

proteins. Normalized abundance ratios of selected proteins were loaded into R (3.6.2) for

processing. Significant differential expression of proteins between samples was determined as $p <$

0.05 using unpaired Welch's t-test with FDR multiple-comparisons correction. No fold change

(FC) cutoff was used as TMT-labeled tagging and subsequent analysis is subject to ratio

compression *(99)*. Reactome pathway *(39)* changes were assessed using gProfiler *(38)* were used

against total mouse protein coding background. Significant Reactome pathways (FDR adj. p value

< 0.10) were loaded into Cytoscape for clustering analysis *(40)*.

Immunofluorescence

Immunofluorescence (IF) was performed as previously described *(33, 100)*. Primary antibodies:

pPERK (1:250, SantaCruz) and NRF2 (1:250; Abcam).

Statistical Analysis

Statistical analyses for all data apart from proteomics were performed using GraphPad Prism 8

(Graph Pad Software, Inc. La Jolla, CA, USA). Results are shown as the mean ± standard error or

standard deviation as described in each figure. Single-variate data were analyzed using unpaired

Student's t-test. Multi-variate data were analyzed with one-way or two-way ANOVA where

appropriate, corrected for multiple comparisons with Tukey post-test analysis unless otherwise

stated. A value of $p < 0.05$ was considered statistically significant except for Reactome pathway

analyses which used an FDR-adjusted p value cutoff of 0.10.

References

1. S. Jadhav, J. Avila, M. Schöll, G. G. Kovacs, E. Kövari, R. Skrabana, L. D. Evans, E. Kontsekova,
B. Malawska, R. de Silva, L. Buee, N. Zilka, A walk through tau therapeutic strategies, *Acta*
Neuropathologica Communications **7**, 22 (2019).

2. M. R. Khanna, J. Kovalevich, V. M.-Y. Lee, J. Q. Trojanowski, K. R. Brunden, Therapeutic
Strategies for the Treatment of Tauopathies: Hopes and Challenges, *Alzheimers Dement* **12**,
1051–1065 (2016).

3. L.-K. Huang, S.-P. Chao, C.-J. Hu, Clinical trials of new drugs for Alzheimer disease, *Journal of*
Biomedical Science **27**, 18 (2020).

4. A. L. Boxer, A. E. Lang, M. Grossman, D. S. Knopman, B. L. Miller, L. S. Schneider, R. S. Doody,
A. Lees, L. I. Golbe, D. R. Williams, J.-C. Corvol, A. Ludolph, D. Burn, S. Lorenzl, I. Litvan, E. D.
Roberson, G. U. Höglinger, M. Koestler, C. R. Jack, V. Van Deerlin, C. Randolph, I. V. Lobach, H.
W. Heuer, I. Gozes, L. Parker, S. Whitaker, J. Hirman, A. J. Stewart, M. Gold, B. H. Morimoto, AL-
108-231 Investigators, Davunetide in patients with progressive supranuclear palsy: a
randomised, double-blind, placebo-controlled phase 2/3 trial, *Lancet Neurol* **13**, 676–685
(2014).

5. R. R. Ramsay, M. R. Popovic-Nikolic, K. Nikolic, E. Uliassi, M. L. Bolognesi, A perspective on
multi-target drug discovery and design for complex diseases, *Clinical and Translational Medicine*
7, 3 (2018).

6. E. C. B. Johnson, E. B. Dammer, D. M. Duong, L. Ping, M. Zhou, L. Yin, L. A. Higginbotham, A.
Guajardo, B. White, J. C. Troncoso, M. Thambisetty, T. J. Montine, E. B. Lee, J. Q. Trojanowski, T.
G. Beach, E. M. Reiman, V. Haroutunian, M. Wang, E. Schadt, B. Zhang, D. W. Dickson, N.
Ertekin-Taner, T. E. Golde, V. A. Petyuk, P. L. De Jager, D. A. Bennett, T. S. Wingo, S. Rangaraju, I.
Hajjar, J. M. Shulman, J. J. Lah, A. I. Levey, N. T. Seyfried, Large-scale proteomic analysis of
Alzheimer's disease brain and cerebrospinal fluid reveals early changes in energy metabolism
associated with microglia and astrocyte activation, *Nat. Med.* **26**, 769–780 (2020).

7. E. C. B. Johnson, E. B. Dammer, D. M. Duong, L. Yin, M. Thambisetty, J. C. Troncoso, J. J. Lah,
A. I. Levey, N. T. Seyfried, Deep proteomic network analysis of Alzheimer's disease brain reveals
alterations in RNA binding proteins and RNA splicing associated with disease, *Mol*
Neurodegener **13**, 52 (2018).

8. B. Bai, X. Wang, Y. Li, P.-C. Chen, K. Yu, K. K. Dey, J. M. Yarbro, X. Han, B. M. Lutz, S. Rao, Y.
Jiao, J. M. Sifford, J. Han, M. Wang, H. Tan, T. I. Shaw, J.-H. Cho, S. Zhou, H. Wang, M. Niu, A.
Mancieri, K. A. Messler, X. Sun, Z. Wu, V. Pagala, A. A. High, W. Bi, H. Zhang, H. Chi, V.
Haroutunian, B. Zhang, T. G. Beach, G. Yu, J. Peng, Deep Multilayer Brain Proteomics Identifies
Molecular Networks in Alzheimer's Disease Progression, *Neuron* **105**, 975-991.e7 (2020).

9. M. Perluigi, E. Barone, F. Di Domenico, D. A. Butterfield, Aberrant protein phosphorylation in
Alzheimer disease brain disturbs pro-survival and cell death pathways, *Biochim. Biophys. Acta*
1862, 1871–1882 (2016).

10. J. M. Axten, J. R. Medina, Y. Feng, A. Shu, S. P. Romeril, S. W. Grant, W. H. H. Li, D. A.
Heerding, E. Minthorn, T. Mencken, C. Atkins, Q. Liu, S. Rabindran, R. Kumar, X. Hong, A. Goetz,
T. Stanley, J. D. Taylor, S. D. Sigethy, G. H. Tomberlin, A. M. Hassell, K. M. Kahler, L. M.
Shewchuk, R. T. Gampe, Discovery of 7-methyl-5-(1-{[3-(trifluoromethyl)phenyl]acetyl}-2,3-
dihydro-1H-indol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine (GSK2606414), a potent and
selective first-in-class inhibitor of protein kinase R (PKR)-like endoplasmic reticulum kinase
(PERK), *J. Med. Chem.* **55**, 7193–7207 (2012).

11. H. Radford, J. A. Moreno, N. Verity, M. Halliday, G. R. Mallucci, PERK inhibition prevents tau-
mediated neurodegeneration in a mouse model of frontotemporal dementia, *Acta Neuropathol*
130, 633–642 (2015).

12. J. A. Moreno, M. Halliday, C. Molloy, H. Radford, N. Verity, J. M. Axten, C. A. Ortori, A. E.
Willis, P. M. Fischer, D. A. Barrett, G. R. Mallucci, Oral treatment targeting the unfolded protein
response prevents neurodegeneration and clinical disease in prion-infected mice, *Sci Transl*
Med **5**, 206ra138 (2013).

13. C. Meng, J. Zhang, B. Dang, H. Li, H. Shen, X. Li, Z. Wang, PERK Pathway Activation Promotes
Intracerebral Hemorrhage Induced Secondary Brain Injury by Inducing Neuronal Apoptosis Both
in Vivo and in Vitro, *Front Neurosci* **12** (2018), doi:10.3389/fnins.2018.00111.

14. T. Vanderweyde, D. J. Apicco, K. Youmans-Kidder, P. E. A. Ash, C. Cook, E. L. da Rocha, K.
Jansen-West, A. A. Frame, A. Citro, J. D. Leszyk, P. Ivanov, J. F. Abisambra, M. Steffen, H. Li, L.
Petrucelli, B. Wolozin, Interaction of tau with the RNA-Binding Protein TIA1 Regulates tau
Pathophysiology and Toxicity, *Cell Rep* **15**, 1455–1466 (2016).

15. V. Sharma, H. Ounallah-Saad, D. Chakraborty, M. Hleihil, R. Sood, I. Barrera, E. Edry, S. K.
Chandran, S. B. T. de Leon, H. Kaphzan, K. Rosenblum, Local Inhibition of PERK Enhances
Memory and Reverses Age-Related Deterioration of Cognitive and Neuronal Properties, *J.*
Neurosci. **38**, 648–658 (2018).

16. H. L. Smith, O. J. Freeman, A. J. Butcher, S. Holmqvist, I. Humoud, T. Schätzl, D. T. Hughes, N.
C. Verity, D. P. Swinden, J. Hayes, L. de Weerd, D. H. Rowitch, R. J. M. Franklin, G. R. Mallucci,
Astrocyte Unfolded Protein Response Induces a Specific Reactivity State that Causes Non-Cell-
Autonomous Neuronal Degeneration, *Neuron* **105**, 855-866.e5 (2020).

17. D. Rojas-Rivera, T. Delvaeye, R. Roelandt, W. Nerinckx, K. Augustyns, P. Vandenabeele, M. J.
M. Bertrand, When PERK inhibitors turn out to be new potent RIPK1 inhibitors: critical issues on
the specificity and use of GSK2606414 and GSK2656157, *Cell Death Differ* **24**, 1100–1110
(2017).

18. M. Mahameed, T. Wilhelm, O. Darawshi, A. Obiedat, W.-S. Tommy, C. Chintha, T. Schubert,
A. Samali, E. Chevet, L. A. Eriksson, M. Huber, B. Tirosh, The unfolded protein response
modulators GSK2606414 and KIRA6 are potent KIT inhibitors, *Cell Death Dis* **10**, 300 (2019).

19. Z. Berger, H. Roder, A. Hanna, A. Carlson, V. Rangachari, M. Yue, Z. Wszolek, K. Ashe, J.
Knight, D. Dickson, C. Andorfer, T. L. Rosenberry, J. Lewis, M. Hutton, C. Janus, Accumulation of
Pathological Tau Species and Memory Loss in a Conditional Model of Tauopathy, *J. Neurosci.* **27**,
3650–3662 (2007).

20. K. SantaCruz, J. Lewis, T. Spires, J. Paulson, L. Kotilinek, M. Ingelsson, A. Guimaraes, M.
DeTure, M. Ramsden, E. McGowan, C. Forster, M. Yue, J. Orne, C. Janus, A. Mariash, M.
Kuskowski, B. Hyman, M. Hutton, K. H. Ashe, Tau Suppression in a Neurodegenerative Mouse
Model Improves Memory Function, *Science* **309**, 476–481 (2005).

21. R. M. Bailey, J. Howard, J. Knight, N. Sahara, D. W. Dickson, J. Lewis, Effects of the C57BL/6
strain background on tauopathy progression in the rTg4510 mouse model, *Molecular*
Neurodegeneration **9**, 8 (2014).

22. S. N. Fontaine, A. Ingram, R. A. Cloyd, S. E. Meier, E. Miller, D. Lyons, G. K. Nation, E.
Mechas, B. Weiss, C. Lanzillotta, F. Di Domenico, F. Schmitt, D. K. Powell, M. Vandsburger, J. F.
Abisambra, Identification of changes in neuronal function as a consequence of aging and
tauopathic neurodegeneration using a novel and sensitive magnetic resonance imaging
approach, *Neurobiol Aging* **56**, 78–86 (2017).

23. R. A. Cloyd, S. A. Koren, J. F. Abisambra, Manganese-Enhanced Magnetic Resonance
Imaging: Overview and Central Nervous System Applications With a Focus on
Neurodegeneration, *Front Aging Neurosci* **10** (2018), doi:10.3389/fnagi.2018.00403.

24. B. T. Bedenk, S. Almeida-Corrêa, A. Jurik, N. Dedic, B. Grünecker, A. J. Genewsky, S. F.
Kaltwasser, C. J. Riebe, J. M. Deussing, M. Czisch, C. T. Wotjak, Mn2+ dynamics in manganese-
enhanced MRI (MEMRI): Cav1.2 channel-mediated uptake and preferential accumulation in
projection terminals, *NeuroImage* **169**, 374–382 (2018).

25. H. Lu, Z.-X. Xi, L. Gitajn, W. Rea, Y. Yang, E. A. Stein, Cocaine-induced brain activation
detected by dynamic manganese-enhanced magnetic resonance imaging (MEMRI), *Proc. Natl.*
Acad. Sci. U.S.A. **104**, 2489–2494 (2007).

26. H. Masumiya, H. Tsujikawa, N. Hino, R. Ochi, Modulation of manganese currents by 1, 4-
dihydropyridines, isoproterenol and forskolin in rabbit ventricular cells, *Pflugers Arch.* **446**, 695–
701 (2003).

27. K. Itoh, M. Sakata, M. Watanabe, Y. Aikawa, H. Fujii, The entry of manganese ions into the
brain is accelerated by the activation of N-methyl-d-aspartate receptors, *Neuroscience* **154**,
732–740 (2008).

28. M. H. Vandsburger, B. A. French, C. M. Kramer, X. Zhong, F. H. Epstein, Displacement-
encoded and manganese-enhanced cardiac MRI reveal that nNOS, not eNOS, plays a dominant
role in modulating contraction and calcium influx in the mammalian heart, *Am. J. Physiol. Heart*
Circ. Physiol. **302**, H412-419 (2012).

29. P. F. Antkowiak, M. H. Vandsburger, F. H. Epstein, Quantitative pancreatic β cell MRI using
manganese-enhanced Look-Locker imaging and two-site water exchange analysis, *Magn Reson*
Med **67**, 1730–1739 (2012).

30. F. M. LaFerla, Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease,
Nat. Rev. Neurosci. **3**, 862–872 (2002).

31. C. Hooper, R. Killick, S. Lovestone, The GSK3 hypothesis of Alzheimer's disease, *J Neurochem*
104, 1433–1439 (2008).

32. M. C. Bell, S. E. Meier, A. L. Ingram, J. F. Abisambra, PERK-opathies: An Endoplasmic
Reticulum Stress Mechanism Underlying Neurodegeneration, *Curr Alzheimer Res* **13**, 150–163
(2016).

33. J. F. Abisambra, U. K. Jinwal, L. J. Blair, J. C. O'Leary, Q. Li, S. Brady, L. Wang, C. E. Guidi, B.
Zhang, B. A. Nordhues, M. Cockman, A. Suntharalingham, P. Li, Y. Jin, C. A. Atkins, C. A. Dickey,
Tau accumulation activates the unfolded protein response by impairing endoplasmic reticulum-
associated degradation, *J. Neurosci.* **33**, 9498–9507 (2013).

34. S. A. Koren, M. J. Hamm, S. E. Meier, B. E. Weiss, G. K. Nation, E. A. Chishti, J. P. Arango, J.
Chen, H. Zhu, E. M. Blalock, J. F. Abisambra, Tau drives translational selectivity by interacting
with ribosomal proteins, *Acta Neuropathol* **137**, 571–583 (2019).

35. S. B. Cullinan, D. Zhang, M. Hannink, E. Arvisais, R. J. Kaufman, J. A. Diehl, Nrf2 is a direct
PERK substrate and effector of PERK-dependent cell survival, *Mol. Cell. Biol.* **23**, 7198–7209
(2003).

36. L. O. Goodwin, E. Splinter, T. L. Davis, R. Urban, H. He, R. E. Braun, E. J. Chesler, V. Kumar, M.
van Min, J. Ndukum, V. M. Philip, L. G. Reinholdt, K. Svenson, J. K. White, M. Sasner, C. Lutz, S.
A. Murray, Large-scale discovery of mouse transgenic integration sites reveals frequent
structural variation and insertional mutagenesis, *Genome Res.* **29**, 494–505 (2019).

37. J. Gamache, K. Benzow, C. Forster, L. Kemper, C. Hlynialuk, E. Furrow, K. H. Ashe, M. D.
Koob, Factors other than hTau overexpression that contribute to tauopathy-like phenotype in
rTg4510 mice, *Nat Commun* **10**, 2479 (2019).

38. U. Raudvere, L. Kolberg, I. Kuzmin, T. Arak, P. Adler, H. Peterson, J. Vilo, g:Profiler: a web
server for functional enrichment analysis and conversions of gene lists (2019 update), *Nucleic
Acids Res* **47**, W191–W198 (2019).

39. B. Jassal, L. Matthews, G. Viteri, C. Gong, P. Lorente, A. Fabregat, K. Sidiropoulos, J. Cook, M.
Gillespie, R. Haw, F. Loney, B. May, M. Milacic, K. Rothfels, C. Sevilla, V. Shamovsky, S. Shorser,
T. Varusai, J. Weiser, G. Wu, L. Stein, H. Hermjakob, P. D'Eustachio, The reactome pathway
knowledgebase, *Nucleic Acids Res.* **48**, D498–D503 (2020).

40. P. Shannon, A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.
Schwikowski, T. Ideker, Cytoscape: a software environment for integrated models of
biomolecular interaction networks, *Genome Res.* **13**, 2498–2504 (2003).

41. C. A. Goodman, T. A. Hornberger, Measuring protein synthesis with SUnSET: a valid
alternative to traditional techniques?, *Exerc Sport Sci Rev* **41**, 107–115 (2013).

42. D. A. Butterfield, T. T. Reed, M. Perluigi, C. De Marco, R. Coccia, J. N. Keller, W. R.
Markesbery, R. Sultana, Elevated Levels of 3-Nitrotyrosine in Brain From Subjects with Amnestic
Mild Cognitive Impairment: Implications for the Role of Nitration in the Progression of
Alzheimer's Disease, *Brain Res* **1148**, 243–248 (2007).

43. D. A. Butterfield, T. Reed, R. Sultana, Roles of 3-nitrotyrosine- and 4-hydroxynonenal-
modified brain proteins in the progression and pathogenesis of Alzheimer's disease, *Free Radic.
Res.* **45**, 59–72 (2011).

44. N. T. Seyfried, E. B. Dammer, V. Swarup, D. Nandakumar, D. M. Duong, L. Yin, Q. Deng, T.
Nguyen, C. M. Hales, T. Wingo, J. Glass, M. Gearing, M. Thambisetty, J. C. Troncoso, D. H.
Geschwind, J. J. Lah, A. I. Levey, A Multi-network Approach Identifies Protein-Specific Co-
expression in Asymptomatic and Symptomatic Alzheimer's Disease, *Cell Syst* **4**, 60-72.e4 (2017).

45. T. E. Golde, S. T. DeKosky, D. Galasko, Alzheimer's disease: The right drug, the right time,
Science **362**, 1250–1251 (2018).

46. L. Ping, D. M. Duong, L. Yin, M. Gearing, J. J. Lah, A. I. Levey, N. T. Seyfried, Global
quantitative analysis of the human brain proteome in Alzheimer's and Parkinson's Disease, *Sci*
Data **5**, 180036 (2018).

47. C. H. van Dyck, H. B. Nygaard, K. Chen, M. C. Donohue, R. Raman, R. A. Rissman, J. B.
Brewer, R. A. Koeppe, T. W. Chow, M. S. Rafii, D. Gessert, J. Choi, R. S. Turner, J. A. Kaye, S. A.
Gale, E. M. Reiman, P. S. Aisen, S. M. Strittmatter, Effect of AZD0530 on Cerebral Metabolic
Decline in Alzheimer Disease: A Randomized Clinical Trial, *JAMA Neurol* (2019),
doi:10.1001/jamaneurol.2019.2050.

48. S. Lovestone, M. Boada, B. Dubois, M. Hüll, J. O. Rinne, H.-J. Huppertz, M. Calero, M. V.
Andrés, B. Gómez-Carrillo, T. León, T. del Ser, ARGO investigators, A phase II trial of tideglusib in
Alzheimer's disease, *J. Alzheimers Dis.* **45**, 75–88 (2015).

49. E. Tolosa, I. Litvan, G. U. Höglinger, D. Burn, A. Lees, M. V. Andrés, B. Gómez-Carrillo, T.
León, T. Del Ser, TAUROS Investigators, A phase 2 trial of the GSK-3 inhibitor tideglusib in
progressive supranuclear palsy, *Mov. Disord.* **29**, 470–478 (2014).

50. M. Medina, An Overview on the Clinical Development of Tau-Based Therapeutics, *Int J Mol*
Sci **19** (2018), doi:10.3390/ijms19041160.

51. R. S. Turner, M. L. Hebron, A. Lawler, E. E. Mundel, N. Yusuf, J. N. Starr, M. Anjum, F. Pagan,
Y. Torres-Yaghi, W. Shi, S. Mulki, D. Ferrante, S. Matar, X. Liu, G. Esposito, F. Berkowitz, X. Jiang,
J. Ahn, C. Moussa, Nilotinib Effects on Safety, Tolerability, and Biomarkers in Alzheimer's
Disease, *Annals of Neurology* **88**, 183–194 (2020).

52. O. Hantschel, U. Rix, G. Superti-Furga, Target spectrum of the BCR-ABL inhibitors imatinib,
nilotinib and dasatinib, *Leuk. Lymphoma* **49**, 615–619 (2008).

53. T. L. Spires, J. D. Orne, K. SantaCruz, R. Pitstick, G. A. Carlson, K. H. Ashe, B. T. Hyman,
Region-specific Dissociation of Neuronal Loss and Neurofibrillary Pathology in a Mouse Model
of Tauopathy, *Am J Pathol* **168**, 1598–1607 (2006).

54. H. E. Holmes, N. Colgan, O. Ismail, D. Ma, N. M. Powell, J. M. O'Callaghan, I. F. Harrison, R. A.
Johnson, T. K. Murray, Z. Ahmed, M. Heggenes, A. Fisher, M. J. Cardoso, M. Modat, S. Walker-
Samuel, E. M. C. Fisher, S. Ourselin, M. J. O'Neill, J. A. Wells, E. C. Collins, M. F. Lythgoe, Imaging
the accumulation and suppression of tau pathology using multiparametric MRI, *Neurobiol Aging*
39, 184–194 (2016).

55. M. A. Busche, S. Wegmann, S. Dujardin, C. Commins, J. Schiantarelli, N. Klickstein, T. V.
Kamath, G. A. Carlson, I. Nelken, B. T. Hyman, Tau impairs neural circuits, dominating amyloid-β
effects, in Alzheimer models in vivo, *Nat Neurosci* **22**, 57–64 (2019).

56. T. Blackmore, S. Meftah, T. K. Murray, P. J. Craig, A. Blockeel, K. Phillips, B. Eastwood, M. J.
O'Neill, H. Marston, Z. Ahmed, G. Gilmour, F. Gastambide, Tracking progressive pathological
and functional decline in the rTg4510 mouse model of tauopathy, *Alzheimer's Research &*
Therapy **9**, 77 (2017).

57. C. Cook, J. H. Dunmore, M. E. Murray, K. Scheffel, N. Shukoor, J. Tong, M. Castanedes-Casey,
V. Phillips, L. Rousseau, M. S. Penuliar, A. Kurti, D. W. Dickson, L. Petrucelli, J. D. Fryer, Severe
amygdala dysfunction in a MAPT transgenic mouse model of frontotemporal dementia,
Neurobiol. Aging **35**, 1769–1777 (2014).

58. J. Lewis, E. McGowan, J. Rockwood, H. Melrose, P. Nacharaju, M. Van Slegtenhorst, K.
Gwinn-Hardy, M. Paul Murphy, M. Baker, X. Yu, K. Duff, J. Hardy, A. Corral, W. L. Lin, S. H. Yen,
D. W. Dickson, P. Davies, M. Hutton, Neurofibrillary tangles, amyotrophy and progressive motor
disturbance in mice expressing mutant (P301L) tau protein, *Nat. Genet.* **25**, 402–405 (2000).

59. H. Lee, G. Casadesus, X. Zhu, R. J. Castellani, A. McShea, G. Perry, R. B. Petersen, V. Bajic, M.
A. Smith, Cell Cycle Re-entry Mediated Neurodegeneration and Its Treatment Role in the
Pathogenesis of Alzheimer's Disease, *Neurochem Int* **54**, 84–88 (2009).

60. A. C. Hradek, H.-P. Lee, S. L. Siedlak, S. L. Torres, W. Jung, A. H. Han, H. Lee, Distinct
chronology of neuronal cell cycle re-entry and tau pathology in the 3×Tg-AD mouse model and
Alzheimer disease patients, *J Alzheimers Dis* **43**, 57–65 (2015).

61. C. Andorfer, C. M. Acker, Y. Kress, P. R. Hof, K. Duff, P. Davies, Cell-cycle reentry and cell
death in transgenic mice expressing nonmutant human tau isoforms, *J. Neurosci.* **25**, 5446–
5454 (2005).

62. M. C. Pace, G. Xu, S. Fromholt, J. Howard, K. Crosby, B. I. Giasson, J. Lewis, D. R. Borchelt,
Changes in proteome solubility indicate widespread proteostatic disruption in mouse models of
neurodegenerative disease, *Acta Neuropathol* **136**, 919–938 (2018).

63. W. Wan, S. Xia, B. Kalionis, L. Liu, Y. Li, The Role of Wnt Signaling in the Development of
Alzheimer's Disease: A Potential Therapeutic Target?, *Biomed Res Int* **2014** (2014),
doi:10.1155/2014/301575.

64. Y. Takaichi, Y. Ano, J. K. Chambers, K. Uchida, A. Takashima, H. Nakayama, Deposition of
Phosphorylated α-Synuclein in the rTg4510 Mouse Model of Tauopathy, *J. Neuropathol. Exp.*
Neurol. **77**, 920–928 (2018).

65. Y. Takaichi, J. K. Chambers, H. Inoue, Y. Ano, A. Takashima, H. Nakayama, K. Uchida,
Phosphorylation and oligomerization of α-synuclein associated with GSK-3β activation in the
rTg4510 mouse model of tauopathy, *Acta Neuropathol Commun* **8**, 86 (2020).

66. D. Twohig, H. M. Nielsen, α-synuclein in the pathophysiology of Alzheimer's disease, *Mol*
Neurodegener **14**, 23 (2019).

67. M. E. Larson, M. A. Sherman, S. Greimel, M. Kuskowski, J. A. Schneider, D. A. Bennett, S. E.
Lesné, Soluble α-synuclein is a novel modulator of Alzheimer's disease pathophysiology, *J.*
Neurosci. **32**, 10253–10266 (2012).

68. J. Modregger, B. Ritter, B. Witter, M. Paulsson, M. Plomann, All three PACSIN isoforms bind
to endocytic proteins and inhibit endocytosis, *J. Cell. Sci.* **113 Pt 24**, 4511–4521 (2000).

69. V. Anggono, K. J. Smillie, M. E. Graham, V. A. Valova, M. A. Cousin, P. J. Robinson, Syndapin I
is the phosphorylation-regulated dynamin I partner in synaptic vesicle endocytosis, *Nat.*
Neurosci. **9**, 752–760 (2006).

70. A. Braun, R. Pinyol, R. Dahlhaus, D. Koch, P. Fonarev, B. D. Grant, M. M. Kessels, B.
Qualmann, EHD proteins associate with syndapin I and II and such interactions play a crucial
role in endosomal recycling, *Mol. Biol. Cell* **16**, 3642–3658 (2005).

71. E.-M. S. Grimm-Günter, M. Milbrandt, B. Merkl, M. Paulsson, M. Plomann, PACSIN proteins
bind tubulin and promote microtubule assembly, *Exp. Cell Res.* **314**, 1991–2003 (2008).

72. Y. Liu, K. Lv, Z. Li, A. C. H. Yu, J. Chen, J. Teng, PACSIN1, a Tau-interacting protein, regulates
axonal elongation and branching by facilitating microtubule instability, *J. Biol. Chem.* **287**,
39911–39924 (2012).

73. F. Edlich, G. Fischer, Pharmacological targeting of catalyzed protein folding: the example of
peptide bond cis/trans isomerases, *Handb Exp Pharmacol* , 359–404 (2006).

74. L. J. Blair, B. A. Nordhues, S. E. Hill, K. M. Scaglione, J. C. O'Leary, S. N. Fontaine, L. Breydo, B.
Zhang, P. Li, L. Wang, C. Cotman, H. L. Paulson, M. Muschol, V. N. Uversky, T. Klengel, E. B.
Binder, R. Kayed, T. E. Golde, N. Berchtold, C. A. Dickey, Accelerated neurodegeneration
through chaperone-mediated oligomerization of tau, *J. Clin. Invest.* **123**, 4158–4169 (2013).

75. L. J. Blair, J. D. Baker, J. J. Sabbagh, C. A. Dickey, The emerging role of peptidyl-prolyl
isomerase chaperones in tau oligomerization, amyloid processing, and Alzheimer's disease, *J.*
Neurochem. **133**, 1–13 (2015).

76. U. K. Jinwal, J. Koren, S. I. Borysov, A. B. Schmid, J. F. Abisambra, L. J. Blair, A. G. Johnson, J.
R. Jones, C. L. Shults, J. C. O'Leary, Y. Jin, J. Buchner, M. B. Cox, C. A. Dickey, The Hsp90
cochaperone, FKBP51, increases Tau stability and polymerizes microtubules, *J. Neurosci.* **30**,
591–599 (2010).

77. J. Koren, U. K. Jinwal, Z. Davey, J. Kiray, K. Arulselvam, C. A. Dickey, Bending tau into shape:
the emerging role of peptidyl-prolyl isomerases in tauopathies, *Mol. Neurobiol.* **44**, 65–70
(2011).

78. J. Oroz, B. J. Chang, P. Wysoczanski, C.-T. Lee, Á. Pérez-Lara, P. Chakraborty, R. V. Hofele, J.
D. Baker, L. J. Blair, J. Biernat, H. Urlaub, E. Mandelkow, C. A. Dickey, M. Zweckstetter, Structure

and pro-toxic mechanism of the human Hsp90/PPIase/Tau complex, *Nat Commun* **9**, 4532
(2018).

79. J. J. Sabbagh, J. C. O'Leary, L. J. Blair, T. Klengel, B. A. Nordhues, S. N. Fontaine, E. B. Binder,
C. A. Dickey, Age-associated epigenetic upregulation of the FKBP5 gene selectively impairs
stress resiliency, *PLoS ONE* **9**, e107241 (2014).

80. C. L. Klaips, G. G. Jayaraj, F. U. Hartl, Pathways of cellular proteostasis in aging and disease,
J. Cell Biol. **217**, 51–63 (2018).

81. M. C. Inda, S. Joshi, T. Wang, A. Bolaender, S. Gandu, J. Koren Iii, A. Y. Che, T. Taldone, P.
Yan, W. Sun, M. Uddin, P. Panchal, M. Riolo, S. Shah, A. Barlas, K. Xu, L. Y. L. Chan, A. Gruzinova,
S. Kishinevsky, L. Studer, V. Fossati, S. A. Noggle, J. R. White, E. de Stanchina, S. Sequeira, K. H.
Anthoney, J. W. Steele, K. Manova-Todorova, S. Patil, M. P. Dunphy, N. Pillarsetty, A. C. Pereira,
H. Erdjument-Bromage, T. A. Neubert, A. Rodina, S. D. Ginsberg, N. De Marco Garcia, W. Luo, G.
Chiosis, The epichaperome is a mediator of toxic hippocampal stress and leads to protein
connectivity-based dysfunction, *Nat Commun* **11**, 319 (2020).

82. Y. Honjo, T. Ayaki, T. Horibe, H. Ito, R. Takahashi, K. Kawakami, FKBP12-immunopositive
inclusions in patients with α-synucleinopathies, *Brain Res.* **1680**, 39–45 (2018).

83. A. Deleersnijder, A.-S. Van Rompuy, L. Desender, H. Pottel, L. Buée, Z. Debyser, V.
Baekelandt, M. Gerard, Comparative analysis of different peptidyl-prolyl isomerases reveals
FK506-binding protein 12 as the most potent enhancer of alpha-synuclein aggregation, *J. Biol.*
Chem. **286**, 26687–26701 (2011).

84. M. Gerard, Z. Debyser, L. Desender, P. J. Kahle, J. Baert, V. Baekelandt, Y. Engelborghs, The
aggregation of alpha-synuclein is stimulated by FK506 binding proteins as shown by
fluorescence correlation spectroscopy, *FASEB J.* **20**, 524–526 (2006).

85. M. Gerard, Z. Debyser, L. Desender, J. Baert, I. Brandt, V. Baekelandt, Y. Engelborghs, FK506
binding protein 12 differentially accelerates fibril formation of wild type alpha-synuclein and its
clinical mutants A30P or A53T, *J. Neurochem.* **106**, 121–133 (2008).

86. M. Gerard, A. Deleersnijder, V. Daniëls, S. Schreurs, S. Munck, V. Reumers, H. Pottel, Y.
Engelborghs, C. Van den Haute, J.-M. Taymans, Z. Debyser, V. Baekelandt, Inhibition of FK506
binding proteins reduces alpha-synuclein aggregation and Parkinson's disease-like pathology, *J.*
Neurosci. **30**, 2454–2463 (2010).

87. F.-L. Liu, P.-H. Liu, H.-W. Shao, F.-L. Kung, The intracellular domain of amyloid precursor
protein interacts with FKBP12, *Biochem. Biophys. Res. Commun.* **350**, 472–477 (2006).

88. U. Kemmerling, P. Muñoz, M. Müller, G. Sánchez, M. L. Aylwin, E. Klann, M. A. Carrasco, C.
Hidalgo, Calcium release by ryanodine receptors mediates hydrogen peroxide-induced
activation of ERK and CREB phosphorylation in N2a cells and hippocampal neurons, *Cell Calcium*
41, 491–502 (2007).

89. S. E. Lehnart, F. Huang, S. O. Marx, A. R. Marks, Immunophilins and coupled gating of
ryanodine receptors, *Curr Top Med Chem* **3**, 1383–1391 (2003).

90. M. Forgac, Vacuolar ATPases: rotary proton pumps in physiology and pathophysiology, *Nat.*
Rev. Mol. Cell Biol. **8**, 917–929 (2007).

91. J. D. Pearson, D. B. DeWald, W. R. Mathews, N. M. Mozier, H. A. Zürcher-Neely, R. L.
Heinrikson, M. A. Morris, W. D. McCubbin, J. R. McDonald, E. D. Fraser, Amino acid sequence
and characterization of a protein inhibitor of protein kinase C, *J. Biol. Chem.* **265**, 4583–4591
(1990).

92. L. Wang, Y. Zhang, H. Li, Z. Xu, R. M. Santella, I. B. Weinstein, Hint1 inhibits growth and
activator protein-1 activity in human colon cancer cells, *Cancer Res.* **67**, 4700–4708 (2007).

93. O. Huber, J. Weiske, Beta-catenin takes a HIT, *Cell Cycle* **7**, 1326–1331 (2008).

94. N. Korsisaari, T. P. Mäkelä, Interactions of Cdk7 and Kin28 with Hint/PKCI-1 and Hnt1
histidine triad proteins, *J. Biol. Chem.* **275**, 34837–34840 (2000).

95. P. Liu, Z. Liu, J. Wang, X. Ma, Y. Dang, HINT1 in Neuropsychiatric Diseases: A Potential
Neuroplastic Mediator, *Neural Plast.* **2017**, 5181925 (2017).

96. M. P. Vawter, J. M. Crook, T. M. Hyde, J. E. Kleinman, D. R. Weinberger, K. G. Becker, W. J.
Freed, Microarray analysis of gene expression in the prefrontal cortex in schizophrenia: a
preliminary study, *Schizophr. Res.* **58**, 11–20 (2002).

97. M. P. Vawter, C. Shannon Weickert, E. Ferran, M. Matsumoto, K. Overman, T. M. Hyde, D. R.
Weinberger, W. E. Bunney, J. E. Kleinman, Gene expression of metabolic enzymes and a
protease inhibitor in the prefrontal cortex are decreased in schizophrenia, *Neurochem. Res.* **29**,
1245–1255 (2004).

98. Y. Fujiki, A. L. Hubbard, S. Fowler, P. B. Lazarow, Isolation of intracellular membranes by
means of sodium carbonate treatment: application to endoplasmic reticulum, *J. Cell Biol.* **93**,
97–102 (1982).

99. M. M. Savitski, T. Mathieson, N. Zinn, G. Sweetman, C. Doce, I. Becher, F. Pachl, B. Kuster,
M. Bantscheff, Measuring and managing ratio compression for accurate iTRAQ/TMT
quantification, *J. Proteome Res.* **12**, 3586–3598 (2013).

100. S. Meier, M. Bell, D. N. Lyons, J. Rodriguez-Rivera, A. Ingram, S. N. Fontaine, E. Mechas, J.
Chen, B. Wolozin, H. LeVine, H. Zhu, J. F. Abisambra, Pathological Tau Promotes Neuronal
Damage by Impairing Ribosomal Function and Decreasing Protein Synthesis, *J Neurosci* **36**,
1001–1007 (2016).

Acknowledgements: We thank the UF Proteomics and Mass Spectrometry core for proteomic

analysis, the University of Kentucky Magnetic Resonance Imaging and Spectroscopy Center

(MRISC) for MRI analysis. We thank Dr. Peter Davies for his generous contribution of the PHF1

antibody. We thank Dr. Fred Schmitt for insightful discussions and crucial intellectual

contributions to this study. We thank Dr. Shelby Meier for her contributions to the work prior to

this study and experimental support in this current project. **Funding:** This work was supported by

the Alzheimer's Association NIRG-14-322441, Department of Defense AZ140097, NIH/NIMHD

L32 MD009205-01, NIH 1R21NS093440, NIH/NINDS 1R01 NS091329-01. **Author**

Contributions: S.A.K., S.N.F., and J.F.A. designed the project. S.A.K., M.J.H., R.C., S.N.F.,

C.L., J.R.R., A.I., M.B., E.J.M., S.M.G.E., N.Z., N.S., D.T., D.P., M.V., T.F., and A.M.S.H.

performed experiments, acquired data, or assisted in analysis. S.A.K., M.J.H., and J.F.A drafted

the manuscript. J.K., J.M.A., and N.J.L. edited the manuscript. **Competing interests:** GSK

manufactured GSK2606414, which was used in this study. Moreover, this study was funded in

part by a contract from GSK. JMA and NL are employed by GSK. **Data and materials**

availability: Raw proteomic data will be accessible in PRIDE.

Figures

Figure 1: Multi-target kinase inhibition with GSK2606414 rescues brain atrophy in rTg4510 tau transgenic mice.

A Timeline of tauopathy phenotype in transgenic (Tg) mice and depiction of drug delivery between 4-5mo.

B-E Representative T2-weighted MR images of non-transgenic (NTg) and Tg mice treated with vehicle (Veh) or GSK2606414 (414).

F Total brain volume quantification (NTg + 414, n = 2; others n = 3). Two-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean ± SEM, **p < 0.01.

1001

Figure 2: GSK2606414 treatment rescues brain dysfunction in tau transgenic mice.

A-D Representative ΔR1-rendered MEMRI images. The insets in panels A-D show half of the hippocampus of each representative image.

E-F Calculated ΔR1 values in the dentate gyrus (DG, **E**) and Cannu ammonis 1 (CA1, **F**). Two-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean ± SEM, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, (n = 4-5).

G Results of open field behavioral task indicating percent of time spent in the periphery, normalized to the first five minutes (n = 6-10). Two-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean relative to NTg + Veh ± SEM, $*p < 0.05$.

1011   **H** Y-maze behavioral task results for number of novel arm entries (n = 5-7). Two-way ANOVA
1012   with Tukey *post hoc* test. Data expressed as the mean ± SEM, *$p < 0.05$, & denotes p = 0.1095.

1013

Figure 3. GSK2606414-mediated cognitive rescue in early-stage tau transgenic mice is independent of tau hyper-phosphorylation and GSK3β activity

A Immunoblots of PHF1 (S396/S414) and total tau in 5mo NTg and Tg animals treated with vehicle or GSK2606414 (414).

B-D Results of PHF1 (**B**), total tau (**C**), and relative phosphorylation (**D**) is normalized to GAPDH (n = 3). Two-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean relative to Tg + Veh ± SEM.

E Immunoblots of relative activated pY216 to total GSK3β level.

F Results of relative phosphorylation is provided normalized to actin (n = 3). Two-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean relative to NTg + Veh ± SEM.

G-J Representative images of active, pT980 PERK (pPERK) staining. All images were post-processed equally and highlights lack of positive staining in 5mo Tg mice. Scale bar = 25 microns.

K Quantification of pPERK staining reveals that 414 treatment has no detectable change in 5mo Tg mice (n = 5-7), but strongly inhibits PERK activity in the CA3 and superior medial cortex (Ctx) in 8mo Tg mice (hatched bars, n = 3). Two-way ANOVA with Sidak *post hoc* test. Data are expressed as the mean relative to NTg + Veh ± SEM, ****$p < 0.0001$.

L Relative hippocampal gene expression of direct mediators of the PERK-UPR pathway in 5mo NTg and Tg mice treated for 30d with Veh or 414 normalized to GAPDH and 18S (n = 4-6). Two-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean relative to NTg + Veh ± SEM, *$p < 0.05$, **$p < 0.01$.

M-N Representative images (**M**) and quantification (**N**) of overlap between NRF2 (red) and nuclear staining (DAPI, blue) in the CA3 of 5mo NTg and Tg mice treated with either Veh or 414 (n = 3). Unpaired Student's t-test. Error bars denote SEM. Scale bar = 25 microns.

1040

Figure 4. Hippocampal proteome signatures of early stage tauopathy model mice.

A Schematic of TMT isobaric tag proteomic pipeline. Mice hippocampal sections were extracted and lysed. Individual isobaric tags were used to chemically label proteins from each individual sample and combined for LC/MS-MS detection. Batch normalization was completed using a shared batch correction standard, resulting in n = 4 per group. All samples were from female mice.

B Heatmap of proteomics results showing 337 significantly altered proteins across all comparisons (FDR adjusted, p < 0.05).

C Summation of the number of significantly altered proteins across each test.

D Representative protein for each test of significant differential abundance.

F-G Network clustering of Reactome pathways found up-regulated (**F**) or down-regulated (**G**) due
to tau transgenic genotype relative to NTg mice (Tg effect). Node size proportionally represents
the protein count in that pathway, whereas node color represents the statistical significance
gradient (FDR-adjusted p value) from p = 0.10 (brown) approaching p = 0 (white). Dashed
connections indicate manually adjusted nodes.

1055

1056

Figure 5. Multi-target kinase inhibition rescues pathways related to cognitive and neuronal function in early-stage tau transgenic mice.

A-B Network clustering of Reactome pathways found up-regulated (**F**) or down-regulated (**G**) in Tg mice and significantly rescued toward NTg + Veh levels in Tg + 414 mice. Node size proportionally represents the protein count in that pathway, whereas node color represents the statistical significance gradient (FDR-adjusted p value) from p = 0.10 (brown) approaching p = 0 (white). Dashed connections indicate manually adjusted nodes.

C-D Puromycin immunostaining of the CA1 region (**D**) and quantified (**E**) relative to total CA1 area (n = 3). One-way ANOVA with Tukey *post hoc* test. Data are expressed as the mean absolute integrated density ± SEM, *p < 0.05. Scale bar = 25 microns.

E-F Immunoblots (e) of total tyrosine nitration and total lane quantified results (**F**) relative to NTg + Veh and normalized to actin (n = 3 - 4). Two-way ANOVA with Tukey post hoc test. Data are expressed as the mean relative to Tg + Veh ± SEM, **p < 0.01.

1070
1071

Figure 6: Proteomic signatures of rTg4510 transgenic mice match human tauopathy and reveal novel candidates for therapeutic intervention.

A-B Significantly altered proteins by Tg effect were compared to human homolog abundances found in two proteomics analyses of human neurodegenerative diseases (**A**, Seyfried et al., 2016; **B**, Ping et al., 2018). Proteins annotated in the "Rescued" column (yellow) were significantly rescued by 414 in Tg animals compared to NTg controls. Human homolog protein abundances in each human disease group were compared to non-demented controls. Statistically different human proteins were annotated by disease group in different colors (Asym AD, pink; late AD, purple; late PD, green). Proteins that were significantly altered in the same direction (increased or decreased) between species were annotated in the "Consistent" column (black). Candidate proteins passing all criteria for inclusion were labeled with their corresponding human gene symbols. Sample abundances were averaged per group and each corresponding sample number was appended to the bottom of each column.

C Venn diagram comparing candidate proteins identified in late AD across both datasets.

1089
1090   **Supplementary Figures:**

SFig. 1

SFig. 2

SFig. 3

1091
1092

1093   **Fig. S1. Total duration spent in novel arm is partially rescued in tau transgenic mice treated**
1094   **with GSK2606414.** Total time was measured during mouse exploration in the Y-maze behavioral
1095   task. Tg + Veh mice exhibited reduced time spent in the novel arm (*p = 0.0433). Tg mice treated
1096   with 414 were only partially rescued compared to Tg + Veh mice (p = 0.0882). Data were
1097   compared using two-way ANOVA with Tukey *post hoc* test.

1098

1099   **Fig. S2. Non-PERK UPR proteins do not have altered transcript levels in 5mo tau transgenic**
1100   **mice.** Other eIF2a kinases do not have altered transcript levels in 5mo Tg mice. *ATF6* transcript
1101   levels decrease in Tg mice compared to NTg controls, though 414 has no effect. Transcript levels
1102   normalized to GAPDH and 18S (n = 4-6). Two-way ANOVA with Tukey *post hoc* test. Data are
1103   expressed as the mean relative to NTg + Veh ± SEM, *p < 0.05, **p < 0.01.

1104

Fig. S3. GSK2606414 treatment for 30d does not cause marked weight loss. Prolonged treatment with GSK2606414 induces weight loss in mice *(12)*. However, treatment for only 30d was not sufficient to alter weight loss using a 100mg/kg treatment paradigm.

Table S1. Top kinase targets of GSK2606414 and relevance to tauopathy.

The top targets of GSK2606414 with experimentally determined IC50 are listed based on reported data *in vitro (10)* or based on cell culture experiments *(17, 18)*. Relevance to tauopathy is briefly summarized.

Table S2. Significantly enriched reactome pathways identified by proteomics. Reactome pathways significantly altered (FDR adjusted $p < 0.10$) by either direction in Tg mice compared to NTg controls along with pathways determined rescued by 414.

Table S3. Human to mouse proteomic comparisons.

Table S4. Patient demographics of human proteomic samples.

Patient demographics and disease-related pathology scores were averaged and compiled from original reports *(44, 46)*.